1   **Vasopressin 1a receptor antagonist disrupts male-male affiliative relationships formed by**
2   **triadic cohabitation in large-billed crows**
3

4   Akiko Seguchi [1,2,3*], Ei-Ichi Izawa [1]
5
6   [1] Department of Psychology, Keio University, Tokyo 108-8345, Japan
7   [2] Art Future Research Field, Tokyo University of the Arts, Tokyo 110-8714, Japan
8   [3] Lead contact
9
*Correspondence:
Akiko Seguchi (akiko.seguchi@keio.jp)
Mita 2-15-45, Minato-ku, Tokyo 108-8345, Japan
Tel: +81-90-5327-3213

Abstract

Same-sex affiliative relationships are common in humans and some social animals, forming one of the

bases of group living. The neuropeptide vasopressin (VP) and its receptors mediate these relationships

and behaviours in mammals and birds with gregarious and colonial social structures. In some species,

affiliative relationships between dominant and subordinate individuals can be maintained while still

retaining strict dominance hierarchies where three or more individuals interact. However, it is unclear

whether triadic interaction promotes these relationships, and whether the VP system is also involved

in such affiliations due to the lack of suitable animal models and experimental settings. This study

addresses these questions with two experiments. In Experiment 1, two-week cohabitation among three

male crows facilitated affiliative relationships in particular dyads within each triad. In Experiment 2,

vasopressin 1a receptor (V1aR) antagonism disrupted affiliative behaviours and led to the resurgence

of agonistic behaviours in affiliated males but not in unaffiliated ones by peripherally administering a

V1aR antagonist. These findings suggest that the VP system might universally mediate same-sex

affiliative relationships, despite differences in inherent aggression levels among individuals. The

triadic cohabitation paradigm established here could advance our understanding of animal societies

and be applied across various species, sexes, and social structures.

Impact statement

This study provides evidence that a triadic interactive environment facilitates the formation of

affiliative relationship between specific dominant and subordinate males in large-billed crows, with

crucial involvement of vasopressin 1a receptor in maintaining this affiliation.

Introduction

Affiliative relationships, such as parent-offspring bonds (Clutton–Brock, 1991; Royle et al., 2012), male-female pair-bonds (Black, 1996; Kleiman, 1977; Bales et al., 2021), and same-sex platonic bonds (Brent et al., 2014), are beneficial for individuals living in groups as they increase the fitness of resource acquisition and, ultimately, reproductive success (Ziegler & Crockford, 2017). Data on the functions and neuroendocrine system of same-sex affiliative relationships, characterised by the exchange of non-sexual affiliative behaviours between same-sex individuals, including social grooming (allogrooming in mammals and allopreening in birds), huddling, spending time together, and sharing food, has predominantly have been obtained from studies on mammals and birds (Brent et al., 2014; Ziegler & Crockford, 2017; Beery, 2019). These relationships are not limited to one sex or kinship, as they have been observed in males and females, as well as in kin and non-kin, both in captivity and wild conditions (Ziegler & Crockford, 2017). Same-sex affiliative relationships are also frequent in humans, existing throughout most of our history and in many societies and cultures (Brent et al., 2014).

The neural mechanisms underlying affiliative relationships might have been co-opted from parent-offspring bonds into more flexible bond structures, such as pair bonds and same-sex affiliative relationships (Broad et al., 2006; Crespi, 2016; Numan & Young, 2016). The role of neuropeptides of the arginine vasopressin family (arginine vasotocin, AVT, in non-mammals) and oxytocin (OT) signals mediated V1aR has been implicated in same-sex affiliative relationships. This has been suggested by previous studies specifically in males and females of gregarious birds and females of voles. These studies primarily used paradigms that investigated "gregariousness" in finches as a preference for the larger of two groups, and partner preference in voles as a choice of familiar over unfamiliar. Goodson and Wang (2006) found that VT neurons in the medial bed nucleus of the stria terminalis were activated, in terms of immediate early gene Fos expression, in response to the exposure of same-sex conspecifics through the wire barrier in males and females of gregarious species, but were not in asocial ones. The authors also indicated that the activation of VT neurons is specifically induced by social stimuli with a positive valence to enhance affiliative behaviour (Goodson & Wang, 2006). Antisense-based knockdown of VT production in the medial bed nucleus of the stria terminalis of male zebra finches, a gregarious species, increased aggressive behaviour and reduced courtship in colony housing situations (Kelly & Goodson, 2013). V1aR antagonism in the LS of male zebra finches reduces flock size in housing conditions (Kelly et al., 2011). A previous study

showed that in female meadow voles (*Microtus pennsylvanicus*), a season-specific social species, the co-administration of OT and a vasopressin 1a receptor (V1aR) antagonist in the LS inhibited female-female partner preference for huddling behaviour with cagemates over unfamiliar individuals (Anacker et al., 2016). Research into the physiological underpinnings of same-sex affiliation in voles is extensive and includes studies on neurotransmitters beyond VP/OT, such as the possible relevance of estradiol in meadow voles (Beery et al., 2008) and dopamine signalling in prairie voles (*M. ochrogaster*) with female-female partner preference (Lee & Beery, 2021; Lee et al., 2023).

However, among same-sex affiliative relationships, there is a type where individuals selectively affiliate with specific individuals despite the presence of multiple partners. Particularly, adult males can simultaneously hold a position within strict dominance relationships and form affiliative relationships between specific dominant and subordinate individuals. Such relationships, including coalitions and alliances, have been observed in various animal species (including humans, *Homo sapiens* (Macfarlan et al., 2014; von Rueden et al., 2008; von Rueden & Jaeggi, 2016; Patton, 2005); chimpanzees, *Pan troglodytes* (Watts, 2002; Muller & Mitani, 2005; Gilby et al., 2013); baboons, *Papio* (Bercovitch, 1988; Noë, 1994); macaques, *Macaca radiata* (Silk, 1999); *M. fuscata* (Kutsukake & Hasegawa, 2005); cetaceans bottlenose dolphins, *Tursiops truncates* (Connor et al., 1992a; Connor et al., 1992b; Connor et al., 1999); spotted dolphins, *Stenella frontalis* (Green et al., 2015); sperm whales, *Physeter macrocephalus* (Kobayashi et al., 2020); carnivores like lions, *Panthera leo* (Bygott et al., 1979); cheetahs, *Acinonyx jubatus* (Caro, 1993; Caro, 1994); birds, corvids (Fraser & Bugnyar, 2012). Most of these studies indicated that male-male affiliative relationships were formed in social interactions involving multiple participants, typically two individuals vs. one individual, in both the wild and captivity. Theoretical research exploring the social factors of coalition formation predicted that social interactions among three individuals differing in "power" and "activity" should be the minimal group size to produce coalition/alliance (Caplow, 1956; Mesterton-Gibbons et al., 2011; Koykka & Wild, 2017). Several previous studies with human participants have examined the combinations of ranks that facilitate the formation of coalitions in game situations (Vinacke & Arkoff, 1957; Kelley & Arrowood, 1960). If replicated in experimental research on social animals, the formation of affiliative relationships between males within these three-individual interactions can facilitate studies on the physiological basis of such relationships.

Crows and related species (i.e., *Corvus* spp) are an ideal model to investigate the ecological factors and physiological basis of male-male affiliative relationships. Non-breeding crows form

groups during foraging, roosting, and socialising, with group sizes ranging from a few to several

hundred or even thousands of birds and with the composition of group members changing during the

day and across contexts (Boucherie et al., 2019; Izawa, 2011; Loretto et al., 2017; Uhl et al., 2019).

Despite the high degree of fission-fusion dynamics, temporarily stable sub-groups can be observed.

For example, certain birds use the same food source or roost repeatedly over consecutive weeks or

years (Braun & Bugnyar, 2012). Additionally, individuals form multiple highly differentiated

affiliative and agonistic relationships with others that comprise various qualities (Fraser & Bugnyar,

2010a). Males form and maintain stable dyadic dominance relationships and, in a group setting, a

strict linear hierarchy, whereas those of females are less robust (Fraser & Bugnyar, 2010a; Izawa &

Watanabe, 2008; Nishizawa et al., 2011; Ode et al., 2015; Takeda et al., 2022).

In both captive and wild populations, affiliative relationships are formed between unpaired adults

and unmatured juveniles by exchanging affiliative behaviour such as allopreening, food giving, and

conflict aiding (Bugnyar, 2012; de Kort et al., 2006; Emery et al., 2007a; Scheid et al., 2008; von

Bayern et al., 2007; Picard et al., 2020). Male-male and male-female dyads tend to form stronger and

more stable affiliative relationships than female-female dyads (Fraser & Bugnyar, 2010a; Miyazawa

et al., 2020). The formation of affiliative relationships between individuals increases the dominance

rank and likelihood of success in competing for food in captive groups of rooks (*Corvus frugilegus*)

(Emery et al., 2007b) and wild common ravens (*C. corax*) (Braun & Bugnyar, 2012; Braun et al.,

2012). These findings suggest an advantage of the affiliative relationship in group lives. Ravens and

rooks reconcile with and console valuable partners (Fraser & Bugnyar, 2010b; Fraser & Bugnyar,

2011; Seed et al., 2007), remember former group mates and their relationships with them over years

(Boeckle & Bugnyar, 2012), understand third-party relations (Massen et al., 2014a), and prevent

others from forming too strong alliances (Massen et al., 2014b). Moreover, ravens cooperate in the

wild; for example, they support each other in conflicts (Fraser & Bugnyar, 2012), hunt cooperatively

(Hendricks & Schlang, 1998) and cooperatively chase away larger predators or dominant male-female

pairs from prey items (Heinrich & Marzluff, 1991; Marzluff et al., 1996). Like other *Corvus* species,

non-breeders of large-billed crows (*C. macrorhynchos*) form fission–fusion societies, with flexible

changes in the social structure depending on the time of day, seasons, and their age (Izawa, 2011;

Islam et al., 2010; Kuroda, 1990). In captive non-breeder groups, males strictly form a linear

dominance hierarchy (Izawa & Watanabe, 2008; Ode et al., 2015) and allopreening often occurs

between males (Miyazawa et al., 2020). This allopreening is primarily initiated by dominants but also

occurs from subordinates (Miyazawa et al., 2020).

A triadic interactive scenario is an ecological factor that facilitates the formation of affiliative

relationships between dominant and subordinate males. Previous studies have suggested the

involvement of V1aR in same-sex affiliative relationships; therefore, in this study, we hypothesised

that cohabitation of three male non-breeder large-billed crows could facilitate the formation of

affiliative relationships between specific dominant and subordinate individuals and that V1aR is

involved in the maintenance of these relationships. To this end, we conducted two experiments to test

our hypotheses.

In Experiment 1, we introduced three non-breeder males of captive large-billed crows, which had

formed dominance relationships but not affiliative relationships prior to this study, to a triadic

cohabitation in an aviary for two weeks. We examined the formation of an affiliative relationship in

one of the three possible dyads by comparing the occurrence of reciprocal allopreening before and

after cohabitation. As we predicted, the two-week triadic cohabitation facilitated the formation of an

affiliative relationship exclusively between two males in the triad. The two-week triadic cohabitation

paradigm developed in this study enabled us to investigate the physiological mechanisms of male-

male affiliative relationships in a laboratory setting.

In Experiment 2, we investigated the effects of peripheral administration of a V1aR antagonist on

the social behaviours of males with affiliative relationships formed via triadic cohabitation. Our

prediction was that blocking V1aR would reduce allopreening and increase aggressive and submissive

behaviours among dominant and subordinate males with affiliative relationships but not among those

without such relationships. We discovered that administering a V1aR antagonist disrupted the

affiliative relationships between males, leading to more aggressive and submissive behaviours from

dominants and subordinates. However, it had no effect on the social behaviour of males without

affiliative relationships. In conclusion, our study suggests that living in a group of three promotes the

formation of close relationships between males and that V1aR is involved in maintaining these

relationships by promoting allopreening and inhibiting aggressive and submissive behaviours based

on dominance.

Results

Experiment 1: The triadic cohabitation facilitated the formation of an affiliative relationship

between two particular males

To determine whether triadic cohabitation among males promotes the formation of affiliative

relationships between specific dyads, we introduced eight groups of three male crows, each already

having established dominance relationships but no affiliative relationships, into an aviary for two

weeks (Figure 1). We compared social behaviour between males before and after cohabitation to

evaluate the formation of affiliative relationships characterised by reciprocal allopreening exclusively

between two of three males in each triad.

Figure 1. Experimental design.

(A) Experiment 1: Triadic cohabitation and formation of affiliative relationships. Step 1: three male crows with already established relative dominance relationships through each dyadic encounter but no affiliative relationships were introduced into an aviary for two weeks. These three crows are referred to as high (orange), middle (green), and low (blue) based on their relative dominance. Step 2: pre-test dyadic cohabitation for 3 days to ensure any affiliative relationships noted were not due to prior dyadic interactions. Step 3: triadic cohabitation for two weeks to observe the formation of affiliative relationships, characterised by reciprocal allopreening between two specific crows in each triad. Step 4: post-test dyadic cohabitation for 3 days to confirm the persistence of these relationships. If no affiliative relationship was formed in any dyad, that triad would not proceed to Experiment 2, the behavioural pharmacological experiment.

(B) Experiment 2: Effects of V1a receptor antagonism on dominant-subordinate affiliative relationships. Steps 5 and 6: V1aR antagonists and control vehicle solutions were peripherally administered to one of the two individuals with established affiliative or unaffiliative relationships in Experiment 1, before their dyadic interaction. For each trial, the vehicle was administered to either the dominant or subordinate, followed by free interaction in an aviary. After completing all vehicle trials, the antagonist was administered under the same protocol. Allopreening, aggressive behaviours and submissive vocalisation, were coded for analysis to discern the influence of the V1aR antagonist on the maintenance of affiliative behaviours and the emergence of aggression or submission.

Before this experiment, dominance relationships of all dyads used in this study were assessed

using a social dyadic encounter paradigm from previous studies (Nishizawa et al., 2011; Takeda et al.,

2022). Briefly, two birds were introduced into an aviary to freely interact for 5 min in three trials

within a 2-day inter-trial interval. In each trial, the individual exhibiting submissive behaviours in

response to aggressive behaviours from the opponent was identified as the loser, while the other as the

winner. Within the dyads, the individual who won all three trials was defined as dominant, while the

others as subordinates. All 26 dyads comprising the eight triads tested in this study showed a clear

asymmetry in win/loss numbers, indicating established dominance relationships in all the dyads

(Table 1). In these dyadic encounter trials, we also measured the number of alloopreening and

confirmed none occurred (Figure 2, Figure 2—source data 1). These results indicated that dominance,

but no affiliative, relationships were established in all the dyads before Experiment 1.

Figure 2. Determination of relative dominance in dyads.

The numbers of aggressive (A), submissive (B) and allopreening behaviours (C) across the three

trials of dyadic social encounters. The numbers of aggressive behaviours (jab, peck, aggressive

vocalisation and displacing approach) by the dominant (thick solid black line) and submissive

behaviours (submissive begging vocalisation and avoidance) by the subordinates (thick dotted

black line) decreased through the three encounters, suggesting the stable formation of dyadic

dominance relationships. Allopreening behaviours were not observed in any dyads. Grey-

coloured solid and dotted lines indicate dominant and subordinate individual data, respectively.

Data are represented with mean ± s.e.

Prior to triadic cohabitation (the pre-test), we housed all potential dyads together in the

experimental aviary for three days to ensure that any affiliative relationships observed were not due to

prior dyadic interactions. This was followed by a 2-week triadic cohabitation to determine whether

such interactive situations would encourage affiliative relationships, evidenced by increased

reciprocal allopreening between two specific birds in each triad. To confirm the persistence of these

relationships, a follow-up dyadic cohabitation (the post-test) was conducted after the initial 2-week

period.

Affiliative relationships were defined by the exchange of reciprocal allopreening between two

birds, with each individual initiating allopreening at least 10 bouts per 30 min in any of the three 3-

day blocks (early, middle and last) during the triadic cohabitation. Dyads not reaching this threshold

were considered unaffiliated. Discriminant analysis, with leave-one-out cross-validation, was

employed to differentiate the numbers of dominant-initiated and subordinate-initiated allopreening

between the pre-test and post-test and to validate the exclusivity of the affiliative relationships formed

during triadic cohabitation.

In six of the eight triads, the 2-week cohabitation resulted in the formation of an affiliative

relationship exclusively in one dyad but not in the other dyads (Figure 3A, Figure 3—source data 1,

Video 1). A specific dyad in each triad (#1–#6), which showed no allopreening in the pre-test,

exhibited an increase in allopreening through the 2-week triadic cohabitation. In the post-test of these

six dyads, allopreening was initiated by both dominants and subordinates, whereas no reciprocal

allopreening occurred in the pre-test, and no such allopreening emerged in any other dyads in either

test (Figure 3A, Figure 3—source data 1). Discriminant analysis based on the number of allopreening

initiated by dominants and subordinates within the dyads yielded two discriminant functions. Using

the leave-one-out method, the two discriminant functions distinguished the affiliated dyads in the

post-test from themselves in the pre-test and from the other unaffiliated dyads with an 83.3% correct

rate (Figure 3B, Figure 3—source data 2, i.e., five of six dyads were classified as affiliated dyads; see

the outputs for Table 2). These results indicate that the 2-week triadic cohabitation facilitates the

formation of affiliative relationships exclusively between two of the three birds, characterised by

reciprocal allopreening.

Due to a physical injury in one bird, five of the six dyads forming an affiliative relationship were

used for the test in Experiment 2 (Table 1).

Figure 3. Triadic cohabitation facilitated the formation of an affiliative relationship between two particular males.

(A) Total number of allopreening behaviours in each dyad across the experimental periods: pre-tests, 2-week triadic cohabitation (early, middle, and last), and post-tests. Data plots are presented as mean ± 1 s.e. Panels indicate triads (#1−#8). Bright-coloured plots (pink, green, purple, blue, orange and red) in the #1−#6 triads represent dyads that showed the formation of affiliated relationships. Darker plots denote the dyads that did not form affiliative relationships.

(B) Scatter plots of allopreening behaviours initiated by dominants (x-axis) versus subordinates (y-axis) of dyads in the pre-tests (opened) and the post-tests (shaded). Coloured plots represent dyads wherein affiliative relationships were formed. Allopreening occurred exclusively in the coloured dyads after the cohabitation (i.e., post-test). The colours correspond to the dyad in Figure 3A. Data are presented as the mean with natural log-transformed values.

Experiment 2: V1a receptor antagonism disrupted dominant-subordinate affiliative relationships

To determine the role of V1a-like receptors in maintaining male-male affiliative relationships, crows

with established affiliative or unaffiliative relationships from Experiment 1 were systematically given

a V1aR antagonist to see how it affected their social interactions with affiliated and unaffiliated crows

(see Materials and Methods Exp2). The antagonist was administered in high and low doses, as well as

a control vehicle solution, through intramuscular injections. For each trial, the antagonist or vehicle

was given to either the dominant or subordinate crow an hour before they were observed in the aviary,

where they interacted freely. Social behaviours such as allopreening, aggressive behaviours, and

submissive vocalisations were recorded from video data for analysis. This procedure aimed to

understand the influence of the V1aR antagonist on the maintenance of established affiliative

behaviours and the emergence of aggression or submission.

Administration of the V1aR antagonist significantly reduced allopreening in the affiliated dyads

involving both dominants and subordinates (Figure 4A, Figure 4—source data 1, Video 2). The best

model of the generalised linear mixed model (GLMM) analysis showed that the drug condition had a

significant effect, with the vehicle condition having a positive coefficient (estimate $\pm$ s.e. = 1.639 $\pm$

0.217, $p < 0.001$) and the high dose condition a negative coefficient (estimate $\pm$ s.e. = -0.724 $\pm$ 0.313,

$p < 0.05$; Table 3). There was no significance of the dominance and the interaction factors. These

results indicate that the number of allopreening bouts decreased dose-dependently among the three

conditions, irrespective of the dominance position in the affiliated dyads. However, no effect was

found on allopreening by V1aR antagonist administration in any birds of the unaffiliated dyads

(Figure 4A). GLMM analysis was not performed on unaffiliated dyads because no allopreening

occurred under any condition or with any bird.

The V1aR antagonist increased aggression from dominant birds to subordinate birds but not from

subordinate birds to dominant birds (Figure 4B, Figure 4—source data 1, Video 2). The most reliable

model from the GLMM analysis on aggressive behaviours in the affiliated dyads indicated a

significant impact of the drug condition, especially the high-dose condition with a positive coefficient

(estimate $\pm$ s.e. = 0.791 $\pm$ 0.251, $p < 0.01$), but no significance for the vehicle condition ($p = 0.54$;

refer to Table 4). This suggests a substantial rise in aggression from dominant birds after receiving a

high dose of the V1aR antagonist. However, there was no similar increase in aggression due to the

V1aR antagonist in subordinate birds (see Figure 4B; GLMM was not conducted due to the absence

of aggression in any condition or with any bird). Furthermore, the V1aR antagonist did not increase

aggression in unaffiliated dyads (*Vehicle*, $p = 0.929$; *V1aRA high-dose*, $p = 0.89$; Figure 4B and Table

5), while in the affiliated dyads it increased the submissive begging vocalisation of the subordinates

but not the dominants (Figure 4C, Figure 4—source data 1, Video 2). The best model of the GLMM

analysis of the submissive vocalisations in affiliated dyads showed a significant effect of the *drug*

condition, specifically the high dose condition with a positive coefficient (estimate ± s.e. = 1.929 ±

0.253, $p < 0.001$), but no significance for the vehicle condition ($p = 0.455$; Table 6). In contrast, no

such increase in submissive vocalisations was found in any condition in the unaffiliated dyads

(*Vehicle*, $p = 0.791$; *V1aRA high-dose*, $p = 0.777$; Figure 4C and Table 7).

Figure 4. V1a receptor antagonism disrupted dominant-subordinate affiliative relationships.

(A) Effects of V1a receptor antagonist (V1aRA) on the number of allopreening behaviours in affiliated and unaffiliated dyads. In affiliated dyads, allopreening initiated by the dominant (DOM, left panel) and the subordinate (SUB, right panel) crows was markedly reduced in the V1aRA *high-dose* conditions and slightly in the *low-dose* condition compared with the *vehicle* condition. In unaffiliated dyads, V1aRA had no effects on allopreening by either the dominant (DOM, left panel) or the subordinate (SUB, right panel).

(B) Effects of V1aRA on the number of aggressive behaviours in affiliated and unaffiliated dyads. In affiliated dyads, aggressive behaviour initiated by the dominant bird (DOM, left panel) markedly increased in the V1aRA *high-dose* condition but not in the *low-dose* condition. This increase in aggression did not occur in any condition for the subordinates (SUB, right panel). No effects were observed in any condition in both the dominants and the subordinates of unaffiliated dyads.

(C) Effects of V1aRA on the number of submissive begging vocalisations in affiliated and unaffiliated dyads. In affiliated dyads, submissive begging calls initiated by the dominant (DOM, left panel) did not change in any condition. However, those initiated by the subordinates (SUB, right panel) increased in the V1aRA *high-dose* condition but not in the *low-dose* condition. In unaffiliated dyads, no effects were observed in any condition in both the dominants and the subordinates.

Plots in (A)–(C) show the individual data for each trial. In each triad, circles represent birds affiliated with another male, while triangles represent those with no affiliation. The top, middle, and bottom of the boxes depict the 75th, 50th and 25th percentiles, respectively, with

whiskers representing the maximum and minimum values. Numbers in the brackets below

the boxes denote the number of samples (no. of birds × no. of trials). Asterisks indicate

significant differences (Wald χ^2 statistics, *** $p < 0.001$). Colours correspond to the affiliated

dyads in Figure 3.

Discussion

The present study suggested that in Experiment 1, triadic cohabitation facilitated the formation of an

affiliative relationship exclusively between specific dominant and subordinate male crows within each

triad. In Experiment 2, the peripheral administration of a V1aR antagonist disrupted these affiliative

relationships, leading to a resurgence of aggressive and submissive/avoidant behaviours depending on

the relative dominance but had no effect on social behaviours within unaffiliated dyads. Our findings

suggest that the triadically interactive environment facilitative the formation of affiliative relationships

between dominant and subordinate males. Moreover, the V1aR, previously implicated in same-sex

affiliative relationships in studies on voles and finches, is essential in maintaining these types of male-

male affiliative relationships.

Experiment 1: The triadic cohabitation facilitated the formation of an affiliative relationship

between two particular males

The results of Experiment 1 suggest that a triadically interactive situation plays a crucial role in

facilitating the formation of affiliative relationships exclusively between two specific dominant and

subordinate males of the three. Allopreening behaviour rarely occurred the dyadic cohabitation (i.e.,

pre-triadic cohabitation) but did frequently and exclusively between two specific males during triadic

cohabitation. This finding suggests that triadic interactions are crucial for facilitating allopreening and

the formation of affiliative relationships. Our finding aligns with previous theoretical studies (Caplow,

1956; Mesterton-Gibbons et al., 2011; Koykka & Wild, 2017), indicating that triad, including

individuals with different power dynamics, are the minimal social groups that promote the formation

of coalitions in the form of two against one.

This two-vs.-one form has been confirmed in previous behavioural observation studies on same-

sex coalitions and alliances of primates, cetaceans, and corvids in the wild and in captivity (Macfarlan

et al., 2014; von Rueden et al., 2008; von Rueden & Jaeggi, 2016; Patton, 2005; Watts, 2002; Muller

& Mitani, 2005; Gilby et al., 2013; Bercovitch, 1988; Noë, 1994; Silk, 1999; Kutsukake & Hasegawa,

2005; Connor et al., 1992a; Connor et al., 1992b; Connor et al., 1999; Green et al., 2015; Kobayashi

et al., 2020; Bygott et al., 1979; Caro, 1993; Caro, 1994; Fraser & Bugnyar, 2012). Our findings in

Experiment 1 provide experimental evidence that a triadically interactive situation, such as 2-week

cohabitation, facilitates the formation of affiliative relationships between dominant and subordinate

males, bridging the gap between previous theoretical and behavioural observation studies.

The function of male-male affiliative relationships formed in Experiment 2 remains to be

elucidated, but several possibilities warrant discussion. Generally, having affiliative partners may be

beneficial in various ways. For large birds like crows, where predation pressure is relatively low

(Braun et al., 2012), these relationships may primarily enhance foraging efficiency (Beauchamp,

2010; Dall & Wright, 2009). Experiments with captive *Corvus* species show that the presence of

bonded individuals affects some behaviours, such as exploring new objects (Stöwe et al., 2006),

paying attention to others (Scheid et al., 2007) and social learning (Schwab et al., 2008). Proximately,

affiliation may serve as a social manoeuvre facilitating resource access and may assess the quality of a

partner in these long-term monogamous birds (Braun & Bugnyar, 2012).

Another possibility is that crows use such affiliative interactions to maintain and intensify

valuable social relationships, enabling coalition formation (de Waal & Luttrell, 1988). Studies in other

Corvus species have shown a significant correlation between preening and agonistic support,

suggesting that partners who engage in mutual preening are more likely to support each other in

aggressive interactions (Scheid et al., 2007). Such behaviour can increase social rank for both

individuals involved (Emery et al., 2007b). Common ravens focus on a few close partners at a time,

aligning with the selective affiliative relationship formation observed in this study. Although agonistic

support and subsequent rank reversals were not observed, this does not entirely rule out the possibility

of coalition functions. Although agonistic support and subsequent rank reversals were not observed in

our study, this does not entirely rule out the possibility of coalition functions. Aggressive support is

much rarer than preening or grooming, making the opportunity to reciprocate support within a day

very low. Ravens have been shown to reciprocate support over two years rather than within a week,

reflecting the likelihood of long-term service exchange (Fraser & Bugnyar, 2012). Thus, coalition-like

behaviours might not have been observed due to our study's specific context and observation period

but might be evident over longer time spans or different contexts. Future experiments could test the

coalition function in male-male affiliative relationships by confronting affiliated dyads formed in

triadic cohabitation with a male-female pair or by introducing them into a captive group to observe

rank changes. The high levels of affiliative interactions maintained during post-dyadic cohabitation

suggest potential for long-term alliances. Long-term collaborations in conflicts, referred to as

alliances, contrast with facultative short-term coalitions (Harcourt & de Waal, 1992). Our study

implies that the affiliative relationships formed in triadic cohabitation may constitute alliances,

warranting further investigation into their stability when a third-party individual is introduced or when

the dyad is integrated into larger groups.

Male-male affiliative relationships in our crows may also function as social bonds similar to those

seen in mother-offspring or pair bonds, potentially reducing social stress. The presence of affiliative

individuals mitigates stress responses, a phenomenon known as social buffering (Kikusui et al., 2006).

Many animals, including humans, exhibit smaller increases in stress hormone (cortisol) levels when

exposed to aversive stimuli in the presence of an affiliative partner compared to when alone (de Vries

et al., 2003). In primates, social grooming reduces heart rate (Aureli et al., 1999), and individuals with

tightly-knit social networks have lower baseline levels of cortisol metabolites in their faeces (Brent et

al., 2011; Crockford et al., 2008). Following concepts developed in primatology (Hinde, 1976), social

bonds in Corvids are defined by the reciprocal exchange of affiliative behaviours over time, observed

across all age classes and between both male-female and male-male dyads (Braun et al., 2012). In

ravens, individuals with high levels of positive social connections within their group show lower

corticosterone levels when within the group and higher levels when separated (Stocker et al., 2016).

Affiliated dyads in our study might have reduced stress from attacks by the dominant and the stress of

living in confined cages through mutual preening. In large-billed crows, methods for measuring

corticosterone in both faeces and saliva using enzyme-linked immunosorbent assay have been

established (Ode et al., 2015; Aota et al., 2023). Future studies should compare corticosterone levels

in saliva or faeces during both the early and last periods of triadic cohabitation to determine whether a

decrease in corticosterone accompanies the formation of affiliative relationships. Additionally,

examining changes in corticosterone levels during separation from and reunion with affiliative

partners would provide further insights into the stress-buffering role of these relationships.

While numerous challenges remain in understanding the functions of male-male affiliative

relationships in large-billed crows, the hypotheses discussed above are not mutually exclusive.

Applying triadic cohabitation experiments as described can provide valuable insights through careful

and systematic testing. Moreover, these experiments can help to identify predictors of affiliative

relationship formation among males. Several studies have concluded that personality similarity or

homophily is a strong predictor of affiliative relationships, potentially facilitating cooperation

(Verspeek et al., 2019; Laakasuo et al., 2020; Noë et al., 2016; Massen & Koski, 2014). By collecting

data on inter-individual behavioural differences or personalities, such as exploration tendencies and

stress vulnerability before triadic cohabitation, we can examine which characteristics, beyond social

rank, kinship or age, predict affiliative relationships. Triadic cohabitation experiments enable

physiological exploration and rigorous testing of predictors, enhancing our understanding of the

mechanisms underlying affiliative relationship formation.

Two of the eight triads tested in the present study did not form affiliative relationships even after

2-week cohabitation. As illustrated in Figure 3, allopreening occurred slightly in two males of the two

triads during the cohabitation period but eventually disappeared. According to the video observation

of behaviours during the cohabitation period, one possible cause for the failure to form affiliative

relationships in these triads is that one male avoided the initiation of allopreening by the other male.

Another possible cause is the intervention by the third-party male when the two males attempted to

engage in socially affiliative interactions. The intervention by third-party individual in affiliative

interactions between other individuals has been found in a captive group of ravens (Massen et al.,

2014b). The questions of what social characteristics of individuals in dyads could facilitate or inhibit

allopreening and what individual and/or social factors could drive the intervention in others' affiliative

behaviour in birds and mammals remain to be addressed. An in-depth analysis of social behaviour

using the triadic cohabitation paradigm used in this study may elucidate the behavioural and

psychological mechanisms for forming male-male affiliative relationships.

V1a receptor antagonism disrupted the dominant-subordinate affiliative relationship

Systemic injection of a V1aR antagonist to males with affiliative relationships caused a dose-

dependent decrease in allopreening and the resurgence of agonistic behaviours associated with

relative dominance, such as aggressive behaviour in the dominant and submissive behaviour (i.e.,

submissive begging calls) in the subordinate. In contrast, dominant and subordinate males without an

affiliated relationship showed no behavioural effects from V1aR blocking. These results suggest that

V1aR is involved in the suppression of dominance-dependent aggressive/submissive behaviour and

facilitating affiliative behaviour, thus playing a crucial role in the formation and maintenance of

affiliated relationships between dominant and subordinate males.

Given the peripheral administration of the drug in this study, the behavioural effects found in

affiliated dyads from V1aR antagonists do not necessarily reflect the role of V1aR in the brain.

Peripheral nonapeptide receptors have been suggested to influence a variety of homeostatic functions,

such as hydromineral balance (Goldstein, 2006) and thermoregulation (Hassinen et al., 1999). For

example, maternal crouching in rodents serves a thermoregulatory function (Jans & Woodside, 1990)

and, is regulated by V1aR (Bosch & Neumann, 2008).

V1aR has also been suggested to be involved in peripheral thermoregulation (Pittman et al., 1998;

Shido et al., 1984; Yang et al., 2006). Thus, the effects of maternal crouching by V1aR blocking in

rodents may be indirectly due to thermoregulation via V1aR antagonism. Similarly, maternal licking

and grooming in rodents may indirectly involve peripheral V1aR (Bosch & Neumann, 2008). Given

that during maternal licking and grooming, mothers often ingest water to maintain the hydromineral

balance (Gubernick & Alberts, 1983), based on peripheral V1aR (Koshimizu et al., 2012), maternal

licking and grooming may be regulated indirectly by peripheral V1aR. Although the distribution of

V1aR (i.e., VT4R) has been examined in the brain and pituitary gland (Leung et al., 2009; Selvam et

al., 2013) but not in peripheral organs in birds, previous studies reporting the involvement of VT in

renal osmoregulation (Goldstein, 2006; Braun & Dantzler, 1984) suggest the presence of V1aR in the

periphery of birds. Thus, we cannot exclude the possibility that peripheral injection of a V1aR

antagonist affected osmoregulation and secondary social behaviour effects. Although we did not

measure the effects of body temperature and the amount of drinking/evacuation following V1aR

antagonist injection, such homeostatic effects on the periphery from V1aR blocking are unlikely to

occur selectively in affiliated dyads. To determine the involvement of the central V1aR in the social

behaviour of affiliated males, it is necessary to examine the effect of central infusion of V1aR

antagonists in future studies.

Differences in functions and inherent aggression in same-sex affiliative relationships: a

comparison of crows and other species

There are various types of same-sex affiliative relationships that differ in sex, kinship and

function. The underlying physiological mechanisms might also be diverse. The extensive body of

research on meadow voles (Anacker et al., 2016) and zebra finches (Goodson & Wang, 2006; Kelly &

Goodson, 2013; Kelly et al., 2011) has established that V1aR plays a crucial role in promoting same-

sex affiliative relationships or behaviours. In this study on large-billed crows, it was revealed that

V1aR is involved, although the function of same-sex affiliation and the inherent agonistic tendencies

between individuals differ from those in voles and finches.

Meadow voles and prairie voles exhibit a behavioural preference for familiar individuals over

unfamiliar ones, engaging in affiliative interactions and forming communal groups with minimal

aggression (Madison & McShea, 1987; Getz et al., 1993). Both voles share common functions in

maintaining body temperature and predator defence through group living, which supports safe

breeding (Madison & McShea, 1987; Getz et al., 1993). Meadow voles prefer familiar individuals

during winter in the wild and under short day winter-like conditions in the lab, huddling together.

Winter social groups, seeded by a female and her undispersed offspring (mother-daughter or sister-

sister relationships), often continue into spring with communal nesting (McShea & Madison, 1984).

Prairie voles, closely related to meadow voles, have been primarily studied for male-female

interactions, but same-sex interactions have also been investigated. Regardless of day length, both

males and females prefer familiar mates and same-sex peers (DeVries et al., 1997; Beery et al., 2018;

Lee et al., 2019). This preference, without seasonal variation and not biased towards females, differs

from that of meadow voles. However, prairie vole units are fundamentally composed of bonded

breeding pairs and undispersed offspring (Getz et al., 1993; Carter & Getz, 1993). These findings

suggest that same-sex affiliative relationships in both species of voles provide not only

thermoregulatory benefits, as commonly demonstrated in rodents (Andrews et al., 1987; Andrews &

Belknap, 1993; Gilbert et al., 2010; Kauffman et al., 2003), but also ensure the safe rearing of

offspring.

Gregarious zebra finches are highly social and tolerant, roosting and breeding together in colonies

for anti-predator defense (Zann, 1996; Schuett & Dall, 2010; Evans et al., 2018), with low-intensity

aggression such as threats and chases (Ruploh et al., 2013). Gregariousness is a term used to describe

species that affiliate with or live in groups (Miller, 1922; Treisman, 1975; Goodson & Kingsbury,

2011; Peterson & Weckerly, 2018). Recent studies of neuropeptide-mediated behaviours have also

focused on spiny mice (*Acomys cahirinus*), group-living communal breeders, which also exhibit a

gregarious phenotype (Kelly & Seifert, 2021; Powell et al., 2023; Fricker & Kelly., 2024). In spiny

mice, both males and females show little aggression in various social contexts and prefer affiliating

with larger groups (Fricker et al., 2023; Gonzalez Abreu et al., 2022).

Research on voles, finches, and spiny mice has shed light on the physiological basis of same-sex

affiliative relationships, including the role of V1aR. However, since these species exhibit low levels

of inter-individual aggression, the exploration of the physiological basis of affiliative relationships

formed between dominant and subordinate individuals has not been investigated. As previously

mentioned, the exact function of the affiliative relationships formed between dominant and

subordinate males in large-billed crows has not yet been clarified. However, in other *Corvus* species,

these relationships are known to function as coalitions and provide social buffering effects (Fraser &

Bugnyar, 2012; Braun & Bugnyar, 2012; Fraser & Bugnyar, 2010b; Fraser & Bygnyar, 2011; Seed et

al., 2007). Therefore, it is highly likely that similar functions exist in our study subjects as well. The

involvement of V1aR demonstrated in this study of male-male affiliative relationships suggests a

universal role for V1aR in same-sex affiliative relationships despite the likely fundamental differences

in function and the levels of inherent aggression between individuals compared to voles and finches.

One of our most intriguing findings was that the V1aR antagonist disrupted the affiliative

relationship, causing the resurgence of aggressive and submissive behaviours depending on the

relative dominance only within affiliated dyads but had no effect within unaffiliated dyads. Given this

result, the role of V1aR in male-male affiliation among crows might be to inhibit the inherent

agonistic tendencies and enable affiliative interactions, such as allopreening and close proximity. This

function of V1aR might have been evolutionarily acquired in crows and, possibly in primates and

dolphins as well. Similar context-specific involvement of V1aR in the control of aggressive behaviour

in birds has been suggested in previous studies. For instance, a previous study on violet-eared

waxbills (*Uraeginthus granatina*) found that peripheral injection of a V1aR antagonist disinhibited

and increased the aggression of subordinate males in a territorial challenge context, whereas V1aR

reduced the aggression of dominant males towards rival males in a mate competition context

(Goodson et al., 2009). However, in this study, male-male interactions were highly antagonistic and

did not form affiliative relationships, making it impossible to verify the connection between affiliation

and dominance relationships. To clarify the role of V1aR and the relevant VP/VT signals in

facilitating affiliative behaviour and/or inhibiting agonistic behaviour in affiliated males, future

studies should investigate the central infusion of a V1aR antagonist and assess V1aR-/VT-neuron

activities using crows. Additionally, recent studies on localisation of V1aR and OTR in chimpanzees

and other primates (Rogers Flattery et al., 2022) suggest promising developments in this field.

In conclusion, we demonstrated that a triadically interactive environment plays a crucial role in

forming male-male affiliative relationships in non-human animals. Our study suggests that dominant-

subordinate affiliative relationships in males share common neural underpinnings with other same-sex

bonds, pair bonds and parent-infant bonds involving the VP/VT systems. This finding provides

insights into social affiliation, indicating that VP/VT may facilitate these relationships by promoting

affiliative behaviours and suppressing aggression. Our results underscore the significance of triadic

interactions in social behaviour, providing a new perspective on how complex social relationships are

formed and sustained. Future research should explore the roles of VP/VT as well as OT/MT and their

receptor subtypes in social affiliation among same-sex individuals to enhance our understanding of

the evolution of inter-individual affiliation in the social lives of various species, including humans.

Acknowledgements

We thank the members of the Animal Psychology Lab, Keio University, for their inspiring scientific

discussions and their support in the daily care of the animals. We also thank Dr Shigeru Watanabe

(Keio University, Japan) for his helpful advice and discussions. This work was financially supported

by JSPS KAKENHI (#23H01058, #23H00074, #23K17645, #20H01787 to E.-I., and #19J22654 to

A.S.) and a Keio University Grant-in-Aid for ICRP (#MKJ1905) to E.-I.

Author contributions

A.S. designed research, conducted the experiments, analysed data, wrote the manuscript, and acquired

funding. E.-I. reviewed and edited the manuscript, supervised the work, and acquired funding.

Declaration of interests

The authors declare no competing interests.

Source data and videos

Source data and videos related to Experiments 1 and 2 can be found online at

https://figshare.com/projects/Seguchi_Izawa_Role_of_Vasopressin1a_on_male-

male_affiliations_in_large-billed_crows/213883

Figure 2—source data 1

: Source data of social behaviours in dyadic encounters before Experiment 1 (Figure 2).

Figure 3—source data 1

566 : Source data of allopreening behaviours in Experiment 1 (Figure 3A).

567 Figure 3—source data 2

568 : Source data of allopreening behaviours used in discriminant analysis (Figure 3B).

569 Figure 4—source data 1

570 : Source data of social behaviours in Experiment 2 (Figure 4).

571 Video 1

572 : Triadic cohabitation.

573 Video 2

574 : Effects of V1aR antagonism.

586 **Materials and Methods**

587 **Key resources table**

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Chemicals		
SR49059	Sigma-Aldrich	CAS# 150375-75-0
Dimethyl sulfoxide	Fujifilm	CAS# 67-68-5
Deposited data		
Data for figures and tables	This paper	https://figshare.com/projects/Seguchi_Izawa_Role_of_Vasopressin1a_on_male-male_affiliations_in_large-billed_crows/213883
Software		
SPSS	IBM Corp. (2021). IBM SPSS Statistics for Windows, Version 28.0. IBM Corp, Armonk, NY, USA. URL https://www.ibm.com/products/spss-statistics.	NA
R, RStudio	RStudio Team (2020). RStudio: Integrated Development for R. RStudio, PBC, Boston, MA, USA. URL http://www.rstudio.com/.	NA

Experimental Model and Subject Details

We used 18 adult male crows aged between 4 and 7 years, with body weights ranging from 640 to 790

g. Sex was determined using DNA from blood samples (Fridolfsson & Ellegren, 1999). The birds

were caught as yearlings in Tokyo and neighbouring areas during 2017–2019, with permission from

the Ministry of the Environment Government of Japan (authorisation nos. 29030001, 30081482,

30030019, and 30224001). After capture, the crows were housed in one of three different groups with

5–10 other crows (Table 8) in outdoor aviaries (W 5 m × D 10 m × H 3 m). Three to six months

before the start of this study, the crows were transported to individual steel-mesh cages (W 57 cm × D

93 cm × H 63 cm) in an animal housing room to prevent the formation of dominance or affiliative

relationships through physical interactions. Individual cages were placed side by side on shelves in

the room, allowing the birds to see and hear each other but preventing physical contact to reduce

physical and social stress due to individual housing. Room temperature was maintained at 21 ± 2 °C,

with a light/dark cycle of 13 hours light and 11 hours dark, starting at 0800. Food (dried dog food and

other supplements) and water were freely available in the cages.

The experiments were conducted from October 2019 to September 2022. Due to the accidental death

of some birds and the restricted use of the experimental facility during the COVID-19 pandemic,

some of the planned experiments were not performed, resulting in unbalanced data. The experimental

and housing protocols adhered to the Japanese National Regulations for Animal Welfare and were

approved by the Animal Care and Use Committee of Keio University (#20036, #A2022-055).

Method Details

Determination of relative dominance in dyads

We assigned 18 birds to 8 triads for the cohabitation experiment (Table 1). Before assigning the triad

groups, we determined the relative dominance between each pair of birds through three trials of

dyadic social encounters in an indoor aviary (W 2 m × D 2 m × H 2 m), using a procedure similar to

that reported in previous studies (Nishizawa et al., 2011; Takeda et al., 2022). In each trial, two birds

were given a 5-min opportunity to interact freely. The winner and loser were determined according to

the behavioural criteria used in previous studies (Nishizawa et al., 2011; Takeda et al., 2022).

Specifically, the loser was identified as the bird exhibiting submissive behaviour (i.e., submissive

begging vocalisation and/or avoidance) in response to aggressive behaviour (i.e., jab, peck, aggressive

vocalisation and approach). Within the dyad, the bird who won all three trials was defined as

dominant, while the other as subordinate. All 26 dyads showed an apparent asymmetry in win/loss

numbers across the three trials, indicating the formation of dominance relationships (Figure 2A, B,

Figure 2—source data 1). Thus, 26 dyads with clear dominance relationships were used in Experiment

1. The social encounter trials were conducted between 10:00 and 15:00, with a 2-day interval between

consecutive trials involving the same dyads. No subjects showed allopreening in any trial, indicating

that no affiliative relationship was formed in any dyad at this stage (Figure 2C, Figure 2—source data

1).

Experiment 1: The triadic cohabitation facilitated the formation of an affiliative relationship

between two particular males

After determining the relative dominance positions of the individuals in the dyads of each triad, we

tested whether a 2-week cohabitation in triads facilitated the formation of affiliative relationships

between two specific birds in the triad (Figure 1A). The formation of an affiliative relationship was

evaluated based on the emergence and increase in reciprocal allopreening exclusively between two

specific birds in each triad. To confirm that affiliative relationships were not formed through dyadic

interactions, all possible dyads of each triad (i.e., three dyads) were housed together in the

experimental aviary on separate occasions for three days before triadic cohabitation (hereafter referred

to as pre-test). This pre-test phase was followed by the 2-week triadic cohabitation to examine

whether triadic interactions could facilitate the formation of affiliative relationships. To ensure that the

relationships observed during the triadic cohabitation were not simply temporary or artefacts of

spatial arrangements, a second dyadic cohabitation period was conducted following the triadic

cohabitation (hereafter referred to as post-test). In the pre- and post-tests, each of the two birds was

transferred from its individual home cage into the aviary at 8:00 am, and after 72 h, it was returned to

the home cage at 8:00 am. On the day following the end of the pre-test period, the three birds were

introduced into the experimental aviary and housed together for two weeks. At the end of the triadic

cohabitation, the birds were transferred back to their individual home cages. At 8:00 am the following

day, each pair of the triad was again transferred into the aviary for 3 days as a post-test on separate

occasions.

The behaviour of the birds during the pre-/post-tests and triadic cohabitation was video recorded

using a ceiling-mounted camera (EX-ZR4100, CASIO, Japan) in the aviary. Recordings were

conducted for 30 min randomly between 9:00 and 17:00 every day in the pre- and post-tests and for

the early (1−3 days), middle (7−9 days), and last (12−14 days) three days of triadic cohabitation.

Therefore, 90-min video data (30 min × 3 days) were used to analyse social interactions in each test,

as well as in the 3 × 3-day blocks during the triadic cohabitation period. In the aviary, three wooden

perches were placed in parallel with 30 cm intervals at a height of 80 cm above the floor. Food, water,

and a bathing tub were consistently provided, except during the 30-min video recording sessions.

Experiment 2: V1a receptor antagonism disrupted the dominant-subordinate affiliative relationship

To investigate the involvement of V1a-like receptors in the maintenance of male-male affiliative relationships, birds forming affiliated relationships in Experiment 1 were systematically administered a V1aR antagonist (SR 49059, Sigma-Aldrich, MO, USA) to examine the effects on social behaviour towards the affiliated counterpart and the unaffiliated male (Figure 1B and Table 1). This V1aR is an effective antagonist of the VT4 receptor in birds (Kuenel et al., 2016). The effect of the V1aR antagonist was tested with high dose (0.3 mL × 1 mg/mL vehicle solution) and low dose (0.3 mL × 0.1 mg/mL vehicle solution), compared with a 0.3 mL vehicle solution as a control, administered via intramuscular injection into the chest of subject crows. The vehicle solution contained 10% dimethyl sulfoxide (Fujifilm, Tokyo, Japan) in a saline solution.

In each trial, the V1aR antagonist (high or low dose) or vehicle was administered to either the dominant or subordinate bird. Each bird was tested in eight trials: three with the high-dose condition, three with the low-dose condition, and two with the vehicle control condition. The trial order of the dominance position of the focal individual (dominant or subordinate) and the drug conditions were counterbalanced within and between dyads. After testing the affiliated dyads, the birds underwent three additional trials of dyadic interactions without drug injection to ensure the maintenance or recovery of the relationship. At least one week after this recovery trial, both birds of affiliated dyads received another set of eight trials with the unaffiliated individual (the third bird in the triadic cohabitation in Experiment 1) as an unaffiliated dyad test with the same procedure as that used in the affiliated dyad test.

For each trial, either the dominant or the subordinate bird received an injection 1-h before the start of the trial (referred to as the focal bird). During the 1-h period, the focal bird was isolated from other birds in an individual cage. The trial began soon after the focal bird was introduced into the indoor aviary (W 2 m × D 2 m × H 2 m), where the other bird had been introduced 1 min earlier. The trial procedure was identical to that of Experiment 1; the birds were allowed to interact freely for 30 min. Based on the typical social behaviour in dyads and interactions between dominant and subordinate large-billed crows (Nishizawa et al., 2011; Takeda et al., 2022), we coded allopreening; aggressive behaviour such as peck, jab, and kick; submissive begging vocalisations from video-recorded data for offline analysis. The inter-trial interval was three and five days, with each bird participating in one trial per day. All trials were conducted between 8:30 and 12:00.

Statistical Analysis

Experiment 1: The triadic cohabitation facilitated the formation of an affiliative relationship

between two particular males

We defined an affiliative relationship as a dyad where reciprocal, but not unidirectional, allopreening occurred between two birds, independent of their dominance relationship. This definition aligns the behavioural characteristics of affiliative relationships in primates and other social animals, such as maintaining close proximity and exchanging affiliative behaviours (Fedurek & Dunbar, 2009). An allopreening bout was counted when a bird passed its bill through the feathers of another bird for longer than 2 s (Miyazawa et al., 2020). Given the absence of allopreening in any dyad before triadic cohabitation, we defined a dyad as having an affiliative relationship if both individuals initiated allopreening in 10 or more bouts per 30 min in any of the 3-day blocks during the triadic cohabitation (Figure 3A, Figure 3—source data 1). Dyads not meeting this behavioural criterion were considered as not having formed an affiliative relationship. To confirm the exclusive formation of affiliative relationships through triadic cohabitation, we performed a discriminant analysis to determine whether the numbers of dominant-initiated and subordinate-initiated allopreening in the pre- and post-triadic cohabitation tests could distinguish affiliated dyads from their pre-relationship state and unaffiliated dyads (Figure 3B, Figure 3—source data 2). Leave-one-out cross-validation was used to calculate the correct discrimination rate. The discriminant analysis was performed using SPSS Statistics version 29. Allopreening behaviours were coded using BORIS v.7 (Frinard & Gamba, 2016).

Experiment 2: V1a receptor antagonism disrupted the dominant-subordinate affiliative relationship

To compare the number of social behaviours in the test trials among the drug conditions and between dominants and subordinates, we applied generalised linear mixed models (GLMMs) with Poisson error distribution and the log link function. We analysed allopreening behaviours (Figure 4A, Figure 4—source data 1 and Table 3), aggressive behaviours (Figure 4B, Figure 4—source data 1Tables 4 and 5), and submissive vocalisations (Figure 4C, Figure 4—source data 1Tables 6 and 7) separately. The model included the number of behaviours per 30 min as the response variable, with *dominance* (dominant or subordinate), *drug condition* (V1aR antagonist (V1aRA) high-dose, V1aRA low-dose or vehicle), and their interaction as explanatory variables. *Trial number* and *identity of focal individual* were included as random factors. If the interaction was significant, we stratified the dataset between the dominant and subordinate GLMMs to examine *drug* effects. The significance of variables in each

model was evaluated based on the Wald χ^2 statistics. All statistical analyses were performed using R

version 4.2.2. with the package 'lme4's (Bates et al., 2015) for fitting GLMMs to the data. The

significance level was set at 5%.

Table 1. Individuals used in Experiments 1 and 2.

Birds OW, YA, VV, BB, #28, and KR are included in two different triads. Triad #6 (RM, YA, VA) did

not receive Experiment 2 (see text for the details). *n.a.* indicates data is not available.

Triads used in Experiment 1				Dyads used in Experiment 2							
Triad No.	Dominance rank			Affiliated	Drug conditions received			Unaffiliated	Drug conditions received		
	High	Middle	Low		Vehicle	V1aRA low-dose	V1aRA high-dose		Vehicle	V1aRA low-dose	V1aRA high-dose
1	OW	PY	VV	PY, VV	✓	✓	✓	OW, VV	✓	✓	✓
2	RR	OW	BB	OW, BB	✓	✓	✓	RR, BB	✓	✓	✓
3	#28	YA	#27	#28, YA	✓	*n.a.*	✓	#28, #27	✓	*n.a.*	✓
4	VV	BB	#28	VV, BB	✓	✓	✓	BB, #28	✓	*n.a.*	✓
5	KR	KY	OBO	KY, OBO	✓	*n.a.*	✓	KR, KY	✓	*n.a.*	✓
6	RM	YA	VA	(RM, YA)	–	–	–	–	–	–	–
7	GK	KG	RM	–	–	–	–	–	–	–	–
8	KV	KR	LK	–	–	–	–	–	–	–	–

727 **Table 2. Outputs of discriminant analysis.**

| | | | | Predicted classification | | | | |
| | | | | Affiliated dyad | | Unaffiliated dyad | | |
				pre-test	post-test	pre-test	post-test	Total
Original	Counts	Affiliated dyad	pre-test	2	0	4	0	6
			post-test	0	6	0	0	6
		Unaffiliated dyad	pre-test	2	0	15	0	17
			post-test	3	0	14	0	17
	%	Affiliated dyad	pre-test	33.3	0	66.7	0	100
			post-test	0	100	0	0	100
		Unaffiliated dyad	pre-test	11.8	0	88.2	0	100
			post-test	17.6	0	82.4	0	100
Cross-validated	Counts	Affiliated dyad	pre-test	0	0	6	0	6
			post-test	1	5	0	0	6
		Unaffiliated dyad	pre-test	2	0	0	15	17
			post-test	3	0	14	0	17
	%	Affiliated dyad	pre-test	0	0	100	0	100
			post-test	16.7	83.3	0	0	100
		Unaffiliated dyad	pre-test	11.8	0	0	88.2	100
			post-test	17.6	0	82.4	0	100

730 **Table 3. Outputs of GLMMs for allopreening behaviours in affiliated dyads.**

variables	estimates	s.e.	z -value	p
intercept	0.854	0.275	3.108	*< 0.001*
Dominance[a]	0.325	0.257	1.266	0.206
Drug condition[b]				
Vehicle	1.639	0.217	7.554	*< 0.001*
V1aRA high-dose	-0.724	0.313	-2.309	*0.021*
Dominance[a]×Vehicle	0.068	0.281	0.241	0.811

731 Italicized indicates statistically significant results.

732 [a] Subordinate was set to 0.

733 [b] V1aRA low-dose was set to 0.

735   **Table 4. Outputs of GLMMs for aggressive behaviours in affiliated dyads.**

variables	estimates	s.e.	z-value	p
intercept	0.094	0.293	0.319	0.749
Drug condition[a]				
Vehicle	-0.193	0.315	-0.612	0.540
V1aRA high-dose	0.791	0.251	3.147	*< 0.01*

736   Italicized indicates statistically significant results.

737   [a] V1aRA low-dose was set to 0.

739 **Table 5. Outputs of GLMMs for aggressive behaviours in unaffiliated dyads.**

variables	estimates	s.e.	z-value	p
intercept	0.080	0.277	0.289	0.773
Drug condition[a]				
Vehicle	-0.031	0.353	-0.089	0.929
V1aRA high-dose	0.045	0.326	0.138	0.890

740 Italicized indicates statistically significant results.

741 [a] V1aRA low-dose was set to 0.

743 **Table 6. Outputs of GLMMs for submissive vocalizations in affiliated dyads.**

variables	estimates	s.e.	z-value	p
intercept	-0.083	0.273	-0.303	0.762
Drug condition[a]				
Vehicle	0.237	0.318	0.747	0.455
V1aRA high-dose	1.929	0.253	7.617	*<0.001*

744   Italicized indicates statistically significant results.

745   [a] V1aRA low-dose was set to 0.

747 **Table 7. Outputs of GLMMs for submissive vocalizations in unaffiliated dyads.**

variables	estimates	s.e.	z-value	p
intercept	0.000	0.289	0.000	1.000
Drug condition[a]				
Vehicle	0.095	0.359	0.266	0.791
V1aRA high-dose	0.095	0.337	0.283	0.777

748   Italicized indicates statistically significant results.

749   [a] V1aRA low-dose was set to 0.

751 **Table 8. Composition of the subjects in the group housing prior to this study and their**

752 **ages.**

ID	Group	Age
VV	Group A	6
RR		6
OW		6
PY		6
BB		6
#28	Group B	6
#27		6
KR	Group C	6
KY		6
KV		6
KG		6
LK	Group D	4
GK		4
YA	Group E	7
OBO	Group F	4
RM	Group G	7
VA	Group H	6

753 Letters indicate individual birds.

References

Anacker A. M., Christensen J. D., LaFlamme E. M., Grunberg D. M., Beery A. K (2016) **Septal oxytocin administration impairs peer affiliation via V1a receptors in female meadow voles** *Psychoneuroendocrinology* **68**: 156–162. https://doi.org/10.1016/j.psyneuen.2016.02.025.

Andrews R. V., and Belknap R. W (1993) **Season affects tolerance of cohabitation by deer mice** *Physiology and Behavior* **53**: 617–620. https://doi.org/10.1016/0031-9384(93)90163-A.

Andrews R. V., Phillips D., Makihara, D (1987) **Metabolic and thermoregulatory consequences of social behaviors between *Microtus townsendii* Comparative biochemistry and physiology** **87**: 345–348. https://doi.org/10.1016/0300-9629(87)90133-2.

Aota I., Yatsuda C., Izawa E.-I (2023) **Salivary corticosterone measurement in large-billed crows by enzyme-linked immunosorbent assay** *Journal of veterinary medical science* **85**: 71–75. https://doi.org/10.1292/jvms.22-0383.

Aureli F., Preston S. D., de Waal, F. B. M (1999) **Heart rate responses to social interactions in free-moving rhesus macaques (*Macaca mulatta*): a pilot study** *Journal of comparative psychology* **113**: 59–65. https://doi.org/10.1037/0735-7036.113.1.59.

Bales K. L., Ardekani C. S., Baxter A., Karaskiewicz C. L., Kuske J. X., Lau A. R., Savidge L. E., Sayler K. R., Witczak L. R (2021) **What is a pair bond?** *Hormones and behavior* **136**: 105062. https://doi.org/10.1016/j.yhbeh.2021.105062.

Bates D., Mächler M., Bolker B., Walker S (2015) **Fitting linear mixed-effects models using lme4** *Journal of statistical software* **67**: 1–48. https://doi.org/10.18637/jss.v067.i01.

Beauchamp G (2010) **Relaxed predation risk reduces but does not eliminate sociality in birds** *Biology letters* **6**: 472–474. https://doi.org/10.1098/rsbl.2009.1063.

Beery A. K (2019) **Frank Beach award winner: Neuroendocrinology of group living** *Hormones and behavior* **107**: 67–75. https://doi.org/10.1016/j.yhbeh.2018.11.002.

Beery A. K., Christensen J. D., Lee N. S., Blandino K. L (2018) **Specificity in sociality: mice and prairie voles exhibit different patterns of peer affiliation** *Frontiers in behavioral neuroscience* **12**: 50. https://doi.org/10.3389/fnbeh.2018.00050.

Beery A. K., Loo T. J., Zucker I (2008) **Day length and estradiol affect same-sex affiliative**

behavior in the female meadow vole *Hormones and behavior* 54: 153–159.

https://doi.org/10.1016/j.yhbeh.2008.02.007.

Bercovitch, F. B (1988) **Coalitions, cooperation and reproductive tactics among adult male**

baboons *Animal Behaviour* 36: 1198–1209. https://doi.org/10.1016/S0003-3472(88)80079-4.

Black J. M (1996) **Partnerships in birds: the study of monogamy**. (Oxford: Oxford University

Press). https://doi.org/10.1093/oso/9780198548614.001.0001.

Boeckle M., Bugnyar T (2012) **Long-Term Memory for Affiliates in Ravens** *Current Biology* **22**:

801–806. https://doi.org/10.1016/j.cub.2012.03.023.

Bosch O. J., Neumann I. D (2008) **Brain vasopressin is an important regulator of maternal**

behavior independent of dams' trait anxiety *Proceedings of the National Academy of Sciences of*

the United States of America **105**: 17139–17144. https://doi.org/10.1073/pnas.0807412105.

Boucherie P. H., Loretto M.-C., Massen J. J. M., Bugnyar T (2019) **What constitutes "social**

complexity" and "social intelligence" in birds? Lessons from ravens *Behavioral ecology and*

sociobiology **73**: 12. https://doi.org/10.1007/s10336-023-02111-6.

Braun A., Bugnyar T (2012) **Social bonds and rank acquisition in raven non-breeder aggregations**

Animal Behaviour **84**: 1507–1515. https://doi.org/10.1016/j.anbehav.2012.09.024.

Braun A., Walsdorff T., Fraser O. N., Bugnyar T (2012) **Socialized sub-groups in a temporary**

stable raven flock? *Journal of ornithology* **153**: 97–104. https://doi.org/10.1007/s10336-011-0810-2.

Braun E. J., Dantzler W. H (1984) **Endocrine regulation of avian renal function** *The Journal of*

experimental zoology **232**: 715–723. https://doi.org/10.1002/jez.1402320341.

Brent L. J., Chang S. W., Gariépy J. F., Platt M. L (2014) **The neuroethology of friendship** *Annals of*

the New York Academy of Sciences **1316**: 1–17. https://doi.org/10.1111/nyas.12315.

Brent L. J. N., Semple S., Dubuc C., et al. (2011) **Social capital and physiological stress levels in**

free-ranging adult female rhesus macaques *Physiology and Behavior* **102**: 76–83.

https://doi.org/10.1016/j.physbeh.2010.09.022.

Broad K. D., Curley J. P., Keverne E. B (2006) **Mother-infant bonding and the evolution of**

mammalian social relationships *Philosophical transactions of the Royal Society of London. Series*

B, Biological sciences **361**: 2199–2214. https://doi.org/10.1098/rstb.2006.1940.

Bygott J. D., Bertram B. C. R., Hanby J. P (1979) **Male lions in large coalitions gain reproductive advantages** *Nature* **282**: 839–841. https://doi.org/10.1038/282839a0.

Caplow T (1956) **A theory of coalitions in the triad** *American Sociological Review* **21**: 489–493. https://doi.org/10.2307/2088718.

Caro T. M (1993) **Behavioral solutions to breeding cheetahs in captivity: insights from the wild** *Zoo Biology* **12**: 19–30. https://doi.org/10.1002/zoo.1430120105.

Caro T. M (1994) **Cheetahs of the Serengeti plains: group living in an asocial species** (Chicago: University of Chicago Press).

Carter C. S., Getz L. L (1993) **Monogamy and the prairie vole: battle of the sexes** *Scientific American* **268**: 100–106. https://doi.org/10.1038/scientificamerican0693-100.

Clutton–Brock T. H (1991) **The evolution of parental care** (Princeton: Princeton University Press).

Connor R. C., Smolker R. A., Richards A. F (1992a) **Dolphin alliances and coalitions** In Coalitions and alliances in humans and other animals, A.H. Harcourt, and F.B.M. de Waal, eds. (Oxford: Oxford University Press). 415–443. https://doi.org/10.1098/rstb.2006.1997.

Connor R. C., Smolker R. A., Richards A. F (1992b) **Two levels of alliance formation among male bottlenose dolphins (*Tursiops sp.*)** *Proceedings of the National Academy of Sciences of the United States of America* **89**: 987–990. https://doi.org/10.1073/pnas.89.3.987.

Connor R. C., Heithaus M. R., Barre L. M (1999) **Superalliance of bottlenose dolphins** *Nature* **397**: 571–572. https://doi.org/10.1098/rspb.2000.1357.

Crespi B. J (2016) **Oxytocin, testosterone, and human social cognition** *Biological reviews of the Cambridge Philosophical Society* **91**: 390–408. https://doi.org/10.1111/brv.12175.

Crockford C., Wittig R. M., Whitten P. L., et al (2008) **Social stressors and coping mechanisms in wild female baboons (*Papio hamadryas ursinus*)** *Hormones and Behavior* **53**: 254–265. https://doi.org/10.1016/j.yhbeh.2007.10.007.

Dall S. R. X., Wright J (2009) **Rich pickings near large communal roosts favor "gang" foraging**

by juvenile common ravens, *Corvus corax* *PLoS ONE* 4: e4530.
https://doi.org/10.1371/journal.pone.0004530.

de Kort S. R., Emery N. J., Clayton N. S (2006) **Food sharing in jackdaws,** *Corvus monedula***:**
What, why and with whom? *Animal Behaviour* **72**: 297–304.
https://doi.org/10.1016/j.anbehav.2005.10.016.

deVries A. C., Glasper E. R., Detillion C. E (2003) **Social modulation of stress responses** *Physiology*
and Behavior **79**: 399–407. https://doi.org/10.1016/s0031-9384(03)00152-5.

deVries A. C., Johnson C. L., Carter C.S (1997) **Familiarity and gender influence social**
preferences in prairie voles (*Microtus ochrogaster***)** *Canadian journal of zoology* **75**: 295–301.
https://doi.org/10.1139/z97-037

de Waal F. B., Luttrell L. M (1988) **Mechanisms of social reciprocity in three primate species:**
Symmetrical relationship characteristics or cognition? *Ethology and sociobiology* **9**: 101–118.
https://doi.org/10.1016/0162-3095(88)90016-7.

Emery N. J., Seed A. M., von Bayern A.M.P., Clayton N. S (2007a) **Cognitive adaptations of social**
bonding in birds *Philosophical transactions of the Royal Society of London. Series B: Biological*
sciences **362**: 489–505. https://doi.org/10.1098/rstb.2006.1991.

Emery N. J., Seed A. M., von Bayern A. M. P., Clayton N. S (2007b) **Cognitive adaptations of social**
bonding in birds *Philosophical transactions of the Royal Society of London. Series B: Biological*
sciences **362**: 489–505. https://doi.org/10.1098/rstb.2006.1991.

Evans J. C., Dall S. R. X., Kight C. R (2018) **Effects of ambient noise on zebra finch vigilance and**
foraging efficiency *PLoS ONE* **13**: e0209471. https://doi.org/10.1371/journal.pone.0209471.

Fedurek P., Dunbar R. I. M (2009) **What does mutual grooming tell us about why chimpanzees**
groom? *Ethology* **115**: 566–575. https://doi.org/10.1111/j.1439-0310.2009.01637.x.

Fraser O. N., Bugnyar T (2010a) **The quality of social relationships in ravens** *Animal Behaviour* **79**:
927–933. https://doi.org/10.1016/j.anbehav.2010.01.008.

Fraser O. N., Bugnyar T (2010b) **Do Ravens Show Consolation? Responses to Distressed Others**
PLoS ONE **5**: e10605. https://doi.org/10.1371/journal.pone.0010605.

Fraser O. N., Bugnyar T (2011) **Ravens reconcile after aggressive conflicts with valuable partners**
PLoS ONE **6**: e18118. https://doi.org/10.1371/journal.pone.0018118.

Fraser O. N., Bugnyar T (2012) **Reciprocity of agonistic support in ravens** *Animal Behaviour* **83**:
171–177. https://doi.org/10.1016/j.anbehav.2011.10.023.

Fricker B. A., Ho D., Seifert A. W., Kelly A. M (2023) **Biased brain and behavioral responses**
towards kin in males of a communally breeding species *Scientific Report* **13**: 17040.
https://doi.org/10.1038/s41598-023-44257-6.

Fricker B. A., Kelly A. M (2024) **From grouping and cooperation to menstruation: Spiny mice**
(*Acomys cahirinus*) are an emerging mammalian model for sociality and beyond *Hormones and*
behavior **158**: 105462. https://doi.org/10.1016/j.yhbeh.2023.105462.

Fridolfsson A. -K., Ellegren H (1999) **A simple and universal method for molecular sexing of non-**
ratite birds *Journal of avian biology* **30**: 116–121. https://doi.org/10.2307/3677252.

Friard O., Gamba M (2016) **BORIS: a free, versatile open-source event-logging software for**
video/audio coding and live observations *Methods in ecology and evolution* **7**: 1325–1330.
https://doi.org/10.1111/2041-210X.12584.

Getz L. L., McGuire B., Pizzuto T., Hofmann J. E., Frase B (1993) **Social organization of the prairie**
vole (*Microtus ochrogaster*) *Journal of mammalogy* **74**: 44–58. https://doi.org/10.2307/1381904.

Gilbert C., McCafferty D., Le Maho Y., Martrette J. -M., Giroud S., Blanc S., and Ancel A (2010)
One for all and all for one: the energetic benefits of huddling in endotherms *Biological reviews of*
the Cambridge philosophical society **85**: 545–569. https://doi.org/10.1111/j.1469-185X.2009.00115.x.

Gilby I. C., Brent L. J., Wroblewski E. E., Rudicell R. S., Hahn B. H., Goodall J., Pusey A. E (2013)
Fitness benefits of coalitionary aggression in male chimpanzees *Behavioral Ecology and*
Sociobiology **67**: 373–381. https://doi.org/10.1007/s00265-012-1457-6.

Goldstein, D.L. (2006). Regulation of the avian kidney by arginine vasotocin. Gen. Comp.
Endocrinol. 147, 78–84. https://doi.org/10.1016/j.ygcen.2005.09.018.

Gonzalez Abreu J. A., Rosenberg A. E., Fricker B. A., Wallace K. J., Seifert A. W., Kelly A. M (2022)
Species-typical group size differentially influences social reward neural circuitry during
nonreproductive social interactions *iScience* 104230. https://doi.org/10.1016/j.isci.2022.104230.

Goodson J. L., Kabelik D., Schrock S. E (2009) **Dynamic neuromodulation of aggression by vasotocin: influence of social context and social phenotype in territorial songbirds** *Biology letters* **5**: 554–556. https://doi.org/10.1098/rsbl.2009.0316.

Goodson J. L., Kingsbury M. A (2011) **Nonapeptides and the evolution of social group sizes in birds** *Frontiers in Neuroanatomy* **5**: 13. https://doi.org/10.3389/fnana.2011.00013.

Goodson J. L., Wang Y (2006) **Valence-sensitive neurons exhibit divergent functional profiles in gregarious and asocial species** *Proceedings of the national academy of sciences* **103**: 17013–17017. https://doi.org/10.1073/pnas.0606278103.

Green M. L., Herzing D. L., Baldwin J. D (2015) **Molecular assessment of mating strategies in a population of Atlantic spotted dolphins** *PLoS ONE* **10**: e0118227. https://doi.org/10.1371/journal.pone.0118227.

Gubernick D. J., Alberts J. R (1983) **Maternal licking of young: resource exchange and proximate controls** *Physiology and behavior* **31**: 593–601. https://doi.org/10.1016/S0031-9384(24)00190-2

Harcourt A., de Waal F. B. M (1992) **Coalitions and Alliances in Humans and other Animals** (Oxford: Oxford University Press).

Hassinen E., Pyörnilä A., Hissa R (1999) **Cardiovascular and thermoregulatory responses to vasotocin and angiotensin II in the pigeon** *Part A, Molecular & integrative physiology* **123**: 279–285. https://doi.org/10.1016/s1095-6433(99)00061-6.

Heinrich B., Marzluff J. M (1991) **Do common ravens yell because they want to attract others?** *Behavioral ecology and sociobiology* **28**: 13–21. https://doi.org/10.1007/BF00172134

Hendricks P., Schlang S (1998) **Aerial attacks by common ravens, *Corvus corax*, on adult feral pigeons, *Columba livia*** *The Canadian field-naturalist* **112**: 702–707. https://doi.org/10.5962/p.358502.

Hinde R. A (1976) **Interactions, relationships and social structure** *Man* **11**: 11–17. https://doi.org/10.2307/2800384.

Islam M. N., Zhu Z. B., Aoyama M., Sugita S (2010) **Histological and morphometric analyses of seasonal testicular variations in the Jungle crow (*Corvus macrorhynchos*)** *Anatomical science*

international **85**: 121–129. https://doi.org/10.1007/s12565-009-0066-6.

Izawa E. -I (2011) **Social ecology of corvids** *Japanese journal of animal psychology* **61**: 55–68.
https://doi.org/10.2502/janip.61.1.5.

Izawa E. -I., Watanabe S (2008) **Formation of linear dominance relationship in captive jungle**
crows (*Corvus macrorhynchos*): implications for individual recognition *Behavioral processes* **78**:
44–52. https://doi.org/10.1016/j.beproc.2007.12.010.

Jans J. E., Woodside B. C (1990) **Nest temperature: effects on maternal behavior, pup**
development, and interactions with handling *Developmental psychobiology* **23**: 519–534.
https://doi.org/10.1002/dev.420230607.

Kauffman A. S., Paul M. J., Butler M. P., Zucker I (2003) **Huddling, locomotor, and nestbuilding**
behaviors of furred and furless Siberian hamsters *Physiology and behavior* 79: 247–256.
https://doi.org/10.1016/S0031-9384(03)00115-X.

Kelley H. H., Arrowood A. J (1960) **Coalitions in the triad: critique and experiment** *Sociometry*
23: 231–244. https://doi.org/10.2307/2785888.

Kelly A. M., Goodson J. L (2013) **Functional significance of a phylogenetically widespread sexual**
dimorphism in vasotocin/vasopressin production *Hormones and behavior* **64**: 840–846.
https://doi.org/10.1016/j.yhbeh.2013.09.006.

Kelly A. M., Kingsbury M. A., Hoffbuhr K., Schrock S. E., Waxman B., Kabelik D., Thompson, R.
R., Goodson J. L (2011) **Vasotocin neurons and septal V1a-like receptors potently modulate**
songbird flocking and responses to novelty *Hormones and behavior* **60**: 12–21.
https://doi.org/10.1016/j.yhbeh.2011.01.012.

Kelly A. M., Seifert A. W (2021) **Distribution of vasopressin and oxytocin neurons in the basal**
forebrain and midbrain of spiny mice (*Acomys cahirinus*) *Neuroscience* **468**: 16–28.
https://doi.org/10.1016/j.neuroscience.2021.05.034.

Kikusui T., Winslow J. T., Mori Y (2006) **Social buffering: relief from stress and anxiety**
Philosophical transactions of the Royal Society of London. Series B: Biological sciences **361**: 2215–
2228. https://doi.org/10.1098/rstb.2006.1941.

Kleiman D. G (1977) **Monogamy in mammals** *The quarterly review of biology* **52**: 39–69.

https://doi.org/10.1086/409721.

1024

1025 Kobayashi H., Whitehead H., Amano M (2020) **Long-term associations among male sperm whales**
1026 (***Physeter macrocephalus***) *PLoS ONE* **15**: e0244204. https://doi.org/10.1371/journal.pone.0244204.

1027

1028 Koshimizu T. A., Nakamura K., Egashira N., Hiroyama M., Nonoguchi H., Tanoue A (2012)
1029 **Vasopressin V1a and V1b receptors: from molecules to physiological systems** *Physiological*
1030 *reviews* **92**: 1813–1864. https://doi.org/10.1152/physrev.00035.2011.

1031

1032 Koykka C., Wild G (2017) **Concessions, lifetime fitness consequences, and the evolution of**
1033 **coalitionary behavior** *Behavioral ecology* **28**: 20–30. https://doi.org/10.1093/beheco/arw126.

1034

1035 Kuenzel W. J., Hancock M., Nagarajan G., Aman N. A., Kang S. W (2016) **Central effect of**
1036 **vasotocin 4 receptor (VT4R/V1aR) antagonists on the stress response and food intake in chicks**
1037 **given neuropeptide Y (NPY)** *Neuroscience letters* **620**: 57–61.
1038 https://doi.org/10.1016/j.neulet.2016.03.036.

1039

1040 Kuroda N (1990) **The jungle crows in Tokyo** (Abiko, Japan: Yamashina Institute for Ornithology).

1041

1042 Kutsukake N., Hasegawa T (2005) **Dominance turnover between an alpha and a beta male and**
1043 **dynamics of social relationships in Japanese macaques** *International journal of primatology* **26**:
1044 775–800. https://doi.org/10.1007/s10764-005-5308-4.

1045

1046 Laakasuo M., Rotkirch A., van Duijn M., Berg V., Jokela M., David-Barrett T., Miettinen A., Pearce
1047 E., Dunbar R (2020) **Homophily in personality enhances group success among real-life friends**
1048 *Frontiers in psychology* **11**: 710. https://doi.org/10.3389/fpsyg.2020.00710.

1049

1050 Lee N. S., Beery A. K (2021) **The role of dopamine signaling in prairie vole peer relationships**
1051 *Hormones and behavior* **127**: 104876. https://doi.org/10.1016/j.yhbeh.2020.104876.

1052

1053 Lee, N.S., Goodwin, N.L., Freitas, K.E., and Beery, A.K. (2019). Affiliation, aggression, and
1054 selectivity of peer relationships in meadow and prairie voles. Front. Behav. Neurosci. 13.
1055 https://doi.org/10.3389/fnbeh.2019.00052.

1056

1057 Lee N. S., Kim C. Y., Beery A. K (2023) **Peer social environment impacts behavior and dopamine**
1058 **D1 receptor density in Prairie Voles (***Microtus ochrogaster***)** *Neuroscience* **515**: 62–70.
1059 https://doi.org/10.1016/j.neuroscience.2023.02.002.

1060

1061   Leung C. H., Goode C. T., Young L. J., Maney D. L (2009) **Neural distribution of nonapeptide**

1062   **binding sites in two species of songbird** *Journal of comparative neurology* **513**: 197–208.

1063   https://doi.org/10.1002/cne.21947.

1064

1065   Loretto M. C., Schuster R., Itty C., Marchand P., Genero F., Bugnyar T (2017) **Fission-fusion**

1066   **dynamics over large distances in raven non-breeders** *Scientific reports* **7**: 380.

1067   https://doi.org/10.1038/s41598-017-00404-4.

1068

1069   Macfarlan S. J., Walker R. S., Flinn M. V., Chagnon N. A (2014) **Lethal coalitionary aggression and**

1070   **long-term alliance formation among Yanomamö men** *Proceedings of the national academy of*

1071   *sciences* **111**: 16662–16669. https://doi.org/10.1073/pnas.1418639111.

1072

1073   Madison D. M., McShea W. J (1987) **Seasonal changes in reproductive tolerance, spacing, and**

1074   **social organization in meadow voles: A Microtine Model** *American zoologist* **27**: 899–908.

1075   https://doi.org/10.1093/icb/27.3.899.

1076

1077   Marzluff J. M., Heinrich B., Marzluff C. S (1996) **Raven roosts are mobile information centers**

1078   *Animal behaviour* **51**: 89–103. https://doi.org/10.1006/anbe.1996.0008.

1079

1080   Massen J. J. M., Koski S. E (2014) **Chimps of a feather sit together: chimpanzee friendships are**

1081   **based on homophily in personality** *Evolution and human behavior* **35**: 1–8.

1082   https://doi.org/10.1016/j.evolhumbehav.2013.08.008.

1083

1084   Massen J. J. M., Pasukonis A., Schmidt J., Bugnyar T (2014a) **Ravens notice dominance reversals**

1085   **among conspecifics within and outside their social group** *Nature communications* **5**: 3679.

1086   https://doi.org/10.1038/ncomms4679.

1087

1088   Massen J. J. M., Szipl G., Spreafico M., Bugnyar T (2014b) **Ravens intervene in others' bonding**

1089   **attempts** *Current biology* **24**: 2733–2736. https://doi.org/10.1016/j.cub.2014.09.073.

1090

1091   McShea W. J., Madison D. M (1984) **Communal nesting between reproductively active females in**

1092   **a spring population of** *Microtus pennsylvanicus* *Canadian journal of zoology* **62**: 344–346.

1093   https://doi.org/10.1139/z84-053.

1094

1095   Mesterton-Gibbons M., Gavrilets S., Gravner J., Akçay E (2011) **Models of coalition or alliance**

1096   **formation** *Journal of theoretical biology* **274**: 187–204. https://doi.org/10.1016/j.jtbi.2010.12.031.

1097

1098   Miyazawa E., Seguchi A., Takahashi N., Motai A., Izawa E.-I. (2020) **Different patterns of**

allopreening in the same-sex and opposite-sex interactions of juvenile large-billed crows (*Corvus macrorhynchos*) *Ethology* **126**: 195–206. https://doi.org/10.1111/eth.12992.

Miller R. C (1922) **The significance of the gregarious habit** *Ecology* **3**: 122–126. https://doi.org/10.2307/1929145.

Muller M. N., Mitani J. C (2005) **Conflict and cooperation in wild chimpanzees** *Advances in the study of behavior* **35**: 275–331. https://doi.org/10.1016/S0065-3454(05)35007-8.

Nishizawa K., Izawa E. I., Watanabe S (2011) **Neural-activity mapping of memory-based dominance in the crow: neural networks integrating individual discrimination and social behaviour control** *Neuroscience* **197**: 307–319. https://doi.org/10.1016/j.neuroscience.2011.09.001.

Noë N., Whitaker R. M., Chorley M. J., Pollet T. V (2016) **Birds of a feather locate together? Foursquare checkins and personality homophily** *Computers in human behavior* **58**: 343–353. https://doi.org/10.1016/j.chb.2016.01.009.

Noë R (1994) **A model of coalition formation among male baboons with fighting ability as the crucial parameter** *Animal behaviour* **47**: 211–213. https://doi.org/10.1006/anbe.1994.1024.

Numan M., Young L. J (2016) **Neural mechanisms of mother-infant bonding and pair bonding: Similarities, differences, and broader implications** *Hormones and behavior* **77**: 98–112. https://doi.org/10.1016/j.yhbeh.2015.05.015.

Ode M., Asaba A., Miyazawa E., Mogi K., Kikusui T., Izawa E.-I (2015) **Sex-reversed correlation between stress levels and dominance rank in a captive non-breeder flock of crows** *Hormones and behavior* **73**: 131–134. https://doi.org/10.1016/j.yhbeh.2015.07.012.

Patton J. Q (2005) **Meat sharing for coalitional support** *Evolution and human behavior* **26**: 137–157. https://doi.org/10.1016/j.evolhumbehav.2004.08.008.

Peterson L. M., Weckerly F. W (2018) **Social behavior and changes in foraging behavior in a gregarious ungulate** *Journal of mammalogy* **99**: 1422–1429. https://doi.org/10.1093/jmammal/gyy129.

Picard A. M., Mundry R., Auersperg A. M., Boeving E. R., Boucherie P. H., Bugnyar T., Dufour V., Emery N. J., Federspiel I. G., Gajdon G. K. et al (2020) Why preen others? **Predictors of allopreening in parrots and corvids and comparisons to grooming in great apes** *Ethology* **126**:

207–228. https://doi.org/10.1007/s10071-023-01794-x.

Pittman Q. J., Chen X., Mouihate A., Hirasawa M., Martin S (1998) **Arginine vasopressin, fever and**
temperature regulation *Progress in brain research* **119**: 383–392. https://doi.org/10.1016/s0079-
6123(08)61582-4.

Powell J. M., Inoue K., Wallace K. J., Seifert A. W., Young L. J., Kelly A. M (2023) **Distribution of**
vasopressin 1a and oxytocin receptor protein and mRNA in the basal forebrain and midbrain of
the spiny mouse (*Acomys cahirinus***)** *Brain structure and function* **228**: 413–431.
https://doi.org/10.1007/s00429-022-02581-z.

Rogers Flattery C. N., Coppeto D. J., Inoue K., Rilling J. K., Preuss T. M., Young L. J (2022)
Distribution of brain oxytocin and vasopressin V1a receptors in chimpanzees (*Pan***
troGglodytes): comparison with humans and other primate species** *Brain structure function* **227**:
1907–1919. https://doi.org/10.1007/s00429-021-02369-7.

Royle N. J., Smiseth P. T., Kölliker M (2012) **The evolution of parental care** (Oxford: Oxford
University Press).

Ruploh T., Bischof H. J., von Engelhardt N (2013) **Adolescent social environment shapes sexual**
and aggressive behaviour of adult male zebra finches (*Taeniopygia guttata***)** *Behavioral ecology*
and sociobiology **67**: 175–184. https://doi.org/10.1007/s00265-012-1436-y.

Scheid C., Range F., Bugnyar T (2007) **When, 2what, and whom to watch? Quantifying attention**
in ravens (*Corvus corax***) and jackdaws (***Corvus monedula***)** *Journal of comparative psychology*
121: 380–386. https://doi.org/10.1037/0735-7036.121.4.380.

Scheid C., Schmidt J., Noë R (2008) **Distinct pattern of food offering and co-feeding in rooks**
Animal behaviour **76**: 1701–1707. https://doi.org/10.1016/j.anbehav.2008.07.023.

Schuett W., Dall S. R. X (2010) **Appearance, "state," and behavior in male zebra finches,**
Taeniopygia guttata *Journal of ethology* **28**: 273–286. https://doi.org/10.1007/s10164-009-0181-9.

Schwab C., Bugnyar T., Schloegl C., Kotrschal K (2008) **Enhanced social learning between siblings**
in common ravens, *Corvus corax* *Animal behaviour* **75**: 501–508.
https://doi.org/10.1016/j.anbehav.2007.06.006.

Seed A. M., Clayton N. S., Emery N. J (2007) **Postconflict third-party affiliation in rooks,** *Corvus*

frugilegus Current biology **17**: 152–158. https://doi.org/10.1016/j.cub.2006.11.025.

Selvam R., Jurkevich A., Kang S. W., Mikhailova M. V., Cornett L. E., Kuenzel W. J (2013) **Distribution of the vasotocin Subtype Four receptor (VT4R) in the anterior pituitary gland of the chicken, *Gallus gallus*, and its possible role in the avian stress response** *Journal of neuroendocrinology* **25**: 56–66. https://doi.org/10.1080/07391102.2018.1464957.

Seyfarth R. M., Cheney D. L (1984) **Grooming, alliances and reciprocal altruism in vervet monkeys** *Nature* **308** 541–543. https://doi.org/10.1038/308541a0.

Shido O., Kifune A., Nagasaka T (1984) **Baroreflexive suppression of heat production and fall in body temperature following peripheral administration of vasopressin in rats** *Japanese journal of physiology* **34**: 397–406. https://doi.org/10.2170/jjphysiol.34.397.

Silk J. B (1999) **Male bonnet macaques use information about third-party rank relationships to recruit allies** *Animal behaviour* **58**: 45–51. https://doi.org/10.1006/anbe.1999.1129.

Stocker M., Munteanu A., Stöwe M., Schwab C., Palme R., Bugnyar T (2016) **Loner or socializer? Ravens' adrenocortical response to individual separation depends on social integration** *Hormones and behavior* **78**: 194–199. https://doi.org/10.1016/j.yhbeh.2015.11.009.

Stöwe M., Bugnyar T., Loretto M. C., Schloegl C., Range F., Kotrschal K (2006) **Novel object exploration in ravens (*Corvus corax*): effects of social relationships** *Behavioral processes* **73**: 68–75. https://doi.org/10.1007/s10336-007-0145-1.

Takeda K., Takahashi N., Izawa E.-I (2022) **Social encounters produce different autonomic response between dominants and subordinates in crows** *Royal society open science* **9**: 220972. https://doi.org/10.1098/rsos.220972.

Treisman M (1975) **Predation and the evolution of gregariousness. I. Models for concealment and evasion** *Animal behaviour* **23**: 779–800. https://doi.org/10.1016/0003-3472(75)90106-2.

Uhl F., Ringler M., Miller R., Deventer S. A., Bugnyar T., Schwab C (2019) **Counting crows: Population structure and group size variation in an urban population of crows** *Behavioral ecology* **30**: 57–67. https://doi.org/10.1093/beheco/ary157.

Vinacke W. E., Arkoff A (1957) **An experimental study of coalitions in the triad** *American sociological review* **22**: 406–414. https://doi.org/10.1177/014616727400100185.

Verspeek J., Staes N., van Leeuwen E. J. C., Eens M., Stevens J. M. G (2019) **Bonobo personality predicts friendship** *Scientific reports* **9**: 19245. https://doi.org/10.1038/s41598-019-55884-3.

von Bayern A. M. P., de Kort S. R., Clayton N. S., Emery N. J (2007) **The role of food- and object-sharing in the development of social bonds in juvenile jackdaws (*Corvus monedula*)** *Behaviour* **144**: 711–733. https://doi.org/10.1163/156853907781347826.

von Rueden C. R., Gurven M., Kaplan H (2008) **The multiple dimensions of male social status in an Amazonian society** *Evolution and human behavior* **29**: 402–415. https://doi.org/10.1016/j.evolhumbehav.2008.05.001.

von Rueden C. R., Jaeggi A. V (2016) **Men's status and reproductive success in 33 nonindustrial societies: effects of subsistence, marriage system, and reproductive strategy** *Proceedings of the national academy of sciences* **113**: 10824–10829. https://doi.org/10.1073/pnas.1606800113.

Watts D. P (2002) **Reciprocity and interchange in the social relationships of wild male chimpanzees** *Behaviour* **139**: 343–370. https://doi.org/10.1163/156853902760102708

Yang J., Yang Y., Xu H. T., Chen J. M., Liu W. Y., Lin B. C (2006) **Arginine vasopressin enhances periaqueductal gray synthesis and secretion of enkephalin and endorphin in the rat** *Brain research bulletin* **71**: 193–199. https://doi.org/10.1016/j.brainresbull.2006.09.003.

Zann R. A (1996) **The zebra finch – a synthesis of field and laboratory study** (Oxford: Oxford University Press).

Ziegler T. E., Crockford C (2017) **Neuroendocrine control in social relationships in non-human primates: Field based evidence** *Hormones and behavior* **91**: 107–121. https://doi.org/10.1016/j.yhbeh.2017.03.004.