Title:

Inhibition of V1a-like receptors disrupts male-male affiliative relationships formed by triadic cohabitation in large-billed crows

Authors and Affiliations:

Akiko Seguchi [a], Ei-Ichi Izawa [a]

[a] Department of Psychology, Keio University, Tokyo 108-8345, Japan

Corresponding to: E-I. Izawa

Mita 2-15-45, Minato-ku, Tokyo 108-8345, Japan.

E-mail: izawa@flet.keio.ac.jp

Tel: +81-3-5418-6400,

Fax: +81-3-5443-3867

Competing interests

We declare that we have no competing interests.

Funding

This work was financially supported by JSPS KAKENHI (#23H01058, #23H00074, #23K17645, #20H01787 to EI, and #19J22654 to AS) and a Keio University Grant-in-Aid for ICRP (#MKJ1905) to E.I.

Acknowledgments

We thank the members of the Animal Psychology Lab, Keio University, for inspiring scientific discussions and their support in the daily care of the animals.

Inhibition of V1a-like receptors disrupts male-male affiliative relationships formed by triadic cohabitation in large-billed crows

Abstract

Affiliative relationships, such as parent-offspring bonds, male-female pair-bonds, and within-sex coalitions/alliances, are beneficial for group-living animals to increase fitness and reproductive success. Male-male affiliative relationship, such as coalitions/alliances, have been reported in social mammals and birds in situations where three or more individuals interact, suggesting that male-male affiliative relationships can be formed via interactive situations involving three (or more) individuals. However, no experimental studies have tested this hypothesis on social animals. Many studies on mammals and birds have revealed that an affiliative relationship is underpinned by the oxytocin (OT)/vasopressin (VP) system. However, the role of the OT/VP system has been studied only in parent-offspring bonds and pair-bonds and is lacking in male-male affiliative relationships. In this study, we found the 2-week cohabitation of three male crows in an aviary facilitated affiliative relationship formation between two of the three crows (Experiment 1). In Experiment 2, we examined the effects of peripheral administration of vasopressin 1a receptor (V1aR) antagonist peripheral administration on the social behaviour of affiliated males. We found that the V1aR antagonist caused the disappearance of affiliative behaviour and the resurgence of agonistic behaviour in affiliated males. Our findings provide the first evidence for V1aR involvement in male-male affiliative relationships in vertebrates.

Keywords:

affiliative relationship, vasotocin, vasopressin, vasopressin 1a receptor, dominance relationship, birds

1. Introduction

Affiliative relationships, such as parent-offspring bonds [1, 2], male-female pair-bonds [3, 4, 5], and within-sex coalitions/alliances [6], are beneficial for individuals living in groups to increase the fitness of resource acquisition and, ultimately, reproductive success. In particular, an affiliative relationship between same-sex individuals, characterised by exchanging non-sexual affiliative behaviours, including social grooming (allogrooming in mammals and allopreening in birds), huddling, spending time together, and sharing food, is typically called a coalition or alliance which is a form of cooperation in agonistic conflicts to counter dominant individuals [6].

Male-male affiliative relationships such as coalitions/alliances have been reported in various mammals (including humans, *Homo sapiens*, [7, 8, 9, 10]; chimpanzees, *Pan troglodytes*, [11, 12, 13]; baboons, *Papio*, [14, 15]; macaques, *Macaca radiata*, [16]; *M. fuscata*, [17]; cetaceans bottlenose dolphins, *Tursiops truncates*, [18, 19, 20]; spotted dolphins, *Stenella frontalis*, [21]; sperm whales, *Physeter macrocephalus*, [22]; carnivore lions, *Panthera leo*, [23]; cheetahs, *Acinonyx jubatus*, [24, 25]; and birds, corvids, [26]). Most of these studies shared the fact that male-male affiliative relationships were formed in social interactions involving multiple participants, typically two individuals *vs* one individual, in both the wild and in captivity. Theoretical research to explore the social factor of coalition formation predicted that the social interaction among three individuals different in "power" and "activity" should be the minimal group size to produce coalition/alliance [27, 28, 29]. Several previous studies with human participants have examined the combinations of ranks that facilitate the formation of coalitions in game situations [30, 31]. However, to date, no studies have experimentally tested in non-human animals whether social interactions among three individuals, especially males, cause affiliative relationships between two of the three individuals.

Neural mechanisms underlying affiliative relationships have been extensively studied in mammals and birds. Neuropeptides of the arginine vasotocin (AVT) family include arginine vasopressin (VP) in mammals, arginine vasotocin (VT) in non-mammalian vertebrates, and oxytocin (OT). In particular, VP/VT has been shown to crucially modulate sexual behaviour, non-sexual affiliative behaviour, and aggression, particularly in males, based on the affiliative relationship with specific individuals such as partner females of pair-bonds [32, 33]. In mammals, the central infusion of VP in male monogamous prairie voles facilitates affiliative behaviour towards partner females [32]. Vasopressin 1a receptor (V1aR) antagonism of the lateral septum (LS) in male prairie voles selectively violates affiliative body contact behaviour with partner females [34]. The role of VT mediated by V1aR in affiliative relationships in birds is poorly understood. A previous study on monogamous zebra finches reported an increase of V1aR mRNA expression in the auditory pallium

associated with pair-bond formation [35]. In contrast, another study on zebra finches showed that antisense-

based VP knockdown had no effect on the formation of pair-bonds in males [36]. In terms of male-female

bonds, previous findings on the role of VT mediated by V1aR in affiliative relationships have been

inconsistent in birds.

The role of VT-mediated V1aR in inter-individual affiliation, other than pair-bonds, has been suggested

by species-comparison studies in estrildid finches with different gregariousness. Goodson and Wang [37]

found that VT neurons in the medial bed nucleus of the stria terminals (BSTm) were activated, in terms of

immediate early gene Fos expression, in response to the exposure of same-sex conspecifics in gregarious

species but did not in asocial ones. It has also been found that the activation of VT neurons is specifically

induced by social stimuli with a positive valence to enhance affiliative behaviour [37]. Antisense-based

knockdown of VT production in the BSTm of male zebra finches, a gregarious species, increased aggressive

behaviour and reduced courtship in colony housing situations [38]. The relevance of social affiliation and VT

signals, specifically in males of gregarious birds, has been suggested to be mediated by V1aR. A previous

study reported that the density of V1aR was higher in gregarious species than in asocial territorial species

[37]. V1aR antagonism in the LS of male zebra finches was found to reduce flock size in housing conditions

[39]. These previous findings suggest a role for VT and V1aR in the social affiliation of males, raising the

hypothesis that VT is mediated by V1aR in the affiliative relationships between males. However, the

involvement of neuroendocrine mechanisms, such as V1aR-mediated VT signals, in male-male affiliative

relationships in birds and mammals is yet to be reported.

Large-billed crows (*Corvus macrorhynchos*) are ideal models for investigating male-male affiliative

relationships. Non-breeder individuals form fission-fusion aggregations in a wide range of areas, behaving as

free floaters, akin to other *Corvus* species [40, 41, 42]. Such social dynamics in *Corvus* species provide an

ecological basis for the formation of affiliative relationships among males as counter-tactics against

dominant birds. Indeed, in various *Corvus* species, affiliative relationships are formed by exchanging

affiliative behaviours such as allopreening [26, 43, 44, 45, 46, 47]. Similar to other *Corvus* species, large-

billed crows, mostly in captive non-breeder groups, have been found to strictly form a linear dominance

hierarchy in males [40, 48, 49, 50] and allopreening often occurs between males. Allopreening is primarily

initiated by dominants but also occurs from subordinates [50].

Given the triadic interactive situation as an ecological factor that facilitates the formation of and

involvement of V1aR in affiliative relationships, we hypothesised that cohabitation in male triads should

facilitate the formation of male-male affiliative relationships in a dyad of non-breeder large-billed crows and

that V1aR should be involved in the maintenance of male-male affiliative relationships. We conducted two experiments to test our hypotheses. In Experiment 1, we introduced three male crows, those which had formed dominance relationships but not affiliative relationships prior to this study, to a triadic cohabitation in an aviary for two weeks and examined the formation of an affiliative relationship in one of the three possible dyads by comparing the occurrence of reciprocal allopreening before and after cohabitation. In Experiment 2, if an affiliated relationship was formed in certain male-male dyads in Experiment 1, we examined the effects of peripheral administration of a V1aR antagonist on social behaviour. We examined whether allopreening was reduced and, aggression by dominants and submission by subordinates increased between males with affiliated relationships.

2. Methods

Subjects and housing

We used 16 adult male crows aged between 4 and 7 years with body weights ranging from 640 to 790 g. Sex was determined using DNA from blood samples [51]. The birds were caught as yearlings in Tokyo and neighbouring areas during 2017–2019 with permission from the Ministry of the Environment Government of Japan (authorisation nos. 29030001, 30081482, 30030019, and 30224001). After being captured, the subject crows were housed in one of three different groups with 5–10 other crows, which were not used in this study (Table S1), in outdoor aviaries (W 5 m × D 10 m × H 3 m). Three to six months before the start of this study, the crows were transported to individual steel-mesh cages (W 57 cm × D 93 cm × H 63 cm) in an animal housing room to prevent the formation of dominance or affiliative relationships through physical interactions. Individual cages were placed side by side on shelves in the room. The birds were visible and audible to each other but without physical contact to reduce physical and social stress due to individual housing. Room temperature was set at $21 \pm 2°C$, and light/dark cycles were 13-h light and 11-h dark with light onset at 0800. Food (dried dog food and other supplements) and water were freely available in the cages. The experiments were conducted from Oct 2019 to Sep 2022. Owing to the accidental death of some birds and the restricted use of the experimental facility during the pandemic situation of COVID-19, a part of the planned experiments were not performed, and the data were unbalanced. The experimental and housing protocols adhered to the Japanese National Regulations for Animal Welfare and were approved by the Animal Care and Use Committee of Keio University (#20036, #A2022-055).

Determination of relative dominance in dyads

Seventeen birds were assigned to eight triads for the cohabitation experiment (Table 1). For the triad group

assignment, the relative dominance between each of the two birds was determined through three trials of

dyadic social encounters in an indoor aviary (W 2 m × D 2 m × H 2 m) using a procedure similar to that

reported in previous studies [48, 52]. In each trial, two birds were given a 5-min opportunity to freely

interact, and the winner and loser were determined according to the behavioural criteria used in previous

studies [40, 48, 52]. Specifically, the loser was identified as an individual exhibiting submissive behaviour

(i.e., submissive begging vocalisation and/or avoidance) in response to aggressive behaviour (i.e., jab, peck,

aggressive vocalisation, and approach). Within the dyad, the individual who won all three trials was defined

as dominant, and the other were defined as subordinates. All 26 dyads showed an apparent asymmetry of

win/loss numbers in the three trials and were considered to have formed a dominance relationship. Therefore,

26 dyads with clear dominance relationships were used in Experiment 1. The social encounter trials were

conducted between 10:00 and 15:00, with a 2-day interval between consecutive trials involving the same

dyads. No subjects showed allopreening in any trial, indicating that no affiliative relationship was formed in

any dyad at this stage.

Experiment 1: Formation of affiliative relationship through the triadic cohabitation

After determining the relative dominance positions of the individuals in the dyads of each triad, we tested

whether a 2-week cohabitation in triads facilitated the formation of affiliative relationships between two

specific birds in the triad. The formation of an affiliative relationship was evaluated based on the emergence

and increase in reciprocal allopreening exclusively in two specific birds in each triad. To examine the

increase in allopreening frequency of dyads, all possible dyads of each triad (i.e., three dyads) were housed

together in the experimental aviary on separate occasions for three days before the start and after the end of

the 2-week triadic cohabitation (hereafter referred to as pre-test and post-test, respectively). In the pre- and

post-tests, each of the two birds was transferred from its individual home cage into the aviary at 8:00 am, and

after 72 h, it was returned to the home cage at 8:00 am. On the day following the end of the pre-test period,

the three birds were introduced into the experimental aviary and housed together for two weeks. At the end

of the triadic cohabitation, the birds were transferred back to their individual home cages. At 8:00 am the

following day, each pair of the triad was again transferred into the aviary for 3 days as a post-test on separate

occasions.

The behaviour of the birds during the pre-/post-tests and triadic cohabitation was video-recorded using a

camera (EX-ZR4100, CASIO, Japan) on the ceiling of the aviary. The recording was conducted for 30 min

randomly between 9:00 and 17:00 every day in the pre- and post-tests and for the early (1−3 day), middle

(7−9 day), and last (12−14 day) three days of triadic cohabitation. Therefore, 90-min video data (30 min × 3

days) were used to analyse social interactions in each test, as well as 3 × 3-day blocks in the triadic

cohabitation period. In the aviary, three wooden perches were placed in parallel with 30 cm intervals at a

hight of 80 cm above the floor. Food, water, and a bathing tub were consistently provided, except during the

30-min video recording session.

Experiment 2: Effects of V1a receptor antagonism on affiliative relationship

To investigate the involvement of V1a-like receptors in the maintenance of male-male affiliative

relationships, the birds forming affiliated relationships in Experiment 1 were systematically administered

V1aR antagonist (SR 49059, Sigma-Aldrich, MO, USA) to examine the effects on social behaviour towards

the affiliated counterpart and the unaffiliated male. This V1aR antagonist was previously shown to be an

effective antagonist of the VT4 receptor in birds [53]. The effect of V1aR antagonist was tested with high

dose (0.3 mL × 1 mg/mL vehicle solution) and low dose (0.3 mL × 0.1 mg/mL vehicle solution), compared

with 0.3 mL vehicle solution as control, via intramuscular injection onto the chest of subject crows. The

vehicle solution contained 10% dimethyl sulfoxide in a saline solution.

In each trial, the V1aR antagonist (high or low dose) or vehicle was administered to either the dominant

or subordinate. Both the dominant and subordinate of each affiliated dyad were tested in 8 trials: 3 trials with

the high-dose condition, 3 trials with the low-dose condition, and 2 trials with the vehicle control condition.

The trial order of the dominance position of the focal individual (dominant or subordinate) and the drug

conditions were counterbalanced within and between dyads. After the affiliated dyad test, the birds

underwent three additional trials of dyadic interactions without drug injection to ensure the maintenance or

recovery of the relationship. At least one week after this recovery trial, both birds of affiliated dyads received

another set of five trials with the unaffiliated individual (the third bird in the triadic cohabitation in

Experiment 1) as an unaffiliated dyad test with the same procedure as that used in the affiliated dyad test.

For each trial, either the dominant or the subordinate individual received an injection 1-h before the start

of the trial (hereafter referred to as the focal individual). During the 1-h period, the focal bird was isolated

from other birds in an individual cage. A trial was started soon after the focal individual was introduced into

the indoor aviary (W 2 m × D 2 m × H 2 m), to which another individual was introduced 1 min beforehand.

The trial procedure was identical to that of Experiment 1, and the birds were allowed to interact freely for 30

min. Given the typical social behaviour in the dyad and interactions between dominant and subordinate

large-billed crows [40, 48, 52], we coded allopreening; aggressive behaviour such as peck, jab, and kick; and

submissive begging vocalisations from video-recorded data for offline analysis. The inter-trial interval was

three and five days. Each bird participated in one daily trial. All trials were conducted between 8:30 and

12:00.

3. Analyses

Experiment 1: Formation of affiliative relationship through the triadic cohabitation

We defined an affiliative relationship as a dyad which exchanged allopreening reciprocally, but not

unidirectionally, between two birds, independent of the dominance relationship. This definition meets the

behavioural characteristics of affiliative relationships in primates and other social animals, such as

maintaining close proximity and exchanging affiliative behaviour [54]. An allopreening bout was counted

when a bird passed its bill through the feathers of another bird for longer than 2 s [51]. Given the absence of

allopreening in any dyad before triadic cohabitation, we defined a dyad as having an affiliative relationship

in which both individuals initiated allopreening in 10 or more bouts per 30 min in either of 3-day block in the

triadic cohabitation. A dyad that did not meet this behavioural criterion was treated as having not formed

affiliative relationship. To confirm the exclusive formation of affiliative relationships through triadic

cohabitation, a discriminant analysis was performed to examine whether the numbers of dominant-initiated

and subordinate-initiated allopreening in the pre- and post-triadic cohabitation tests could distinguish

affiliated dyads from those before forming their affiliative relationships and from unaffiliated dyads. Leave-

one-out cross validation was used to calculate the correct discrimination rate. The discriminant analysis was

performed using SPSS Statistics version 29. Allopreening was coded using BORIS v.7 [55].

Experiment 2: Effects of V1a receptor antagonism on affiliative relationship

To compare the number of social behaviours in the test trials among the drug conditions and between

dominants and subordinates, we applied generalised linear mixed models (GLMMs) with Poisson error

distribution and the log link function separately for allopreening behaviours, aggressive behaviours, and

submissive vocalisations. The model included the number of behaviours per 30 min as the response variable;

dominance (dominant or subordinate), *drug condition* [V1aR antagonist (V1aRA) high-dose, V1aRA low-

dose, or vehicle], and their interaction as explanatory variables; and *trial number* and *identity of focal*

individual as random factors. If the interaction was significant, we stratified the dataset between the

dominant and subordinate GLMMs to examine *drug* effects. The significance of variables in each model was

evaluated based on the Wald χ^2 statistics. All statistical analyses were performed using R version 4.2.2. with

the package 'lme4' [56] for fitting GLMMs to the data. The significance level was set at 5%.

4. Results

Experiment 1: Formation of affiliative relationship through the triadic cohabitation

In six of the eight triads, the 2-week cohabitation resulted in the formation of an affiliative relationship

exclusively in one dyad but not in the other dyads (Figure 1). A specific dyad in each triad (#1 – #6), which

showed no allopreening in the pre-test, exhibited exclusively increased allopreening through the 2-week

triadic cohabitation. In the post-test of these six dyads, allopreening was initiated by both dominants and

subordinates, whereas they did not exhibit reciprocal allopreening in the pre-test, and no such allopreening

emerged in any other dyads in either test (Figure 2). Discriminant analysis based on the number of

allopreening initiated by dominants and subordinates within the dyads yielded two discriminant functions.

Using the leave-one-out method, the two discriminant functions could distinguish the affiliated dyads in the

post-test from themselves in the pre-test and the other unaffiliated dyads with an 83.3% correct rate (i.e., five

of six dyads were classified as affiliated dyads; see the outputs for Table S2). These results indicate that the

2-week cohabitation in the triad facilitated the formation of the affiliative relationship exclusively in two of

the three birds characterised by the exchange of allopreening.

Because one bird with a physical injury was discarded in the subsequent experiment, five of the six dyads

forming an affiliated relationship were used for the test in Experiment 2.

Experiment 2: Effects of V1a receptor antagonism on affiliative relationship

Allopreening in the affiliated dyads was severely reduced by the administration of the V1aR antagonist to

the dominants and subordinates (Figure 3A). The best model of the GLMM analysis on the allopreening in

the affiliated dyads showed a significant effect of the *drug condition*, namely the *vehicle* condition with a

positive coefficient (estimate ± s.e. = 1.639 ± 0.217, $p < 0.001$) and the *high dose* condition with a negative

coefficient (estimate ± s.e. = -0.724 ± 0.313, $p < 0.05$; Table 2) but no significance of the dominance and the

interaction factors. These results indicate that the numbers of allopreening decreased dose-dependently

among the three conditions, irrespective of the relative dominance position in the affiliated dyads. In

contrast, no effect was found on allopreening by V1aR antagonist administration in any birds of the

unaffiliated dyads (Figure 3B). The GLMM analysis was not performed on unaffiliated dyads because no

allopreening occurred under any condition or with any bird.

Administration of the V1aR antagonist in the affiliated dyads elevated aggression from dominants to

subordinates, but not from subordinates to dominants (Figure 4A). The best model of the GLMM analysis on

the aggressive behaviours in the affiliated dyads showed a significant effect of the *drug condition*,

specifically the high dose condition with a positive coefficient (estimate ± s.e. = 0.791 ± 0.251, $p < 0.01$) but

no significance of the vehicle condition ($p = 0.54$; Table 3), indicating the significant increase of dominants'

aggression upon the administration of a high dose of the V1aR antagonist. However, no such increase in

aggression by the V1aR antagonist was found in subordinates (Figure 4A; GLMM was not performed owing

to no aggression in any condition or with any bird). Administration of the V1aR antagonist did not increase

aggression in unaffiliated dyads (*Vehicle*, $p = 0.929$; *V1aRA high-dose*, $p = 0.89$; Figure 4B and Table 4).

Administration of the V1aR antagonist in the affiliated dyads increased the submissive begging

vocalisation of the subordinates but not the dominants in the affiliated dyads (Figure 5A). The best model of

the GLMM analysis of the submissive vocalisations in affiliated dyads produced a significant effect of the

drug condition, namely the high dose condition with a positive coefficient (estimate ± s.e. = 1.929 ± 0.253, *p*

< 0.001) but no significance of the vehicle condition ($p = 0.455$; Table 5). In contrast, no such increase in

submissive vocalisation was found in any condition in the unaffiliated dyads (*Vehicle*, $p = 0.791$; *V1aRA*

high-dose, $p = 0.777$; Figure 5B and Table 6).

5.   Discussion

The present study suggested that in Experiment 1, triadic cohabitation facilitated the formation of an

affiliative relationship exclusively between two specific birds of each triad and that in Experiment 2, the

peripheral administration of a V1aR antagonist destroyed the affiliated relationship by the resurgence of

aggressive and submissive/avoiding behaviour depending on the relative dominance but had no effect on

social behaviour within unaffiliated dyads. These findings represent the first evidence not only in birds but

also in mammals and other taxa, suggesting the facilitative role of the triadically interactive environment in

the formation of male-male affiliative relationships and the crucial involvement of V1aR in the maintenance

of male-male affiliation as has been suggested in mother-infant bonds and pair-bonds.

Formation of affiliative relationship through the triadic cohabitation

The results of Experiment 1 suggest that the triadically interactive situation plays a crucial role in

facilitating the formation of male–male affiliative relationships exclusively between two of the three in the

triad. Our finding is in accordance with the ideas of previous theoretical studies [27-29], which explained

that the triad, including individuals with different powers, is the minimal social group that promotes the

formation of a coalition in the form of two against one. This two-vs-one form has been confirmed in the

previous behavioural observation studies regarding the same-sex alliance and coalition of primates,

cetaceans, and corvids both in the wild and captivity [7-26]. Our findings in Experiment 1 provide the first

experimental evidence that a triadically interactive situation, such as 2-week cohabitation in a triad, indeed

facilitates the formation of a male-male affiliative relationship, which fills the gap between previous

theoretical and behavioural observation studies.

Two of the eight triads tested in the present study did not form affiliative relationships, even after 2-week

cohabitation. As shown in Figure 1, allopreening occurred slightly in two males of the two triads during the

cohabitation period but finally disappeared. According to the video observation of behaviour during the

cohabitation period, one possible cause for the failure of affiliative relationship formation in those triads is

the avoidance response of one male to the approach to initiate allopreening by the other. Another possible

cause is intervention by the third-party male in attempts of socially affiliative interactions between two

males. Third-party individual's intervention in affiliative interactions between individuals was found in a

captive group of ravens [57]. There is still an open question as to what social characteristics of individuals in

dyads could facilitate or inhibit allopreening and what individual and/or social factors could drive the

intervention of others' affiliative behaviour in birds and mammals. An in-depth analysis of social behaviour

using the triadic cohabitation paradigm used in this study may allow us to elucidate the behavioural and

psychological mechanisms for forming male-male affiliative relationships.

Effects of V1a receptor antagonism on affiliative relationship

Systemic injection of a V1aR antagonist to males with affiliative relationships caused a dose-dependent

decrease in allopreening and the resurgence of agonistic behaviour associated with relative dominance, such

as aggressive behaviour of the dominant and submissive behaviour (i.e., avoiding and/or submissive begging

calls) of the subordinate. In contrast, dominant and subordinate males without an affiliated relationship

showed no behavioural effects by V1aR blocking. These results suggest that V1aR may be involved in the

suppression of dominance-dependent aggressive/submissive behaviour and facilitate affiliative behaviour,

thus playing a crucial role in the formation and maintenance of affiliated relationships between males.

Given the peripheral administration of the drug in this study, the behavioural effects found in affiliated

dyads by V1aR antagonists do not necessarily reflect the role of V1aR in the brain. Peripheral nonapeptide

receptors have been suggested to influence a variety of homeostatic functions such as hydromineral balance

[58] and thermoregulation [59]. For example, maternal crouching in rodents serves a thermoregulatory

function [60] and, is regulated by V1aR [62]. V1aR has also been suggested to be involved in peripheral

thermoregulation [62, 63, 64]. Thus, the effects of maternal crouching by V1aR blocking in rodents may be

indirectly caused by thermoregulation via V1aR antagonism. Similarly, maternal licking and grooming in

rodents may indirectly involve peripheral V1aR [62]. Given the fact that during maternal licking and

grooming, mothers often ingest water to maintain the hydromineral balance [65] based on peripheral V1aR

[66], maternal licking and grooming may be regulated indirectly by peripheral V1aR. Although the

distribution of V1aR (i.e., VT4R) has been examined in the brain and pituitary gland [67, 68] but not in

peripheral organs in birds, previous studies reporting the involvement of VT in renal osmoregulation [58, 69]

suggest the presence of V1aR in the periphery of birds. Thus, we cannot exclude the possibility that

peripheral injection of a V1aR antagonist affected osmoregulation and secondary social behaviour effects.

Although we did not measure the effects of body temperature and the amount of drinking/evacuation

following V1aR antagonist injection, such homeostatic effects on the periphery by V1aR blocking are

unlikely to occur selectively in affiliated dyads. To determine the involvement of the central V1aR in the

social behaviour of affiliated males, it will be necessary to examine the effect of central infusion of V1aR

antagonists in future studies.

322   The present findings on the involvement of V1aR in promoting inter-individual affiliation, such as male-

female pair-bonds and group formation, are consistent with earlier findings in in mammals and birds. In

monogamous prairie voles, the injection of V1aR antagonism into the lateral septum of males selectively

reduced body-contact affiliation to partner females compared to stranger females [34]. In zebra finches, the

formation of pair bonds was found to be accompanied by increased activity in terms of immediate early gene

ZENK expression in neurons with V1aR mRNA expression in the auditory pallium [35]. Regarding group

formation in gregarious finch species, VT neurons in males and females are activated in response to

exposure to a conspecific stimulus [37]. It has also been found that intraseptal infusions of a V1aR antagonist

reduced gregariousness in male zebra finches [39]. Although these previous studies did not reveal the

modulatory role of V1aR in specific affiliative behaviours as a facilitative factor for the promotion of male-

conspecific affiliation, our finding of male-male affiliation in crows could support the role of V1aR in the

formation and/or facilitation of inter-individual affiliation.

334   Given the results of Experiment 2, the role of V1aR in the inter-individual affiliation might be effective

in inhibiting agonistic behaviour (i.e., aggression and submission), specifically in affiliated males and not in

unaffiliated ones. Similar context-specific V1aR involvement in the control of aggressive behaviour in males

has been suggested in previous studies on birds. A previous study on violet-eared waxbills (*Uraeginthus*

granatina) found that peripheral injection of a V1aR antagonist disinhibited and increased the aggression of

subordinate males in a territorial challenge context, although V1aR decreased the aggression of a dominant

male toward rival males in a mate competition context [70]. To clarify the role of V1aR and the relevant

VP/VT signals in facilitating affiliative behaviour and/or inhibiting agonistic behaviour for the male-male

affiliation, central infusion of a V1aR antagonist and assessment of V1aR-/VT-neuron activities should be

investigated in future studies using crows.

In summary, the present study suggests, for the first time in social vertebrates, that male-male affiliative

relationships share neural underpinnings, such as the VP/VT systems, with inter-individual social

affiliations, such as pair and mother-infant bonds, which have been extensively studied in mammals and

birds. Future research should be directed to investigate the role of VP/VT as well as OT/MT and their

receptor subtypes in social affiliation between individuals of the same sex, which will allow us to understand

inter-individual affiliation in the social lives of animals, including humans.

References

1. Clutton-Brock TH. The evolution of parental care. Princeton: Princeton University Press; 1991.

2. Royle NJ, Smiseth PT, Kölliker M. The evolution of parental care. Oxford, UK: Oxford University Press; 2012.

3. Black JM. Partnerships in birds: the study of monogamy. Oxford, UK: Oxford University Press; 1996.

4. Kleiman DG. Monogamy in mammals. Q Rev Biol. 1977;52(1):39-69. doi: 10.1086/409721, PMID 857268.

5. Bales KL, Ardekani CS, Baxter A, Karaskiewicz CL, Kuske JX, Lau AR, Savidge LE, Sayler KRm Witczak LR. What is a pair bond? Horm Behav. 2021;136:105062. doi: 10.1016/j.yhbeh.2021.105062, PMID 34601430.

6. Brent LJ, Chang SW, Gariépy JF, Platt ML. The neuroethology of friendship. Ann N Y Acad Sci. 2014;1316(1):1-17. doi: 10.1111/nyas.12315, PMID 24329760.

7. Macfarlan SJ, Walker RS, Flinn MV, Chagnon NA. Lethal coalitionary aggression and long-term alliance formation among Yanomamö men. Proc Natl Acad Sci U S A. 2014;111(47):16662-9. doi: 10.1073/pnas.1418639111, PMID 25349394.

8. von Rueden CR, Gurven M, Kaplan H. The multiple dimensions of male social status in an Amazonian society. Evol Hum Behav. 2008;29(6):402-15. doi: 10.1016/j.evolhumbehav.2008.05.001, PMID 19884954.

9. von Rueden CR, Jaeggi AV. Men's status and reproductive success in 33 nonindustrial societies: effects of subsistence, marriage system, and reproductive strategy. Proc Natl Acad Sci U S A. 2016;113(39):10824-9. doi: 10.1073/pnas.1606800113, PMID 27601650.

10. Patton JQ. Meat sharing for coalitional support. Evol Hum Behav. 2005;26(2):137-57. doi: 10.1016/j.evolhumbehav.2004.08.008.

11. Watts DP. Reciprocity and interchange in the social relationships of wild male chimpanzees. Behaviour. 2002;139(2):343-70. doi: 10.1163/156853902760102708.

12. Muller MN, Mitani JC. Conflict and cooperation in wild chimpanzees. Asv Study Behav. 2005;35:275-331. doi: 10.1016/S0065-3454(05)35007-8.

13. Gilby IC, Brent LJ, Wroblewski EE, Rudicell RS, Hahn BH, Goodall J, Pusey AE. Fitness benefits of coalitionary aggression in male chimpanzees. Behav Ecol Sociobiol. 2013;67(3):373-81. doi: 10.1007/s00265-012-1457-6, PMID 23459197.

14. Bercovitch FB. Coalitions, cooperation and reproductive tactics among adult male baboons. Anim
Behav. 1988;36(4):1198-209. doi: 10.1016/S0003-3472(88)80079-4.

15. Noë R. A model of coalition formation among male baboons with fighting ability as the crucial
parameter. Anim Behav. 1994;47(1):211-3. doi: 10.1006/anbe.1994.1024.

16. Silk JB. Male bonnet macaques use information about third-party rank relationships to recruit allies.
Anim Behav. 1999;58(1):45-51. doi: 10.1006/anbe.1999.1129, PMID 10413539.

17. Kutsukake N, Hasegawa T. Dominance turnover between an alpha and a beta male and dynamics of
social relationships in Japanese macaques. Int J Primatol. 2005;26(4):775-800. doi: 10.1007/s10764-005-
5308-4.

18. Connor RC, Smolker RA, Richards AF. Dolphin alliances and coalitions. In: Harcourt AH, de Waal
FBM, editors. Coalitions and alliances in humans and other animals. Oxford, UK: Oxford University
Press; 1992a. p. 415-43.

19. Connor RC, Smolker RA, Richards AF. Two levels of alliance formation among male bottlenose
dolphins (*Tursiops* sp.). Proc Natl Acad Sci USA. 1992b;89(3):987-90. doi: 10.1073/pnas.89.3.987,
PMID 11607275.

20. Connor RC, Heithaus MR, Barre LM. Superalliance of bottlenose dolphins. Nature.
1999;397(6720):571-2. doi: 10.1038/17501.

21. Green ML, Herzing DL, Baldwin JD. Molecular assessment of mating strategies in a population of
Atlantic spotted dolphins. PLOS ONE. 2015;10(2):e0118227. doi: 10.1371/journal.pone.0118227, PMID
25692972.

22. Kobayashi H, Whitehead H, Amano M. Long-term associations among male sperm whales (*Physeter
macrocephalus*). PLOS ONE. 2020;15(12):e0244204. doi: 10.1371/journal.pone.0244204, PMID
33362216.

23. Bygott JD, Bertram BCR, Hanby JP. Male lions in large coalitions gain reproductive advantages. Nature.
1979;282(5741):839-41. doi: 10.1038/282839a0.

24. Caro TM. Behavioral solutions to breeding cheetahs in captivity: insights from the wild. Zoo Biol.
1993;12(1):19-30. doi: 10.1002/zoo.1430120105.

25. Caro TM. Cheetahs of the Serengeti plains: group living in an asocial species. Chicago: University of
Chicago Press; 1994.

26. Fraser ON, Bugnyar T. Reciprocity of agonistic support in ravens. Anim Behav. 2012;83(1):171-7. doi:
10.1016/j.anbehav.2011.10.023, PMID 22298910.

27. Caplow T. A theory of coalitions in the triad. American Sociological Review. 1956;21(4):489-93. doi:

10.2307/2088718.

28. Mesterton-Gibbons M, Gavrilets S, Gravner J, Akçay E. Models of coalition or alliance formation. J

Theor Biol. 2011;274(1):187-204. doi: 10.1016/j.jtbi.2010.12.031, PMID 21195717.

29. Koykka C, Wild G. Concessions, lifetime fitness consequences, and the evolution of coalitionary

behavior. Behav Ecol. 2017;28(1):20-30. doi: 10.1093/beheco/arw126.

30. Vinacke WE, Arkoff A. An experimental study of coalitions in the triad. American Sociological Review.

1957;22(4):406-14. doi: 10.2307/2089158.

31. Kelley HH, Arrowood AJ. Coalitions in the triad: critique and experiment. Sociometry. 1960;23(3):231-

44. doi: 10.2307/2785888.

32. Lim MM, Young LJ. Vasopressin-dependent neural circuits underlying pair bond formation in the

monogamous prairie vole. Neuroscience. 2004;125(1):35-45. doi: 10.1016/j.neuroscience.2003.12.008,

PMID 15051143.

33. Goodson JL, Thompson RR. Nonapeptide mechanisms of social cognition, behavior and species-specific

social systems. Curr Opin Neurobiol. 2010;20(6):784-94. doi: 10.1016/j.conb.2010.08.020, PMID

20850965.

34. Liu Y, Curtis JT, Wang Z. Vasopressin in the lateral septum regulates pair bond formation in male

prairie voles (*Microtus ochrogaster*). Behav Neurosci. 2001;115(4):910-9. doi: 10.1037/0735-

7044.115.4.910, PMID 11508730.

35. Tomaszycki ML, Atchley D. Pairing increases activation of V1aR, but not OTR, in auditory regions of

zebra finches: the importance of signal modality in nonapeptide-social behavior relationships. Integr

Comp Biol. 2017;57(4):878-90. doi: 10.1093/icb/icx043, PMID 28992311.

36. Kelly AM, Goodson JL. Hypothalamic oxytocin and vasopressin neurons exert sex-specific effects on

pair bonding, gregariousness, and aggression in finches. Proc Natl Acad Sci U S A. 2014;111(16):6069-

74. doi: 10.1073/pnas.1322554111, PMID 24711411.

37. Goodson JL, Wang Y. Valence-sensitive neurons exhibit divergent functional profiles in gregarious and

asocial species. Proc Natl Acad Sci U S A. 2006;103(45):17013-7. doi: 10.1073/pnas.0606278103,

PMID 17071744.

38. Kelly AM, Goodson JL. Functional significance of a phylogenetically widespread sexual dimorphism in

vasotocin/vasopressin production. Horm Behav. 2013;64(5):840-6. doi: 10.1016/j.yhbeh.2013.09.006,

PMID 24100197.

39. Kelly AM, Kingsbury MA, Hoffbuhr K, Schrock SE, Waxman B, Kabelik D, Thompson RR, Goodson JL. Vasotocin neurons and septal V1a-like receptors potently modulate songbird flocking and responses to novelty. Horm Behav. 2011;60(1):12-21. doi: 10.1016/j.yhbeh.2011.01.012, PMID 21295577.

40. Izawa E-I, Watanabe S. Formation of linear dominance relationship in captive jungle crows (*Corvus macrorhynchos*): implications for individual recognition. Behav Processes. 2008;78(1):44-52. doi: 10.1016/j.beproc.2007.12.010, PMID 18294782.

41. Loretto MC, Schuster R, Itty C, Marchand P, Genero F, Bugnyar T. Fission-fusion dynamics over large distances in raven non-breeders. Sci Rep. 2017;7(1):380. doi: 10.1038/s41598-017-00404-4, PMID 28336913.

42. Boucherie PH, Loretto MC, Massen JJM, Bugnyar T. What constitutes "social complexity" and "social intelligence" in birds? Lessons from ravens. Behav Ecol Sociobiol. 2019;73(1):12. doi: 10.1007/s00265-018-2607-2, PMID 30930524.

43. Emery NJ, Seed AM, von Bayern AMP, Clayton NS. Cognitive adaptations of social bonding in birds. Philos Trans R Soc Lond B Biol Sci. 2007;362(1480):489-505. doi: 10.1098/rstb.2006.1991, PMID 17255008.

44. Scheid C, Schmidt J, Noë R. Distinct patterns of food offering and co-feeding in rooks. Anim Behav. 2008;76(5):1701-7. doi: 10.1016/j.anbehav.2008.07.023.

45. Emery N, de Kort S, von Bayern A, Clayton N. The role of food- and object-sharing in the development of social bonds in juvenile jackdaws (*Corvus monedula*). Behaviour. 2007;144(6):711-33. doi: 10.1163/156853907781347826.

46. Braun A, Bugnyar T. Social bonds and rank acquisition in raven nonbreeder aggregations. Anim Behav. 2012;84(6):1507-15. doi: 10.1016/j.anbehav.2012.09.024, PMID 23264693.

47. Morales Picard A, Mundry R, Auersperg AM, Boeving ER, Boucherie PH, Bugnyar T, Dufour V, Emery NJ, Federspiel IG, Gajdon GK, et al. Why preen others? Predictors of allopreening in parrots and corvids and comparisons to grooming in great apes. Ethology. 2020;126(2):207-28. doi: 10.1111/eth.12999.

48. Nishizawa K, Izawa EI, Watanabe S. Neural-activity mapping of memory-based dominance in the crow: neural networks integrating individual discrimination and social behaviour control. Neuroscience. 2011;197:307-19. doi: 10.1016/j.neuroscience.2011.09.001, PMID 21939742.

49. Ode M, Asaba A, Miyazawa E, Mogi K, Kikusui T, Izawa E-I. Sex-reversed correlation between stress levels and dominance rank in a captive non-breeder flock of crows. Horm Behav. 2015;73:131-4. doi: 10.1016/j.yhbeh.2015.07.012, PMID 26193673.

50. Miyazawa E, Seguchi A, Takahashi N, Motai A, Izawa E-I. Different patterns of allopreening in the

same-sex and opposite-sex interactions of juvenile large-billed crows (*Corvus macrorhynchos*).

Ethology. 2020;126(2):195-206. doi: 10.1111/eth.12992.

51. Fridolfsson A-K, Ellegren H. A simple and universal method for molecular sexing of non-ratite birds. J

Avian Biol. 1999;30(1):116-21. doi: 10.2307/3677252.

52. Takeda K, Takahashi N, Izawa EI. Social encounters produce different autonomic response between

dominants and subordinates in crows. R Soc Open Sci. 2022;9(10):220972. doi: 10.1098/rsos.220972,

PMID 36300140

53. Kuenzel WJ, Hancock M, Nagarajan G, Aman NA, Kang SW. Central effect of vasotocin 4 receptor

(VT4R/V1aR) antagonists on the stress response and food intake in chicks given neuropeptide Y (NPY).

Neurosci Lett. 2016;620:57-61. doi: 10.1016/j.neulet.2016.03.036, PMID 27016389.

54. Fedurek P, Dunbar RIM. What does mutual grooming tell us about why chimpanzees groom? Ethology.

2009;115(6):566-75. doi: 10.1111/j.1439-0310.2009.01637.x.

55. Friard O, Gamba M. BORIS: a free, versatile open-source event-logging software for video/audio coding

and live observations. Methods Ecol Evol. 2016;7(11):1325-30. doi: 10.1111/2041-210X.12584.

56. Bates D, Mächler M, Bolker B, Walker S. Fitting linear mixed-effects models using lme4. J Stat Soft.

2015;67(1):1-48. doi: 10.18637/jss.v067.i01.

57. Massen JJ, Szipl G, Spreafico M, Bugnyar T. Ravens intervene in others' bonding attempts. Curr Biol.

58. Goldstein DL. Regulation of the avian kidney by arginine vasotocin. Gen Comp Endocrinol.

2006;147(1):78-84. doi: 10.1016/j.ygcen.2005.09.018, PMID 16288753.

59. Hassinen E, Pyörnilä A, Hissa R. Cardiovascular and thermoregulatory responses to vasotocin and

angiotensin II in the pigeon. Comp Biochem Physiol A Mol Integr Physiol. 1999;123(3):279-85. doi:

10.1016/s1095-6433(99)00061-6, PMID 10501020.

60. Jans JE, Woodside BC. Nest temperature: effects on maternal behavior, pup development, and

interactions with handling. Dev Psychobiol. 1990;23(6):519-34. doi: 10.1002/dev.420230607, PMID

2272408.

61. Bosch OJ, Neumann ID. Brain vasopressin is an important regulator of maternal behavior independent of

dams' trait anxiety. Proc Natl Acad Sci U S A. 2008;105(44):17139-44. doi: 10.1073/pnas.0807412105,

PMID 18955705.

62. Pittman QJ, Chen X, Mouihate A, Hirasawa M, Martin S. Arginine vasopressin, fever and temperature

regulation. Prog Brain Res. 1998;119:383-92. doi: 10.1016/s0079-6123(08)61582-4, PMID 10074801.

63. Shido O, Kifune A, Nagasaka T. Baroreflexive suppression of heat production and fall in body

temperature following peripheral administration of vasopressin in rats. Jpn J Physiol. 1984;34(3):397-

406. doi: 10.2170/jjphysiol.34.397, PMID 6492492.

64. Yang J, Yang Y, Xu HT, Chen JM, Liu WY, Lin BC. Arginine vasopressin enhances periaqueductal

gray synthesis and secretion of enkephalin and endorphin in the rat. Brain Res Bull. 2006;71(1-3):193-9.

doi: 10.1016/j.brainresbull.2006.09.003, PMID 17113946.

65. Gubernick DJ, Alberts JR. Maternal licking of young: resource exchange and proximate controls.

Physiol Behav. 1983;31(5):593-601. PMID 6665051.

66. Koshimizu TA, Nakamura K, Egashira N, Hiroyama M, Nonoguchi H, Tanoue A. Vasopressin V1a and

V1b receptors: from molecules to physiological systems. Physiol Rev. 2012;92(4):1813-64. doi:

10.1152/physrev.00035.2011, PMID 23073632.

67. Leung CH, Goode CT, Young LJ, Maney DL. Neural distribution of nonapeptide binding sites in two

species of songbird. J Comp Neurol. 2009;513(2):197-208. doi: 10.1002/cne.21947, PMID 19132730.

68. Selvam R, Jurkevich A, Kang SW, Mikhailova MV, Cornett LE, Kuenzel WJ. Distribution of the

vasotocin Subtype Four receptor (VT4R) in the anterior pituitary gland of the chicken, *Gallus gallus*, and

its possible role in the avian stress response. J Neuroendocrinol. 2013;25(1):56-66. doi: 10.1111/j.1365-

2826.2012.02370.x, PMID 22849330.

69. Braun EJ, Dantzler WH. Endocrine regulation of avian renal function. J Exp Zool. 1984;232(3):715-23.

doi: 10.1002/jez.1402320341, PMID 6394708.

70. Goodson JL, Kabelik D, Schrock SE. Dynamic neuromodulation of aggression by vasotocin: influence

of social context and social phenotype in territorial songbirds. Biol Lett. 2009;5(4):554-6. doi:

10.1098/rsbl.2009.0316, PMID 19493876.

     Figure captions

     Figure 1. Total number of allopreening behaviours in each dyad across the experimental periods; pre-tests, 2-week

     triadic cohabitation (early, middle, and last), and post-tests. Data plots are presented with mean ± 1 s.e. Panels

     indicate triads (#1 − #8). Bright-coloured plots (pink, green, purple, blue, orange, and red) in the #1 − #6 triads

     represent dyads that showed the formation of affiliated relationships. Darker plots denote the dyads that did not

     form affiliative relationships.

     Figure 2. Scatter plots of allopreening behaviours initiated by dominants (x-axis) versus subordinates (y-axis) of

     dyads in the pre-tests (opened) and the post-tests (shaded). Coloured plots represent dyads wherein affiliative

     relationships were formed. Allopreening occurred exclusively in the coloured dyads after the cohabitation (i.e., post-

     test). Colour corresponds to the dyad in Figure 1. Data are presented as the mean with natural log-transformed.

     Figure 3. Effects of V1a receptor antagonist (V1aRA) on the number of allopreening behaviour. (A) Effects of

     V1aRA in affiliated dyads. Allopreening initiated by the dominant (DOM, left panel) and the subordinate (SUB,

     right panel) birds was reduced drastically in the V1aRA *high-dose* conditions and slightly in the *low-dose* condition,

     compared with the *vehicle* condition. (B) Effects of V1aRA in unaffiliated dyads. No effect of V1aRA on

     allopreening either by the dominant (DOM, left panel) or the subordinate (SUB, right panel) was found in

     unaffiliated dyads. Plots show the individual data at each trial. In each triad, circles represent birds affiliated with

     another male, while triangles represent those with no affiliation. The top, middle, and bottom of the boxes depict

     the 75, 50, and 25 percentiles, respectively, with whiskers representing the maximum and minimum values.

     Numbers in the brackets below the boxes denote the number of samples (no. of birds × no. of trials). Asterisks

     indicate significant differences (Wald χ^2 statistics, *** $p < 0.001$). Colours correspond to the affiliated dyads in

     Figure 1.

     Figure 4. Effects of V1aRA on the number of aggressive behaviours. (A) Effects of V1aRA in affiliated dyads.

     Aggressive behaviour initiated by the dominant bird (DOM, left panel) drastically increased in the V1aRA *high-*

     dose condition but not in the *low-dose* condition. Such increase of aggression occurred in all conditions of the

     subordinates (SUB, right panel). (B) No effects were observed in any condition in both the dominants and the

     subordinates of unaffiliated dyads. Plots show the individual data at each trial. In each triad, circles represent birds

     affiliated with another male, while triangles represent those with no affiliation. The top, middle, and bottom of the

boxes depict the 75, 50, and 25 percentiles, respectively, with whiskers representing the maximum and minimum

values. Numbers in the brackets below the boxes denote the number of samples (no. of birds × no. of trials).

Asterisks indicate significant differences (Wald χ^2 statistics, *** $p < 0.001$, ** $p < 0.01$). Colours correspond to the

affiliated dyads in Figure 1.

Figure 5. Effects of V1aRA on the number of submissive begging vocalizations. The number of submissions were

compared between the vehicle control and V1aRA *low-dose* and *high-dose* groups. Dom (left panel)/Sub (right

panel) refer to initiation of dominants and subordinates. (A) Effects of V1aRA in affiliated dyads. Submissive

begging calls initiated by the dominant (DOM, left panel) did not change in any conditions; however, those initiated

by the subordinates (SUB, right panel) increased in the V1aRA *high-dose* condition but not in the *low-dose* condition.

(B) No effects were observed in any condition in both the dominants and the subordinates of unaffiliated dyads.

Plots show the individual data at each trial. In each triad, circles represent birds affiliated with another male, while

triangles represent those with no affiliation. The top, middle, and bottom of the boxes depict the 75, 50, and 25

percentiles, respectively, with whiskers representing the maximum and minimum values. Numbers in the brackets

below the boxes denote the number of samples (no. of birds × no. of trials). Asterisks indicate significant differences

(Wald χ^2 statistics, *** $p < 0.001$, ** $p < 0.01$). Colours correspond to the affiliated dyads in Figure 1.

Figure 1.

Figure 2.

Figure 3.

Figure 4.

Figure 5.

Table 1. Individuals used in the experiment 1 and 2

| | Triads used in Experiment 1 | | | Dyads used in experiment 2 | | | | | | | |
| | Dominace rank | | | Affiliated | Drug conditions received | | | Unaffiliated | Drug conditions received | | |
Triad No.	High	Middle	Low		Saline	V1aRA low-dose	V1aRA high-dose		Saline	V1aRA. low-dose	V1aRA high-dose
1	OW	PY	VV	PY, VV	✓	✓	✓	OW, VV	✓	✓	✓
2	RR	OW	BB	OW, BB	✓	✓	✓	RR, BB	✓	✓	✓
3	#28	YA	#27	#28, YA	✓	n.a.	✓	#28, #27	✓	n.a.	✓
4	VV	BB	#28	VV, BB	✓	✓	✓	BB, #28	✓	n.a.	✓
5	KR	KY	OBO	KY, OBO	✓	n.a.	✓	KR, KY	✓	n.a.	✓
6	RM	YA	VA	(RM, YA)	−	−	−	−	−	−	−
7	GK	KG	RM	−	−	−	−	−	−	−	−
8	KV	KR	LK	−	−	−	−	−	−	−	−

Birds OW, YA, VV, BB, #28, KR included in two different triads. Trio #6 (RM, YA, VA) did not receive the experiment 2 (see text for the details). *n.a.* indicates data is not available.

Table 2. Outputs of GLMMs for allopreening behaviours in affiliated dyads.

variables	estimates	s.e.	z-value	p
intercept	0.854	0.275	3.108	*<0.001*
Dominance[a]	0.325	0.257	1.266	0.206
Drug condition[b]				
Vehicle	1.639	0.217	7.554	*<0.001*
V1aRA high-dose	-0.724	0.313	-2.309	*0.021*
Dominance × Vehicle	0.068	0.281	0.241	0.811

Italicized indicates statistically significant results.

[a] *Subordinate* was set to 0.

[b] V1aRA low-dose was set to 0.

Table 3. Outputs of GLMMs for aggressive behaviours in affiliated dyads.

variables	estimates	s.e.	z-value	p
intercept	0.094	0.293	0.319	0.749
Drug condition[a]				
Vehicle	-0.193	0.315	-0.612	0.540
V1aRA high-dose	0.791	0.251	3.147	*<0.01*

Italicized indicates statistically significant results.

[a] V1aRA low-dose was set to 0.

Table 4. Outputs of GLMMs for aggressive behaviours in unaffiliated dyads.

variables	estimates	s.e.	z-value	p
intercept	0.080	0.277	0.289	0.773
Drug condition[a]				
Vehicle	-0.031	0.353	-0.089	0.929
V1aRA high-dose	0.045	0.326	0.138	0.890

Italicized indicates statistically significant results.

[a] V1aRA low-dose was set to 0.

Table 5. Outputs of GLMMs for submissive vocalizations in affiliated dyads.

variables	estimates	s.e.	z-value	p
intercept	-0.083	0.273	-0.303	0.762
Drug condition[a]				
Vehicle	0.237	0.318	0.747	0.455
V1aRA high-dose	1.929	0.253	7.617	*<0.001*

Italicized indicates statistically significant results.

[a] V1aRA low-dose was set to 0.

Table 6. Outputs of GLMMs for submissive vocalizations in unaffiliated dyads.

variables	estimates	s.e.	z-value	p
intercept	0.000	0.289	0.000	1.000
Drug condition[a]				
Vehicle	0.095	0.359	0.266	0.791
V1aRA high-dose	0.095	0.337	0.283	0.777

Italicized indicates statistically significant results.

[a] V1aRA. low-dose was set to 0.