Title: Basic and translational science of cyst wall proteins of the eye pathogen *Acanthamoeba* (87 characters +

spaces)

Short title: *Acanthamoeba* cyst wall proteins (32 characters + spaces)

Bharath Kanakapura Sundararaj[1], Manish Goyal[1], and John Samuelson[1]*

[1] Department of Molecular and Cell Biology, Boston University Goldman School of Dental Medicine, Boston,

Massachusetts, United States of America

*jsamuels@bu.edu

Abstract

The cyst wall of *Acanthamoeba castellanii* (Ac) contains cellulose and has outer ectocyst and inner endocyst layers connected by conical ostioles. Previously, we used mass spectrometry of purified Ac walls to identify families of abundant lectins (Jonah, Luke, and Leo). With the goal of identifying and characterizing novel target proteins for the diagnosis of cysts in patients with *Acanthamoeba* keratitis (AK), we localized three additional GFP-tagged proteins (a multicopper oxidase (AKA laccase) and additional Jonah and Leo lectins) to the ectocyst layer of the Ac wall. Although all walls proteins are expressed only during encystation, the precise timing of expression plays a decisive factor in their subcellular localization. Promoter swaps experiments showed early expression causes proteins to localize to the ectocyst layer, while later expression causes proteins to localize to the endocyst layer and ostioles. Ac wall proteins originate from bacteria by horizontal gene transfer (β-helical folds of Jonah and three cupredoxin-like domains of the laccase), share a common ancestry with wall proteins of slime molds (β-jelly-roll folds of Luke), or are unique to *Acanthamoebae* (four disulfide knots of Leo). β-jelly-roll folds of Luke and disulfide knots of Leo both share a linear array of aromatic amino acids, which are necessary for binding cellulose and proper localization of these proteins in the cyst wall, a feature consistent with convergent evolution. Purified rabbit antibodies to recombinant proteins bound to the ectocyst layer (Jonah and laccase) and/or to the ostioles (Luke and Leo) of cysts of 10 of 11 *Acanthamoeba* isolates tested, including five T4 genotypes that cause most cases of AK. In summary, we combined basic science experiments to understand better the localization, origin, and structure of Ac cyst wall proteins with translational studies to test target proteins for diagnosis of cysts in eye infections.

Author summary

Acanthamoebae are free-living amoeba in the soil and water that cause *Acanthamoeba* keratitis in under-

resourced countries, where water for washing hands may be scarce. *Acanthamoeba* is an emerging pathogen

in the United States, because of its association with contact lens use. While cysts are the primary target for

diagnosing *Acanthamoeba* keratitis by microscopy of corneal scrapings, there are presently no diagnostic

antibodies to characterize cyst wall proteins. Here we show that two Jonah lectins, a Leo lectin with a large

spacer, and a laccase are present in the outer layer of the *Acanthamoeba* cyst wall, while Luke lectin and Leo

without a spacer are present in the inner layer and the conical ostioles that connect the layers. We used structural

predictions to identify the aromatic amino acids of Luke and Leo necessary for binding cellulose in the wall and

to identify domains of Jonah and laccase useful for making recombinant proteins to immunize rabbits. Rabbit

antibodies to Jonah, Luke, Leo, and laccase efficiently detected cysts of ten of 11 *Acanthamoeba* isolates,

including five T4 genotypes that cause most keratitis cases. We conclude that four abundant wall proteins are

each good targets for diagnostic anti-cyst antibodies.

Introduction

Acanthamoeba keratitis (AK), which leads to scarring and blindness, if not successfully treated, is caused by free-living amoebae, named for acanthopods (spikes) on the surface of trophozoites [1-3]. In immunocompromised patients, Acanthamoebae may cause encephalitis, which is frequently fatal [4]. AK is associated with corneal trauma in the Middle East, South Asia, and Africa, where water for handwashing is often scarce [5-9]. AK is associated with contact lens use in the US, Europe, and Australia and so Acanthamoeba is on the NIAID list of Emerging Infectious Diseases/Pathogens [3, 10-12]. Regardless of the place or the cause of infection, 18S RNA sequences show that the T4 genotype most often causes AK, while other genotypes less frequently or never cause AK [13-16]. The whole genome of the Neff strain of Acanthamoeba castellanii (Ac), a T4 genotype that is broadly studied, was sequenced, and proteins were predicted and deposited in AmoebaDB [17, 18]. Recently, the Neff genome has been re-sequenced with long reads to produce 33 chromosome-size assemblies, while protein prediction has been improved by transcriptomes of trophozoites and organisms encysting for eight hours [19, 20].

Cyst walls, which form when trophozoites are starved of nutrients, protect free-living Acanthamoebae from osmotic shock in freshwater or drying in the air [21, 22]. The cyst wall also makes Acanthamoebae resistant to disinfectants used to clean surfaces, alcohol-based hand sanitizers, sterilizing agents in contact lens solutions, and/or antibiotics applied to the eye [23-27]. While trophozoites cause damage to corneal epithelial cells, AK is most often diagnosed by microscopic identification of cysts in corneal scrapings [1, 2, 28]. Acanthamoeba cysts are detected using calcofluor white (CFW) or wheat germ agglutinin (WGA), which bind to cellulose and chitin, respectively, in their walls [29, 30]. Because CFW and WGA also react with walls of plants, fungi, and other protists, they are suboptimal reagents that may lead to incorrect diagnosis of AK [31-33]. The translational goal of the present study, therefore, is to identify abundant cyst wall proteins, which might be targets for diagnostic antibodies. Antibodies to the Acanthamoeba wall might also be useful to detect cysts in concentrated water samples, monitor contamination in contact lens solutions, test reagents for cleaning reusable contact lenses, and/or test drugs that inhibit encystation.

Nearly 50 years ago, cellulose (β-1,4-linked glucose) was identified in Acanthamoeba cyst walls, which has an outer ectocyst layer and an inner endocyst layer connected by conical ostioles [34, 35]. Because

Acanthamoeba has a chitin synthase and WGA binds to cyst walls, it is also likely that chitin (β-1,4-linked

GlcNAc) is also present [36]. No cyst wall proteins were known until we purified cyst walls from the Neff strain

of *Acanthamoeba castellanii* (Ac), which is a T4 genotype, and identified three families of cellulose-binding

lectins that we named Jonah, Luke, and Leo [29]. When a representative protein from each family was tagged

with green fluorescent protein (GFP) and expressed under its own promoter in transfected Ac, Jonah localized

to the ectocyst layer that is made early, while Luke and Leo localized to the endocyst layer and ostioles that

are made later [37, 38]. Anti-GFP antibodies showed Jonah-GFP is more accessible than Luke-GFP and Leo-

GFP, suggesting Jonah and other ectocyst proteins might be better targets for anti-cyst antibodies to diagnosis

AK by fluorescence microscopy of eye scrapings [29].

Here we performed basic science experiments to understand better the localization, origin, and

structure of Jonah, Luke, and Leo lectins, as well as an abundant laccase, as well as translational science

experiments to test their value as targets for antibodies to diagnose cysts in patients with AK. First, we

identified three additional Ac proteins in the ectocyst layer and used RT-PCR and promoter swaps to begin to

understand why proteins are located there versus the endocyst layer and ostioles. Second, we used AlphaFold

and Foldseek to better understand the structure and origin of wall proteins [39-41], and we made alanine

mutations to show linear arrays aromatic amino acids of Luke and Leo bind cellulose [42-47]. Third, we tested

rabbit antibodies (rAbs) to recombinant Ac cyst wall proteins against cysts of 10 other *Acanthamoeba* strains

and species, including five T4 genotypes that cause most cases of AK [13-16].

Results

Localization of Jonah-3, Leo-S, and laccase-1 to the ectocyst layer of mature walls of Ac

The goal here was to identify additional abundant Ac cyst wall proteins that localize to the ectocyst layer, which

is likely more accessible to diagnostic antibodies than the endocyst layer. Jonah family lectins are highly

abundant in cyst walls and have an N-terminal signal peptide followed by one or three choice-of-anchor A (CAA)

domains, which form β-helical folds (BHFs) (Fig 1A). Jonah-1 (ACA1_164810), which has a long, unstructured

Thr-rich domain in addition to a single BHF, is expressed early during encystation and localizes to the ectocyst

layer when tagged with GFP and expressed under its own promoter in transformed Ac of the Neff strain (ATCC

30010) (Fig 1A, 1B, and 1E) (Supporting Information S1 and S2) [29, 37-41]. The ectocyst layer is labeled with

WGA, which binds to chitin, while the endocyst layer and ostioles are labeled with CFW, which binds to cellulose

and/or chitin. Here we tested the localization of Jonah-3 (ACA1_157320), which contains three BHFs and was

very abundant in purified cyst walls [17, 18, 29]. While other cyst wall proteins contain a signal peptide but no

TMHs, Jonah-3-AmoebaDB contains 12 transmembrane helices (TMHs) (Fig 1A) [48, 49]. GFP-tagged Jonah-

3-AmoebaDB, which was expressed under its own promoter, localizes to punctate rings around each ostiole (Fig

1B and 1C). To test whether this ring-like localization of Jonah-3-AmoebaDB might be an artifact of incorrect

protein prediction, we used an unfinished transcriptome of encysting Ac to make Jonah-3(c) (corrected), which

contains three BHFs separated by unstructured, Ser-rich spacers (Fig 1A and 1F and Supporting Information

S1). An alignment of Jonah-3(c) and Jonah-3-AmoebaDB shows 12 TMHs in the latter derive from a series of

frameshifts rather than deletions or insertions. Further, Jonah-3(c)-GFP localizes to the ectocyst layer with an

identical appearance to that of Jonah-1-GFP (Fig 1D).

Luke-2 (ACA1_377670) and Luke-3 (ACA1_245650), which have two or three β-jelly-roll folds (BJRFs),

respectively, that are separated by unstructured, Ser-rich spacer(s), are expressed later during encystation and

localize to the endocyst layer and ostioles when tagged with GFP and expressed under their own promoters (Fig

2A, 2B, 2D, and 2E). Leo-A (ACA1_074730), which contains two adjacent sets of four disulfide knots (4DKs), is

also expressed later in encystation and localizes to the endocyst layer and ostioles when tagged with GFP (Fig

3A, 3B, and 3G). Here we tested the localization of Leo-S lectin (ACA1_188350), which contains two sets of

4DKs separated by a long, unstructured, Thr-rich spacer (Figs 3A and 3I). We used for a second time the

unfinished Ac transcriptome to make Leo-S(c) that corrects an eight amino acid deletion in the N-terminal 4DK,

which was identified by an alignment with other Leo-S lectins (Supporting Information file S1). Unlike Leo-A-

GFP, which localizes to the endocyst layer and forms a flat ring around the ostioles, Leo-S(c)-GFP localizes to

the ectocyst layer of mature cyst walls in a pattern like those of Jonah-1 and Jonah-3(c) (Fig 3F). Time course

studies of Leo-S(c)-GFP during the encystation showed that after 12 hours encystation, Leo-S(c)-GFP is present

in a dense set of secretory vesicles (SV in Fig 3D) that fill the cytosol of cells, which are rounded but lack a wall

(as shown by a failure to label with WGA or CFW). However, after 24 hours encystation, Leo-S(c)-GFP forms a

patchy distribution on a single-layered wall, which is labeled with both WGA and CFW (Fig 3E). After 48 hours,

Leo-S(c)-GFP has a homogeneous distribution in the ectocyst layer (data not shown), which is the same as Leo-S(c)-GFP in mature cysts at 72 hours (Fig 3F).

The last protein tested was laccase-1 (ACA1_068450), which is a multicopper oxidase with three cupredoxin-like domains (CuRO-1, CuRO-2 and CuRO-3) that is abundant in purified cyst walls of Ac (Figs. 4A and 4E) [29, 50, 51]. Laccase-2(c) (ACA1_006180), the sequence of which we corrected using the unfinished transcriptome, is much less abundant and so was not localized. GFP-tagged laccase-1 under its own promoter localizes to the ectocyst layer of mature Ac walls with an appearance like those of GFP-tagged Jonah-1, Jonah-3(c), and Leo-S(c) (Fig 4D). Like Leo-S(c)-GFP, laccase-1-GFP fills secretory vesicles of Ac encysting for 18 hours, when there is no wall labeled with WGA or CFW (Fig 4B). N.B. that the precise timing of secretory vesicles and wall formation may vary by up to 6 hours, depending on the exact condition of trophozoites prior to placement in encystation medium. Unlike Leo-S(c)-GFP, after 24 hours encystation laccase-1-GFP is homogeneously distributed in the single layered wall of Ac, which becomes the ectocyst layer of mature cysts at 72 hours (Fig 4C and 4D).

In summary, we localized three new proteins to the ectocyst layer of the Ac cyst wall, one of which was correctly predicted by AmoebaDB (laccase-1) and two of which needed to be corrected using an unfinished transcriptome (Jonah-3(c) and Leo-S(c)). We also showed that Leo-S(c) and the laccase-1 are each made early during encystation in a dense set of secretory vesicles, which are released onto the single-layered wall in a fashion like that of Jonah-1 [29].

RT-PCR and promoter swaps show timing of expression does not just correlate with but causes protein localization in the ectocyst layer (earlier) or the endocyst layer and ostioles (later)

The first goal here was to test the hypothesis that early expression does not just correlate with but causes Ac cyst wall proteins to localize to the ectocyst layer, while later expression causes cyst wall proteins to localize to the endocyst layer and ostioles [29]. The second goal was to determine whether proteins expressed under the same promoter have the similar or different localizations in the Ac cyst wall, which would suggest similar or different binding specificity, respectively, for cellulose and/or chitin, the two glycopolymers that form fibers in the wall [35, 36]. For this, we performed qRT-PCR analysis to first determine the expression profile (timing of expression) of individual lectins during encystation. RT-PCR with mRNA isolated from trophozoites and Neff

strain encysting for 96 hours, confirmed that Jonah-1 and laccase-1 are transcribed earlier during encystation, while Luke-2 and Leo-A are transcribed later (Fig 5A). Note here that qRT-PCR products were normalized by calreticulin, which is part of the *N*-glycan-dependent quality control of protein folding in the ER and is highly expressed in trophozoites and encysting Ac [52]. The qRT-PCR results agree with those of the unfinished transcriptome, which also suggests the low level of Leo-A is likely caused by a suboptimal choice of primers for RT-PCR. Next, we performed the promoter swap experiments, in which we used GFP-tagged proteins that were either made earlier and localized to the ectocyst layer (Jonah-1 and laccase-1), and/or made later and localize to the endocyst layer and ostioles (Luke-2 and Leo-A) (Fig 1B, 2B, 3D to 3F, and 4B to 4D) [29].

In the first promoter swap experiment, we expressed Jonah-1-GFP (earlier) and Luke-2-GFP (later) each under its own promoter or under the other promoter and then determined their localizations in mature Ac cyst walls [29, 37, 38]. Jonah-1-GFP expressed under the later Luke-2 promoter moves from the ectocyst layer to the ostioles, which are densely coated and increase in number from ~9 to as many as 16, while the endocyst layer is weakly labeled (Fig 5B and 5C). The localization of Jonah-1-GFP under Luke-2 promoter is distinct from those of GFP-tagged Luke-2 and Leo-A under their own promoters, which show much greater localization to the endocyst layer, suggesting Jonah-1 binds to different glycopolymers (Fig 1B and 2B). Conversely, Luke-2-GFP expressed under the earlier Jonah-1 promoter moves from the endocyst layer and ostioles to lightly decorate the ectocyst layer and form flat, narrow rings around the ostioles (Fig 5D and 5E). These narrow rings, which match those of Luke-2-GFP under its own promoter, suggest that the glycopolymer in the ostiole to which Luke-2 is binding is available both earlier and later during encystation. The localization of Luke-2-GFP in the ectocyst layer is distinct from those of GFP-tagged Jonah-1, Jonah-3(c), Leo-S(c), or laccase-1, which, under their own promoters, coat the ectocyst layer and form dimples where ostioles are located (Fig 1B, 1D, 3F, and 4D). This result suggests that glycopolymers bound by Luke-2 are likely not the same as those bound by proteins that normally localize to the ectocyst layer.

In the second promoter swap experiment, we expressed laccase-1-GFP (earlier) and Leo-A-GFP (later) each under its own promoter or the other promoter and determined their localizations in mature cyst walls. Laccase-1-GFP expressed under the later Leo-A promoter moves from the ectocyst layer to the ostioles, which are densely coated (Fig 5F and 5G). Leo-A-GFP expressed under the earlier laccase-1 promoter, moves from

the endocyst layer and ostioles to decorate lightly the ectocyst layer and form flat rings around the ostioles (Fig

5H and 5I). Of note, the locations of Jonah-1-GFP and laccase-1-GFP under both promoters resemble each

other but are distinct from locations of Luke-2-GFP and Leo-A-GFP under both promoters.

In summary, qRT-PCR support previous conclusions based upon GFP-tagged proteins that Jonah-1 and

laccase-1 in the ectocyst layer proteins are made earlier, while Luke-2 and Leo-A in the endocyst layer and

ostioles are made later [29]. Promoter swaps support our hypothesis that earlier expression does not just

correlate with but causes proteins to localize to the ectocyst layer and later expression causes proteins to localize

to the endocyst layer and ostioles. Promotor swaps also show that all four wall proteins can locate to ectocyst

and endocyst layers and ostioles, which has relevance to complicated patterns of binding of rAbs to Luke-2 and

Leo-A to cyst walls (see below). Finally, Jonah-1 and laccase-1 appear to bind to different glycopolymers than

Luke-2 and Leo-A.

Structures of abundant wall proteins reveal the complex ancestry of Ac cyst wall proteins and identify

linear arrays of three aromatic amino acids in Luke-2 and Leo-A, which bind cellulose and direct proteins

to the Ac cyst wall

Previously, we used sequence-based searches to show that Jonah lectins contain one or three choice of anchor-

A (CAA) domains like a spore coat protein of *Bacillus anthracis*, while Luke lectins contain two or three domains

similar to those of cellulose-binding proteins of *Dictyostelium discoideum* to carbohydrate-binding modules of

bacteria (CBM2) and plant (CBM49) endocellulases [29, 32, 42-45, 53-56]. Further, the Ac laccase-1 has three

cupredoxin-like domains like those of bacterial and fungal enzymes, while 8-Cys domains of Leo appear to be

unique to Ac [50, 51]. Here we used AlphaFold and Foldseek to 1) predict structures of abundant cyst wall

proteins and compare them to proteins with proven structures, 2) identify cellulose-binding sites for Luke and

Leo lectins, and 3) select domains in Jonah-1 and laccase-1 for immunizing rabbits to produce antibodies that

detect cysts by immunofluorescence microscopy [39-41].

Luke-2 and Luke-3 have two and three β-jelly-roll folds (BJRFs), respectively, each of which has a

disulfide bond linking its beginning to its end. The BJRFs of Luke lectins are separated by short (33 to 47-aa)

unstructured domains enriched in Ser (yellow) or Thr (blue) (Fig 2D and 2E). Slimes molds including

Dictyostelium, Tieghemostelium, Heterostelium, Cavenderia, and *Polysphondylium* also have cell wall proteins

with one to five similar BJRFs, consistent their shared ancestry with Luke lectins [32, 55]. Further, the BJRFs of Luke-2 and a predicted *Dictyostelium* cellulose-binding protein (Q86KB6_DICDI) each contain a linear array of three aromatic amino acids that are also present in CBM2 of an endocellulase of a cellulolytic soil bacterium *Cellulomonas fimi,* the structure of which has been solved (PDB 1EXH) (pink arrows in Fig 2F and 2G) [57, 58]. Ala mutations of these three aromatic amino acids in CBM2 and in CBM49 of a plant endocellulase, which is closely related, show they are the binding sites for cellulose [42, 47]. Here we muted to Ala two sets of three aromatics (W35, W73, and W88 and W187, W228, and F244) of Luke-2, which was fused to maltose-binding protein (MBP) and expressed in the periplasm of *E. coli* where, like the endoplasmic reticulum of eukaryotic cells, disulfide bonds are formed [29, 59, 60]. We purified MBP-Luke-2 +/- Ala mutations on amylose resins and tested their binding to Avicel (crystalline) cellulose. Western blots showed WT Luke-2 binds well to Avicel cellulose, while MBP alone and Luke-2 plus Ala-mutations fail to bind to cellulose (Fig 2J). Further, Luke-2-GFP plus Ala mutations, which was expressed under the same promoter as WT Luke-2-GFP, no longer localizes in the endocyst layer and ostioles (Fig 2B and 2C). The binding of MBP-WT Luke-2 to Avicel cellulose argues for its proper folding, so we used it to immunize rabbits.

Leo lectins have two sets of four disulfide knots (4DKs), which are adjacent (Leo-A) or separated by a long, unstructured, Thr-rich domain (Leo-S(c)) (Fig 3G and 3I). Although there are numerous carbohydrate-binding modules composed of sets of 4DKs (e.g., CBM18 of WGA, CBM19 of a *Saccharomyces cerevisiae* GH18 chitinase, and CBM55 of *Entamoeba histolytica* GH18 chitinase), we were unable with Foldseek to identify any shared structures with the set of 4DKs of Leo [43, 46, 61-64]. Remarkably, 4DKs of Leo-A and Leo-S(c) contain linear arrays of three aromatic residues, even though these 4DKs have no relationship to BJRFs of Luke or to CBM2 and CBM49 of bacterial and plant endocellulases (Fig 3H). Ala mutations to two sets of three aromatics (Y46, Y63, and Y77 and Y134, Y151, and Y165) caused an MBP-Leo-A fusion to no longer bind to Avicel cellulose (Fig 2J). Similarly, the Ala mutant of Leo-A-GFP, which was expressed under the same promoter as WT Leo-A-GFP, no longer localizes to the endocyst layer and ostioles (Fig 3B and 3C). As above, the binding of MBP-WT Leo-A to Avicel cellulose argues for its proper folding, so we used it to immunize rabbits.

Jonah-1 has a single three-sided BHF, a pair of α-helices of unknown function, and a long, unstructured, Thr-rich domain, while Jonah-3(c) has three BHFs and three long, unstructured, Ser-rich domains (Fig 1E and

1F). The BHF of Jonah-1 closely resembles the three-sided BHF of an antifreeze protein of an Antarctic sea ice

bacterium *Colwella sp*., the structure of which has been solved (PDB 3WP9) (Figs 1E to 1H) [65]. Although

Jonah-1 and laccase-1 (next paragraph) contained numerous aromatic amino acids on their surface, none was

in linear arrays, so we did not perform Ala mutations to test cellulose-binding. Finally, we chose the BHF of

Jonah-1 to make an MBP-fusion for immunizing rabbits, because we wanted to avoid unstructured, Thr-rich

regions, which might interfere with proper folding of the protein and/or contain *O*-linked glycans that would

obscure antigenic sites.

The laccase-1 of Ac has three cupredoxin-like domains (CuRO-1, CuRO-2 and CuRO-3) but lacks

unstructured, Ser- or Thr-rich domains, which distinguishes it from other ectocyst layer proteins (Jonah-1, Jonah-

3(c), and Leo-S(c)) (Fig 4E). Instead, laccase-1 has a positively charged loop between CuRO-2 and CuRO-3,

which is also present in spore coat proteins of bacteria. Ac laccase-1 shares a 44% identity with spore coat

protein A of *Caldicoprobacter faecalis*, which is present in sewage sludge. Ac laccase-1 shows slightly lower

positional identities with laccases of archaea (e.g., 40% with *Halalkalicoccus paucihalophilus*, fungi (e.g., 38%

identity with *Penicillium nalgiovense*), and plants (e.g., 36% with *Diphasiastrum complanatum* (36%). *Pelomyxa*

schiedti (36% identity) is the only Amoebazoa with a laccase, which is absent from humans and most metazoa

but are present in a few arthropods (e.g., 28% identity with *Rotaria sp.*). While the high positional identity with

bacterial laccases suggests the Ac laccase-1 may be active, we did not determine what substrates are oxidized,

and we did not knock it out to determine its function. The CuRO-1 of laccase-1, which we chose to make an

MBP-fusion for immunizing rabbits, closely resembles that of the spore coat protein A of *Bacillus subtilis*, the

structure of which has been solved (PDB 4YVN) (Fig. 4F) [66]. A possible concern here is the presence of a

10-aa sequence (NVYAGLAGFY) near the C-terminus, which is also present in laccases of some bacteria,

archaea, fungi, and plants and so might lead to cross-reacting antibodies.

In summary, BJRFs of Luke lectins share recent common ancestry with wall proteins of slime molds and

distant ancestry with CBM2 and CBM49 of bacterial and plant endocellulases, while 4DKs of Leo are unique.

BHFs of Jonah lectins and three CuRO domains of laccase closely resemble those of bacterial proteins and so

likely derive by horizontal gene transfer, which was not proven here. Although the structures of the BJRFs of

Luke-2 and the set of 4DKs of Leo-A show no resemblance, the linear arrays of three aromatic amino acids,

which are involved in binding cellulose and localizing proteins in the endocyst layer and ostioles, are the same, consistent with convergent evolution. Finally, while the BHF of Jonah-1, BJRFs of Luke-2, and 4DKs of Leo-A have no well-conserved sequences that might lead to cross-reacting antibodies with bacteria or fungi in eye scrapings, CuRO-1 of laccase-1 has a well-conserved 10-aa sequence that may be problematic.

Rabbit antibodies to Jonah-1, Luke-2, Leo-A, and laccase-1 all efficiently detect cysts of 10 of 11 *Acanthamoeba* **isolates, including five T4 genotypes that cause most cases of AK.**

The goals here were to 1) use rAbs to recombinant wall proteins to visualize native proteins by western blots and confocal microscopy and 2) test how well each antibody detected CFW-labeled cysts of 11 *Acanthamoeba* species/strains.

Rabbits were immunized with MBP-fused to WT Luke-2, WT-Leo-A, BHF of Jonah-1, and CuRO-1 of the laccase-1 in complete Freund's adjuvant and boosted three times with incomplete Freund's adjuvant at Cocalico Biologics, Inc., and rabbit IgGs were purified using Protein-A Sepharose [59, 60]. Western blots and confocal microscopy showed that pre-bleeds from rabbits do not react with trophozoites or cysts of Ac (Fig. 6A). While rAbs to MBP-fusions with Jonah-1, Luke-2, Leo-A and laccase-1 do not bind to trophozoite proteins, each rAb binds well to cyst wall proteins of Neff strain of Ac. Anti-Jonah-1 rAbs bind to a protein of the expected size of ~55-kDa, as well as to two lower mol wt bands. The latter may result from proteolytic cleavage of Jonah-1 before or during isolation of cyst proteins and/or by cross-reaction with smaller Jonah-1 proteins, which share multiple conserved domains. Anti-Luke-2 rAbs bind to a thick ~50-kDa band, which is greater than the 27-kDa expected size of Luke-2. We suspect that the increased size of Luke-2 is caused by extensive glycosylation of its five *N*-glycan sites and ~20 *O*-glycan sites in the low complexity, Ser-rich spacer [67]. Anti-Leo-A rAbs bind to a thick ~13-kDa band, which is slightly smaller than the expected size of 17-kDa, perhaps due to proteolytic cleavage. Finally, anti-laccase-1 rAbs bind to a 75-kDa band, which is slightly bigger than the expected size of 64-kDa, as well as to a less abundant 55-kDa band. Again, addition of *N*-glycans or *O*-glycans may explain the increase in size, while proteolytic cleavage may explain the lower mol wt band.

We tested rAbs versus the model Neff strain of Ac, as well as *Acanthamoeba* species/strains from the ATCC collected by Monica Crary, who studies their binding to contact lenses [24, 27]. Segments of 18S RNA genes were amplified from trophozoites, cloned, and sequenced to confirm the genotype of 11 *Acanthamoeba*

species/strains prior to encystation and testing with rAbs [13-16]. Four T4 genotypes from the ATCC included *A. castellanii* Neff stain (30010), MA strain (50370), and unnamed strain (50324) and *A. polyphaga* (30872), while a fifth T4 (MEEI 0184) and another AK isolate (Esbc4) came from Noorjahan Panjwani, who discovered the mannose-binding protein on the surface of trophozoites [68]. Other genotypes from ATCC included T7 (*A. astronyxis* 30137), T11 (*A. hatchetti* PRA-115) and T18 (*A. byersi* PRA-411 and *Acanthamoeba sp. 13* 50655).

High-power confocal microscopy showed rAbs to the BHF of Jonah-1 bound in a somewhat patchy distribution to the ectocyst layer (surface) of cysts of Neff strain and 10 other *Acanthamoeba* species and/or strains (Fig 6A and 7A). This patchy distribution, which is in contrast to the homogeneous distribution of Jonah-1-GFP (Fig 1B), may result from masking of the BHF of Jonah-1 by its large, unstructured Thr-rich domain, which likely contains *O*-linked glycans, and/or masking by other ectocyst wall proteins (Fig 1E). In contrast, rAbs to CuRO-1 of laccase-1 bound in a homogenous pattern to the Neff strain and to other *Acanthamoeba* species/strains, which is similar to that of laccase-1-GFP, suggesting there is less masking of laccase-1 (Fig 4D, 6E, and 7D). While rAbs to Luke-2 and Leo-A densely labeled the ostioles and weakly labeled the endocyst layer of many cysts (Figs. 6C and 6D), other cysts were labeled on the endocyst layer and/or ectocyst layer (Fig. 7B and 7C). Of note, rAbs to Luke-2 and Leo-A often bound in similar patterns to a particular *Acanthamoeba* isolate, which is reminiscent of similar appearances of Luke-2-GFP and Leo-A-GFP in the promoter swap experiments (Fig 5). There are two explanations, which are not mutually exclusive, for the variable distribution in the cyst walls of rAbs to Luke-2 and Leo-A. First, these proteins may be made at different times by different *Acanthamoeba* species/strains, so that Luke-2 and Leo-A localize to different places. Second, these rAbs may bind to relatively small amounts of Luke-2 and Leo-A in the ectocyst layer, which was not visualized by confocal microscopy, because of the large amounts of GFP-tagged Luke-2 and Leo-A in the endocyst layer and ostioles (Fig 2B and 3B).

To determine which wall proteins might be the best targets for diagnosis of cysts in AK, we counted in randomly selected low power confocal fields the number of CFW-labeled cysts detected by rAbs to Jonah-1, Luke-2, Leo-A, and laccase-1 (Fig 6F to 6I). In two separate experiments, >100 cysts were counted for each rAb and each *Acanthamoeba* species/strain, and the averages +/- SEM were plotted in Fig 6J. We found that rAbs to Luke-2, Leo-A, and laccase-1 each detected >95% of CFW-labeled cysts of 10 of 11 *Acanthamoeba* species

and strains tested. In contrast, anti-Jonah-1 rAbs showed >95% detection for 7 of 11 *Acanthamoeba* species but detected somewhat fewer cysts of *A. byersi* (PRA-115) (T11) (91%), *Acanthamoeba* sp. 13 (ATCC 50655) (T18) (89%), and *A. polyphaga* (ATCC 30872) (T4) (85%). Why this is the case is not clear, but the result, which suggests that the BHF of Jonah-1 might not be quite as good a target as the BJRFs of Luke-2, the 4DKs of Leo-A, or CuRO-1 of laccase-1, was reproduced in separate experiments. In addition, all four rAbs struggled to detect *A. mauritaniensis* (T4) (ATCC 50676), suggesting that there is a problem with encysting these trophozoites under the conditions used here. While counts were performed for two sets of cysts labeled with Protein-A Sepharose-purified antibodies, we obtained similar results with rabbit sera diluted 1:300. Most important, we confirmed that rAbs to all four proteins were also visible with a conventional Zeiss fluorescence microscope, which is more similar to those present in clinical labs or in ophthalmologists' offices. While the success with rAbs to Jonah-1 and laccase-1 was expected, because these proteins are on the ectocyst layer (surface) of cysts, the success of rAbs to Luke-2 and Leo-A was unexpected because we previously had trouble locating Luke-2-GFP or Leo-A-GFP using anti-GFP antibodies [29]. Finally, regardless of whether the rAbs are binding to the ectocyst layer or endocyst layer, the presence of ~10 ostioles makes *Acanthamoeba* cysts look distinct from fungal walls, which may at most have a few bud scars [29, 31, 34].

Discussion

Incorrect protein prediction of Jonah-3 in AmoebaDB led to its art factual but interesting localization in rings around ostioles

While Ac proteins sequences available in AmoebaDB made it possible for us to perform mass spectrometry of cyst walls and for others to discover dozens of proteins of interest, its predictions are based upon an incomplete genome of Neff strain of Ac and algorithms that identify exons in genes with a high density of introns rather than transcriptomic data [17, 18, 29]. We used an unfinished transcriptome to show TMHs between three BHFs, which incorrectly localized GFP-tagged Jonah-3-AmoebaDB to a ring around ostioles, were caused by frameshifts. Instead, Jonah-3(c) has low-complexity, Ser-rich spacers like those of Luke-2 and localizes to the ectocyst layer in a pattern matching that of Jonah-1. The unfinished transcriptome also allowed us to correct an eight amino acid long deletion in Leo-S-AmoebaDB that would have disabled its N-terminal 4DK, which is necessary for binding cellulose and localizing the protein to the ectocyst layer. Because AmoebaDB predicts

337 14,000 proteins, the errors described here in Jonah-3, Leo-S, and laccase-2 are mere anecdotes but suggest a

338 need for a reannotated proteome based upon a well-constructed transcriptome. Finally, despite being an artifact,

339 rings around the ostioles formed by Jonah-3-AmoebaDB may be useful for studying ostiole formation during

340 encystation and ostiole breakdown during encystation. Similarly, the heavy labeling of ostioles by Jonah-1 and

341 laccase-1 under the later Luke-2 and Leo-A promoters, respectively, may also facilitate studies of ostiole

342 formation and degradation.

343 **Timing of expression and properties of abundant wall proteins determine their localization in the**

344 **ectocyst layer, endocyst layer, and/or ostioles**

345 Because whole cyst walls were purified by Percoll gradients and examined by mass spectrometry, we expressed

346 selected proteins with a GFP-tag under their own promoters to precisely localize them in the two layered-wall

347 connected by ostioles [29, 37, 38]. Here we increased the number of ectocyst layer proteins from one (Jonah-1)

348 to three (Jonah-3(c), Leo-S(c), and laccase-1. Laccase-1 and Leo-S(c) were also localized to a massive set of

349 secretory vesicles early in encystation, as was previously shown for Jonah-1. Rabbit antibodies confirmed the

350 presence of Jonah-1 and laccase-1 in the ectocyst layer of untransformed cysts of the model Neff strain of Ac,

351 as well as cysts of 10 other *Acanthamoeba* species/strains. Promoter swaps showed Jonah-1 and laccase-1

352 switch to the endocyst layer and ostioles when expressed under later promoters of Luke-2 and Leo-A,

353 respectively. Conversely, Luke-2 and Leo-A switch from the endocyst layer and ostioles to the ectocyst layer

354 when expressed under the early promoters of Jonah-1 and laccase-1, respectively. Because the localizations of

355 Jonah-1 and laccase-1 resemble each other under early and later promoters, it is likely that they are binding to

356 the same glycopolymer(s), which we did not determine here. Similar localizations of Luke-2 and Leo-A under

357 both promoters suggest these lectins are also binding to the same glycopolymer(s). This result is not surprising,

358 as Luke-2 and Leo-A contain similar arrays of aromatic amino acids that bind cellulose.

359 Our conclusion that early expression causes proteins to localize in the ectocyst layer, and later expression

360 causes proteins to localize in the endocyst layer and ostioles leaves many questions unanswered. As above, we

361 do not know how ostioles are formed, and we have not yet precisely localized cellulose and chitin in cyst walls.

362 Further, we have not identified sequences in regions 5' to the start ATG that bind transcription factors specific

for earlier or later encystation-specific expression, nor have we figured out why some encystation-specific

promoters are stronger than others and so produce greater quantities of transcripts for wall proteins.

While its 4DKs are unique, Leo lectins share many properties with wall proteins of Ac and other

organisms, arguing for convergent evolution

Although the four families of Ac wall proteins show no relationship to each other, three have been previously

identified and characterized in walls of protists, fungi, plants, and bacteria. Luke lectins have two or three BJRFs,

which share recent ancestry with wall proteins of *Dictyostelium* and distant ancestry with CBM2 and CBM49 of

bacterial and plant endocellulases [32, 42, 47, 55, 58]. Laccases, which are most similar to those of bacteria,

are widely distributed in walls of plants, fungi, and oomycetes [50, 51, 66]. Although their function in Ac, which

proliferate in temperate climates, is not clear, BHFs similar to those in Jonah lectins protect Artic bacteria from

freezing [56, 65]. While its 4DKs that bind cellulose are unique, Leo lectins share properties with wall proteins of

Ac and other organisms, suggesting the importance of convergent evolution [69]. Chitin-binding domains of

Saccharomyces chitinases (CBM19), *Entamoeba* chitinases (CBM55), and WGA (CBM18) also contain unique

4DKs, while chitin-binding domains of Jacob lectins contain unique 3DKs [43, 46, 62-64]. Even though 4DKs of

Leo and BJRFs of Luke have no structural similarity, each contains linear arrays of three aromatic amino acids,

which Ala mutations showed bind cellulose. Similar linear arrays of aromatics that bind cellulose are present in

endocellulases of bacteria (CBM2) and plants (CBM49), as well as in CBM63s of expansins (proteins that unfold

cellulose) of plants, bacteria, and Ac [42, 44, 47]. Reinvention by convergent evolution of these linear arrays of

aromatics strongly suggests they are the best means for binding cellulose, which forms flat ribbons with

alternating glucose residues facing opposite surfaces.

Like Leo-A with two adjacent 4DKs and Leo-S with 4DKs separated by a large Thr-rich spacer,

Entamoeba Jacob-1 has two adjacent 3DKs, while Jacob-2 has a third 3DK separated by a long, unstructured,

Ser-rich spacer [70]. Unstructured Ser- or Thr-rich domains, which are also present in Jonah and Luke lectins,

are likely modified by *O*-linked glycans, which protect glycoproteins from being degraded by bacterial proteases,

as shown for wall proteins of *Entamoeba* and *Cryptosporidium* [71, 72]. Due to constraints on resources and

time, we did not identify *O*-linked glycans on Ac cyst wall proteins, nor did we confirm the AlphaFold structures

by crystallizing any of the four wall proteins studied here. This seems less of an issue for the BHFs of Jonah-1

and -3, BJRFs of Luke-2 and Luke-3, and CuRO-1 domain of laccase-1, the structures of which match those of crystallized proteins [58, 65, 66]. A goal of future studies will be to solve the crystal structure of the unique 4DK of Leo +/- cellulose.

Abundant wall proteins do not have to be in the ectocyst layer to be good targets for diagnostic antibodies

While it is rare for us (and other parasitologists) to pair basic and translational science experiments, it worked well here for the following reasons. The BHF of Jonah-1, the BJRFs of Luke-2, the 4DKs of Leo-A, and CuRO-1 domain of laccase-1 each expressed well as MBP-fusions in the periplasm of bacteria, where disulfide bonds are formed [59, 60]. In contrast to anti-peptide rAbs to Jonah-1 and Leo-A, which bound to Western blots but not to fixed cysts [29], rAbs to MBP-fusions of all four Ac wall proteins bound well to Western blots and to cysts of Neff strain of Ac, as well as to cysts of 9 of 10 other isolates of *Acanthamoeba*. While we expected that rAbs to Jonah-1 and laccase-1, which are both in the accessible ectocyst layer, to efficiently detect CFW-labeled cysts, anti-laccase-1 performed slightly better than anti-Jonah-1. Even though we doubted whether rAbs to Luke-2 and Leo-A, which are in the relatively inaccessible endocyst layer and ostioles, would detect CFW-labeled cysts efficiently, anti-Luke-2 and anti-Leo-A rAbs both performed very well. Although the labeling of the ectocyst layer by rAbs to Luke-2 and Leo-A is not easy to explain, efficient detection is what matters for a diagnostic reagent. Finally, the signal from the secondary goat antibody binding to rAbs to Jonah-1, Luke-2, Leo-A, and laccase-1 is strong enough, so that cysts were easily detected with a conventional fluorescence microscope, not just by an expensive confocal microscope used to get three-dimensional, high-resolution images of the ectocyst layer, endocyst layer, and ostioles.

Because these rAbs cross-react with MBP that was not removed prior to vaccination, they are not ready for testing on corneal scrapings that likely contain numerous bacteria. Although we are confident that Luke-2, Leo-A, and Jonah-1 antigens are each unique to Ac, a 10-aa sequence in laccase-1 may lead to antibodies that cross-react with walls of bacteria, fungi, or plants. Cysts examined here were made by starving cultured *Acanthamoebae* and so might not be the same as those made in the corneal epithelium, soil, or water. Diagnostic anti-cyst antibodies will complement monoclonal antibodies to the mannose-binding proteins on trophozoites [68, 73], as well as antibodies to transporters and secreted proteins of trophozoites and cysts [74-76]. Anti-cyst

antibodies may also supplement Loop-mediated Isothermal Amplification (LAMP) assays for diagnosing AK [77,

78].

METHODS

Ethics statement

Culture and manipulation of *Acanthamoebae* under BSL-2 protocols were approved by the Boston University

Institutional Biosafety Committee. Production of custom rabbit antibodies was approved by the Institutional

Animal Care and Use Committee of Cocalico Biologics, Inc., Denver PA.

Summary of new methods to study Ac wall proteins.

Many methods are the same that we used in our mass spectrometric characterization of proteins in purified cyst

walls of Ac, which were described in detail [29]. New here is use of unfinished transcriptome to correct protein

prediction in AmoebaDB for Jonah-3(c) and Leo-S(c). Confocal microscopy replaced structured illumination

microscopy (SIM), because it is quicker and allowed us to examine many more parasites. Two pairs of promoter

swaps were used to show that localization in the two layers of the wall is not just correlated with timing of

expression but is caused by timing of expression. AlphaFold was used to predict structures for each cyst wall

protein and test with Ala mutations aromatic amino acids involved in binding cellulose. Foldseek was used to

suggest the origin of domains in cyst wall proteins that could not be identified using sequence-based searches.

Recombinant proteins, rather than peptides, were used to make rAbs to abundant wall proteins, which supported

localizations of four GFP-tagged constructs under their own promoters. Binding of the rAbs to cysts of numerous

Acanthamoeba genotypes, which were confirmed by PCR of 18S RNA genes, supported their use as targets for

diagnostic antibodies.

***Acanthamoeba* species and strains, culture, and cyst preparation**

A. castellanii Neff strain (ATCC 30010) trophozoite parasites were obtained from the American Type Culture

collection (ATCC). Trophozoites of other strains of *Acanthamoeba*, originally derived from human corneal

infections and granulomatous encephalitis infections, were acquired from Dr. Monica Crary, Alcon Research,

LLC, Fort Worth, TX, United States or from Noorjahan Panjwani of Tufts University Medical School [24, 68]. All

experiments were performed using *A. castellanii* Neff strain until otherwise mentioned.

Ac trophozoites were grown and maintained in axenic culture at 30ºC in T-75 tissue culture flasks in 10 ml ATCC medium 712 (PYG plus additives) with antibiotics (Pen-Strep) (Sigma-Aldrich Corporation, St. Louis, MO) as described previously [21, 29, 79]. Adherent log-phase trophozoites from stationary culture were detached with the help of a cell scraper and pelleted down by centrifugation at 500x g for 5 min followed by two washes with 1X phosphate buffered saline (PBS). Cysts were prepared from trophozoites by incubating them with encystation medium (EM, 20 mM Tris-HCl [pH 8.8], 100 mM KCl, 8 mM $MgSO_4$, 0.4 mM $CaCl_2$, and 1 mM $NaHCO_3$) [29, 79]. In brief, ~10^7 trophozoites obtained from a confluent flask were washed with 1x PBS and subsequently incubated with EM in a T-75 tissue culture flask for 120 hours. The mature cysts were released with the help of a cell scrapper, harvested, and washed with 1x PBS by centrifugation at 1,500x g for 10 min. The harvested cysts were either used immediately or stored at 4ºC or -20ºC depending upon the need of the experiment.

In silico sequence analysis of candidate cyst wall lectins

The full-length coding sequence of different cyst wall proteins of _A. castellanii_ namely Jonah-1 (ACA1_16481), Jonah-3 (ACA1_157320), Luke-2 (ACA1_377670), laccase-1 (ACA1_068450), Leo-A (ACA1_074730) and Leo-S (ACA1_188350) were obtained from AmoebaDB, a functional genomic database useful for genetic studies of the Neff strain and ten other _Acanthamoeba_ strains (https://amoebadb.org/amoeba/app) [80]. The AmoebaDB database was also used to predict introns and identify paralog proteins and upstream promoters (nucleotide sequence) of different cyst wall proteins. The full-length amino acid sequences of candidate wall proteins were further analyzed, reannotated, and corrected based on homologs proteins using BLAST server in other species [53]. Other _in-silico_ tools used to functionally characterize the cyst wall proteins include the CDD database (conserved domain identification) [81], SignalP 4.1 [49] and DeepTMHMM [82] (signal peptides and transmembrane helices), CAZy and InterPro databases (Carbohydrate-binding modules) [43, 46, 54] [48] and big-PI (Glycosylphosphatidylinositol anchors) [83], respectively. Finally, we used an unfinished transcriptome of trophozoites and encysting Ac, which will be described elsewhere when completed, to check and correct protein predictions.

In silico structural prediction of candidate cyst wall proteins

The 3D structure models of different cyst wall lectins were predicted by AlphaFold2 using the Colab server

(https://colab.research.google.com/github/sokrypton/ColabFold/blob/main/AlphaFold2.ipynb) [39]. The

molecular visualization of active site residues, 3D structures, and structural superimposition was performed with

the help of PyMOL Molecular Graphics System, Version 2.0 Schrödinger, LLC [84]. Additionally, the predicted

3D structures were used to perform structural comparison and similarity search with large structure sets

(AlphaFold database, PDB database, etc.) using Foldseek server (https://search.foldseek.com/search) [40].

Expression and localization of candidate cysts wall lectins during encystation in *A. castellanii*

RT-PCR studies were performed to check the expression of cyst wall lectins in encysting protists. In brief,

adherent log-phase trophozoites of Ac were washed twice with PBS and subsequently stimulated to encyst by

incubation at 30ºC with encystation media. The amoeba cells were collected at various time points including 0,

6, 12, 18, 24, 36, 48, 72, and 96 hours post incubation. The cells were pelleted down by centrifugation, washed

with PBS, and stored in Trizol reagent at -80ºC for further use. Once all the time points were collected, the Trizol

cell pellets were thawed on ice, and the cells were lysed by bead beating (BioSpec mini bead beater) operated

in a cold room at 4ºC with 1 min beating and cooling off for 5 mins for 3 cycles. Total RNA was extracted using

Directzol RNA minprep kit (Zymo Research), and the concentration was checked using Nanodrop 2000/2000c

(Thermo Fisher Scientific). cDNA was synthesized using AMV reverse transcriptase (New England Biolabs,

Ipswich, MA) as per the manufacturer instructions. The expression of abundant cyst wall proteins during

encystation was analyzed using Realtime-PCR (Biorad) using Power SYBR green PCR master mix (Applied

Biosystems). Calreticulin was used as housekeeping gene to normalize the expression profile of various

candidate cyst wall lectins. The primers used for real time PCR analysis are listed in Supporting Information S1.

To verify the expression and subcellular localization of cyst wall proteins during encystation, we

expressed individual cyst wall lectins from an episomal plasmid in Neff strain of Ac under their own promoter.

Primers for making constructs are listed in Supporting Information S1, while sequences of promoters and

proteins are listed in Supporting Information S2. For all constructs, the pGAPDH plasmid was used, which

harbors a neomycin resistance gene (for G418 drug selection) and a glyceraldehyde 3-phosphate

dehydrogenase (GAPDH) promoter for constitutive expression of a C-terminus GFP fusion chimera [37]. In our

in silico structural studies, we observed a comparable binding site topography for Luke-2 and Leo-A cellulose-

binding module (CBM), which consists of a linear array of aromatic amino acids, as previously observed phenomenon for other CBMs. To confirm the involvement of these aromatics in Luke-2 and Leo-A CBMs these aromatic amino acid residues were mutated to alanine. Tryptophan (W35, W73, W88, W187, W228 and phenylalanine (F244) of Luke-2 were replaced with alanine, while tyrosine (Y46, Y63, Y66, Y134, Y151, and Y165) of Leo-A were replaced with alanine. The construct used for Jonah-1, Leo-A (WT), and Luke-2 (WT) were taken from a previously published study [29]. While the full-length coding sequences of Leo-A mutant, Luke-2 mutant, laccase-1, and Leo-S gene were codon optimized and custom synthesized from the Twist Biosciences along with their respective promoter sequences. For the promoter swap experiment, the Luke-2 promoter was replaced with Jonah-1, while the laccase-1 promoter was replaced with the Leo-A promoter and vice versa. The 5' upstream sequence (400-600 bp) (Supporting Information S2) for each gene was PCR amplified from Ac genomic DNA (used as a promoter to drive their expression) to swap the respective gene promoter in the pGAPDH plasmid. For construct preparation, we used a restriction-free cloning strategy using NEBuilder HiFi DNA assembly master Mix from New England Biolabs, Ipswich, MA. The identities of final constructs were sequence verified using Oxford nanopore sequencing technology from Plasmidsaurus. The primers used for cloning of different promoters, and cyst wall lectin genes are listed in Supporting Information S1.

Transfection

Transfection in *A. castellanii* was performed routinely in the lab using Lipofectamine™ Transfection Reagent (Thermo Fisher Scientific), as per the manufacturer's instruction. Briefly, for each transfection ~5 x 10^5 log-phase trophozoites were seeded in a T-25 flask in ATCC medium 712 and incubated for 30 min at 30ºC. After the incubation, the adherent trophozoites were washed, and the media was replaced with 500 µl of encystation media (EM). Prior to transfection, the Lipofectamine™ 3000 reagent was diluted in EM (7.5 µL Lipofectamine™ 3000 was diluted in 125 µL EM). The master mix of DNA was prepared by diluting 8 µg of plasmid DNA with EM (to achieve a final volume of 117 µL) and subsequent addition of 8 µL P3000™ reagent. Diluted Lipofectamine™ 3000 and DNA master mix both were added in a 1:1 (125 + 125 µl) ratio and incubated at room temperature for an additional 15 min. Following incubation, the DNA-lipofectamine complex (250 µL) mixture was added directly onto the adherent trophozoites in 1 mL EM (1250 µL). The trophozoites were further incubated for an additional 4 hours at 30ºC. After this, the EM and transfection mixture was removed by pipetting and 10 ml fresh media

was added into each well. After 24h of transfection, the used medium was replaced with fresh media containing

G418 (12.5 µg/mL). After 48h of transfection, cells were transferred from 6 well plates to T-75 flasks with ATCC

medium 712 plus G418 (25 µg/mL). The used medium was changed routinely with fresh media containing G418

(25 µg/mL) on every fourth day. After 2 to 4 weeks, the transfectants began growing robustly in the presence of

G418.

Confocal microscopy

To check the expression of the abundant cyst wall proteins during the encystation, the transgenic Ac trophozoites

(which express GFP-tagged chimera cyst wall proteins) were induced to encyst with EM. The encysting amoebas

were collected at various time points (0, 6, 12, 18, 24, 36, 48, 72, and 96 hours), centrifuged, washed (1x PBS),

and fixed by incubating them with 4% paraformaldehyde (PFA) for 15 min at room temperature. Following fixation

cells were pelleted down, washed, and resuspended in PBS. The fixed cells were stained with Wheat Germ

Agglutinin 1:10 (WGA, 1 mg/ml, ThermoFisher Scientific) conjugated to Alexa Flour 647 and Calcofluor White

1:20 (CFW, 1mg/ml, Sigma Aldrich) in PBS for 30 min at room temperature. The cells were washed 3 times with

1x PBS and mounted in VECTASHIELD® Antifade Mounting Medium (Vector Laboratories, Newark, CA).

Samples were illuminated using 380 nm (CFW), 488 nm (GFP chimera), and 647 nm (WGA) laser excitation. .

The fluorescence images were captured using CFI Plan Apochromat VC 60XC WI Plan APO 60x oil objective of

Nikon Ni2 AX inverted confocal microscope equipped with NIR imaging system. We deconvolved 0.1 µm optical

sections using NIS elements (Version: AR5.41.02) imaging software. All confocal images shown were 3D

reconstructions using dozens of z-stacks. Size bars were based upon 2D cross-sections.

Overexpression and purification of candidate cyst wall proteins as MBP-fusion

The MBP-fusion constructs of Jonah-1, laccase-1, Luke-2 (WT and mutant), and Leo-A (WT and mutant) were

prepared by cloning the codon-optimized (*E.coli* expression) synthetic gene fragments into pMAL-p2x vector

(New England Biolabs) for periplasmic expression in BL21-CodonPlus(DE3)-RIPL (Agilent Technologies,

Lexington, MA) [60]. The primers used for cloning synthetic gene fragments and the length of different domains

of cyst wall proteins (devoid of signal sequence) are listed in Supporting Information S1 and S2. The expression

of each MBP-fusion proteins was induced by the addition of 0.1 mM of IPTG for 16 h at 16°C. The bacterial cell

pellet was lysed, and supernatant was collected as per the manufacturer's specifications (New England Biolabs).

AKTA start was used to purify the MBP tagged protein using an MBPtrap HP column (Cytiva). The MBPtrap

column was equilibrated with equilibration or wash buffer (20 mM Tris-HCl (pH 8), 200 mM NaCl, 1 mM EDTA).

Later the supernatant mixture was passed through the column for the MBP-tagged protein to bind to the column

at 0.5 mL/min flow rate. The column washed with 20 column volumes of wash buffer, and the purified protein

was eluted with elution buffer (20 mM Tris-HCl (pH 8), 200 mM NaCl, 1 mM EDTA, 10 mM maltose). The identity

and purity of recombinant purified MBP-fusion proteins were confirmed by SDS-PAGE and Western

immunoblotting analyses. The wild-type recombinant Jonah-1, Luke-2, Leo-A, and laccase-1 MBP-fusions were

used to raise custom polyclonal antibodies (Cocalico Biologicals). Total IgG was purified from plasma samples

of pre-immune and post-immunized rabbits via Protein A affinity chromatography (Pierce™ Protein A Agarose,

Thermo Fisher Scientific, USA) as per manufacturer instructions. In brief, for Protein A Sepharose purification,

plasma was diluted 2-fold with binding buffer (1x Tris-Buffered Saline, pH 7.4) and loaded onto a column

containing Protein A Agarose beads. The diluted plasma was passed through the column twice, and beads were

washed with 1x TBS (20-fold column volume). The bound IgG was eluted with 0.1 M glycine-HCl (pH 2.7), into

the neutralizing buffer (1 M Tris-HCl, pH 9.0) and concentrated, and the buffer exchanged into PBS using a 30

kDa Amicon Ultra centrifugal filter (Millipore, USA).

Cellulose binding assay using WT and Mutant Luke-2 and Leo-A

To check the cellulose binding activity of WT and mutant Luke-2 and Leo-A, we performed an *in vitro* cellulose

binding assay. In this assay, recombinant purified MBP fusion Luke-2 (WT and Ala mutant) and Leo-A (WT and

Ala mutant) proteins were used. A total of 1 µg MBP-fusion protein (in 100 µl of 1% NP40) was incubated with

0.5 µg Avicel microcrystalline cellulose (Sigma-Aldrich) for 3 hours at 4°C with rocking. Following binding,

microcrystalline cellulose fibers were pelleted down by centrifugation, while the supernatant was collected in a

separate tube (unbound fraction). The microcrystalline cellulose fibers (bound fractions) were washed three

times with 1% NP40. The input material (total) unbound (U), and bound (B) fractions were boiled in the SDS

sample buffer. Soluble proteins were separated on SDS-PAGE (4-15%), blotted to nitrocellulose membranes,

blocked in 5% BSA, and detected using anti-MBP antibodies (New England Biolabs). As a negative control,

microcrystalline beads were incubated with MBP alone.

Detection of candidate cyst wall lectins in *A. castellanii* trophozoites and cysts

To detect the cyst wall proteins in Ac, log-phase trophozoites parasites and 120 h post-encysting mature cysts

were harvested and lysed in SDS sample buffer. The lysates were separated on SDS-PAGE gel (4-15%),

transferred on nitrocellulose membrane, and blocked in 5% BSA in PBS. The blots were probed with primary

rabbit polyclonal antibodies (1:5000)/purified rabbit IgG (1:1000) raised against the different abundant cyst wall

proteins. Anti-rabbit HRP conjugated IgG (Thermo Fisher Scientific) was used as the secondary antibody. Rabbit

pre-immune serum or anti-rabbit IgG were used as control. Super Signal West Pico PLUS (Thermo Fisher

Scientific) substrate was used for chemiluminescent detection. Blots were imaged using GE ImageQuant LAS

4000 gel imager.

Immuno-staining of the mature cysts with purified rabbit polyclonal antibodies

To check the applicability of custom raised polyclonal rabbit antibodies against detection of Jonah-1, Luke-2,

Leo-A, and laccase paralogs*,* we performed IFA imaging using mature cysts from other *Acanthamoeba*

species/strains. For IFA imaging, ~0.5 to 1.0 x 10^7 mature cysts (120 h post encystation) from different strains

of *Acanthamoeba* were washed in PBS and fixed in 4% paraformaldehyde for 15 minutes at room temperature.

Following fixation, cysts were washed three times with PBS and blocked with 1% BSA for 1 h at room

temperature. The cysts were further incubated with the rabbit polyclonal antibody (1:200 dilution) for 1h at room

temperature. Subsequently these cysts were washed three times with PBS and incubated with secondary anti-

rabbit IgG conjugated with Alexa flour 488 (1:300) and Calcofluor White 1:20 (CFW, 1 mg/ml, Sigma Aldrich) for

30 min at room temperature. The cysts were washed 3 times with 1x PBS and mounted in VECTASHIELD®

Antifade Mounting Medium (Vector Laboratories, Newark, CA). Samples were illuminated using 380nm (CFW),

and 488 nm (Alexa flour 488) laser excitation. The stained cysts were imaged using confocal microscope as

detailed above. For each rAb and each *Acanthamoeba* isolate, we counted at least 100 cysts in random fields

to determine what percentage of CFW-labeled cysts were detected with the rAb. This experiment was repeated

twice and counts for the two experiments were averaged. Finally, we examined the same set of slides using a

Plan-APO Chromat 100x oil immersion lens on a Zeiss Axio Observer Z1 microscope equipped with Axio Cam

ERc5s to show that a confocal microscope was not needed to detect rAb-labeled cysts.

 Figure legends.

Fig 1. Localization and structure of Jonah lectins. (A) Drawings show Jonah-1 contains a signal peptide

(purple), followed by a Thr-rich spacer (light blue), and a single BHF (green). Jonah-3-AmoebaDB has twelve

TMHs (red) and three BHFs, while Jonah-3(c) has three BHFs, three Ser-rich domains (yellow), and no TMHs.

(B) Confocal microscopy shows Jonah-1 localizes to the ectocyst layer (yellow arrows, stained with WGA). The

endocyst layer is marked with CFW (pink arrow), while ostioles appear as dimples (purple arrows). **(C)** Jonah-3-

AmoebaDB with 12 TMHs forms rings around ostioles (green arrows) when expressed under its own promoter

with a GFP tag. **(D)** Jonah-3(c) localizes to the ectocyst layer, like Jonah-1. Confocal micrographs here and in

Fig 2 to 7 were shot with a 60x objective, and 3D reconstructions were made from 0.1 μm optical sections. **(E)**

AlphaFold structure with confidence colored shows the Jonah-1 lectin is composed of a single three-sided, β-

helical fold (BHF), as well as two α-helices (function unknown), and a 170-aa, unstructured domain rich in Thr

(light blue). **(F)** Jonah-3-(c) has three BHFs (red, blue, and yellow) and 132-aa, 267-aa and 308-aa long,

unstructured domains rich in Ser (yellow). **(G)** End-view with confidence colored shows the Jonah-1 BHF is three-

sided (BHFs may also be two-sided). **(H)** Foldseek analysis shows the Jonah-1 BHF (white) closely matches the

BHF of *Colwella sp.* (blue), the structure of which has been solved (PDB 3WP9). The E-value is 5.10e-4, and

the RMSD is 9.21, despite a sequence identity of just 12% over a 245-aa overlap. Scale bars for **B** to **D** are each

5 μm.

Fig 2. Structure and localization and structure of Luke-2. A. Drawings show Luke-2 and Luke-3 contain a

signal peptide (purple) and two or three β-jelly roll folds (BJRFs) (orange), respectively, which are separated by

Ser-rich spacers (yellow). B. Confocal microscopy shows WT Luke-2-GFP localizes to the endocyst layer (pink

arrows) and forms a flat ring around ostioles (blue-green arrows) (see also Fig. 5D). C. When five Tyr and a Phe

are mutated to Ala (see H and I below), Ala-mut Luke-2-GFP no longer localizes to endocyst layer. D. AlphaFold

shows Luke-2 contains two BJRFs, which are separated by a 47-aa unstructured, spacer rich in Ser (yellow). E.

Luke-3 has three BHFs separated by 42-aa and 33-aa long, unstructured spacers that are also Ser-rich. F and

G. Foldseek shows the N-half BJRF of Luke-2 shares a similar structure with a BJRF of *Dictyostelium* wall protein

and with CBM2 of a *Cellulomonas fimi* endoglucanase (blue), the structure of which has been solved (PDB

1EXH). The E-value for the BJRFs of Luke-2 and *D. discoideum* wall protein is 4.99e-7, and the RMSD is 1.79,

strongly suggesting common ancestry. The E-value for the N-terminal BJRF of Luke-2 and the CBM2 of *C. fimi*

is 9.72e-5, and the RMSD is 5.01, despite a 16% identity over a 105-aa overlap. The N-terminal BJRF of Luke-

2 has three linearly arranged aromatics (orange), which match those of *D. discoideum* and *C. fimi* (pink arrows).

H and I. Surface views of Luke-2-N and Luke-2-C highlight linear arrays of Trp (blue) and Phe (pink), which were

mutated to make Ala-mut of Luke-2-GFP that no longer localizes to the endocyst layer and ostioles (see B

above). J. Western blots show that MBP alone fails to bind to Avicel cellulose, while MBP-Luke-2 and MBP-

Leo-A fusion-proteins (described in Fig. 3) each bind well to Avicel cellulose. In contrast, cellulose binding is lost

when six aromatics are mutated to Ala in both Luke-2 and Leo-A. These experiments support the idea that Ala-

mutants of Leo-A and Luke-2 mislocalize in cyst walls, because of the loss of cellulose binding. Scale bars for **B**

and **C** are each 5 µm.

Fig 3. Structure and localization of Leo lectins. (A) Drawings show Leo-A contains a signal peptide (purple)

and two adjacent sets of four disulfide knots (4DKs) (blue), while Leo-S contains a Thr-rich spacer (grey-blue)

between sets of 4DKs. **(B)** Confocal microscopy shows WT Leo-A localizes to the endocyst layer (pink arrows)

and forms a flat ring around ostioles (blue-green arrows) (see also Fig 5D). **(C)**. When six Tyr are mutated to Ala

(see H below), Ala-mut Leo-A (grey arrow) no longer localizes to endocyst layer. **(D)** Early in encystation (12

hours), secretory vesicles (SV) containing Leo-S(c) fill the cytosol of Ac, which lack a cyst wall as shown by

failure to label with WGA or CFW. **(E)** Later in encystation (24 hours), Leo-S(c) has a somewhat patchy

distribution in the single-layered wall, which now labels with WGA and CFW. **(F)** In mature cyst walls (72 hours),

Leo-S(c) has a homogeneous distribution in the ectocyst layer (yellow arrows). Note that CFW predominantly

labels the endocyst layer of mature cysts. **(G)** A ribbon diagram of Leo-A highlights 4DKs (red spheres), which

connect short, parallel loops. **(H)** The surface of Leo-A shows three Tyr residues (green) linearly arrayed in each

4DK, which were mutated to Ala to test the effects on localization (see C above) and cellulose-binding (see Fig

2J). **(I)** The surface of Leo-S-corrected shows a 252-aa long unstructured, spacer rich in Thr (blue) between

4DKs, each of which contains linearly arrayed Tyr residues. Foldseek had no hit with the structure of Leo-A.

Scale bars for **B** to **F** are each 5 µm.

Fig 4. Localization and structure of an abundant laccase. (A) The Ac Laccase-1 has signal peptide (purple) and three copper oxidase domains (red), separated by short, unstructured spacers lacking either Ser- or Thr (white). **(B)** Confocal microscopy shows that early in encystation (18 hours) secretory vesicles (SV) containing laccase-GFP fill the cytosol of Ac, which lack a cyst wall as shown by failure to label with WGA or CFW. **(C)** Later in encystation (24 hours), laccase-GFP has a homogeneous distribution in the single-layered wall, which now labels with lightly with WGA and somewhat more with CFW. **(D)** In mature cyst walls (72 hours), laccase-GFP has the same homogeneous distribution in the ectocyst layer (yellow arrows) with ostioles as dimples (purple arrows). **(E)** AlphaFold structure shows CuRO-1 (blue), CuRO-2 (red), and CuRO-3 (yellow) domains of the laccase, all of which are predicted with confidence (not shown). **(G)** Typical for an enzyme, the E-value is 1.32e-59 and the RMSD = 2.75 with a sequence identity of 39% over a 505-aa overlap. Scale bars for **B** to **D** are each 5 μm.

Fig 5. Promoter swaps between either Jonah-1 and Luke-2 or laccase-1 and Leo-A. (A) RT-PCR shows that ectocyst proteins (Jonah-1 and laccase-1) are made earlier than endocyst proteins (Luke-2 and Leo-A). Note values for each time points are normalized with calreticulin, which is highly expressed in trophozoites and encysting Ac. **(B)** In the first promoter swap, Jonah-1-GFP expressed under its own earlier promoter localizes to the ectocyst layer (yellow arrow) (see also Fig 1B). **(C)** Expression under the later Luke-2 promoter causes Jonah-1-GFP to localize to the ostioles (blue-green arrows), which increase in thickness and number, while there is minimal localization of Jonah-1-GFP to the endocyst layer (pink arrows). **(D)** Conversely, Luke-2-GFP under its own later promoter localizes to the endocyst layer and forms a narrow ring around ostioles (see also Fig 2B) **(E)** Expression under the earlier Jonah-1 promoter causes Luke-2 to relocate to the ectocyst layer. Remarkably, the narrow ring of Luke-2-GFP around the ostioles appears the same under either its own or the Jonah-1 promoter, suggesting its target glycopolymer is available early and later during encystation. **(F)** In the second promoter swap, laccase-1-GFP expressed under its own earlier promoter localizes to the ectocyst layer (see also Fig 4D). **(G)** Expression under the later Leo-A promoter causes laccase-1-GFP to localize to the ostioles with minimal localization to the endocyst layer. **(H)** Conversely, Leo-A-GFP under its own later promoter localizes to the endocyst layer and forms a narrow ring around ostioles. **(I)** Expression under the earlier laccase promoter causes Leo-A to localize to the ectocyst layer and forms a narrow ring around ostioles. These experiments show

that under both early and late promoters, Jonah-1 and Laccase-1 have similar localizations in the cyst wall,

suggesting that each bind to the same glycopolymer(s). Luke-2 and Leo-A also have similar localizations under

early and late promoters, but the localization is distinct from that of Jonah-1 and Laccase-1, suggesting pairs of

wall proteins are binding to different glycopolymers. Scale bars for **B** to **I** are each 5 µm.

Fig. 6. Binding of rAbs to cyst wall proteins and to Neff strain Ac. (A) Western blots show pre-bleeds from

rabbits (negative control) fail to bind to proteins of trophozoites or cysts, while rAbs fail to bind to trophozoite

proteins (a second negative control) but bind well to cyst proteins of Neff strain. As discussed in detail in the

Results, rAbs to the BHF of Jonah-1 binds to the expected 55-kDa band and to two lower mol wt bands. Anti-

Luke-2 rAbs bind to a heavy and broad 50-kDa band, which is greater than the expected size of 27-kDa. Anti-

Leo rAbs bind to a 13-kDa band, which is slightly smaller than the expected size of 17-kDA. Anti-laccase rAbs

bind to a 75-kDa band, which is slightly bigger than the expected size of 64-kDa, as well as to a less abundant

55-kDa band. **(B)** High power confocal microscopy shows rAbs to Jonah-1 bind in a patchy distribution to the

surface of Neff strain of Ac and other *Acanthamoeba* species, which are shown in Fig 7A. **(C)** Anti-Luke-2 rAbs

bind strongly to ostioles and weakly to endocyst layer of some but not all Neff strain Ac. An asterisk is attached

to "ostioles" because anti-Luke-2 rAbs can also bind to ectocyst layer, as shown in Fig 7B. **(D)** Anti-Leo-A rAbs

also bind strongly to ostioles with a variable binding to the endocyst layer of some but not all of Neff strain Ac.

Again, an asterisk indicates that anti-Leo-A rAbs can also bind to the ectocyst layer, as shown in Fig 7C. **(E)**

Anti-laccase-1 rAbs bind in a homogenous manner to cysts of Neff strain, as well as to cyst of other

Acanthamoeba species, which are shown in Fig 7D. **(F to I)**. Low power confocal micrographs shows rAbs to

Jonah-1, Luke-2, Leo-A and laccase-1 detect well CFW-labeled cysts of the Neff strain. **(J)** A plot shows the

percentage of CFW-labeled cysts detected by four rAbs in two independent experiments (average plus SEM), in

which 100+ cysts were counted in random low power fields. Rabbit antibodies to Luke-2, Leo-A, and Laccase-1

each detect >95% of cysts 10 of 11 *Acanthamoeba* isolates tested, while rAbs to Jonah-1 are slightly less

efficient in detecting three isolates of *Acanthamoeba*. The exception is *A. mauritaniensis*, which was poorly

detected with all four rAbs, suggesting a problem with cyst formation under the conditions used here. Single

scale bar for **B** to **E** is 5 µm.

Fig 7. Binding of rAbs to Jonah-1, Luke-2, Leo-A, and Laccase to a diverse set of *Acanthamoeba Spp.* cysts. High power confocal micrographs show binding of rAbs to cysts ten *Acanthamoeba* isolates, the identities of which were confirmed by sequencing 18S RNAs. Column **(A)** Anti-Jonah-1 rAbs bind in a patchy distribution to the ectocyst layer (surface) of most cysts of *Acanthamoeba* isolates. Column **(B)** Anti-Luke-2 rAbs bind to the endocyst layer and ostioles of some cysts (e.g., *A. polyphaga, A. hatchetii, A, byersi*, and ESBC4) and to the ectocyst layer of other cysts (e.g., three *A. castellanii* and *A. astronyxis*). Asterisks indicate heterogeneous distribution of rAbs among the same *Acanthamoeba* isolate. Column **(C)** Anti-Leo-A rAbs also have a heterogeneous distribution on the ectocyst layer, endocyst layer, and/or ostioles. In some cases anti-Leo-A distribution matches that of anti-Luke-2 (e.g., *A. astronyxis, A. byersi*, and *A. sp. 13*), in the same way that Leo-GFP and Luke-GFP match in the promoter swap experiments (Fig 5). Column **(D)** Anti-Jonah-1 rAbs bind in a homogeneous distribution to the ectocyst layer (surface) of the vast majority of cysts of *Acanthamoeba* isolates. Counts of CFW-labeled cysts by each antibody are shown in Fig 6J. Single scale bar for **A** to **D** is 50 µm.

Acknowledgments

Thanks to Jonathan Stefely (Massachusetts General Hospital), Sarah Calvo (Broad Institute and Howard Hughes Medical Institute), and Brian Haas (Broad Institute) for access to the unfinished transcriptome analysis of *Acanthamoeba castellanii* Neff strain. Thanks to Monica Crary of Alcon, Inc. and Noorjahan Panjwani of Tufts University for cultures of *Acanthamoeba*. Thanks to Cataldo Leone, Dean of BUSDM for financial support for this work. This paper is dedicated to Jude, the cousin of Jonah, Luke, and Leo.

Supporting Information S1 lists the primers used for making constructs and for performing qRT-PCR.

Supporting Information S2 lists promoter sequences and coding sequences of the abundant cyst wall proteins used for making GFP-tagged proteins. Also listed are the coding sequences used to make MBP-fusions for testing cellulose binding and for immunizing rabbits.

References

1.   Azzopardi M, Chong YJ, Ng B, Recchioni A, Logeswaran A, Ting DSJ. Diagnosis of Acanthamoeba Keratitis: Past, Present and Future. Diagnostics (Basel). 2023;13(16). Epub 2023/08/26. doi: 10.3390/diagnostics13162655. PubMed PMID: 37627913; PubMed Central PMCID: PMCPMC10453105.

2.   Bouten M EH. Diagnosis and Management of Acanthamoeba Keratitis: A Continental Approach. Parasitologia. 2022;2:167-97.

3.   Carnt N, Hoffman JM, Verma S, Hau S, Radford CF, Minassian DC, et al. Acanthamoeba keratitis: confirmation of the UK outbreak and a prospective case-control study identifying contributing risk factors. Br J Ophthalmol. 2018;102(12):1621-8. Epub 2018/09/21. doi: 10.1136/bjophthalmol-2018-312544. PubMed PMID: 30232172.

4.   Satlin MJ, Graham JK, Visvesvara GS, Mena H, Marks KM, Saal SD, et al. Fulminant and fatal encephalitis caused by Acanthamoeba in a kidney transplant recipient: case report and literature review. Transpl Infect Dis. 2013;15(6):619-26. Epub 2013/09/10. doi: 10.1111/tid.12131. PubMed PMID: 24010955.

5.   Dos Santos DL, Kwitko S, Marinho DR, de Araujo BS, Locatelli CI, Rott MB. Acanthamoeba keratitis in Porto Alegre (southern Brazil): 28 cases and risk factors. Parasitol Res. 2018;117(3):747-50. Epub 2018/01/15. doi: 10.1007/s00436-017-5745-y. PubMed PMID: 29332157.

6.   Ghaderifar S, Najafpoor AA, Zarrinfar H, Esmaily H, Hajialilo E. Isolation and identification of Acanthamoeba from pond water of parks in a tropical and subtropical region in the Middle East, and its relation with physicochemical parameters. BMC Microbiol. 2018;18(1):139. Epub 2018/10/21. doi: 10.1186/s12866-018-1301-x. PubMed PMID: 30340522; PubMed Central PMCID: PMCPMC6195741.

7.   Henriquez FL, Campbell SJ, Sundararaj BK, Cano A, Muench SP, Roberts CW. The Acanthamoeba shikimate pathway has a unique molecular arrangement and is essential for aromatic amino acid biosynthesis. Protist. 2015;166(1):93-105. Epub 2015/01/13. doi: 10.1016/j.protis.2014.12.001. PubMed PMID: 25576842.

8.   Pittet D, Allegranzi B, Boyce J, World Health Organization World Alliance for Patient Safety First Global Patient Safety Challenge Core Group of E. The World Health Organization Guidelines on Hand Hygiene in Health Care and their consensus recommendations. Infect Control Hosp Epidemiol. 2009;30(7):611-22. Epub 2009/06/11. doi: 10.1086/600379. PubMed PMID: 19508124.

9.   Zhang Y, Xu X, Wei Z, Cao K, Zhang Z, Liang Q. The global epidemiology and clinical diagnosis of Acanthamoeba keratitis. J Infect Public Health. 2023;16(6):841-52. Epub 2023/04/09. doi: 10.1016/j.jiph.2023.03.020. PubMed PMID: 37030037.

10.   Brown AC, Ross J, Jones DB, Collier SA, Ayers TL, Hoekstra RM, et al. Risk Factors for Acanthamoeba Keratitis-A Multistate Case-Control Study, 2008-2011. Eye Contact Lens. 2018;44 Suppl 1:S173-S8. Epub 2017/01/19. doi: 10.1097/ICL.0000000000000365. PubMed PMID: 28099282.

11.   Carnt N, Robaei D, Minassian DC, Dart JKG. Acanthamoeba keratitis in 194 patients: risk factors for bad outcomes and severe inflammatory complications. Br J Ophthalmol. 2018;102(10):1431-5. Epub 2018/01/05. doi: 10.1136/bjophthalmol-2017-310806. PubMed PMID: 29298778.

12.   Lorenzo-Morales J, Martin-Navarro CM, Lopez-Arencibia A, Arnalich-Montiel F, Pinero JE, Valladares B. Acanthamoeba keratitis: an emerging disease gathering importance worldwide? Trends Parasitol. 2013;29(4):181-7. Epub 2013/02/26. doi: 10.1016/j.pt.2013.01.006. PubMed PMID: 23433689.

13.   Diehl MLN, Paes J, Rott MB. Genotype distribution of Acanthamoeba in keratitis: a systematic review. Parasitol Res. 2021;120(9):3051-63. Epub 2021/08/06. doi: 10.1007/s00436-021-07261-1. PubMed PMID: 34351492; PubMed Central PMCID: PMCPMC8339388.

14.   Fuerst PA. The Status of Molecular Analyses of Isolates of Acanthamoeba Maintained by International Culture Collections. Microorganisms. 2023;11(2). Epub 2023/02/26. doi: 10.3390/microorganisms11020295. PubMed PMID: 36838260; PubMed Central PMCID: PMCPMC9961329.

15.   Fuerst PA, Booton GC, Crary M. Phylogenetic analysis and the evolution of the 18S rRNA gene typing
system of Acanthamoeba. J Eukaryot Microbiol. 2015;62(1):69-84. Epub 2014/10/07. doi: 10.1111/jeu.12186.
PubMed PMID: 25284310.
16.   Rayamajhee B, Sharma S, Willcox M, Henriquez FL, Rajagopal RN, Shrestha GS, et al. Assessment of
genotypes, endosymbionts and clinical characteristics of Acanthamoeba recovered from ocular infection. BMC
Infect Dis. 2022;22(1):757. Epub 2022/09/30. doi: 10.1186/s12879-022-07741-4. PubMed PMID: 36175838;
PubMed Central PMCID: PMCPMC9520893.
17.   Amos B, Aurrecoechea C, Barba M, Barreto A, Basenko EY, Bazant W, et al. VEuPathDB: the eukaryotic
pathogen, vector and host bioinformatics resource center. Nucleic Acids Res. 2022;50(D1):D898-D911. Epub
2021/11/01. doi: 10.1093/nar/gkab929. PubMed PMID: 34718728; PubMed Central PMCID:
PMCPMC8728164.
18.   Clarke M, Lohan AJ, Liu B, Lagkouvardos I, Roy S, Zafar N, et al. Genome of Acanthamoeba castellanii
highlights extensive lateral gene transfer and early evolution of tyrosine kinase signaling. Genome Biol.
2013;14(2):R11. Epub 2013/02/05. doi: 10.1186/gb-2013-14-2-r11. PubMed PMID: 23375108; PubMed Central
PMCID: PMCPMC4053784.
19.   Bernard C, Locard-Paulet M, Noel C, Duchateau M, Giai Gianetto Q, Moumen B, et al. A time-resolved
multi-omics atlas of Acanthamoeba castellanii encystment. Nat Commun. 2022;13(1):4104. Epub 2022/07/15.
doi: 10.1038/s41467-022-31832-0. PubMed PMID: 35835784; PubMed Central PMCID: PMCPMC9283445.
20.   Matthey-Doret C, Colp MJ, Escoll P, Thierry A, Moreau P, Curtis B, et al. Chromosome-scale assemblies
of Acanthamoeba castellanii genomes provide insights into Legionella pneumophila infection-related
chromatin reorganization. Genome Res. 2022;32(9):1698-710. Epub 2022/09/16. doi: 10.1101/gr.276375.121.
PubMed PMID: 36109147; PubMed Central PMCID: PMCPMC9528979.
21.   Jensen T, Barnes WG, Meyers D. Axenic cultivation of large populations of Acanthamoeba castellanii
(JBM). J Parasitol. 1970;56(5):904-6. Epub 1970/10/01. PubMed PMID: 5504530.
22.   Neff RJ, Ray SA, Benton WF, Wilborn M. Induction of synchronous encystment (differentiation) in
Acanthamoeba spp. Methods in Cell Biology. 1964;1:55-83. doi:10.1016/S0091-679X(08)62086-5.
23.   Aqeel Y, Rodriguez R, Chatterjee A, Ingalls RR, Samuelson J. Killing of diverse eye pathogens
(Acanthamoeba spp., Fusarium solani, and Chlamydia trachomatis) with alcohols. PLoS Negl Trop Dis.
2017;11(2):e0005382. Epub 2017/02/10. doi: 10.1371/journal.pntd.0005382. PubMed PMID: 28182670;
PubMed Central PMCID: PMCPMC5321442.
24.   Campolo A, Pifer R, Walters R, Thomas M, Miller E, Harris V, et al. Acanthamoeba spp. aggregate and
encyst on contact lens material increasing resistance to disinfection. Front Microbiol. 2022;13:1089092. Epub
2023/01/06. doi: 10.3389/fmicb.2022.1089092. PubMed PMID: 36601401; PubMed Central PMCID:
PMCPMC9806144.
25.   Coulon C, Collignon A, McDonnell G, Thomas V. Resistance of Acanthamoeba cysts to disinfection
treatments used in health care settings. J Clin Microbiol. 2010;48(8):2689-97. Epub 2010/06/04. doi:
10.1128/JCM.00309-10. PubMed PMID: 20519477; PubMed Central PMCID: PMCPMC2916629.
26.   Dupuy M, Berne F, Herbelin P, Binet M, Berthelot N, Rodier MH, et al. Sensitivity of free-living amoeba
trophozoites and cysts to water disinfectants. Int J Hyg Environ Health. 2014;217(2-3):335-9. Epub
2013/08/13. doi: 10.1016/j.ijheh.2013.07.007. PubMed PMID: 23932411.
27.   Walters R, Campolo A, Miller E, McAnally C, Gabriel M, Shannon P, et al. Differential Antimicrobial
Efficacy of Preservative-Free Contact Lens Disinfection Systems against Common Ocular Pathogens. Microbiol
Spectr. 2022;10(1):e0213821. Epub 2022/02/10. doi: 10.1128/spectrum.02138-21. PubMed PMID: 35138157;
PubMed Central PMCID: PMCPMC8826922.
28.   Wang Y, Jiang L, Zhao Y, Ju X, Wang L, Jin L, et al. Biological characteristics and pathogenicity of
Acanthamoeba. Front Microbiol. 2023;14:1147077. Epub 2023/04/24. doi: 10.3389/fmicb.2023.1147077.
PubMed PMID: 37089530; PubMed Central PMCID: PMCPMC10113681.

29.   Magistrado-Coxen P, Aqeel Y, Lopez A, Haserick JR, Urbanowicz BR, Costello CE, et al. The most
abundant cyst wall proteins of Acanthamoeba castellanii are lectins that bind cellulose and localize to distinct
structures in developing and mature cyst walls. PLoS Negl Trop Dis. 2019;13(5):e0007352. Epub 2019/05/17.
doi: 10.1371/journal.pntd.0007352. PubMed PMID: 31095564; PubMed Central PMCID: PMCPMC6541295.
30.   Wilhelmus KR, Osato MS, Font RL, Robinson NM, Jones DB. Rapid diagnosis of Acanthamoeba keratitis
using calcofluor white. Arch Ophthalmol. 1986;104(9):1309-12. Epub 1986/09/01. doi:
10.1001/archopht.1986.01050210063026. PubMed PMID: 2428343.
31.   Orlean P. Architecture and biosynthesis of the Saccharomyces cerevisiae cell wall. Genetics.
2012;192(3):775-818. Epub 2012/11/09. doi: 10.1534/genetics.112.144485. PubMed PMID: 23135325;
PubMed Central PMCID: PMCPMC3522159.
32.   West CM. Comparative analysis of spore coat formation, structure, and function in Dictyostelium. Int
Rev Cytol. 2003;222:237-93. Epub 2002/12/31. doi: 10.1016/s0074-7696(02)22016-1. PubMed PMID:
12503851.
33.   Zablackis E, Huang J, Muller B, Darvill AG, Albersheim P. Characterization of the cell-wall
polysaccharides of Arabidopsis thaliana leaves. Plant Physiol. 1995;107(4):1129-38. Epub 1995/04/01. doi:
10.1104/pp.107.4.1129. PubMed PMID: 7770522; PubMed Central PMCID: PMCPMC157245.
34.   Bowers B, Korn ED. The fine structure of Acanthamoeba castellanii (Neff strain). II. Encystment. J Cell
Biol. 1969;41(3):786-805. Epub 1969/06/01. doi: 10.1083/jcb.41.3.786. PubMed PMID: 5768875; PubMed
Central PMCID: PMCPMC2107820.
35.   Potter JL, Weisman RA. Cellulose synthesis by extracts of Acanthamoeba castellanii during encystment.
Stimulation of the incorporation of radioactivity from UDP-(14C)glucose into alkali-soluble and insoluble beta-
glucans by glucose 6-phosphate and related compounds. Biochim Biophys Acta. 1976;428(1):240-52. Epub
1976/03/25. doi: 10.1016/0304-4165(76)90125-2. PubMed PMID: 1260020.
36.   Morozov AA, Likhoshway YV. Evolutionary history of the chitin synthases of eukaryotes. Glycobiology.
2016;26(6):635-9. Epub 2016/02/19. doi: 10.1093/glycob/cww018. PubMed PMID: 26887391.
37.   Bateman E. Expression plasmids and production of EGFP in stably transfected Acanthamoeba. Protein
Expr Purif. 2010;70(1):95-100. Epub 2009/10/20. doi: 10.1016/j.pep.2009.10.008. PubMed PMID: 19836453;
PubMed Central PMCID: PMCPMC2822096.
38.   Peng Z, Omaruddin R, Bateman E. Stable transfection of Acanthamoeba castellanii. Biochim Biophys
Acta. 2005;1743(1-2):93-100. Epub 2005/03/22. doi: 10.1016/j.bbamcr.2004.08.014. PubMed PMID:
15777844.
39.   Jumper J, Evans R, Pritzel A, Green T, Figurnov M, Ronneberger O, et al. Highly accurate protein
structure prediction with AlphaFold. Nature. 2021;596(7873):583-9. Epub 2021/07/16. doi: 10.1038/s41586-
021-03819-2. PubMed PMID: 34265844; PubMed Central PMCID: PMCPMC8371605 have filed non-provisional
patent applications 16/701,070 and PCT/EP2020/084238, and provisional patent applications 63/107,362,
63/118,917, 63/118,918, 63/118,921 and 63/118,919, each in the name of DeepMind Technologies Limited,
each pending, relating to machine learning for predicting protein structures. The other authors declare no
competing interests.
40.   van Kempen M, Kim SS, Tumescheit C, Mirdita M, Lee J, Gilchrist CLM, et al. Fast and accurate protein
structure search with Foldseek. Nat Biotechnol. 2023. Epub 2023/05/09. doi: 10.1038/s41587-023-01773-0.
PubMed PMID: 37156916.
41.   Varadi M, Anyango S, Deshpande M, Nair S, Natassia C, Yordanova G, et al. AlphaFold Protein Structure
Database: massively expanding the structural coverage of protein-sequence space with high-accuracy models.
Nucleic Acids Res. 2022;50(D1):D439-D44. Epub 2021/11/19. doi: 10.1093/nar/gkab1061. PubMed PMID:
34791371; PubMed Central PMCID: PMCPMC8728224.

42. Boraston AB, Bolam DN, Gilbert HJ, Davies GJ. Carbohydrate-binding modules: fine-tuning
polysaccharide recognition. Biochem J. 2004;382(Pt 3):769-81. Epub 2004/06/25. doi: 10.1042/BJ20040892.
PubMed PMID: 15214846; PubMed Central PMCID: PMCPMC1133952.
43. Consortium CA. Ten years of CAZypedia: a living encyclopedia of carbohydrate-active enzymes.
Glycobiology. 2018;28(1):3-8. Epub 2017/10/19. doi: 10.1093/glycob/cwx089. PubMed PMID: 29040563.
44. Cosgrove DJ. Microbial Expansins. Annu Rev Microbiol. 2017;71:479-97. Epub 2017/09/10. doi:
10.1146/annurev-micro-090816-093315. PubMed PMID: 28886679.
45. Gilbert HJ, Knox JP, Boraston AB. Advances in understanding the molecular basis of plant cell wall
polysaccharide recognition by carbohydrate-binding modules. Curr Opin Struct Biol. 2013;23(5):669-77. Epub
2013/06/19. doi: 10.1016/j.sbi.2013.05.005. PubMed PMID: 23769966.
46. Lombard V, Golaconda Ramulu H, Drula E, Coutinho PM, Henrissat B. The carbohydrate-active enzymes
database (CAZy) in 2013. Nucleic Acids Res. 2014;42(Database issue):D490-5. Epub 2013/11/26. doi:
10.1093/nar/gkt1178. PubMed PMID: 24270786; PubMed Central PMCID: PMCPMC3965031.
47. Urbanowicz BR, Catala C, Irwin D, Wilson DB, Ripoll DR, Rose JK. A tomato endo-beta-1,4-glucanase,
SlCel9C1, represents a distinct subclass with a new family of carbohydrate binding modules (CBM49). J Biol
Chem. 2007;282(16):12066-74. Epub 2007/02/27. doi: 10.1074/jbc.M607925200. PubMed PMID: 17322304.
48. Hallgren J. TKD, Pedersen M.D., Almagro Armenteros J.J., Marcatili P., Nielsen H., Krogh A., Winther O.
DeepTMHMM predicts alpha and beta transmembrane proteins using deep neural networks. bioRxiv. 2022.
doi: doi: 10.1101/2022.04.08.487609.
49. Nielsen H. Predicting Secretory Proteins with SignalP. Methods Mol Biol. 2017;1611:59-73. Epub
2017/04/30. doi: 10.1007/978-1-4939-7015-5_6. PubMed PMID: 28451972.
50. Enguita FJ, Martins LO, Henriques AO, Carrondo MA. Crystal structure of a bacterial endospore coat
component. A laccase with enhanced thermostability properties. J Biol Chem. 2003;278(21):19416-25. Epub
2003/03/15. doi: 10.1074/jbc.M301251200. PubMed PMID: 12637519.
51. Kiiskinen LL, Palonen H, Linder M, Viikari L, Kruus K. Laccase from Melanocarpus albomyces binds
effectively to cellulose. FEBS Lett. 2004;576(1-2):251-5. Epub 2004/10/12. doi: 10.1016/j.febslet.2004.08.040.
PubMed PMID: 15474046.
52. Banerjee S, Vishwanath P, Cui J, Kelleher DJ, Gilmore R, Robbins PW, et al. The evolution of N-glycan-
dependent endoplasmic reticulum quality control factors for glycoprotein folding and degradation. Proc Natl
Acad Sci U S A. 2007;104(28):11676-81. Epub 2007/07/04. doi: 10.1073/pnas.0704862104. PubMed PMID:
17606910; PubMed Central PMCID: PMCPMC1905923.
53. Altschul SF, Madden TL, Schaffer AA, Zhang J, Zhang Z, Miller W, et al. Gapped BLAST and PSI-BLAST: a
new generation of protein database search programs. Nucleic Acids Res. 1997;25(17):3389-402. Epub
1997/09/01. doi: 10.1093/nar/25.17.3389. PubMed PMID: 9254694; PubMed Central PMCID:
PMCPMC146917.
54. Finn RD, Attwood TK, Babbitt PC, Bateman A, Bork P, Bridge AJ, et al. InterPro in 2017-beyond protein
family and domain annotations. Nucleic Acids Res. 2017;45(D1):D190-D9. Epub 2016/12/03. doi:
10.1093/nar/gkw1107. PubMed PMID: 27899635; PubMed Central PMCID: PMCPMC5210578.
55. Wang Y, Slade MB, Gooley AA, Atwell BJ, Williams KL. Cellulose-binding modules from extracellular
matrix proteins of Dictyostelium discoideum stalk and sheath. Eur J Biochem. 2001;268(15):4334-45. Epub
2001/08/08. doi: 10.1046/j.1432-1327.2001.02354.x. PubMed PMID: 11488929.
56. Xu Y, Liang X, Chen Y, Koehler TM, Hook M. Identification and biochemical characterization of two
novel collagen binding MSCRAMMs of Bacillus anthracis. J Biol Chem. 2004;279(50):51760-8. Epub
2004/10/01. doi: 10.1074/jbc.M406417200. PubMed PMID: 15456768.
57. Davies G, Henrissat B. Structures and mechanisms of glycosyl hydrolases. Structure. 1995;3(9):853-9.
Epub 1995/09/15. doi: 10.1016/S0969-2126(01)00220-9. PubMed PMID: 8535779.

58.   Xu GY, Ong E, Gilkes NR, Kilburn DG, Muhandiram DR, Harris-Brandts M, et al. Solution structure of a
cellulose-binding domain from Cellulomonas fimi by nuclear magnetic resonance spectroscopy. Biochemistry.
1995;34(21):6993-7009. Epub 1995/05/30. PubMed PMID: 7766609.
59.   Duong-Ly KC, Gabelli SB. Affinity Purification of a Recombinant Protein Expressed as a Fusion with the
Maltose-Binding Protein (MBP) Tag. Methods Enzymol. 2015;559:17-26. Epub 2015/06/23. doi:
10.1016/bs.mie.2014.11.004. PubMed PMID: 26096500; PubMed Central PMCID: PMCPMC4612625.
60.   Kapust RB, Waugh DS. Escherichia coli maltose-binding protein is uncommonly effective at promoting
the solubility of polypeptides to which it is fused. Protein Sci. 1999;8(8):1668-74. Epub 1999/08/19. doi:
10.1110/ps.8.8.1668. PubMed PMID: 10452611; PubMed Central PMCID: PMCPMC2144417.
61.   Chatterjee A, Ghosh SK, Jang K, Bullitt E, Moore L, Robbins PW, et al. Evidence for a "wattle and daub"
model of the cyst wall of entamoeba. PLoS Pathog. 2009;5(7):e1000498. Epub 2009/07/07. doi:
10.1371/journal.ppat.1000498. PubMed PMID: 19578434; PubMed Central PMCID: PMCPMC2698119.
62.   Harata K, Nagahora H, Jigami Y. X-ray structure of wheat germ agglutinin isolectin 3. Acta Crystallogr D
Biol Crystallogr. 1995;51(Pt 6):1013-9. Epub 1995/11/01. doi: 10.1107/S0907444995004070. PubMed PMID:
15299769.
63.   Kuranda MJ, Robbins PW. Chitinase is required for cell separation during growth of Saccharomyces
cerevisiae. J Biol Chem. 1991;266(29):19758-67. Epub 1991/10/15. PubMed PMID: 1918080.
64.   Van Dellen K, Ghosh SK, Robbins PW, Loftus B, Samuelson J. Entamoeba histolytica lectins contain
unique 6-Cys or 8-Cys chitin-binding domains. Infect Immun. 2002;70(6):3259-63. Epub 2002/05/16. doi:
10.1128/IAI.70.6.3259-3263.2002. PubMed PMID: 12011021; PubMed Central PMCID: PMCPMC127964.
65.   Hanada Y, Nishimiya Y, Miura A, Tsuda S, Kondo H. Hyperactive antifreeze protein from an Antarctic
sea ice bacterium Colwellia sp. has a compound ice-binding site without repetitive sequences. FEBS J.
2014;281(16):3576-90. Epub 2014/06/19. doi: 10.1111/febs.12878. PubMed PMID: 24938370.
66.   Liu Z, Xie T, Zhong Q, Wang G. Crystal structure of CotA laccase complexed with 2,2-azinobis-(3-
ethylbenzothiazoline-6-sulfonate) at a novel binding site. Acta Crystallogr F Struct Biol Commun. 2016;72(Pt
4):328-35. Epub 2016/04/07. doi: 10.1107/S2053230X1600426X. PubMed PMID: 27050268; PubMed Central
PMCID: PMCPMC4822991.
67.   Schiller B, Makrypidi G, Razzazi-Fazeli E, Paschinger K, Walochnik J, Wilson IB. Exploring the unique N-
glycome of the opportunistic human pathogen Acanthamoeba. J Biol Chem. 2012;287(52):43191-204. Epub
2012/11/10. doi: 10.1074/jbc.M112.418095. PubMed PMID: 23139421; PubMed Central PMCID:
PMCPMC3527907.
68.   Garate M, Cubillos I, Marchant J, Panjwani N. Biochemical characterization and functional studies of
Acanthamoeba mannose-binding protein. Infect Immun. 2005;73(9):5775-81. Epub 2005/08/23. doi:
10.1128/IAI.73.9.5775-5781.2005. PubMed PMID: 16113295; PubMed Central PMCID: PMCPMC1231072.
69.   Samuelson J, Bushkin GG, Chatterjee A, Robbins PW. Strategies to discover the structural components
of cyst and oocyst walls. Eukaryot Cell. 2013;12(12):1578-87. Epub 2013/10/08. doi: 10.1128/EC.00213-13.
PubMed PMID: 24096907; PubMed Central PMCID: PMCPMC3889564.
70.   Ghosh SK, Van Dellen KL, Chatterjee A, Dey T, Haque R, Robbins PW, et al. The Jacob2 lectin of the
Entamoeba histolytica cyst wall binds chitin and is polymorphic. PLoS Negl Trop Dis. 2010;4(7):e750. Epub
2010/07/24. doi: 10.1371/journal.pntd.0000750. PubMed PMID: 20652032; PubMed Central PMCID:
PMCPMC2907411.
71.   Haserick JR, Klein JA, Costello CE, Samuelson J. Cryptosporidium parvum vaccine candidates are
incompletely modified with O-linked-N-acetylgalactosamine or contain N-terminal N-myristate and S-
palmitate. PLoS One. 2017;12(8):e0182395. Epub 2017/08/10. doi: 10.1371/journal.pone.0182395. PubMed
PMID: 28792526; PubMed Central PMCID: PMCPMC5549699.
72.   Van Dellen KL, Chatterjee A, Ratner DM, Magnelli PE, Cipollo JF, Steffen M, et al. Unique
posttranslational modifications of chitin-binding lectins of Entamoeba invadens cyst walls. Eukaryot Cell.

2006;5(5):836-48. Epub 2006/05/10. doi: 10.1128/EC.5.5.836-848.2006. PubMed PMID: 16682461; PubMed
Central PMCID: PMCPMC1459681.
73.   Kang AY, Park AY, Shin HJ, Khan NA, Maciver SK, Jung SY. Production of a monoclonal antibody against
a mannose-binding protein of Acanthamoeba culbertsoni and its localization. Exp Parasitol. 2018;192:19-24.
Epub 2018/07/22. doi: 10.1016/j.exppara.2018.07.009. PubMed PMID: 30031120.
74.   Kim DY, Son DH, Matin A, Jung SY. Production of a monoclonal antibody against a galactose-binding
protein of Acanthamoeba castellanii and its cytotoxicity. Parasitol Res. 2021;120(11):3845-50. Epub
2021/09/26. doi: 10.1007/s00436-021-07321-6. PubMed PMID: 34561750.
75.   Kim MJ, Chu KB, Lee HA, Quan FS, Kong HH, Moon EK. Detection of Acanthamoeba spp. using
carboxylesterase antibody and its usage for diagnosing Acanthamoeba-keratitis. PLoS One.
2022;17(1):e0262223. Epub 2022/01/06. doi: 10.1371/journal.pone.0262223. PubMed PMID: 34986189;
PubMed Central PMCID: PMCPMC8730387.
76.   Weber-Lima MM, Prado-Costa B, Becker-Finco A, Costa AO, Billilad P, Furst C, et al. Acanthamoeba spp.
monoclonal antibody against a CPA2 transporter: a promising molecular tool for acanthamoebiasis diagnosis
and encystment study. Parasitology. 2020;147(14):1678-88. Epub 2020/09/22. doi:
10.1017/S0031182020001778. PubMed PMID: 32951614; PubMed Central PMCID: PMCPMC10317748.
77.   Aykur M, Karakavuk M, Akil M, Can H, Doskaya M, Guruz AY, et al. Development and Sensitivity
Determination of 18S rRNA Gene-specific Fast Loop-mediated Isothermal Amplification (LAMP) Assay for Rapid
Detection of Acanthamoeba. Turkiye Parazitol Derg. 2023;47(3):129-35. Epub 2023/09/19. doi:
10.4274/tpd.galenos.2023.46362. PubMed PMID: 37724360.
78.   Mewara A, Khurana S, Yoonus S, Megha K, Tanwar P, Gupta A, et al. Evaluation of loop-mediated
isothermal amplification assay for rapid diagnosis of Acanthamoeba keratitis. Indian J Med Microbiol.
2017;35(1):90-4. Epub 2017/03/18. doi: 10.4103/ijmm.IJMM_16_227. PubMed PMID: 28303825.
79.   Samuelson J, Robbins PW. Effects of N-glycan precursor length diversity on quality control of protein
folding and on protein glycosylation. Semin Cell Dev Biol. 2015;41:121-8. Epub 2014/12/06. doi:
10.1016/j.semcdb.2014.11.008. PubMed PMID: 25475176; PubMed Central PMCID: PMCPMC4452448.
80.   Aurrecoechea C, Barreto A, Brestelli J, Brunk BP, Caler EV, Fischer S, et al. AmoebaDB and
MicrosporidiaDB: functional genomic resources for Amoebozoa and Microsporidia species. Nucleic Acids Res.
2011;39(Database issue):D612-9. Epub 2010/10/27. doi: 10.1093/nar/gkq1006. PubMed PMID: 20974635;
PubMed Central PMCID: PMCPMC3013638.
81.   Marchler-Bauer A, Derbyshire MK, Gonzales NR, Lu S, Chitsaz F, Geer LY, et al. CDD: NCBI's conserved
domain database. Nucleic Acids Res. 2015;43(Database issue):D222-6. Epub 2014/11/22. doi:
10.1093/nar/gku1221. PubMed PMID: 25414356; PubMed Central PMCID: PMCPMC4383992.
82.   Krogh A, Larsson B, von Heijne G, Sonnhammer EL. Predicting transmembrane protein topology with a
hidden Markov model: application to complete genomes. J Mol Biol. 2001;305(3):567-80. Epub 2001/01/12.
doi: 10.1006/jmbi.2000.4315. PubMed PMID: 11152613.
83.   Eisenhaber B, Bork P, Eisenhaber F. Prediction of potential GPI-modification sites in proprotein
sequences. J Mol Biol. 1999;292(3):741-58. Epub 1999/09/25. doi: 10.1006/jmbi.1999.3069. PubMed PMID:
10497036.
84.   Rigsby RE, Parker AB. Using the PyMOL application to reinforce visual understanding of protein
structure. Biochem Mol Biol Educ. 2016;44(5):433-7. Epub 2016/06/01. doi: 10.1002/bmb.20966. PubMed
PMID: 27241834.
006

Fig 4.

Fig 4

Fig 5.

A. Western blots of nAbs binding to trophozoites and cysts

Fig 6.

Fig 7.

Fig 7

Fig 1.

Fig 1

Fig 2.

Fig 2

Fig 3.