1   The ubiquitin ligase HUWE1 enhances WNT signaling by antagonizing destruction complex-

2   mediated β-catenin degradation and through a mechanism independent of β-catenin stability

4   Short Title: HUWE1 enhances WNT signaling via two mechanisms

Joseph K. McKenna[1,¶], Yalan Wu[1,¶], Praveen Sonkusre[1], Raj Chari[2] and Andres M. Lebensohn[1,*]

[1] Laboratory of Cellular and Molecular Biology, Center for Cancer Research, National Cancer

Institute, National Institutes of Health, Bethesda, Maryland, United States of America.

[2] Genome Modification Core, Laboratory Animal Sciences Program, Frederick National Lab for

Cancer Research, Frederick, Maryland, United States of America.

[*] Corresponding author

E-mail: andres.lebensohn@nih.gov (AML)

[¶] These authors contributed equally to this work and are listed in the order in which they started

working on this project.

Abstract:

WNT/β-catenin signaling is mediated by the transcriptional coactivator β-catenin (CTNNB1). CTNNB1 abundance is regulated by phosphorylation and proteasomal degradation promoted by a destruction complex composed of the scaffold proteins APC and AXIN1 or AXIN2, and the kinases CSNK1A1 and GSK3A or GSK3B. Loss of CSNK1A1 increases CTNNB1 abundance, resulting in hyperactive WNT signaling. Previously, we demonstrated that the HECT domain ubiquitin ligase HUWE1 is necessary for hyperactive WNT signaling in HAP1 haploid human cells lacking CSNK1A1. Here, we investigate the mechanism underlying this requirement. In the absence of CSNK1A1, GSK3A/GSK3B still phosphorylated a fraction of CTNNB1, promoting its degradation. HUWE1 loss enhanced GSK3A/GSK3B-dependent CTNNB1 phosphorylation, further reducing CTNNB1 abundance. However, the reduction in CTNNB1 caused by HUWE1 loss was disproportionately smaller than the reduction in WNT target gene transcription. To test if the reduction in WNT signaling resulted from reduced CTNNB1 abundance alone, we engineered the endogenous *CTNNB1* locus in HAP1 cells to encode a CTNNB1 variant insensitive to destruction complex-mediated phosphorylation and degradation. HUWE1 loss in these cells reduced WNT signaling with no change in CTNNB1 abundance. Genetic interaction and overexpression analyses revealed that the effects of HUWE1 on WNT signaling were not only mediated by GSK3A/GSK3B, but also by APC and AXIN1. Regulation of WNT signaling by HUWE1 required its ubiquitin ligase activity. These results suggest that in cells lacking CSNK1A1, a destruction complex containing APC, AXIN1 and GSK3A/GSK3B downregulates WNT signaling by phosphorylating and targeting CTNNB1 for degradation. HUWE1 enhances WNT signaling by antagonizing this activity. Therefore, HUWE1 enhances WNT/CTNNB1

signaling through two mechanisms, one that regulates CTNNB1 abundance and another that is

independent of CTNNB1 stability. Coordinated regulation of CTNNB1 abundance and an

independent signaling step by HUWE1 would be an efficient way to control WNT signaling

output, enabling sensitive and robust activation of the pathway.

Author Summary

The WNT pathway is a conserved signaling system with diverse functions in embryonic

development and adult tissue homeostasis. Dysregulation of WNT signaling drives many types

of cancer. Over four decades of research have revealed a great deal about how the core

components of the WNT pathway regulate signaling, but much less is known about additional

regulatory layers superimposed on the core signaling module. In this study we present an

example of such regulation by the ubiquitin ligase HUWE1. Phosphorylation of the

transcriptional co-activator β-catenin by a protein complex called the destruction complex targets

β-catenin for degradation. This is considered the main regulated step in WNT signaling. We

demonstrate that HUWE1 enhances WNT signaling through two distinct mechanisms. First,

HUWE1 antagonizes the phosphorylation and degradation of β-catenin by the destruction

complex. Second, HUWE1 enhances WNT signaling through a mechanism independent from

control of β-catenin stability. The effects of HUWE1 on WNT signaling require its ubiquitin

ligase activity, suggesting there is a HUWE1 substrate awaiting discovery. Our work therefore

reveals a new role for HUWE1 controlling the main regulated step in WNT signaling – β-catenin

phosphorylation by the destruction complex – and most likely a downstream mechanism.

<u>Introduction:</u>

During embryonic development and tissue homeostasis, WNT/β-catenin signaling orchestrates

cellular processes that control tissue patterning and morphogenesis, cell fate specification, and

stem cell self-renewal among many other functions [1, 2]. Mutations in WNT signaling pathway

components can drive tumorigenesis of many cancer types, most notably colorectal cancer [3, 4].

At the heart of the WNT/β-catenin signaling pathway, the destruction complex (DC) controls the

abundance of the transcriptional coactivator β-catenin (CTNNB1) by regulating its degradation

through the ubiquitin/proteasome system. The DC is comprised of a set of core components,

including the scaffold proteins APC and AXIN1 or AXIN2, and the kinases casein kinase 1α

(CSNK1A1) and glycogen synthase kinase 3α (GSK3A) or β (GSK3B) [5]. In the absence of

signals initiated by secreted WNT ligands, CSNK1A1 phosphorylates CTNNB1 at serine (S) 45

[6, 7], priming it for further sequential phosphorylation of threonine (T) 41, S37 and S33 by

GSK3A and/or GSK3B [6, 8] (we refer to residues S33, S37, T41 and S45 as the CTNNB1

phosphodegron). When phosphorylated, residues S33 and S37 create a recognition site for the

ubiquitin ligase complex SCF$^{\beta TrCP}$ [9, 10], which ubiquitylates CTNNB1 and targets it for

proteasomal degradation [5]. Therefore, CTNNB1 abundance is kept low and WNT-dependent

transcriptional programs are repressed. Binding of WNT ligands to the cell surface receptors

frizzled (FZD) and LDL receptor related proteins 5 (LRP5) or 6 (LRP6) triggers the recruitment

of dishevelled (DVL) and at least some DC components to FZD and LRP5/6. Formation of this

receptor complex, or signalosome, downregulates the DC [11, 12] and results in accumulation of

non-phosphorylated CTNNB1. CTNNB1 enters the nucleus, where it forms a complex with

transcription factors of the TCF/LEF family and other coactivators to drive WNT target gene

transcription [13].

This description of WNT/CTNNB1 signaling omits additional regulatory mechanisms

superimposed on the core pathway that control the abundance, interactions, and subcellular

localization of many components of the pathway. Such additional regulatory mechanisms tune

WNT responses in diverse biological contexts, expand the functional repertoire of the pathway,

and may represent potential sites of therapeutic intervention in WNT-driven cancers. Classical

genetic approaches have been very successful at discovering new regulation in WNT signaling

[14]. In a previous study, we sought to uncover new regulatory mechanisms in WNT signaling

by performing forward genetic screens in HAP1-7TGP cells, a derivative of the haploid human

cell line HAP1 harboring a fluorescent reporter of WNT signaling [15]. HAP1 cells are

especially well suited for genetic screens due to the presence of a single allele of most genes in

their near-haploid genome, which can be disrupted by mutagenesis to generate true genetic null

cells [16]. We previously reported a comprehensive set of forward genetic screens designed to

identify positive, negative and attenuating regulators of WNT/CTNNB1 signaling, as well as

regulators of R-spondin (RSPO) signaling and suppressors of hyperactive WNT signaling

induced by loss of distinct DC components, including APC and CSNK1A1 [15]. These screens

recovered hits implicated at several levels of the pathway, including WNT and RSPO reception

at the plasma membrane, cytosolic signal transduction, and transcriptional regulation.

Comparative analyses of the screens enabled us to infer genetic interactions based on distinct

patterns of hits identified by the different screens. The screens for suppressors of hyperactive

signaling induced by loss of APC or CSNK1A1 suggested potential candidates for targeting

oncogenic WNT signaling.

An unexpected outcome of the *APC* and *CSNK1A1* suppressor screens was that we

observed only a partial overlap between significant hits in the two screens [15]. The phenotypic

selection parameters used in both screens were the same and the cell lines used for the two

screens were isogenic except for the mutations in *APC* or *CSNK1A1* we introduced by

CRISPR/Cas9-mediated genome editing. Therefore, we expected that the hits identified in the

two suppressor screens would be the same. After all, if APC and CSNK1A1 regulate

WNT/CTNNB1 signaling through a single common function in the DC phosphorylating

CTNNB1, we assumed that hyperactivating the pathway by knocking out one or the other would

be functionally equivalent, and the complement of downstream regulators would be shared.

While there were indeed many common hits with high significance scores in both suppressor

screens, including established downstream regulators of WNT/CTNNB1 signaling such as

CTNNB1 and *CREBBP*, there were also many hits unique to the *APC* suppressor or the

CSNK1A1 suppressor screen [15]. These results suggested that the hyperactive signaling state

resulting from loss of these two DC components was not equivalent. We hypothesized that the

difference in potential downstream regulators in the two genetic backgrounds in which the

screens were conducted – *APC* knock-out (KO) or *CSNK1A1* KO – could reflect additional roles

of APC or CSNK1A1 in WNT/CTNNB1 signaling beyond their shared function regulating

CTNNB1 stability.

HUWE1, the gene encoding the eponymous ubiquitin ligase, was the most striking

example of a hit that was highly significant in the *CSNK1A1* suppressor but not the *APC*

suppressor screen [15]. HUWE1 is a very large, 482 kilodalton (kDa) HECT domain ubiquitin

ligase that has been implicated in many cellular processes, including transcriptional regulation,

DNA replication and repair, cell cycle arrest, cell adhesion, cell migration, cell proliferation and

differentiation, proteotoxic stress, ribosome biogenesis, mitochondrial maintenance, autophagy,

apoptosis and WNT signaling [17-20]. *HUWE1* was the third most significant hit in the

CSNK1A1 suppressor screen, surpassed only by *CTNNB1* and *CREBBP*, which encode two of

the main components of the TCF/LEF transcription complex and are therefore central players in

the WNT pathway [15]. However, *HUWE1* was not a significant hit in the *APC* suppressor

screen (rank number 8040 out of 11022 genes with mapped gene-trap integrations), and it was

not among the most significant hits in any of the screens performed in wild-type (WT) HAP1-

7TGP cells, designed to identify positive regulators of WNT3A- and RSPO1-induced signaling.

These results suggested that HUWE1 might be involved in a regulatory mechanism that is most

evident in the CSNK1A1^{KO} genetic background (for brevity, HAP1-7TGP cell lines in which

genes were disrupted will be referred to by the name of the protein encoded by the targeted gene

or genes followed by a "KO" superscript). In follow-up studies, we had confirmed that HUWE1

loss reduced WNT target gene transcription – and to a smaller extent CTNNB1 abundance – in

CSNK1A1^{KO} but not in APC^{KO} cells [15]. We had also shown that microinjection of *HUWE1*

mRNA into *Xenopus laevis* embryos promoted body axis duplication, a hallmark of ectopic

WNT signaling [15]. These experiments established a few biological contexts in which HUWE1

acts as a positive regulator of WNT/CTNNB1 signaling, but the underlying mechanism remained

unclear and the reason why HUWE1 loss selectively reduced WNT/CTNNB1 signaling in

CSNK1A1^{KO} cells remained unknown.

Here we extend our genetic analyses to show that HUWE1 enhances WNT/CTNNB1

signaling through two different mechanisms. First, HUWE1 reduces phosphorylation of the

CTNNB1 phosphodegron by antagonizing the activity of a DC composed of GSK3A/GSK3B,

APC and AXIN1, therefore increasing CTNNB1 abundance. Second, HUWE1 enhances WNT

signaling through a mechanism that is independent from the control of CTNNB1 stability.

 Results:

HUWE1 enhances WNT signaling in CSNK1A1^{KO} cells by antagonizing GSK3A/GSK3B-

dependent phosphorylation of the CTNNB1 phosphodegron and increasing CTNNB1

abundance

We previously reported that HUWE1 loss in CSNK1A1^{KO} cells caused a substantial, 80-90%

reduction in WNT reporter activity and endogenous WNT target gene expression that was

accompanied by a smaller, 20-32% reduction in soluble CTNNB1 abundance [15]. Soluble

CTNNB1 is a proxy for the signaling CTNNB1 pool because it excludes the more stable, plasma

membrane-associated junctional CTNNB1 pool. We readily reproduced these results in the

current study: HUWE1 loss in CSNK1A1^{KO} cells reduced WNT reporter activity by 89% and

soluble CTNNB1 abundance by 36% (Figs 1A and 1B, and S1A Fig). These results raised the

possibility that in CSNK1A1^{KO} cells, HUWE1 loss reduces WNT signaling solely by reducing

CTNNB1 abundance, but that a non-linear relationship between changes in CTNNB1 abundance

and transcriptional activity results in a disproportionately greater reduction in WNT target gene

expression than CTNNB1 abundance. Alternatively, HUWE1 could regulate both CTNNB1

abundance and another process, which when disrupted together following HUWE1 loss result in

a greater reduction in WNT target gene expression than in CTNNB1 abundance. To distinguish

between these possibilities, we thought it was important to first determine the mechanism

underlying the reduction in CTNNB1 abundance caused by HUWE1 loss.

The main mechanism regulating CTNNB1 abundance is phosphorylation of the CTNNB1

phosphodegron by the DC [5]. In CSNK1A1^{KO} cells we did not expect the phosphodegron to be

phosphorylated by GSK3A/GSK3B at residues S33, S37 and T41 because phosphorylation of

these residues generally requires the priming phosphorylation of residue S45 by CSNK1A1 [6,

7]. Nevertheless, we tested whether the reduction in CTNNB1 abundance caused by HUWE1

loss in CSNK1A1^{KO} cells was due to changes in phosphorylation of the CTNNB1

phosphodegron. CTNNB1 phosphorylated at S33, S37 and T41 can be measured directly by

Western blot, but due to the rapid proteasomal degradation of this species, treatment with

proteasome inhibitors is usually required to make such measurements [7]. Since any effects of

HUWE1 on WNT signaling could conceivably also depend on proteasomal degradation of a

HUWE1 substrate, which would be disrupted by proteasome inhibitors, we opted for a different

way to evaluate phosphorylation of the CTNNB1 phosphodegron. We instead quantified

CTNNB1 that is *not* phosphorylated at residues S33, S37 and T41 (we refer to this species as

non-phospho-CTNNB1, but it is also known as active CTNNB1 [21]) from whole cell extracts

(WCE) (Figs 1C and S1B Fig). As a control, we also measured total CTNNB1 from WCE (S1B

and S1C Figs). Non-pospho-CTNNB1 abundance in the various conditions tested was different

from and exhibited larger changes than total CTNNB1 abundance (Fig 1C, and S1B and S1C

Figs). This indicated that non-phospho-CTNNB1 only represents a fraction of the total CTNNB1

in WCE and is likely to accurately reflect changes in phosphorylation of the CTNNB1

phosphodegron.

HUWE1 loss in CSNK1A1^{KO} cells reduced non-phospho-CTNNB1 abundance by 37%, a

reduction that correlated closely with the 36% reduction in soluble CTNNB1 abundance caused

by HUWE1 loss in the same cell line (Figs 1B and 1C, and S1A and S1B Figs). This correlation

suggested that the reduction in CTNNB1 abundance caused by HUWE1 loss was due to

increased CTNNB1 phosphorylation at S33, S37 and T41, presumably mediated by

GSK3A/GSK3B. If this were the case, inhibiting GSK3A/GSK3B should reverse the reduction

in both soluble and non-phospho-CTNNB1 abundance caused by HUWE1 loss. Treatment of

CSNK1A1^{KO}; HUWE1^{KO} cells with the GSK3A/GSK3B inhibitor CHIR-99021 indeed increased

the abundance of soluble CTNNB1 by 2.4-fold and the abundance of non-phospho-CTNNB1 by

3.2-fold (Figs 1B and 1C, and S1A and S1B Figs), entirely reversing the reductions caused by

HUWE1 loss. Furthermore, GSK3A/GSK3B inhibition in CSNK1A1^{KO}; HUWE1^{KO} cells

increased WNT reporter activity by 10.9-fold, restoring signaling to a comparable level to that in

DMSO vehicle-treated CSNK1A1^{KO} cells (Fig 1A). These results indicate that even in the

absence of CSNK1A1, phosphorylation of residues S33, S37 and T41 by GSK3A/GSK3B can

regulate CTNNB1 abundance, and that HUWE1 loss reduces CTNNB1 abundance and WNT

signaling by promoting the phosphorylation of these residues.

Since HUWE1 loss in CSNK1A1^{KO} cells increased GSK3A/GSK3B-dependent

phosphorylation of the CTNNB1 phosphodegron, we wondered whether in CSNK1A1^{KO} cells

containing HUWE1, residues S33, S37 and T41 in the phosphodegron might be partially

phosphorylated by GSK3A/GSK3B despite the absence of CSNK1A1. CSNK1A1^{KO} cells had a

relatively high abundance of soluble and non-phospho-CTNNB1, as well as high WNT reporter

activity, compared to basal levels in unstimulated WT HAP1-7TGP cells (Figs 1A-C, and S1A

and S1B Figs). However, GSK3A/GSK3B inhibition with CHIR-99021 in CSNK1A1^{KO} cells

increased the abundance of soluble CTNNB1 by 1.6-fold and the abundance of non-phospho-

CTNNB1 by 1.8-fold (Figs 1B and 1C, and S1A and S1B Figs). WNT reporter activity also

increased 1.7-fold following treatment of CSNK1A1^{KO} cells with CHIR-99021 (Fig 1A).

Therefore, in the absence of CSNK1A1, residual GSK3A/GSK3B-dependent phosphorylation of

the CTNNB1 phosphodegron can still take place. This is presumably followed by ubiquitylation

and proteasomal degradation of phosphorylated CTNNB1.

In summary, in CSNK1A1^{KO} cells, CTNNB1 is still phosphorylated by GSK3A/GSK3B at

residues S33, S37 and S41 in the CTNNB1 phosphodegron, and the reduction in soluble

CTNNB1 abundance caused by HUWE1 loss is due to increased GSK3A/GSK3B-dependent

phosphorylation of these residues. We conclude that when present, HUWE1 antagonizes the

GSK3A/GSK3B-dependent phosphorylation and ensuing degradation of CTNNB1, thereby

increasing CTNNB1 abundance and promoting WNT signaling.

Our results raise two important questions. First, is control of CTNNB1 phosphorylation and

abundance the only mechanism whereby HUWE1 enhances WNT signaling, or is there another

mechanism distinct from the control of CTNNB1 stability? Second, is the GSK3A/GSK3B-

dependent regulation of CTNNB1 abundance by HUWE1, and potentially any other mechanisms

by which HUWE1 enhances WNT signaling, also mediated by other components of the DC in

addition to GSK3A/GSK3B? We addressed both these questions.

HUWE1 enhances WNT signaling through a mechanism independent of CTNNB1 stability

We next sought to determine if HUWE1 could promote WNT signaling through additional

mechanisms distinct from control of CTNNB1 phosphorylation and abundance. Mutations in the

CTNNB1 phosphodegron that prevent phosphorylation by CSNK1A1 and GSK3A/GSK3B

render CTNNB1 insensitive to degradation by the DC [6, 7, 22]. We reasoned that introducing

such mutations into the single *CTNNB1* allele of HAP1-7TGP cells would enable us to decouple

control of CTNNB1 abundance from any other mechanism by which HUWE1 enhances WNT

signaling.

We used CRISPR/Cas9-induced homology directed repair (HDR) to edit the codons

encoding CSNK1A1 and GSK3A/GSK3B phosphorylation sites in the phosphodegron of the

single endogenous *CTNNB1* locus in HAP1-7TGP cells. We introduced mutations encoding

alanine (A) substitutions in the codon encoding S45, which is phosphorylated by CSNK1A1, and

in the codons encoding T41 and S37, which are sequentially phosphorylated by GSK3A/GSK3B

(S1 File and S2A Fig). We were unable to mutate S33, the third GSK3A/GSK3B target site.

However, recognition of CTNNB1 by $SCF^{\beta TrCP}$ requires phosphorylation of both S33 and S37 [9,

10], and therefore the mutations we introduced still prevented DC-dependent CTNNB1

degradation, as we demonstrate below. We called the resulting HAP1-7TGP derivative cell line

$CTNNB1^{ST-A}$. The mutations in the *CTNNB1* locus of $CTNNB1^{ST-A}$ cells indeed increased

soluble CTNNB1 abundance 42-fold compared to unstimulated WT HAP1-7TGP cells (Fig 2A

and S2B Fig), and promoted constitutive WNT signaling as judged by WNT reporter activity and

endogenous WNT target gene (*AXIN2 [23]*, *RNF43 [23]*, *NKD1 [24]*, *TNFRSF19 [25]*)

expression (Figs 2B-F). Furthermore, CTNNB1 abundance, WNT reporter activity and WNT

target gene expression in $CTNNB1^{ST-A}$ cells were substantially higher than in WT HAP1-7TGP

cells treated with a near-saturating dose of WNT3A conditioned media (CM) (Figs 2A-F, and

S2B Fig). Stimulation of $CTNNB1^{ST-A}$ cells with WNT3A CM did not significantly increase

total CTNNB1 abundance or WNT target gene expression (S2C-E Figs). These results confirmed

that the mutations we introduced into $CTNNB1^{ST-A}$ cells rendered CTNNB1 insensitive to

degradation by the DC, and therefore abolished the control of CTNNB1 stability by WNT

ligands.

We then knocked out *HUWE1* in CTNNB1^{ST-A} cells (S1 File and S2B Fig) and measured the

effect on CTNNB1 abundance and WNT signaling. HUWE1 loss in multiple independent clonal

cell lines (CTNNB1^{ST-A}; HUWE1^{KO}) did not affect soluble CTNNB1 abundance (Fig 2A and

S2B Fig), but significantly reduced WNT reporter activity (Fig 2B and S2F Fig) and the

expression of some WNT target genes (Figs 2C-F). These results demonstrate that HUWE1 loss

reduces WNT signaling in part through a mechanism independent from the control of CTNNB1

stability. We also note that the 49% reduction in WNT reporter activity, 45% reduction in *AXIN2*

expression and 31% reduction in *RNF43* expression caused by HUWE1 loss in CTNNB1^{ST-A}

cells (Figs 2B-D and S2F Fig) were smaller than the 89% reduction in WNT reporter activity,

67% reduction in *AXIN2* expression and 73% reduction in *RNF43* expression caused by HUWE1

loss in CSNK1A1^{KO} cells (Figs 1A and 3C-D). This difference could be because in CSNK1A1^{KO}

cells, HUWE1 loss caused a 31-36% reduction in soluble CTNNB1 abundance (Figs 1B and 3A-

B, and S1A Fig) in addition to the reduction in signaling caused by the second regulatory

mechanism that is independent from changes in CTNNB1 abundance, whereas no corresponding

reduction in CTNNB1 abundance was observed following HUWE1 loss in CTNNB1^{ST-A} cells

(Fig 2A and S2B Fig).

In summary, we distinguished two mechanisms whereby HUWE1 loss reduces WNT

signaling. In CSNK1A1^{KO} cells containing WT CTNNB1, HUWE1 loss caused a moderate

reduction in CTNNB1 abundance and a comparable increase in GSK3A/GSK3B-dependent

phosphorylation of the CTNNB1 phosphodegron, as well as a much larger GSK3A/GSK3B-

dependent reduction in WNT reporter activity (Fig 1). In CTNNB1^{ST-A} cells containing WT

CSNK1A1 but a mutated CTNNB1 phosphodegron, HUWE1 loss did not alter CTNNB1

abundance but still caused a significant reduction in WNT reporter activity and WNT target gene

294 expression (Fig 2). We conclude that HUWE1 enhances WNT signaling through two distinct

295 mechanisms, one that increases CTNNB1 abundance and one that is independent of CTNNB1

296 stability.

298 **HUWE1 enhances WNT signaling through mechanisms mediated by APC**

300 Having defined two mechanisms whereby HUWE1 regulates WNT signaling, a

301 GSK3A/GSK3B-dependent mechanism that controls CTNNB1 phosphorylation and abundance,

302 and another mechanism that is independent from the control of CTNNB1 stability, we wondered

303 whether these mechanisms were also mediated by other DC components. HUWE1 was one of

304 the most significant hits in a *CSNK1A1* suppressor screen but was not a significant hit in an *APC*

305 suppressor screen [15]. Consistent with the results of these screens, HUWE1 loss substantially

306 reduced WNT reporter activity in CSNK1A1^{KO} cells but did not affect WNT reporter activity in

307 APC^{KO} cells [15]. Based on these results, we hypothesized that APC may be required to mediate

308 the effects of HUWE1 on WNT signaling. If APC is required for the reduction in WNT signaling

309 caused by HUWE1 loss in CSNK1A1^{KO} cells, then eliminating APC function in CSNK1A1^{KO};

310 HUWE1^{KO} cells, like inhibiting GSK3A/GSK3B activity (Fig 1), should reverse said reduction.

311 To the best of our knowledge, there are no pharmacological inhibitors that we could use to

312 acutely inhibit APC. We were also unable to knock out *APC* in CSNK1A1^{KO}; HUWE1^{KO} cells,

313 as we found that knocking out additional genes by CRISPR/Cas9-mediated genome editing in

314 this cell line yielded very few viable clones. Instead, we first made cell lines lacking both APC

315 and CSNK1A1, and then tested the effects of HUWE1 loss in these cells, comparing them to

316 cells lacking CSNK1A1 alone.

CSNK1A1 single KO clonal HAP1-7TGP cell lines were generated and characterized

previously [15] (CSNK1A1^{KO-1} and CSNK1A1^{KO-2}; we note that CSNK1A1^{KO-2} is a loss-of-

function allele containing a two amino acid deletion). We generated two new *APC* single KO

clonal HAP1-7TGP cell lines (APC^{KO-1} and APC^{KO-2}) as well as two new *APC* and *CSNK1A1*

double KO clonal HAP1-7TGP cell lines (APC^{KO-1}; CSNK1A1^{KO-1} and APC^{KO-1}; CSNK1A1^{KO-}

^{2}) using CRISPR/Cas9-mediated genome editing. We validated these cell lines by sequencing

each targeted locus (S1 File), and by Western blot analysis (Fig 3A). CSNK1A1^{KO}, APC^{KO} and

APC^{KO}; CSNK1A1^{KO} cells all exhibited elevated soluble CTNNB1 abundance several-fold

higher than unstimulated WT HAP1-7TGP cells (Figs 3A and 3B). All these clonal cell lines

exhibited constitutive expression of WNT target genes several-fold higher than the level of gene

expression in unstimulated WT HAP1-7TGP cells and in WT HAP1-7TGP cells stimulated with

a near-saturating dose of WNT3A CM (Figs 3C-F). Consistent with our results demonstrating

that in CSNK1A1^{KO} cells residual phosphorylation of the CTNNB1 phosphodegron by

GSK3A/GSK3B results in some CTNNB1 degradation (Fig 1), both soluble CTNNB1

abundance and WNT target gene expression were higher in APC^{KO} and APC^{KO}; CSNK1A1^{KO}

cells than in CSNK1A1^{KO} cells (Figs 3A-F). These results support the notion that in HAP1 cells

CSNK1A1 is partially dispensable for CTNNB1 phosphorylation by GSK3A/GSK3B.

We then knocked out *HUWE1* in CSNK1A1^{KO}, APC^{KO} and APC^{KO}; CSNK1A1^{KO} cells to

generate three CSNK1A1^{KO}; HUWE1^{KO}, three APC^{KO}; HUWE1^{KO} and three APC^{KO};

CSNK1A1^{KO}; HUWE1^{KO} clonal cell lines, which we validated by sequencing the targeted

HUWE1 locus (S1 File) and by Western blot analysis (Fig 3A). HUWE1 loss in CSNK1A1^{KO}

cells substantially reduced the expression of all WNT target genes tested (Figs 3C-F) and, to a

lesser extent, soluble CTNNB1 abundance (Figs 3A and 3B). In contrast, HUWE1 loss in APC^{KO}

cells resulted in a variable but not statistically significant reduction in WNT target gene

expression (Figs 3C-F) and did not reduce soluble CTNNB1 abundance (Figs 3A and 3B),

consistent with our previous finding that HUWE1 loss in APC[KO] cells had no effect on WNT

reporter activity [15]. Finally, HUWE1 loss in APC[KO]; CSNK1A1[KO] cells yielded equivalent

results to those in APC[KO] cells, showing no significant reduction in WNT target gene expression

(Figs 3C-F) or soluble CTNNB1 abundance (Figs 3A and 3B). These results indicate that, like

GSK3A/GSK3B inhibition, APC loss precludes the reduction in WNT target gene expression

and CTNNB1 abundance caused by HUWE1 loss in CSNK1A1[KO] cells. We conclude that APC

mediates the effects of HUWE1 on WNT signaling.

HUWE1 enhances WNT signaling through mechanisms mediated by a subset of DC

components including APC, AXIN1 and GSK3A or GSK3B

We extended the same logic as for APC (Fig 3) to test the role of every core component of the

DC in mediating the functions of HUWE1 in WNT signaling. We first knocked out components

of the DC individually or in certain combinations (Table 1) so we could then test the effects of

HUWE1 loss on WNT signaling in each of these mutant genetic backgrounds. While we had

already established the role of GSK3A/GSK3B and APC in mediating HUWE1 functions (Figs 1

and 3), we included them in our analysis to confirm those results and, in the case of GSK3A and

GSK3B, test their roles individually. We used CRISPR/Cas9-mediated genome editing to

generate HAP1-7TGP clonal cell lines lacking the desired DC components (Table 1). We

confirmed that each targeted genomic locus had been successfully mutated (S1 File) and that the

encoded protein had been eliminated (S4A Fig).

Table 1. HUWE1 enhances WNT signaling through mechanisms mediated by a subset of DC components including APC, AXIN1 and GSK3A or GSK3B.

Genotype	Aggregate WNT target gene expression, average of HUWE1 sgRNA1 and sgRNA2 (% of SCR sgRNA control)	Significance of change in aggregate WNT target gene expression for HUWE1 sgRNA 1/2 relative to SCR sgRNA control
WT - WNT3A	101	n.s./n.s.
WT + WNT3A	70	*/*
$CSNK1A1^{KO}$	47.5	**/***
APC^{KO}	86.5	n.s./n.s.
APC^{KO}; $CSNK1A1^{KO}$	86	n.s./n.s.
$AXIN1^{KO}$; $AXIN2^{KO}$	91.5	n.s./n.s.
$CSNK1A1^{KO}$; $AXIN1^{KO}$; $AXIN2^{KO}$	103	n.s./n.s.
$CSNK1A1^{KO}$; $AXIN1^{KO}$	95	n.s./n.s.
$CSNK1A1^{KO}$; $AXIN2^{KO}$	45.5	**/**
$GSK3A^{KO}$; $GSK3B^{KO}$	88	n.s./n.s.
$CSNK1A1^{KO}$; $GSK3A^{KO}$; $GSK3B^{KO}$	141.5	*/n.s.
$CSNK1A1^{KO}$; $GSK3A^{KO}$	53	**/*
$CSNK1A1^{KO}$; $GSK3B^{KO}$	40	**/**

Summary of effects of CRISPRi-mediated HUWE1 KD on aggregate WNT target gene expression (Fig 4A). For the genotypes treatments indicated in green, both HUWE1 sgRNAs used resulted in a significant reduction in aggregate WNT target gene expression relative to the SCR sgRNA control.

    To test the effects of HUWE1 loss in each of these mutant cell lines, we adopted a different experimental strategy. We had previously quantified the effect of HUWE1 loss on WNT signaling through an experimental scheme that we refer to as clonal analysis. In this scheme, we used CRISPR/Cas9-mediated genome editing to target *HUWE1*. We isolated multiple independent clonal cell lines in which *HUWE1* had been knocked out, and multiple clonal cell lines that remained WT at the targeted locus to use as controls. We then compared several KO

and WT clones for WNT reporter activity or other parameters of interest (Figs 2 and 3, and S2F

Fig). While clonal analysis enables comparisons in true genetic null conditions, it is subject to

substantial inter-clonal variability, requiring the laborious isolation of many independent clones

to achieve statistical significance. Isolation of multiple clones harboring *HUWE1* mutations in

each of the 12 different genetic backgrounds (Table 1) in which we wanted to test the effect of

HUWE1 loss was unfeasible. Therefore, we implemented a CRISPR interference (CRISPRi)-

mediated knock-down (KD) strategy [26] that enabled us to measure the outcome of knocking

down HUWE1 in polyclonal cell populations rather than in multiple individual clonal cell lines.

We used a lentivirus to deliver the CRISPRi machinery together with sgRNAs targeting *HUWE1*

in the various cell lines we had generated lacking DC components (Table 1 and S4A Fig). Based

on Western blot measurements (S5A and S5B Figs), lentiviral delivery of either of two different

sgRNAs targeting *HUWE1* (HUWE1 sgRNA1 or sgRNA2) followed by antibiotic selection of

transduced cells resulted in a consistent 59-95% KD of HUWE1 compared to a control,

scrambled (SCR) sgRNA. We refer to polyclonal cell populations in which we knocked down

HUWE1 using CRISPRi as HUWE1^{KD}, in contrast to HUWE1^{KO} clonal cell lines in which we

knocked out *HUWE1* using CRISPR/Cas9-mediated genome editing.

To validate the CRISPRi KD strategy, we tested whether knocking down HUWE1 in

CSNK1A1^{KO}, APC^{KO}, and APC^{KO}; CSNK1A1^{KO} cell populations produced equivalent results to

those we had observed when we knocked out HUWE1 and conducted clonal analysis in these

same cell lines (Fig 3). Consistent with our clonal analysis, HUWE1 KD in CSNK1A1^{KO} cells

significantly reduced the expression of four WNT target genes compared to CSNK1A1^{KO} cells

transduced with SCR sgRNA (Figs 4A-E and Table 1). However, this reduction was smaller than

that caused by complete HUWE1 loss in CSNK1A1^{KO}; HUWE1^{KO} cells (Figs 3C-F),

presumably owing to some residual HUWE1 protein present in CSNK1A1^{KO}; HUWE1^{KD} cells

(S5A and S5B Figs). Also consistent with our clonal analysis, HUWE1 KD in APC^{KO} and in

APC^{KO}; CSNK1A1^{KO} cells did not cause a statistically significant reduction in WNT target gene

expression (Figs 4A-E and Table 1). These results validate CRISPRi-mediated HUWE1 KD in

polyclonal cell populations as a reliable alternative to the more laborious clonal analysis of

multiple individual HUWE1^{KO} clonal cell lines. We also knocked down HUWE1 in WT HAP1-

7TGP cells (S5A and S5B Figs), in which we had previously reported that HUWE1 KO did not

cause a significant reduction in WNT reporter activity or *AXIN2* expression induced by a near-

saturating dose of WNT3A [15]. In agreement with those results, HUWE1 KD did not reduce

WNT3A-induced expression of *AXIN2* (Fig 4B). However, HUWE1 KD in WT HAP1-7TGP

cells did reduce the expression of other WNT target genes, including *RNF43*, *NKD1* and

TNFRSF19, but to a smaller extent than in CSNK1A1^{KO} cells (Figs 4A-E and Table 1). Together

with our analysis in CTNNB1^{ST-A} cells (Fig 2), these results demonstrate that the contribution of

HUWE1 to WNT signaling is not limited to cells lacking CSNK1A1.

We then asked whether other DC components mediate the function of HUWE1. As we had

done for APC, we tested whether knocking out AXIN1 and AXIN2 eliminated the reduction in

WNT signaling caused by HUWE1 loss. In WT HAP1-7TGP cells, AXIN1 and AXIN2 are

functionally redundant in their capacity to suppress WNT signaling, presumably by regulating

CTNNB1 abundance as scaffolds in the DC: eliminating either AXIN1 or AXIN2 has no effect

on WNT reporter activity, whereas eliminating both promotes constitutive pathway activation

[15]. We initially assumed that a possible role of AXIN1 and AXIN2 in mediating the effects of

HUWE1 may also be redundant, so we knocked out both paralogs in HAP1-7TGP cells (S1 File

and S4A Fig) and tested their contribution following HUWE1 KD (S5A and S5B Figs). HUWE1

KD in AXIN1^{KO}; AXIN2^{KO} cells did not reduce WNT target gene expression (Figs 4A-E and

Table 1), suggesting that AXIN1, AXIN2 or both mediate the effects of HUWE1 on WNT

signaling, similarly to what we had observed for APC (Fig 3). As we had done for

GSK3A/GSK3B (Fig 1) and APC (Fig 3), we also tested whether the combined loss of AXIN1

and AXIN2 eliminated the reduction in WNT signaling caused by HUWE1 KD in CSNK1A1^{KO}

cells. Indeed, knocking down HUWE1 in CSNK1A1^{KO}; AXIN1^{KO}; AXIN2^{KO} cells did not

reduce WNT target gene expression as it did in CSNK1A1^{KO} cells (Figs 4A-E and Table 1).

These results confirmed that AXIN1, AXIN2 or both mediate the effects of HUWE1 on WNT

signaling in CSNK1A1^{KO} cells. While AXIN1 and AXIN2 are redundant in their capacity to

suppress WNT signaling in WT HAP1-7TGP cells [15], it was conceivable that they may not be

redundant in mediating the function of HUWE1 in CSNK1A1^{KO} cells. To test for individual

contributions of AXIN1 or AXIN2, we knocked each of them out individually in CSNK1A1^{KO}

cells (S1 File and S4A Fig) and then knocked down HUWE1 (S5A and S5B Figs). AXIN1 loss

in CSNK1A1^{KO} cells eliminated the reduction in WNT signaling caused by HUWE1 KD, but

AXIN2 loss did not (Figs 4A-E and Table 1). These results suggest that, in contrast to its

redundant function with AXIN2 in suppressing WNT signaling in WT HAP1-7TGP cells [15],

AXIN1 plays a unique role in mediating HUWE1-dependent effects on WNT signaling that is

not redundant with AXIN2.

Given these results, we wondered whether GSK3A and GSK3B are redundant in mediating

the functions of HUWE1 in WNT signaling. To answer this question, we did an equivalent series

of experiments as the one we did to determine the individual roles of AXIN1 and AXIN2 in

mediating HUWE1 function. Like AXIN1 and AXIN2, GSK3A and GSK3B are functionally

redundant in their capacity to suppress WNT signaling in WT HAP1-7TGP cells: eliminating

either GSK3A or GSK3B has no effect on WNT reporter activity, whereas eliminating both

promotes constitutive pathway activation (S1 File, and S4B and S4C Figs). HUWE1 KD in

GSK3A^{KO}; GSK3B^{KO} cells (S5A and S5B Figs) did not reduce WNT target gene expression

(Figs 4A-E and Table 1), suggesting that GSK3A, GSK3B or both mediate the effects of

HUWE1 on WNT signaling. Next, we tested whether the combined loss of GSK3A and GSK3B

eliminated the reduction in WNT signaling caused by HUWE1 KD in CSNK1A1^{KO} cells.

Knocking down HUWE1 in CSNK1A1^{KO}; GSK3A^{KO}; GSK3B^{KO} cells (S5A and S5B Figs) did

not reduce – and in fact increased – WNT target gene expression (Figs 4A-E and Table 1). These

results confirmed that GSK3A, GSK3B or both mediate the effects of HUWE1 on WNT

signaling in CSNK1A1^{KO} cells. However, unlike their combined loss, loss of GSK3A or GSK3B

individually in CSNK1A1^{KO} cells did not eliminate the reduction in WNT signaling caused by

HUWE1 KD (Fig 4A-E and Table 1). We conclude that the role of GSK3A and GSK3B in

mediating HUWE1-dependent effects on WNT signaling is redundant, similarly to their role

suppressing WNT signaling in WT HAP1-7TGP cells (S4C Fig). Therefore, only the combined

loss of GSK3A and GSK3B eliminates the reduction in WNT signaling caused by HUWE1 KD

in CSNK1A1^{KO} cells (Fig 4A-E and Table 1).

We considered the possibility that the distinct outcomes of knocking down HUWE1 in the

various genetic backgrounds we tested (Table 1) could be due to differences in the steady state

abundance of HUWE1 caused by loss of some DC complex components but not others, rather

than due to other effects of distinct DC components in mediating HUWE1 function. Standard

Western blot analysis did not reveal obvious differences in steady state HUWE1 abundance

among the various genetic backgrounds in which we knocked down HUWE1 (S4A Fig). We

corroborated this result by quantitative dot blot analysis (see Materials and methods) and did not

detect significant differences in HUWE1 abundance among the different genetic backgrounds

(S4D Fig).

In conclusion, a subset of DC components, including APC, AXIN1 and GSK3A or GSK3B,

but not CSNK1A1 or AXIN2, mediates the function of HUWE1 in WNT signaling. Since

HUWE1 enhances WNT signaling by increasing CTNNB1 abundance (Fig 1) and through

another mechanism independent from the control of CTNNB1 stability (Fig 2), a DC composed

of APC, AXIN1 and GSK3A/GSK3B must mediate the effects of HUWE1 on one or both

mechanisms.

HUWE1 enhances WNT signaling by antagonizing the DC

The results presented so far are consistent with the following hypothesis: 1. In CSNK1A1^{KO}

cells, APC, AXIN1 and GSK3A/GSK3B are part of a DC that can partially suppress WNT

signaling by phosphorylating the CTNNB1 phosphodegron and targeting CTNNB1 for

proteasomal degradation; 2. HUWE1 enhances WNT signaling by antagonizing CTNNB1

phosphorylation and degradation mediated by this DC, and through another mechanism

independent of CTNNB1 stability. Whether the second mechanism is also mediated by the DC

remains unclear. Since all the data presented above were from loss-of-function genetic

experiments, we tested this hypothesis further through overexpression experiments. Based on

this hypothesis, we predicted that overexpressing the DC scaffold AXIN1 in CSNK1A1^{KO} cells

should increase DC activity and therefore have similar effects as knocking out HUWE1: it

should reduce WNT signaling by promoting GSK3A/GSK3B-dependent phosphorylation and

degradation of CTNNB1, and possibly by promoting the second mechanism independent of

CTNNB1 stability. Furthermore, since AXIN1 loss in CSNK1A1^{KO} cells eliminated the

reduction in WNT signaling caused by HUWE1 loss (Figs 4A-E and Table 1), we reasoned that

overexpressing AXIN1 in CSNK1A1^{KO}; HUWE1^{KO} cells should have the opposite effect and

synergize with HUWE1 loss to reduce WNT signaling. To test these predictions, we stably

overexpressed human AXIN1 in CSNK1A1^{KO} and in CSNK1A1^{KO}; HUWE1^{KO} cells through

lentiviral delivery of *AXIN1* cDNA followed by antibiotic selection. We obtained polyclonal cell

populations (CSNK1A1^{KO}; AXIN1^{OE} and CSNK1A1^{KO}; HUWE1^{KO}; AXIN1^{OE}, respectively) in

which AXIN1 abundance was at least 2-fold higher than that in the respective parental cell lines

(S1D Fig).

AXIN1 overexpression in CSNK1A1^{KO} cells indeed reduced WNT reporter activity by 80%,

which was comparable to the 89% reduction caused by HUWE1 loss in CSNK1A1^{KO} cells (Fig

1A). AXIN1 overexpression combined with HUWE1 loss in CSNK1A1^{KO} cells reduced WNT

reporter activity by 98%, nearly down to the basal level of unstimulated WT HAP1-7TGP cells

(Fig 1A). Therefore, AXIN1 overexpression in CSNK1A1^{KO} cells phenocopied HUWE1 loss,

and AXIN1 overexpression in CSNK1A1^{KO}; HUWE1^{KO} cells synergized with HUWE1 loss to

reduce WNT signaling. We conclude that HUWE1 and AXIN1 exert opposing effects on WNT

signaling.

To test whether the reduction in WNT signaling caused by AXIN1 overexpression and by

HUWE1 loss was due to the same underlying mechanisms, we measured the abundance of

soluble and non-phospho-CTNNB1 in CSNK1A1^{KO}; AXIN1^{OE} and CSNK1A1^{KO}; HUWE1^{KO};

AXIN1^{OE} cells, as we had done in WT HAP1-7TGP, CSNK1A1^{KO} and CSNK1A1^{KO};

HUWE1^{KO} cells (Figs 1B and 1C, and S1A and S1B Figs). AXIN1 overexpression in

CSNK1A1^{KO} cells caused a 45% reduction in soluble CTNNB1 abundance and a 64% reduction

in non-phospho-CTNNB1 abundance (Figs 1B and 1C, and S1A and S1B Figs). These

reductions were comparable to and greater than the respective 36% and 37% reductions caused

by HUWE1 loss in CSNK1A1^{KO} cells (Figs 1B and 1C, and S1A and S1B Figs). AXIN1

overexpression combined with HUWE1 loss in CSNK1A1^{KO} cells reduced soluble CTNNB1

abundance by 57% and non-phospho-CTNNB1 by 62% (Figs 1B and 1C, and S1A and S1B

Figs). These results indicate that HUWE1 and AXIN1 have opposing functions regulating a

common mechanism: AXIN1 promotes CTNNB1 phosphodegron phosphorylation and the

resulting reduction in CTNNB1 abundance, while HUWE1 antagonizes both.

If HUWE1 and AXIN1 exert opposing effects on WNT signaling by regulating the same

GSK3A/GSK3B-dependent processes – CTNNB1 phosphorylation and abundance, and

potentially another mechanism independent of CTNNB1 stability – then GSK3A/GSK3B

inhibition should reverse the effects of AXIN1 overexpression in CSNK1A1^{KO} cells, as it

reverses the effects of HUWE1 loss (Figs 1A-C, and S1A and S1B Figs). Therefore, we tested

whether the changes in WNT reporter activity, soluble CTNNB1 abundance and CTNNB1

phosphodegron phosphorylation caused by AXIN1 overexpression alone or combined with

HUWE1 loss were dependent on GSK3A/GSK3B activity. Treatment of CSNK1A1^{KO}; AXIN1^{OE}

cells with the GSK3A/GSK3B inhibitor CHIR-99021 reversed the effects of AXIN1

overexpression, increasing WNT reporter activity as well as soluble and non-phospho-CTNNB1

abundance to levels higher than those measured in DMSO vehicle treated-CSNK1A1^{KO} cells,

and comparable to those measured in CHIR-99021-treated CSNK1A1^{KO} cells (Figs 1A-C, and

S1A and S1B Figs). GSK3A/GSK3B inhibition with CHIR-99021 also reversed the synergistic

reduction in WNT reporter activity, as well as the reduction in soluble and non-phospho-

CTNNB1 abundance, caused by combined AXIN1 overexpression and HUWE1 loss in

CSNK1A1^{KO} cells (Figs 1A-C, and S1A and S1B Figs). These results demonstrate that in

CSNK1A1^{KO} cells, HUWE1 enhances and AXIN1 inhibits WNT signaling by opposing

mechanisms mediated by GSK3A/GSK3B. Altogether, our results support the hypothesis that

AXIN1, acting as a scaffold in the DC, promotes GSK3A/GSK3B-dependent CTNNB1

phosphorylation and degradation, even in the absence of CSNK1A1. HUWE1 enhances WNT

signaling by antagonizing this DC activity.

Regulation of WNT signaling by HUWE1 requires its ubiquitin ligase activity

HUWE1 is a very large 482 kDa ubiquitin ligase with many protein-protein interaction domains

in addition to its catalytic HECT domain [27]. Therefore, it was important to determine whether

the ubiquitin ligase activity of HUWE1 was required for its functions enhancing WNT signaling.

HECT domain ubiquitin ligases form a covalent intermediate between a catalytic cysteine (C)

residue in the HECT domain and ubiquitin before ubiquitin is transferred to the substrate [28].

We used CRISPR-mediated base editing [29] to engineer the endogenous *HUWE1* locus of

CSNK1A1^{KO} cells, introducing a single point mutation that replaced the catalytic C4341 residue

with arginine (R). We isolated three independent clonal cell lines (CSNK1A1^{KO}; HUWE1$^{C4341R-}$

^{1}, CSNK1A1^{KO}; HUWE1$^{C4341R-2}$ and CSNK1A1^{KO}; HUWE1$^{C4341R-3}$) in which we confirmed by

sequencing that the intended point mutation had been introduced (S1 File). We compared the

effects of eliminating the catalytic activity of HUWE1 to those of knocking out HUWE1 on

WNT signaling. All three CSNK1A1^{KO}; HUWE1^{C4341R} clonal cell lines exhibited a substantial

89-94% reduction in WNT reporter activity and a 79-88% reduction in the expression of three

WNT target genes, equivalent to what we observed in CSNK1A1^{KO}; HUWE1^{KO} cells (Figs 5A-

D). The C4341R point mutation did not affect HUWE1 protein stability as determined by

Western blot analysis of the three CSNK1A1^{KO}; HUWE1^{C4341R} cell lines (Fig 5E). In contrast, no

HUWE1 protein was detected in CSNK1A1^{KO}; HUWE1^{KO} cells (Fig 5E). Therefore, the

reduction in WNT signaling measured in CSNK1A1^{KO}; HUWE1^{C4341R} cells was not due to loss

of HUWE1 protein, but rather due to the elimination of its catalytic activity. We conclude that

the ubiquitin ligase activity of HUWE1 is required for its functions enhancing WNT signaling.

Discussion:

In this study we probed the mechanisms underlying the requirement for the HECT domain ubiquitin ligase HUWE1 to sustain hyperactive WNT/CTNNB1 signaling [15]. We demonstrate that HUWE1 enhances WNT/CTNNB1 signaling through two distinct mechanisms: by antagonizing DC-mediated CTNNB1 phosphorylation and degradation, and through another mechanism independent of CTNNB1 stability. These results are significant for two main reasons. First, they reveal a new mechanism that controls CTNNB1 stability, the main regulated step in WNT/CTNNB1 signaling. Second, by controlling another downstream step in the pathway, HUWE1 adds a new layer of regulation superimposed on the core WNT/CTNNB1 signaling module. Importantly, the coordinated regulation of CTNNB1 abundance and an independent signaling step in the pathway by HUWE1 would be an efficient way to control multiple processes that determine WNT signaling output. This may enable sensitive and robust activation of the pathway.

In CSNK1A1^{KO} cells, GSK3A/GSK3B still phosphorylate a fraction of CTNNB1 at residues S33, S37 and T41 in the phosphodegron, which reduces CTNNB1 abundance (Fig 1). HUWE1 enhances signaling by counteracting DC-dependent phosphorylation of these residues, since HUWE1 loss in CSNK1A1^{KO} cells increases phosphorylation and reduces both CTNNB1 abundance and WNT signaling activity (Fig 1). However, the reduction in CTNNB1 abundance caused by HUWE1 loss appears insufficient to account for the larger reduction in WNT target gene expression (Fig 1), suggesting that HUWE1 also enhances WNT/CTNNB1 signaling through another mechanism. In CTNNB1^{ST-A} cells containing mutations in the CTNNB1 phosphodegron that render CTNNB1 abundance insensitive to regulation by WNT ligands and

the DC, HUWE1 enhances WNT target gene expression through a mechanism distinct from the

control of CTNNB1 stability (Fig 2). Furthermore, regulation of WNT/CTNNB1 signaling by

HUWE1 is mediated by a subset of DC components, including APC, AXIN1 and GSK3A or

GSK3B, but excluding CSNK1A1 and AXIN2 (Figs 1, 3 and 4). HUWE1 promotes WNT

signaling by antagonizing the activity of this DC (Fig 1). The ubiquitin ligase activity of

HUWE1 is required to enhance WNT signaling (Fig 5), suggesting that a substrate of HUWE1

mediates its function.

One of the mechanisms whereby HUWE1 enhances WNT/CTNNB1 signaling is by

antagonizing phosphorylation of the CTNNB1 phosphodegron by the DC complex, thereby

increasing CTNNB1 abundance, but surprisingly this happens in the absence of CSNK1A1.

These results demonstrate that in HAP1 cells, CSNK1A1 is not absolutely required for

GSK3A/GSK3B-dependent phosphorylation of residues S33, S37 and T41 in the CTNNB1

phosphodegron, either because GSK3A/GSK3B can phosphorylate these residues without the

priming phosphorylation of S45 by CSNK1A1, or because other kinases phosphorylate S45 in

the absence of CSNK1A1. While priming of S45 by CSNK1A1 is generally considered a

requirement for phosphorylation of S33, S37 and T41 by GSK3A/GSK3B [6, 7], some reports

suggest it is not [30, 31].

We also show there is another mechanism whereby HUWE1 enhances WNT/CTNNB1

signaling that is independent of CTNNB1 stability. HUWE1 could potentially regulate CTNNB1

subcellular localization or its interactions with the TCF/LEF transcription complex, or it could

regulate other downstream steps in the pathway. Elucidating this second mechanism and whether

it is also mediated by a subset of DC complex components, like HUWE1-dependent regulation

of CTNNB1 abundance, will be crucial to understand the full scope of how HUWE1 regulates

WNT signaling.

Intriguingly, only a subset of DC components, including APC, AXIN1 and GSK3A or

GSK3B, but not CSNK1A1 or AXIN2, mediate the function of HUWE1 in WNT signaling (Figs

1, 3 and 4). We were surprised to find that AXIN1 was required to mediate the effects of

HUWE1 but AXIN2 was not. In WT HAP1-7TGP cells, AXIN1 and AXIN2 are redundant in

their capacity to suppress WNT signaling: eliminating either AXIN1 or AXIN2 has no effect on

WNT reporter activity, whereas eliminating both results in constitutive pathway activation [15].

Yet, in CSNK1A1^{KO} cells, only AXIN1 loss eliminated the reduction in WNT target gene

expression caused by HUWE1 KD (Fig 4 and Table 1). These results suggest that AXIN1 and

AXIN2 are not redundant in their capacity to mediate the effects of HUWE1, at least in the

absence of CSNK1A1. This finding is unexpected given that mouse AXIN1 and AXIN2 proteins

have been reported to be functionally equivalent *in vivo* [32], and will require further

investigation.

The ubiquitin ligase activity of HUWE1 is required to promote WNT signaling in

CSNK1A1^{KO} cells (Fig 5). What are the relevant ubiquitylated HUWE1 substrates, and how do

they regulate WNT signaling? Does HUWE1-dependent ubiquitylation target putative substrates

for proteasomal degradation or does it regulate their activity? Since a subset of DC components

mediates the effects of HUWE1 on WNT signaling, is the abundance or activity of a DC

component regulated by HUWE1-dependent ubiquitylation or are there other ubiquitylated

substrates that indirectly impinge on DC abundance or activity? Identification of the relevant

HUWE1 substrates should help answer these questions.

Previous reports have implicated HUWE1 as a *negative* regulator of WNT signaling [33-36].

This is the opposite of what we find in WT HAP1-7TGP, CSNK1A1^{KO} and CTNNB1^{ST-A} cells,

in which HUWE1 is a *positive* regulator of the pathway: eliminating HUWE1 or its catalytic

activity in these cells substantially reduces WNT/CTNNB1 signaling (Figs 1-5). HUWE1 has

been reported to polyubiquitylate DVL and prevent DVL multimerization [33], which is required

to form a functional signalosome and transduce WNT signals [37, 38]. HUWE1 has also been

reported to ubiquitylate CTNNB1 and promote CTNNB1 degradation [34]. The latter mechanism

is in fact the opposite of what we find in CSNK1A1^{KO} cells, in which HUWE1 loss reduces

CTNNB1 abundance (Fig 1). Based on both reported mechanisms, HUWE1 loss would be

expected to promote rather than reduce WNT signaling. Therefore, we do not think that DVL or

CTNNB1 are the relevant ubiquitylated substrates that mediate the effects of HUWE1 on WNT

signaling in HAP1 cells. These disparate results could reflect differences in experimental

systems, since the previous reports primarily studied HUWE1 in *C. elegans* and HEK293T cells

[33, 34], while the experiments presented in the current study were conducted in HAP1 cells.

Identifying the substrate of HUWE1 that mediates its role as a positive regulator of

WNT/CTNNB1 signaling in HAP1 cells should help explain these differences.

We demonstrate that HUWE1 loss reduces WNT signaling in cells containing mutations in

some WNT pathway components but not in others (Figs 1-5). These results raise the possibility

of targeting the signaling mechanisms by which HUWE1 enhances WNT signaling selectively in

tumors harboring mutations in specific WNT pathway components. Eliminating or reducing the

activity of HUWE1 itself, which reduces WNT/CTNNB1 signaling in WT HAP1-7TGP,

CSNK1A1^{KO} and CTNNB1^{ST-A} cells, is unlikely to be a viable therapeutic strategy due to the

pleiotropic effects of HUWE1 on cell physiology, including tumor suppressor functions [39].

However, if the relevant ubiquitylated target of HUWE1 is identified, there may be other ways to

phenocopy the effects of HUWE1 loss on WNT signaling more specifically. Phenocopying the

effects of HUWE1 loss may not be effective in tumors driven by APC truncations, given that in

APC^{KO} cells HUWE1 loss does not reduce WNT signaling due to the role of APC itself in

mediating the effects of HUWE1. However, in tumors containing activating mutations in

CTNNB1 like those engineered into our CTNNB1^{ST-A} cell line, or mutations in the ZNRF3 or

RNF43 tumor suppressors, all of which result in hyperactive WNT signaling in the presence of a

functional DC, phenocopying the effects of HUWE1 loss may reduce WNT signaling enough to

provide a therapeutic benefit.

We recognize that all the experiments presented in the Results section of this manuscript

were conducted in HAP1 cells or derivatives thereof, which could raise concerns about the

generality and specificity of our conclusions. We also studied HUWE1 in other cell lines

commonly used in WNT signaling research, but our attempts to knock out HUWE1 yielded only

partial KOs. We targeted *HUWE1* by CRISPR/Cas9-mediated genome editing in HEK293T-7TG

and HEK293T-7TG CSNK1A1^{KO} cells (see Materials and methods). Out of 113 independent

clonal cell lines in which we identified mutations in all *HUWE1* alleles, at least one allele had

been repaired in frame to encode WT HUWE1 protein (S6A and S6B Figs). This is probably

because HUWE1 is a common essential gene as per DEPMAP classification

(https://depmap.org/portal/gene/HUWE1?tab=overview), so complete loss of HUWE1 may be

lethal in HEK293T cells. However, we have previously shown that microinjection of *HUWE1*

mRNA into *Xenopus* embryos results in body axis duplication [15], consistent with a more

general role of HUWE1 as a positive regulator of WNT signaling beyond HAP1 cells.

Furthermore, we designed many of our experiments so as to minimize the possibility of non-

specific or pleiotropic effects. We knocked out HUWE1 in two independent CSNK1A1^{KO} cell

lines with comparable results (Fig 3). We used two different sgRNAs for CRISPR/Cas9-

mediated HUWE1 KO in multiple clonal cell lines (Figs 1-3), a different sgRNA for CRISPR

base editing of the HUWE1 catalytic residue in multiple CSNK1A1^{KO}; HUWE1^{C4341R} clonal cell

lines (Fig 5), and another two different sgRNAs for CRISPRi-mediated HUWE1 KD in

polyclonal cell populations (Fig 4). In all cases, we found reproducible reductions in

WNT/CTNNB1 signaling. We also saw consistent effects of HUWE1 loss in three different

genetic backgrounds: WT HAP1-7TGP, CSNK1A1^{KO} and CTNNB1^{ST-A} cells (Figs 1-5). We

measured the effects of HUWE1 loss on three or four endogenous WNT target genes and on an

7TGP, an established WNT transcriptional reporter, with comparable outcomes (Figs 1-5). The

effects of HUWE1 on WNT signaling could be reversed completely by a relatively short and

specific pharmacological treatment with the GSK3A/GSK3B inhibitor CHIR-99021 (Fig 1), and

by introducing mutations in some DC components but not others (Figs 3 and 4). Altogether,

these results make it very unlikely that the effects of HUWE1 loss are non-specific or due to

pleiotropic downregulation of unrelated cellular functions that affect WNT signaling.

Our study also highlights the remarkable potential of HAP1 haploid cells to dissect complex

genetic networks in a cell line of human origin [16] through a combination of genome-wide

forward genetic screens, loss-of-function and site-directed mutagenesis analyses, and genetic

interaction analyses. Despite great advances in CRISPR/Cas-based genome editing technologies

during the last decade [40], it remains challenging to knock out two or more alleles of multiple

genes and to introduce targeted homozygous point mutations at scale in diploid primary cells,

stem cells, and polyploid immortalized cell lines. We could readily do both in HAP1 cells

because they have a single allele of most genes. This enabled us to conduct loss-of-function

genetic analyses in multiple genetic backgrounds by comparing several HUWE1 KO and control

WT clonal cell lines to obtain highly quantitative phenotypic data that confirmed and extended

the results of our initial genetic screens (Fig 3). Using CRISPR/Cas9-mediated HDR, we

generated a CTNNB1 variant in which we mutated three key phosphorylation sites in the

phosphodegron at the single endogenous *CTNNB1* locus, and in a second round of

CRISPR/Cas9-mediated genome editing we generated multiple HUWE1 KO and WT cell lines

to demonstrate that regulation of WNT signaling by HUWE1 has a component that is

independent of CTNNB1 stability (Fig 2). Using CRISPR-mediated base editing, we generated

three clonal cell lines containing a point mutation in the catalytic residue of HUWE1 at the single

endogenous *HUWE1* locus, which enabled us to demonstrate that the ubiquitin ligase activity of

HUWE1 is required for its function in WNT signaling (Fig 5). Finally, we generated single,

double, and triple KO mutants for all components of the DC, alone and in certain combinations

(11 distinct mutant genetic backgrounds in total) (Table 1 and S4 Fig). Combined with a

CRISPRi strategy, this enabled us to carry out an extensive genetic interaction analysis and

demonstrate that positive regulation of WNT signaling by HUWE1 is mediated by a subset of

DC components (Fig 4 and Table 1). These kinds of genetic analyses would have been

practically impossible to conduct in any diploid or polyploid human cell line. We hope this study

will inspire other researchers to take advantage of haploid human cell lines, of which there are

now many available [41, 42], to unravel other signaling pathways or biological processes in

similar ways.

HUWE1 has emerged as an important ubiquitin ligase with many cellular functions [17-20].

Here we show another role for HUWE1 regulating WNT/CTNNB1 signaling. Regulation of

CTNNB1 abundance by the DC is the central step in WNT/CTNNB1 signaling. Our discovery

that HUWE1 enhances WNT signaling by antagonizing DC-dependent CTNNB1

phosphorylation, thereby increasing CTNNB1 abundance, demonstrates that this crucial step in

WNT/CTNNB1 signaling is subject to more nuanced regulation than previously thought. The

second mechanism by which HUWE1 enhances WNT signaling independently of CTNNB1

stability is an intriguing additional layer of regulation that remains to be elucidated. Both

mechanisms provide new insights into WNT signaling and ubiquitin biology, bridging two

research fields that already have many intimate connections.

Materials and methods:

The following Materials and methods relevant to this manuscript have been described previously

[15]: cell lines and growth conditions, preparation of WNT3A conditioned media and

construction of the HAP1-7TGP WNT reporter haploid cell line.

Tissue culture media

Complete growth medium (CGM) 1 contains Dulbecco's Modified Eagles Medium (DMEM)

with High Glucose, without L-Glutamine and Sodium Pyruvate (GE Healthcare Life Sciences

Cat. # SH30081.01); 1X GlutaMAX-I (Thermo Fisher Scientific Cat. # 35050079); 1X MEM

Non-Essential Amino Acids (Thermo Fisher Scientific Cat. # 11140050); 1 mM Sodium

Pyruvate (Thermo Fisher Scientific Cat. # 11360070); 40 Units/ml Penicillin, 40 mg/ml

Streptomycin (Thermo Fisher Scientific Cat. # 15140122); 10% Fetal Bovine Serum (FBS).

CGM 2 contains Iscove's Modified Dulbecco's Medium (IMDM) with L-glutamine, with

HEPES, without Alpha-Thioglycerol (GE Healthcare Life Sciences Cat. # SH30228.01); 1X

GlutaMAX-I; 40 Units/ml Penicillin, 40 mg/ml Streptomycin; 10% FBS.

Plasmids

pX330-U6-Chimeric_BB-CBh-hSpCas9 (pX330) (Addgene plasmid # 42230) was a gift from

Feng Zhang; pCMV_ABEmax_P2A_GFP (Addgene plasmid # 112101) was a gift from David

Liu; MLM3636 (Addgene plasmid # 43860) was a gift from Keith Joung; Lenti-(BB)-EF1a-

KRAB-dCas9-P2A-BlastR (Addgene plasmid # 118154) was a gift from Jorge Ferrer;

LentiCRISPRv2-mCherry (Addgene plasmid # 99154) was a gift from Agata Smogorzewska;

pMDLg/pRRE (Addgene plasmid # 12251), pRSV-Rev (Addgene plasmid # 12253) and

pMD2.G (Addgene plasmid # 12259) were a gift from Didier Trono; pCS2-YFP was a gift from

Henry Ho; pmCherry was a gift from Jan Carette; pX458-mCherry was generated as described

previously [43].

The following plasmids were purchased: pLenti6.2/V5-DEST (Thermo Fisher Scientific Cat.

V36820); pENTR2B (Thermo Fisher Scientific Cat. # A10463); MGC Human AXIN1

Sequence-verified cDNA (Clone ID 5809104) (Horizon Cat. # MHS6278-202833071).

To generate pCMV_ABEmax_P2A_mCherry, mCherry was amplified by PCR from plasmid

pmCherry using primers pCMV_ABEmax_P2A_mCherry_Fw (5'-GAA GCA GGC TGG AGA

CGT GGA GGA GAA CCC TGG ACC TAT GGT GAG CAA GGG CGA GGA-3') and

pCMV_ABEmax_P2A_mCherry_Rv (5'-CAG ACT TGT ACA GCT CGT CCA TGC CG-3'),

designed to include BsmBI and BsrGI restriction sites, respectively. The PCR product was

digested with BsmBI and BsrGI and ligated into pCMV_ABEmax_P2A_GFP digested with the

same enzymes to replace GFP with mCherry.

To generate pLenti6.2-V5-EXP-N-TERM-S-FLAG-N-hAXIN1, human AXIN1 was

amplified by PCR from MGC Human AXIN1 Sequence-verified cDNA (Clone ID 5809104)

using primers pENTR2B_SalI_S-FLAG-N_hAXIN1_pcr_fw (5'-GCG CCG GAA CCA ATT

CAG TCG ACC CTG CAG GAT GGA TTA CAA GGA CGA CGA TGA CAA GGG CGG

CCG CAT GAA TAT CCA AGA GCA GGG TTT CCC CTT GGA CC-3'), containing an N-

terminal SalI restriction site followed by a FLAG tag sequence flanked by SbfI and NotI

restriction sites, and pENTR2B_XhoI_hAXIN1_pcr_rv (5'-AAA GCT GGG TCT AGA TAT

CTC GAG TCA GTC CAC CTT CTC CAC TTT GCC GAT GA-3'), containing a C-terminal

XhoI restriction site. The product was digested with SalI and XhoI, and subcloned into

pENTR2B digested with the same enzymes. One clone was sequenced completely and subcloned

into pLenti6.2/V5-DEST using the Gateway LR Clonase II Enzyme mix.

All constructs were confirmed by sequencing.

Antibodies

Primary antibodies: purified mouse anti-β-catenin (Clone 14/Beta-Catenin) (1:1000, BD

Biosciences Cat. # 610154), rabbit mAb anti-non-phospho (active) β-catenin (Ser33-37-Thr41)

(D13A1) (1:1000, Cell Signaling Technology Cat. # 8814), mouse anti-GAPDH (1:4000, Santa

Cruz Biotechnology, Cat. # sc-47724), recombinant rabbit anti-Sodium Potassium (Na^+/K^+)

ATPase [EP1845Y] (1:4000, Abcam Cat. # ab76020), rabbit anti-Lasu1/Ureb1 (HUWE1)

(1:1000, Bethyl Laboratories Cat. # A300-486A), rabbit mAb anti-AXIN1 (C76H11) (1:1000,

Cell Signaling Technology Cat. # 2087), rabbit mAb anti-AXIN2 (76G6) (1:500, Cell Signaling

Technology Cat. # 2151), rabbit mAb anti-GSK-3α/β (D75D3) (1:2000, Cell Signaling

Technology Cat. # 5676), mouse anti-APC (NT, clone Ali 12.28) (1:1000, Millipore Sigma, Cat.

MAB3785), rabbit anti-APC (1:1000, Biorbyt Cat. # orb213564), mouse anti-CSNK1A1

(1:250, Santa Cruz Biotechnology, Cat. # sc-74582).

Secondary antibodies: IRDye 800CW donkey anti-mouse IgG (H+L) (1:10000, Li-Cor Cat. #

926-32212), IRDye 680RD donkey anti-rabbit IgG (H+L) (1:10000, Li-Cor Cat. # 925-68073),

peroxidase AffiniPure donkey anti-goat IgG (H+L) (1:5000, Jackson ImmunoResearch

Laboratories Cat. # 705-035-003), peroxidase AffiniPure goat anti-rabbit IgG (H+L) (1:10000,

Jackson ImmunoResearch Laboratories Cat. # 111-035-003), peroxidase AffiniPure donkey anti-

mouse IgG (H+L) (1:5000, Jackson ImmunoResearch Laboratories Cat. # 715-035-150), goat

anti-mouse IgG (H+L) HRP conjugate (1:10000, Bio-Rad Cat. # 1706516).

Primary and secondary antibodies used for detection with the Li-Cor Odyssey imaging

system were diluted in a 1 to 1 mixture of Odyssey Intercept Blocking Buffer (Li-Cor Cat. #

927–40000) and TBST (Tris buffered saline (TBS) + 0.1% Tween-20), and those used for

detection by chemiluminescence were diluted in TBST + 5% skim milk. All primary antibody

incubations were done overnight at 4°C, and secondary antibody incubations were done for 1 hr

at room temperature (RT).

Construction of mutant HAP1-7TGP cell lines by CRISPR/Cas9-mediated genome editing

Oligonucleotides encoding single guide RNAs (sgRNAs) (S2 File) were selected from a

published library [44], or designed using either of two online CRISPR design tools [45, 46] and

cloned into either pX330 or pX458-mCherry according to a published protocol [47].

Clonal HAP1-7TGP cell lines were established by transient transfection with either pX330 or

pX458-mCherry containing the sgRNA followed by single cell sorting as follows. A transfection

mix was prepared by diluting 450 ng of pX330 and 50 ng of pmCherry (used as a cotransfection

marker for FACS sorting) or 500 ng of pX458-mCherry in 48 µl Opti-MEM I, adding 2 µl of X-

tremeGENE HP and incubating for 20 min at RT. HAP1-7TGP cells or derivatives thereof were

reverse-transfected in a well of a 24-well plate by overlaying 0.5 ml of CGM 2 (without

antibiotics) containing 6×10^5 cells over the 50 µl of transfection mix. Cells were passaged to a

10 cm dish ~24 hr post-transfection, using 150 µl of Trypsin-EDTA (0.25%) (Thermo Fisher

Scientific Cat. # 25200056) to detach them (reverse-transfection of HAP1 cells caused unusually

high adherence, hence the higher trypsin concentration). Three to four days post-transfection,

single transfected (mCherry[+]) cells were sorted into 96-well plates containing 200 µl of CGM 2

per well and grown undisturbed for 16 to 18 days. Single colonies were passaged to 24-well

plates, and a small number of cells was reserved for genotyping.

For genotyping, genomic DNA was extracted by adding 4 volumes of QuickExtract DNA

Extraction Solution (Epicentre Cat. # QE09050) to the cells. Extracts were incubated 10 min at

65°C, 3 min at 98°C, and 5 µl were used as input for PCR amplification of the genomic locus

containing the sgRNA target site in 15 µl reactions containing 1X LongAmp Taq reaction buffer,

300 mM of each dNTP, 400 nM of each of the flanking primers indicated in S2 File (most of

them designed using the Primer-BLAST online tool from the NCBI) and 0.1 units/µl of

LongAmp Taq DNA polymerase (NEB Cat. # M0323L). Amplification of the genomic locus

containing the sgRNA target site was confirmed by analysis of the PCR products on a 1%

agarose gel and the presence of desired mutations was confirmed by sequencing the amplicons

using the primers indicated in S2 File. Given that most engineered cell lines remained haploid,

sequencing results were usually unequivocal. Sequencing results for all the clonal cell lines used

in the study is presented in S1 File, and for selected clonal cell lines, immunoblot analysis

confirmed the absence of the protein products.

Whenever possible, multiple independent mutant cells lines, often generated using two

different sgRNAs (see S1 File), were expanded and used for further characterization. For some

of the comparisons between WT and mutant cells, multiple individual cell lines confirmed by

sequencing to be WT at the sgRNA target site were also expanded and used as controls. To

generate double and triple mutant cell lines, a single clonal cell line with the first desired

mutation was used in a subsequent round of transfection with either pX330 or pX458-mCherry

containing the second and, if applicable, third sgRNAs. Alternatively, WT HAP1-7TGP cells

were directly transfected with a combination of pX330 or pX458-mCherry constructs targeting

two genes simultaneously.

Construction of CTNNB1ST-A cell line by CRISPR/Cas9-mediated HDR

Oligonucleotides encoding sgRNAs complementary to exon 3 of *CTNNB1* (S2 File) were

designed using either of two online CRISPR design tools [45, 46] and cloned into pX458-

mCherry using a published protocol [47].

Clonal CTNNB1ST-A cell lines were established by transient transfection of HAP1-7TGP

cells with pX458-mCherry containing the sgRNA, and a single stranded oligonucleotide

(ssODN) donor template encoding the desired mutations, called CTNNB1 (ST-A mutant) donor

(5'-ATT TGA TGG AGT TGG ACA TGG CCA TGG AAC CAG ACA GAA AAG CGG CTG

TTA GTC ACT GGC AGC AAC AGT CTT ACC TGG ACG CTG GAA TCC ATG CTG GTG

CCA CTG CCA CAG CTC CTG CTC TGA GTG GTA AAG GCA ATC CTG AGG AAG

AGG ATG TGG ATA CCT CCC AAG TCC TGT ATG AGT GGG AAC AGG GAT TTT CTC

AG-3'). A transfection mix was prepared by diluting 500 ng pX458-mCherry-CTNNB1-Ex3-

sgRNA and 500 ng (8 pmol) ssODN in 48 µl Opti-MEM I. 2 µl of X-tremeGENE HP were

added, and the mix was vortexed and incubated for 20 min at RT. The 50 µl mix was placed in

an empty well of a 24-well plate and 0.5 ml of CGM 2 containing 6×10^5 cells was seeded onto

the mix. The cells were passaged the following day to a 10 cm dish and grown for 3 additional

days. Single cells exhibiting high EGFP fluorescence from the 7TGP WNT reporter, presumably

due to successful mutagenesis of the CTNNB1 phosphodegron, were sorted, expanded, and

genotyped as described above. A single clonal cell line containing point mutations in three of the

four targeted sites in the phosphodegron (S2A Fig and S1 File) was used for all subsequent

experiments.

Construction of HUWE1 catalytic mutant CSNK1A1^{KO}; HUWE1^{C4341R} cell lines by base

editing

An oligonucleotide encoding an sgRNA complementary to exon 83 of *HUWE1* (S2 File) was

designed to include the targeted nucleotide within the editing window of the base editor

ABEmax (positions 4-8 in the protospacer) using BE-Hive (https://www.crisprbehive.design), an

online base editing sgRNA design tool [48], and cloned into MLM3636 according to a published

protocol (Joung Lab gRNA cloning protocol:

https://media.addgene.org/data/plasmids/43/43860/43860-attachment_T35tt6ebKxov.pdf). A

transfection mix was prepared by diluting 750 ng pCMV-ABEmax-P2A-mCherry and 250 ng

MLM3636-HUWE1-C4341R-sgRNA1 in 50 µl Opti-MEM I, adding 2 µl of X-tremeGENE HP

and incubating for 20 min at RT. CSNK1A1^{KO} cells were reverse-transfected in a well of a 24-

well plate by overlaying 0.5 ml of CGM 2 (without antibiotics) containing 6 x 10^5 cells over the

50 µl of transfection mix. Cells were passaged to a 6 cm dish ~24 hr post-transfection, using 150

µl of Trypsin-EDTA (0.25%) (Thermo Fisher Scientific Cat. # 25200056) to detach them. Three

days post-transfection, single transfected (mCherry$^+$) cells were sorted into 96-well plates

containing 200 µl of CGM 2 per well and grown undisturbed for 16 to 18 days. Cells were

expanded and genotyped as described above.

Targeting *HUWE1* by CRISPR/Cas9 in HEK293T-7TG and HEK293T-7TG CSNK1A1^{KO}

cells

Oligonucleotides HUWE1-IVT-2503-F and HUWE1-IVT-2503-R encoding sgRNAs

complementary to exon 6 of *HUWE1* (S2 File) were designed using sgRNA Scorer 2.0 [49] and

cloned into LentiCRISPRv2-mCherry previously digested with BsmBI. HEK293T-7TG is a

clonal cell line derived from HEK293T cells that contains a fluorescent WNT reporter.

HEK293T-7TG CSNK1A1^{KO} is a clonal cell line derived from HEK293T-7TG cells in which

CSNK1A1 has been knocked out. Construction of both cell lines will be described elsewhere.

Clonal HEK293T-7TG and HEK293T-7TG CSNK1A1^{KO} cell lines in which HUWE1 was

targeted by CRISPR/Cas9 were established by transient transfection with LentiCRISPRv2-

mCherry containing the sgRNAs followed by single cell sorting. ~24 hr before transfection, 8 x

10^4 HEK293T-7TG or HEK293T-7TG CSNK1A1^{KO} cells per well were seeded in 24-well plates

and grown in CGM 1. On the day of transfection, CGM 1 was replaced with 450 µl of antibiotic-

free CGM 1. 50 µl of a transfection mixture containing 500 ng LentiCRISPRv2-mCherry and 1

µl of X-tremeGENE™ HP DNA Transfection Reagent (Millipore Sigma, Cat # 06366236001)

prepared in OptiMEM were added dropwise. ~24 hr post-transfection, cells were transferred to a

6 cm dish, and ~72 hr post-transfection, single transfected (mCherry$^+$) cells were sorted into 96-

well plates containing 200 µl of CGM 1 media per well and grown undisturbed for 16 days.

Single colonies were expanded by passaging to 24-well plates, and 10 µl of cell suspension were

reserved for genotyping.

For genotyping, genomic DNA was extracted by adding 4 volumes of QuickExtract DNA

Extraction Solution (Epicentre, Cat # QE09050) to the cells. Extracts were incubated for 10 min

at 65°C, 3 min at 98°C, and 5 µl were used as input for PCR amplification of the *HUWE1* target

site in 15 µl reactions containing 1X LongAmp Taq reaction buffer, 300 mM of each dNTP, 400

nM of each of the flanking primers PS1057-NGS-F and PS1057-NGS-R (S2 File) and 0.1

units/µl of LongAmp Taq DNA polymerase (NEB Cat. # M0323L). In a second amplification

step, complete Illumina adapter sequences (F: 5'-AAT GAT ACG GCG ACC ACC GAG ATC

TAC AC <8 bp barcode> AC ACT CTT TCC CTA CAC GAC GCT CTT CCG ATC* T-3' ; R:

5'-CAA GCA GAA GAC GGC ATA CGA GAT <8 bp barcode> G TGA CTG GAG TTC AGA

CGT GTG CTC TTC CGA TC*T-3'; * indicates a phosphorothioate (PTO) linked base) were

added and the amplicons were sequenced on the MiSeq system (Illumina). FASTQ sequencing

files were analyzed using the branch 1.1 version [50] of a previously described analysis pipeline

(https://github.com/rajchari2/ngs_amplicon_analysis). Total (dark blue) and out-of-frame (light

blue) mutation rates were calculated and plotted (S6 Fig).

Preparation of lentivirus, lentiviral transduction, and selection of HUWE1 KD and AXIN1-

overexpressing polyclonal cell populations

The transfer plasmid used to generate HUWE1 KD cell lines by CRISPRi was Lenti-(BB)-EF1a-

KRAB-dCas9-P2A-BlastR. The transfer plasmid used to generate cell lines overexpressing

AXIN1 was pLenti6.2-V5-EXP-N-TERM-S-FLAG-N-hAXIN1. ~24 hr before transfection, 21 x

10^6 HEK293T cells were plated in 20 ml of CGM 1 without antibiotics in a T-175 flask. A

transfection mixture was prepared by diluting 9.3 µg of transfer plasmid, 7 µg of pMDLg/pRRE,

7 µg of pRSV-Rev, 4.66 µg of pMD2.G, 1.05 µg pCS2-YFP (as a cotransfection marker), and

87.15 µl of 1 mg/ml polyethylenimine (PEI) in a final volume of 1 ml serum-free DMEM. The

mixture was incubated for 20 min at RT and added to the culture media in the flasks. The day

after transfection, the media was replaced with 18 ml of CGM1 containing a total of 20% FBS

without antibiotics. ~48 hr after transfection, the media was collected (first viral harvest),

centrifuged at 1000 x g for 5 min to remove cell debris, and the supernatant was reserved at 4°C.

18 ml of fresh media were added to the flask of cells. ~72 hr after transfection, the media was

collected (second viral harvest), centrifuged as before, and the supernatant was pooled with the

first viral harvest. The pooled supernatant was filtered through 0.45 µm filters (Acrodisc syringe

filters with 0.45 µm Supor membrane, Pall Corporation Cat. # 4654). The filtered media

containing lentiviral particles was aliquoted, snap frozen in liquid nitrogen, and stored at -80°C.

For smaller scale preparations of the lentivirus used for AXIN1 overexpression, the above

protocol was followed but the lentivirus was prepared using 293FT cells in in T-25 flasks and all

quantities and volumes were scaled down by ~1/7. 3 x 10^6 293FT cells were plated in 5 ml of

CGM 1 without antibiotics in a T-25 flask. A transfection mixture was prepared by diluting 1.33

µg of transfer plasmid, 1 µg of pMDLg/pRRE, 1 µg of pRSV-Rev, 0.66 µg of pMD2.G, 0.15 µg

pCS2-YFP (as a cotransfection marker), and 69.4 µg/ml PEI in a final volume of 180 µL serum-

free DMEM. The mixture was incubated for 20 min at RT and added to the culture media in the

flasks. The day after transfection, the media was replaced with 2.5 ml of CGM 1 containing 20%

FBS without antibiotics and the viral supernatants were collected and processed as described

above.

Approximately 24 hr before transduction, 2.5 x 10^5 HAP1-7TGP cells or derivatives thereof

were seeded in a 6-well plate. Cells were transduced by adding 1 ml of lentivirus-containing

supernatant mixed with 1 ml of CGM 2 and 4.4 µg/ml polybrene. ~24 hr post-transduction, cells

were passaged to 10 cm dishes and selected with 8 µg/ml blasticidin in CGM 2 for ~96 hr.

Untransduced cells from each genetic background were treated in parallel with 8 µg/mL

blasticidin to ensure that all cells were killed by the time selection of transduced cells was

complete.

Analysis of WNT reporter fluorescence

To measure WNT reporter activity in HAP1-7TGP cells or derivatives thereof, ~24 hr before

treatment cells were seeded in 24-well plates at a density of 8×10^4 per well and grown in 0.5 ml

of CGM 2. Cells were treated for 24 hr with the indicated concentrations of WNT3A CM diluted

in CGM 2. Cells were washed with 0.5 ml PBS, harvested in 150 µl of Trypsin-EDTA (0.05%)

(Thermo Fisher Scientific Cat. # 25300054), resuspended in 450 µl of CGM 2, and EGFP

fluorescence was measured by FACS on either a SA3800 Spectral Cell Analyzer (Sony

Biotechnology) or a CytoFLEX S Flow Cytometer (Beckman Coulter). Typically, fluorescence

data for 5,000–50,000 singlet-gated cells was collected and, unless indicated otherwise, the

median EGFP fluorescence ± standard error of the median (SEM = 1.253 s/n, where s = standard

deviation and n = sample size) was used to represent the data.

To measure WNT reporter activity in cells treated with the GSK3A/B inhibitor CHIR-99021,

~24 hr before treatment cells were seeded in 6-well plates at a density of 0.5×10^6 per dish. Cells

were treated the following day with 10 µM CHIR-99021 (CT99021) (Selleckchem Cat. # S2924)

or an equivalent volume of DMSO vehicle diluted in CGM 2 for 48 hr, replacing the media with

fresh CHIR-99021 or DMSO in CGM 2 after 24 hr of treatment. Cells were washed with 2 ml

PBS, harvested in 0.5 ml of 0.05% Trypsin-EDTA, resuspended in 1.5 ml of CGM 2, and EGFP

fluorescence was measured as described above.

Quantitative (q)RT-PCR analysis

Approximately 24 hr before treatment, cells were seeded in 24-well plates at a density of 3×10^5

per well and grown in 0.5 ml of CGM 2. Cells were treated for 24 hr with 50% WNT3A CM in

CGM 2 where indicated. The medium was removed, cells were washed once with PBS and

harvested in 400 µl of TRIzol Reagent (Thermo Fisher Scientific Cat. # 15596018). Extracts

were processed according to the manufacturer's protocol, taking the appropriate precautions to

avoid contamination with nucleases, and total RNA was resuspended in 20 µl of DEPC-treated

water (Thermo Fisher Scientific Cat. # AM9920). To synthesize cDNA, 125 ng of RNA were

diluted in 2 µl DEPC-treated water and incubated with 0.25 µl 10X ezDNase buffer and 0.25 µl

ezDNase enzyme for 5 min at 37°C to digest DNA contaminants. After DNase treatment, 1 µl of

DEPC-treated water and 1 µl of SuperScript^{TM} IV VILO^{TM} MM (Invitrogen Cat. # 11766500)

were added and the reaction was incubated for 10 min at 25°C, 10 min at 50°C, and 5 min at

85°C. For each primer pair, a cDNA dilution series from a representative sample was analyzed to

ensure that target amplification was linear across a sufficiently broad range of cDNA

concentrations. cDNA was diluted 1:100 in water, and 5 µl were mixed with 5 µl of Power

SYBR Green PCR Master Mix (Applied Biosystems Cat. # 4367659) containing 400 nM each of

forward and reverse primer (S2 File). Triplicate reactions for each cDNA and primer pair were

prepared in a MicroAmp Optical 384-well Reaction Plate (Thermo Fisher Scientific Cat. #

4309849), sealed with MicroAmp Optical Adhesive Film (Thermo Fisher Scientific Cat. #

4311971) and run using standard parameters in a QuantStudio 5 Real-Time PCR System

(Applied Biosystems). Thermo Fisher cloud design and analysis software (DA2) was used to

calculate the average relative abundance of *AXIN2*, *RNF43*, *TNFRSF19*, or *NKD1* mRNA

normalized to *HPRT1* mRNA, and fold-changes in mRNA abundance were calculated as the

quotient between the experimental and reference samples, with appropriate error propagation of

the respective standard deviations (SD).

1008

Immunoblot analysis and quantification of soluble CTNNB1 from membrane-free

supernatants (MFS)

Approximately 24 hr before treatment, cells were seeded in 6 cm dishes at a density of 2.5 x 10^6

per dish and grown in 5 ml of CGM 2. Cells were treated for 24 hr with 50% WNT3A CM in

CGM 2 where indicated. Cells were harvested, lysed by hypotonic shock, and extracts were

prepared as follows, with all handling done at 4°C. Cells were washed twice with ~5 ml cold

PBS and twice with ~5 ml cold 10 mM HEPES pH 7.4. Residual buffer was removed, and 100 µl

of ice-cold SEAT buffer (10 mM triethanolamine/acetic acid pH 7.6, 250 mM sucrose, 1X

SIGMAFAST Protease Inhibitor Cocktail Tablets EDTA-free (Sigma-Aldrich Cat. # S8830), 25

µM MG132 (Sigma-Aldrich Cat. # C2211), 1X PhosSTOP (Roche Cat. # 04906837001), 1 mM

NaF, 1 mM Na_3VO_4, 1 mM dithiothreitol (DTT), 62.5 U/ml Benzonase Nuclease (EMD

Millipore Cat. # 70664), 1 mM $MgCl_2$) were added to the cells. Cells were scraped using a cell

lifter (Corning Cat. # 3008), transferred to 2-ml centrifuge tubes and disrupted mechanically by

triturating 10 times. Crude extracts were centrifuged for 20 min at 20,000 x g to pellet

membranes and other insoluble cellular material, and the MFS was carefully removed, avoiding

contamination from the pellet. The MFS was flash-frozen in liquid nitrogen and stored at -80°C

until further processing.

Extracts were thawed quickly at RT and transferred to ice. The protein concentration in the

MFS was quantified with the Pierce BCA Protein Assay Kit (Thermo Fisher Scientific Cat. #

23225), using BSA as a standard, and samples were normalized by dilution with SEAT buffer.

The MFS was diluted with 4X LDS sample buffer (Thermo Fisher Scientific Cat. # NP0007)

supplemented with 50 mM tris(2-carboxyethyl)phosphine (TCEP), incubated for 45 min at RT or

heated at 95°C for 10 min, and 30 µg of total protein were electrophoresed alongside Precision

Plus Protein All Blue Prestained Protein Standards (Bio-Rad Cat. # 1610373) in 4-15% TGX

Stain-Free protein gels (BioRad, various Cat. numbers) at 75 V for 15 min and 100 V for 1 hr 15

min using 1X Tris/Glycine/SDS running buffer (BioRad Cat. # 1610772). Following

electrophoresis, the gel was briefly activated with UV light using a Chemidoc imager (BioRad)

to covalently label proteins in the gel with Stain-Free fluorochromes.

Proteins were transferred to PVDF membranes in a Criterion Blotter apparatus (Bio-Rad Cat.

1704071) at 60 V for 2 hr using 1X Tris/Glycine transfer buffer (BioRad Cat. # 1610771)

containing 20% methanol. Following transfer, the membrane was imaged using the ChemiDoc

imager, and the total protein in each lane was quantified. Membranes were cut, blocked with

Odyssey Blocking Buffer (Li-Cor Cat. # 927–40000), incubated with mouse anti-β-catenin or

mouse anti-GAPDH primary antibodies, washed with TBST, incubated with IRDye 800CW

donkey anti-mouse IgG secondary antibody, washed with TBST followed by TBS, and imaged

using a Li-Cor Odyssey imaging system. Acquisition parameters in the manufacturer's Li-Cor

Odyssey Image Studio Lite software were set so as to avoid saturated pixels in the bands of

interest, and bands were quantified using background subtraction. The integrated intensity for

CTNNB1 was normalized to the total protein (or in some cases to the average of total protein and

the integrated intensity for GAPDH) in the corresponding lane. The average ± SD normalized

CTNNB1 intensity from duplicate blots was used to represent the data.

For CHIR-99021 or DMSO vehicle treated cells, ~24 hr before treatment cells were seeded

in 6 cm dishes at a density of 2×10^6 per dish and treated the following day with 10 µM CHIR-

99021 or DMSO for 48 hr. The media was replaced with fresh CHIR-99021 or DMSO in CGM 2

after 24 hr of treatment.

1054

Immunoblot analyses of soluble HUWE1, APC and CSNK1A1 from MFS

Some of the same membranes used to blot for soluble CTNNB1 were cut and used to blot for other proteins as indicated in the same figures. Blots were incubated with rabbit anti-Lasu1/Ureb1 (HUWE1), mouse anti-APC and mouse anti-CSNK1A1 primary antibodies. The following secondary antibodies were used: for HUWE1, IRDye 680RD donkey anti-rabbit IgG, for APC, IRDye 800CW donkey anti-mouse IgG (both were imaged using a Li-Cor Odyssey imaging system) and for CSNK1A1, peroxidase AffiniPure goat anti-mouse IgG (developed using SuperSignal West Femto Maximum Sensitivity Substrate (Thermo Fisher Scientific Cat. # 34095)).

Immunoblot analysis of total HUWE1, APC, CTNNB1, AXIN1, AXIN2, GSK3A/B and CSNK1A1 from whole cell extracts (WCE)

Approximately 72 hr before harvest, cells were seeded in 10 cm dishes at a density of 3×10^6 per dish and grown in 10 ml of CGM 2. Cells were harvested in 2 ml Trypsin-EDTA (0.05%) and resuspended in 6 ml CGM 2. 10×10^6 cells were centrifuged at 400 x g for 5 min, washed in 5 ml PBS, and centrifuged at 400 x g for 5 min. The supernatant was aspirated, and the cell pellets were flash-frozen in liquid nitrogen and stored at -80°C. Pellets were thawed quickly at RT and transferred to ice. All subsequent steps were done on ice. The cell pellets were resuspended in 150 μl of ice-cold RIPA lysis buffer (50 mM Tris-HCl pH 8.0, 150 mM NaCl, 2% NP-40, 0.25% deoxycholate, 0.1% SDS, 1X SIGMAFAST protease inhibitors, 1 mM $MgCl_2$, 62.5 U/ml Benzonase Nuclease, 1 mM DTT, 10% glycerol), sonicated in a Bioruptor Pico sonication device (Diagenode) 4 x 30 s in the ultra-high setting, centrifuged 10 min at 20,000 x g and the supernatant (WCE) was recovered.

The protein concentration in the WCE was quantified using the Pierce BCA Protein Assay

Kit. Samples were normalized by dilution with RIPA lysis buffer, further diluted with 4X LDS

sample buffer supplemented with 50 mM TCEP, incubated for 45 min at RT, and 30 µg of total

protein were electrophoresed alongside Precision Plus Protein All Blue Prestained Protein

Standards in 4-15% Criterion TGX Stain-Free protein gels at 75 V for 15min, and 100 V for 1 hr

and 15 min using 1X Tris/Glycine/SDS running buffer.

Proteins were transferred at 60 V for 2 hr to PVDF membranes using 1X Tris/Glycine

transfer buffer containing 20% methanol, and membranes were cut and blocked with either

Odyssey Intercept Blocking Buffer or TBST, 5% skim milk. Blots were incubated with rabbit

anti-Lasu1/Ureb1 (HUWE1), rabbit anti-APC, mouse anti-β-catenin, rabbit anti-AXIN1, rabbit

anti-AXIN2, rabbit anti-GSK3A/B, mouse anti-CSNK1A1 and mouse anti-GAPDH (as a loading

control) primary antibodies, washed with TBST, incubated with Peroxidase AffiniPure anti-

rabbit or anti-mouse secondary antibodies, washed with TBST followed by TBS, and developed

with SuperSignal West Femto.

1092

Immunoblot analysis and quantification of non-phospho-CTNNB1 (S33/S37/T41) and total

CTNNB1 from WCE

Approximately 24 hr before treatment, cells were seeded in 6 cm dishes at a density of 2 x

10^6 per dish and treated the following day with 10 µM CHIR-99021 or an equivalent volume of

DMSO vehicle for 48 hr. The media was replaced with fresh CHIR-99021 or DMSO in CGM 2

after 24 hr of treatment. Cells were harvested in 1 ml Trypsin-EDTA (0.05%) and resuspended in

3 ml CGM 2. Cells were centrifuged at 400 x g for 5 min, washed in 5 ml PBS, and centrifuged

at 400 x g for 5 min. The above protocol for immunoblot analysis of total proteins from WCE

was followed, except that protein samples were heated at 95°C for 10 min prior to

electrophoresis, and the total protein in each lane was quantified as follows and used for

normalization. Following electrophoresis, the gel was briefly activated with UV light using a

Chemidoc imager (BioRad) to covalently label proteins in the gel with Stain-Free fluorochromes.

Following transfer, the membrane was imaged using the ChemiDoc, and the total protein in each

lane was quantified. The blots were incubated with rabbit non-phospho (active) β-catenin (Ser33-

37-Thr41), mouse anti-β-catenin or mouse anti-GAPDH (as a loading control) primary

antibodies, IRDye 680RD donkey anti-rabbit IgG or IRDye 800CW donkey anti-mouse IgG

secondary antibodies, and imaged using a Li-Cor Odyssey imaging system.

1110

Quantitative dot blot of HUWE1 from WCE

3 µl WCE containing 8 µg protein were spotted onto nitrocellulose membrane for each sample in

triplicate. The membrane was allowed to dry for 15 min prior to staining with Revert 520 Total

Protein Stain (Li-Cor Cat. # 926-10010) according to the manufacturer's protocol

(https://www.licor.com/documents/1o8anlg26tnwqkj135ki6bo61fy4ztmi). The membrane was

imaged on the Li-Cor Odyssey M imaging system using the 520 nm channel to obtain a total

protein quantification for normalization. The membrane was then blocked with Odyssey

Blocking Buffer, incubated with rabbit anti-Lasu1/Ureb1 (HUWE1), washed with TBST,

incubated with IRDye 680RD donkey anti-rabbit IgG secondary antibody, washed with TBST

followed by TBS, and imaged using the Li-Cor Odyssey M imaging system. Acquisition

parameters in the manufacturer's Li-Cor Odyssey Image Studio Lite software were set so as to

avoid saturated pixels in the dots of interest, and dots were quantified using background

subtraction. The integrated intensity for HUWE1 was normalized to that for Revert 520 Total

Protein Stain in the same blot, and the average ± SD from triplicate dot blots was used to

represent the data. The specificity of the HUWE1 signal was confirmed by comparing dot blots

of WCE from CSNK1A1^{KO} and CSNK1A1^{KO}; HUWE1^{KO} cells. All normalized HUWE1

intensity values were within the linear range of a standard curve prepared from dot blots of a

serial dilution of WCE from CSNK1A1^{KO} cells.

1129

Preparation of figures and statistical analysis

Figures were prepared using PowerPoint (Microsoft). Table 1, S1 and S2 Files were prepared

using Excel (Microsoft). Graphs were prepared and statistical analysis performed using Prism 6

(GraphPad) or Excel. Details of the statistical tests used are given in the figure legends.

Significance is indicated as ****($p<0.0001$), *** ($p<0.001$), ** ($p<0.01$), * ($p<0.05$) or n.s. (not

significant). Pictures of immunoblots were adjusted for contrast and brightness only when

necessary for clarity using Image Studio Lite (LiCor), and were arranged in PowerPoint.

1137

Acknowledgments:

AML was supported by the Intramural Research Program of the National Institutes of Health, National Cancer Institute, Center for Cancer Research. We thank Caleb K. Sinclear for help with experiments, Lebensohn Lab members and members of the Laboratory of Cellular and Molecular Biology for input on the project, Rajat Rohatgi for feedback on the manuscript, and the Center for Cancer Research Genomics Core and the Laboratory of Genome Integrity Flow Cytometry Core for assistance and services. The authors declare no conflicting financial interests.

1147   **<u>References:</u>**

1148

1.   Nusse R, He X, Amerongen Rv. Wnt signaling : a subject collection from Cold Spring

Harbor Perspectives in Biology. Cold Spring Harbor, N.Y.: Cold Spring Harbor Laboratory

Press; 2013. vii, 454 p. p.

2.   Hoppler S, Moon RT. Wnt signaling in development and disease : molecular mechanisms

and biological functions. Hoboken, New Jersey: Wiley Blackwell; 2014. ix, 459 pages p.

3.   Nusse R, Clevers H. Wnt/beta-Catenin Signaling, Disease, and Emerging Therapeutic

Modalities. Cell. 2017;169(6):985-99.

4.   Anastas JN, Moon RT. WNT signalling pathways as therapeutic targets in cancer. Nat

Rev Cancer. 2013;13(1):11-26.

5.   Stamos JL, Weis WI. The beta-catenin destruction complex. Cold Spring Harb Perspect

Biol. 2013;5(1):a007898.

6.   Liu C, Li Y, Semenov M, Han C, Baeg GH, Tan Y, et al. Control of beta-catenin

phosphorylation/degradation by a dual-kinase mechanism. Cell. 2002;108(6):837-47.

7.   Amit S, Hatzubai A, Birman Y, Andersen JS, Ben-Shushan E, Mann M, et al. Axin-

mediated CKI phosphorylation of beta-catenin at Ser 45: a molecular switch for the Wnt

pathway. Genes Dev. 2002;16(9):1066-76.

8.   Yost C, Torres M, Miller JR, Huang E, Kimelman D, Moon RT. The axis-inducing

activity, stability, and subcellular distribution of beta-catenin is regulated in Xenopus embryos

by glycogen synthase kinase 3. Genes Dev. 1996;10(12):1443-54.

9.   Orford K, Crockett C, Jensen JP, Weissman AM, Byers SW. Serine phosphorylation-

regulated ubiquitination and degradation of beta-catenin. J Biol Chem. 1997;272(40):24735-8.

10.   Wu G, Xu G, Schulman BA, Jeffrey PD, Harper JW, Pavletich NP. Structure of a beta-

TrCP1-Skp1-beta-catenin complex: destruction motif binding and lysine specificity of the

SCF(beta-TrCP1) ubiquitin ligase. Mol Cell. 2003;11(6):1445-56.

11.   Gammons M, Bienz M. Multiprotein complexes governing Wnt signal transduction. Curr

Opin Cell Biol. 2018;51:42-9.

12.   Colozza G, Koo BK. Wnt/beta-catenin signaling: Structure, assembly and endocytosis of

the signalosome. Dev Growth Differ. 2021;63(3):199-218.

13.   Anthony CC, Robbins DJ, Ahmed Y, Lee E. Nuclear Regulation of Wnt/beta-Catenin

Signaling: It's a Complex Situation. Genes (Basel). 2020;11(8).

14.   Wiese KE, Nusse R, van Amerongen R. Wnt signalling: conquering complexity.

Development. 2018;145(12).

15.   Lebensohn AM, Dubey R, Neitzel LR, Tacchelly-Benites O, Yang E, Marceau CD, et al.

Comparative genetic screens in human cells reveal new regulatory mechanisms in WNT

signaling. Elife. 2016;5.

16.   Carette JE, Guimaraes CP, Wuethrich I, Blomen VA, Varadarajan M, Sun C, et al.

Global gene disruption in human cells to assign genes to phenotypes by deep sequencing. Nat

Biotechnol. 2011;29(6):542-6.

17.   Giles AC, Grill B. Roles of the HUWE1 ubiquitin ligase in nervous system development,

function and disease. Neural Dev. 2020;15(1):6.

18.   Gong X, Du D, Deng Y, Zhou Y, Sun L, Yuan S. The structure and regulation of the E3

ubiquitin ligase HUWE1 and its biological functions in cancer. Invest New Drugs.

2020;38(2):515-24.

19. Kao SH, Wu HT, Wu KJ. Ubiquitination by HUWE1 in tumorigenesis and beyond. J

Biomed Sci. 2018;25(1):67.

20. Qi L, Xu X, Qi X. The giant E3 ligase HUWE1 is linked to tumorigenesis,

spermatogenesis, intellectual disability, and inflammatory diseases. Front Cell Infect Microbiol.

2022;12:905906.

21. Staal FJ, van Noort M, Strous GJ, Clevers HC. Wnt signals are transmitted through N-

terminally dephosphorylated beta-catenin. EMBO Rep. 2002;3(1):63-8.

22. Aberle H, Bauer A, Stappert J, Kispert A, Kemler R. beta-catenin is a target for the

ubiquitin-proteasome pathway. EMBO J. 1997;16(13):3797-804.

23. Boonekamp KE, Heo I, Artegiani B, Asra P, van Son G, de Ligt J, Clevers H.

Identification of novel human Wnt target genes using adult endodermal tissue-derived organoids.

Dev Biol. 2021;474:37-47.

24. Yan D, Wallingford JB, Sun TQ, Nelson AM, Sakanaka C, Reinhard C, et al. Cell

autonomous regulation of multiple Dishevelled-dependent pathways by mammalian Nkd. Proc

Natl Acad Sci U S A. 2001;98(7):3802-7.

25. Schon S, Flierman I, Ofner A, Stahringer A, Holdt LM, Kolligs FT, Herbst A. beta-

catenin regulates NF-kappaB activity via TNFRSF19 in colorectal cancer cells. Int J Cancer.

2014;135(8):1800-11.

26. Qi LS, Larson MH, Gilbert LA, Doudna JA, Weissman JS, Arkin AP, Lim WA.

Repurposing CRISPR as an RNA-guided platform for sequence-specific control of gene

expression. Cell. 2013;152(5):1173-83.

27. Hunkeler M, Jin CY, Ma MW, Monda JK, Overwijn D, Bennett EJ, Fischer ES. Solenoid

architecture of HUWE1 contributes to ligase activity and substrate recognition. Mol Cell.

2021;81(17):3468-80 e7.

28. Bernassola F, Karin M, Ciechanover A, Melino G. The HECT family of E3 ubiquitin

ligases: multiple players in cancer development. Cancer Cell. 2008;14(1):10-21.

29. Koblan LW, Doman JL, Wilson C, Levy JM, Tay T, Newby GA, et al. Improving

cytidine and adenine base editors by expression optimization and ancestral reconstruction. Nat

Biotechnol. 2018;36(9):843-6.

30. Moshkovsky AR, Ma W, Kirschner MW. The self-regulated response of the Wnt

pathway to an oncogenic mutation in β-catenin. bioRxiv. 2021:2021.05.05.442673.

31. Wang Z, Vogelstein B, Kinzler KW. Phosphorylation of beta-catenin at S33, S37, or T41

can occur in the absence of phosphorylation at T45 in colon cancer cells. Cancer Res.

2003;63(17):5234-5.

32. Chia IV, Costantini F. Mouse axin and axin2/conductin proteins are functionally

equivalent in vivo. Mol Cell Biol. 2005;25(11):4371-6.

33. de Groot RE, Ganji RS, Bernatik O, Lloyd-Lewis B, Seipel K, Sedova K, et al. Huwe1-

mediated ubiquitylation of dishevelled defines a negative feedback loop in the Wnt signaling

pathway. Sci Signal. 2014;7(317):ra26.

34. Dominguez-Brauer C, Khatun R, Elia AJ, Thu KL, Ramachandran P, Baniasadi SP, et al.

E3 ubiquitin ligase Mule targets beta-catenin under conditions of hyperactive Wnt signaling.

Proc Natl Acad Sci U S A. 2017;114(7):E1148-E57.

35.   Vandewalle J, Langen M, Zschatzsch M, Nijhof B, Kramer JM, Brems H, et al. Ubiquitin

ligase HUWE1 regulates axon branching through the Wnt/beta-catenin pathway in a Drosophila

model for intellectual disability. PLoS One. 2013;8(11):e81791.

36.   Dominguez-Brauer C, Hao Z, Elia AJ, Fortin JM, Nechanitzky R, Brauer PM, et al. Mule

Regulates the Intestinal Stem Cell Niche via the Wnt Pathway and Targets EphB3 for

Proteasomal and Lysosomal Degradation. Cell Stem Cell. 2016;19(2):205-16.

37.   Fiedler M, Mendoza-Topaz C, Rutherford TJ, Mieszczanek J, Bienz M. Dishevelled

interacts with the DIX domain polymerization interface of Axin to interfere with its function in

down-regulating beta-catenin. Proc Natl Acad Sci U S A. 2011;108(5):1937-42.

38.   Schwarz-Romond T, Fiedler M, Shibata N, Butler PJ, Kikuchi A, Higuchi Y, Bienz M.

The DIX domain of Dishevelled confers Wnt signaling by dynamic polymerization. Nat Struct

Mol Biol. 2007;14(6):484-92.

39.   Su C, Wang T, Zhao J, Cheng J, Hou J. Meta-analysis of gene expression alterations and

clinical significance of the HECT domain-containing ubiquitin ligase HUWE1 in cancer. Oncol

Lett. 2019;18(3):2292-303.

40.   Wang JY, Doudna JA. CRISPR technology: A decade of genome editing is only the

beginning. Science. 2023;379(6629):eadd8643.

41.   Horii T, Hatada I. Genome Editing Using Mammalian Haploid Cells. Int J Mol Sci.

2015;16(10):23604-14.

42.   Li Y, Shuai L. A versatile genetic tool: haploid cells. Stem Cell Res Ther. 2017;8(1):197.

43.   Dubey R, van Kerkhof P, Jordens I, Malinauskas T, Pusapati GV, McKenna JK, et al. R-

spondins engage heparan sulfate proteoglycans to potentiate WNT signaling. Elife. 2020;9.

44.   Shalem O, Sanjana NE, Hartenian E, Shi X, Scott DA, Mikkelson T, et al. Genome-scale

CRISPR-Cas9 knockout screening in human cells. Science. 2014;343(6166):84-7.

45.   Doench JG, Hartenian E, Graham DB, Tothova Z, Hegde M, Smith I, et al. Rational

design of highly active sgRNAs for CRISPR-Cas9-mediated gene inactivation. Nat Biotechnol.

2014;32(12):1262-7.

46.   Hsu PD, Scott DA, Weinstein JA, Ran FA, Konermann S, Agarwala V, et al. DNA

targeting specificity of RNA-guided Cas9 nucleases. Nat Biotechnol. 2013;31(9):827-32.

47.   Cong L, Ran FA, Cox D, Lin S, Barretto R, Habib N, et al. Multiplex genome

engineering using CRISPR/Cas systems. Science. 2013;339(6121):819-23.

48.   Arbab M, Shen MW, Mok B, Wilson C, Matuszek Z, Cassa CA, Liu DR. Determinants

of Base Editing Outcomes from Target Library Analysis and Machine Learning. Cell.

2020;182(2):463-80 e30.

49.   Chari R, Yeo NC, Chavez A, Church GM. sgRNA Scorer 2.0: A Species-Independent

Model To Predict CRISPR/Cas9 Activity. ACS Synth Biol. 2017;6(5):902-4.

50.   Gooden AA, Evans CN, Sheets TP, Clapp ME, Chari R. dbGuide: a database of

functionally validated guide RNAs for genome editing in human and mouse cells. Nucleic Acids

Res. 2021;49(D1):D871-D6.

1273

 <u>Figure legends:</u>

Fig 1. HUWE1 and AXIN1 reciprocally regulate WNT signaling by modulating

GSK3A/GSK3B-dependent CTNNB1 phosphorylation and abundance.

(A-C). We note that the data for WT HAP-7TGP, CSNK1A1^{KO} and CSNK1A1^{KO}; HUWE1^{KO}

cells is discussed in the first section of the results, while the data for CSNK1A1^{KO}; AXIN1^{OE} and

CSNK1A1^{KO}; HUWE1^{KO}; AXIN1^{OE} cells is discussed in a later section of the results subtitled

"HUWE1 enhances WNT signaling by antagonizing the DC." Cells were treated with DMSO

vehicle or 10 µM of the GSK3A/GSK3B inhibitor CHIR-99021 for 48 hr as indicated. (A) WNT

reporter activity (median EGFP fluorescence from 10,000 singlets was measured for triplicate

wells and the average ± standard deviation (SD) of the three measurements is depicted), relative

to WT HAP1-7TGP cells treated with DMSO. Significance was determined by unpaired t-test

with Welch's correction. (B) Soluble CTNNB1 abundance (CTNNB1 intensity normalized to

total protein, average ± SD from duplicate immunoblots shown in S1A Fig) in membrane-free

supernatants (MFS) of the indicated cell lines, relative to WT HAP1-7TGP cells treated with

DMSO. (C) Non-phospho-CTNNB1 (S33/S37/T41) abundance (non-phospho-CTNNB1

intensity normalized to total protein, average ± SD from duplicate immunoblots shown in S1B

Fig) in whole cell extracts (WCE) of the indicated cell lines, relative to WT HAP1-7TGP cells

treated with DMSO.

1293

S1 Fig. HUWE1 and AXIN1 reciprocally regulate WNT signaling by modulating

GSK3A/GSK3B-dependent CTNNB1 phosphorylation and abundance.

(A-C) We note that the data for WT HAP-7TGP, CSNK1A1^{KO} and CSNK1A1^{KO}; HUWE1^{KO}

cells is discussed in the first section of the results, while the data for CSNK1A1^{KO}; AXIN1^{OE} and

CSNK1A1^{KO}; HUWE1^{KO}; AXIN1^{OE} cells is discussed in a later section of the results subtitled

"HUWE1 enhances WNT signaling by antagonizing the DC." Cells were treated with DMSO

vehicle or 10 µM of the GSK3A/GSK3B inhibitor CHIR-99021 for 48 hr as indicated. (A)

Immunoblots of soluble CTNNB1 from MFS, used for quantification in Fig 1B. (B)

Immunoblots of non-phospho-CTNNB1 (S33/S37/T41) and total CTNNB1 from WCE, used for

quantification in Fig 1C and S1C Fig, respectively. (C) Total CTNNB1 abundance (CTNNB1

intensity normalized to total protein, average ± SD from duplicate immunoblots shown in S1B)

in WCE of the indicated cell lines, relative to WT HAP1-7TGP cells treated with DMSO. (D)

Immunoblot analysis of total AXIN1 from WCE of the indicated cell lines used in A-C, and in

Fig 1. The polyclonal cell populations overexpressing AXIN1 were generated as described in

Materials and methods. AXIN1 abundance (AXIN1 intensity normalized to GAPDH intensity),

relative to CSNK1A1^{KO} cells, is indicated below the blots.

1310

Fig 2. HUWE1 enhances WNT signaling through a mechanism independent of CTNNB1

stability.

(A-F) Each circle represents a unique clonal cell line (determined by genotyping, S1 File). The

same cell lines were used in A-F. A single value for the parental WT HAP1-7TGP cell line, and

the average value from 3 independent clonal cell lines for each of the other genotypes, all

relative to the untreated WT HAP1-7TGP sample, are indicated by a horizontal line and

quantified above each group of circles. WT HAP1-7TGP cells were treated with 50% WNT3A

CM for 24 hr where indicated. Significance was determined by unpaired t-test with Welch's

correction. (A) Relative soluble CTNNB1 abundance (CTNNB1 intensity normalized to total

protein and GAPDH intensity, average from duplicate immunoblots shown in S2B Fig) in MFS

of the indicated cell lines. (B) Relative WNT reporter activity (median EGFP fluorescence from

100,000 singlets). (C-F) Relative WNT target gene expression (average quantification of *AXIN2*,

RNF43, *TNFRSF19*, or *NKD1* mRNA normalized to *HPRT1* mRNA, each measured in triplicate

reactions).

1325

S2 Fig. HUWE1 enhances WNT signaling through a mechanism independent of CTNNB1

stability.

(A) Genomic nucleotide and corresponding amino acid sequences comprising the CTNNB1

phosphodegron of WT HAP1-7TGP and CTNNB1^{ST-A} cells. The kinases that phosphorylate S or

T residues in the phosphodegron are indicated. Nucleotides and amino acids in red indicate

mutations. (B) Immunoblots of soluble HUWE1 and CTNNB1 from MFS of the indicated cell

lines. The CTNNB1 immunoblots were used for quantification in Fig 2A. (C-E) Treatment of

CTNNB1^{ST-A} cells with WNT3A does not promote further accumulation of soluble CTNNB1

and does not further increase WNT target gene expression. Cells were treated with 50% WNT3A

CM for 24 hr where indicated. (C) Immunoblots of total CTNNB1 from WCE used for

quantification in D. (D) Total CTNNB1 abundance (CTNNB1 intensity normalized to total

protein and GAPDH intensity, average ± SD from duplicate lanes of the immunoblots shown in

C) in WCE of CTNNB1^{ST-A} cells treated with WNT3A CM, relative to untreated CTNNB1^{ST-A}

cells. Significance was determined by unpaired t-test with Welch's correction. (E) mRNA

abundance (average ± SD *AXIN2*, *RNF43*, *TNFRSF19*, or *NKD1* mRNA normalized to *HPRT1*

mRNA, each measured in triplicate reactions) in CTNNB1^{ST-A} cells treated with WNT3A CM,

reported as percentage of the mRNA abundance in untreated CTNNB1^{ST-A} cells. (F) WNT

reporter activity (median EGFP fluorescence from 5,000 singlets) for the indicated cell lines,

relative to the average for CTNNB1^{ST-A} cells. Each circle represents a unique clonal cell line

(determined by genotyping, S1 File), and the average of 9-12 independent clones for each

genotype is indicated by a horizontal line and quantified above each group of circles.

Significance was determined by unpaired t-test with Welch's correction.

1348

Fig 3. HUWE1 enhances WNT signaling through mechanisms mediated by APC.

(A-F) The same cell lines were used in A-F. WT cells were treated with 50% WNT3A CM for

24 hr where indicated. (A) Immunoblot analysis of soluble proteins from MFS of the indicated

clonal cell lines. We note that CSNK1A1^{KO-2} cells contain a loss-of-function mutation resulting

in a 2-amino acid deletion (S1 File), and hence the protein product is still present. The "a" and

"b" superscripts next to the protein names indicate which of two membranes the corresponding

strips were cut from. * indicates a non-specific band observed with the mouse anti-APC

antibody. (B-F) Each circle represents a unique clonal cell line (determined by genotyping, S1

File). A single value for the parental WT HAP1-7TGP cell line, and the average value from 2-3

independent clonal cell lines for each of the other genotypes, all relative to the untreated WT

HAP1-7TGP sample, are indicated by a horizontal line and quantified above each group of

circles. Significance was determined by unpaired t-test with Welch's correction. (B) Relative

soluble CTNNB1 abundance (CTNNB1 intensity normalized to total protein, average from

duplicate immunoblots) in MFS of the indicated cell lines. (C-F) Relative WNT target gene

expression (average quantification of *AXIN2*, *RNF43*, *TNFRSF19*, or *NKD1* mRNA normalized

to *HPRT1* mRNA, each measured in triplicate reactions).

Fig 4. HUWE1 enhances WNT signaling through mechanisms mediated by a subset of DC components including APC, AXIN1 and GSK3A or GSK3B.

(A-E) The same cell lines are used in A-E. WT HAP1-7TGP cells were treated with 50% WNT3A CM for 24 hr where indicated. (A) Aggregate WNT target gene expression (average ± SD of all four target genes, calculated from the individual average quantifications of *AXIN2*, *RNF43*, *TNFRSF19* or *NKD1* mRNA normalized to *HPRT1* mRNA, each measured in triplicate reactions) in polyclonal cell populations targeted with HUWE1 sgRNAs, reported as percentage of aggregate WNT target gene expression in polyclonal cell populations targeted with SCR sgRNA control. Significance was determined by unpaired t-test with Welch's correction. (B-E) WNT target gene expression (average ± SD *AXIN2*, *RNF43*, *TNFRSF19*, or *NKD1* mRNA normalized to *HPRT1* mRNA, each measured in triplicate reactions) in polyclonal cell populations targeted with HUWE1 sgRNAs, reported as percentage of WNT target gene expression in polyclonal cell populations targeted with SCR sgRNA control.

S4 Fig. HUWE1 enhances WNT signaling through mechanisms mediated by a subset of DC components including APC, AXIN1 and GSK3A or GSK3B.

(A) Immunoblot analysis of total protein from WCE of the indicated clonal cell lines used for CRISPRi-mediated HUWE1 KD in Fig 4 and S5 Fig. The AXIN1 and AXIN2 immunoblots of CSNK1A1^{KO}; AXIN1^{KO} and CSNK1A1^{KO}; AXIN2^{KO} cells, respectively, exhibited bands of lower abundance and molecular weight than their respective counterparts in WT HAP1-7TGP cells. These bands may represent residual truncated protein products, but in both cases frameshift mutations in the single allele of the respective genes (S1 File) predicted the absence of full-

length, WT proteins. * indicates a non-specific band observed with the rabbit anti-APC antibody.

The "a" and "b" superscripts next to the protein names indicate which of two membranes the

corresponding strips were cut from. Dashed vertical lines indicate a rearrangement of samples

within the same blot. (B, C) GSK3A and GSK3B are functionally redundant in WNT signaling

in HAP1 cells. The same cell lines were used in B and C. (B) Immunoblot analysis of total

GSK3A and GSK3B from WCE of the indicated cell lines. (C) WNT reporter activity (median

EGFP fluorescence from 50,000 singlets was measured for biological duplicates of a single

clone, and the average ± SD of the two measurements was calculated) relative to untreated WT

HAP1-7TGP cells. (D) HUWE1 abundance, quantified by dot blots, in the clonal cell lines used

for CRISPRi-mediated HUWE1 KD in Fig 4 and S5 Fig. Total HUWE1 abundance (HUWE1

intensity normalized to total protein, average ± SD from triplicate dot blots) in WCE of the

indicated cell lines, relative to WT HAP1-7TGP cells. Significance was determined by unpaired

t-test with Welch's correction, and the difference in HUWE1 abundance between each mutant

cell line and WT HAP1-7TGP cells was not significant (not depicted).

1402

S5 Fig. Quantification of CRISPRi-mediated HUWE1 KD in various genetic backgrounds.

(A-B) Two polyclonal cell populations targeted with HUWE1 sgRNAs (1 and 2) and one

polyclonal cell population targeted with SCR sgRNA were derived for each genotype as

described in Materials and methods. (A) Immunoblots of total HUWE1 from WCE used for

quantification in B. The "a" and "b" superscripts next to the protein names indicate which of two

duplicate membranes the corresponding strips were cut from. Dashed vertical lines indicate a

rearrangement of samples within the same blot. (B) HUWE1 abundance (average HUWE1

intensity normalized to either Na$^+$/K$^+$ ATPase or GAPDH intensity from duplicate immunoblots

shown in A) in WCE of cell populations targeted with HUWE1 sgRNAs, reported as percentage

of HUWE1 abundance in WCE of cell populations targeted with SCR sgRNA control.

1413

Fig 5. Regulation of WNT signaling by HUWE1 requires its ubiquitin ligase activity.

(A) WNT reporter activity (median ± standard error of the median (SEM) EGFP fluorescence

from 37,000-50,000 cells) of one CSNK1A1^{KO}; HUWE1^{KO} clonal cell line and three catalytic

mutant CSNK1A1^{KO}; HUWE1^{C4341R} clonal cell lines, relative to CSNK1A1^{KO} cells. (B-D) WNT

target gene expression (average ± SD *AXIN2*, *RNF43* or *TNFSRF19* mRNA normalized to

HPRT1 mRNA, each measured in triplicate reactions) of the indicated clonal cell lines, relative

to CSNK1A1^{KO} cells. (E) Immunoblot analysis of total HUWE1 protein in WCE of the same cell

lines used in A-D.

1422

S6 Fig. Quantification of CRISPR/Cas9-mediated *HUWE1* mutations in HEK293T-7TG

and HEK293T-7TG CSNK1A1^{KO} cells.

(A, B) Sequencing reads of the *HUWE1* locus targeted by CRISPR/Cas9 in individual clonal cell

lines derived from HEK293T-7TG (A) or HEK293T-7TG CSNK1A1^{KO} (B) cells were quantified

for mutations. The X-axis shows individual clones, and the Y-axis indicates the percentage of

reads containing mutations. Bars in dark blue indicate the percentage of reads containing any

kind of mutation (total mutations) at the targeted locus in each clone, and bars in light blue

indicate the percentage of reads containing out-of-frame mutations at the same locus. In all 113

clones in which ~100% of the reads contained mutations (indicating all *HUWE1* alleles had been

successfully targeted), some of those mutations were always in frame, strongly suggesting that at

least one WT *HUWE1* allele is required for cell viability in HEK293T cells.

1434

 Supporting Information:

 S1 File. CRISPR/Cas9-engineered clonal cell lines used in this study.

 Single-mutant clones in which a single gene was targeted using CRISPR/Cas9 and double- or

 triple-mutant clones in which multiple genes were targeted using CRISPR/Cas9 are described in

 two separate spreadsheets labeled accordingly. When more than one clone was generated using

 the same CRISPR guide, the 'Clone Name' column indicates the generic name used throughout

 the manuscript to describe the genotype, and the 'Clone #' column identifies an individual clone.

 The 'HDR Donor' column indicates the name of the ssODN donor template used to generate

 some of the clonal cell lines (see Materials and Methods). The 'CRISPR guide' column indicates

 the name of the guide used, which is the same as that of the oligos encoding sgRNAs (see

 Materials and methods, and S2 File). The 'Genomic Sequence' column shows 80 bases of

 genomic sequence (5' relative to the gene is to the left) surrounding the target site. For each

 group of clones made using the same CRISPR guide (separated by gray spacers), the 'Genomic

 Sequence' column is headlined by the reference WT genomic sequence (obtained from RefSeq),

 with the guide sequence colored blue. The site of the double strand cut made by Cas9 is between

 the two underlined bases. Sequencing results for individual clones are indicated below the

 reference sequence. Some clones that remained WT at the targeted locus are indicated as such

 and were used as controls. For mutant clones, mutated bases are colored red (dashes represent

 deleted bases, three dots are used to indicate that a deletion continues beyond the 80 bases of

 sequence shown, and large insertions are indicated in brackets), and the nature of the mutation

 and the resulting genotype are described in the columns labeled accordingly. The figures in

 which each clone was used are also indicated. For double- and triple-mutant clones, the CRISPR

1458   guide used, the genomic sequence, the mutation and the genotype pertaining to each of the two

1459   or three targeted loci are designated '1', '2' and '3' in the column headings, and are shown under

1460   green, orange and purple spacers, respectively.

1461

S2 File. Oligonucleotides and primers used in this study.

Oligonucleotides and primers used for generation and characterization of clonal cell lines

engineered using CRISPR/Cas9 nuclease (CRISPRn), base editing, and CRISPRi, as well as

those used for qRT-PCR, are described in separate spreadsheets labeled accordingly. CRISPRn,

base editing and CRISPRi: the names and sequences of pairs of oligonucleotides encoding

sgRNAs, which were cloned into the respective vectors for each application as described in

Materials and methods, are shown in columns A and B, respectively. Additionally, for CRISPRn

and base editing the names and sequences of pairs of forward and reverse primers used to

amplify corresponding genomic regions flanking sgRNA target sites are shown in columns C and

D, respectively, and where applicable, the names and sequences of individual primers used to

sequence the amplified target sites are shown in columns E and F, respectively. qRT-PCR: the

names and sequences of pairs of forward and reverse primers used for qRT-PCR are shown in

columns A and B, respectively.

1475

Fig 1

A

B

C

S1 Fig

A

	WT	CSNK1A1^{KO}	CSNK1A1^{KO}; AXIN1^{OE}	CSNK1A1^{KO}; HUWE1^{KO}	CSNK1A1^{KO}; HUWE1^{KO}; AXIN1^{OE}	WT	CSNK1A1^{KO}	CSNK1A1^{KO}; AXIN1^{OE}	CSNK1A1^{KO}; HUWE1^{KO}	CSNK1A1^{KO}; HUWE1^{KO}; AXIN1^{OE}
DMSO	+	+	+	+	+	-	-	-	-	-
CHIR-99021	-	-	-	-	-	+	+	+	+	+

B

	WT	CSNK1A1^{KO}	CSNK1A1^{KO}; AXIN1^{OE}	CSNK1A1^{KO}; HUWE1^{KO}	CSNK1A1^{KO}; HUWE1^{KO}; AXIN1^{OE}	WT	CSNK1A1^{KO}	CSNK1A1^{KO}; AXIN1^{OE}	CSNK1A1^{KO}; HUWE1^{KO}	CSNK1A1^{KO}; HUWE1^{KO}; AXIN1^{OE}
DMSO	+	+	+	+	+	-	-	-	-	-
CHIR-99021	-	-	-	-	-	+	+	+	+	+

C

D

Relative AXIN1: 1.00 2.68 1.29 2.61

Fig 2

A

CTNNB1 phosphodegron

CTNNB1^{WT}:   CTG GAC TCT GGA ATC CAT TCT GGT GCC ACT ACC ACA GCT CCT TCT CTG
CTNNB1^{ST-A}: CTG GAC TCT GGA ATC CAT GCT GGT GCC ACT GCC ACA GCT CCT GCT CTG

CTNNB1^{WT} : L31,D32,S33,G34,I35,H36,S37,G38,A39,T40,T41,T42,A43,F44,S45,L36
CTNNB1^{ST-A} : L31,D32,S33,G34,I35,H36,A37,G38,A39,T40,A41,T42,A43,F44,A45,L36

Kinase: GSK3A/B   GSK3A/B   GSK3A/B   CSNK1A1

B

C

D

E

F

Fig 3

Fig 4

A

B

Fig 5

S6 Fig

A HEK293T-7TG

B HEK293T-7TG CSNK1A1^{KO}