PPARγ mediated enhanced lipid biogenesis fuels *Mycobacterium tuberculosis* growth in a drug-tolerant hepatocyte environment.

Binayak Sarkar[1], Jyotsna Singh[1], Mohit Yadav[1], Priya Sharma[2], Raman Deep Sharma[2], Shweta Singh[3], Aakash Chandramouli[4], Kritee Mehdiratta[5], Ashwani Kumar[3], Siddhesh S. Kamat[4], Devram S. Ghorpade[1], Debasisa Mohanty[1], Dhiraj Kumar[2], Rajesh S. Gokhale[1,4, #]

1 Immunometabolism laboratory, National Institute of Immunology, New Delhi, India, 2 Cellular Immunology Group, International Centre for Genetic Engineering and Biotechnology, New Delhi, India 3 CSIR-Institute of Microbial Technology, Chandigarh, India, 4 Indian Institute of Science Education and Research, Pune, Maharashtra, India.5 Council Of Scientific and Industrial Research–Institute of Genomics and Integrative Biology (CSIR-IGIB), New Delhi 110025, India, Academy of Scientific and Innovative Research (AcSIR), Ghaziabad 201002, India

Current address: Indian Institute of Science Education and Research, Pune, Maharashtra, India.

Keywords: Liver, Mtb, hepatocytes, PPARγ, anti-TB drugs, lipid droplets.

Corresponding author: Rajesh S. Gokhale (rsg@nii.ac.in)

Abstract:

Mycobacterium tuberculosis (Mtb) infection of the lungs, besides producing prolonged cough with mucus, also causes progressive fatigue and cachexia with debilitating loss of muscle mass. While anti-tuberculosis (TB) drug therapy is directed toward eliminating bacilli, the treatment regimen ignores the systemic pathogenic derailments that probably dictate TB-associated mortality and morbidity. Presently, it is not understood whether Mtb spreads to metabolic organs and brings about these impairments. Here we show that Mtb creates a replication-conducive milieu of lipid droplets in hepatocytes by upregulating transcription factor PPARγ and scavenging lipids from the host cells. In hepatocytes, Mtb shields itself against the common anti-TB drugs by inducing drug-metabolizing enzymes. Infection of the hepatocytes in the *in vivo* aerosol mice model can be consistently observed post-week 4 along with enhanced expression of PPARγ and drug-metabolizing enzymes. Moreover, histopathological analysis indeed shows the presence of Mtb in hepatocytes along with granuloma-like structures in human biopsied liver sections. Hepatotropism of Mtb during the chronic infectious cycle results in immuno-metabolic dysregulation that could magnify local and systemic pathogenicity, altering clinical presentations.

Keywords: *Mycobacterium tuberculosis*, PPARγ, anti-TB drugs, lipid droplets.

.

Introduction

Mycobacterium tuberculosis (Mtb), the causative agent of human tuberculosis, remains the leading infectious killer globally with an estimated death of 1.3 million in 2022[1]. Despite progressive work on designing new anti-TB therapeutics and implementing vaccination programs in TB-endemic countries, it has a high global case fatality rate and a poor treatment success rate, along with a rising number of drug-resistant infections[2, 3]. Emerging paradigms in infectious diseases advocate tackling pathogen-driven ailments as one-dimensional problems, where the entire emphasis is given to pathogen elimination. A holistic understanding of how the host systems respond to the infection, vaccination, and treatment is key to TB management programs[4]. Recent widespread and severe physiological derangements associated with COVID-19 patients, even after the elimination of the virus, have brought back the focus on identifying novel strategies that are inclusive of modulating the host immune-metabolic axis[5, 6].

The clinical symptoms of pulmonary TB encompass localized manifestations like prolonged cough with mucus, pleuritic chest pain, hemoptysis, and lung damage. Besides, systemic outcomes like cachexia, progressive fatigue, oxidative stress, altered microbiota and glucose intolerance result in organ-wide disruptions[7, 8]. Pulmonary TB patients often suffer from progressive and debilitating loss of muscle mass and function with severe weight loss, this TB-associated cachexia cannot be reversed by conventional nutritional support[9, 10]. Besides, numerous epidemiological studies indicate that hyperglycemia may occur during active tuberculosis, which can compromise insulin resistance and glucose tolerance, although the mechanisms are unclear[7, 11, 12]. Both the localized and systemic pathophysiology of TB infection indicate an alteration in the

host immuno-metabolic axis. It is somewhat bewildering that the engagement of the liver

during the Mtb infection cycle is not considered, despite its central role in balancing the

immune and metabolic functions of the body[13]. The crosstalk between the liver and lung

has been largely overlooked in tuberculosis (TB), even though acute phase proteins

(APPs) are used as predictive biomarkers in pulmonary tuberculosis[14]. A robust hepatic

APR response in mice, mediated by key hepatocyte transcription factors, STAT3 and NF-

κB/RelA, has been known to trigger pulmonary host defenses for survival during

pneumonia and sepsis[15]. In TB, the active phase of the disease is associated with

heightened expression of various genes that modulate flux in the lipid metabolic

pathways[16-18].

The liver is involved in a wide variety of functions – synthesis of plasma proteins, secretion

of various hepatokines, degradation of xenobiotic compounds, and storage of lipids,

glucose, vitamins, and minerals [19, 20]. De-novo lipogenesis, secretion of acute phase

proteins, hepatokine production, etc. are all directly or indirectly controlled by the

hepatocytes, thereby communicating with almost all the organs of the body[19, 21].

Moreover, the liver is actively involved in triacylglycerol synthesis and storage under the

intricate regulation of various hormones like insulin, glucagon, and thyroid hormone[22].

All these functions uniquely position the liver as the central regulator of lipid metabolism.

To avert organ damage, the liver maintains tolerogenic properties rendering it an attractive

target for various pathogenic microorganisms. Although several studies have indicated

the role of both -Mtb virulence components and host factors (immune activation, nitric

oxide, IFNγ, intracellular pH and hypoxia) in the generation of Mtb drug tolerance, the role

of liver, being the principal center for xenobiotic metabolism needs careful

investigation[23-27]. Liver is the hub of both phase I and phase II drug modifying enzymes

(DMEs). The levels as well as the activity of both type of DMEs play significant roles in

determining the pharmacokinetics and efficacy of various drugs across multiple diseases,

from infection to malignancy[28].

In this study, we demonstrate the active involvement of the liver in a murine aerosol TB

infection model during the chronic phase and establish hepatocytes as a new replicative

niche for Mtb. Using a variety of *in vivo*, *ex vivo*, and *in vitro* techniques we show how Mtb

perturbs biological functions within hepatocytes remodeling intracellular growth,

localization, and drug sensitivity. Cellular and mass spectrometric studies demonstrate

Mtb infection-mediated enhanced fatty acid biogenesis and TAG biosynthesis in the

hepatocytes regulated by PPARγ. We propose that infection of hepatocytes by Mtb during

the chronic phase can contribute to significant changes in disease progression, TB

treatment, and development of infection-induced metabolic diseases.

Results

Human pulmonary tuberculosis patients harbor Mtb in the liver.

Mtb infects lungs, and other organs like lymph nodes, pleura, bones, and meninges.

There are also isolated case reports of hepatic TB, without providing many

pathophysiological consequences[29, 30]. To gain further insights into the involvement of

the liver in Mtb infections, we acquired human autopsied liver samples from individuals

with miliary tuberculosis and analyzed them for the presence of Mtb bacilli. Hematoxylin

and eosin (H and E) staining showed the presence of distinct immune cell infiltration and

granuloma-like structures in the infected samples **(Fig 1A and Fig S1C)**. We examined

the liver specimens with Mtb specific Ziehl-Nielsen (Z-N) acid-fast and auramine O-rhodamine B stain (**Fig 1B, C**). Both these stains showed distinct positive signals with characteristic rod-shaped bacilli that could be visualized by the acid-fast staining (as indicated by arrows) **(Fig 1B).** We further corroborated our findings by performing fluorescence in-situ hybridization (FISH) using Mtb-specific 16s rRNA probes, where specific signals were observed **(Fig 1D).** The specific staining in FISH eliminates the possibility of non-tuberculous mycobacteria (NTMs) in the tissue specimen and confirms the presence of Mtb infection in the liver. Similar staining in the uninfected liver sections from other individuals did not show any signal. **(Fig S1 A, B**). Although hepatic granuloma is the characteristic histological feature for both local and miliary forms of hepatic TB, the precise involvement of the different cells like Kupffer cells, hepatocytes, stellate cell, liver sinusoidal endothelial cells, hepatic stellate cells, and other cell types have not been studied in detail[31]. Multiplex immunostaining with β-actin antibody and Mtb-specific Ag85B antibody shows the presence of Ag85B signals within the human hepatocytes further confirming the presence of Mtb within hepatocytes (**Fig 1E and Fig S1D)**, indicated with yellow arrows). Hepatocytes are morphologically distinct, large polygonal cells (20-30 μm) with round nuclei, many of which are double-nucleated and mainly positioned at the center of the cytoplasm[32]. The presence of Ag85B signals near the nucleus, as depicted in (**Fig 1E and S1 D**), further supports our assertion. Moreover, hepatic granulomas in the human samples showed localized clustering of the immune cells **(Fig S1D).** Furthermore, to establish a correlation between liver Mtb load and Mtb burden at the primary infection site, the lung, we conducted H&E staining and acid-fast staining on lung sections from the same individuals. H&E staining revealed distinct

granulomas, while acid-fast staining confirmed an elevated bacterial load, both indicative

of a high degree of infection **(Fig S1E, F).** Using auramine O-rhodamine B, Ziehl-Nielsen

acid-fast staining, and FISH, Mtb presence was detected in several lung specimens **(Fig**

S1G). These results indicate Mtb infection in the liver of human subjects and suggest the

localization of Mtb within hepatocytes. These findings are quite intriguing considering very

few studies have discussed the lung-liver cross talk or the involvement of liver in

pulmonary TB.

Mtb infection of mice via the aerosol route leads to significant infection of the liver

and primary hepatocytes.

To investigate whether the liver harbors Mtb during mice aerosol infection, we infected

C57BL/6 mice with 200 CFU of Mtb *H37Rv* and scored for the bacterial load in the

conventional niche- the lung as well as in the liver. Mtb could be detected in the liver

consistently across several experiments in 4 weeks post-infection and the bacterial load

increased till week 10 **(Fig 2A and B).** Consistent with the Mtb burden, phalloidin and

hematoxylin, and eosin (H and E) staining of the infected liver at 8 weeks post-infection

showed localized cellular aggregation forming ectopic granulomas forming a granuloma-

like structure **(Fig 2C and S2C).** Further staining the liver sections with CD45.2 antibody

shows clustering of immune cells in Mtb infected mice liver **(Fig S2D).** To assess whether

liver infection leads to deranged liver function in the aerosol model, we analyzed the levels

of liver functional enzymes like albumin aspartate transaminase (AST), alanine

transaminase (ALT), and gamma-glutamyl transpeptidase (GGT) in the sera. Till week 10

post-infection, sera levels of these markers remain almost unchanged at the different time

points post-infection **(Fig S2B).** Since hepatocytes, the principal parenchymal cells of the

liver constitute 70-80 percent of the liver by weight, we hypothesized whether hepatocytes

could harbor Mtb, in the murine model of infection. To this end, we isolated primary

hepatocytes from the mice at different time points post-infection. The purity of the isolated

hepatocytes was validated by multiple methods- morphologically hepatocytes can be

identified by their distinct hexagonal architecture, round nucleus, some of which are

binucleated as observed in Alexa fluor phalloidin 488 and DAPI stained hepatocytes **(Fig**

2E). Flow cytometry staining with antibody against hepatocyte specific marker

asialoglycoprotein receptor 1 (ASGR1) protein, confirmed that almost 100 perfect of the

isolated cells are primary hepatocytes as seen in **(Fig 2F)**. It was further validated by

confocal microscopy as all the cultured cells specifically stained for ASGR1 protein **(Fig**

2G). After thorough confirmation of the identity and the purity of the hepatocytes, the cells

were lysed and plated. Like the whole liver CFU, Mtb infected PHCs at 4 weeks post-

infection with substantial bacterial load at week 6 and week 8 **(Fig 2H).** Staining with

antibody for Mtb-specific Ag85B protein in both the infected tissue sections and cultured

hepatocytes isolated from the *in vivo* infected mice revealed the presence of Mtb within

hepatocytes **(Fig 2D and 2I)**. Besides the aerosol route, infection intraperitoneally also

led to hepatocyte infection, also led to hepatocyte infection as early as day 10 **(Fig S2A).**

Further, to prove the spread of Mtb to the liver in other model organisms, we infected

guinea pigs with 200 CFU of Mtb and analyzed the bacterial load in the lung, liver, and

spleen at week 4 and week 8 post-infection **(Fig S3A)**. At both the time points we could

observe robust liver infection with granulomatous structure in the liver, proving that the

dissemination of Mtb to the liver occurs across multiple model animals **(Fig S3B).**

Hepatocytes provide a replicative niche to Mtb.

Consistent data from human TB patients and aerosol murine model prompted us to develop an in vitro model of Mtb-hepatocyte. We examined Mtb infection in several hepatocyte cell lines of mouse and human origin along with primary murine hepatocytes. Infection studies with fluorescently labeled Mtb *H37Rv* were carried out with primary murine hepatocyte cells (PHCs), HepG2, Huh-7, and AML-12 **(Fig 3A, B, and C).** The multiplicity of infection was titrated both in PHCs and HepG2 with the MOI of 1, 2.5, 5, and 10. MOI 10 consistently showed an infectivity of more than 60% in both HepG2 and PHCs **(Fig S4 A, B, and C).** While we acknowledge that MOI 10 is moderately high, several well-cited studies have used MOI 10 for various conventional phagocytes and other non-conventional cell types like mesenchymal stem cells, adipocytes and human lymphatic endothelial cells[33-37]. Hence MOI 10 was selected for further experiments Macrophage cell lines RAW 264.7, THP-1, and mouse bone marrow-derived macrophages (BMDMs) were used as positive controls. Even though PHCs, HepG2, Huh-7, and AML-12 are not considered to be classical phagocytic cells, all cells showed infectivity of more than 60 percent after 24 hours, comparable to RAW 264.7, THP-1 and BMDMs **(Fig 2A, B).** Analysis of bacterial load in the PHCs post-infection showed colony-forming units (CFU) like RAW 264.7, THP-1, and BMDMs, supporting microscopic observations **(Fig 3C).** Mtb in macrophages is known to remodel the intracellular environment to survive within phagosomes. We studied Mtb growth kinetics within hepatocytes using GFP-labeled Mtb *H37Rv* in PHCs and HepG2 **(Fig 3D and F).** Mean fluorescent intensity measurements showed a consistent increase in GFP intensity in both PHCs and HepG2 with increasing time **(Fig 3E and G).** Fold change in replication

dynamics with respect to 5 HPI showed that, while bacterial growth in macrophages

plateaus after 48 hours post-infection, Mtb continues to grow in both HepG2 and PHCs

(Fig 3H). Our studies thus establish that hepatocytes, besides being robustly infected by

Mtb, also provide a favorable replicative niche for Mtb.

Transcriptomics of infected hepatocytes reveal significant changes in key

metabolic pathways.

To understand Mtb-induced changes in the hepatocytes and the underlying mechanisms

of how hepatocytes provide a favorable environment to the pathogen, we performed

transcriptomic analysis of the infected and sorted HepG2 cells at 0 hours (5 hours post

incubation) and 48 hours post-infection. Sorting before RNA isolation specifically enriches

the infected cellular population thus eliminating cellular RNA from uninfected cells **(Fig**

4A). Unsupervised clustering segregated the data into 4 distinct groups on the PC1 with

a variance of 27 percent, showing good concordance within the replicates **(Fig 4B).** The

close spatial clustering for the two 0-hour time points corresponding to uninfected and

infected is indicative of relatively less transcriptomic changes. On the other hand, the

spatial segregation of the 48 hours datasets suggests clear differences between the RNA

transcripts of uninfected and infected cells. The differentially expressed genes were

calculated using DE seqR with fold change >0.5 and a false discovery rate of <0.2. Gene

ontology (GO) enrichment analysis for the differentially regulated pathways at both the

early (0 hours) and the late (48 hours) infection time points is shown in **(Fig 4C).** At 0

hours post-infection, the immediate stress response pathway of the cell, involving ROS

generation, intracellular receptor signaling pathways, and response to xenobiotic stresses

got activated, while at a late time point, Mtb modulated some of the key immuno-metabolic

pathways like macroautophagy, cellular respiration, proteasomal degradation pathway,

response to type I interferon, IκB kinase/NF-κB signaling, etc **(Fig 4D).** Major alterations

in the vacuolar and vesicular transport at 48 hours are indicative of the dynamic changes

in the phagosome maturation pathway **(Fig 4D).** Volcano plot analysis showed greater

relative changes in the gene expression pattern at 48 hours compared to 0 hours, with

many genes like *CXCL10, CXCL11, IDO, CCL5* etc being greatly upregulated **(Fig 4E,**

4F). Interestingly, our RNA sequencing data indicated major changes in various facets of

lipid metabolic pathways like fatty acid biosynthesis pathway, glycerolipid and

glycerophospholipid metabolism, cholesterol biosynthesis pathways, etc. Several key

genes like *FASN, DGAT1, DGAT2, HMGCR*, etc. were upregulated, indicating the

possibility of greater synthesis of neutral lipids. Thus, transcriptomic studies shed light on

several of the key Mtb-induced changes in the hepatocytes.

To understand the difference and similarities between the pathways that get affected in

macrophages compared to hepatocytes during Mtb infection, we compared the gene

expression analysis data from Mtb-infected HepG2, 48 hours post-infection with THP-1

infected macrophages, 48 post-infection, taken from Kalam et al.,[35] **(Fig S5 A, B).** The

comparison provides insights regarding how Mtb modulates different pathways

depending on the type of the host cell infected. In THP-1, most of the altered pathways

are related to mounting an effective immune response to the bacteria like type 1 interferon

signalling pathway, regulation of cytokine secretion, leukocyte chemotaxis, response to

zinc, etc while in HepG2 the pathways that are getting altered are related to metabolism

like response to xenobiotic stimulus, macroautophagy, glycerophospholipid and

glycerolipid metabolism, and cellular respiration. This comparative analysis is indicative

of Mtb's ability to harness the metabolic richness of hepatocytes probably as a source for

nutrients. Moreover, being a non-immune cell type hepatocytes might lack a robust innate

immune pathway like the macrophages and hence be less likely to clear mycobacterial

infection.

Increased fatty acid synthesis drives Mtb growth in hepatocytes.

Mtb survival in foamy macrophages is driven by nutrient acquisition from the lipid

droplets[36]. Transcriptomic studies of Mtb-infected cells also showed upregulated

pathways for lipid metabolism. Examination of Lipid droplets in both PHCs and HepG2

revealed an increase in the number of lipid droplets at 24 hours post-infection **(Fig 5A**

and B). Time kinetic analysis of BODIPY intensity in the infected HepG2 at different days

post-infection indicated a concomitant increase in lipid droplets with the progress of

infection **(Fig 5C).** Moreover, in PHCs, GFP labeled Mtb showed a high degree of

colocalization with the lipid droplets **(Fig 5D).** Lipid droplets are single membrane-bound

depots consisting mainly of neutral lipids like diacylglycerols (DAGs), triacylglycerols

(TAGs), and cholesterol esters (CEs)[38].

Mass-spectrometric analysis of the infected and uninfected HepG2 cells at 24 hours post-

infection, showed an increase in both TAGs and DAGs and CEs with a decrease in the

levels of free cholesterols, indicating Mtb-induced changes in the neutral lipid

biosynthesis **(Fig 5E).** To understand whether Mtb utilizes host lipid droplets as a source

of nutrients in hepatocytes, we treated Mtb-infected hepatocytes with specific inhibitors

of de-novo fatty acid biosynthesis (C75) and TAG biosynthesis (T863) **(Fig S6 A).**

Interestingly, inhibiting both de-novo fatty acid biosynthesis as well as TAG biosynthesis

reduced the bacterial load in both PHCs and HepG2 by almost 1.0-1.5 log fold **(Fig 5G**

and H). In THP-1 macrophage, although C75 reduced bacterial load by 0.5 log fold but

T863 did not affect the bacterial load **(Fig 5I).** To get better insights into the nutritional

dependency of intracellular Mtb on hepatocyte lipid source, we metabolically labelled

HepG2 with 7.5 µg/ml of fluorescently tagged fatty acid (BODIPY 558/568 C_{12}) which

subsequently accumulated into host lipid droplets. After 16-20 hours, the labelled cells

were treated with T863 and C75. Although T863 significantly reduced the load of the

intracellular TAGs, C75 had little effect on the level of accumulated TAGs and showed a

phenotype like the DMSO control **(Fig S6 C and D).** We infected these three set of cells

with Mtb (MOI:10) for 24 hours. The Mtb isolated from DMSO and C75 treated HepG2

became distinctly labelled with highly fluorescent lipid bodies, but the fluorescence signal

in Mtb derived from T863 treated cells was quite low and sparse **(Fig 5F and S7 A).**

Considering that host-derived fatty acids are converted to triacylglycerols (TAGs) in the

bacterial cytoplasm as a source of carbon and energy, we wanted to check whether the

TAG synthesis machinery was modulated in the hepatocyte-infected Mtb. To this end, we

isolated Mtb from HepG2, 48 hours post-infection, and analysed for key genes involved

in TAG biosynthesis. Mtb grown in DMEM was used as a control. Interestingly, *Tgs1,*

Tgs4, Rv 1760, etc were upregulated by 6-8-fold, indicating an active Mtb transcriptional

change in Mtb to utilize and store host-derived lipids **(Fig S6 B).** Cumulatively, our studies

thus demonstrate fatty acid biosynthesis and TAG formation to be important for Mtb

growth in hepatocytes.

To assess the status of the neutral lipids in the liver, BODIPY staining was conducted in

the liver cryosections of uninfected and infected mice (8 weeks post-infection). The liver

of Mtb-infected mice bears more lipid bodies **(Fig S8 A).** Co-immunostaining of BODIPY

and Ag85B also indicated a certain degree of colocalization of Mtb and Lipid bodies in the

liver **(Fig S8 B).** Next, we stained the liver of Mtb-infected mice at 2-, 4- and 8-weeks

post-infection with LipidTOX neutral red dye. Surprisingly we found an elevated signal

intensity of the dye at 8 weeks post-infection. This shows that in mice, Lipid droplets might

correlate with Mtb burden **(Fig S8 C, D).** Our data comprehensively establishes infection-

induced lipid droplet accumulation as a pathogenic outcome of Mtb involvement in the

liver during the chronic stage of infection.

PPARγ upregulation in Mtb-infected hepatocytes leads to augmented lipid

biogenesis.

To understand the molecular mechanism behind the accumulation of lipid droplets in the

Mtb-infected liver at 8 weeks post-infection, we investigated the expression patterns of

the transcription factors involved in the regulation of the genes of lipid biogenesis in our

RNA-seq data. At 48 hours post-infection, the transcript levels of Peroxisome proliferator-

activated receptor-gamma (PPARγ) were upregulated by 5-6-fold. To validate, that we

performed quantitative real-time PCR analysis of the *PPAR γ* gene in the infected HepG2

at 48 hours post-infection. With respect to the uninfected control, *PPAR γ* was upregulated

by 3-4-fold. Besides, downstream adipogenic genes that are directly or indirectly

controlled by *PPARγ* like *MGAT1*, *FSP27*, *FASN*, *DGAT1*, *DGAT2*, *ACAT1, ACAT2,*

ADIPSIN etc were all upregulated by more than 2-fold in the infected cells. **(Fig S9 A).**

Immunoblot also revealed a greater level of PPARγ protein in primary mouse hepatocytes

at 24 hours and 48 hours post-infection in the Mtb infected cells **(Fig 6C, D)** .In the in-vivo

Mtb aerosol infection model, *Pparγ* expression levels showed an intriguing trend, at week

2 post-infection, the expression level was comparable to the uninfected control, while at

week 4 post-infection, the expression level spiked to 1.5-2-fold, reaching 3-4 fold at week

8 **(Fig 6B).** The pattern of expression of *Ppary* in the infected liver correlated with the

bacterial load in the liver, where we see the induction at week 4 when Mtb reaches the

liver and maximum expression at week 8 when the load of Mtb is considerably high.

Besides, PPARγ, some of the critical enzymes involved in fatty acid biosynthesis, TAG

biosynthesis, cholesterol esterification like *Fasn, Dgat1, Dgat2, Mgat1, Acat1, Acat2* etc

showed a temporal increase in expression levels at the later time points post-infection

(Fig 6A). We then used a specific inhibitor of PPARγ, GW9662 (20 µM) and agonist of

PPARγ, rosiglitazone (20µM) in infected HepG2 and quantified the bacterial load after 48

hours. Surprisingly, inhibition of PPARγ decreased the bacterial load by almost 2-fold,

while chemically inducing PPARγ increased the bacterial load considerably **(Fig 6 E and**

F, and S9B). Moreover, LD number in cells also directly correlated with the levels of

PPARγ **(Fig 6G and H).** To investigate whether, PPARγ expression was also induced in

the liver of infected mice, we examined PPARγ in the liver post infection. Interestingly, we

found enhanced expression PPARγ in the liver of the mice at 8 weeks post infection **(Fig**

S9 C and D). Moreover, PPAR-γ intensity in hepatoctyte was also high in the infected

liver **(Fig S9 E).** Thus, PPARγ activation resulting in lipid droplets formation by Mtb might

be a mechanism of prolonging survival within hepatocytes.

Hepatocyte resident Mtb displays a drug-tolerant phenotype.

The success of Mtb as a pathogen is partially attributed to its ability to survive and adapt to antibiotics[39]. Again Mtb-induced lipid accumulation in the liver might influence the activity of the various anti-TB drugs[40]. Hepatocytes contain both phase I and phase II drug metabolizing enzymes (DMEs). Further, the pharmacological potency of lipophilic drugs is determined by the rate at which these drugs are metabolized to inactive products[41, 42]. Cytochrome P450 monooxygenases (cyp450s) system is the key phase I DMEs and known to interact with rifampicin. We therefore analysed whether Mtb infection influences the cyp genes in hepatocytes. *CYP3A4* and *CYP3A43* respectively, both of which metabolize anti-TB drug rifampicin were upregulated in Mtb infected HepG2 by approximately 4-fold and 2-fold respectively **(Fig S10A).** Moreover, *NAT2* gene responsible for N-acetylation of isoniazid was also upregulated in the Mtb infected HepG2 by almost 2-fold) **(Figure S10 A).** We therefore argued that hepatocyte resident Mtb may display higher tolerance to rifampicin. Towards this, we treated Mtb-infected HepG2 and PHCs with different concentrations of Rifampicin (0.1, 0.5, 5 µg/ml) for 24 hours and CFU enumerated the bacterial after lysis. RAW 264.7 was kept as macrophage control with the similar experimental setup. The percentage of bacteria which survived the drugs was the drug tolerant population. Both HepG2 and PHCs resident Mtb were significantly tolerant to (25-30%) to rifampicin, as compared to the macrophages **(Fig 7A and B).** Almost 10 percent of the bacterial population in HepG2 display a tolerogenic phenotype at the highest antibiotic concentration **(Fig 7B).**

We also examined Mtb susceptibility to isoniazid (INH), which is predominantly metabolized (50–90%) via N-acetylation of its hydrazine functionality by arylamine N-

acetyltransferase 2 (*NAT2*)[43]. Interestingly, KEGG analysis of transcriptomic data suggested several genes in this pathway to be upregulated in hepatocytes infected with Mtb **(Fig S10 B).** Experimental studies indeed showed higher tolerance of Mtb to INH in both primary hepatocytes and HepG2 at different concentrations (0.1, 0.5, 5 µg/ml) **(Fig 7A and C).** Moreover, transcript levels of some of the key DMEs (were upregulated in the liver of the infected mice, 8 weeks post-infection like *Nat2, Cyp2e1, Slco1b2, Ces1, Ces2, and Aadac* (upregulated by almost 2-3-fold) **(Fig 7F).** To corroborate the CFU data, we repeated the same experiment **(Fig 7A)** and looked at the bacterial load within each PHC/BMDM using high-resolution confocal microscopy. Mouse BMDMs were as a macrophage control. Here also we have calculated the % tolerance as a ratio by measuring the mean fluorescent intensity of GFP-Mtb per hepatocyte treated with drug to MFI of GFP-Mtb per hepatocyte treated with DMSO (control). Extensive analysis of more than 200 cells showed distinctly presence of more Mtb within PHCs treated with both rifampicin and isoniazid than in BMDMs treated with the same concentration of the drugs **(Fig 7D, E and S11A).**

Drug tolerance is the phenomenon of Mtb surviving drug treatment for longer durations in the absence of any resistance mechanisms. It is a bacterial property influenced by both bacterial and host-induced factors[23, 44]. To check whether Mtb derived from hepatocytes, post-drug exposure has altered drug sensitivity or not, we conducted a resazurin microtitre assay with different dilutions of both rifampicin and isoniazid with Mtb derived from PHCs treated with DMSO, Mtb derived from PHCs post-exposure to rifampicin and Mtb derived from PHCs post-exposure to isoniazid. Mtb from all three different setups did not show any alterations in their MIC values, displaying no change in

drug sensitivity (**Fig S12 A**). The increased tolerance of Mtb within the hepatocytes can

also be attributed to the increased activity of the drug efflux pumps. Drug efflux pumps

like Rv1258c, Rv1410, and Rv1819c are known to get upregulated under drug stress

conditions, preventing the accumulation of the drugs and thereby decreasing drug

sensitivity. To this end, we isolated intracellular Mtb RNA from HepG2 treated with DMSO,

0.5 µg/ml rifampicin, and 0.5 µg/ml isoniazid and checked for the expression of key drug

efflux pumps like *Rv 1258c. Rv1410, Rv1819* etc in these three Mtb populations **(Fig S12**

B, C and D). Interestingly, quantitative RT-PCR analyses of these Mtb genes did not show

any upregulation of these transporters in Mtb derived from HepG2 exposed to rifampicin

and isoniazid, indicating that drug tolerance phenotype can be better attributed to host

intrinsic factors rather than Mtb efflux pumps. Our data show that Mtb-infection of

hepatocytes induces DMEs, and this host-specific extrinsic activation may result in

decreased bioavailability or increased inactivation of anti-TB drugs.

Discussion

Over the years, various anatomical and cellular environments conducive to Mtb infection

have been identified, particularly concerning its latency[45, 46]. Pulmonary TB typically

presents with classical symptoms such as persistent coughing and mucus production due

to lung involvement[7]. While systemic manifestations like weight loss, fatigue, and loss

of appetite are commonly linked to TB progression, the underlying factors driving these

metabolic derangements remain unexplored. Our study sheds light on hepatocytes as a

previously unrecognized site for Mtb survival and replication. We have observed that Mtb

infection disrupts hepatocyte lipid metabolism via PPARγ, leading to increased levels of

fatty acids and triglycerides that the bacteria utilize for sustenance and proliferation. We

hypothesize that liver involvement in chronic TB, characterized by lipid imbalance, may

contribute to TB-associated metabolic complications. Intriguingly, a recent investigation

demonstrated that hepatic PPARγ activation induces the expression of growth

differentiation factor 15 (GDF-15), a crucial regulator of weight loss observed in ketogenic

diets[47].

In a chronic disease like tuberculosis (TB), which persists over an extended period, it is

quite surprising that Mtb infection of hepatocytes has been ignored thus far. While

previous research has explored liver infection by Bacillus Calmette-Guérin (BCG) using

an intra-venous model in mice, the bacilli were predominantly found within tissue-resident

macrophages or Kupffer cells at 15 minutes and 2 days post-infection[48]. However, our

study consistently identified the presence of Mtb within hepatocytes in a murine aerosol

infection model after the fourth week. Similarly, in a guinea pig aerosol infection model,

liver infection was evident, marked by distinct granulomas by week 4. Furthermore,

analysis of human biopsy liver samples from pulmonary TB patients revealed ectopic

granuloma-like structures within the liver hepatocytes, with the presence of Mtb-specific

Ag85B signals within hepatocytes. Together, these findings underscore hepatocytes as a

novel niche for Mtb persistence, shedding new light on the pathogenesis of TB.

Interestingly, in infected armadillos, *M. leprae* is shown to infect hepatocytes and hijack

liver homeostatic, regeneration pathways to promote *de novo* organogenesis[49].

The increased expression of PPARγ induced by Mtb in hepatocytes results in elevated

levels of cholesterol esters (CE 16:0, 18:0, and 18:1), diacylglycerols (DAGs 36:1, 36:2,

and 34:1), and triacylglycerols (TAGs 18:1/36:2, 18:0/36:2, and 18:0/36:1), leading to

enhanced lipid droplets. Alongside the colocalization of labeled Mtb with lipid droplets in

primary hepatocytes, the upregulation of key TAG biosynthesis genes such as *Tgs1*,

Tgs4, and *Rv1760* in Mtb residing within hepatocytes underscores a transcriptional

rewiring within the pathogen to assimilate lipids. Using PPARγ agonists and antagonists,

corresponding decreases or increases in bacterial burden were observed, underscoring

the pivotal role of PPARγ in Mtb survival within hepatocytes. The lipid remodeling induced

by Mtb infection in hepatocytes is recapitulated in the murine aerosol model, where an

increased number of lipid droplets was observed at week 8, accompanied by localized

accumulation of immune cells and granuloma-like structures.

Previous studies have demonstrated that Mtb-infected macrophage populations can

induce PPARγ expression through the mannose receptor-dependent pathway leading to

enhanced lipid biosynthesis, production of anti-inflammatory mediators, and prevention

of host cell apoptosis[50, 51]. Recent findings have further underscored heightened levels

of PPARγ in peripheral blood mononuclear cells (PBMCs) from TB patients with elevated

cortisol levels and increased disease severity[52]. Moreover, what distinguishes

hepatocytes is their pivotal role in systemic lipid metabolism by regulating de novo fatty

acid synthesis, TAG synthesis, fatty acid oxidation, and the regulation of LDL and

VLDL[22, 53]. Mtb-mediated perturbation of these vital processes can lead to a sequela

of metabolic disorders such as non-alcoholic fatty liver disease (NAFLD), dyslipidemia

and insulin resistance. Accumulation of lipids within hepatocytes offers advantageous

conditions for Mtb survival. In macrophages, heightened intracellular lipid levels have

been observed to impede autophagy and the acidification of phagolysosomes, both

crucial for bacterial eradication[54]. Despite these parallels, comparative transcriptomic

analyses of infected HepG2 cells with THP1 cells highlight differences in pathways such

as vacuolar and vesicular transport, xenobiotic metabolism, macroautophagy, and cellular

respiration. Furthermore, hepatocytes secrete hepatokines, a class of proteins serving as

signaling molecules, with diverse roles in metabolism, inflammation, and energy

homeostasis, thereby influencing the host-pathogen interplay[13].

Additionally, we suggest that Mtb infection of hepatocytes creates a drug-tolerant

environment in the liver due to activation of DMEs, many of which are earlier shown to

metabolize the two frontline drugs, isoniazid and rifampicin. Moreover, lipid accumulation

in the liver might also indirectly alter the levels of the drug metabolizing enzymes as

reported in previous studies[40, 55]. Upregulation of transcripts is not only observed in

infected cells *in vitro*, but also in the livers of mice eight weeks after Mtb infection.

Particularly noteworthy is the upregulation of NAT-2, which controls the rate-limiting step

of acetylating isoniazid into acetylisoniazid. This metabolite is further processed into

acetylhydrazine and isonicotinic acid, ultimately reducing the drug's effectiveness against

Mtb. Similarly, key rifampicin-metabolizing esterase genes such as Ces1, Ces2, Aadac,

and transporter Slco1b2 exhibit upregulation, potentially influencing the drug's distribution

and metabolism in the body. The activation of DMEs can significantly modify the

pharmacokinetics and pharmacodynamics of anti-TB medications, resulting in suboptimal

drug concentrations and the emergence of drug-resistant strains[56].

Although animal models offer valuable insights into Mtb infection and TB pathology, they

fall short of fully replicating the complexities of human TB. Several recent studies in

COVID-19 patients have shown that hepatocytes get infected with SARs-CoV-2, thereby

increasing gluconeogenesis and hyperglycemia [57, 58].) With compelling reports

associating hepatic steatosis with the onset of type 2 diabetes mellitus (T2DM), we

speculate that TB-induced perturbations in lipid metabolism might predispose chronic TB

patients to T2DM and vice versa[59, 60]. Through our investigation, we propose that

future studies in human TB patients might scrutinize this metabolically rich hepatocyte

niche to understand multiple organ-wide derangements in TB pathogenesis.

Materials and Methods

Ethical clearances

Human studies:

Paraffinized non-Mtb infected, and Mtb-infected human autopsied liver specimens were

obtained from the Postgraduate Institute of Medical Education and Research,

Chandigarh, India. The Institutional Ethics Committee of CSIR-IMTech (Council of

Scientific and Industrial Research-Institute of Microbial Technology) approved all research

experiments involving human samples (Approval no [IEC (May 2021) #6]). The samples

utilized in the study are from a library of autopsied human specimens in PGIMER

(Postgraduate Institute of Medical Education and Research). The consent is provided by

the relatives of the deceased person while its submission to the PGIMER.

Animal studies:

All the mice (C57BL/6) were housed in the animal house at the National Institute of

Immunology (NII) before being transported to the TACF-ICGEB (Tuberculosis Aerosol

Challenge Facility), for subsequent animal infection studies. The animal experiments

were performed adhering to the institutional guidelines (Approval number: Institutional

Animal Ethical Committee, IAEC 543/20.). Female Hartley Dunkan guinea pigs were used

for infection studies in TACF-ICGEB in accordance with the institutional guidelines

(Approval number: Institutional Animal Ethical Committee, IEAC #440/17)

Confocal microscopy and Immunofluorescence measurements:

Lipid droplets were stained by BODIPY 493/503 dye. Primary mouse hepatocytes and

HepG2 cells were grown on 12 mm coverslips at respective densities as per the

experimental requirement. Cells were washed with 1X PBS (HiMedia M1452-500G), fixed

by 4% paraformaldehyde, and incubated at room temperature for 15 minutes. After

incubation, the cells were washed thrice with 1X PBS followed by staining with 10 µM

BODIPY dye in 1XPBS for 45 minutes at room temperature. After the staining, the excess

dye was removed by washing thrice with 1X PBS. To check for the acidified compartments

within the cells, LysoTracker Red DND-99 (L7528) was added to the cells at a

concentration of 500 nM for 30 minutes at 37° Celsius, followed by washing thrice with

1X PBS to remove the residual dye. The cells were fixed with 4% paraformaldehyde as

previously mentioned. For staining with various antibodies, the cells were permeabilized

with 0.2% Triton-x 100 (X-100-1L) for 15 minutes, followed by proper washing with 1X

PBS. The cells were then blocked by 2% BSA in 1X PBST for 1 hour at room temperature.

Post blocking, the cells were treated with primary antibody overnight at 4°Celsius,

followed by proper washing with 1X PBS. Secondary antibody (1:500 dilution) was added

to the cells for 1 hour at room temperature. Three washes with 1X PBS were given. The

nucleus was stained with DAPI (Sigma D9542-5MG) at 1ug/ml concentration for 20

minutes at room temperature. The excess DAPI was washed with 1X PBS followed by

mounting with ProLong Gold Antifade mountant (P36930). Images were acquired by Zeiss

LSM 980 Laser scanning confocal microscope.

Image Analysis:

Image Analysis:

Analysis was done using Image J and Zen blue software. Mean fluorescent intensity/ cell

was calculated by corrected total cell fluorescence (CTCF) = Integrated Density – (Area

of Selected Cell x Mean Fluorescence of Background readings). Signal intensity in tissue

sections were normalized to respective areas. For colocalization of Mtb with various intra-

cellular markers, each data point represents a single field with 6-8 infected cells. More

than 100 cells across 4 independent experiments were analyzed. Serial confocal sections

were acquired with Z-stacks spanning 10-15 uM. The percentage of Mtb in the respective

markers were calculated by calculating the Mtb voxels colocalizing in the respective

markers. The image was analyzed by Zen Blue software (Zeiss) and Image J.

For percentage tolerance of Mtb within hepatocytes, we have calculated the ratio by

measuring the mean fluorescent intensity of GFP-Mtb per hepatocyte treated with drug

to MFI of GFP-Mtb per hepatocyte treated with DMSO (control). BMDMs were used as

the macrophage control. More than 20 fields, each consisting of more than 4 infected

cells have been used for analysis. MFI/ cell was calculated using the Zeiss ZEN blue

image analysis software.

Tissue Immunofluorescence staining:

Paraffin sections

Five-micron thick sections of paraffin-embedded tissue sections were taken in poly-L-

Lysine coated slides (P0425-72EA). Deparaffinization was performed by heating the

slides at 50°C for 20 seconds (3 times) till the wax melts, followed by the subsequent

steps, 100% xylene (Merck, CAS-1330-20-7) for 10 minutes (3 times), xylene and

absolute ethanol (Merck, CAS- 64-17-5) for 10 minutes, 100% ethanol for 10 minutes,

70% percent ethanol for 5 minutes (2 times), 50% ethanol, distilled water for 5 minutes

(2minutes) and a final wash in 1x PBS for 5 minutes (2 times). Antigen retrieval was

performed in an antigen retrieval buffer (10mM Sodium Citrate, 0.05% tween-20, pH: 6)

by heating the slides at 60°C for 15 minutes. After antigen retrieval, permeabilization was

performed with 0.4% Triton-X 100 in 1X PBS for 20 minutes followed by proper washing

with 1x PBS. Blocking was done with 5% BSA for 1 hour. Sequential addition of primary

antibody was performed at 1:100 dilution at 4°C overnight. Primary antibody was washed

with 1X PBS followed by counter stain with DAPI nuclear stain at 1 ug/ml concentration.

Mounting was done with a drop of vectashield (sigma-aldrich, F6182-20ml). The slides

were visualised in Zeis LSM 980 confocal microscopy at 40X (oil) magnification.

Cryosections

Seven-micron thick cryosections were taken in poly-L-Lysine coated slides (P0425-

72EA). The sections were washed with 1X PBS, 3 times for 5 mins each. Permeabilisation

was done with 0.25% Triton-X 100 in 1X PBS for 15 minutes followed by washing with 1X

PBS. Blocking was done with 5% BSA for 1 hour followed by incubation with primary

antibody overnight at 4°C. After that the slides were washed with 1X PBS 2 times for 5

minutes each followed by incubation with fluorophore conjugated secondary antibody for

45 minutes. The slides were washed with 1X PBS followed by counterstain with DAPI at

1ug/ml concentration. Mounting was done with a drop of Vectashield (Sigma-Aldrich,

F6182-20ml). The dilution of the dyes and antibodies used in the staining are mentioned

in the table.

For staining with BODIPY, the sections were incubated with 15 ums of BODIPY for 40

minutes at room temperature. The slides were visualized in Zeis LSM 980 confocal

microscopy at 40X (oil) magnification.

Fluorescence in-situ hybridization:

FISH was used to detect Mtb in infected human liver following published protocols (1-3).

Briefly, the paraffinized human liver tissue sections were initially deparaffinized using a

serial washing step with xylene and ethanol, following which the sections were treated

with 1 mg/ml Proteinase K and 10 mg/ml Lysozyme in 10 mM Tris (pH 7.5) at 37°C for 30

min. Next, the samples were incubated in the Prehybridization buffer at 37°C for 1 h.

Prehybridization buffer is composed of 20% 2X Saline sodium citrate (SSC), 20% Dextran

sulfate, 30% Formamide, 1% 50X Denhardt's reagent, 2.5% of 10 mg/ml PolyA, 2.5% of

10 mg/ml salmon sperm DNA, 2.5% of 10 mg/ml tRNA. The slides were thoroughly

washed with a 2X SSC buffer. The sections were then incubated in hybridization buffer at

95°C for 10 min and then chilled on ice for 10 min. Further hybridization was allowed at

37°C overnight. Hybridization buffer is composed of prehybridization buffer plus 16S Mtb-

H_{37} Rv probe (5′ FITC – CCACACCGCTAAAG – 3′), which is specific for the 16S rRNA

of Mtb at a final concentration of 1 ng/μl. The liver tissue sections were next subjected to

a series of washing steps with 1X SSC at room temperature for 1 min, 1X SSC at 55°C

for 15 min, 1X SSC at 55°C for 15 min, 0.5X SSC at 55°C for 15 min, 0.5X SSC at 55°C

for 15 min, 0.5X SSC at room temperature for 10 min. Coverslips were mounted on glass

slides and visualized using Nikon A1R confocal microscope with a 488 nm laser.

Acid Fast Staining and Auramine O and Rhodamine B staining in liver sections.

Acid-fast staining was performed using a ZN Acid Fast Stains-Kit (K005L-1KT, HIMEDIA). Prior to staining, the paraffinized samples were deparaffinized using a serial washing step with xylene and ethanol 1. 100% Xylene for 6 min, 2. Xylene: Ethanol 1:1 for 3 min, 3. 100% Ethanol for 3 min, 4. 95% Ethanol for 3 min, 5. 70% Ethanol for 3 min, 6. 50% Ethanol for 3 min, 7. Distilled water. The glass slides were flooded with Carbol Fuchsin stain and heated to steam for 5 min with a low flame. The glass slides were allowed to stand for 5 min without further heating. The glass slides were then washed in running tap water. The glass slides were decolorized with acid-fast decolourizer for 2 min. 5. Washed with tap water. 6. Counterstain for 30 sec with Methylene Blue Washed with tap water, dried in air, and examined under 100x objective with oil immersion. The presence or absence of bacteria in infected and uninfected samples was checked through staining with Phenolic Auramine O-Rhodamine B dye (4) (1/3 dilution of stock solution) (Auramine O-861020-25gm, Sigma) (Rhodamine B-R6626-100gm, Sigma). Coverslips were mounted on glass slides and visualized using CLSM with a 488 nm laser.

Bacterial cultures and in-vitro experiments:

Virulent laboratory strains of H37Rv, BCG, and GFP-H37Rv bacterial cultures were cultivated on 7H9 medium (BD Difco) supplemented with 10% Oleic Acid-Albumin-Dextrose-Catalase (OADC, BD, Difco),0.05% glycerol and 0.05% tween 80 under shaking at 37°C conditions for in vitro assays. The cultures were then incubated in an orbital shaker at 100 rpm and 37 °C until the mid-log phase. pMN437-GFPm2 vector (Addgene, 32362) was used to electroporate the virulent H37Rv strain to create GFP-H37Rv, which

was then maintained in 50 μg/ml hygromycin 7H9-OADC medium. pMSP12: mCherry

plasmid (Addgene No. 30169) was electroporated in H37RV to generate Mtb-H37Rv-

mCherry.To prepare the single-cell suspension needed for infection tests, bacterial

cultures were passed through a sequence of different gauge needles five times through

23-gauge, 26-gauge, and three times through 30-gauge.

Human monocytic cell line THP-1 were obtained from American Type Culture Collection

(ATCC) and cultured in RPMI-1640 medium with 10% FBS at 37 °C and 5% CO_2

incubator. THP-1 derived macrophages were obtained by incubating THP-1 cells with 20

ng/ml phorbol 12-myristate 13-acetate (PMA, sigma) for 24 hours followed by washing

and maintenance in complete media. The cells were kept in non-PMA containing

complete media for 16-20 hours before infection with Mtb. HepG2, Huh-7, RAW 264.7

and PHCS were grown in DMEM under with 10% FBS at 37 °C and 5% CO_2 incubator.

AML-12 cells were cultured in DMEM media containing 1X insulin-transferrin-selenium

supplement (ITS-G), under the above-mentioned conditions.

In-vitro and ex-vivo infection experiments in primary cells and different cell lines were

performed at a Multiplicity of infection of 10 (MOI: 10) for both CFU enumeration and

confocal microscopy. The macrophage experiments involving THP-1 and RAW 264.7

involved incubating the cells with Mtb-H37Rv for 5-6 hours followed by washing with 1X

PBS and amikacin treatment (200 ug/ ml for 2 hours) to remove the extracellular bacteria.

The cells were kept for the designated time points for 24 hours, 48 hours and so on and

then lysed with lysis buffer (0.05 % SDS in 1X PBS) followed by plating in 7H11 plates.

For primary mouse hepatocytes and AML-12 cells, the cells were infected with the Mtb at

a multiplicity of 10 for 8 hours followed by amikacin treatment (200 ug/ ml for 2 hours).

For HepG2 and Huh-7, the time of incubation was 5- 6 hours followed by washing with

1X PBS and amikacin treatment as previously mentioned to remove the extracellular

bacteria.

The percent drug tolerant population was calculated using the following formula (A/B X

100) where A is the CFU in the drug treated group, B is the CFU in the untreated group.

The cells were morphologically checked for signs of cell death before proceeding with the

plating. For the inhibitor and inducer experiments, the cells were treated with the

respective drugs for 48 hours post uptake of Mtb. After that they were lysed and Mtb CFU

was enumerated.

Standardization of Multiplicity of infection (MOI) for PHCs and HepG2:

To standardise the MOI, HepG2 and BMDMs were infected with the following MOI of GFP-

Mtb as per the standardized protocol: 1, 2.5, 5 and 10. 6 hours post-infection, the cells

were trypsinised and fixed with 2% PFA. The percentage uptake was calculated by flow

cytometry. For, PHCs, the cells were infected at the above-mentioned MOI and

percentage infection was calculated by microscopy as (# of cells with GFP-Mtb/Total # of

cells *100)

C57BL/6 aerosol challenge:

The mice infection experiments were conducted in the Tuberculosis Aerosol Challenge

Facility (TACF, ICGEB, New Delhi, India). C57BL/6 mice were placed in individual

ventilated cages within the enclosure, maintaining a temperature of 20-25ºC, 30-60 %

humidity and 12h-12h of light-dark cycle. Following the standardized protocol, mice were

infected with 200 CFUs of H37Rv in a Wisconsin-Madison chamber. To ensure proper

establishment of infection, two animals were euthanized 24 hours post aerosol challenge.

The lungs were harvested and homogenised in 1X PBS and plated in Middlebrook 7H11

agar plates (Difco) supplemented with 10% OADC and 0.5% glycerol. CFU enumeration

was done three weeks post plating.

C57BL/6 peritoneal infection:

The mice were injected with 10^6 CFUs of H37Rv in 0.2 ml of 1X PBS. Following infection,

on different days post infection, the lung, spleen, and the liver was harvested and

homogenized and plated in Middlebrook 7H11 agar plates (Difco) supplemented with 10%

OADC and 0.5% glycerol. CFU enumeration was done three weeks post plating.

List of antibodies and dyes used in the study:

Serial number	Antibodies and dyes	Catalogue number	Company	Application and dilution
1.	Rab5	2143T	CST	IF/ICC (1:200)
2.	Rab7	D9F2	CST	IF/ICC (1:200)
3.	Cathepsin D	Ab75852	Abcam	IF/ICC (1:150)
4.	LAMP1	9091T	CST	IF/ICC (1:200)
5.	EEA1	3288T	CST	IF/ICC (1:200)

6.	ASPGR1	PA5-32030	Invitrogen	IF/ICC (1:500)
7.	PPAR γ	PA3-831A	Invitrogen	IF/ICC (1:150) WB (1:1000)
5.	Ag85B	Ab43019	Abcam	IF/ICC (1:800)
6	CD45-APC (clone 104)	558702	BD biosciences	1:300
7.	β-actin rabbit polyclonal Antibody.	4967	CST	IF/ICC (1:100) WB (1:2000)
8.	Goat anti-rabbit IgG (H+L) secondary Ab Alexa fluor 555 Plus	A32732	Thermo fischer Scientific	IF/ICC (1:400)
9.	Goat anti-rabbit IgG (H+L) Secondary Antibody, Alexa Fluor 488	A11008	Thermo fischer Scientific	IF/ICC (1:400)
10.	Goat anti-rabbit IgG (H+L) Secondary Antibody, Alexa Fluor 405	A-31556	Thermo fischer Scientific	IF/ICC (1:400)

11.	Lysotracker Red	L7528	Invitrogen	500 nM
12.	BODIPY-493/503	D3922	Invitrogen	Staining (10-15 µM)
13.	BODIPY 558/568 C12	D3835	Invitrogen	Metabolic Labelling (7.5 µg/ml)
14.	Phalloidin-Alexa fluor 488	A12379	Invitrogen	IF/ICC 1 unit/slide
15.	Phalloidin-Alexa Fluor 555 plus	A30106	Invitrogen	IF/ICC 1 unit/slide
16.	HCS LipidTOX Red Neutral Lipid Stain	H34476	Invitrogen	IF/ICC (1:400)
17.	DAPI	D9542	Sigma-aldrich	1 ug/ml
18.	ZN Acid Fast Stains	K005L-1KT	Himedia	
19.	Auramine O	861020-25gm	Sigma-Aldrich	

19	Rhodamine B	R6626-100gm	Sigma-Aldrich	

List of reagents in the study:

Serial no	Reagents	Catalogue number
1.	C75	C5490-5MG, Sigma-Aldrich
2.	OA-BSA	O3008-5ML, Sigma-Aldrich
5.	GW9662	M6191-25MG, Sigma-Aldrich
6.	Rosiglitazone	R2408-10MG, Sigma-Aldrich
7.	T863	SML0539-5MG, Sigma-Aldrich
8.	Rifampicin	557303-1G, Merck
9.	Isoniazid	I3377-5G, Merck
10.	PMA (Phorbol 12-myristate 13-acetate)	P8139-1MG, Merck
11.	Mouse MCSF (macrophage colony stimulating factor)	130-101-706 (25 µg), Miltenyi Biotec

12.	Insulin-Transferrin-Selenium (ITS - G) (100X)	41400045, Gibco

Sorting of labelled H37Rv cells

668 The infected hepatocytes were trypsined and washed with 1X PBS followed by passing

669 through 40µm cell strainer to make single cell suspension. After that the cells positive for

670 mCherry signals were sorted using the BD FACS Aria at TACF, ICGEB. Approximately 1

671 million cells were used for RNA isolation per sample.

Primary hepatocyte Isolation:

After euthanizing the mice, the peritoneal cavity was opened, and the liver was perfused

with 1X HBSS solution till exsanguination was complete. After that collagenase solution

(17 mg in 30 ml of 1X HBSS) was passed for digestion. The liver was cut into small pieces

and kept in 25 ml of DMEM media followed by lysing the tissues with a glass pestle and

sieve. The cells were then passed through a 70 µM cell strainer. 25 ml of Percoll was

added to the cells with proper mixing. It was centrifuged at 1000 rpm for 5 minutes. The

floating cells were removed, and a brownish layer of pure hepatocytes pelted at the

bottom. The cells were counted and plated according to the experimental need on a

collagen coated plate.

Isolation of Bone marrow derived macrophages (BMDMs) from mouse: BMDMs

used for Mtb infection studies were isolated following the published protocol with minor

modifications[61]. Briefly, the epiphysis of the femurs and tibia from C57BL/6 mice were

cut and the bone marrow were gently flushed into BMDM supplemented with 10% FBS

and 1% penicillin streptomycin. The bone marrow cells were centrifuged at 200 X g for 5

minutes at 4° C. The cell pellet was aspirated with 1X PBS and treated with RBC Lysis

buffer (for 3 minutes on ice, followed by the addition of complete media. The cells were

resuspended in 5 ml of the media and passed through a 70 μm cell strainer. The cells

were seeded in complete DMEM supplemented with 10ng/ml of MCSF. On the third day,

half of the media volume was replaced by fresh DMEM supplemented with 10ng/ml of

MCSF. On the 7th day, the cells were trypsinised, counted and seeded according to

experimental requirements.

Lipid Extraction protocol and Mass spectrometry:

5 million HepG2 cells were infected with Mtb-H37Rv at MOI 10 and kept for 24 hours and

48 hours post-infection. An equal number of uninfected cells were taken. The cells were

scrapped, and the procedure of Bligh and Dyer was followed[62]. In brief, the cells were

lysed in 1% Triton X-100 after being rinsed twice with 1X PBS. Following lysis, the lysate

was vortexed and four volumes of methanol-chloroform (2:1) were added. After that, one

volume each of water, chloroform, and 50 mM citric acid was added and vortexed.

Following a 10-minute centrifugation at 10,000 rpm at 4° Celsius, the lower organic phase

was collected and dried using liquid nitrogen. All semi-quantitative lipid measurements

were done using previously reported high-resolution MS/MS methods and

chromatographic techniques on an Agilent 6545 QTOF instrument. All sterols were

resolved using a Gemini 5U C-18 column (Phenomenex) while DAGs/TAGs were

resolved using a Luna 5U C-5 column (Phenomenex) using established solvent systems.

Cell lysate preparation and Western blotting:

The cells were washed twice with 1X PBS followed by lysis with SDS-RIPA buffer (50 mM

Tris-HCl pH 7.5, 150 mM NaCl, 1 mM EDTA, 0.1% SDS, 1% Triton-X 100, 1 mM DTT, 1X

Proteinase inhibitor). The cells were incubated with the buffer for 30 minutes in ice

followed by vortexing for 5 minutes. The supernatant fraction was collected by

centrifuging at 10000 rpm for 20 minutes at $4^{O}C$. The protein concentration was

determined by bicinchoninic acid (BCA) protein estimation kit (Thermo Fisher Scientific,

Waltham, Massachusetts, USA, 23227) following the manufacturer's protocol. 60-80 ug

of protein was resolved in SDS-PAGE followed by transferring onto a PVDF membrane.

Blocking was done in 5% skimmed milk in 1X TBST followed by incubation with primary

antibodies overnight at $4^{o}C$. The membranes were washed 3 times with 1x TBST for 10

minutes each followed by incubation with the HRP-conjugated secondary antibody for 1

hour. Immobilon HRP substrate was used to develop the blots and ImageQuant

Chemiluminescent imaging system (LAS 500) was used to acquire the images. Band

intensities were measured by using ImageJ.

MIC determination of Mtb isolated from primary hepatocytes and primary
hepatocytes treated with Rifampicin and isoniazid:

Mtb was isolated from primary hepatocytes, primary hepatocytes treated with 0.5µg/ml of

rifampicin and primary hepatocytes treated with 0.5µg/ml of isoniazid. These bacteria

were grown to a logarithmic phase of OD_{600}= 0.6 in Middlebrook 7H9 broth supplemented

with 10% OADC, 0.05% glycerol and 0.05% tween 80 under shaking conditions. Single-

cell suspension of the Mtb was made as previously described. Two-fold serial dilutions of

both rifampicin and isoniazid was prepared in 0.1 ml 7H9-OADC (without Tween 80) (The

concentration and the dilution of the drugs have been mentioned in the figure) in 96-well

flat bottom microplates. Approximately 5×10^4 Mtb cells were added to each well in a

volume of 0.1 ml. Control wells containing (Mtb only, medium+ inhibitor and only medium)

were included in the plate setup. The plate was incubated at 37°C 5 days, followed by the

addition of 20 µl of 0.02% resazurin for 24 hours. Visually, minimum inhibitory

concentration (MIC) was noted as the lowest drug concentration that prevented the

change of colour from blue to pink.

qRT-PCR of intracellular Mtb isolated from hepatocytes:

HepG2 cells were infected with Mtb at a MOI of 10 and at the respective time points post-

infection, the intracellular Mtb was isolated. In all experiments, the cells were treated with

amikacin to remove the extracellular bacteria, followed by subsequent washing by 1X

PBS. 4M Guanidine isocyanate was added to the flasks in equal amount to the media

and the cells were scrapped. The suspension was centrifuged at 3500 rpm for 7 minutes

and the pellet was resuspended in 350 µl of the RNA Lysis buffer of MN Kit (740955.250),

followed by beat beating. RNA was isolated as per the manufacturer's protocol. Isolated

RNA was treated with Dnase (PGM052, Puregene) for 1 hour to remove genomic DNA

contamination. RNA integrity was checked by running it on a 2% agarose gel with RNA

loading dye.

1 µg of RNA was reverse transcribed to cDNA using the Takara cDNA synthesis kit

(6110A) as per the manufacturer's protocol. Gene expression analysis by quantitative

real-time PCR was performed PowerUp SYBR Green PCR (Thermo fischer scientific,

(A25742) master mix in ABI 7500 FAST instrument. 16S rRNA was used as the

normalizing control and comparative Ct method was used for quantification.

Fluorescent fatty acid labelling of HepG2 and isolation:

HepG2 cells were incubated with 7.5 µg/ml of fluorescently tagged fatty acid (BODIPY

558/568 C_{12}) for 24 hours. The unincorporated fatty acids were removed by washing it

with 1X PBS, three times. After that the cells were treated with DMSO or inhibitors (T863

and C75) for 24 hours, followed by infection with Mtb at a MOI of 10, following the

standardized protocol of infection. A separate set of cells before infection were analysed

for the effect of the inhibitors on LD formation by microscopy.

After, 48 hours post infection, the cells were washed with 1X PBS and treated with Triton

X -100 (0.05% v/v in water), probe sonicated, and Mtb from the cells were isolated by

centrifuging at 3500 RPM for 8 minutes. Isolated Mtb cells were washed with 1X PBS and

fixed with 4% PFA and stained with DAPI (1mg/ml). The cells were mounted on poly-L-

lysine coated slides with antifade agent. The cells were visualised in Zeiss 980 LSM

confocal microscope at 100X, magnification.

Gene expression studies

RNA isolation

Total RNA was isolated from HepG2, PHCs and liver sections using the MN-NucleoSpin

RNA isolation kit (740955.250) following the manufacturer's protocol. For RNA

sequencing, an equal number of cells was used during isolation. For Liver tissue,

approximately 10mg was tissue was used. The quality of the RNA was verified by running

it on a 1.5% agarose gel and by monitoring 260/280 ratios. All the RNA samples were

frozen together in -80°C. For RNA sequencing, 3-5 ug of RNA was shipped in sodium

acetate buffer (3 M Sodium acetate, pH 5.2) with 100% Ethanol. 4 biological replicates

from each time point were sent for sequencing. The RNA samples with RNA integrity

number (RIN > 8.5) were used for library preparation.

RNA samples (for four biological replicates) were subjected to pair-end RNA sequencing

after rRNA depletion on the Illumina platform Novaseq-6000 at CCMB, Hyderabad, India.

Quality control and sequence trimming was performed using fastp (v0.23.2). The trimmed

paired-end reads were aligned to the human genome (GRCh38) using the HISAT2

(v2.2.1) pipeline. Reads were assembled into transcripts using StringTie (v2.2.1).

Annotation was conducted using aligned sequences and a GTF annotation file. The

mapped reads were then used for generating the count table using StringTie

(v2.2.1), genes lacking an Ensembl ID annotation were excluded. We arrived at a list of

62694 genes which were used for further analysis **(available through GEO accession-**

GSE256184). The differential count was performed by DEseq2 (R package) using the

uninfected samples of respective time points. Pathway enrichment was performed using

the GO database and clusters were visualized using the R package ClusterProfiler.

Further pathways of interest were analysed by GAGE and visualized using Pathview in

KEGG view.

cDNA synthesis and RT-qPCR

1ug of RNA was reverse transcribed to cDNA using the Takara cDNA synthesis kit (6110A) as per the manufacturer's protocol. Gene expression analysis by quantitative real-time PCR was performed PowerUp SYBR Green PCR (Thermo fischer scientific, (A25742) master mix in ABI 7500 FAST instrument. Beta-actin was used as the normalizing control and comparative Ct method was used for quantification.

List of primers used in the study.

GENE NAMES	FORWARD PRIMER (5'------------>3')	REVERSE PRIMER (5'------------>3')	
β-actin	ATGGAGGGGAATACAGCCC	TTCTTTGCAGCTCCTTCGTT	Mouse
ppar-γ	CTCTGGGAGATTCTCCTGTTGA	GGTGGGCCAGAATGGCATCT	Mouse
ppar-α	AGAGCCCCATCTGTCCTCTC	ACTGGTAGTCTGCAAAACCAAA	Mouse
fasn	CTGCGTGGCTATGATTATGG	AGGTTGCTGTCGTCTGTAGT	Mouse
acat1	CAGGAAGTAAGATGCCTGGAAC	TTCACCCCCTTGGATGACATT	Mouse
acat2	CCCGTGGTCATCGTCTCAG	GGACAGGGCACCATTGAAGG	Mouse
dgat1	TGCCCTGACAGAGCAGATGG	CAGGTTGACATCCCGGTAGG	Mouse
dgat2	TGGCGCTACTTCCGAGACTAC	TGCTGACTTCAGTAGCCTCTGTG	Mouse
nat2	ACACTCCAGCCAATAAGTACAGC	GGTAGGAACGTCCAAACCCA	Mouse
Slco1b2	GGGAACATGCTTCGTGGGATA	GGAGTTATGCGGACACTTCTC	Mouse
aadac	TACCGCTTCCAGATGCTATTGA	ACTGATTCCCAAAAGTTCACCAA	Mouse
cyp2e1	CATCACCGTTGCCTTGCTTG	GGGGCAGGTTCCAACTTCT	Mouse
ces1c	GCACTACGCTGGGTCCAAGAT	AAAGATGGTCACGGAATCCG	Mouse
ces2c	GCTGAATGCTGGGTTCTTCG	GCTGCCTTGGATCTGTCCTGT	Mouse
β-actin	AAGGCCAACCGCGAGAAGAT	GCCAGAGGCGTACAGGGATA	Human

pparγ	GGTGAAACTCTGGGAGATTCT	CTCTGTGTCAACCATGGTCA	Human
fasn	AGTACACACCCAAGGCCAAG	GTGGATGATGCTGATGATGG	Human
acat1	TACCAGAAGTAAAGCAGCATGG	TCATTCAGTGTACTGGCATTGG	Human
acat2	CTTTAGCACGGATAGTTTCCTGG	GCTGCAAAGGCTTCATTGATTTC	Human
dgat1	CAATCTGACCTACCGCGATCT	TCGATGATGCGTGAGTAGTCC	Human
dgat2	GAATGGGAGTGGCAATGCTAT	CCTCGAAGATCACCTGCTTGT	Human
nat2	CCAGAAGGGGTTTACTGTTTGG	CAGGTTTGGGCACGAGATTTC	Human
mgat1	CGCAAGTTCCAGGGCTACTAC	CTTCAGCAGCGGATAGGTGG	Human
fsp27	ATGGACTACGCCATGAAGTCT	CGGTGCTAACACGACAGGG	Human
Rv1258c	CTACGAGGCGATCCTCAACC	CGAGGATGGACAACCCGAAT	Mtb
Rv1819c	ATCGGGGTTTTCAGCGTGAT	GTCGAGCCAGTCTTGTGTGA	Mtb
Rv1218c	ATCGAGATTCGCGGACTGAC	ACATCGCCTGGAACATAGGC	Mtb
Rv1410c	ATGGTGACGCTGGTTGATGT	CGGTAGGGCGATAAGGAACC	Mtb
tgs1 (Rv3130c)	CCTTCTTATCGTCGCTCGCT	GCCAAGATCGAAGTCGGGAT	Mtb
tgs4 (Rv3088)	ACGGTCTCGTTCCTCAACAC	CCGGTGAATCGGAGAGGAAG	Mtb
lipY (Rv3097c)	TCTGCGCTCGAAACTCACTT	GCTCATCCCGTCATAGGTGG	Mtb
acrA1 (Rv3391)	ATGGCGGTCAACTACTTCGG	CCTTGGTGGGCAGATACGAG	Mtb

807 **List of reagents in the study**

Reagent	Company	Catalogue number
Immobilon Forte Western HRP substrate	Millipore	WBLUF0500
Amersham Hybond Western blotting membranes, PVDF	Merck	GE10600023
PowerUP SYBR	Thermo Fischer Scientific	A25742
Vectashield	Sigma-Aldrich	F6182
Primescript 1st strand cDNA synthesis kit	Takara Bio	6110A
NucleoSpin RNA isolation Kit	Macherey-Nagel	74106
ProLong Gold antifade reagent	Thermo Fischer Scientific	P36930
PolyFreeze Tissue Freezing medium	Sigma- Aldrich	SHH0026

Microscope slides	Himedia	BG005
Poly-Prep slides	Sigma- Aldrich	P0425-72EA
2X RNA loading dye	Thermo Scientific	R0641

Softwares

All the graphs have been generated using GraphPad Prism 10 and Microsoft Excel. The schematics have been drawn with the help of Biorender.com.

Acknowledgements

We thank Dr. Lakshyaveer Singh, Tuberculosis Aerosol Challenge Facility (TACF), ICGEB for mice experiments. We thank Dr. Neerja Wadhwa for helping in the NII Central Confocal facility, Mr. Birendra Nath Roy for the preparation of the Cryosections, and the NII animal facility for providing us with the animals. We thank the Next Generation Sequencing (NGS) facility at CSIR-CCMB for transcriptomic support.

Author Contribution

BS and RSG conceptualized the study and analysed the data. BS, DSG, JS, MY, PS, RDS performed the experiments. Human liver sections were stained by SS under the guidance from AK. Transcriptomic analysis was done by JS. Sorting was done by PS and RDS under the guidance from DK. Mass spectrometry analysis was conducted by AC under the supervision of SSK.KM conducted the guinea pig infection experiments. BS and RSG wrote the manuscript. BS, DM, DSG, AK, SSK, DK and RSG reviewed and edited the manuscript. RSG supervised the project.

825   **Declaration of Interests:** The authors declare no conflicts of interests.

826   **References**

827   1.   *Global Tuberculosis Report* 2023, World Health Organisation (WHO).
828   2.   Seung, K.J., S. Keshavjee, and M.L. Rich, *Multidrug-Resistant Tuberculosis and Extensively Drug-*
829   *Resistant Tuberculosis.* Cold Spring Harb Perspect Med, 2015. **5**(9): p. a017863.
830   3.   Husain, A.A., et al., *Current perspective in tuberculosis vaccine development for high TB endemic*
831   *regions.* Tuberculosis (Edinb), 2016. **98**: p. 149-58.
832   4.   Eckhardt, M., et al., *A systems approach to infectious disease.* Nat Rev Genet, 2020. **21**(6): p.
833   339-354.
834   5.   Davis, H.E., et al., *Long COVID: major findings, mechanisms and recommendations.* Nat Rev
835   Microbiol, 2023. **21**(3): p. 133-146.
836   6.   Lippi, G., F. Sanchis-Gomar, and B.M. Henry, *COVID-19 and its long-term sequelae: what do we*
837   *know in 2023?* Pol Arch Intern Med, 2023. **133**(4).
838   7.   Luies, L. and I. du Preez, *The Echo of Pulmonary Tuberculosis: Mechanisms of Clinical Symptoms*
839   *and Other Disease-Induced Systemic Complications.* Clin Microbiol Rev, 2020. **33**(4).
840   8.   Loddenkemper, R., M. Lipman, and A. Zumla, *Clinical Aspects of Adult Tuberculosis.* Cold Spring
841   Harb Perspect Med, 2015. **6**(1): p. a017848.
842   9.   Morley, J.E., D.R. Thomas, and M.M. Wilson, *Cachexia: pathophysiology and clinical relevance.*
843   Am J Clin Nutr, 2006. **83**(4): p. 735-43.
844   10.   Chang, S.W., et al., *Gut hormones, appetite suppression and cachexia in patients with pulmonary*
845   *TB.* PLoS One, 2013. **8**(1): p. e54564.
846   11.   Bisht, M.K., et al., *The cause-effect relation of tuberculosis on incidence of diabetes mellitus.*
847   Front Cell Infect Microbiol, 2023. **13**: p. 1134036.
848   12.   Niazi, A.K. and S. Kalra, *Diabetes and tuberculosis: a review of the role of optimal glycemic*
849   *control.* J Diabetes Metab Disord, 2012. **11**(1): p. 28.
850   13.   Jensen-Cody, S.O. and M.J. Potthoff, *Hepatokines and metabolism: Deciphering communication*
851   *from the liver.* Mol Metab, 2021. **44**: p. 101138.
852   14.   Kumar, N.P., et al., *Acute Phase Proteins Are Baseline Predictors of Tuberculosis Treatment*
853   *Failure.* Front Immunol, 2021. **12**: p. 731878.
854   15.   Quinton, L.J., A.J. Walkey, and J.P. Mizgerd, *Integrative Physiology of Pneumonia.* Physiol Rev,
855   2018. **98**(3): p. 1417-1464.
856   16.   Shim, D., H. Kim, and S.J. Shin, *Mycobacterium tuberculosis Infection-Driven Foamy Macrophages*
857   *and Their Implications in Tuberculosis Control as Targets for Host-Directed Therapy.* Front
858   Immunol, 2020. **11**: p. 910.
859   17.   Park, J.H., et al., *Understanding Metabolic Regulation Between Host and Pathogens: New*
860   *Opportunities for the Development of Improved Therapeutic Strategies Against Mycobacterium*
861   *tuberculosis Infection.* Front Cell Infect Microbiol, 2021. **11**: p. 635335.
862   18.   Gago, G., L. Diacovich, and H. Gramajo, *Lipid metabolism and its implication in mycobacteria-host*
863   *interaction.* Curr Opin Microbiol, 2018. **41**: p. 36-42.
864   19.   Wang, F., et al., *Organ-organ communication: The liver's perspective.* Theranostics, 2021. **11**(7):
865   p. 3317-3330.
866   20.   Alper, C.A., et al., *Human C'3: evidence for the liver as the primary site of synthesis.* Science,
867   1969. **163**(3864): p. 286-8.

21. Zhou, Z., M.J. Xu, and B. Gao, *Hepatocytes: a key cell type for innate immunity.* Cell Mol
Immunol, 2016. **13**(3): p. 301-15.

22. Zhang, D., et al., *Important Hormones Regulating Lipid Metabolism.* Molecules, 2022. **27**(20).

23. Datta, D., et al., *Actionable mechanisms of drug tolerance and resistance in Mycobacterium*
tuberculosis. FEBS J, 2024.

24. Deb, C., et al., *A novel in vitro multiple-stress dormancy model for Mycobacterium tuberculosis*
generates a lipid-loaded, drug-tolerant, dormant pathogen. PLoS One, 2009. **4**(6): p. e6077.

25. Liu, Y., et al., *Immune activation of the host cell induces drug tolerance in Mycobacterium*
tuberculosis both in vitro and in vivo. J Exp Med, 2016. **213**(5): p. 809-25.

26. Samuels, A.N., et al., *Understanding the contribution of metabolism to Mycobacterium*
tuberculosis drug tolerance. Front Cell Infect Microbiol, 2022. **12**: p. 958555.

27. Santucci, P., et al., *Intracellular localisation of Mycobacterium tuberculosis affects efficacy of the*
antibiotic pyrazinamide. Nat Commun, 2021. **12**(1): p. 3816.

28. Wu, K.C. and C.J. Lin, *The regulation of drug-metabolizing enzymes and membrane transporters*
by inflammation: Evidences in inflammatory diseases and age-related disorders. J Food Drug
Anal, 2019. **27**(1): p. 48-59.

29. Wu, Z., et al., *Diagnosis and treatment of hepatic tuberculosis: report of five cases and review of*
literature. Int J Clin Exp Med, 2013. **6**(9): p. 845-50.

30. Turhan, N., et al., *Hepatic granulomas: a clinicopathologic analysis of 86 cases.* Pathol Res Pract,
2011. **207**(6): p. 359-65.

31. Hickey, A.J., et al., *A systematic review of hepatic tuberculosis with considerations in human*
immunodeficiency virus co-infection. BMC Infect Dis, 2015. **15**: p. 209.

32. Celton-Morizur, S., et al., *Polyploidy and liver proliferation: central role of insulin signaling.* Cell
Cycle, 2010. **9**(3): p. 460-6.

33. Jain, N., et al., *Mesenchymal stem cells offer a drug-tolerant and immune-privileged niche to*
Mycobacterium tuberculosis. Nat Commun, 2020. **11**(1): p. 3062.

34. Lerner, T.R., et al., *Mycobacterium tuberculosis cords within lymphatic endothelial cells to evade*
host immunity. JCI Insight, 2020. **5**(10).

35. Kalam, H., M.F. Fontana, and D. Kumar, *Alternate splicing of transcripts shape macrophage*
response to Mycobacterium tuberculosis infection. PLoS Pathog, 2017. **13**(3): p. e1006236.

36. Peyron, P., et al., *Foamy macrophages from tuberculous patients' granulomas constitute a*
nutrient-rich reservoir for M. tuberculosis persistence. PLoS Pathog, 2008. **4**(11): p. e1000204.

37. Beigier-Bompadre, M., et al., *Mycobacterium tuberculosis infection modulates adipose tissue*
biology. PLoS Pathog, 2017. **13**(10): p. e1006676.

38. Wolk, M. and M. Fedorova, *The lipid droplet lipidome.* FEBS Lett, 2024. **598**(10): p. 1215-1225.

39. Day, N.J., P. Santucci, and M.G. Gutierrez, *Host cell environments and antibiotic efficacy in*
tuberculosis. Trends Microbiol, 2023.

40. Fisher, C.D., et al., *Hepatic cytochrome P450 enzyme alterations in humans with progressive*
stages of nonalcoholic fatty liver disease. Drug Metab Dispos, 2009. **37**(10): p. 2087-94.

41. Almazroo, O.A., M.K. Miah, and R. Venkataramanan, *Drug Metabolism in the Liver.* Clin Liver Dis,
2017. **21**(1): p. 1-20.

42. Ahmed, S., et al., *Pharmacogenomics of Drug Metabolizing Enzymes and Transporters: Relevance*
to Precision Medicine. Genomics Proteomics Bioinformatics, 2016. **14**(5): p. 298-313.

43. Sotsuka, T., et al., *Association of isoniazid-metabolizing enzyme genotypes and isoniazid-induced*
hepatotoxicity in tuberculosis patients. In Vivo, 2011. **25**(5): p. 803-12.

44. Rocha-Granados, M.C., et al., *The social network: Impact of host and microbial interactions on*
bacterial antibiotic tolerance and persistence. Cell Signal, 2020. **75**: p. 109750.

45.   Cosma, C.L., D.R. Sherman, and L. Ramakrishnan, *The secret lives of the pathogenic*
mycobacteria. Annu Rev Microbiol, 2003. **57**: p. 641-76.
46.   Bussi, C. and M.G. Gutierrez, *Mycobacterium tuberculosis infection of host cells in space and*
time. FEMS Microbiol Rev, 2019. **43**(4): p. 341-361.
47.   Lu, J.F., et al., *GDF15 is a major determinant of ketogenic diet-induced weight loss.* Cell Metab,
2023. **35**(12): p. 2165-2182 e7.
48.   Seiler, P., et al., *Limited mycobacterial infection of the liver as a consequence of its*
microanatomical structure causing restriction of mycobacterial growth to professional
phagocytes. Infect Immun, 2001. **69**(12): p. 7922-6.
49.   Hess, S., et al., *In vivo partial reprogramming by bacteria promotes adult liver organ growth*
without fibrosis and tumorigenesis. Cell Rep Med, 2022. **3**(11): p. 100820.
50.   Arnett, E., et al., *PPARgamma is critical for Mycobacterium tuberculosis induction of Mcl-1 and*
limitation of human macrophage apoptosis. PLoS Pathog, 2018. **14**(6): p. e1007100.
51.   Rajaram, M.V., et al., *Mycobacterium tuberculosis activates human macrophage peroxisome*
proliferator-activated receptor gamma linking mannose receptor recognition to regulation of
immune responses. J Immunol, 2010. **185**(2): p. 929-42.
52.   Díaz, A., et al., *Studies on the contribution of PPAR Gamma to tuberculosis physiopathology.*
Front Cell Infect Microbiol, 2023. **13**: p. 1067464.
53.   Bechmann, L.P., et al., *The interaction of hepatic lipid and glucose metabolism in liver diseases.* J
Hepatol, 2012. **56**(4): p. 952-64.
54.   Lovewell, R.R., C.M. Sassetti, and B.C. VanderVen, *Chewing the fat: lipid metabolism and*
homeostasis during M. tuberculosis infection. Curr Opin Microbiol, 2016. **29**: p. 30-6.
55.   Rey-Bedon, C., et al., *CYP450 drug inducibility in NAFLD via an in vitro hepatic model:*
Understanding drug-drug interactions in the fatty liver. Biomed Pharmacother, 2022. **146**: p.
112377.
56.   Dookie, N., et al., *Evolution of drug resistance in Mycobacterium tuberculosis: a review on the*
molecular determinants of resistance and implications for personalized care. J Antimicrob
Chemother, 2018. **73**(5): p. 1138-1151.
57.   Barreto, E.A., et al., *COVID-19-related hyperglycemia is associated with infection of hepatocytes*
and stimulation of gluconeogenesis. Proc Natl Acad Sci U S A, 2023. **120**(21): p. e2217119120.
58.   Mercado-Gomez, M., et al., *The spike of SARS-CoV-2 promotes metabolic rewiring in*
hepatocytes. Commun Biol, 2022. **5**(1): p. 827.
59.   Hazlehurst, J.M., et al., *Non-alcoholic fatty liver disease and diabetes.* Metabolism, 2016. **65**(8):
p. 1096-108.
60.   Dharmalingam, M. and P.G. Yamasandhi, *Nonalcoholic Fatty Liver Disease and Type 2 Diabetes*
Mellitus. Indian J Endocrinol Metab, 2018. **22**(3): p. 421-428.
61.   Toda, G., et al., *Preparation and culture of bone marrow-derived macrophages from mice for*
functional analysis. STAR Protoc, 2021. **2**(1): p. 100246.
62.   Bligh, E.G. and W.J. Dyer, *A rapid method of total lipid extraction and purification.* Can J Biochem
Physiol, 1959. **37**(8): p. 911-7.

Main figure legends:

Figure 1: Presence of Mtb in human pulmonary tuberculosis patients: (A) Hematoxylin and eosin (H and E) staining of the human autopsied liver tissue sections showing the presence of granuloma-like structure in the Mtb infected liver. **(B)** Acid-fast staining showing the presence of Mtb in the liver section of pulmonary TB patients. (arrows point to the presence of Mtb in the enlarged image) **(C-D)** Auramine O-Rhodamine B (A-O) staining and fluorescence in-situ hybridization (FISH) with Mtb specific 16s rRNA primer further confirms the presence of Mtb in human liver tissue sections. Data shown here is a representative of 5 autopsied specimens having Mtb infection. **(E).** Dual staining of β-actin (green) and Ag85B (red) using respective antibodies shows the presence of Mtb in hepatocytes of human biopsied liver sections.

Figure 2: Involvement of the liver and hepatocytes in a mouse aerosol model of Mtb infection. **(A).** Schematic denoting the flow of experimental set up of studying the liver at tissue level and cellular level (generated by Biorender.com). **(B)** C57BL/6 mice were infected with 200 CFU of H37Rv through the aerosol route and the bacterial burden of the lung and liver was enumerated at different time points post-infection in lung and spleen **(C).** Alexa Fluor 488 Phalloidin and DAPI staining of uninfected and infected lungs at 8 weeks post-infection. Images were taken in 20X, and 40 X magnification as mentioned (scale bar is 20 μm). **(D).** Immunofluorescence staining of Alexa Fluor 488 Phalloidin, DAPI, and Ag85B in the infected mice liver at 8 weeks post-infection shows the presence of Ag85B signals within hepatocytes (magnified image) (scale bar is 20 μm). **(E)**. Isolated primary hepatocytes stained with phalloidin green and DAPI show the typical polygonal shape with binucleated morphology. **(F).** Anti-asialoglycoprotein receptor

(ASGR1) antibody staining specifically stains primary hepatocytes isolated from infected

mice as validated by the contour plot in flow cytometry. **(G).** Anti-asialoglycoprotein

receptor (ASGR1) antibody staining specifically stains primary hepatocytes isolated from

mice as visualized through confocal microscopy. **(H).** Primary hepatocytes were isolated

from the infected mice, lysed and CFU enumeration was done at the mentioned time

points. **(I).** Ag85B staining of cultured primary hepatocytes, isolated from mice at 8 weeks

post Mtb infection. (scale bar is 20 μm). **(G)**. Animal litter size (n)= 6-7 mice in each group.

Figure 3: Mtb uses hepatocytes as a replicative niche: (A and B) Representative

microscopic images showing the infection of primary hepatocytes (PHCs), AML-12,

HepG2, and Huh-7 with labelled Mtb-*H37Rv* strains and subsequent quantification of

percentage infectivity in the respective cell types. RAW 264.7, THP-1 and murine BMDMs

were used as macrophage controls. The scale bar in all images is 10 μm except in HepG2,

which is 5 μm **(C)** CFU enumeration of Mtb-*H37Rv* in different hepatic cell lines. **(D and**

E) Representative confocal microscopy images of Mtb-GFP-*H37Rv* infected PHCs at the

respective time points post-infection and bar blot depicting mean fluorescent intensity

(MFI) / cell at the respective time points. **(F and G)** Representative confocal microscopy

images of Mtb-*H37Rv*-GFP infected HepG2 at the respective time points post-infection

and bar blot depicting MFI/cell at the respective time points**,** n= 4 independent

experiments, with each dot representing 5 fields of 30-60 cells. Scale bar 10 μm. Growth

kinetics of Mtb-*H37Rv* in **(H)** Fold change of growth rate of Mtb within HepG2, PHCs,

RAW 264.7, THP-1 and BMDMs.

Figure 4: RNA sequence analysis of infected and sorted hepatocytes at 0 hours and

48 hours post-infection (A) Experimental setup for infection of HepG2 with Mb-*H37Rv*-

mCherry at 10 MOI with histogram of mCherry signals at 0- and 48-hours post-infection(

Schematic depicting experimental set up generated with Biorender.com) **(B)** Principal

component analysis (PCA) plot illustrating the HepG2 transcriptome, identified through

global transcriptomic analysis of 0 hours uninfected and infected and 48 hours uninfected

and infected. **(C)** GO pathway enrichment analysis was done for DEGs with adjusted p-

value < 0.05 at 0- and 48-hours post-infection. The bubble plot depicts the enrichment of

pathways on the mentioned time points post-infection, where the coordinate on the x-axis

represents the gene ratio, the bubble size represents the gene count and colour

represents the p-value. **(D and E)** Volcano plot depicting the fold change of different

genes in 0 hours and 48 hours post-infection, the red dots represent the significantly

upregulated and downregulated genes.

Figure 5: **Increased fatty acid biogenesis drives Mtb growth in hepatocytes**: **(A)**

Increased number of LDs/cells in HepG2 and PHCs post-Mtb infection **(B).** Quantification

of the number of lipid droplets in the infected HepG2 and PHCs with their uninfected

control, 50-70 cells were analyzed from 3 independent experiments. **(C).** Increase in the

BODIPY intensity at different days post-infection in infected hepatocytes **(D)**. A high

degree of colocalization of Mtb-*H37Rv*-GFP (green) with lipid droplets (red) within PHCs

(E) Mass spectrometric quantification of the different species of DAGs, TAGs, and

cholesterol esters in the Mtb infected hepatocytes. **(F)**. Confocal images showing puncta

of fluorescently labelled fatty acid in Mtb derived from metabolically labelled (with BODIPY

558/568 C_{12}) HepG2**.** Relative CFU of Mtb under the administration of different inhibitors

of the lipid metabolic pathway in **(G)** PHCs **(H)** HepG2 and **(I)** THP-1. Representative data

from 3 independent experiments. Data were analyzed using the two-tailed unpaired

Student's t-test in B and one-way ANOVA in C, F, G, and H.*p < 0.05, **p < 0.005, ***p <

0.0005, ****p < 0.0001. ns=non-significant.

Figure 6: PPARγ driven lipid biogenesis drives Mtb growth in hepatocytes: (A) Heat

map showing the fold change of the genes involved in the lipid biosynthesis and LD

biogenesis in liver across weeks 2, 4 and 8, post infection. **(B)** Kinetic increase in the

gene expression of *Pparγ* transcript levels across different weeks post infection **(C)**

Immunoblot showing increased PPARγ in PHCs (MOI: 10) at 5 ,24 and 48 hours post-

infection. **(D)** Bar plot showing the increased band intensity of PPARγ in the infected

PHCs at the mentioned time points. **(E)** Representative confocal microscopy images

showing HepG2 infected with Mtb-mCherry-*H37Rv* and treated with GW9662 and

rosiglitazone **(F)** MFI/cell of in Mtb-mCherry-*H37Rv* DMSO treated, GW9662 and

rosiglitazone treated HepG2 cells. **(G)** Representative confocal images of uninfected,

infected, GW9662 and rosiglitazone treated infected HepG2 showing changes in the

number of LDs. **(H)** Plot depicting the quantification of LDs/cell in the mentioned

conditions. Representative data from n=4 biological replicates. Data were analysed by

using the two tailed unpaired Student's t test in D and by one way ANOVA in B, F and H.

*p < 0.05, **p < 0.005, ***p < 0.0005, ****p < 0.0001, ns=non-significant

Figure 7: Hepatocytes provide a drug-tolerant niche to Mtb: (A). Experimental

scheme of deducting the percentage drug tolerance in hepatocytes (generated with

iorender.com) **(B).** Percentage tolerance of Mtb-*H37Rv* against Rifampicin within RAW

264.7, PHCs and HepG2 at different time points post-infection. **(C).** Percentage tolerance

of Mtb-H37Rv against Isoniazid within RAW 264.7, PHCs, and HepG2 at different time

points post-infection. Representative data from 3 independent experiments. **(D).**

Percentage tolerance of Mtb-*H37Rv* against rifampicin within BMDMs, PHCs, and HepG2

as measured microscopically (**E).** Percentage tolerance of Mtb-H37Rv against isoniazid

within BMDMs, PHCs, and HepG2 measured microscopically. Representative data from

3 independent experiments. Each dot represents a single field having more than 4

infected cells. 20 such fields were analyzed **(F)**. Transcript levels of the various DMEs

involved in Rifampicin and Isoniazid metabolism in mice liver, 8 weeks post-infection (n=4

mice), the fold change has been calculated by considering the expression in the

uninfected mice to be 1. Data were analyzed by using 2-way ANOVA in (B) and (C), one-

way ANOVA in (D) and (E) and the two-tailed unpaired Student's t-test in (F) and

Representative data from n=4 biological replicates *p < 0.05, **p < 0.005, ***p < 0.0005,

****p < 0. 0001.ns=non-significant.

1061

1062

1063

A.

Uninfected
10X 60X

Infected
10X 60X

Acid-fast staining

B.

C.

A-O staining   Merged

D.

FISH   Merged

E.

DAPI
Ag85B
β-actin

20 μm

Figure 1

A. Aerosol infection of 200 CFU

Lung

Liver

Tissue level analysis

Isolation of Primary Hepatocytes

B. Lung, Liver

log10_CFU/ tissue

Day 1, Week 3, Week 4, Week 6, Week 8, Week 10

C. 20X, 40X

Uninfected

Infected

D. DAPI, Phalloidin, Ag85B, Merged

20X

40X

20 µm

E.

20 µm

F. ASGR1+ :100%

ASGR1+ 100.0

FSC-A

ASGR1

G.

50 µm

H. Log_10 CFU/ 10 million hepatocytes

Day 1, Week 3, Week 4, Week 6, Week 8, Week 10

I. PHCs-Ag85B, PHCs-Ag85B

10 µm

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7