Downscaling mutualistic networks from species to individuals reveals consistent interaction niches and roles within plant populations

Elena Quintero[1,2], Blanca Arroyo-Correa[1], Jorge Isla[1], Francisco Rodríguez-Sánchez[2§], Pedro Jordano[1,2§]

[1] Integrative Ecology Group, Estación Biológica de Doñana, EBD-CSIC, Av. Américo Vespucio 26, E-41092 Sevilla, Spain.

[2] Departamento de Biología Vegetal y Ecología, Facultad de Biología, Universidad de Sevilla, Av. Reina Mercedes, E-41012 Sevilla, Spain.

Corresponding authors: Pedro Jordano and Elena Quintero

Email: jordano@ebd.csic.es, equintero1@us.es

§ joint senior authors

Author Contributions:

All authors conceptualized the study and provided data, E.Q., B.A.C., and J.I. curated data, E.Q. and F.R.S. developed software, E.Q. performed formal analyses and visualization, F.R.S. and P.J. supervised the study, P.J. provided funding acquisition and project management, E.Q. led writing, all authors contributed to writing, reviewing and editing.

Competing Interest Statement: The authors declare no competing interest.

Keywords: individual-based networks, interaction niche, frugivory, mutualism.

Abstract

Species-level networks emerge as the combination of interactions spanning multiple individuals, and their study has received considerable attention over the past 30 years. However, less is known about the structure of interaction configurations within species, even though individuals are the actual interacting units in nature.

We compiled 46 empirical, individual-based, interaction networks on plant-animal seed dispersal mutualisms, comprising 1037 plant individuals across 29 species from various regions. We compared the structure of individual-based networks to that of species-based networks and, by extending the niche concept to interaction assemblages, we explored individual plant specialization. Using a Bayesian framework to account for uncertainty derived from sampling, we examined how plant individuals "explore" the interaction niche of their populations.

Both individual-based and species-based networks exhibited high variability in network properties, lacking remarkable structural and topological differences between them. Within populations, frugivores' interaction allocation among plant individuals was highly heterogeneous, with one to three frugivore species dominating interactions. Regardless of species or bioregion, plant individuals displayed a variety of interaction profiles across populations, with a consistently small percentage of individuals playing a central role and exhibiting high diversity in their interaction assemblage. Plant populations showed variable mid to low levels of niche specialization; and individuals' interaction niche "breadth" accounted for 70% of the population interaction diversity, on average.

Our results highlight how downscaling from species to individual-based networks helps understanding the structuring of interactions within ecological communities and provide an empirical basis for the extension of niche theory to complex mutualistic networks.

Significance Statement

Ecological interactions in nature occur between individual partners rather than species, and their outcomes determine fitness variation. By examining among-individual variation in interaction niches, we can bridge evolutionary and ecological perspectives to understand interaction biodiversity. This study investigates individual plant variation in frugivore assemblages worldwide, exploring how plant individuals "build" their interaction profiles with animal frugivores. The structure of networks composed of individuals was surprisingly similar to networks composed of species. Within populations, only a few plants played a key role in attracting a high diversity of frugivores, making them central to the overall network structure. Individuals actually interacted with a substantial diversity of partners, with individual niche "breadth" accounting for up to 70% of total interaction diversity, on average.

Introduction

Species are a fundamental unit of study in most ecological research, resulting in numerous theoretical and methodological approaches to assess how their interactions support ecosystem functions. Network ecology based on graph theory has emerged as a useful framework to study these multi-species interactions simultaneously and assess the complexity of natural ecosystems (Solé & Valverde 2004, Fortuna & Bascompte 2008, Fontaine et al. 2011). Starting with food webs (Cohen 1978), network theory expanded its versatility to other ecological interaction modes such as mutualisms (Jordano 1987, Memmott 1999). Since then, abundant literature has revealed emergent and global properties of ecological networks, highlighting surprisingly similar architecture in the way they are assembled (McCann et al. 1998, Mora et al. 2018). Among ecological networks, mutualistic networks represent mutually beneficial interactions, and their structure and topology have been extensively explored (Bascompte & Jordano 2007). Plant–animal mutualistic networks are highly heterogeneous (i.e., most species have few interactions while a minority of species are much more connected) and nested (i.e., specialists interact with subsets of the species with which generalists interact), leading to asymmetric dependences among species (Jordano et al. 2003, Bascompte & Jordano 2007). Yet, it is not clear to what extent these properties emerge from networks at lower levels of organization, such as those composed of individual interactions.

Although interaction patterns are usually summarized at the species level, ecological interactions actually occur as encounters between individuals rather than species (Clark et al. 2011). For instance, while we say blackbirds (*Turdus merula*) consume fruits and disperse raspberry (*Rubus idaeus*) seeds, it's actually individual plants and birds interacting within a local population. By missing this individual-level resolution we miss two important opportunities: 1) the ability to effectively link individual trait variation with interaction outcomes (fitness effects) and thus connect ecological and evolutionary perspectives; and 2) to bridge the gap between niche theory and

complex interaction networks, i.e, to assess how individual-based interactions scale up into complex

interaction networks.

Classic studies of animal-mediated seed dispersal interactions have been plant-focused (e.g., Snow

& Snow 1988), and provide a useful framework to zoom-in into the interactions established

between a particular plant species and a set of animal frugivores. By considering individual-based

networks, in which one set of nodes is composed of plant individuals, and the other set is composed

of animal species, we can examine individual variation in "interactions build-up", as well as its

subsequent implications, in e.g., fitness (Gómez & Perfectti 2012, Rodríguez-Rodríguez et al. 2017,

Arroyo-Correa et al. 2021). This is helpful not just for building a proper bridge between interaction

ecology and demographic consequences (e.g., Quintero et al. 2023), but also for bridging network

ecology with evolutionary consequences (Guimarães et al. 2011, Segar et al. 2020).

Network structure may not be consistent across hierarchical scales of organization (Tur et al. 2014,

Wang et al. 2021). The similarity in the set of partners available to individuals of the same species

will be higher than that to different species. That is, the physical and phenological traits of

conspecific individuals tend to be more similar than those among species (Siefert et al. 2015),

discouraging major forbidden interactions (but see Albert et al. 2010, González-Varo & Traveset

2016) and likely increasing overall network connectance. Thus, we might expect individual-based

networks to exhibit architectural and structural properties different to those found in species-based

networks; yet, this remains an underexplored question.

Downscaling the study of interactions to individuals allows us to observe how the variation among

individuals in their partner use is distributed in the population (Fig. 1A-B). Since its origins, the niche

concept has provided an ideal framework for studying individual variation in resource use (Grinnell

1917, Van Valen 1965, Bolnick et al. 2003). Even so, most previous work has focused on antagonistic

interactions such as predator–prey trophic niches (Bolnick et al. 2003, Araujo et al. 2011, Costa-

Pereira et al. 2018, Costa-Pereira et al. 2019). It was only very recently that niche theory was applied

for understanding individual variation in mutualistic interactions (Tur et al. 2014, Albrecht et al.

2018, Phillips et al. 2020, Koffel et al. 2021, Arroyo-Correa et al. 2023, Gómez et al. 2023). However,

the study on how a plant individual's interaction niche is distributed within a population remains

largely unexplored, even less has been compared across different ecological systems. For this study,

we rely on the concept of 'interaction niche' as the space defined by the set of species with which a

population, or a plant individual, can interact (Fig. 1C) (Ponisio et al. 2019).

Interaction probabilities between plant individuals and animal species (i.e., probability of

interspecific encounter, PIE; Hurlbert 1971, Chase & Knight 2013) are influenced by a myriad of

factors such as population abundances, accessibility of resources, individual preferences or

physiological needs (e.g., optimal foraging theory) as well as required matching in traits and

phenology (Guimarães 2020). Intraspecific trait variability, neighborhood attributes and spatio-

temporal context drive animal preferences for certain plant individuals, which will govern the

establishment of interactions between plant individuals and their mutualistic partner species

(Sallabanks 1993, Snell et al. 2019, Isla et al. 2023). In mutualistic systems such as pollination or seed

dispersal, variation in the patterns of interaction or exploitation of niches (available partners) can

play a determining role, as mutualists directly affect the reproductive outcome of individuals,

influencing fitness variation and trait selection, which act as raw material for coevolution

(Thompson 1999), as well as population dynamics and community assembly (Simmons et al. 2020,

Arroyo-Correa et al. 2023).

Quantifying individual variation in interaction niche, and particularly niche partitioning, can shed

light on the coexistence and stability of mutualistic communities. For instance, individuals in a

population can behave as specialists or generalists when exploiting their interaction niche, and this

may influence how these individuals are affected by interspecific competition and how partner

diversity is promoted, determining, e.g., degree distributions in interaction networks (Bascompte &

Jordano 2014). The extent to which individuals behave as specialists or generalists in a population

can be elucidated by partitioning niche variation into its between- (BIC) and within-individual (WIC)

components. Thus, this approach can prove useful to predict niche-shifts or interaction niche

expansion (Roughgarden, 1972, Bolnick et al. 2007). The levels of individual specialization in

individual-based networks can be estimated as the proportion of the total niche width in the

population (TNW; total partner diversity) due to within-individual variation (WIC; average partner

diversity of individuals). Thus, the distribution of frugivore–partner species richness and interaction

allocation among plant individuals can be highly variable in local populations (e.g., Jordano &

Schupp 2000, Guerra et al. 2017, Miguel et al. 2018, Jácome-Flores et al. 2020, Quintero et al. 2023).

By studying plant individual specialization and how frugivores distribute interactions among plants,

we aim to understand variation in mutualistic interactions within plant populations (Fig. 1).

Examining how interaction niches are partitioned globally can expand the concept of niche variation

to mutualistic interactions and pave the way for future hypotheses.

A variety of node-level metrics for complex networks can provide insight into an individual's

strategy within its population (Dormann 2011, Poisot 2012). Several studies have used node-level

metrics to characterize individuals' positioning in the network, informing us about their role and

significance in their population (e.g., Gómez & Perfectti 2012, Guerra et al. 2017, Rodríguez-

Rodríguez et al. 2017, Crestani et al. 2019, Vissoto et al. 2022, Isla et al. 2023, Arroyo-Correa et al.

2023). However, most of these studies have used a single or several metrics separately to

understand the interaction profile of individuals and for single populations. By using a combination

of node-level metrics, we aim to characterize interaction profiles of plant individuals with frugivore

species and the distribution and frequencies of roles among and within populations in different

geographic regions. Given contrasted differences in e.g., frugivore diversity and life histories across

biogeographic realms, we could expect plant individuals from certain regions to exhibit similar

interaction profiles, markedly different from those of individuals from other species and/or regions.

Conversely, if life history or context-dependent effects were not determinant in structuring

interactions between plant individuals and their frugivore partners, we could expect consistent

individual interaction profiles across populations, irrespective of geographic location or biome type.

Obtaining an insightful picture of a mutualistic network structure from field data is a challenging

task. When sampling is limited, the inferred network structure can be noisy, even biased, and thus

subject to sampling fluctuations. This issue becomes particularly relevant when comparing

networks from different studies (Brimacombe et al. 2023). Here, we build upon Young et al. (2021)

Bayesian framework for reconstructing mutualistic networks to infer each pairwise interaction in

individual-based networks, accounting for sampling completeness and the inherent stochasticity of

field observation data. We then propagate the uncertainty of all pairwise interactions down through

niche specialization and network interaction profiles.

The overarching goal of this study is to investigate the role played by individuals in the assembly,

structure and functioning of complex ecological interaction networks. To do so, we combine

network and niche theory to characterize the interaction profile of plant individuals in mutualistic

seed dispersal systems across different bioregions aiming to illustrate the wide diversity of plant

populations considered. We outline three main objectives: 1) examine whether networks composed

of individuals exhibit different architectural and structural properties than those found in species-

based networks, 2) understand how variation in frugivory interactions takes place at the plant

population level by quantifying individual niche-partitioning and frugivore interaction allocation,

and 3) characterize interaction profiles of plant individuals with frugivore species and assess the

distribution and frequencies of roles among populations.

Results

<u>Structure of individual versus species-based networks</u>

We assembled a total of 46 individual-based plant-frugivore networks and compared them with 59 species-based networks using six network metrics (connectance, nestedness, modularity, assortativity, centralization and interaction evenness). Networks showed a remarkable overlap in all metrics at both resolution scales (Fig. 2A, Fig. S4, Table S3). Species-based and individual-based networks presented wide variation in their structure. Overall, when controlled by network relative size and interaction abundances, networks were less connected, nested and even, and more modular, assortative and central than their corresponding null models, although some of these differences were very subtle and consistent for networks at the two scales (Fig. 2B, Fig. S4). Comparing across scales, individual-based networks were slightly less connected and less modular than species-based networks relative to random expectations. However, these differences remained subtle and highly overlapping (e.g., 0.05 average difference in connectance and 0.04 in modularity; Table S3).

<u>Plant individuals' specialization in interaction niche</u>

The 46 populations included a total of 1037 plant individuals: 373 individuals (from 9 species) from the Mediterranean (Iberian Peninsula), 389 (17 species) from Tropical regions in Asia and America, and 275 (3 species) from Southern Temperate regions (Southern Brazil and Argentina). Our aim in the compilation of individual-based networks was to showcase the diversity of environments and their associated variations; this analysis revealed major gaps in data coverage on the African continent and in the Australasian region (see Table S1), so a proper comparison among biogeographic areas awaits more complete data. Most plant populations studied presented low to medium levels of individual specialization (mean WIC/TNW = 0.68; 90% CI = 0.32 - 0.95). Individual-based networks from the Mediterranean region showed a higher proportion of generalized plant

individuals (Fig. 3; mean WIC/TNW = 0.82), whereas southern temperate and tropical populations

presented higher levels of individual specialization (mean WIC/TNW = 0.56 and 0.65 respectively, p-

value < 0.01). Plant populations interacting with higher numbers of frugivore species had a wider

interaction niche (TNW, i.e., Shannon diversity index, r = 0.53, p < 0.01), but not necessarily higher

levels of individual specialization (WIC/TNW) (Fig. S5). The degree of individual specialization did

not increase as TNW increased, because WIC increased in the same proportion as TNW (slope of

log-log model = 1.000, SE = 0.005). That is, differences in the degree of individual specialization

between bioregions were achieved via changes in both the BIC and the WIC components.

<u>Frugivore interactions within plant populations</u>

Across all plant populations and regions, we found that just a reduced subset of frugivore species

(generally between one and three) usually accumulated most of the interactions, while the rest of

frugivore species contributed a minor proportion. On average, more than half of these interactions

were contributed by less than 20% of frugivores (SD = 8.2%), regardless of the total number of

frugivore species in the plant population (Fig. S6). The frugivores that contributed most interactions

also tended to interact with a higher number of plant individuals (Spearman's rho = 0.81, Fig. 4).

Remarkably, frugivore species with smaller contributions interacted with a variable proportion of

plant individuals, such proportion being higher in Mediterranean networks and lower in Southern

Temperate networks (Fig. 4). Frugivores' body mass was not correlated with interaction

contribution (rho = -0.08) nor with the proportion of plant individuals they interacted with (rho = -

0.13).

<u>Plant individuals' interaction profiles</u>

The multidimensional principal component analysis space occupied by all plant individuals and

defined by node-level metrics (normalized degree, strength, specificity, overlap and weighted

closeness) did not produce distinct clusters by bioregion or population. Instead, individuals from

different populations spread across the multidimensional space, suggesting ample within-

population heterogeneity in plant individuals' interaction profiles (Fig. 5, Fig. S8). The first principal

component (PC1), explaining more than half of the variation (51%), was mainly related to

interaction degree and specificity, thus capturing individual variation in frugivore richness and

composition of the plant individual's assemblages. The second component (PC2) explained 24% of

the variation and was correlated with niche overlap and interaction strength; these metrics are

related to plant individuals' interaction patterns in relation to their conspecifics and affected by

interaction frequency (link weight). Plant individuals with more unique frugivore assemblages were

positioned in the upper area of the PCA space, while many plants with highly-overlapping frugivore

assemblages within their populations were positioned towards the bottom area. The third

component (PC3; 10% variation explained; Table S4) was strongly related to weighted closeness, a

measure of how strongly and well connected (i.e., central) individuals are within the network.

Few plants were highly central in the interaction network (high weighted closeness) and important

for frugivore dependence (high species strength) (i.e., points in the upper-left area of the

multivariate space). Most plant individuals showed uneven dependencies on frugivore species

and/or medium-high frugivore overlap with other plants in the population. Yet plants with strong

dependencies on one or few frugivore species tended to show lower overlap with other individuals

in the frugivore assemblage, suggesting a trade-off between partner specialization and partner

sharing (lower-right Fig. 5). Overall, plant individuals from the Mediterranean tended to have more

similar frugivore assemblages (higher niche overlap), while plant individuals from Southern

Temperate regions presented less overlapping and more specialized frugivore assemblages (Fig.

S9).

A cluster analysis further revealed consistent interaction profiles within plant populations,

regardless of bioregion or species. All networks presented a variable but substantial proportion

(mean = 42%; 90% CI = 8 - 77%) of plant individuals with "neutral" interaction profiles, which did not

excel in any node-level metric (Fig. 5, Fig. S9). The clusters "specialized" (mean = 28%; 90% CI = 0 -

67%) and "overlapping" (mean = 22%; 90% CI = 0 - 86%) included a variable but substantial

proportion of plants, most likely reflecting either distinct frugivore assemblages (e.g., in the tropical

biomes) or more overlapping in composition (mostly non-tropical), respectively. Notably, in most

populations (36 out of 46), we detected a low proportion (mean = 8%, 90% CI = 0 - 24%) of

"keystone" individuals showing high strength and weighted closeness. We termed these plant

individuals as keystone since they accounted for a high proportion of interactions within their

respective populations, playing a central role in plant-frugivore interaction structuring.

Discussion

Our study highlights how downscaling from species to individuals uncovers consistent network

structures across biological levels, mutualistic partner allocation among plants globally, and interaction

profile similarities within populations, regardless of species or ecological context. This reveals new

aspects of ecological interactions and network assembly at the individual level.

<u>Effects of downscaling resolution on network structure</u>

The structure of plant–animal mutualistic networks revealed fundamental heterogeneity across

networks and resolution scales. We did not find major deviations in the assembly patterns of

interactions as we zoomed in from species to an individual-based scale. Previous research exploring

the consequences of downscaling on network architecture found significant shifts in the structure of

pollination networks (Tur el al. 2014, Wang et al. 2021): individual-based networks were less

connected and individuals were more specialized than species. These studies examined one

community using a single methodology. Our approach, however, involved diverse communities and

populations from different regions with varying methodologies, capturing broader interaction patterns.

Surprisingly, we found no significant differences in network connectance across scales, contrary to our

initial expectations. The slightly larger connectance of individual networks after Bayesian modeling

(average connectance = 0.4 ± 0.2 vs 0.3 ± 0.1 in raw networks) suggests that undersampled

interaction matrices could be missing interactions. On the other hand, the slightly lower modularity

in individual-based networks could stem from a decrease in the number of forbidden links as

frugivores can potentially interact with virtually all the plant individuals within populations (see

González-Varo & Traveset 2016). In contrast, species-based networks involve much more

heterogeneous sets of plant species. The addition of new species or individuals with new traits

provides new link possibilities in a network, yet in the case of species, potential interactions must

undergo stronger trait and phenological matching filters than in individual-based networks

(Sazatornil et al. 2016). Simply put, a given frugivore species may interact with a broader range of

partners within a plant population than when interacting with the full range of taxonomically-

diverse available plant species in a community. The former set imposes less constraints to

interactions by including much more homogenous conspecific partners. In this way, downscaling

from species to individuals fundamentally alters the probabilistic distribution of interactions among

partners (Poisot et al. 2016).

Aside from minor differences in certain network metrics, the overall topology and structure of

frugivory networks at different resolution scales were not sufficient to make clear distinctions. This

convergence in network structure may be driven by factors like the probability distribution of

interspecific encounters (PIEs), influencing network configurations consistently across scales. We argue

that numerical effects are likely at the base of these emergent properties, governing interactions

distribution across nodes and asymmetric interactions (Jordano 1987, Vázquez et al. 2007,

Schleuning et al. 2011, Guimarães 2020). These numerical effects can be caused by varied organism

abundances in the case of species, or traits and genotypes in the case of individuals, that modulate

the attractiveness of plants to frugivores, such as crop size, plant height, or phenology (Snell et al.

2019).

<u>Individual specialization in the interaction niche</u>

Individuals' interaction niches were narrower than those of their populations, supporting that

individual specialization is substantial and common in nature (Bolnick et al. 2003), even in

mutualisms. Plant individuals' specialization levels were similar to levels reported in other animal

taxa (Araujo et al. 2011). Interestingly, the degree of individual specialization varied across

biogeographic regions, yet most plant species showed WIC/TNW ratios >50%, which indicates

moderate generalization among plant individuals. Broader and more overlapping frugivore

assemblages in Mediterranean regions versus higher specialization and variability in southern

temperate and tropical networks could not be attributed to differences in taxonomic diversity as all

regions presented similar total niche width (Fig. S5). Instead, plant individuals in southern

temperate regions presented smaller relative niches. Tropical populations consisted of frugivore

assemblages of variable diversity and highly variable individual specialization, yet limited area

coverage of the available individual-based networks hindered comparisons. No significant niche

breadth differences were found across bioregions, aligning with studies of terrestrial food webs,

challenging the latitude-niche breadth hypothesis that predicts narrower niches in tropical regions

(Cirtwill et al. 2015, although see Araujo & Costa-Pereira 2013). The large variation in individual

specialization within bioregions may be pointing to the role of fine-scale factors such as the local

habitat, neighborhood effects, or the influences of individual phenological variation. Further

research is needed to evaluate the ecological correlates of plant individual interaction niche

utilization and its consequences.

Different levels of individual specialization can have implications for population stability (Arroyo-

Correa et al. 2024) and niche expansion (Van Valen 1965). According to the niche variation

hypothesis, populations experiencing niche expansion achieve it through increasing their inter-

individual variation (Bolnick et al. 2007). By diversifying its resources, plant individuals would be

able to exploit novel and underutilized frugivores escaping competition from conspecifics. Niche

shifts and expansion have become exceptionally important for adaptation to changing climate

conditions (Hallfors et al 2023) as well as changes in frugivore assemblages and fluctuating

abundances (Campos-Celada et al. 2022). Therefore, the variation we found among populations in

frugivore assemblage specialization will likely have an impact on the adaptation of plant–frugivore

mutualistic interaction niche in current and future scenarios of global change.

In all plant populations just a few frugivore species, even within diversified assemblages,

consistently perform most of the mutualistic interactions (Fig. S6; Rother et al. 2016, Guerra et al.

2017, Isla et al. 2023, Rehling et al. 2023, Thiel et al. 2023). Although frugivore body mass did not

prove to be a good indicator of their contribution to interactions (although see Valenzuela-Ospina &

Kattan 2021), it may play a role in seed dispersal effectiveness due to its positive correlation with

the number of fruits consumed per visit or the frequency of long-distance seed dispersal events

(Snow & Snow 1988, Jordano et al. 2007, Godínez-Álvarez et al. 2020). These highly uneven

interaction patterns will result in asymmetric dependencies between plant individuals and frugivore

species, where the main frugivore shows low specificity for specific plants, while most plant

individuals rely mostly on the main frugivores' service (Quintero et al. 2023). Asymmetric

dependency between partners also emerges at species–species interaction level (Jordano 1987,

Vázquez & Aizen 2014, Bascompte et al. 2006); further downscaling into individual-individual

interactions would help elucidate if asymmetry remains consistent across scales. Finally, our

analysis reveals a consistent trend for frugivory and seed dispersal service in a given plant

population (estimated from the proportion of plant individuals with which a frugivore species

interacts) to increase with the overall contribution to the total number of interactions by each

frugivore species (i.e., link weights in the individual-based networks, Fig. 4). Thus, central frugivores

interact with a wide range of plant individuals, most likely an emergent result of the interaction

asymmetry discussed above.

<u>Consistency of individual plants' interaction profiles across regions and populations</u>

Plant individuals' interaction profiles were not explained by bioregion or species, pointing to

fundamental architectural patterns in the assemblage of mutualistic interactions that are not

strongly constrained by phylogeny or geographic location but rather by the interplay between traits

and numerical effects (Jordano 1987, Carnicer et al. 2009, Albrecht et al. 2018, Guimarães 2020).

Remarkably, we found a consistent distribution of plant interaction profiles within populations, with

most individuals acting in an average manner, a variable fraction standing out for their

specialization or redundancy and only very few individuals having a central role, high diversity of

interactions, and strong frugivores' dependence on them ("keystone" plant individuals). Similar

results are reported in food webs, where a core group of species shares ecological roles, while

peripheral species have unique interaction profiles (Mora et al. 2018). It is likely that within frugivory

networks these key individuals present unique phenotypic traits, such as abundant fruit crops or

advantageous locations that make them reliable to many frugivores (Snell et al. 2019, Isla et al.

2023).

Although some of the plant species considered in this study were generalists within their

community, individuals in their populations showed variable interaction niche breadths (Guerra et

al. 2017), with populations consisting of both generalist and specialist individuals (Arroyo-Correa et

al. 2023, Bolnick et al. 2007). This mix creates species that appear broadly interactive but actually

include individuals with varied interaction patterns, from wide generalization to specific partner

preferences. This highlights the complexity of species' ecological roles within communities.

(Guimarães 2020).

<u>Concluding remarks</u>

We found consistent patterns of interaction assembly across biological scales using a set of

biologically informative network metrics. On top of the absence of a clear hierarchy differentiation

in network structure between individuals and species, we found that almost every individual-based

network included a similar representation of individual interaction profiles, evidencing a common

backbone in the way interactions are organized (Mora et al. 2018). Conducting future analyses on

interaction types or motifs of individual-based networks may provide us with new insights, as these

approaches have proven effective in distinguishing networks between and within ecological

systems (Mora et al. 2018, Michalska-Smith et al. 2022, Pichon et al. 2023).

Intraspecific variation appears as a central ingredient in the configuration of complex networks of

mutualistic interactions, driven by the widespread interaction profiles of frugivore species with

plant individuals. High levels of intraspecific variation have been shown to confer greater stability to

mutualistic systems (Arroyo-Correa et al. 2023). By zooming in on ecological interactions this study

provides valuable insights into how mutualistic interactions are similarly structured at the individual

level and reveal underlying, consistent, patterns of role assignment within populations and across

bioregions.

Methods

<u>Dataset acquisition</u>

We compiled frugivory ecological networks , both at the species and the plant individual level.

Species-based networks were gathered from 40 published studies at the community scale (see

Table S1). For individual-based networks, which are scarcer, we compiled phyto-centric studies

(plant-based), with quantitative information on frugivore visitation on plant individuals within

populations. We combined published studies with unpublished datasets, gathering data for 21

different study systems, including datasets from our own field studies with different Mediterranean

species (n = 9). Some of the studies selected presented more than one network from different

communities (in species-based studies) or populations (in individual-based studies) (Table S1).

These datasets document interactions between plant species and animal frugivore species (in

species-based studies) or interactions between plant individuals of a single species that coexist in a

local population and animal frugivore species (in individual-based studies). None of the datasets

collected are nested, that is, individual-based networks are not sampled within the same study as a

species-based network, which prevents a direct structure dependence between the two scales of

the datasets. Data was entered as adjacency matrices, where rows represent plant species (or

individuals) and columns represent animal species, with matrix elements a_{ij} indicating interaction

frequency (visitation frequency to plants). In order to ensure networks were sufficiently sampled to

robustly characterize their structure and interaction profiles, we only kept those that were

reasonably complete. We checked for sampling coverage of individual-based networks using iNext

R package (Hsieh et al. 2016) (Table S2). To do so, we converted the adjacency matrix data to an

incidence frequency-data and considered plant individuals as sampling units and the number of

frugivore species detected at each plant (species richness). We discarded networks in which the

number of interacting nodes (plants and frugivore species) was less than 15 or plants were less than

six (n = 13 networks). Our final dataset consisted of 105 networks with an average size of 384

potential links or cells (range = 65 - 2904) and 90 unique interactions (range = 22 - 419). Forty-six

were individual-based networks and 59 were species-based networks (Table S1). When possible we

referred the interaction value to the coarsest level, that is, frugivore visitation events, otherwise

number of fruits consumed.

<u>Network-level metrics</u>

For both the individual and species-based networks, we calculated several network-level metrics,

using R packages bipartite (Dormann et al. 2008) and igraph (Csárdi & Nepusz 2006). All network

metrics were calculated using standardized matrices to minimize the impact of differences in

sampling effort and study-site characteristics. Specifically, for all those individual-based networks

where sampling effort across plant individuals was heterogeneous within the population (21 out of

46 networks), we divided plant visits by the amount of time observed and/or the area sampled in

each plant, so that interaction counts were comparable. Subsequently, all networks (individual and

species-based) were scaled by dividing the weight of each pairwise interaction by the total number

of interactions in the matrix (grand total standardization; Quintero et al. 2022). In this way, the

interaction values (matrix cells) represent the relative frequency of a plant individual (in individual-

based networks) or a plant species (in species-based networks) interacting with a given frugivore

species, and the sum of all relative frequencies equals one.

We selected a representative set of metrics that had suitable biological interpretation and were not

highly correlated (Variance Inflation Factor < 3) and/or not strongly affected by the number of

species/individuals sampled or overall network size (Fig. S1, S2, S3).

Selected network-level metrics were:

1.   Connectance *(topology)*. This metric gives the proportion of realized over potential links in the

network. Calculated as the sum of realized links (unweighted) divided by the number of

cells in the matrix. Values range from 0 (no links) to 1 (fully connected networks where

all nodes interact among them) (Dunne et al. 2002).

2.   Weighted nestedness wNODF *(structure)*. It informs on the way interactions are organized. A

highly nested structure is one in which nodes with fewer connections tend to interact with a

subset of highly connected nodes that in turn interact with the highly connected ones

(Bascompte et al. 2003). Values of 0 indicate non-nestedness, those of 100 perfect nesting

(Almeida-Neto & Ulrich 2011).

3.   Assortativity *(topology)*. This metric indicates the level of homophily among nodes in the

graph. It ranges from -1 to 1, when high it means that nodes tend to connect with nodes of

similar degree; when low, nodes of low-degree connect with nodes of high-degree

(disassortative) (Newman 2002, Barabasi 2016).

4.   Modularity *(structure - clustering)*. It reflects the tendency of a network to be organized in

distinct clusters or modules (Bascompte & Jordano 2014). This metric ranges from 0 (no

clusters) to 1, where nodes within a module interact more among them than with nodes from

other modules (highly compartmentalized network; Newman 2006).

5.   Eigenvector centralization *(centrality)*. This metric quantifies how centralized or decentralized

the distribution of eigenvector centrality scores is across all nodes in a network (Freeman et al.

1979). The eigenvector centrality of a given node in a network is a measure of the influence of

that node, taking into account both the node's direct connections and the connections of its

neighbors. Nodes with high eigenvector centrality are connected to other nodes that are also

central, giving them a higher score (de Oliveira Lima et al. 2020). The network-level

eigenvector centralization provides a measure of the extent to which a few nodes dominate the

network in terms of influence. In a network with low centralization, the centrality scores are

relatively evenly distributed among the nodes, suggesting a more decentralized structure

where many nodes contribute to the overall connectivity of the network, and therefore to the

interaction services. On the other hand, a network with high centralization indicates that only a

small number of nodes have a higher centrality, suggesting a more centralized structure where

a few nodes play a crucial role in the network's overall connectivity. We normalized this

measure to ensure that the centralization value is relative to the maximum centralization for a

network of a given size.

6.   Interaction evenness (*interaction diversity*). Also known as Pielou's evenness, it quantifies how

balanced the distribution of interactions is across nodes (Blüthgen et al. 2008). This metric

considers all links as species, and their weight as a measure of their abundance. It quantifies

link diversity using Shannon index and divides it by the theoretical maximum richness of all

possible links (i.e., logarithm of Shannon). It ranges from 0 to 1, where low values indicate high

unevenness and high values approach perfect evenness in interaction distribution.

<u>Comparison across network scales</u>

We used the aforementioned metrics to compare the structure and assemblage of species-based

versus individual-based networks. Additionally, to account for network size and inherent

characteristics of each study, we assessed metric deviations against 1000 randomizations using null

models. We use a Patefield algorithm (Patefield 1981, Dormann et al. 2009) that maintains network

marginal totals. For each network and metric, we calculated the difference between the observed

values and the randomizations. This difference indicates both the magnitude and direction in which

observed values deviate from what could be expected at random.

<u>Bayesian modeling of individual-based network structure</u>

Reconstructing mutualistic network structure from field data is a challenging task. Interaction data

are hard to collect and typically require large sampling efforts, particularly to characterize

infrequent interactions. Inferred network structure is highly sensitive to sampling design, effort, and

completeness (Jordano 2016). Comparing networks from different studies without accounting for

these sampling effects may lead to mistaken inferences (Brimacombe et al. 2023). Here, we build

upon Young et al. (2021) Bayesian framework for reconstructing mutualistic networks to infer the

posterior probability of each pairwise interaction in individual-based networks, accounting for

sampling completeness and the inherent stochasticity of field observation data.

Following Young et al. (2021), pairwise interaction counts between plants and animals can be

modeled as following a Poisson distribution whose mean (μ_{ij}) is determined by the sampling effort

spent on each plant (C_i), the relative interaction abundance of each plant and animal in the

population (σ_i and τ_j), the (inferred) existence of an interaction link among both partners (B_{ij}), and

the "preference" parameter r which represents the difference in the average number of visits or

interactions when there is a connection between mutualistic partners (i.e., when a frugivory

preferentially interacts with a given plant, $B_{ij} = 1$) compared to when there is no connection ($B_{ij} = 0$).

Thus,

$$\mu_{ij} = C_i \sigma_i \tau_j (1 + r B_{ij})$$

The preference parameter r accommodates the expectation that the number of interactions will be

higher when a frugivore prefers a given plant. When there is no preference the number of

interactions can still be higher than zero, but lower in principle. Here we adapted the model

proposed by Young et al. (2021) to allow for varying preferences (i.e., interaction counts) among

different frugivore species, so that some frugivores may bring many more visits than other

mutualists. Rather than having a single preference parameter r for all species, each frugivore has

their own preference r_j which is drawn from an exponential distribution with rate parameter beta =

0.01. In order to facilitate modeling and incorporate these modifications, we developed an R

package called 'BayesianWebs' (Rodríguez-Sánchez 2024a) which relies on Stan (Stan Development

Team 2024) and cmdstanr (Gabry et al. 2024) for parameter estimation, and contains functions to

facilitate modeling of bipartite mutualistic networks including data preparation, model fitting,

checking and visualization. Using this package, we fitted a Bayesian model to each individual-based

network. As model output, we obtained 1000 posterior samples of the expected count for each

pairwise interaction. Posterior predictive checks showed the expected increase of uncertainty in

networks with limited sampling, and overall confirmed the good match between the observed data

and predicted counts.

To ensure comparability across networks, posterior interaction counts were rounded to integer

values and then standardized following the same procedure explained above. Networks with

heterogeneous sampling effort across plant individuals were adjusted by dividing the posterior

counts by the individual relative observation time or sampled area. Subsequently, all networks were

standardized dividing each interaction value by the total number of interactions (grand total

standardization, Quintero et al. 2022), so that matrix cell values reflected the relative frequency of

each interaction within the population.

The resulting distributions of pairwise interaction counts generated by the Bayesian model for each

network (1000 matrices for each network or population) were then propagated all the way down to

niche specialization and individual node-level metrics (see below).

<u>Niche specialization</u>

We estimated populations' niche specialization using the Shannon approximation of the WIC/TNW

index for discrete data (Roughgarden 1979, Bolnick et al. 2002) implemented in the 'network.tools'

R package (Rodríguez-Sánchez 2024b). In this case, we define as a niche-resource the available

coterie of visiting frugivore species in a given population. This index computes the relative degree of

individual specialization as the proportion of total niche width (TNW) explained by within-individual

variation (WIC). Total niche width (TNW) is calculated as the total diversity of frugivore species

visiting the plant population, using Shannon index,

533
$$TNW = -\sum_j q_j ln(q_j),$$

534   where j represents an animal species, and q_j is the proportion of interactions contributed by

535   frugivore species j to the total number of interactions in the plant population.

536   The within-individual variation (WIC) is calculated as the average Shannon diversity of frugivores for

537   each plant individual, weighted by the relative proportion of all frugivore interactions in the

538   population that are used by each individual,

$$WIC = \sum_i p_i \left(-\sum_j p_{ij} ln(p_{ij}) \right),$$

where *i* presents a plant individual, p_i is the proportion of interactions contributed by plant *i* to the

total number of interactions in the population, and p_{ij} the proportion of interactions that animal

species *j* contributed to plant individual *i*.

Finally, WIC is divided by TNW. Values closer to 1 indicate a population composed of generalist

individuals that are using most of the niche available to the population. On the contrary, values

closer to 0 indicate a population of specialist individuals using small subsets of the population niche,

with large differences in resource-use among them. WIC and TNW estimates were calculated for

each of the posterior counts (n = 1000) generated from Bayesian models for each network,

rendering a credibility interval of individual specialization for each plant population. To test

differences in individual specialization (i.e., WIC/TNW) between different bioregions (n = 3) we fitted

a mixed-effects linear model with a Normal distribution where the network and study were present

as random factors (Bates et al. 2015).

<u>Node-level metrics</u>

To characterize plant individuals' interaction profiles in their populations, we computed a set of

node-level indices for each plant individual using R package bipartite (Dormann et al. 2008).

Additionally, we calculated average niche overlap using Bray-Curtis dissimilarity index (vegan R

package; Oksanen et al. 2022). We computed these metrics using the full posterior distribution of

each individual-based network. That is, for each of the 1000 posterior matrices generated for each

network we calculated node-level metrics, therefore we obtained 1000 values for each plant

individual and metric. We selected a representative set of metrics that had suitable biological

interpretation for assessing the individuals' interaction profiles and were not highly correlated nor

affected by the number of individuals sampled (Fig. S7).

Selected node-level metrics were:

1. Normalized degree *(interaction diversity)*. It represents the richness of partners for a given node and is scaled relative to the rest of nodes in the network. This metric ranges from 0 to 1, where a plant individual would score 1 if it interacts with all the frugivore species available (Dormann 2011).

2. Species specificity index *(interaction diversity)*. This metric informs about the variation in the distribution of interactions with frugivore species partners. It is estimated as a coefficient of variation of interactions for each plant individual, normalized to range between 0 and 1 (Julliard et al. 2006, Poisot et al. 2012). High values indicate higher variation in dependence on frugivore species. Plants with high dependence on few or a single frugivore species yield values close to 1, and plants that distribute their interactions equally with many frugivore species show low values.

3. Species strength *(interaction intensity)*. It quantifies the dependence of the community on a given node (Dormann 2011), in this case, the frugivores represent the community and the plant individuals the nodes. It is calculated as the sum of the dependencies of each frugivore species (i.e., the fraction of all visits to a given plant individual coming from a particular frugivore species) (Bascompte et al. 2006).

4. Weighted closeness *(node position)*. This metric provides an index of the magnitude to which a given node has short connection paths to all other nodes in the network (Opsahl et al. 2010). It is influenced by the intensity and number of links and indicates to what extent a node is in the "center" of the connections of the graph. This metric is calculated on the unipartite projection of the individual-based network for the plant individuals, with links between plant individuals representing the number of frugivore interactions shared. The weighted closeness of a plant individual is estimated as the inverse of the sum of all path lengths (link weights) between this plant individual and all other plant individuals in the unipartite network. Individuals with higher

values of weighted closeness are strongly connected with more plant individuals in the

population through shared frugivore species.

5.   Mean interaction overlap using Bray-Curtis index *(node similarity)*. This measure of interaction

overlap informs on the average similarity in frugivore use between pairs of plant individuals.

This metric indicates how different the frugivore assemblage of a given plant individual is

compared to the rest of the population (e.g., Gómez et al. 2010). Higher values (i.e., higher

overlap) indicate a higher similarity in interaction assemblage for a given plant individual with

respect to other individuals in the population.

<u>Comparison across individual-based networks</u>

In order to determine variation distribution in interaction structuring and node topology we

performed a principal component analysis (PCA). Previous studies have used PCA for comparing

network metrics (e.g., Sazima et al. 2010, Mora et al. 2018, Medeiros et al. 2018, Burin et al. 2021,

Acevedo-Quintero et al 2023). In order to compare plant individuals' interaction profiles, we

performed a PCA using the full set of node-level metric distributions estimated for plant individuals

within their population (i.e., individual node metrics derived from the full Bayesian posterior of each

individual). This resulted in an ordination of elements (1000 points per each plant individual, single

values sampled from the full posterior distributions of each individual-based network) in relation to

the multivariate space defined by node-level metrics. Thus, such PCA provides an exploratory

analysis of how plant individuals span the multivariate space of network metrics, where the location

of each individual characterizes its interaction profile.

With the aim of identifying distinct plant individual profiles we performed a discriminant analysis of

principal components (DAPC) using the adegenet R package (Jombart et al. 2023). We performed

1000 DAPCs using node-level metrics derived from the 1000 posterior samples of the individual-

based networks. We selected four groups or clusters as the minimal number that would facilitate

interpretation of each plant individual performance given its position in the network. In each of the

1000 analysis runs, plant individuals were assigned to one of the four groups given the highest

probability provided by the DAPC. This analysis was based on the variation captured by the first 3

PCs which together explained 86% of the variance, with eigenvalues >0.7. Cluster groups were

named post-hoc according to plants' relative position in the PCA into 'Keystone', 'Overlapping',

'Specialist', and 'Neutral' interaction profile types. We subsequently quantified the percentage of

individuals assigned to these four groups for each plant population in each of the 1000 repetitions.

<u>Data availability and code</u>

All datasets and code used to generate this study are available for download in Zenodo

(https://doi.org/10.5281/zenodo.10630818; Quintero et al. 2024) and GitHub repository

(https://github.com/PJordano-Lab/MS_individual-based_networks).

Acknowledgments

We are grateful to Miguel Jácome, Gemma Calvo, Pablo Homet, Pablo Villaba, Juan Miguel Arroyo,

Irene Mendoza and Eva Moracho, for support in data collection and processing with some of the

individual-based networks which was fundamental for this manuscript, as well as, discussion and

comments for improving the quality of this work. We are thankful to Daniel Stouffer for his help and

discussion on the network centralization metric, Giacomo Puglielli for encouraging ideas on PCA

analysis and Enrico Bazzicalupo for his support and insights on cluster analysis. We thank the

logistic and facilities support form ICTS-RBD Doñana and the Doñana National Park for the access

authorizations to collect data from many of the individual networks here presented. We are

thankful to two anonymous reviewers and the managing editor for their suggestions, which

substantially helped to improve the manuscript. Funding from LIFEWATCH-2019-09-CSIC-13

project and PID2022-136812NB-I00, Spanish Ministry of Science and Innovation, as well as funding

from the VI Plan Propio de Investigación of the Universidad de Sevilla (VI PPIT-US) supported this

research (PJ). EQ was supported by "la Caixa" Foundation fellowship LCF/BQ/DE18/1167000. BAC

received funding from the Ministry of Universities of the Spanish Government (Ref. FPU19_02552).

JI was supported by FPI grant PRE2018-085916 from the Spanish Ministry of Science and

Innovation. FRS was supported by VI PPIT-US and grants US-1381388 from Universidad de

Sevilla/Junta de Andalucía/FEDER-UE and CNS2022-135839 funded by

MICIU/AEI/10.13039/501100011033 and by European Union NextGenerationEU/PRTR.

References

1. Acevedo-Quintero, J. F., Zamora-Abrego, J. G., & Mancera-Rodríguez, N. J. (2023). Structural but not functional resistance of frugivore-plant interaction networks to the defaunation process. *Food Webs*, e00322. https://doi.org/10.1016/j.fooweb.2023.e00322

2. Alatalo, R. V. (1981). Problems in the measurement of evenness in ecology. *Oikos*, *37*(2), 199. https://doi.org/10.2307/3544465

3. Albert, C. H., Thuiller, W., Yoccoz, N. G., Soudant, A., Boucher, F., Saccone, P., & Lavorel, S. (2010). Intraspecific functional variability: Extent, structure and sources of variation. *Journal of Ecology*, *98*(3), 604–613. https://doi.org/10.1111/j.1365-2745.2010.01651.x

4. Albrecht, J., Classen, A., Vollstädt, M. G. R., Mayr, A., Mollel, N. P., Schellenberger Costa, D., Dulle, H. I., Fischer, M., Hemp, A., Howell, K. M., Kleyer, M., Nauss, T., Peters, M. K., Tschapka, M., Steffan-Dewenter, I., Böhning-Gaese, K., & Schleuning, M. (2018). Plant and animal functional diversity drive mutualistic network assembly across an elevational gradient. *Nature Communications*, *9*(1), 3177. https://doi.org/10.1038/s41467-018-05610-w

5. Almeida-Neto, M., & Ulrich, W. (2011). A straightforward computational approach for measuring nestedness using quantitative matrices. *Environmental Modelling & Software*, *26*(2), 173–178. https://doi.org/10.1016/j.envsoft.2010.08.003

6. Araújo, M. S., Bolnick, D. I., & Layman, C. A. (2011). The ecological causes of individual specialisation. *Ecology Letters*, *14*(9), 948–958. https://doi.org/10.1111/j.1461-0248.2011.01662.x

7. Araújo, M. S., & Costa-Pereira, R. (2013). Latitudinal gradients in intraspecific ecological diversity. Biology Letters, *9*(6), 20130778. https://doi.org/10.1098/rsbl.2013.0778

8. Arroyo-Correa, B., Bartomeus, I., & Jordano, P. (2021). Individual-based plant–pollinator networks are structured by phenotypic and microsite plant traits. Journal of Ecology, 109(8), 2832-2844.

9. Arroyo-Correa, B., Jordano, P., & Bartomeus, I. (2023). Intraspecific variation in species interactions promotes the feasibility of mutualistic assemblages. *Ecology Letters*, *24*, 448–459. https://doi.org/10.1111/ele.14163

10. Arroyo-Correa, B., Bartomeus, I., Jordano, P., Cagua, E. F., & Stouffer, D. B. (2024). Integrating intraspecific variation into population dynamics reveals how interacting species persist in mutualistic communities. bioRxiv, 2024-09. https://doi.org/10.1101/2024.09.12.612448

11. Barabási, A. L. (2016). *Network science*. Cambridge University Press.

12. Bascompte, J., & Jordano, P. (2007). Plant-animal mutualistic networks: The architecture of biodiversity. *Annual Review of Ecology, Evolution, and Systematics*, *38*(1), 567–593. https://doi.org/10.1146/annurev.ecolsys.38.091206.095818

13. Bascompte, J., & Jordano, P. (2014). *Mutualistic networks*. Princeton University Press.
https://doi.org/10.1515/9781400848720

14. Bascompte, J., Jordano, P., Melian, C. J., & Olesen, J. M. (2003). The nested assembly of plant-
animal mutualistic networks. *Proceedings of the National Academy of Sciences*, *100*(16), 9383–
9387. https://doi.org/10.1073/pnas.1633576100

15. Bascompte, J., Jordano, P., & Olesen, J. M. (2006). Asymmetric coevolutionary networks
facilitate biodiversity maintenance. *Science*, *312*, 431–433.
https://doi.org/10.1126/science.1123412

16. Bates, D., Mächler, M., Bolker, B., & Walker, S. (2015). Fitting linear mixed-effects models using
lme4. *Journal of Statistical Software*, *67*, 1–48. https://doi.org/10.18637/jss.v067.i01

17. Blüthgen, N., Frund, J., Vázquez, D. P., & Menzel, F. (2008). What do interaction network
metrics tell us about specialization and biological traits? *Ecology*, 89(12), 3387–3399.
https://doi.org/10.1890/07-2121.1

18. Bolnick, D. I., Amarasekare, P., Araújo, M. S., Bürger, R., Levine, J. M., Novak, M., Rudolf, V. H.
W., Schreiber, S. J., Urban, M. C., & Vasseur, D. A. (2011). Why intraspecific trait variation
matters in community ecology. *Trends in Ecology & Evolution*, *26*(4), 183–192.
https://doi.org/10.1016/j.tree.2011.01.009

19. Bolnick, D. I., Svanbäck, R., Araújo, M. S., & Persson, L. (2007). Comparative support for the
niche variation hypothesis that more generalized populations also are more heterogeneous.
Proceedings of the National Academy of Sciences, *104*(24), 10075–10079.
https://doi.org/10.1073/pnas.0703743104

20. Bolnick, D. I., Svanbäck, R., Fordyce, J. A., Yang, L. H., Davis, J. M., Hulsey, C. D., & Forister, M.
L. (2003). The ecology of individuals: Incidence and implications of individual specialization.
The American Naturalist, *106*(1), 1–28.

21. Bolnick, D. I., Yang, L. H., Fordyce, J. A., Davis, J. M., & Svanbäck, R. (2002). Measuring
individual-level resource specialization. *Ecology*, *83*(10), 2936–2941.
https://doi.org/10.1890/0012-9658(2002)083[2936:MILRS]2.0.CO;2

22. Brimacombe, C., Bodner, K., Michalska-Smith, M., Poisot, T. & Fortin, M.-J. (2023).
Shortcomings of reusing species interaction networks created by different sets of researchers.
PLoS Biol, 21, e3002068. doi: 10.1371/journal.pbio.3002068

23. Burin, G., Guimarães Jr, P. R., & Quental, T. B. (2021). Macroevolutionary stability predicts
interaction patterns of species in seed dispersal networks. *Science*, *372*, 733–737.
https://doi.org/10.1126/science.abf0556

24. Campo-Celada, M., Jordano, P., Benítez-López, A., Gutiérrez-Expósito, C., Rabadán-González,
J., & Mendoza, I. (2022). Assessing short and long-term variations in diversity, timing and body
condition of frugivorous birds. *Oikos*, *2022*, e08387. https://doi.org/10.1111/oik.08387

25. Carnicer, J., Jordano, P., & Melián, C. J. (2009). The temporal dynamics of resource use by
frugivorous birds: A network approach. *Ecology*, *90*(7), 1958–1970. https://doi.org/10.1890/07-
1939.1

26. Chase, J. M., & Knight, T. M. (2013). Scale-dependent effect sizes of ecological drivers on
biodiversity: Why standardised sampling is not enough. *Ecology Letters*, *16*(s1), 17–26.
https://doi.org/10.1111/ele.12112

27. Cirtwill, A. R., Stouffer, D. B., & Romanuk, T. N. (2015). Latitudinal gradients in biotic niche
breadth vary across ecosystem types. *Proceedings of the Royal Society B: Biological Sciences*,
282(1819), 20151589. https://doi.org/10.1098/rspb.2015.1589

28. Clark, J. S., Bell, D. M., Hersh, M. H., Kwit, M. C., Moran, E., Salk, C., Stine, A., Valle, D., & Zhu,
K. (2011). Individual-scale variation, species-scale differences: Inference needed to understand
diversity: Individual-scale variation, species-scale differences. Ecology Letters, 14(12), 1273–
1287. https://doi.org/10.1111/j.1461-0248.2011.01685.x

29. Cohen, J. E. (1978). *Food webs and niche space*. Princeton University Press.

30. Costa-Pereira, R., Araújo, M. S., Souza, F. L., & Ingram, T. (2019). Competition and resource
breadth shape niche variation and overlap in multiple trophic dimensions. *Proceedings of the*
Royal Society B: Biological Sciences, *286*(1902), 20190369.
https://doi.org/10.1098/rspb.2019.0369

31. Costa-Pereira, R., Rudolf, V. H. W., Souza, F. L., & Araújo, M. S. (2018). Drivers of individual
niche variation in coexisting species. *Journal of Animal Ecology*, *87*(5), 1452–1464.
https://doi.org/10.1111/1365-2656.12879

32. Crestani, A. C., Mello, M. A. R., & Cazetta, E. (2019). Interindividual variations in plant and fruit
traits affect the structure of a plant-frugivore network. *Acta Oecologica*, *95*, 120–127.
https://doi.org/10.1016/j.actao.2018.11.003

33. Csárdi, G., & Nepusz, T. (2006). The igraph software package for complex network research
<http://igraph.org>. *InterJournal Complex Systems*, *1695*, 1–9.

34. de Oliveira Lima, G., Leite, A. V., Souza, C. S., Castro, C. C., & de Santana Bezerra, E. L. (2020).
A multilayer network in an herbaceous tropical community reveals multiple roles of floral
visitors. *Oikos*, *129*(8), 1141–1151. https://doi.org/10.1111/oik.06565

35. Dormann, C. F. (2011). How to be a specialist? Quantifying specialisation in pollination
networks. *Network Biology*, *1*(1), 1–20.

36. Dormann, C. F., Frund, J., Bluthgen, N., & Gruber, B. (2009). Indices, graphs and null models:
Analyzing bipartite ecological networks. *The Open Ecology Journal*, *2*(1), 7–24.
https://doi.org/10.2174/1874213000902010007

37. Dormann, C. F., Gruber, B., & Fruend, J. (2008). Introducing the bipartite package: Analysing
ecological networks. *R News*, *8*(2), 8–11.

38. Dunne, J. A., Williams, R. J., & Martinez, N. D. (2002). Food-web structure and network theory:
The role of connectance and size. *Proceedings of the National Academy of Sciences*, *99*(20),
12917–12922. https://doi.org/10.1073/pnas.192407699

39. Fontaine, C., Guimarães Jr, P. R., Kéfi, S., Loeuille, N., Memmott, J., van der Putten, W. H., van
Veen, F. J. F., & Thébault, E. (2011). The ecological and evolutionary implications of merging
different types of networks. *Ecology Letters*, *14*(11), 1170–1181. https://doi.org/10.1111/j.1461-
0248.2011.01688.x

40. Fortuna, M. A., & Bascompte, J. (2008). The network approach in ecology. In *Unity in diversity:*
Reflections on ecology after the legacy of Ramon Margalef. Fundación BBVA.

41. Freeman, L. C., Roeder, D., & Mulholland, R. R. (1979). Centrality in social networks: II.
Experimental results. *Social Networks*, *2*(2), 119–141. https://doi.org/10.1016/0378-
8733(79)90002-9

42. Gabry J, Češnovar R, Johnson A, Bronder S (2024). cmdstanr: R Interface to 'CmdStan'. R
package version 0.8.1, https://discourse.mc-stan.org, https://mc-stan.org/cmdstanr/

43. Godínez-Alvarez, H., Ríos-Casanova, L., & Peco, B. (2020). Are large frugivorous birds better
seed dispersers than medium- and small-sized ones? Effect of body mass on seed dispersal
effectiveness. *Ecology and Evolution*, *10*(12), 6136–6143. https://doi.org/10.1002/ece3.6285

44. Gómez, J. M., Abdelaziz, M., Lorite, J., Jesús Muñoz-Pajares, A., & Perfectti, F. (2010). Changes
in pollinator fauna cause spatial variation in pollen limitation. *Journal of Ecology*, *98*(5), 1243–
1252. https://doi.org/10.1111/j.1365-2745.2010.01691.x

45. Gómez, J. M., González-Megías, A., Armas, C., Narbona, E., Navarro, L., & Perfectti, F. (2023).
The role of phenotypic plasticity in shaping ecological networks. *Ecology Letters*, *26*(S1), S47–
S61. https://doi.org/10.1111/ele.14192

46. Gómez, J. M., & Perfectti, F. (2012). Fitness consequences of centrality in mutualistic
individual-based networks. *Proceedings of the Royal Society B: Biological Sciences*, *279*(1734),
1754–1760. https://doi.org/10.1098/rspb.2011.2244

47. González-Varo, J. P., & Traveset, A. (2016). The labile limits of forbidden interactions. *Trends in*
Ecology & Evolution, *31*(9), 700–710. https://doi.org/10.1016/j.tree.2016.06.009

48. Grinnell, J. (1917). The niche-relationships of the California thrasher. *The Auk*, *34*(4), 427–433.
https://doi.org/10.2307/4072271

49. Guerra, T. J., Dayrell, R. L. C., Arruda, A. J., Dáttilo, W., Teixido, A. L., Messeder, J. V. S., &
Silveira, F. A. O. (2017). Intraspecific variation in fruit–frugivore interactions: Effects of fruiting

neighborhood and consequences for seed dispersal. *Oecologia*, *185*(2), 233–243.
https://doi.org/10.1007/s00442-017-3943-z

50. Guimarães, P. R. (2020). The structure of ecological networks across levels of organization.
Annual Review of Ecology, Evolution, and Systematics, *51*(1), 433–460.
https://doi.org/10.1146/annurev-ecolsys-012220-120819

51. Guimarães, P. R., Jordano, P., & Thompson, J. N. (2011). Evolution and coevolution in
mutualistic networks. *Ecology Letters*, *14*(9), 877–885. https://doi.org/10.1111/j.1461-
0248.2011.01649.x

52. Hällfors, M. H., Heikkinen, R. K., Kuussaari, M., Lehikoinen, A., Luoto, M., Pöyry, J., Virkkala,
R., Saastamoinen, M., & Kujala, H. (2023). Recent range shifts of moths, butterflies, and birds
are driven by the breadth of their climatic niche. *Evolution Letters*, qrad004.
https://doi.org/10.1093/evlett/qrad004

53. Hill, M. O. (1973). Diversity and evenness: A unifying notation and its consequences. *Ecology*,
54(2), 427–432. https://doi.org/10.2307/1934352

54. Hsieh, T. C., Ma, K. H., & Chao, A. (2016). iNEXT: An R package for rarefaction and
extrapolation of species diversity (Hill numbers). *Methods in Ecology and Evolution*, *7*(12), 1451–
1456. https://doi.org/10.1111/2041-210X.12613

55. Hurlbert, S. H. (1971). The nonconcept of species diversity: A critique and alternative
parameters. Ecology, 52(4), 577–586. https://doi.org/10.2307/1934145

56. Isla, J., Jácome-Flores, M., Arroyo, J. M., & Jordano, P. (2023). The turnover of plant–frugivore
interactions along plant range expansion: Consequences for natural colonization processes.
Proceedings of the Royal Society B: Biological Sciences, *290*(1999), 20222547.
https://doi.org/10.1098/rspb.2022.2547

57. Jácome-Flores, M. E., Jordano, P., Delibes, M., & Fedriani, J. M. (2020). Interaction motifs
variability in a Mediterranean palm under environmental disturbances: The mutualism–
antagonism continuum. *Oikos*, *129*(3), 367–379. https://doi.org/10.1111/oik.06688

58. Jombart, T., Kamvar, Z. N., Collins, C., Lustrik, R., Beugin, M.-P., Knaus, B. J., Solymos, P.,
Mikryukov, V., Schliep, K., Maié, T., Morkovsky, L., Ahmed, I., Cori, A., Calboli, F., Ewing, R. J.,
Michaud, F., DeCamp, R., Courtiol, A., Clark, L. V., & Dimens, P. (2023). *adegenet: Exploratory
Analysis of Genetic and Genomic Data* (2.1.10) [Computer software]. https://cran.r-
project.org/web/packages/adegenet/index.html

59. Jordano, P. (1987). Patterns of mutualistic interactions in pollination and seed dispersal:
Connectance, dependence asymmetries, and coevolution. *The American Naturalist*, *129*(5),
657–677. https://doi.org/10.1086/284665

60. Jordano, P., Bascompte, J., & Olesen, J. M. (2003). Invariant properties in coevolutionary
networks of plant-animal interactions: Invariant properties in coevolutionary networks. *Ecology*
Letters, *6*(1), 69–81. https://doi.org/10.1046/j.1461-0248.2003.00403.x

61. Jordano, P., García, C., Godoy, J. A., & García-Castaño, J. L. (2007). Differential contribution of
frugivores to complex seed dispersal patterns. *Proceedings of the National Academy of Sciences*,
104(9), 3278–3282. https://doi.org/10.1073/pnas.0606793104

62. Jordano, P., & Schupp, E. W. (2000). Seed disperser effectiveness: The quantity component
and patterns of seed rain for *Prunus mahaleb*. *Ecological Monographs*, *70*(4), 591–615.
https://doi.org/10.2307/2657187

63. Jordano, P. (2016). Sampling networks of ecological interactions. *Functional Ecology*, *30*(12),
1883–1893. https://doi.org/10.1111/1365-2435.12763

64. Julliard, R., Clavel, J., Devictor, V., Jiguet, F., & Couvet, D. (2006). Spatial segregation of
specialists and generalists in bird communities. *Ecology Letters*, *9*(11), 1237–1244.
https://doi.org/10.1111/j.1461-0248.2006.00977.x

65. Koffel, T., Daufresne, T., & Klausmeier, C. A. (2021). From competition to facilitation and
mutualism: A general theory of the niche. *Ecological Monographs*, *91*(3), e01458.
https://doi.org/10.1002/ecm.1458

66. McCann, K., Hastings, A., & Huxel, G. R. (1998). Weak trophic interactions and the balance of
nature. *Nature*, *395*(6704), Article 6704. https://doi.org/10.1038/27427

67. Medeiros, L. P., Garcia, G., Thompson, J. N., & Guimarães, P. R. (2018). The geographic mosaic
of coevolution in mutualistic networks. *Proceedings of the National Academy of Sciences*,
115(47), 12017–12022. https://doi.org/10.1073/pnas.1809088115

68. Mello, M. A. R., Marquitti, F. M. D., Guimarães, P. R., Kalko, E. K. V., Jordano, P., & De Aguiar,
M. A. M. (2011). The missing part of seed dispersal networks: Structure and robustness of bat-
fruit interactions. *PLoS ONE*, *6*(2), e17395. https://doi.org/10.1371/journal.pone.0017395

69. Memmott, J. (1999). The structure of a plant-pollinator food web. *Ecology Letters*, *2*(5), 276–
280. https://doi.org/10.1046/j.1461-0248.1999.00087.x

70. Michalska-Smith, M., Song, Z., Spawn-Lee, S. A., Hansen, Z. A., Johnson, M., May, G., Borer, E.
T., Seabloom, E. W., & Kinkel, L. L. (2022). Network structure of resource use and niche overlap
within the endophytic microbiome. *The ISME Journal*, *16*(2), 435–446.
https://doi.org/10.1038/s41396-021-01080-z

71. Miguel, M. F., Jordano, P., Tabeni, S., & Campos, C. M. (2018). Context-dependency and
anthropogenic effects on individual plant-frugivore networks. *Oikos*, *127*(7), 1045–1059.
https://doi.org/10.1111/oik.04978

72. Mora, B. B., Gravel, D., Gilarranz, L. J., Poisot, T., & Stouffer, D. B. (2018). Identifying a
common backbone of interactions underlying food webs from different ecosystems. *Nature*
Communications, *9*(1), 2603. https://doi.org/10.1038/s41467-018-05056-0

73. Müller, C. B., Adriaanse, I. C. T., Belshaw, R., & Godfray, H. C. J. (1999). The structure of an
aphid–parasitoid community. *Journal of Animal Ecology*, *68*(2), 346–370.
https://doi.org/10.1046/j.1365-2656.1999.00288.x

74. Newman, M. E. J. (2002). Assortative mixing in networks. *Physical Review Letters*, *89*(20),
208701. https://doi.org/10.1103/PhysRevLett.89.208701

75. Newman, M. E. J. (2006). Modularity and community structure in networks. *Proceedings of the*
National Academy of Sciences, *103*(23), 8577–8582. https://doi.org/10.1073/pnas.0601602103

76. Oksanen, J., Simpson, G. L., Blanchet, F. G., Kindt, R., Legendre, P., Minchin, P. R., O'Hara, R.
B., Solymos, P., Stevens, M. H. H., Szoecs, E., Wagner, H., Barbour, M., Bedward, M., Bolker,
B., Borcard, D., Carvalho, G., Chirico, M., Caceres, M. D., Durand, S., ... Weedon, J. (2022).
vegan: Community Ecology Package (2.6-4) [Computer software]. https://cran.r-
project.org/web/packages/vegan/index.html

77. Opsahl, T., Agneessens, F., & Skvoretz, J. (2010). Node centrality in weighted networks:
Generalizing degree and shortest paths. *Social Networks*, *32*(3), 245–251.
https://doi.org/10.1016/j.socnet.2010.03.006

78. Patefield, W. M. (1981). Algorithm AS 159: An afficient method of generating random R × C
tables with given row and column totals. Journal of the Royal Statistical Society. Series C
(Applied Statistics), 30(1), 91–97. https://doi.org/10.2307/2346669

79. Phillips, R. D., Peakall, R., Van Der Niet, T., & Johnson, S. D. (2020). Niche perspectives on
plant–pollinator interactions. *Trends in Plant Science*, *25*(8), 779–793.
https://doi.org/10.1016/j.tplants.2020.03.009

80. Pichon, B., Goff, R. L., Morlon, H., & Perez-Lamarque, B. (2023). Telling mutualistic and
antagonistic ecological networks apart by learning their multiscale structure. *bioRxiv*.
https://doi.org/10.1101/2023.04.04.535603

81. Poisot, T., Canard, E., Mouquet, N., & Hochberg, M. E. (2012). A comparative study of
ecological specialization estimators. *Methods in Ecology and Evolution*, *3*(3), 537–544.
https://doi.org/10.1111/j.2041-210X.2011.00174.x

82. Poisot, T., Cirtwill, A. R., Cazelles, K., Gravel, D., Fortin, M.-J., & Stouffer, D. B. (2016). The
structure of probabilistic networks. *Methods in Ecology and Evolution*, *7*(3), 303–312.
https://doi.org/10.1111/2041-210X.12468

83. Ponisio, L. C., Valdovinos, F. S., Allhoff, K. T., Gaiarsa, M. P., Barner, A., Guimarães, P. R.,
Hembry, D. H., Morrison, B., & Gillespie, R. (2019). A network perspective for community
assembly. *Frontiers in Ecology and Evolution*, *7*, 103. https://doi.org/10.3389/fevo.2019.00103

84. Quintero, E., Isla, J., & Jordano, P. (2022). Methodological overview and data-merging
approaches in the study of plant–frugivore interactions. *Oikos*, *e08379*, 1–18.
https://doi.org/10.1111/oik.08379

85. Quintero, E., Rodríguez-Sánchez, F., & Jordano, P. (2023). Reciprocity and interaction
effectiveness in generalised mutualisms among free-living species. *Ecology Letters*, *26*(1), 132–
146. https://doi.org/10.1111/ele.14141

86. Quintero, E., Arroyo-Correa, B., Isla, J., Rodriguez-Sánchez, F., & Jordano, P. (2024).
Downscaling mutualistic networks from species to individuals reveals consistent interaction
niches and roles within plant populations. [Dataset]. Zenodo.
https://doi.org/10.5281/zenodo.12724898

87. Rehling, F., Jongejans, E., Schlautmann, J., Albrecht, J., Fassbender, H., Jaroszewicz, B.,
Matthies, D., Waldschmidt, L., Farwig, N., & Schabo, D. G. (2023). Common seed dispersers
contribute most to the persistence of a fleshy-fruited tree. *Communications Biology*, *6*(1), 330.
https://doi.org/10.1038/s42003-023-04647-y

88. Rodríguez-Rodríguez, M. C., Jordano, P., & Valido, A. (2017). Functional consequences of plant-
animal interactions along the mutualism-antagonism gradient. *Ecology*, *98*(5), 1266–1276.
https://doi.org/10.1002/ecy.1756

89. Rodríguez-Sánchez, F. (2024a). *BayesianWebs: Bayesian Modelling of Bipartite Networks*.
https://github.com/Pakillo/BayesianWebs

90. Rodriguez-Sanchez, F (2024b). network.tools: Tools to analyse and visualise bipartite
networks. R package version 0.0.5, https://pakillo.github.io/network.tools

91. Rother, D. C., Pizo, M. A., & Jordano, P. (2016). Variation in seed dispersal effectiveness: The
redundancy of consequences in diversified tropical frugivore assemblages. *Oikos*, *125*(3), 336–
342. https://doi.org/10.1111/oik.02629

92. Roughgarden, J. (1972). Evolution of niche width. *The American Naturalist*, *106*(952), 683–718.

93. Roughgarden, J. (1979). *Theory of population genetics and evolutionary ecology: An introduction*.
MacMillan.

94. Sallabanks, R. (1993). Hierarchical mechanisms of fruit selection by an avian frugivore. *Ecology*,
74(5), 1326–1336. https://doi.org/10.2307/1940063

95. Sazatornil, F. D., Moré, M., Benitez-Vieyra, S., Cocucci, A. A., Kitching, I. J., Schlumpberger, B.
O., Oliveira, P. E., Sazima, M., & Amorim, F. W. (2016). Beyond neutral and forbidden links:
Morphological matches and the assembly of mutualistic hawkmoth–plant networks. *Journal of*
Animal Ecology, *85*(6), 1586–1594. https://doi.org/10.1111/1365-2656.12509

96. Sazima, C., Guimarães, P. R., Dos Reis, S. F., & Sazima, I. (2010). What makes a species central
in a cleaning mutualism network? *Oikos*, *119*(8), 1319–1325. https://doi.org/10.1111/j.1600-
0706.2009.18222.x

97. Schleuning, M., Blüthgen, N., Flörchinger, M., Braun, J., Schaefer, H. M., & Böhning-Gaese, K.
(2011). Specialization and interaction strength in a tropical plant–frugivore network differ
among forest strata. *Ecology*, *92*(1), 26–36. https://doi.org/10.1890/09-1842.1

98. Segar, S. T., Fayle, T. M., Srivastava, D. S., Lewinsohn, T. M., Lewis, O. T., Novotny, V.,
Kitching, R. L., & Maunsell, S. C. (2020). The role of evolution in shaping ecological networks.
Trends in Ecology & Evolution, *35*(5), 454–466. https://doi.org/10.1016/j.tree.2020.01.004

99. Siefert, A., Violle, C., Chalmandrier, L., Albert, C. H., Taudiere, A., Fajardo, A., Aarssen, L. W.,
Baraloto, C., Carlucci, M. B., Cianciaruso, M. V., de L. Dantas, V., de Bello, F., Duarte, L. D. S.,
Fonseca, C. R., Freschet, G. T., Gaucherand, S., Gross, N., Hikosaka, K., Jackson, B., … Wardle,
D. A. (2015). A global meta-analysis of the relative extent of intraspecific trait variation in plant
communities. *Ecology Letters*, *18*(12), 1406–1419. https://doi.org/10.1111/ele.12508

100.   Simmons, B. I., Wauchope, H. S., Amano, T., Dicks, L. V., Sutherland, W. J., & Dakos, V.
(2020). Estimating the risk of species interaction loss in mutualistic communities. *PLOS*
Biology, *18*(8), e3000843. https://doi.org/10.1371/journal.pbio.3000843

101.   Snell, R. S., Beckman, N. G., Fricke, E., Loiselle, B. A., Carvalho, C. S., Jones, L. R., Lichti, N.
I., Lustenhouwer, N., Schreiber, S. J., Strickland, C., Sullivan, L. L., Cavazos, B. R., Giladi, I.,
Hastings, A., Holbrook, K. M., Jongejans, E., Kogan, O., Montaño-Centellas, F., Rudolph, J., …
Schupp, E. W. (2019). Consequences of intraspecific variation in seed dispersal for plant
demography, communities, evolution and global change. *AoB PLANTS*, *11*(4), plz016.
https://doi.org/10.1093/aobpla/plz016

102.   Snow, B. K., & Snow, D. (1988). *Birds and Berries*. T & AD Poyser.

103.   Solé, R. V., & Valverde, S. (2004). Information theory of complex networks: On evolution
and architectural constraints. In E. Ben-Naim, H. Frauenfelder, & Z. Toroczkai (Eds.), *Complex*
Networks (Vol. 650, pp. 189–207). Springer Berlin Heidelberg. https://doi.org/10.1007/978-3-
540-44485-5_9

104.   Stan Development Team. (2024). Stan Modeling Language Users Guide and Reference
Manual, 2.32.2. https://mc-stan.org

105.   Thiel, S., Willems, F., Farwig, N., Rehling, F., Schabo, D. G., Schleuning, M., Shahuano Tello,
N., Töpfer, T., Tschapka, M., Heymann, E. W., & Heer, K. (2023). Vertically stratified frugivore
community composition and interaction frequency in a liana fruiting across forest strata.
Biotropica, *55*, 650–664. https://doi.org/10.1111/btp.13216

106.   Thompson, J. N. (1999). The raw material for coevolution. *Oikos*, *84*(1), 5.
https://doi.org/10.2307/3546861

107.   Tur, C., Vigalondo, B., Trøjelsgaard, K., Olesen, J. M., & Traveset, A. (2014). Downscaling
956   pollen-transport networks to the level of individuals. *Journal of Animal Ecology*, *83*(1), 306–317.
957   https://doi.org/10.1111/1365-2656.12130

958   108.   Valenzuela-Ospina, L., & Kattan, G. H. (2022). Downscaling plant-frugivore interaction
959   networks in an assemblage of fig consumers. *Biota Colombiana*, *23*(1), e1011.
960   https://doi.org/10.21068/2539200X.1011

961   109.   Van Valen, L. (1965). Morphological variation and width of ecological niche. *The American
962   Naturalist*, *99*(908), 377–390.

963   110.   Vázquez, D. P., & Aizen, M. A. (2004). Asymmetric specialization: A pervasive feature of
964   plant–pollinator interactions. *Ecology*, *85*(5), 1251–1257. https://doi.org/10.1890/03-3112

965   111. Vázquez, D. P., Melián, C. J., Williams, N. M., Blüthgen, N., Krasnov, B. R., & Poulin, R. (2007).
966   Species abundance and asymmetric interaction strength in ecological networks. *Oikos*, *116*(7),
967   1120–1127. https://doi.org/10.1111/j.0030-1299.2007.15828.x

968   112.   Vissoto, M., Vizentin-Bugoni, J., Sendoya, S. F., Gomes, G. C., & Dias, R. A. (2022). Plant
969   height and spatial context influence individual connectivity and specialization on seed
970   dispersers in a tree population. *Oecologia*, *198*(3), 721–731. https://doi.org/10.1007/s00442-022-
971   05142-7

972   113. Wang, L.-L., Yang, Y.-P., & Duan, Y.-W. (2021). Pollinator individual-based networks reveal the
973   specialized plant–pollinator mutualism in two biodiverse communities. *Ecology and Evolution*,
974   *11*(23), 17509–17518. https://doi.org/10.1002/ece3.8384

975   114.   Wheelwright, N. T., & Orians, G. H. (1982). Seed dispersal by animals: Contrasts with pollen
976   dispersal, problems of terminology, and constraints on coevolution. *The American Naturalist*,
977   *119*(3), 402–413.

978   115.   Young, J.-G., Valdovinos, F. S., & Newman, M. E. J. (2021). Reconstruction of plant–
979   pollinator networks from observational data. Nature Communications, 12(1), 3911.
980   https://doi.org/10.1038/s41467-021-24149-x

Individual-based networks references:

1.   Crestani, A. C., Mello, M. A. R., & Cazetta, E. (2019). Interindividual variations in plant and fruit
traits affect the structure of a plant-frugivore network. *Acta Oecologica*, *95*, 120–127.
https://doi.org/10.1016/j.actao.2018.11.003

2.   Friedemann, P., Côrtes, M. C., de Castro, E. R., Galetti, M., Jordano, P., & Guimarães Jr, P. R.
(2022). The individual-based network structure of palm-seed dispersers is explained by a
rainforest gradient. *Oikos*, *2022*, e08384. https://doi.org/10.1111/oik.08384

3. Gopal, A., Mudappa, D., Raman, T. R. S., & Naniwadekar, R. (2020). Forest cover and fruit crop
size differentially influence frugivory of select rainforest tree species in Western Ghats, India.
Biotropica, *52*(5), 871–883. https://doi.org/10.1111/btp.12810

4. Guerra, T. J., Dayrell, R. L. C., Arruda, A. J., Dáttilo, W., Teixido, A. L., Messeder, J. V. S., &
Silveira, F. A. O. (2017). Intraspecific variation in fruit–frugivore interactions: Effects of fruiting
neighborhood and consequences for seed dispersal. *Oecologia*, *185*(2), 233–243.
https://doi.org/10.1007/s00442-017-3943-z

5. Isla, J., Jácome-Flores, M., Arroyo, J. M., & Jordano, P. (2023). The turnover of plant–frugivore
interactions along plant range expansion: Consequences for natural colonization processes.
Proceedings of the Royal Society B: Biological Sciences, *290*(1999), 20222547.
https://doi.org/10.1098/rspb.2022.2547

6. Jácome-Flores, M. E., Jordano, P., Delibes, M., & Fedriani, J. M. (2020). Interaction motifs
variability in a Mediterranean palm under environmental disturbances: The mutualism–
antagonism continuum. *Oikos*, *129*(3), 367–379. https://doi.org/10.1111/oik.06688

7. Jayanth, A., Isvaran, K., & Naniwadekar, R. (2024). Drivers of intraspecific variation in seed
dispersal can differ across two species of fleshy-fruited savanna plants. *Biotropica*, *56*, e13322.
https://doi.org/10.1111/btp.13322.

8. Jordano, P. (1995). Frugivore-mediated selection on fruit and seed size: Birds and St. Lucie's
cherry, *Prunus mahaleb*. *Ecology*, *76*(8), 2627–2639. https://doi.org/10.2307/2265833

9. Jordano, P. (2024). *Material for the course: Curso Posgraduação Frugivoria 2016, UNESP Rio
Claro, Brazil*(course_materials) [dataset]. Zenodo. https://doi.org/10.5281/zenodo.10478535

10. Jordano, P., & Schupp, E. W. (2000). Seed disperser effectiveness: The quantity component
and patterns of seed rain for *Prunus mahaleb*. *Ecological Monographs*, *70*(4), 591–615.
https://doi.org/10.2307/2657187

11. Lamperty, T., Karubian, J., & Dunham, A. E. (2021). Ecological drivers of intraspecific variation
in seed dispersal services of a common neotropical palm. *Biotropica*, *53*(4), 1226–1237.
https://doi.org/10.1111/btp.12966

12. Miguel, M. F., Jordano, P., Tabeni, S., & Campos, C. M. (2018). Context-dependency and
anthropogenic effects on individual plant-frugivore networks. *Oikos*, *127*(7), 1045–1059.
https://doi.org/10.1111/oik.04978

13. Quintero, E., Rodríguez-Sánchez, F., & Jordano, P. (2023). Reciprocity and interaction
effectiveness in generalised mutualisms among free-living species. *Ecology Letters*, *26*(1), 132–
146. https://doi.org/10.1111/ele.14141

14. Ramaswami, G., Somnath, P., & Quader, S. (2017). Plant-disperser mutualisms in a semi-arid
habitat invaded by Lantana camara L. *Plant Ecology*, *218*(8), 935–946.
https://doi.org/10.1007/s11258-017-0741-y

15. Rodriguez-Sanchez, F. (2010). *An integrative framework to investigate species responses to*
climate change: Biogeography and ecology of relict trees in the Mediterranean. Universidad de
Sevilla.

16. Thiel, S., Willems, F., Farwig, N., Rehling, F., Schabo, D. G., Schleuning, M., Shahuano Tello,
N., Töpfer, T., Tschapka, M., Heymann, E. W., & Heer, K. (2023). Vertically stratified frugivore
community composition and interaction frequency in a liana fruiting across forest strata.
Biotropica, *55*, 650–664. https://doi.org/10.1111/btp.13216

17. Vergara-Tabares, D. L., Blendinger, P. G., Tello, A., Peluc, S. I., & Tecco, P. A. (2022). Fleshy-
fruited invasive shrubs indirectly increase native tree seed dispersal. *Oikos*, *2022*(2).
https://doi.org/10.1111/oik.08311

18. Villalva, P., Arroyo-Correa, B., Calvo, G., Homet, P., Isla, J., Mendoza, I., Moracho, E.,
Quintero, E., Rodríguez-Sánchez, F., & Jordano, P. (2023). *FRUGIVORY CAMTRAP: A dataset*
of plant-animal interactions recorded with camera traps [Text/csv]. DIGITAL.CSIC.
https://doi.org/10.20350/DIGITALCSIC/15623

19. Vissoto, M., Vizentin-Bugoni, J., Sendoya, S. F., Gomes, G. C., & Dias, R. A. (2022). Plant
height and spatial context influence individual connectivity and specialization on seed
dispersers in a tree population. *Oecologia*, *198*(3), 721–731. https://doi.org/10.1007/s00442-
022-05142-7

1043

Species-based network references:

1. Albrecht, J., Bohle, V., Berens, D. G., Jaroszewicz, B., Selva, N., & Farwig, N. (2015). Variation
in neighbourhood context shapes frugivore-mediated facilitation and competition among co-
dispersed plant species. *Journal of Ecology*, *103*(2), 526–536. https://doi.org/10.1111/1365-
2745.12375
2. Alves, K. J. F. (2008). *Composição da avifauna e frugivoria por aves em um mosaico sucessional*
na mata Atlântica. Universidade Estadual Paulista.
3. Andrade, P. C., Mota, J. V. L., & de Carvalho, A. A. F. (2011). Interações mutualísticas entre
aves frugívoras e plantas em um fragmento urbano de Mata Atlântica, Salvador, BA. *Revista*
Brasileira de Ornitologia, *19*(1), 63–73.
4. Athiê, S. (2009). *Composição da avifauna e frugivoria por aves em um mosaico de vegetação*
secundária em Rio Claro, região centro-leste do estado de São Paulo. Universidade Federal de
São Carlos.
5. Baird, J. W. (1980). The selection and use of fruit by birds in an eastern forest. *Wilson Bulletin*,
92(1), 63–73.
6. Beehler, B. (1983). Frugivory and polygamy in birds of paradise. *The Auk*, *100*(1), 1–12.
https://www.jstor.org/stable/4086272
7. Burns, K. C. (2013). What causes size coupling in fruit–frugivore interaction webs? *Ecology*,
94(2), 295–300. https://doi.org/10.1890/12-1161.1

8.   Castro, E. R. de. (2007). *Fenologia reprodutiva do palmito* Euterpe edulis *(Arecaceae) e sua*
influência na abundância de aves frugívoras na floresta atlântica. Universidade Estadual
Paulista.
9.   Correia, J. M. de S. (1997). *Utilização de espécies frutíferas de mata Atlântica na alimentação da*
avifauna da reserva biológica de Poço das Antas, RJ. Universidade de Brasília.
10.   Crome, F. (1975). The ecology of fruit pigeons in tropical Northern Queensland. *Wildlife*
Research, 2(2), 155–185. https://doi.org/10.1071/WR9750155
11.   Fadini, R. F., & de Marco Jr, P. (2004). Interações entre aves frugívoras e plantas em um
fragmento de mata atlântica de Minas Gerais. *Ararajuba*, 12(2), 97–103.
12.   Frost, P. G. H. (1980). Fruit-frugivore interactions in a South African coastal dune forest. In R.
Nöhring (Ed.), *Acta XVII Congressus Internationalis Ornithologici* (Vol. 2, pp. 1179–1184). Verlag
der Deutschen Ornithologen-Gesellschaft.
13.   Galetti, M., & Pizo, M. A. (1996). Fruit eating by birds in a forest fragment in southeastern
Brasil. *Ararajuba*, 4(2), 71–79.
14.   Garcia, Q. S., Rezende, J. L. P., & Aguiar, L. M. S. (2000). Seed dispersal by bats in a disturbed
area of Southeastern Brazil. *Revista de Biología Tropical*, 48(1), 125–128.
15.   García-Castaño, J. L. (2001). *Consecuencias demográficas de la dispersión de semillas por aves y*
mamíferos frugívoros en la vegetación Mediterránea de montaña. Universidad de Sevilla.
16.   Gorchov, D. L., Cornejo, F., Ascorra, C. F., & Jaramillo, M. (1995). Dietary overlap between
frugivorous birds and bats in the Peruvian Amazon. *Oikos*, 74(2), 235.
https://doi.org/10.2307/3545653
17.   Guitián, J. (1984). Ecología de una comunidad de passeriformes en un bosque montano de la
Cordillera Cantábrica Occidental [Http://purl.org/dc/dcmitype/Text, Universidade de Santiago
de Compostela]. In *Ecología de una comunidad de passeriformes en un bosque montano de la*
Cordillera Cantábrica Occidental.
https://investigacion.usc.es/documentos/5d1df66029995204f766a2ba?lang=es
18.   Hasui, E. (1994). *O papel das aves frugívoras na dispersão de sementes em um fragmento de*
floresta estacional semidecídua secundária, em São Paulo, SP [Universidade de São Paulo].
https://repositorio.usp.br/item/000742034
19.   Heleno, R. H., Olesen, J. M., Nogales, M., Vargas, P., & Traveset, A. (2013). Seed dispersal
networks in the Galápagos and the consequences of alien plant invasions. *Proceedings of the*
Royal Society B: Biological Sciences, 280(1750), 20122112.
https://doi.org/10.1098/rspb.2012.2112
20.   Hernández-Montero, J. R., Saldaña-Vázquez, R. A., Galindo-González, J., & Sosa, V. J. (2015).
Bat-fruit interactions are more specialized in shaded-coffee plantations than in tropical
mountain cloud forest fragments. *PLOS ONE*, 10(5), e0126084.
https://doi.org/10.1371/journal.pone.0126084
21.   Lopez, J. E., & Vaughan, C. (2004). Observations on the role of frugivorous bats as seed
dispersers in Costa Rican secondary humid forests. *Acta Chiropterologica*, 6(1), 111–119.
https://doi.org/10.3161/001.006.0109
22.   Menke, S., Böhning-Gaese, K., & Schleuning, M. (2012). Plant-frugivore networks are less
specialized and more robust at forest-farmland edges than in the interior of a tropical forest.
Oikos, 121(10), 1553–1566. https://doi.org/10.1111/j.1600-0706.2011.20210.x

23. Noma, N., & Yumoto, T. (1997). Fruiting phenology of animal-dispersed plants in response to
winter migration of frugivores in a warm temperate forest on Yakushima Island, Japan.
Ecological Research, *12*(2), 119–129. https://doi.org/10.1007/BF02523777

24. Olesen, J. M., Bascompte, J., Dupont, Y. L., Elberling, H., Rasmussen, C., & Jordano, P. (2011).
Missing and forbidden links in mutualistic networks. *Proceedings of the Royal Society B:*
Biological Sciences, *278*(1706), 725–732. https://doi.org/10.1098/rspb.2010.1371

25. Palmeirim, J. M., Gorchov, D. L., & Stoleson, S. (1989). Trophic structure of a neotropical
frugivore community: Is there competition between birds and bats? *Oecologia*, *79*(3), 403–411.
https://doi.org/10.1007/BF00384321

26. Passos, F. C., Silva, W. R., Pedro, W. A., & Bonin, M. R. (2003). Frugivoria em morcegos
(Mammalia, Chiroptera) no Parque Estadual Intervales, sudeste do Brasil. *Revista Brasileira de*
Zoologia, *20*(3), 511–517. https://doi.org/10.1590/S0101-81752003000300024

27. Pedro, W. A. (1992). *Estrutura de uma taxocenose de morcegos da reserva do Panga*
(Uberlandia, MG), com ênfase nas relações tróficas em Phyllostomidae (Mammalia: Chiroptera).
Universidade Estadual de Campinas.

28. Pizo, M. A. (2004). Frugivory and habitat use by fruit-eating birds in a fragmented landscape of
southeast brazil. *Ornitologia Neotropical*, *15*(Suppl.), 117–126.

29. Poulin, B., Wright, S. J., Lefebvre, G., & Calderón, O. (1999). Interspecific synchrony and
asynchrony in the fruiting phenologies of congeneric bird-dispersed plants in Panama. *Journal*
of Tropical Ecology, *15*(2), 213–227. https://doi.org/10.1017/S0266467499000760

30. Ribeiro Da Silva, F., Montoya, D., Furtado, R., Memmott, J., Pizo, M. A., & Rodrigues, R. R.
(2015). The restoration of tropical seed dispersal networks: Restoration of tropical seed
dispersal networks. *Restoration Ecology*, *23*(6), 852–860. https://doi.org/10.1111/rec.12244

31. Robinson, V. (2015). *Índice de importância de diferentes espécies de plantas na atração de aves*
para uma área reflorestada em Piracicaba. Universidade Estadual Paulista.

32. Rodrigues, S. B. M. (2015). *Rede de interações entre aves frugívoras e plantas em uma área de*
mata Atlântica no sudeste do Brasil. Universidade Federal de São Carlos.

33. Saavedra, F., Hensen, I., Beck, S. G., Böhning-Gaese, K., Lippok, D., Töpfer, T., & Schleuning,
M. (2014). Functional importance of avian seed dispersers changes in response to human-
induced forest edges in tropical seed-dispersal networks. *Oecologia*, *176*(3), 837–848.
https://doi.org/10.1007/s00442-014-3056-x

34. Sarmento, R., Alves-Costa, C. P., Ayub, A., & Mello, M. A. R. (2014). Partitioning of seed
dispersal services between birds and bats in a fragment of the Brazilian Atlantic Forest.
Zoologia (Curitiba), *31*(3), 245–255. https://doi.org/10.1590/S1984-46702014000300006

35. Schleuning, M., Blüthgen, N., Flörchinger, M., Braun, J., Schaefer, H. M., & Böhning-Gaese, K.
(2011). Specialization and interaction strength in a tropical plant–frugivore network differ
among forest strata. *Ecology*, *92*(1), 26–36. https://doi.org/10.1890/09-1842.1

36. Silva, R. F. de M. (2011). *Interações entre plantas e aves frugívoras no campus da Universidade*
Federal Rural do Rio de Janeiro. Universidade Federal Rural do Rio de Janeiro.

37. Snow, B. K., & Snow, D. (1988). *Birds and Berries*. T & AD Poyser.

38. Snow, B. K., & Snow, D. W. (1971). The feeding ecology of tanagers and honeycreepers in
trinidad. *The Auk*, *88*(2), 291–322. https://doi.org/10.2307/4083882

39. Stiebel, H., & Bairlein, F. (2008). Frugivorie mitteleuropäischer Vögel I: Nahrung und
Nahrungserwerb. *Vogelwarte*, *46*, 1–23.

40. Yang, S., Albert, R., & Carlo, T. A. (2013). Transience and constancy of interactions in a plant-
frugivore network. *Ecosphere*, 4(12), art147. https://doi.org/10.1890/ES13-00222.1

1152

Figure legends

Figure 1. A) Schematic example of a species-based interaction network between four

ornithochorous plant species (P_1 - P_4) and their frugivore assemblage with five animal species (A_1 -

A_5) (top). B) A zoom in on the individual-based network of plant species P_1 depicting the interactions

of its plant individuals (p_{11} - p_{15}) with five animal species, exemplifying the study focus of this paper.

C) Different plant individuals (p_{11} - p_{15}) interact with frugivore assemblages of variable diversity,

illustrating their individual interaction niches. Plant niches are exemplified by the five colored niche

utilization curves within the inset which indicate the relative interaction frequency with each of the

five animal species in the assemblage.

1163

Figure 2. Network metrics comparison across network scales (Ind = individual-based, n = 46; and Sp

= species-based networks, n = 59). Panel A shows observed network-level metrics and panel B

deviation of observed metric values from their corresponding null model, which represents a

network of the same size and relative abundances (generated using Patefield's algorithm). Points

represent the mean and thick and thin lines represent 0.6 and 0.9 confidence intervals respectively.

For each network metric, deviation from observed represents the difference between the observed

value and the null expectation.

1171

Figure 3. A schematic representation of two plant populations (top): the one on the left presents a

population composed of highly specialized individuals (low WIC; low WIC/TNW) and the one on the

right presents a population composed of highly generalized individuals (high WIC; high WIC/TNW).

Values of WIC/TNW closer to 1 represent populations with generalized individuals where plants use

most of the available interaction niche. On the other extreme, values closer to 0 indicate

populations with specialized individuals that use a smaller subset of the available interaction niche

(in this case plants do not tend to interact with the same frugivore species). Bottom panel: Values of

individual specialization (WIC/TNW) for all plant populations studied (n = 46 networks). The total

niche width (TNW) represents the interaction niche of the population, calculated as the Shannon

diversity of interactions with frugivore species at the population level, i.e., aggregating across

individuals. The within-individual component (WIC) is the average Shannon diversity of interactions

1183   with frugivores found within individual plants. Each point-interval in the graph represents an

1184   individual-based network (i.e., population), where the point indicates the mean WIC/TNW, and the

1185   thick and thin lines span the 60% and 90% credibility interval, respectively of each network

1186   posterior distribution. Colors represent the bioregions of the study site (see Table S1 for each

1187   network metadata).

1188

Figure 4. Relationship between the overall contribution to the total number of interactions by each

frugivore species (i.e., link weights in the individual-based networks) and the proportion of plant

individuals with which they interacted (i.e., animal degree). Each point represents a frugivore

species in their respective plant population (individual-based network). Point color indicates the

bioregion and point size is proportional to the frugivore's body mass relative to other frugivores

present in its population (z-score).

Figure 5. Principal Component Analysis for node-level metrics of plant individuals in all plant populations with the most common cluster attribution for each individual depicted in color. Each plant individual (points) is represented by the centroid of its 1000 node-metric values calculated with the full posterior distribution (Fig. S8). The panel below shows different plant populations as bars with a number code above (see Table S1). Each stacked bar is composed of 1000 thinner stacked bars that represent the cumulative proportion of individuals attributed to each cluster or interaction profile originated from the 1000 reanalysis using the posterior samples.