Comprehensive analysis of the microbial consortium in the culture of flagellate *Monocercomonoides exilis*

Alejandro Jiménez-González[1,2], Sebastian Cristian Treitli[1], Priscila Peña-Diaz[1], Anna Janovská[3], Vladimír Beneš[4], Petr Žáček[3], Vladimír Hampl[1]

Affiliations:

[1] Charles University, Faculty of Science, Department of Parasitology, BIOCEV, Vestec 252 50, Czech Republic

[2] Uppsala Biomedicine Centre, Department of Cell and Molecular Biology, Molecular Evolution program, Uppsala University, Uppsala, Sweden

[3] Charles University, Faculty of Science, BIOCEV, Vestec 252 50, Czech Republic

[4] Genome Biology Unit, European Molecular Biology Laboratory (EMBL), Heidelberg, Germany.

Corresponding author: Vladimir Hampl, vlada@natur.cuni.cz

Abstract

Monocercomonoides exilis is the only known amitochondriate eukaryote, making it an excellent model for studying the implications of mitochondrial reduction from a cellular and evolutionary point of view. Although *M. exilis* is an endobiotic heterotroph, it can grow *in vitro*, albeit with an uncharacterized and complex prokaryotic community. All attempts to grow *M. exilis* axenically have been unsuccessful. Here, we use metagenomic sequencing at different time points during culture growth to describe the composition and dynamics of this community. We assembled genomes of 24 from at least the 30 different bacterial species within. Based on DNA read abundances, *M. exilis* represents less than 1.5%, and the representation of dominant bacterial members changes over time. Genome-scale metabolic reconstruction, differential expression analysis and measurements of metabolites in the media showed that the community depends on organic carbon oxidation, fermentation, and hydrogen production without methanogenesis. This is consistent with the rapid decline of amino acids, nucleotides, glyceraldehyde, lactate, fatty acids, and alcohols in the media. The community depends on recycling the external supply of amino acids since it has a limited capacity to fix nitrogen gas and lacks ammonia oxidizers to close the nitrogen cycle. With the senescence of the culture, we observe changes in the expression of several metabolic pathways in *M. exilis*, particularly those adapting to starvation. We do not reveal any clear metabolic link to explain the dependence of *M. exilis* on prokaryotes.

Introduction

The gut microbiota refers to the community of microorganisms inhabiting the gastrointestinal system. Several studies have proved the influence of this community on human health (Fan and Pedersen, 2021; Jandhyala et al., 2015; Thursby and Juge, 2017), but these have mainly focused on the prokaryotic component since this represents the majority of cells and biomass. Yet, the animal gut microbiota also comprises viruses and eukaryotes, including unicellular protists. The effect of protists on the function and processes in the intestine, as well as their metabolic interactions with prokaryotes, are essentially unknown, except for some medically and veterinary important parasites, e.g.*, Entamoeba histolytica, Giardia intestinalis, Cryptosporidium* spp., *Spironucleus salmonicida, Blastocystis* spp. (Partida-Rodriguez et al., 2021; von Huth et al., 2021). However, these represent only a fraction of the intestinal protist diversity and their presence often leads to pathological states that significantly depart from the conditions found in the healthy gut.

Many intestinal protists can be successfully maintained in xenic cultures using a medium such as TYSGM-9 (Diamond, 1982; Hamann et al., 2016) and for many, this is the only feasible method of *in vitro* maintenance, given that the process of axenisation, i.e., removal of other unwanted organisms, has not been achieved. Xenic cultivation depends on establishing a stable community in culture, typically derived from the original sample, i.e., gut content or stool, which can be maintained for years by regular transfers. Whereas the community may be derived from the sampled environment, its eventual composition will most probably reduce as it adapts to the culture conditions (nutrient source, oxygen concentration, etc.) and thus will only distantly resemble the situation in the host gut. Still, the culture represents a simple and tractable model system from which information about these complex communities may be obtained. To our knowledge, such attempts are rare (Hamann et al., 2017).

In the present study, we aim to characterize the culture community derived from the

stool of *Chinchilla lanigera*. It contains a single eukaryote, the commensal bacterivorous

flagellate *Monocercomonoides exilis*, for which a quality genome draft is available and

functionally annotated (Karnkowska et al., 2019, 2016; Treitli et al., 2021). This species

belongs to Oxymonadida (Preaxostyla, Metamonada), known as inhabitants of animal

intestines (Hampl, 2017; Treitli et al., 2018). The whole group, including *M. exilis*, has lost

mitochondria, which is unique among eukaryotes and raises interest in their biochemistry and

physiology (Karnkowska et al., 2016; Novák et al., 2023). While other oxymonad species

harbor symbionts (Hampl, 2017) and specific roles and interactions have been partially

elucidated, e.g. *Streblomastix strix*, a protist that takes part in the complex community of the

hindgut of termites and is involved in cellulose digestion (Treitli et al., 2019), no symbiotic

interaction is known for *M. exilis*. Light and electron microscopy has demonstrated that

species of *Monocercomonoides* actively feed on prokaryotes (Treitli et al., 2018), hence

acting as predators within the intestinal microbiota. The effect and scale of this grazing on the

prokaryotic community and the level of prey selectivity are unknown.

We used a multi-omic approach to obtain a first insight into the processes of this

multipartite culture community over five days of growth. Namely, we determined the species

composition and changes in species abundances through time, predicted the metabolism

pathways, followed gene expression changes of the most abundant members as the culture

aged, and finally, measured the composition and changes of selected metabolites in the

media. With all these data in hand, we attempted to reconstruct the major community

biochemical processes and searched for potential interactions between its members.

Results

Growth of the culture and sampling of DNA, RNA and media

The development of the culture was followed for seven days in an experiment

performed in triplicates (Figure 1). An aliquot of the bacterized medium, i.e., medium in

which *Citrobacter portucalensis* grew overnight, representing day 0, and aliquots of days 2,

3, and 5 representing the exponential, stationary, and decline phases of *M. exilis* growth, were

chosen to sample metagenomic, metatranscriptomic, and metabolomic data.

Community composition

Using metagenomic reads generated from triplicates of each sampling point, we

assembled 24 metagenome-assembled genomes (MAGs) corresponding to the prokaryotic

members of the culture community (Figure 2, Supplementary Table 1). In addition to the

assembled MAGs, we identified six additional bacterial species based on the presence of 16S

rRNA genes. However, their read abundances were too low to assemble them into clean

MAGs (Supplementary Table 1). No viral or plasmid DNA was identified in the

metagenomic data. This suggests that the community consists of one eukaryote and at least

30 bacterial species, representing seven phyla, without any archaeal or viral component.

The quality of the 24 assembled MAGs varied, with the lowest completeness being

around 83% and the vast majority having genome completeness greater than 95%

(Supplementary Table 1). The estimated contamination levels are overall lower than 5%,

except for *Deltaproteobacteria* sp. MAG, whose contamination was estimated to be 11.8%

(Supplementary Table 1). This MAG was the least abundant during the whole experiment.

We calculated the relative read abundance of each bacterium and *M. exilis* for days 2, 3, and

5 of culture. While *M. exilis* represented only 0.25% to 1.23% of the reads, the seven most

abundant bacterial species comprised 80 – 90% of the reads, and the remaining bacterial

species never exceeded 1% of the reads per species (Figure 2 and Supplementary Table 1).

We observed a switch in the dominant groups throughout the culture growth. Not considering

Citrobacter portucalensis (Gammaproteobacteria), which was present in the medium before

inoculation of the community ("bacterization" of the medium), the most abundant bacterium

on day two was *Fusobacterium varium* (Fusobacteriota). On days three and five, the

abundance of this bacterium decreased, and representatives of the phylum Bacteroidota

dominated. *M. exilis* reached its peak of 1.23% on day 3 (Figure 2).

Monocercomonoides exilis **metabolic capacities**

Our analysis of *M. exilis* metabolism identified 933 enzymatic reactions, 31 transport

and 148 enzymatic pathways (Supplementary Figure 1). As previously published

(Karnkowska et al., 2019, 2016; Treitli et al., 2021), *M. exilis* is equipped with a limited

repertoire for energy metabolism relying on glycolysis to produce pyruvate, which is

converted into acetyl-coenzyme A and further to acetate and ethanol. Our analysis showed

that *M. exilis* can obtain glucose from the degradation of glycogen, maltose or starch but can

also digest chitin and (1,3)- α-D-glucans present in fungal cell walls and peptidoglycan

present in bacterial cell walls, which corroborates its bacterivorous feeding mode

(Supplementary Table 2). The biosynthesis of amino acids and nucleotides is limited and

dependent on conversion and salvage pathways from compounds acquired from the diet or

via transporters. However, *M. exilis* can synthesize dNTPs from NTPs using ribonucleoside-

triphosphate reductase (Karnkowska et al., 2019; Novák et al., 2023).

Bacterial community metabolic capacities

To better understand the metabolic potential of the 24 assembled bacterial genomes,

we reconstructed the metabolism of each MAG using EggNOG- mapper and Pathway-Tools.

Our results showed a higher number of pathways and reactions compared to *M. exilis*, with

considerable diversity in the number of enzymatic reactions, transporters and pathways (from

443 in *Citrobacter portucalensis* to 218 in *Peptostreptococcus russellii*) (Supplementary

Table 1). The metabolic maps of the seven most abundant MAGs are provided in

Supplementary Figures 2 – 8. The number of reactions and pathways identified in closely

related MAGs was very similar, even if the assembly quality differed between MAGs. This

suggests that our metabolic analyses were not heavily affected by the level of fragmentation

of the MAG assemblies.

We analyzed the roles of community members within the carbon, nitrogen, sulfur, and

iron cycles of the community. The results showed that the community likely depends on the

oxidation of organic carbon compounds (mainly amino acids and complex carbon

compounds), fermentation, and H_2 generation (Figure 3). The community does not perform

methanogenesis, and the produced H_2 is probably oxidized to water. Our analyses identified

Phascolarctobacterium sp. as the only community member able to fix N_2 to ammonium.

However, the abundance of this species is low (Figure 2 and Supplementary Table 1) so the

N_2 fixation most likely does not significantly contribute to the ammonium pool. Another

potential source of ammonium could be the Dissimilatory Nitrate Reduction to Ammonium

(DNRA) pathway (Kaviraj et al., 2024). Components of this pathway, periplasmic nitrate

reductase (NapAB) or the nitrate reductase A (NarGHI) and cytochrome c552 nitrite

reductase (NrfAH and NrfABCDEF), are present in multiple members of the community

(Figure 3, Supplementary Table 3, Supplementary Figures 2, 4 – 7). However, we failed to

identify the source of the nitrates and nitrites. These anions are not added to the cultivation

medium, and we did not identify enzymes producing nitrate or nitrite either by oxidation of

ammonia derived from the amino acid and nucleotide decay or by the biosynthesis of nitrite

from aspartate (Sugai et al., 2016). In conformity with the above, the expression of the

DNRA pathway in most bacteria, for which the transcriptomic data are available, is

extremely low and in the two species, which express it (*B. thetaiotaomicron* and

Parabacteroides sp.), it is down-regulated in later days (Supplementary Table 3,

Supplementary Figures 2, 4 – 7). We conclude that the community does not run a complete

nitrogen cycle but depends on the ammonium derived from amino acids and nucleotides

supplemented to the medium or originating from dead cells with a small addition through the

fixation of nitrogen gas.

In contrast to the carbon and nitrogen cycles, the sulfur cycle is complete. It is

maintained by *Desulphovibrio* sp, the only member able to reduce sulfate and oxidize sulfite.

We identified eight species that encode the sulfide:quinone oxidoreductase, suggesting that

they accumulate sulfur in globules within the cytoplasm (Arieli et al., 1994). The reduction of

iron is another important activity in the community, with 13 species involved in the process

(Figure 3). In contrast, iron oxidation activity was identified only for *Saezia sanguinis*.

(Figure 3).

The consumption and production of compounds by the community

Triplicates of the cell-free medium were sampled at each time point and their

composition was analyzed by non-targeted metabolomics. LC-MS/MS and GCxGC-MS

covered a broad range of structural classes of metabolites, detecting 270 distinctive

compounds altogether (Supplementary Table 4). Of these, 171 take part in the biological

reactions listed on the MetaCyc database (Caspi et al., 2014); hence, they were the only ones

used for the follow-up analyses (Supplementary Table 4).

On the day of inoculation (day 0), the medium was already affected by an overnight

growth of *C. portucalensis*. Yet it was relatively abundant in nucleotides and their derivates,

glyceraldehyde, lactate, fatty acids, alcohols and amino acids, except asparagine, aspartate,

cysteine, glutamine and serine, which were below the detection limit at this time point

(Supplementary Table 4). On day 2, a drastic evanesce of some compounds was registered,

specifically nucleotides and their derivates, most amino acids, glyceraldehyde, glyceric acid,

malate, citric acid, and citrulline, suggesting that the community had consumed them. On the

contrary, a spectrum of short-chain fatty acids and aromatic compounds putatively derived

from the degradation of amino acids, such as tyrosine, phenylalanine and tryptophan, was

observed in the media on this day (Supplementary Table 4). The measurements displayed

little changes between days 2, 3, and 5, suggesting that the composition of cell-free media

stabilized.

We integrated the synthesis and consumption of compounds into an *in silico*

metabolic capacity prediction, both for *M. exilis* and the bacterial community, using the

Metage2Metabo (m2m) pipeline (Belcour et al., 2020). In these predictions, the culture days

were compared pairwise, i.e., day 0 vs. day 2, day 2 vs. day 3 and day 3 vs. day 5. We then

classified compounds either as seeds if detected on the first day in pairwise comparison or as

targets if they were detected only or their concentration increased on the second sampling day

of that pair. This was performed using measured differences in media composition,

presence/absence and concentration changes of compounds between sampling days. These

results are summarized in Figure 4. From the fact that *M. exilis* represents less than 1.5% of

the community metagenomic reads, we infer that its metabolic power is low and the

prokaryotic component converts most of the metabolites. Nevertheless, in subsequent

analyses, we sought to stress out the metabolic capabilities of this protozoan, which was our

main point of interest. Between day 0 and day 2, 137 compounds were classified as seeds and

92 as targets, of which *M. exilis* can synthesize 63 by itself. However, the metabolic

capacities of *M. exilis* are not limited to the target compounds, as it can synthesize other

important compounds like pyruvate, amino acids, medium-chain fatty acids, and vitamins. An

additional 29 target compounds can be synthesized via metabolic interactions of *M. exilis* and

the bacterial community (Figure 4). Among these compounds, there are several amino acids.

In this pairwise comparison, m2m did not identify any bacteria as an essential member of the

community. However, two Bacteroidota bacteria were classified as key members (species

occurring in at least one of the minimal community models) (Figure 4). Between day 2 and

day 3, we identified 141 compounds as seeds and 72 as targets. According to the m2m

pipeline, *M. exilis* can synthesize 68 targets independently and 51 compounds through

metabolic interaction with the community (Figure 4). In this comparison, no bacterial species

were considered essential or key members of the community. The last comparison, day 3 vs.

day 5, showed a similar result. *M. exilis* could synthesize 54 of the 55 compounds identified

as targets using 140 compounds as seeds. The interaction between *M. exilis* and the bacterial

community allows the synthesis of 49 more compounds and none of the bacteria were

considered essential or key members (Figure 4).

Monocercomonoides exilis **differential expression analysis**

To investigate how *M. exilis* metabolism changes during the growth curve, we also

compared the expression of transcripts between days 2, 3 and 5, again pairwise. Most

significant changes in gene expression were observed between days 2 and 5, while

comparisons between consecutive days revealed less significant changes (i.e., day 3 vs. day 2

and day 5 vs. day 3). 2,236 genes were up-regulated and 2,431 down-regulated between days

2 and 5 (Supplementary Table 1), of which the majority, 1703 and 1,3 81 genes, respectively,

are annotated as hypothetical; hence their function could not be deciphered.

The changes in expression of all functionally annotated metabolic enzymes and

pathways are shown in Supplementary Figure 1 and those discussed below are shown in

Figures 5 and 6 in more detail. We observed an increase in the expression of importers of

sugars such as α-D-xylose and *sn*-glycerol-3-phosphate towards day 5. Connected to this,

there was an increment in the expression of enzymes responsible for the degradation of

starch, (1->4)-α-glucan, chitin and other polysaccharides (Figure 5 and Supplementary Table

2). Furthermore, the interconversion between glucose-1-P and glucose-6-P was up-regulated,

which may relate to the increased trehalose biosynthesis from glucose-6-P. Trehalose is often

used as carbohydrate storage and as a stress protector (Iturriaga et al., 2009). All three

pathways for trehalose synthesis were up-regulated on day 5, including the synthesis directly

from α-maltose, performed by both α- and β-amylases. The synthesis of glycogen, another

storage polysaccharide, was also up-regulated. The expression of most genes involved in

carbohydrate metabolism and glycolysis remained unchanged. However, some, such as

phosphoenolpyruvate carboxykinase (PEPCK), exhibit down-regulation, suggesting that

glycolysis is regulated to produce pyruvate via the up-regulated pyruvate kinase (PK). Later,

this pyruvate is converted to acetyl-CoA by pyruvate-ferredoxin oxidoreductase (PFOR) and

the up-regulated hydrogenases (HYD) re-oxidize ferredoxins, allowing the cycle to continue.

The expression of enzymes of extended glycolysis and ethanol fermentation, i.e., acetate

synthase (ACS) producing acetate and aldehyde and alcohol dehydrogenases (ALDH, Adh)

producing ethanol, decreases greatly on day 5 for all but one paralogue of ALDH

(Supplementary Table 5). ACS1 was consistently the most expressed of these enzymes in all

three time points and replicates. Overall, these expression data suggest that acetate and

hydrogen are the preferred metabolic end products at all three time points. In contrast to

glycolysis, the pentose phosphate pathway may be down-regulated due to the repression of

the ribulose-phosphate 3-epimerase (RPE) (Figure 5).

Genes responsible for the phosphorylation of (d)NDP, DNA and RNA

polymerization, and protein ubiquitination were up-regulated on day 5 (Supplementary

Figure 1). In contrast, nucleoside import, and the salvage pathway of adenosine were down-

regulated. Similarly, the mevalonate pathway for the synthesis of isoprenoids from

acetoacetyl-CoA was down-regulated (Figure 5 and Supplementary Figure 1). Most amino

acid importers displayed lower expression or showed no significant change in expression

(Figure 6). This relates to the fact that amino acid content in the media significantly

decreased from day 2. Asparagine, serine, and cysteine biosynthesis were down-regulated, as

was the ATP-producing arginine deaminase pathway (Figure 6).

Another system that showed significant changes during cultivation was the

endomembrane system. Among the proteins involved in the regulation of endomembrane

transport, some show significant changes in expression (Supplementary Table 6). The

proteins involved in the early secretory pathway show little change, but several proteins

involved in the late steps (Law et al., 2022; Shimizu and Uemura, 2022; Tanaka et al., 2017)

were significantly up-regulated on day 5 (Supplementary Table 6). These include proteins

involved in trans-Golgi-network (TGN) to plasma membrane trafficking (DSCR3A and

exocyst complex components Exo84, Sec5 and Sec15, DSCR), multivesicular body

formation and sorting (Vsp2, 4a-c, 23, 24a and 60b, charged multivesicular body protein 7),

plasma membrane uptake (AP-2a, TSET complex protein TPLATE subunit B),

endocytic/digestive pathway (Vps34bc, Vps41b) and endosomes to TGN transport (Vps51b,

TRAPP component Bet3 paralogue A, B, syntaxin 6D and 16). On the contrary, four Rab7

paralogues were consistently down-regulated, implying a decrease in lysosome biogenesis on

day 5 (Bucci et al., 2000).

At the end of the growth curve, we observed a decline in the number of *M. exilis* cells.

Our data showed that most of the enzymes involved in the apoptosis pathway (Teulière et al.,

2020) do not show significant expression change (Supplementary Table 7). This suggests that

the decline in cell density in later days is not due to programmed cell death but most likely

due to uncontrolled decay combined with slow growth caused by nutrient deficiencies.

Bacterial community differential expression analysis

Our data only allowed to perform differential expression analysis for the seven most

abundant bacteria: *B. fragilis*, *B. thetaiotaomicron*, *B. intestinalis*, *Parabacteroides* sp., *K.*

gyiorum, *Fusobacterium varium* and *C. portucalensis*. These species represent 80-90% of the

community DNA reads, a proxy measure of their abundance. These bacteria showed a

variable number of up- or down-regulated genes (Supplementary Table 1); for example, only

9% of *B*. intestinalis. transcripts changed in expression between day 2 vs. day 5, while in

Parabacteroides sp., it was 51%. Changes in the expression of metabolic enzymes and

pathways of these bacteria are indicated in Supplementary Figures 2 – 8.

We focused specifically on pathways absent in *C. portucalensis* but present in other

members of the community, as these may explain why we failed to grow *M. exilis* with only

C. portucalensis as food. First, *F. varium* and *Tissierella* sp. can ferment lysine to acetate,

butanoate and ammonia, while in *C. portucalensis*, this pathway is incomplete. Furthermore,

B. intestinalis, S. sanguinis, K. gyiorum, Phascolarctobacterium sp. and *Tissierella* sp.

encode genes for the synthesis of the essential polyamine spermidine and the phospholipid

phosphatidylcholine. While phosphatidylcholine is one of the major components of cell

membranes (Zhukov and Popov, 2023), spermidine is involved in an array of biological

processes, such as maintenance of membrane potential, control of intracellular pH, lipid

metabolism and regulation of cell growth, proliferation, and death (Michael, 2016). In both

cases, our metabolic analysis fails to identify a pathway that can synthesize these compounds

de novo in *M. exilis*, suggesting that the eukaryote may depend on the uptake of these

compounds from these prokaryotes.

Iron uptake is another relevant function for which no pathway is known in *M. exilis*.

Several bacteria encode for transporters of Fe^{2+}, the anaerobic Feo import system, the aerobic

Efe system and the Sit system (Braun and Hantke, 2013) (Supplementary Table 8). Our data

showed that in all bacteria with expression data, only the Sit system in *K. gyiorum* is up-

regulated between day 2 and day 5. In the rest of the bacteria, all the Fe^{+2} transporters are

down-regulated or not significantly differentially expressed between these days

(Supplementary Figures 2 – 8). Of importance is also another type of molecules used for iron

chelation in bacteria known as siderophores. These iron-chelating molecules are secreted into

the environment and their iron-bound forms are subsequently selectively imported (Kramer et

al., 2020). Our data showed that some, but not all, community members produce

siderophores, *C. portucalensis* and *S. sanguinis* synthesize enterobactin, while *F. varium*

synthesizes staphylopine, and *K. gyiorum* a periplasmic iron-binding protein. Interestingly, all

bacteria, but not *M. exilis*, encode transport systems for siderophores or transferrin. Of these,

we identified components of the Ton uptake system (TonB, ExbB and ExbD) (Braun and

Hantke, 2013) in most gram negative bacteria except for ExbB absent from *Morganella*

morganii (Supplementary Table 8). Consistently, we identified several TonB-dependent

receptors and transporters, namely the enterobactin transporters FepA and FbpABC and the

transferrin receptor TbpA in many members of the community. The presence of these

proteins and protein complexes makes the transport of siderophores or heme groups across

the outer membrane possible (Braun and Hantke, 2013). Our data showed that the Ton uptake

system and the receptor TbpA are up-regulated in *F. varium, K. gyiorum* and members of the

Bacteroidota in later days of the culture (Supplementary Figures 3 – 5, 7,8). The described

pattern of presence and absence suggests that the bacterial community contains only a few

species able to synthesize siderophores and all the other members are "cheaters" that uptake

siderophores produced by others (Butaite et al., 2017; Leventhal et al., 2019). From this point

of view, the siderophores' producers are key players in the prokaryotic community but do not

facilitate the uptake of iron into *M. exilis*. Once iron is imported into the cell, it cannot be

kept free in the cytoplasm. One solution would be its coordination within different kinds of

porphyrins (Bryant et al., 2020; Dailey et al., 2017; Layer, 2021). We observed two main

porphyrins, siroheme and heme *b*. Heme *b* can be synthesized aerobically or in an oxygen-

independent manner, while siroheme is synthesized only independently of oxygen (Dailey et

al., 2017; Layer, 2021). Our data showed that *C. portucalensis* can synthesize both

compounds. Still, under anaerobic conditions, the synthesis of heme *b* is down-regulated,

while that of siroheme is up-regulated on day 5 (Supplementary Figure 2). The opposite

situation was observed in *Bacteroides fragilis,* where oxygen-independent synthesis of heme

b was up-regulated, yet siroheme synthesis underwent down-regulation (Supplementary

Figure 4). Some bacteria in the community can synthesize heme *b* from siroheme. However,

we lacked transcriptome data for their assessment due to their low abundance.

341   **Discussion**

The only study of chinchilla gut microbiota composition (O' Donnell et al., 2017)

reports Betaproteobacteria as the dominant phylum and *Parabacteroides* and *Barnesiella* as

the dominant genera. Our study of an *M. exilis* culture derived from the gut of chinchilla

identified 30 bacterial species, of which *Fusobacterium varium* was the most abundant at the

beginning of the experiment, steadily replaced by the Bacteroidetes group, represented by

Bacteroides fragilis, *Parabacteroides* sp. and *B. thetaiotaomicron* on day 5. *Parabacteroides*

sp. was the second most abundant bacteria on day 5, yet we failed to detect any *Barnesiella*

species. Only two Betaproteobacteria species were identified in our study, but they are not

among the most abundant species. At the level of bacterial phyla, Bacteroidota and Bacillota

are the most abundant in our culture, which, on a broad scale, resemble the composition of

the gut microbiota of Guinea pigs (Crowley et al., 2017). These comparisons indicate similar

compositions of the prokaryotic community between the gut of caviomorph rodents and our

M. exilis culture, yet the culture community has changed due to long-term cultivation.

Given that *M. exilis* represents around 1% of the read percentage, the biochemical

processes in the community are driven by prokaryotes. Of the three main cycles, only the

sulfur cycle seems to be complete and contains reactions converting all basic inorganic sulfur

compounds (Figure 3). Our analyses infer that the community dominantly oxidizes organic

carbon substrates, performs fermentation and generates hydrogen, which is further oxidized

to water (Figure 3). Since the ability to fix carbon from CO_2 is present only in *Eubacterium*

maltosivorans, the least abundant species, the community depends on the supply of organic

carbon. This is consistent with our observation that organic carbon sources (glyceraldehyde,

lactate, fatty acids, alcohols and most amino acids) present on day 0 are quickly consumed.

The community does not run a complete nitrogen cycle. While there is a small level of

N_2 fixation due to the contribution of *Phascolarctobacterium* sp., this bacterium is among the

less abundant members of the community, and most likely, its N_2 fixation does not cover the

demands for ammonium. Another pathway for ammonium synthesis, the DNRA pathway

using the periplasmic nitrate reductase or the nitrate reductase A, and the membrane-bound

nitrate reductase nrf, is encoded by some bacterial species (Kaviraj et al., 2024). This

pathway uses nitrate as electron acceptor and allows building membrane potential (Cabello et

al., 2019). However, in all the bacteria with transcriptomic data, the DNRA is not expressed

or downregulated in later days. The nitrogen sources for building a biomass thus are amino

acids and nucleobases, which are components of the growth medium and are measurable on

the day 0 with a small supplement of ammonia synthesized by *Phascolarctobacterium* sp.

The metabolomic analyses of the soluble media clearly showed that on day 2, when

the population of *M. exilis* only began to grow, essentially all soluble carbon sources,

nucleotides and most amino acids had been already consumed (Supplementary Table 4). The

fact that *M. exilis* maintains active growth after day 2 strongly suggests that the protist uses

bacterial prey as the major source of carbon, nitrogen, and energy to build its biomass. A

putative exception could be arginine, which was detected in later days and may serve as a

potential substrate for energy metabolism (Schofield et al., 1992; Yarlett et al., 1996).

Grazing on bacteria has been previously documented by electron microscopy (Treitli et al.,

2018) and the fact that the protist encodes lysozymes amongst a broad array of glycosyl

hydrolases (Supplementary Table 2) also provides molecular tools for feeding on bacteria.

There is no consistent pattern that would indicate up- or down-regulation of lysozymes as the

culture ages. The protist, however, displays a significant up-regulation of several genes

involved in the late pathways of the endomembrane system on day 5. At the same time

expression of many genes encoding for proteins regulating these processes remained

unchanged and, notably, four paralogues of Rab7 small GTPase, known for its role in

lysosome activation (Bucci et al., 2000), were significantly down-regulated on day 5

(Supplementary Table 6). We interpret this pattern as an indication of up-regulation of

exocytosis and secretion rather than bacterial feeding, which is also consistent with the

decrease in *M. exilis* cell counts at this stage of the culture.

Most *M. exilis* genes that exhibit differential expression completely lack functional

annotation, providing us with a rather large transcriptomic dark matter consisting of approx.

three thousand genes that may convey some functional and adaptive value to the protist, but

we know absolutely nothing about it. Among the known metabolic genes, we observed an up-

regulation of the importers of sugars, α-D-xylose and *sn*-glycerol-3-phosphate, which is

likely the reaction of the cell to the absence of these substrates. Another functional category

of genes up-regulated on day 5 is the enzymes responsible for starch degradation and

synthesis of trehalose and glycogen, indicating the onset of starvation. The degradation of

nucleosides and nucleotides and the biosynthesis of 5-phospho-α-D-ribose 1-diphosphate

(PRPP) are up-regulated, suggesting an increment in the storage of nucleotides and phosphate

donors. On the other hand, glycolysis, extended glycolysis, fermentation, pentose phosphate

pathway, amino acids biosynthesis, arginine deaminase pathway and mevalonate pathway

involved in isoprenoid biosynthesis are generally down-regulated on day 5, with few

exceptions. The same holds for the synthesis of very long and ultra-long fatty acids and the

aminoacyl-tRNA synthetases, indicating the decline of proteosynthesis.

Although we hoped to find some metabolic dependencies between *M. exilis* and

prokaryotes, potentially explaining the inability to grow the flagellate axenically using

nutrients dissolved in the medium, we did not reveal any clear link. This obviously does not

mean that some have remained unobserved or unnoticed, as the amount and complexity of the

data are significant. The *in silico* metabolic capacity prediction by Metage2Metabo (Figure 4)

has shown that the compounds detected in the media at the time points can be synthesized by

the reactions and pathways found in the community. Importantly, none of the community

members has been found to be essential at any time point, though some may provide key

compounds for others, e.g., siderophores. This suggests that the community is based on broad

redundant foundations, which certainly contributes to its long-term stability. However, we

infer that carbon and nitrogen could be limited factors that can disturb this tendency.

Material and methods

Cell culturing

700 mL of TYSGM-9 medium in 2x 1 L glass bottles were inoculated with 500 μL of

Citrobacter portucalensis culture before the inoculation of the whole community. This strain

of *C. portucalensis* was previously isolated from the original culture during the isolation of

M. exilis. After inoculation, the flasks were closed and allowed to grow standing at 37°C. The

next day, the two flasks were merged in a 2 L bottle and mixed thoroughly. 45 mL of culture

were removed and used to isolate DNA, RNA and metabolomic analyses of day 0. The rest of

the medium was inoculated with approx. 125 mL of *M. exilis* culture (2.5×10^5 cells.mL^{-1}).

After inoculation, the media was thoroughly mixed and divided into 50-mL tubes containing

45 mL of media, closed, and incubated at 37°C. Each day, three tubes were removed, one for

each replicate. From each tube, 20 mL of culture were used for DNA isolation, 20 mL for

RNA isolation and 5 mL for metabolomics. Before each nucleic acid isolation, *M. exilis* cells

were counted using a Neubauer cytometer.

DNA isolation

All DNA isolations were performed using the Dneasy Blood and Tissue Kit (Qiagen).

Each culture was centrifuged at 1500 x*rcf* for 10 min at 4°C. The supernatant was separated

into a fresh 50 mL tube and centrifuged again at 6000 x*rcf* for 10 min at 4°C. The pellet from

the first centrifugation was resuspended in 200 µL of PBS and DNA was isolated using the

cultured cells protocol of the Dneasy Blood and Tissue Kit (Qiagen). The second pellet,

obtained after centrifugation at 6000 x*rcf*, was resuspended in 200 µL of PBS and DNA was

isolated using the gram-positive protocol of the Dneasy Blood and Tissue Kit (Qiagen). For

each isolation, the DNA was eluted in 100 µL of elution buffer. The DNA from the two

pellets was merged into one DNA sample. The quality of the DNA was estimated using

Nanodrop and the concentration was measured using the QuantiFluor ONE dsDNA System

(Promega).

RNA isolation

Each culture was centrifuged at 1500 x*rcf* for 10 min at 4°C. The supernatant was

separated into a fresh 50 mL tube and centrifuged again at 6000 x*rcf* for 10 min at 4°C. Both

pellets were resuspended together in 1 mL Tri-Reagent (Sigma-Aldrich) and the total RNA

was isolated according to the manufacturer's procedure. The obtained RNA pellet was

resuspended in 60 µL of RNA-grade H_2O and heated at 37°C for 10 min to allow full

resuspension.

The total RNA was further DNAse-treated. For this purpose, we added 6 µL of

rDNAse buffer and 0.6 µL rDNAse (Macherey-Nagel) per sample and incubated at 37°C for

10 min. Samples were subsequently re-isolated with Tri-Reagent (Sigma-Aldrich) as before

and finally resuspended in 30 µL of RNA-grade H_2O. RNA concentration was measured

using Nanodrop.

Identification of compounds within the growth medium

For metabolomics, cells were pelleted as described for DNA and RNA isolation (see

above). The supernatant obtained in the second centrifugation was filtered (Whatman

polycarbonate filter, 3 µm pore size)-sterilized and stored at -80^{o}C. Metabolites released to

the media were determined using a non-targeted metabolomics approach. To capture both

volatiles and non-volatiles, a combination of two techniques was employed – Liquid

Chromatography analysis in connection with Orbitrap Fusion mass spectrometer (LC-

MS/MS; Orbitrap Fusion, Q-OT-qIT, Thermo Scientific) and two-dimensional

comprehensive gas chromatography in connection with mass spectrometer (GCxGCTOF-MS;

Pegasus 4D, Leco Corp.).

For LC-MS/MS, media samples harvested in triplicates were precipitated with

acetonitrile in a ratio of 4:1 (ACN:media) and centrifugated for 20 min at 16000 xrcf at 5☐C.

100 µL of each supernatant was collected, fully evaporated and dissolved in 50 µL 10% (v/v)

ACN. 15 µL were injected into a ProntoSIL column, 150 x 3.0 mm, 3 µm (Bischoff

Chromatography). Compounds were eluted under the following conditions: 0 - 3 min 100%

A, 3 - 18 min linear gradient to 100% B, 18 - 20 min 100% B using mobile phases. A: 2%

(v/v) ACN, 10 mM $HCOONH_4$, 0.1% (v/v) FA; B: 99% (v/v) ACN, 0.1% (v/v) FA with a

flow of 0.4 mL.min^{-1}. The separated compounds were ionized in an electrospray ion source in

positive polarity mode. Master scans of precursors from 80 to 1000 m/z were performed in

Orbitrap at 120 K resolution with an intensity threshold filter 2.0 x 10^4. Tandem MS was

performed by isolation in the quadrupole, HCD fragmentation with a stepped collision energy

of 15, 30, and 45% and an isolation window of 1.6 m/z. The MS2 ions were analysed and

detected in Orbitrap with a set resolution of 30k and a max injection time of 300 ms. The

dynamic exclusion duration occurred every 20 s with 10 ppm tolerance around the selected

precursor. The data processing, including chromatographic peaks alignment, was performed

using Compound Discoverer 3.3 (Thermo Scientific). Quality control samples were employed

for the data correction. The mzCloud libraries were used as a tool for a fragmentation spectra

comparison and the compound structure assignment.

Volatiles were analyzed using GCxGC-MS and a headspace solid phase

microextraction (HS-SPME) on fiber (DVB/CAR/PDMS_grey; Supelco, USA). A volume of

1 mL of media samples was placed in a 20 mL glass vial and 10 µL of an internal standard

(2,2,2-trifluoro ethanol, Fluka, 0.11 mg.mL^{-1}) was added. Samples were incubated for 10 min

at 40^{o}C before extraction. The extraction was carried out for 5 minutes. The volatiles were

analyzed using a combination of polar and mid-polar separation columns for the separation

(primary column: Stabilwax-DA (30 m x 0.25 mm, Restek, USA); secondary column BPX-

50 (1.38 m x 0.1 mm, SGE, Australia)). Other parameters were set as follows: inlet

temperature 220^{o}C, split 5 modes, constant He flow 1 mL.min^{-1}, modulation time 3 s (hot

pulse 0.7 s), modulation temperature offset with respect to the secondary oven 20^{o}C. The

temperature program applied on the primary oven was 30^{o}C (hold 1.5 min), then increased to

110^{o}C (8^{o}C.min^{-1}), followed by an increase to 250^{o}C (25^{o}C.min^{-1}) to 250^{o}C (hold 5 min). The

temperature offset applied on the secondary column was +5^{o}C. Transferline temperature was

held at 280^{o}C. The mass spectrometer was equipped with an Electron Ionization ion source

(energy of 70eV was applied),and Time-of-Flight analyser enabling a united mass resolution.

The scanned mass range was 29 – 400 *m/z*. The ion source chamber was held at 250^{o}C.

To extend the range of compounds analyzable by gas chromatography to polar non-

volatiles, the media samples underwent two consecutive derivatization procedures –

oximation (25 mg.mL^{-1} of methoxyamine hydrochloride in anhydrous pyridine

hydrochloride, Sigma-Aldrich) and silylation. Two different agents were used for silylation:

N,O-Bis(trimethylsilyl)trifluoroacetamide with trimethylchlorosilane (BSTFA + 1%

(v/v)TMSC) and N-tert-Butyldimethylsilyl-N-methyl trifluoroacetamide (MTBSTFA), both

from Sigma-Aldrich.

For silylation using BSTFA + 1% (v/v) TMSC, 50 µL were transferred to a 2 mL

glass vial and 2 µL of internal standard were added (Adonitol, 424 µL.mL^{-1}). The liquid was

dried in a vacuum concentrator and reconstituted back by adding 30 µL of anhydrous

pyridine and 30 µL of an oximation agent. The samples were then incubated in a

Thermomixer (Eppendorf) at 40ºC for 2 hours at constant shaking (1500 rpm), after which 40

µL of neat solution of BSTFA + 1% (v/v) TMSC mixture were added to the samples and

were further incubated at 70^{O}C for 30 min at constant shaking (1500 rpm). Before analysis,

400 µL of hexane was added to the sample. For silylation using MTBSTFA, the

derivatization process is essentially the same as the derivatization using BSTFA + TMSC

except that 30 µL of media sample are used, plus 5 µL of nor-Valine (550 µL.mL^{-1}) as an

internal standard and 20 µL of oximation agent.

Metabolites were analyzed as oximated trimethylsilylated or tert-butyl silylated

derivatives. A combination of nonpolar and mid-polar separation columns was used for the

separation; a primary Rxi-5SIL MS column (30 m x 0.25 mm, Restek, USA) and a secondary

BPX-50 column (1.39 m x 0.1 mm, SGE, Australia). Other parameters were set as follows:

inlet temperature 290^{o}C, splitless mode, constant He flow 1 mL.min^{-1}, modulation time 3s

(hot pulse 0.7s), modulation temperature offset with respect to the secondary oven 15^{o}C. The

temperature program applied to the primary oven was 50^{o}C (hold 1 min), then increased

(8^{o}C.min^{-1}) to 320^{o}C (hold 2 min). The temperature offset applied on the secondary column

was +5^{o}C. Transferline temperature was held on 280^{o}C. The scanned mass range was 85 –

800 *m/z*. The ion source chamber was held at 280^{o}C. Data were processed in ChromaTOF

v4.5 software. Detected analytes were quantified relatively after normalization to the internal

standards. Metabolites detected by all the methods used were identified by comparison of

their mass spectra with those available in the main NIST mass library, Fiehn´s mass library of

silylated compounds, and in-house-built libraries. The retention index was determined using

linear hydrocarbons. In the cases where the same compounds were detected in both

derivatizations, the highest signal was taken to the final table.

Library preparation and sequencing

For the DNA samples, sequencing libraries were prepared from 1 µg of gDNA using

the Illumina TruSeq DNA PCR free (Illumina) library preparation kit according to the

manufacturer's protocol. The prepared libraries were sequenced on an Illumina HiSeq 4000

with 2x150 bp reads.

For RNA, 1 µg of total RNA per sample was used for library preparation. Ribosomal

RNA depletion was performed using the NEBNext rRNA Depletion Kit for Bacteria (New

England Biolabs), spiked with a custom oligo pool. The oligo pool was designed based on

bacterial and eukaryotic ribosomal sequences that were identified from previous sequencing

projects of *M. exilis* (Karnkowska et al., 2016; Treitli et al., 2021). After rRNA depletion,

libraries were prepared using the NEBNext® Ultra™ Directional RNA Library Prep Kit

(New England Biolabs) according to the manufacturer's protocol. The prepared libraries were

sequenced using Illumina NextSeq 500 with 2x75bp reads.

Metagenome assembly and binning

The metagenomics raw reads were quality-checked with FASTQC v0.11.5 and

trimmed using Trimmonatic v0.39 (Bolger et al., 2014) (ILLUMINACLIP:TruSeq3-PE-

2.fa:2:30:10, LEADING:20, TRAILING:20, SLIDINGWINDOW:5:20 MINLEN:50). To

reduce complexity during assembly, the eukaryotic reads were removed by mapping the reads

against the *M. exilis* genome (Treitli et al., 2021) using BBMap (minidentity=0.98,

idfilter=0.98) included in BBTools v38.90 (https://sourceforge.net/projects/bbmap/) and

taking only unmapped reads.

The filtered reads were assembled independently for each sample using metaSPAdes

v3.14.0 (Nurk et al., 2017) with k-mers 21,33,55,77,99,127. A reiterative process was used

for binning metagenome-assembled genomes (MAGs) within the samples. First, contigs

above 2kbp were binned using MaxBin v2.2.7 (Wu et al., 2016) (minimum probability 0.9,

marker set 107) and tetraESOM (Dick et al., 2009). Then, bins were manually checked based

on the markers identified by MaxBin using Blastp against nr database (April 2021). For each

potential MAG, rRNA genes were identified using Prokka v1.14.6 automatic annotation

pipeline (Seemann, 2014) and used as controls together with the marker genes. Lastly, reads

corresponding to trustful bins (markers and rRNA pointing to the same species) were

removed from the data set by mapping the reads back to the identified bins using BBSplit

(minidentity=0.98, idfilter=0.98) included in BBTools v38.90. This process was iterated and

stopped when no new clean MAG was created. In total, 24 prokaryotic MAGs were

assembled. The contribution of each prokaryotic MAG and *M. exilis* to the whole community

per sample was calculated as the percentage of reads mapped to each genome assembly.

To identify possible viral particles and plasmids in the culture, the filtered reads were

assembled independently for each sample using metaviralSPAdes (Antipov et al., 2020) and

metaplasmidSPAdes (Antipov et al., 2019), respectively, with k-mers 21,33,55,77,99,127.

Candidates contigs were verified using viralVerify tool (github.com/ablab/viralVerify.git)

and Prokka v1.14.6.

Identification of prokaryotic MAGs

The taxonomic classification of the 24 prokaryotic MAGs was computed using

classify_wf, included in GTDB-Tk (Chaumeil et al., 2022; Eddy, 2011; Harris et al., 2020;

Hyatt et al., 2010; Matsen et al., 2010; Ondov et al., 2016; Parks et al., 2020; Price et al.,

2010; Shaw and Yu, 2023; Sukumaran and Holder, 2010) with standard parameters. The

completeness and contamination of every MAG were estimated based on the

presence/absence and number of copies of the 107 marker genes used by MaxBin (see

above).

Annotation and metabolic capacities of prokaryotic MAGs

The annotation of the 24 prokaryotic MAGs was performed using Prokka v1.14.6.

The functional annotation of each MAG was made using EggNOG-mapper v1.0.3-35 on

emapper DB 2.0 (Cantalapiedra et al., 2021; Huerta-Cepas et al., 2019) (standard parameter

except minimum % of query coverage: 50 and minimum % of subject coverage: 50).

EggNOG-mapper predictions and Prokka annotations were combined using emapper2gbk,

included in the Metage2Metabo pipeline (Belcour et al., 2020), to create a compatible input

for Pathway-Tools v24.5 (Karp et al., 2020).

A metabolic database was created for each MAG using Pathologic (Karp et al., 2011),

included in Pathway-Tools (standard parameters except name matching that was turned off).

All databases were manually curated using the Assign Probable Enzymes, Transport

Inference Parser and Rescore Pathways tools included in Pathologic. Predicted pathways

were manually checked to remove highly incomplete pathways.

Identification of metabolic capacities of *Monocercomonoides exilis*

The genome of *M. exilis* was downloaded from NCBI. The metabolic capacities of

this species were predicted using Pathologic, included in Pathway-Tools, with standard

parameters. The resulting database was manually curated using the Assign Probable

Enzymes, Transport Inference Parser and Rescore Pathways tools included in Pathologic.

Information from previous curations (Karnkowska et al., 2019, 2016; Treitli et al., 2021) and

EggNOG-mapper was included at this step.

Carbohydrate-metabolizing enzymes were identified by computing searches against

CAZy databases using METABOLIC-G with default parameters (Zhou et al., 2022).

Transcriptome processing and identification of differentially expressed genes

Meta-transcriptomic raw reads were quality-checked with FASTQC v0.11.5 and

trimmed using Trimmonatic v0.39 (ILLUMINACLIP:TruSeq3-PE-2.fa:2:30:15,

LEADING:15, TRAILING:15, SLIDINGWINDOW:4:15 MINLEN:50), resulting in 47

million read pairs per sample on average (except for sample D5R3 where the number of read

pairs was 77 million). Reads were mapped and the expression of transcripts of prokaryotes

and *M. exilis* was quantified using Salmon (Patro et al., 2017) with default parameters.

Due to the low mapping rate and a low number of reads, the differential expression

was assessed only for *M. exilis* and the seven most abundant prokaryotic genomes:

Bacteroides fragilis, Bacteroides thetaiotaomicron, Bacteroides intestinalis, *Parabacteroides*

sp*., Kerstersia gyiorum, Fusobacterium varium* and *Citrobacter portucalensis*. Raw reads

were re-mapped onto the eight selected genomes using Bowtie2 (Langmead and Salzberg,

2012). Reads mapping to protein-coding sequences were counted with the FeatureCounts

program of the subread package v2.0.3 (Liao et al., 2014). Read counts were normalized

using the DESeq2 package v1.32.0 (Love et al., 2014) in R v4.1.1 (R Core Team, 2021).

Normalisation was based on taxon-specific scaling (Christel et al., 2018; Klingenberg and

Meinicke, 2017).

Identification of the major biochemical cycles in the community

625   The metabolic capacities of the prokaryotic community were also analyzed using

626   METABOLIC-C (Zhou et al., 2022) with default parameters, except tax genus, to understand

627   their roles within the community's major biochemical cycling. We manually verified the

628   results concerning the carbon, nitrogen, sulfur, and iron cycle by combining the results from

629   METABOLIC-C and EggNOG-mapper from previous analyses.

Identification of the metabolic relationships between *M. exilis* and the prokaryotic

community

All identified reactions in *M. exilis* and the 24 prokaryotic MAGs metabolic database

were exported as SBML format from Pathway-Tools. The Metacom tool implemented in the

metage2metabo (m2m) pipeline was used to identify metabolic interactions between *M. exilis*

and the rest of the community. In this analysis, the sampling days were compared pairwise,

with *M. exilis* defined as the host. In each pairwise comparison, compounds that could be

identified in the first sampling point were defined as seeds. Meanwhile, those whose

concentration increased in the second sampling point were defined as targets.

640 Acknowledgment

We thank Professor Joel Dacks for his advice on the analysis of the differentially

expressed protein related to the endomembrane system. This project has received funding

from the European Research Council (ERC) under the European Union's Horizon 2020

research and innovation program (grant agreement No. 771592 to VH) and the Centre for

Research of Pathogenicity and Virulence of Parasites (registration no.

CZ.02.1.01/0.0/0.0/16_019/0000759). The metagenome assemblies were computed using the

resources provided by the e-INFRA CZ project (ID:90254), supported by the Ministry of

Education, Youth and Sports of the Czech Republic.

650 Competing Interest

651   The authors declare that they have no conflict of interest.

653 Data Availability Statement

654   DNA reads, RNA reads and the complete 24 MAGs are available under a BioProject

655 with accession number PRJEB70623. MAGs are available with accession numbers from

656 ERS17699982 to ERS17699999.

658 References

Antipov D, Raiko M, Lapidus A, Pevzner PA. 2020. Metaviral SPAdes: Assembly of viruses

from metagenomic data. *Bioinformatics* **36**:4126–4129.

doi:10.1093/bioinformatics/btaa490

Antipov D, Raiko M, Lapidus A, Pevzner PA. 2019. Plasmid detection and assembly in

genomic and metagenomic data sets. *Genome Res* **29**:961–968.

doi:10.1101/gr.241299.118

Arieli B, Shahak Y, Taglicht D, Hauska G, Padan E. 1994. Purification and characterization

of sulfide-quinone reductase, a novel enzyme driving anoxygenic photosynthesis in

Oscillatoria limnetica. *J Biol Chem* **269**:5705–5711. doi:10.1016/S0021-9258(17)37518-

X

Belcour A, Frioux C, Aite M, Bretaudeau A, Hildebrand F, Siegel A. 2020. Metage2metabo,

microbiota-scale metabolic complementarity for the identi1cation of key species. *Elife*

9:1–38. doi:10.7554/eLife.61968

Bolger AM, Lohse M, Usadel B. 2014. Trimmomatic: a flexible trimmer for Illumina

sequence data. *Bioinformatics* **30**:2114–2120. doi:10.1093/bioinformatics/btu170

Braun V, Hantke K. 2013. The Tricky Ways Bacteria Cope with Iron Limitation In:

Chakraborty R, Braun V, Hantke K, Cornelis P, editors. Iron Uptake in Bacteria with

Emphasis on E. Coli and Pseudomonas. Dordrecht: Springer Netherlands. pp. 31–66.

doi:10.1007/978-94-007-6088-2_2

Bryant DA, Hunter CN, Warren MJ. 2020. Biosynthesis of the modified tetrapyrroles—the

pigments of life. *J Biol Chem* **295**:6888–6925. doi:10.1074/jbc.REV120.006194

Bucci C, Thomsen P, Nicoziani P, McCarthy J, Van Deurs B. 2000. Rab7: A key to lysosome

biogenesis. *Mol Biol Cell* **11**:467–480. doi:10.1091/mbc.11.2.467

Butaite E, Baumgartner M, Wyder S, Kümmerli R. 2017. Siderophore cheating and cheating

resistance shape competition for iron in soil and freshwater Pseudomonas communities.

Nat Commun **8**. doi:10.1038/s41467-017-00509-4

Cabello P, Luque-Almagro VM, Roldán MD, Moreno-Vivián C. 2019. Nitrogen cycle.

Encycl Microbiol 301–310. doi:10.1016/B978-0-12-809633-8.20706-1

Cantalapiedra CP, Hernández-Plaza A, Letunic I, Bork P, Huerta-Cepas J. 2021. eggNOG-

mapper v2: Functional Annotation, Orthology Assignments, and Domain Prediction at

the Metagenomic Scale. *Mol Biol Evol* **38**:5825–5829. doi:10.1093/molbev/msab293

Caspi R, Altman T, Billington R, Dreher K, Foerster H, Fulcher CA, Holland TA, Keseler

IM, Kothari A, Kubo A, Krummenacker M, Latendresse M, Mueller LA, Ong Q, Paley

S, Subhraveti P, Weaver DS, Weerasinghe D, Zhang P, Karp PD. 2014. The MetaCyc

database of metabolic pathways and enzymes and the BioCyc collection of

Pathway/Genome Databases. *Nucleic Acids Res* **42**:459–471. doi:10.1093/nar/gkt1103

Chaumeil PA, Mussig AJ, Hugenholtz P, Parks DH. 2022. GTDB-Tk v2: memory friendly

classification with the genome taxonomy database. *Bioinformatics* **38**:5315–5316.

doi:10.1093/bioinformatics/btac672

Christel S, Herold M, Bellenberg S, Buetti-Dinh A, Hajjami M El, Pivkin I V., Sand W,

Wilmes P, Poetsch A, Vera M, Dopson M. 2018. Weak iron oxidation by Sulfobacillus

thermosulfidooxidans maintains a favorable redox potential for chalcopyrite

bioleaching. *Front Microbiol* **9**:1–12. doi:10.3389/fmicb.2018.03059

Crowley EJ, King JM, Wilkinson T, Worgan HJ, Huson KM, Rose MT, McEwan NR. 2017.

Comparison of the microbial population in rabbits and guinea pigs by next generation

sequencing. *PLoS One* **12**. doi:10.1371/journal.pone.0165779

Dailey HA, Dailey TA, Gerdes S, Jahn D, Jahn M, O'Brian MR, Warren MJ. 2017.

Prokaryotic Heme Biosynthesis: Multiple Pathways to a Common Essential Product.

Microbiol Mol Biol Rev **81**:1–62. doi:10.1128/mmbr.00048-16

Diamond LS. 1982. A new liquid medium for xenic cultivation of *Entamoeba histolytica* and

other lumen-dwelling protozoa. *J Parasitol* **68**:958–9.

Dick GJ, Andersson AF, Baker BJ, Simmons SL, Thomas BC, Yelton AP, Banfield JF. 2009.

Community-wide analysis of microbial genome sequence signatures. *Genome Biol* **10**.

doi:10.1186/gb-2009-10-8-r85

Eddy SR. 2011. Accelerated profile HMM searches. *PLoS Comput Biol* **7**.

doi:10.1371/journal.pcbi.1002195

Fan Y, Pedersen O. 2021. Gut microbiota in human metabolic health and disease. *Nat Rev*

Microbiol **19**:55–71. doi:10.1038/s41579-020-0433-9

Hamann E, Gruber-Vodicka H, Kleiner M, Tegetmeyer HE, Riedel D, Littmann S, Chen J,

Milucka J, Viehweger B, Becker KW, Dong X, Stairs CW, Hinrichs KU, Brown MW,

Roger AJ, Strous M. 2016. Environmental Breviatea harbour mutualistic *Arcobacter*

epibionts. *Nature* **534**:254–258. doi:10.1038/nature18297

Hamann E, Tegetmeyer HE, Riedel Di, Littmann S, Ahmerkamp S, Chen J, Hach PF, Strous

M. 2017. Syntrophic linkage between predatory *Carpediemonas* and specific

prokaryotic populations. *ISME J* **11**:1205–1217. doi:10.1038/ismej.2016.197

Hampl V. 2017. Preaxostyla In: Archibald JM, Simpson AGB, Slamovits CH, editors.

Handbook of the Protists. Cham: Springer International Publishing. pp. 1139–1174.

doi:10.1007/978-3-319-28149-0_8

Harris CR, Millman KJ, van der Walt SJ, Gommers R, Virtanen P, Cournapeau D, Wieser E,

Taylor J, Berg S, Smith NJ, Kern R, Picus M, Hoyer S, van Kerkwijk MH, Brett M,

Haldane A, del Río JF, Wiebe M, Peterson P, Gérard-Marchant P, Sheppard K, Reddy

T, Weckesser W, Abbasi H, Gohlke C, Oliphant TE. 2020. Array programming with

NumPy. *Nature* **585**:357–362. doi:10.1038/s41586-020-2649-2

Huerta-Cepas J, Szklarczyk D, Heller D, Hernández-Plaza A, Forslund SK, Cook H, Mende

DR, Letunic I, Rattei T, Jensen LJ, von Mering C, Bork P. 2019. eggNOG 5.0: a

hierarchical, functionally and phylogenetically annotated orthology resource based on

5090 organisms and 2502 viruses. *Nucleic Acids Res* **47**:D309–D314.

doi:10.1093/nar/gky1085

Hyatt D, Chen GL, LoCascio PF, Land ML, Larimer FW, Hauser LJ. 2010. Prodigal:

Prokaryotic gene recognition and translation initiation site identification. *BMC*

Bioinformatics **11**. doi:10.1186/1471-2105-11-119

Iturriaga G, Suárez R, Nova-Franco B. 2009. Trehalose metabolism: From osmoprotection to

signaling. *Int J Mol Sci* **10**:3793–3810. doi:10.3390/ijms10093793

Jandhyala SM, Talukdar R, Subramanyam C, Vuyyuru H, Sasikala M, Reddy DN. 2015. Role

of the normal gut microbiota. *World J Gastroenterol* **21**:8836–8847.

doi:10.3748/wjg.v21.i29.8787

Karnkowska A, Treitli SC, Brzoň O, Novák L, Vacek V, Soukal P, Barlow LD, Herman EK,

Pipaliya S V., Pánek T, Žihala D, Petrželková R, Butenko A, Eme L, Stairs CW, Roger

AJ, Eliáš M, Dacks JB, Hampl V, Battistuzzi FU. 2019. The Oxymonad Genome

Displays Canonical Eukaryotic Complexity in the Absence of a Mitochondrion. *Mol*

Biol Evol **36**:2292–2312. doi:10.1093/molbev/msz147

Karnkowska A, Vacek V, Zubáčová Z, Treitli SC, Petrželková R, Eme L, Novák L, Žárský

V, Barlow LD, Herman EK, Soukal P, Hroudová M, Doležal P, Stairs CW, Roger AJ,

Eliáš M, Dacks JB, Vlček Č, Hampl V. 2016. A eukaryote without a mitochondrial

organelle. *Curr Biol* **26**:1274–1284. doi:10.1016/j.cub.2016.03.053

Karp PD, Latendresse M, Caspi R. 2011. The pathway tools pathway prediction algorithm.

Stand Genomic Sci **5**:424–429. doi:10.4056/sigs.1794338

Karp PD, Paley SM, Midford PE, Krummenacker M, Billington R, Kothari A, Ong WK,

Subhraveti P, Keseler IM, Caspi R. 2020. Pathway Tools version 24.0: Integrated

Software for Pathway/Genome Informatics and Systems Biology. *arXiv e-prints*.

Kaviraj M, Kumar U, Snigdha A, Chatterjee S. 2024. Nitrate reduction to ammonium: a

phylogenetic, physiological, and genetic aspects in Prokaryotes and eukaryotes. *Arch*

Microbiol **206**:297. doi:10.1007/s00203-024-04009-0

Klingenberg H, Meinicke P. 2017. How to normalize metatranscriptomic count data for

differential expression analysis. *PeerJ* **2017**. doi:10.7717/peerj.3859

Kramer J, Özkaya Ö, Kümmerli R. 2020. Bacterial siderophores in community and host

interactions. *Nat Rev Microbiol* **18**:152–163. doi:10.1038/s41579-019-0284-4

Langmead B, Salzberg SL. 2012. Fast gapped-read alignment with Bowtie 2. *Nat Methods*

9:357–359. doi:10.1038/nmeth.1923

Law KC, Chung KK, Zhuang X. 2022. An Update on Coat Protein Complexes for Vesicle

Formation in Plant Post-Golgi Trafficking. *Front Plant Sci* **13**:1–9.

doi:10.3389/fpls.2022.826007

Layer G. 2021. Heme biosynthesis in prokaryotes. *Biochim Biophys Acta - Mol Cell Res*

1868:118861. doi:10.1016/j.bbamcr.2020.118861

Leventhal GE, Ackermann M, Schiessl KT. 2019. Why microbes secrete molecules to modify

their environment: The case of iron-chelating siderophores. *J R Soc Interface* **16**.

doi:10.1098/rsif.2018.0674

Liao Y, Smyth GK, Shi W. 2014. featureCounts: an efficient general purpose program for

assigning sequence reads to genomic features. *Bioinformatics* **30**:923–930.

doi:10.1093/bioinformatics/btt656

Love MI, Huber W, Anders S. 2014. Moderated estimation of fold change and dispersion for

RNA-seq data with DESeq2. *Genome Biol* **15**:550. doi:10.1186/s13059-014-0550-8

Matsen FA, Kodner RB, Armbrust EV. 2010. pplacer: linear time maximum-likelihood and

Bayesian phylogenetic placement of sequences onto a fixed reference tree. *BMC*

Bioinformatics **11**:538. doi:10.1186/1471-2105-11-538

Michael AJ. 2016. Polyamines in eukaryotes, bacteria, and archaea. *J Biol Chem* **291**:14896–

14903. doi:10.1074/jbc.R116.734780

Novák LVF, Treitli SC, Pyrih J, Hałakuc P, Pipaliya S V., Vacek V, Brzoň O, Soukal P, Eme

L, Dacks JB, Karnkowska A, Eliáš M, Hampl V. 2023. Genomics of Preaxostyla

Flagellates Illuminates the Path Towards the Loss of Mitochondria. *PLoS Genet* **19**:1–

30. doi:10.1371/journal.pgen.1011050

Nurk S, Meleshko D, Korobeynikov A, Pevzner PA. 2017. MetaSPAdes: A new versatile

metagenomic assembler. *Genome Res* **27**:824–834. doi:10.1101/gr.213959.116

O' Donnell MM, Harris HMB, Ross RP, O'Toole PW. 2017. Core fecal microbiota of

domesticated herbivorous ruminant, hindgut fermenters, and monogastric animals.

Microbiologyopen **6**. doi:10.1002/mbo3.509

Ondov BD, Treangen TJ, Melsted P, Mallonee AB, Bergman NH, Koren S, Phillippy AM.

2016. Mash: Fast genome and metagenome distance estimation using MinHash. *Genome*

Biol **17**:1–14. doi:10.1186/s13059-016-0997-x

Parks DH, Chuvochina M, Chaumeil PA, Rinke C, Mussig AJ, Hugenholtz P. 2020. A

complete domain-to-species taxonomy for Bacteria and Archaea. *Nat Biotechnol*

38:1079–1086. doi:10.1038/s41587-020-0501-8

Partida-Rodriguez O, Nieves-Ramirez M, Laforest-Lapointe I, Brown EM, Parfrey L,

Valadez-Salazar A, Thorson L, Morán P, Gonzalez E, Rascon E, Magaña U, Hernandez

E, Rojas-Velázquez L, Torres J, Arrieta MC, Ximenez C, Finlay BB. 2021. Exposure to

Parasitic Protists and Helminths Changes the Intestinal Community Structure of

Bacterial Communities in a Cohort of Mother-Child Binomials from a Semirural Setting

in Mexico. *mSphere* **6**. doi:10.1128/msphere.00083-21

Patro R, Duggal G, Love MI, Irizarry RA, Kingsford C. 2017. Salmon provides fast and bias-

aware quantification of transcript expression. *Nat Methods* **14**:417–419.

doi:10.1038/nmeth.4197

Price MN, Dehal PS, Arkin AP. 2010. FastTree 2 – approximately maximum-likelihood trees

for large alignments. *PLoS One* **5**.

R Core Team. 2021. R: A language and environment for statistical computing.

Schofield PJ, Edwards MR, Matthews J, Wilson JR. 1992. The pathway of arginine

catabolism in *Giardia intestinalis*. *Mol Biochem Parasitol* **51**:29–36. doi:10.1016/0166-

6851(92)90197-R

Seemann T. 2014. Prokka: rapid prokaryotic genome annotation. *Bioinformatics* **30**:2068–

2069. doi:10.1093/bioinformatics/btu153

Shaw J, Yu YW. 2023. Fast and robust metagenomic sequence comparison through sparse

chaining with skani. *Nat Methods* **20**:1661–1665. doi:10.1038/s41592-023-02018-3

Shimizu Y, Uemura T. 2022. The sorting of cargo proteins in the plant trans-Golgi network.

Front Plant Sci **13**:1–10. doi:10.3389/fpls.2022.957995

Sugai Y, Katsuyama Y, Ohnishi Y. 2016. A nitrous acid biosynthetic pathway for diazo

group formation in bacteria. *Nat Chem Biol* **12**:73–75. doi:10.1038/nchembio.1991

Sukumaran J, Holder MT. 2010. DendroPy: A Python library for phylogenetic computing.

Bioinformatics **26**:1569–1571. doi:10.1093/bioinformatics/btq228

Tanaka T, Goto K, Iino M. 2017. Diverse Functions and Signal Transduction of the Exocyst

Complex in Tumor Cells. *J Cell Physiol*. doi:10.1002/jcp.25619

Teulière J, Bernard G, Bapteste E. 2020. The Distribution of Genes Associated With

Regulated Cell Death Is Decoupled From the Mitochondrial Phenotypes Within

Unicellular Eukaryotic Hosts. *Front Cell Dev Biol* **8**:1–8. doi:10.3389/fcell.2020.536389

Thursby E, Juge N. 2017. Introduction to the human gut microbiota. *Biochem J* **474**:1823–

1836. doi:10.1042/BCJ20160510

Treitli SC, Kolisko M, Husník F, Keeling PJ, Hampl V. 2019. Revealing the metabolic

capacity of *Streblomastix strix* and its bacterial symbionts using single-cell

metagenomics. *Proc Natl Acad Sci* **116**:19675–19684. doi:10.1073/pnas.1910793116

Treitli SC, Kotyk M, Yubuki N, Jirounková E, Vlasáková J, Smejkalová P, Šípek P, Čepička

I, Hampl V. 2018. Molecular and Morphological Diversity of the Oxymonad Genera

Monocercomonoides and *Blattamonas* gen. nov. *Protist* **169**:744–783.

doi:10.1016/j.protis.2018.06.005

Treitli SC, Peña-Diaz P, Hałakuc P, Karnkowska A, Hampl V. 2021. High quality genome

assembly of the amitochondriate eukaryote *Monocercomonoides exilis*. *Microb*

Genomics **7**. doi:10.1099/mgen.0.000745

von Huth S, Thingholm LB, Kofoed P-E, Bang C, Rühlemann MC, Franke A, Holmskov U.

2021. Intestinal protozoan infections shape fecal bacterial microbiota in children from

Guinea-Bissau. *PLoS Negl Trop Dis* **15**:e0009232. doi:10.1371/journal.pntd.0009232

Wu Y-W, Simmons BA, Singer SW. 2016. MaxBin 2.0: an automated binning algorithm to

recover genomes from multiple metagenomic datasets. *Bioinformatics* **32**:605–607.

doi:10.1093/bioinformatics/btv638

Yarlett N, Martinez MP, Moharrami MA, Tachezy J. 1996. The contribution of the arginine

dihydrolase pathway to energy metabolism by *Trichomonas vaginalis*. *Mol Biochem*

Parasitol **78**:117–125. doi:10.1016/S0166-6851(96)02616-3

Zhou Z, Tran PQ, Breister AM, Liu Y, Kieft K, Cowley ES, Karaoz U, Anantharaman K.

2022. METABOLIC: high-throughput profiling of microbial genomes for functional

traits, metabolism, biogeochemistry, and community-scale functional networks.

Microbiome **10**. doi:10.1186/s40168-021-01213-8

Zhukov A, Popov V. 2023. Eukaryotic Cell Membranes: Structure, Composition, Research

Methods and Computational Modelling. *Int J Mol Sci* **24**. doi:10.3390/ijms241311226

Figure legends

Figure 1: Growth curve of *Monocercomonoides exilis* and sampling points. Concentration

of *M. exilis* (cells/mL^{-1}) grown in TYSGM-9 medium for seven days. Samples for thorough

analysis were taken on days 0, 2, 3 and 5, as highlighted by red squares. Error bars provide

standard deviations of the values from three measurements.

Figure 2: Relative read abundance of the members of the community. Each member of the community is represented as relative read abundance (%) corresponding to its genome on each day and replicate.

Figure 3: Potential contribution of the bacterial community to the biochemical cycling processes of carbon, nitrogen, iron and sulfur. Labels represent the main steps. Present steps are identified in green. Absent steps are identified in light grey. MAGs: number of MAGs responsible for a step.

Figure 4: Schematic summary of changes in the culture medium. Chemical classification of significant compounds of identified compounds per day (Supplementary Table 4), number of compounds exhibiting an increase in concentration to the previous day (targets), number of targets that *M. exilis* can synthesize independently, and number of compounds that can be synthesized because of the interaction between *M. exilis* and the bacterial community are shown. Key bacterial species identified by the m2m pipeline are shown for each pairwise comparison.

Figure 5: Differential expression of genes involved in carbohydrate metabolism and transport on *Monocercomonoides exilis* between day 5 and day 2. Identified sugar transporters and glycolysis, pentose phosphate pathways, and related reactions (adapted from Karnkowska et al. 2019). Up-regulated genes and processes are marked with a green arrow pointing up. Down-regulated genes and processes are marked with a red arrow pointing down. Non-differentially expressed genes and processes are marked with an orange equal symbol. adhE: Bifunctional acetaldehyde-CoA/alcohol dehydrogenase; ALDH: Aldehyde dehydrogenase; ACS: Acetyl-CoA synthetase (ADP-forming); ACCT/PycB: putative malonyl-CoA:pyruvate transcarboxylase; ACAT: Acetyl-CoA acetyltransferase; PFOR: Pyruvate-ferredoxin oxidoreductase; HYD: [FeFe]-hydrogenase; NADP-ME: NADP-dependent malic enzyme; PK: Pyruvate kinase; PPDK: Pyruvate phosphate dikinase; PEPCK:

Phosphoenolpyruvate carboxykinase; AAT: Aspartate aminotransferase; ASNA: Asparagine

synthetase; ENO: Enolase; iPGM: 2,3-bisphosphoglycerate independent phosphoglycerate

mutase; PGK: Phosphoglycerate kinase; GAPDH: Glyceraldehyde-3-phosphate

dehydrogenase; GAPN: NAD(P)-dependent glyceraldehyde-3-phosphate dehydrogenase;

TPI: Triose phosphate isomerase; DERA: Deoxyribose-phosphate aldolase; RBKS:

Ribokinase; FBA: Fructose-bisphosphate aldolase; PFP: Phosphofructokinase

(pyrophosphate-based); GPI: Glucose-6-phosphate isomerase; PGM: Phosphoglucomutase;

BMY: β-amylase; AMY: α-amylase; treS: Maltose alpha-D-glucosyltransferase/ alpha-

amylase; HXK: Hexokinase; RPDK; ribose-phosphate diphosphokinase; RPI: Ribose-5-

phosphate isomerase; RPE: Ribulose-phosphate 3-epimerase; TKT: Transketolase; TAL:

Transaldolase.

Figure 6: Differential expression of genes involved in amino acid metabolism and

transport on *Monocercomonoides exilis* between D5 and D2. Identified amino acid

importers, amino acid biosynthesis and arginine catabolism. Up-regulated genes and

processes are marked with a green arrow pointing up. Down-regulated genes and processes

are marked with a red arrow pointing down. Non-differentially expressed genes and

processes are marked with an orange equal symbol. Spontaneous reactions are shown in blue.

PGDH: Phosphoglycerate dehydrogenase; PSAT: Phosphoserine transaminase; PSPH:

putative phosphoserine phosphatase; SAT: Serine O-acetyltransferase; SEPHS:

Selenophosphate synthetase; SEPSECS: O-phospho-L-seryl-tRNA[Sec]:L-selenocysteinyl-

tRNA synthase; THRC: Threonine synthase: ASNA: Asparagine synthetase; ALAT: Alanine

aminotransferase; ACCT/PycB: putative malonyl-CoA:pyruvate transcarboxylase; AAT:

Aspartate aminotransferase; ASNA: Asparagine synthetase; DHE: Glutamate dehydrogenase;

GLNA: Glutamine synthetase: ADI: Arginine deiminase: OTC: Ornithine transcarbamylase:

CK: Carbamate kinase; OAT: Lysine/Ornithine aminotransferase.