Title: BRD4-mediated epigenetic regulation of endoplasmic reticulum-mitochondria contact sites is governed by the mitochondrial complex III

Authors: Brandon Chen[1,2], Drew C. Stark[2], Pankaj V. Jadhav[3,4], Theophilus M. Lynn-Nguyen[5,6,7], Benjamin S. Halligan[5,6] , Nicholas J. Rossiter[1,2], Nicole Sindoni[8], Myunsun Shin[8], Joao A. Paulo[9], Matthew Chang[3,4], Imhoi Koo[10], Sergei Koshkin[11], , Sanjana Eyunni[12-14], Paolo Ronchi[15], Michelle T. Paulsen[16], Pietro Morlacchi[17], David A. Hanna[18], Jason Lin[2], Rachel M. Guerra[19-22], David J. Pagliarini[19-22], Ruma Banerjee[18], Abhijit Parolia[12,13,21,23], Mats E. Ljungman[16], Andrew D. Patterson[10], Joseph D. Mancias[8], Shyamal Mosalaganti[3,4,25], Jonathan Z. Sexton[5,6,7,26,27], Tito Calì[28-30], Costas A. Lyssiotis[1,2,6,24,*], Yatrik M. Shah[1,2,6,24,*]

[1]Cellular and Molecular Biology Program, University of Michigan Medical School, Ann Arbor, MI 48109, USA

[2]Department of Molecular & Integrative Physiology, University of Michigan, Ann Arbor, MI 48109, USA.

[3]Life Sciences Institute, University of Michigan, Ann Arbor, Michigan, 48109, United States.

[4]Department of Cell and Developmental Biology, University of Michigan, Ann Arbor, Michigan, 48109, United States.

[5]Department of Medicinal Chemistry, College of Pharmacy, University of Michigan, Ann Arbor, MI 48109

[6]Department of Internal Medicine, Division of Gastroenterology and Hepatology, Michigan Medicine at the University of Michigan, Ann Arbor, MI 48109

[7]College of Pharmacy, University of Michigan, Ann Arbor, MI, 48109, USA

[8]Division of Radiation and Genome Stability, Department of Radiation Oncology, Dana-Farber Cancer Institute, Harvard Medical School, Boston, MA, USA.

[9]Department of Cell Biology, Harvard Medical School, Boston, MA, USA

[10]Department of Veterinary and Biomedical Sciences, The Pennsylvania State University, University Park, PA, USA

[11]The Huck Institutes of the Life Sciences, The Pennsylvania State University, University Park, PA, USA.

[12]Michigan Center for Translational Pathology, University of Michigan, Ann Arbor, MI, 48109

[13]Department of Pathology, University of Michigan, Ann Arbor, MI, 48109

[14]Molecular and Cellular Pathology Program, University of Michigan, Ann Arbor, MI, 48109

[15]Electron Microscopy Core Facility, European Molecular Biology Laboratory, 69117 Heidelberg, Germany

[16]Department of Radiation Oncology, University of Michigan, Ann Arbor, MI, 48109, USA

[17]Agilent Technologies, Inc., Lexington, MA, 02421, USA

[18]Department of Biological Chemistry, University of Michigan Medical School, Ann Arbor, MI, 48109, USA

[19]Departament of Genetics, Washington University School of Medicine, St. Louis, MO, 63110, USA

[20]Department of Biochemistry and Molecular Biophysics, Washington University School of Medicine, St. Louis, MO ,63110, USA

[21]Department of Cell Biology and Physiology, Washington University School of Medicine, St. Louis, MO 63110, USA

[22]Howard Hughes Medical Institute, St. Louis, MO, 63110, USA

[23]Department of Urology, University of Michigan, Ann Arbor, MI, 48109

[24]Rogel Cancer Center, University of Michigan, Ann Arbor, MI, USA

[25]Department of Biophysics, College of Literature, Science and the Arts, University of Michigan, Ann Arbor, MI, 48109, United States.

[26]Center for Drug Repurposing, University of Michigan, Ann Arbor, MI, 48109, USA
[27]Michigan Institute for Clinical and Health Research, University of Michigan, Ann Arbor, MI
48109, USA
[28]Department of Biomedical Sciences, University of Padova, 35131, Padova, Italy
[29]Padova Neuroscience Center (PNC), University of Padova, 35131, Padova, Italy
[30]Center for Neurodegenerative Disease Research (CESNE), University of Padova, 35131,
Padova, Italy
* To whom correspondence should be addressed: shahy@umich.edu, clyssiot@umich.edu

Keywords: Endoplasmic reticulum-mitochondria contact sites, bromodomain protein, high-
throughput screening, mitochondrial electron transport chain.

Abstract:

Inter-organellar communication is critical for cellular metabolic homeostasis. One of the most abundant inter-organellar interactions are those at the endoplasmic reticulum and mitochondria contact sites (ERMCS). However, a detailed understanding of the mechanisms governing ERMCS regulation and their roles in cellular metabolism are limited by a lack of tools that permit temporal induction and reversal. Through unbiased screening approaches, we identified fedratinib, an FDA-approved drug, that dramatically increases ERMCS abundance by inhibiting the epigenetic modifier BRD4. Fedratinib rapidly and reversibly modulates mitochondrial and ER morphology and alters metabolic homeostasis. Moreover, ERMCS modulation depends on mitochondria electron transport chain complex III function. Comparison of fedratinib activity to other reported inducers of ERMCS revealed common mechanisms of induction and function, providing clarity and union to a growing body of experimental observations. In total, our results uncovered a novel epigenetic signaling pathway and an endogenous metabolic regulator that connects ERMCS and cellular metabolism.

Main Texts:

The dynamic regulation of organelle networks and inter-organellar communication is critical for cellular metabolism and physiology. Direct inter-organellar communication occurs at membrane contact sites, which directly regulate metabolite transport, protein complex organization, signaling and organellar function[1,2]. One of the most well studied and abundant inter-organellar contact sites are endoplasmic reticulum-mitochondria contact sites (ERMCS). While a wealth of research has defined the unique proteome at ERMCS, the precise spatiotemporal regulation by various signaling pathways, and evolutionarily conserved molecular tethering complexes that structurally support ER-to-mitochondria interactions remains still elusive. Functional genetic and biochemical studies have revealed essential roles of ERMCS, including phospholipid synthesis[3], calcium buffering[4], mitochondrial dynamics[5,6], mitochondrial DNA distribution[7] and autophagosome formation[8].

Dysregulation of ERMCS and tethers contributes to the etiology of various disease, including neurodegeneration[9], obesity[10], cancer[11], diabetes[12], and inborn errors of metabolism[13]. Understanding the fundamental mechanisms underlying ERMCS dysregulation is essential for identifying potential drug targets to restore ERMCS function in disease. However, significant gaps persist in our basic understanding of the molecular drivers governing the remodeling and organization of ERMCS. Moreover, the exact mechanisms by which the microenvironment, signal transduction, and molecular cues initiate, maintain, and modify adaptive responses of ERMCS are yet to be fully elucidated.

In this study, we have identified a novel epigenetic response and a metabolic redox state that regulate the formation and dynamics of ERMCS. We characterized an FDA-approved compound, fedratinib, which induces ERMCS formation via BRD4-mediated histone recognition and described it as a new tool for temporal, reversible control of ERMCS. Fedratinib treatment establishes selective ER wrapping around mitochondria with cristae structure defects, membrane potential loss, and metabolic rewiring. ERMCS induction requires an intact mitochondrial complex III, which is a major site of coenzyme Q (CoQ) oxidation. Attempts to modulate mitochondrial CoQ redox state suggest a potential mechanism where an increase in the reduced-to-oxidized CoQ ratio could block ERMCS. Importantly, our study demonstrates that the transcriptional and metabolic requirements for ERMCS induction by fedratinib are common to other cellular stressors known to triggers ERMCS and underscores the existence of a conserved regulatory network governing ERMCS.

High throughput pharmacogenomic screening identifies novel regulators of ERMCS. To follow ERMCS in living cells in real-time, we generated a panel of eleven isogenic gene-edited cell lines with a novel reversible split fluorescent ERMCS reporter, termed split-GFP based

contact site sensor (SPLICS) [14–16]. This genetically encoded sensor is stably integrated with a
doxycycline-inducible promoter for titrating reporter expression[17]. In addition, we engineered a
mitochondria matrix reporter (mitoTagRFP) to control for gene expression and monitor
mitochondrial abundance[18] (Fig. 1a). Single cell clones were profiled for those with inducible
and reversible green fluorescence as well as stable red fluorescence. Probe induction did not
have a deleterious effect on cell growth for 7 days, nor on oxygen consumption (OCR) or
extracellular acidification rate (ECAR) at 72 hours post reporter induction (Extended Data Fig.
1a and b). To demonstrate the sensitivity, accuracy, and specificity of the probe, SPLICS lines
were treated with a known inducer of ERMCS, the ER stress inducer thapsigargin[19] (Fig 1b), or
by over-expressing an artificial ER-mitochondria linker (ER-MT tether). Both treatments
increased the SPLICS signal significantly (Fig. 1c). Moreover, using control pairs of splitGFP
probes: mKate2- β_{11} + cytosolic GFP_{1-10}, mKate2- β_{11} + outer mitochondria membrane (OMM)
GFP_{1-10}, ER_{short} + cytosolic GFP_{1-10}, we did not see a remarkable increase in any of these probe
pairs in cells treated with thapsigargin[20] (Extended Data Fig. 1e). Using this inducible SPLICS
system, we generated eleven isogenic human and mouse cell lines in investigate ERMCS
regulation.
To discover novel mediators and chemical tools of ERMCS induction, we developed a
high-content phenotypic imaging screen with machine learning categorization approaches using
our SPLICS lines and assayed an FDA-approved drug repurposing library (Fig. 1d). Consistent
with known roles of ER stress in ERMCS, we identified proteasome inhibitors bortezomib and
carfilzomib[21]. Moreover, we identified microtubule inhibitors that modulate organelle movement
within in cells[22] (Supplementary Table 1 and Zenodo DOI: 10.5281/zenodo.13963526). We
decided to focus on fedratinib as it induced high levels of ERMCS, and its annotated
mechanisms of action have not been associated with ERMCS regulation (Fig. 1d). Fedratinib
inhibits Janus kinase 2 (JAK2) as well as bromodomain-containing protein 4 (BRD4)[23]. The
increase in ERMCS following fedratinib treatment was observed in a large panel of human and
mouse cell lines (Fig 1e). We focused on SW480 and HT1080, as these had the highest fold
change by fedratinib. ERMCS induced by fedratinib was both dose- and time-dependent with
estimated EC_{50} and ET_{50} values of 666.8 nM/4.7hr (SW480) and 404.7 nM/7.8 hr (HT1080,
which was evident using both the SPLICS reporter lines as well as by an orthogonal proximity
labeling method (Fig. 1f and g; Extended Data Fig. 1h, Supplementary Movies 1 and 2).
Importantly, the increase in ERMCS was reversible following a washout of fedratinib from the
culture media, highlighting the capability of SPLICS reporters to dynamically and reversibly
measure organelle contact sites in living cells (Fig. 1h). In addition, fedratinib did not
significantly increase fluorescence of our control splitGFP probe pairs, nor did it induce
autofluorescence when SPLICS is not activated (Extended Data Fig. 1f). Collectively, this
screen identified the first FDA-approved drug with potent bioactivity in increasing ERMCS
across cell lines with distinct tissue and organisms of origin.
148
Bromodomain and Extra-Terminal protein (BET)-dependent transcription response is
required for ERMCS induction. Fedratinib was first described as a JAK2-specific inhibitor for
treating myelodysplastic syndromes[24]. Thus, we tested the ERMCS-inducing activity of other
JAK2 inhibitors. We found that the JAK2 inhibitor Ruxolitinib did not induce ERMCS formation at
a concentration that inhibits JAK2-dependent STAT3 phosphorylation (Fig. 2a and Extended
Data Fig. 2a). In addition, cells with CRISPR interference (CRISPRi) mediated JAK2 knockdown
(KD) still responded to fedratinib-dependent ERMCS induction (Extended Data Fig. 2c and d).
Therefore, both genetic and pharmacological approaches suggest JAK2 is not the main target of
fedratinib for inducing ERMCS.
Fedratinib has other targets, including BRD4, a member of the bromodomain and extra-
terminal (BET) epigenetic reader family[25–27]. Fedratinib binds to the acetylated-lysine recognition
pocket of BRD4 and prevents BRD4 interaction with acetylated histones. Indeed, we observed

that canonical BRD4-hypersensitive targets, such as c-MYC, were decreased in expression
upon fedratinib treatment (Extended Data Fig. 2b and m). Similar to fedratinib, other inhibitors of
BRD4 from our small molecule screen also increased ERMCS (i.e. OTX-015 and IBET-762;
Zenodo DOI: 10.5281/zenodo.13963526.). To directly assess BRD4, we utilized the PROTAC
degrader of BRD4, dBET6, and siRNA-mediated KD, both of which increased ERMCS (Fig. 2a,
b, and c; Extended Data Fig. 2a).
Next, we directly investigated whether both fedratinib and thapsigargin works through
BRD4 inhibition. First, we conducted chromatin fractionation assay and found that fedratinib and
the positive control BET inhibitor JQ1 displaced BRD4 from chromatin, but not thapsigargin
(Extended Data Fig. 2d). This suggests that fedratinib inhibits BRD4 function by directly altering
its chromatin occupancy, but thapsigargin does not. Second, we investigated how fedratinib and
thapsigargin differentially affect genome wide BRD4 occupancy via chromatin
immunoprecipitation (ChIP). BRD4 ChIP-seq conducted from SW480 cells identified that
fedratinib has fewer BRD4 bound genomic loci in both promoter and non-promoter regions than
thapsigargin (Extended Data Fig. 2f and g), and the BRD4 bound regions are mostly promoters,
introns, or intergenic regions (Extended Data Fig. 2h). We then confirmed with ChIP-qPCR in
HT1080 cells that fedratinib led to loss of BRD4 occupancy at the PVT enhancer regions that
controls BRD4-hypersensitive gene MYC (Extended Data Fig. 2i). Therefore, we confirm
fedratinib acts through BRD4 inhibition to induce ERMCS while thapsigargin's mechanism is
BRD4 independent.
To identify whether BRD4-dependent ERMCS induction is transcriptionally regulated, we
inhibited transcription kinases cyclin-dependent kinases 8 and 9 (CDK8 and CDK9). Inhibition of
CDK8 and CDK9 both suppressed ERMCS induction (Fig. 2l and Extended Data Fig. 2e), thus
pointing out the importance of an active transcriptional program for ERMCS regulation. To
further test this we performed a pulse-chase experiment with RNA polymerase II (RNA pol II)
inhibitor Actinomycin D (ActD). Fedratinib- and thapsigargin-induced ERMCS were suppressed
when transcription activity was rapidly blocked (Extended Data Fig. 2f and g), whereas the
mitochondrial RFP signal was stable following acute transcriptional blockade.
To characterize the early transcriptional response upon ERMCS induction, we
conducted bromouridine (BrU) labeling of nascent mRNA sequencing, BrU-Seq, at 30 min and 2
hr after fedratinib treatment (Supplemental table 3). BRD4 hypersensitive target gene MYC
decreased its nascent mRNA transcription within 30 mins of fedratinib treatment (Fig. 2d).
Through gene set enrichment analyses (GSEA), we identified potent upregulation of genes
associated with the integrated stress response (ISR) and the unfolded protein response (UPR).
In addition, many of the ISR genes were increased at the protein level when measured by
quantitative proteomics (Extended Data Fig. 3a, b, and c and Supplementary Table 2). While
thapsigargin is thought to increase ERMCS via inducing ER stress, fedratinib did not increase
UPR markers phospho-PERK, IRE1α, CCPG1, ATF6, spliced XBP1, CHOP and BiP (Extended
Data Fig. 5a). Moreover, inhibiting ER stress propagation with chemical chaperone
tauroursodeoxycholic acid (TUDCA) did not reverse ERMCS induction (Extended Data Fig. 5d).
Therefore, we believe the induction of ERMCS by fedratinib is independent of the UPR.
However, we observed increased levels of ISR markers ATF4 and phosphorylated-eIF2α
at Ser51 upon fedratinib treatment (Extended Data Fig. 3d). To identify which eIF2α kinase arm
drives this ISR signature, we treated our cells with inducers of eIF2AK1/2/3/4 (GCN2, PKR,
PERK, HRI). The most robust inducer of ERMCS was the PKR activator BEPP, and inhibition of
PKR with its inhibitor PKR-IN-C16 reversed ERMCS induction and ATF4 expression (Extended
Data Fig.3e, g, h, and i). ISR inhibitor ISRIB decreased ATF4 expression by thapsigargin and
fedratinib treatment, yet it did not suppress ERMCS (Extended Data Fig. 3f). We, therefore,
posit that fedratinib induces ERMCS via PKR activation, but the ISR is unlikely to be a direct
activator of ERMCS. Additionally, fedratinib did not activate UPR^mito, and activating

mitochondrial ISR via oligomycin or CCCP was not sufficient to induce ERMCS formation
(Extended Data Fig. 3k and 6c).
We also observed that genes associated with autophagy increased with fedratinib
treatment. BRD4 inhibition has been shown to induce autophagy[28], and autophagosomes have
been shown to emerge from ERMCS[8]. Indeed, we confirmed increased autophagy with
fedratinib treatment by immunoblotting for lipidated LC3B (Extended Data Fig. 4a). Additionally,
when we combined fedratinib with an autophagy inhibitor chloroquine (CQ), we observed
enhanced LC3B lipidation, suggesting an elevated autophagic flux (Extended Data Fig. 4a).
Fedratinib treatment also increased lysosomes measured by lysotracker intensity in a dose
dependent manner (Extended Data Fig. 4b). However, even after inhibition of autophagy with
CQ or V-ATPase pump inhibitor Bafilomycin A (BafA), we still observed ERMCS formation
(Extended Data Fig. 4c). Moreover, colocalization studies of lysosomes and autophagosomes
with SPLICS did not show overlap at ERMCS (Extended Data Fig. 4d and e), indicating ERMCS
was not an early step of mitophagosome formation. Therefore, these data suggest autophagy
induction is a result of fedratinib treatment but not an inducer of ERMCS formation.
To identify the BRD4 co-regulated complex and transcription factors that coordinate
ERMCS modulation, we performed an unbiased CRISPR screen coupled with fluorescent
activated cell sorting (FACS), using a nuclear genome library. Following transduction with the
CRISPR library, which contained genes annotated with nuclear localization or roles in
epigenetic modification, we induced ERMCS formation with fedratinib, and sorted cells that
failed to induce ERMCS (bottom 10% SPLICS[low]/mitoTagRFP) (Fig. 2e). Through gene set
enrichment analyses (GSEA), we found a potent signature from components of the mediator
complex (Fig. 2f, g, and h; Supplementary Table 4). The mediator complex is a hetero-
multimeric protein complex that recruits other transcription factors, co-regulators, and RNA pol II
machineries along with BRD4. We generated knockdowns of mediator complex components
MED16 and MED24 KD via CRISPR interference (CRISPRi), which failed to induce ERMCS in
response to fedratinib or thapsigargin (Fig. 2j and k and Extended Data Fig. 2j and k).
Taken together, here we have demonstrated that Fedratinib induces ERMCS via
suppression of BRD4, and this activity is transcriptionally controlled. Moreover, induction of
ERMCS via the ER stressor thapsigargin also depends on an active gene expression program,
indicating that transcriptional rewiring to be common for other modes of ERMCS induction.
242
Fedratinib induces ER-wrapping of mitochondria at ERMCS. Organelle structures and inter-
organellar organization networks inform the metabolic state of cells. An increase in ERMCS
drives adaptive oxidative phosphorylation and supports ER homeostasis. Therefore, we
postulate that understanding how fedratinib impacts ERMCS architecture will provide valuable
information for a deeper understanding of the metabolic factors, regulation, and consequences
of ERMCS induction. To address this, we utilized various electron microscopy imaging
approaches to sequentially characterize the ERMCS architecture (Fig 3a).
From our transmission electron microscopy (TEM) on SW480 and HT1080 cells. We
observed structural changes, such as the ER membrane enveloping the mitochondria and the
collapse of cristae, we also noted the compartmentalization of structures within the inner
mitochondrial membrane (IMM) (Fig. 3b). In addition, approximately 15-20% of mitochondria
exhibited complete loss cristae and 40-45% demonstrated compartmentalization of the IMM (Fig.
3b). Although we did not observe consistent changes in circular cross-section, area, or aspect
ratio, a higher variability of these structural indices were seen in fedratinib-treated cells
(Extended Data Fig. 8d).
To further corroborate our observations and specifically probe ERMCS, we performed
correlative light and electron microscopy (CLEM) using the SPLICS reporter that precisely
marks ERMCS. We collected tilt-series guided by fluorescence from ERMCS in SW480 cells
reconstructed a three-dimensional tomogram (Fig. 3c and d). Fedratinib-treated cells, but not

vehicle controls, exhibited direct juxtaposition of the ER and outer mitochondrial membrane
(OMM) at specific SPLICS[Hi] locations (Fig. 3d), confirming our initial TEM observations (Fig. 3b).
Furthermore, we observed near-complete collapse of the inner mitochondrial membrane/cristae
at these contact sites. Typically, ERMCS are characterized by segments of the ER contacting
the OMM rather than complete ER envelopment of mitochondria. However, following fedratinib
treatment, we observed striking ER-mitochondria engulfment. Thus, fedratinib treatment not
only induced enhanced ER-mitochondria contact but also promoted unanticipated ultrastructural
reorganization at these sites.
Our TEM and CLEM approaches revealed ER-enclosure structures around select
mitochondria. To obtain global higher-resolution volumetric information of ERMCS, we used
high pressure freezing succeeded by freeze-substitution and focused ion beam-scanning
electron microscopy (FIB-SEM) on fedratinib- or vehicle-treated HT1080 cells. This approach
preserves cellular components in their near native state, avoiding chemical fixation artefacts that
might disrupt their delicate organization. Imaging through half a cell volume at 4 nm isotropic
pixel size. We trained a deep-learning model to distinguish mitochondria and ER. We
segmented these organelles in our dataset and identified mitochondria ensconced by ER in
fedratinib-treated cells (Fig. 3e). Consistent with our light microscopy data, we observed a range
of mitochondrial and cristae structure with varying degrees of ERMCS in fedratinib-treated cells
(Fig. 3f). Overall, we did not observe significant global changes in mitochondrial volume,
ERMCS distance, or ERMCS surface area (Fig. 3g-i). However, analysis of individual
mitochondria revealed a substantial increase in percent contact area (Fig. 3j). In conclusion, by
combining multiple EM modalities, we identified a novel ER-wrapped mitochondrial morphology
in fedratinib-treated cells, characterized by a heterogeneous mitochondrial population with
varying degrees of ERMCS.
286
Metabolic and lipidomic changes associated with selective depolarization of ER-wrapped
mitochondria. To visualize ERMCS in a larger cellular context, intact mitochondria and ER
networks, dynamics, and morphology in the whole cell were imaged with light microscopy (LM).
We found that a subset of mitochondria were rounded and swollen in fedratinib-treated cells,
and colocalized with the ERMCS SPLICS reporter sites (Fig. 4a), suggesting that the very close
proximity with the ER is needed to promote the above-mentioned ultrastructural re-organization
of mitochondria. We termed mitochondria with high ERMCS SPLICS[Hi] and those are not
associated with ERMCS as SPLICS[Lo]. Meanwhile, the SPLICS[Lo] mitochondria with a higher
ERMCS distance still form normal network-like morphology. However, OMM proteins (MFN2,
DRP1, and phospho-DRP1[S616/637]), IMM organization proteins (OPA1, MICOS complex
component MIC60 and MIC25), and mitochondrial electron transport chain (ETC) expression did
not change in whole cell western blots (Extended Data Fig. 6a and b). In addition, expression of
the ER sheet and tubule shaping proteins CLIMP63 (CKAP4) and RTN4 did not change
(Extended Data Fig. 6c). These data suggest that the structural changes of ERMCS induction
by fedratinib did not induce changes by altering levels of the canonical mitochondria dynamics
or ER structure shaping proteins. However, we cannot rule out the role of membrane lipid
remodeling that contributes to mitochondrial outer membrane and inner membrane structural
changes.
To further characterize mitochondria function, we profiled mitochondrial mass,
membrane potential (MMP/$\Delta\Psi$m), oxygen consumption rate (OCR), extracellular acidification
rate (ECAR), mitochondria reactive oxygen species (ROS), total cellular ROS, mitochondrial
calcium response, and mitochondrial DNA (mtDNA) content. Again, we did not observe
consistent differences between fedratinib versus control-treated cells in bulk measurements
from whole cells (Extended Data Fig. 6c-g, i-k and Extended Data Fig. 7). From the seahorse
OCR analysis, we analyzed basal respiration, maximal respiration, ATP-linked respiration, spare
respiratory capacity (SRC), and proton leak. We observed a consistent decrease in maximal

respiration ATP-linked respiration and SRC from both cell lines (Extended Data Fig. 7c, d, and
e), suggesting ERMCS induction by fedratinib could limit adaptive respiratory changes. Since
these mitochondrial phenotyping approaches report changes in the average behavior at a whole
cell level changes, they could mask phenotypic differences since not all mitochondria are
wrapped by ER in fedratinib-treated cells.
With these data in mind, we next employed single mitochondria functional profiling with
fluorescence imaging, which revealed intra-cellular heterogeneity in mitochondria membrane
potential. Mitochondria wrapped with ER (SPLICS^{Hi}) have lower ΔΨm compared to those
lacking ERMCS (SPLICS^{Lo}) (Fig. 4a-c; Extended Data Fig. 8b). Orthogonal validation with single
mitochondria flow cytometry further confirmed lower ΔΨm when ERMCS is induced that is
absent at the whole cell level (Fig. 4d). Of note, we did not observe changes in mitochondrial
ROS at the single mitochondrial or whole cell level (Extended Data Fig. 8a and c). Additionally,
FACS analysis at the single mitochondrial level revealed higher mitochondrial calcium in
SPLICS^{Hi} compared to SPLICS^{Lo} mitochondria (Extended Data. Fig. 6l).
These points notwithstanding, we investigated the impact of fedratinib on whole cell
metabolism using metabolomics and lipidomics. Through dimensional reduction approaches to
identify variability in metabolite and lipid profiles, we found that the metabolome is significantly
altered upon induction of ERMCS (Fig.4e; Extended Data Fig. 9a-c; Supplementary Table 5).
Indeed, the changes to the lipidome were even more profound when we performed lipidomics
on mitochondria isolated from fedratinib-treated cells as we observed a dramatic decrease in
mitochondrial lipids (Fig. 4f; Supplementary Table 6). Since loss of cardiolipin, a mitochondrial
inner membrane lipid, or alteration in the phospholipid saturation state can directly affect cristae
organization[30]. From our mitochondrial lipidomics that compared lipidome between fedratinib
and vehicle treated cells, we identified a global decrease in lipids, particularly the structural
phospholipid phosphatidylcholine (PtdCho) (Fig. 4f).
338
Fedratinib induces subcellular proteomic changes specific to ER-enveloped
mitochondria. To further explore mitochondrial heterogeneity within cells, we hypothesized that
distinct mitochondrial populations, with varying degree of ER contact can be detected by single
mitochondrial flow cytometry assay. Although flow cytometric analysis of ERMCS at the whole
cell level revealed continuous distribution, analysis at the individual mitochondria level revealed
two distinct population with high and low levels of SPLICS intensity. Specifically, 36.98% of
mitochondria in SW480 cells and 46.30% in HT1080 cells are SPLICS^{Hi}, whereas over 50% of
the mitochondria exhibited low SPLICS signal (SPLICS^{Lo}) (Fig. 5a). These data align with our
microscopy data, which indicated only a subset of mitochondria were enveloped by the ER.
Additionally, using forward scatter (FSC) as a proxy of particle size, we also validated that
SPLICS^{Hi} mitochondria are larger than those not associated with the ER (Fig. 5b and c).
To further identify mitochondrial level proteomic heterogeneity, we conducted
fluorescence activated mitochondria sorting (FAMS) to isolate SPLICS^{Hi} and SPLICS^{Lo}
mitochondria in fedratinib-treated cells relative to vehicle treated control (Fig. 5d)[31]. We
successfully enriched for 90.8% of SPLICS^{Hi} in SW480 and 85.3% in HT1080 cells (Fig. 5e and
f). We then conducted quantitative proteomics on the FAMS-sorted mitochondria from SW480
cells. Compared to the whole cell proteome, we enriched for mitochondrial protein by 3.8-fold
(Fig. 5g). Fedratinib treatment increased ER localized protein around 2-fold, and all FAMS-
sorted mitochondria have significant mitochondria enrichment (Fig. 5h and i). When comparing
SPLICS^{Hi} to SPLICS^{Lo} mitochondria, we further enriched for ER proteins by approximately 4.8-
fold (Fig. 5j), supporting higher degree of ER engagement with SPLICS^{Hi} mitochondria. Gene
ontology (GO) analysis of SPLICS^{Hi} mitochondria identified significant enrichment of terms
associated with mitochondria, membranes, and organelles (Fig. 5k).
When focusing on SPLICS^{Hi} mitochondria, we identified a strong enrichment of N-linked
glycosylation machineries, including the catalytic subunits of the oligosaccharyltransferase (OST)

complex STT3A and STT3B and other OST components such as RPN1 and MAGT1 (Fig. 5l).
Additionally, mitochondrial proteins selectively enriched in SPLICS^{Hi} mitochondria, included
voltage dependent anion channels (VDAC) VDAC1, 2, and 3, which are known ERMCS tethers.
Interestingly, we also identified enrichment of enzymes involved in reductive proline
biosynthesis genes ALDH18A1/P5CS, PYCR1, and PYCR2, as well as fatty acid oxidation
(FAO) proteins CPT1A, ACADVL, HADHA/B, ACAT1, and SLC25A20. This suggests that
mitochondria in contact with ER could be involved in distinct metabolic processes compared to
those that are not interacting with the ER (Figure 5l). Through our single mitochondrial level
analyses, we have identified unique proteome associated with mitochondria in physical contact
with the ER.
374
ETC Complex III function is necessary for ERMCS formation. Based on the drastic
mitochondrial structural and less pronounced but significant metabolic changes associated with
ERMCS induction, we hypothesized that an optimal metabolic state would be required to sustain
metabolic demands for ERMCS formation. To test this model, we employed a variety of
mitochondrial perturbations, and found that hypoxia reduced ERMCS (Fig. 6a). In addition to
directly inhibiting oxygen dependent-reactions[32], hypoxia also rewires cellular metabolism
through hypoxia inducible factors (HIFs). In contrast to hypoxia, HIF induction with the PDH
inhibitor FG-4592 did not similarly attenuate fedratinib-induced ERMCS, indicating that the
hypoxia effect is independent of HIF activation but oxygen dependent (Fig. 6a, extended figure
10a).
385   Since mitochondria are the largest oxygen consumers within the cell, we individually
inhibited the electron transport chain (ETC) complexes I-V, or depolarized membrane potential,
and then asked whether a specific ETC component is required for ERMCS induction (Fig. 6b).
Only complex III and IV inhibition suppressed the formation of ERMCS induced by fedratinib
(Fig. 6c). We also report that CIII inhibition with Antimycin A reversed mitochondrial PtdCho and
lipid depletion from fedratinib treatment (Fig 6e and f), connecting consistent regulatory patterns
between ERMCS and mitochondrial lipid regulation.
392   Unlike ETC inhibitors that restrict oxygen consumption and generate mitochondrial ROS,
393   inhibiting complex III or IV suppresses the ability to carry out CoQ/ubiquinol oxidation. Inhibition
394   of complex III and IV leads to the buildup of CoQH$_2$, which leads to a reductively shifted
395   CoQH$_2$/CoQ pool, which has been associated with reverse electron transport (RET) at complex I
396   and II, generating ROS and reducing fumarate to succinate[33], respectively.
397   We found that neither mitochondrial-specific ROS inducer mito-paraquat (mitoPQ) nor a
complex I inhibitor that increases oxidized CoQ pool suppressed ERMCS formation (Extended
Data Fig. 10b). Moreover, blocking electron leak and superoxide generation at complex III Q$_0$
site with S3QEL-2 did not rescue antimycin dependent ERMCS suppression (Extended Data Fig.
6h). Therefore, we hypothesized that complex III inhibition could suppress ERMCS formation
from accumulation of reduced CoQH$_2$, but not the generation of ROS. To relieve CoQH$_2$ buildup,
we genetically expressed a *Ciona intestinalis* alternative oxidase (AOX)[33], which oxidizes
CoQH$_2$ to CoQ in an antimycin/myxothiazol-insensitive manner (Fig. 6d). If CoQH$_2$ accumulation
drives ERMCS suppression, AOX expression should restore ERMCS induction by maintaining
an oxidized CoQ pool. Notably, cells expressing AOX retain fedratinib-induced ERMCS when
simultaneously subjected to complex III inhibitors (Fig. 6g; Extended Data Fig. 10c). This
regulation is not limited to fedratinib as thapsigargin induced ERMCS are also regulated by
similar mechanisms (Extended Data Fig. 10d and e). Surprisingly, fedratinib did not directly alter
the CoQH$_2$/CoQ redox ratio or CoQ$_{10}$ abundance at the whole cell or mitochondria level when
measured by mass spectrometry and high performance liquid chromatography (HPLC)
(Extended Data Fig. 10h and i). This suggest that perhaps maintaining the ability for complex III
to carry out CoQ oxidation, rather than the total levels, is crucial for ERMCS induction. Through
confocal microscopy, we also showed that ER-wrapped mitochondrial are enriched with CoQ

biosynthetic enzyme COQ9, indicating a potential role in local CoQ channeling capabilities
(Extended Data Fig. 10j).
In addition to expressing AOX, we also utilized an orthogonal method to alter electron
flow to complex III. Under standard conditions, complex I contributes the majority of electrons to
the ETC via NADH. Co-inhibiting complex I and complex III limits $CoQH_2$ generation, thereby
decreasing the $CoQH_2/CoQ$ ratio. Indeed, inhibition of complex I and III restored ERMCS
inducibility (Fig. 6h). We attempted to rescue the impact of complex IV inhibitors, H_2S and
hypoxia, by similarly inhibiting complex I. However, piericidin A was unable to restore ERMCS
levels (Extended Data Fig. 10f and g). This may be attributed to the dominant role of SQOR-
mediated CoQ reduction as the main electron donor under H_2S conditions, as well as the
inhibition of NADH-dependent electron flux into the ETC during hypoxia.
These findings support that complex III-dependent functions such as CoQ oxidation
could influence the dynamic induction of ERMCS formation. More importantly, this regulation is
not limited to fedratinib induced ERMCS, as ERMCS induced by thapsigargin also responds to
methods that bypass Complex III or CoQ oxidation state manipulation. Consequently, we can
conclude that the Complex III regulated CoQ oxidation could act as a reliable indicator of
ERMCS dynamics and is a novel metabolic pre-requisite for ERMCS induction.
432
Discussion:
Inter-organellar contact sites mediate local signaling events, whole cell metabolism and
tissue physiology. The dysregulation of ERMCS has been implicated in diseases including
cancer, fragile X syndrome[34], and Friedrich's ataxia[13]. Despite their critical role in maintaining
metabolic homeostasis, the mechanisms governing the dynamics of ERMCS remain elusive, at
least in part due to the lack of potent inducers with temporal control. We identified fedratinib, an
FDA-approved drug for myelodysplastic syndrome that robustly induced the formation of
ERMCS across diverse cell lines. We demonstrated that the relevant molecular target of
fedratinib was the epigenetic regulator BRD4, and inhibition initiated an activating gene
expression program that coordinated with mediator complex to induce ERMCS. Using iterative
EM approaches, starting with TEM, CLEM, and then cryo-preserved FIBSEM, we identified that
ERMCS exhibited novel ultrastructural changes in response to fedratinib. These changes
included a tight and complete envelopment of the entire mitochondrial membrane at a subset of
mitochondria.This is similar to ER-mitochondrial interactions observed following infection with
human cytomegalovirus and SARS-CoV-2, metastatic melanoma cells, and fasted hepatocytes
in liver[35–37].
Despite these structural alterations, no consistent changes were observed in various ER
and mitochondrial functions at the whole-cell level. This is remarkable, given that approximately
30% of mitochondria exhibited massive structural alternations, indicating that "normal"
mitochondria may compensate to support cell homeostasis when other mitochondria are
dysfunctional. To dissect mitochondrial level heterogeneity, we optimized a single mitochondrial
sorting pipeline to distinguish different mitochondrial populations. As we observed from our
single mitochondria proteomics, mitochondria with abundant contact sites with the ER may
potentially be dedicated to support specific metabolic pathways, including proline biosynthesis
or beta oxidation. These results underscore the significance of considering mitochondrial
heterogeneity in the study of ERMCS. Moreover, through probing ETC function, we
demonstrated that ERMCS formation is dependent on the activity of mitochondria complex III/IV.
Methods bypassing their function restored ERMCS abundance, suggesting that non-affected
mitochondria adapt or persist when complex III function is suppressed, whereas the affected
mitochondria are targeted for ER engulfment.
Previous studies of BRD4 predominantly focused on its ability to act as a positive
regulator of transcription[38]. In contrast, our findings demonstrate that de-repression of its target
gene expression is required for ERMCS activity. Inhibition of BRD4 activates autophagy-related

genes and corrects for mitochondrial disease phenotype by increasing mitochondrial biogenesis
[28,39]. However, the direct transcriptional target(s) of BRD4 that regulate ERMCS are not clear.
Through transcriptomics we identified activation of the ISR in both cells with high basal ERMCS
and following fedratinib treatment. ISR activation relies on inducing PERK, HRI, GCN2, and
PKR[40]. Recent work has demonstrated that mitochondrial stress and sublethal activation of
mitochondrial outer membrane permeabilization (MOMP) can release cytochrome c and
subsequently activate ISR[41]. Interestingly, the loss of cristae at ERMCS resembles the effect
induced by MOMP activation[42]. Moreover, the dynamic nature of ERMCS is involved in
apoptosis regulation and sensitivity[43,44]. Since the UPR can activate the ISR, we did extensive
comparison of fedratinib to thapsigargin. We did not observe UPR activation after fedratinib
treatment nor changes in UPR targets (IRE1α, PERK, ERO1L, TMX1) that have been
associated with ERMCS induction[45–47]. However, direct activation of the ISR kinase PKR
induced ERMCS, while inhibiting PKR suppressed fedratinib induced ERMCS. Interestingly, ISR
inhibitor ISRIB decreased fedratinib-dependent ATF4 upregulation but failed to suppress
ERMCS. These findings reveal compelling connections between ISR signaling and ERMCS
regulation, underscoring the complex interactions among eIF2α kinases and distinct cellular
effects from early versus latent ISR activation. Discerning the relationship between ISR and
ERMCS warrants further study.
The ratio of oxidized CoQ to reduced $CoQH_2$ plays a critical signaling role that reflects
cellular metabolic states. While recognized as an electron carrier in the ETC [48], CoQ has
emerged as a multifunctional lipid involved in ferroptosis[49,50], pyrimidine biosynthesis[51], H_2S
detoxification[52], glycerol phosphate shuttle, and proline catabolism[53]. Accumulation of $CoQH_2$
under hypoxia or sulfide exposure is essential to maintain tissue homeostasis by utilizing
fumarate as a terminal electron acceptor instead of oxygen[33,52]. In physiological or pathological
hypoxia, tissues rewire metabolism via the canonical HIF oxygen sensing pathway and
reorganize organellar networks to maintain metabolism and other aspects of cellular
homeostasis. Indeed, we demonstrated that ETC complex III and IV could sustain a critical
function needed for establishing ERMCS as inhibiting complex III and IV inhibited ERMCS.
Utilizing genetic methods to bypass complex III inhibition via ectopic expression of alternative
oxidase (AOX) restored ERMCS abundance under complex III inhibition. Surprisingly, limiting
electron entry from complex I reversed the inhibitory response of antimycin or myxothiazol and
re-activated ERMCS induction. Based on the AOX and complex I/III co-inhibition results, we
speculate that there could potentially be a CoQ redox sensing mechanism as both rescue
approaches should prevent $CoQH_2$ accumulation under complex III inhibition alone. Further
studies to profile CoQ oxidation state on the subcellular level and spatial distribution will be
critical for drawing definitive conclusions with the inconsistent change in CoQ redox state and
CoQ abundance. Thus, we provide evidence to support that modulation of oxygen tension re-
organizes the subcellular organellar landscape and Complex III function could potentially be an
endogenous metabolic sensor for ERMCS formation. Importantly, while we made this discovery
with fedratinib, we also illustrated that other ERMCS inducers, like the ER stressor thapsigargin,
require the same molecular and metabolic mechanisms for ERMCS. This evidence suggests the
mechanisms described here are broadly applicable to other ERMCS regulatory modalities.
In conclusion, our study underscores the intricate interplay between molecular regulators
and metabolic cues that dictate the formation and functionality of ERMCS. Moreover, we
characterize a tool compound that can temporally and reversibly modulate ERMCS laying the
foundation for potential therapeutics targeting inter-organellar communication in diverse disease
contexts.

Materials and Methods:
Cell culture: HT1080, SW480 HCT116, DLD1, SW480, HeLa, U2OS, A459, PA-TU8902, K562,
Jurkat, B16, KPC7940 cells were maintained at 37°C in 5% CO_2 and 21% O_2. HT1080, SW480

HCT116, DLD1, SW480, HeLa, U2OS, A459, PA-TU8902, B16, and KPC7940 cells were
cultured in Dulbecco's Modified Eagle Medium (DMEM) supplemented with 10% FBS and 1%
antibiotic/antimycotic, whereas Jurkat and K562 cells were cultured in Roswell Park Memorial
Institute (RPMI) supplemented with 10% FBS and 1% antibiotic/antimycotic.
520
Generation of SPLICS stable cell lines: The SPLICS reporter is composed of a split GFP_{1-10}
barrel localized to the mitochondrial outer membrane, a P2A self-cleavage signal for equimolar
expression, and the remaining $beta_{11}$ fragment targeted to the ER membrane. Upon ERMCS
formation within the 8-10 nm distance, the split-GFP reporter forms intact GFP. Doxycycline-
inducible SPLICS reporter stable cell lines were generated via a three-plasmid PiggyBac
transposase system. Cells were co-transfected with the plasmids using Lipofectamine 2000
transfection reagent, then the next day selected with 2 mg/ml of G418 (Geneticin) for 7 days.
After selection, 100 ng/ml of doxycycline were used to induce reporter expression, and GFP
positive cells were sorted on Bigfoot Spectral Cell Sorter (Invitrogen) to isolate GFP positive
clones. Clones were cultured and screened for accurate localization of SPLICS reporter to
mitochondria and ER via live cell imaging and for normal mitochondrial oxygen consumption
rate. For ERMCS analysis with SPLICS, cells will be induced with 100 ng/ml of doxycycline for
minimally 24 hr before treatment or analysis.
534
Generation of genetically modified cell lines: sgRNAs (oligonucleotide sequences are indicated
in Supplementary Table 5) were ligated into BsmBI-linearized lentiCRISPR-v2 (for CRISPR KO)
or pLV hU6-sgRNA hUbC-dCas9-KRAB-T2a-Puro (pLV hU6-sgRNA hUbC-dCas9-KRAB-T2a-
Puro was a gift from Charles Gersbach (Addgene plasmid # 71236;
http://n2t.net/addgene:71236 ; RRID:Addgene_71236)) with T4 ligase (NEB). AOX
overexpression plasmid was achieved with pCW57.1_AOX-FLAG (pCW57.1_AOX-FLAG was a
gift from David Sabatini & Jessica Spinelli (Addgene plasmid # 177984;
http://n2t.net/addgene:177984 ; RRID:Addgene_177984)). MitoTagRFP plasmid was acquired
from Addgene (pclbw-mitoTagRFP was a gift from David Chan (Addgene plasmid # 58425;
http://n2t.net/addgene:58425 ; RRID:Addgene_58425). BFP and BFP-ER-mitochondria tether
were generated via Vectorbuilder (pLV[Exp]-Bsd-CMV>{new mTagBFP2} and pLV[Exp]-Bsd-
CMV>{new mTagBFP2 ER mito tether}). Lentiviral vectors expressing sgRNAs were transfected
into HEK293T cells with lentiviral packaging vectors CMV VSV-G and psPAX2 using
XtremeGene 9 transfection reagent (Roche). After 24 h, media was aspirated and replaced by
fresh media. The virus-containing supernatant was collected 48 h after transfection and filtered
using a 0.45 mm filter to eliminate cells. Cells to be transduced were plated in 6-well tissue
culture plates and infected in media containing virus and 8 μg/mL of polybrene. Cells were spin
infected by centrifugation at 1,100 g for 1.5 h. After transduction, media was changed, and cells
were selected with puromycin (for sgRNA lentiviral vector) for 72 hrs.
554
555 siRNA mediated knockdown: siRNAs for BRD4 were transfected using a reverse transfection
556 protocol and Lipofectamine RNAiMAX. siRNA was purchased from Horizon Discovery with the
557 following information: siGENOME Human BRD4 (23476) siRNA - SMARTpool, 5 nmol.
558
Western Blots: Whole-cell lysate preparations were described previously (Anderson et al., 2013).
Whole cell lysates were prepared from cell lines by RIPA buffer. Homogenates were incubated
in RIPA buffer for 15 min on ice followed by 13,000 rpm centrifugation for 15 min. Supernatants
were transferred to a new tube and mixed with 5× Laemmli buffer and boiled for 5 min. Lysates
containing 30–40 μg of protein per well were separated by SDS-PAGE, transferred onto
nitrocellulose membranes, and immunoblotted overnight at 4°C with indicated antibodies. All the
primary antibodies were used at a dilution of 1:1000. HRP-conjugated secondary antibodies

used were anti-rabbit and anti-mouse at a dilution of 1: 2000 and immunoblots were developed
using the iBright Imaging System and iBright Analysis Software (Thermofisher).
568

Antigen	Company	Catalog number
Vinculin (clone: E1E9V)	Cell Signaling Technology	13901
MFN2 (clone: D2D10)	Cell Signaling Technology	9482
DRP1 (clone: D6C7)	Cell Signaling Technology	8570
pS616-DRP1	Cell Signaling Technology	3455
pS637-DRP1	Cell Signaling Technology	4867
Mitofilin/MIC60/IMMT	Proteintech	10179-1-AP
MIC25/CHCHD6	Proteintech	20639-1-AP
RTN4/NOGO	Proteintech	10740-1-AP
COQ9	Proteintech	14874-1-AP
CLIMP63/CKAP4 (clone: G1/296)	Enzo Life Science	ENZ-ABS669-0100
OPA1 (clone: D6U6N)	Cell Signaling Technology	80471
ATF4 (clone: E4Q4E)	Cell Signaling Technology	97038
BiP/HSPA5 (clone: C50B12)	Cell Signaling Technology	3177
CHOP/DDIT3 (clone: L63F7)	Cell Signaling Technology	2895
BRD4 (clone: E2A7X)	Cell Signaling Technology	13440
BRD4 (For fractionation)	Bethyl	BL-149-2H5
c-MYC (clone: E5Q6W)	Cell Signaling Technology	18583
STAT3 (clone: 123H6)	Cell Signaling Technology	9139
pY705-STAT3 (clone: D3A7)	Cell Signaling Technology	9145
IP3R-I/II/III (clone: B-2)	Santa Cruz Biotechnology	SC-377518
TOM20 (9D8T4N)	Cell Signaling Technology	42406
CCPG1 (E3C5G)	Cell Signaling Technology	80158
PERK (C33E10)	Cell Signaling Technology	3192
IRE1α (14C10)	Cell Signaling Technology	3294
ATF6 (D4Z8V)	Cell Signaling Technology	65880
XBP1s (E9V3E)	Cell Signaling Technology	40435
LC3B (D11) XP	Cell Signaling Technology	3868
HIF1α (D2U3T)	Cell Signaling Technology	14179
JAK2 (D2E12) XP	Cell Signaling Technology	3230

569
CRISPR Screen: For the epigenetics library screen in SW480 reporter cells, the human sgRNA
library was described in this paper[54] was used. The titer of lentiviral supernatants was
determined by infecting targets cells at several amounts of virus in the presence of polybrene
(10 ug/mL), counting the number of drug resistant infected cells after 3 days of selection. 40
million target cells were infected at an MOI of ~0.5 and selected with puromycin (1 ug/mL) 72 h
after infection. An initial pool of 30 million cells was harvested for genomic DNA extraction. The
remaining cells were cultured for 6 days, treated with fedratinib for 24 hr, then 25 million cells
were sorted from the top and bottom 10% of SPLICS/mitoTagRFP fluorescent ratio on day 8.
After which cells were harvested for genomic DNA extraction. sgRNA inserts were PCR
amplified (primers listed in Supplementary Table 7, purified and sequenced on a NextSeq
(Illumina). Sequencing reads were mapped and the abundance of each sgRNA was tallied.
Gene score is defined as the median log2 fold change in the abundance between the initial and
final population of all sgRNAs targeting that gene. Full result of the screen with sequencing
primers can be found in the supplemental data.
584

<u>CRISPR screen analysis</u>: sgRNA read counts were generated by aligning sequencing reads to
the sgRNA library sequences, with only exact matches allowed. sgRNA abundances were
calculated for both sorted populations by first adding a pseudocount of 1, then dividing by the
total number of reads in each given sample. For each sgRNA, a differential score was
calculated as the log2 fold-change in sgRNA abundance between the GFP-high (top 10%) and
GFP-low (bottom 10%) cell populations. Gene level differential scores were generated by taking
the mean score of all sgRNAs targeting the same gene. The set of differential gene scores was
then standardized by subtracting the mean from each gene score then dividing by the standard
deviation (Z-score).
594
595 <u>Gene set enrichment analysis (GSEA) for CRISPR screen and proteomics</u>: Gene set
596 enrichment analysis was performed on the set of standardized differential gene scores using the
597 clusterProfiler (v4.2.2) package in R. The gseGO() function was called on the data using all
598 ontologies, min and max GSSsizes set to 10 and 500, respectively, FDR as the p-adjustment
599 method, and an FDR cutoff of 0.01.
600
<u>Imaging based drug repurposing screen</u>: 384-well, black, optically clear flat-bottom, tissue-
culture treated plates (Perkin Elmer cat# 6057302) were coated with 0.1mg/ml of poly-D-lysine
hydrobromide (Sigma-Aldrich cat# P6407-5MG) at 37°C for 1hr. Then each plate is washed with
molecular grade sterile water once, and then dried at room temperature for 1 hr. Each plate was
then dry-spotted with 50nL of 2mM drug solution in DMSO using a Beckman Echo 650 acoustic
liquid handler. In each plate, we have 2 columns of vehicle control (Dimethylsulfoxide/DMSO)
and 2 columns of Thapsigargin (50 nM) as positive control for inducing endoplasmic reticulum-
mitochondrial contact sites. With the plates coated and dry-spotted with drugs, 5000 SW480
colorectal cancer cells were plated in each well with complete DMEM media with a final drug
concentration at 2µM. After incubation with the drug for 24 hr, cells were washed with PBS 3
times then fixed with 4% paraformaldehyde (PFA) in 1x PBS (Thermofisher, cat#043368.9M) at
room temperature for 15 min in the dark. PFA was removed with 3 PBS washes and the fixed
cells were staining with Hoechst 33342 (Invitrogen, cat# H3570, final concentration: 5 µg/ml)
and HCS CellMask Deep Red Stain (Invitrogen, cat# H32721, final concentration: 1 µg/ml) for 1
hr at 37°C for 1hr. After staining, the cells were washed with PBS 3 times, and were left in PBS
in the dark at 4°C until imaging. Plates were imaged on a Yokogawa CellVoyager 8000 using a
40X/1.0NA water immersion objective lens and nine fields per well were imaged across three
channels to visualize Hoechst, GFP, Mitotag RFP and HCS CellMask Deep Red. Cellpose was
used to identify nuclei and a custom CellProfiler pipeline was used to delineate the whole cell,
mitochondria, GFP spots and per-cell feature extraction was performed. Endoplasmic
reticulum–mitochondria contact sites (ERMCS) were identified by counting GFP spots within the
cell. A XGBoost machine learning model was trained on centered/scaled cell-level features
trained against the negative control (DMSO) and thapsigargin treated positive control wells to
enable hit picking. XGBoost models were trained on the screening data controls from the entire
screen with an 80%/20% train/test split. Model performance was excellent with a R2>0.95. Z-
prime factors were calculated for each plate and plates below 0.5 were rejected and re-
screened. High correlation was found between the XGBoost score and the GFP spot count per-
cell, however, the XGBoost score exhibited significantly higher separation between the controls
and led to higher Z-primes.
630
A liberal hit threshold of 3 standard deviations above the mean of the negative control (µ+3σ)
was used to maximize hit detection despite an increase in the false positive rate. False positives
were eliminated in dose-response confirmation where 64 compounds were tested in 10-point:2-
fold dilution from 10uM to 20nM in triplicate wells. Of the 64 compounds tested, 33 were found
to be dose-responsive (52% confirmation rate). This is typical for a bioactive compound library

where the compound potencies range from sub-nanomolar to micromolar range and, often,
potent hits are obscured by cytotoxicity when screening at low micromolar concentrations. Hits
were cherry picked and confirmed with triplicate 10-point/2-fold dilution and EC50 values were
fit. Images and details of the screen were deposited to Zenodo and are publicly available in the
repository as part of this record: DOI: 10.5281/zenodo.13963526.

Oxygen Consumption Rate (Seahorse): Cells were seeded at 2×10^4 cells/well in 80 µl/well
of normal growth media (DMEM with 25 mM Glucose and 2 mM Glutamine) in an Agilent
XF96 V3 PS Cell Culture Microplate (#101085-004). To achieve an even distribution of cells
within wells, plates were incubated on the bench top at room temperature for 1 h before
incubating at 37 °C, 5% CO2 overnight. To hydrate the XF96 FluxPak (#102416-100),
200 µL/well of sterile water was added and the entire cartridge was incubated at 37 °C, no
CO2 overnight. The following day, 1 h prior to running the assay, 60 µL/well of growth media
was removed from the cell culture plate, and cells were washed twice with 200 µL/well of assay
medium (XF DMEM Base Medium, pH 7.4 (#103575-100) containing 25 mM glucose
(#103577-100) and 2 mM glutamine (#103579-100)). After washing, 160 µL/well of assay
medium was added to the cell culture plate for a final volume of 180 µL/well. Cells were then
incubated at 37 °C, without CO2 until analysis. One hour prior to the assay, water from the
FluxPak hydration was exchanged for 200 µL/well of XF Calibrant (#100840-000) and the
cartridge was returned at 37 °C, without CO2 until analysis. Oligomycin (100 µM), FCCP
(100 µM), and Rotenone/Antimycin (50 µM) from the XF Cell Mito Stress Test Kit (#103015-
100) were re-constituted in assay medium to make the indicated stock concentrations. Twenty
microliters of Oligomycin was loaded into Port A for each well of the FluxPak, 22 µL of FCCP
into Port B, and 25 µL of Rotenone/Antimycin into Port C. Port D was left empty. The final
FCCP concentration was optimized to achieve maximal respiration in each condition. The Mito
Stress Test was conducted on an XF96 Extracellular Flux Analyzer and OCR was analyzed
using Wave 2.6 software. Following the assay, OCR was normalized to cell number utilizing
Agilent Cytation5 live cell imaging with Gen5 software.

MT-DNA Quantification: Cells were pelleted, washed 1X in PBS, and then lysed in buffer
(25 mM NaOH, 0.2 mM EDTA) for 15 minutes at 95°C. Lysis was neutralized with buffer
(40 mM Tris-HCl) and centrifuged at 16,000 x g for 10 minutes at 4°C. Supernatant containing
MT-DNA was quantified on with nanodrop. Primers amplifying the MT-DNA marker D-Loop and
the nuclear DNA marker β-Globin were used and the relative MT-DNA to nuclear DNA was
quantified in each sample[55].

Mitochondria isolation: SW480 and HT1080 cells were scraped and pelleted at 1000 x g for 5
minutes. Cell pellets were washed in 1X PBS and pelleted at 1000 x g for 5 minutes. Cells were
re-suspended in isolation buffer (200 mM sucrose, 10 mM Tris HCl, 1 mM EGTA/Tris, pH 7.4
(adjusted with 1M HEPES)), transferred into a 27-gauge needle and homogenized by 10
repeated resuspensions. Homogenates were transferred into tubes and spun at 600 x g for 10
minutes at 4°C. Supernatants were moved to a new tube and centrifugation was repeated. Then,
supernatants were moved to a new tube and pelleted at 7,000 x g for 10 minutes at 4°C. Pellets
were washed in isolation buffer and centrifugation was repeated two more times. For
biochemical analyses, mitochondria were aliquoted in 50 µg pellets.

Flow cytometry for cells: HT1080 and SW480 cells were seeded 48 hr to 72 hr prior to the
experiment at 50K and 75K cells density to achieve final 70% confluency by the time of analysis.
For analyzing SPLICS and mitoTagRFP intensity, cells were washed, trypsinized, and filtered
through a cell strainer prior to analyzing. For MitoSOX Deep Red(Dojindo, cat#: MT14-12),
MitoTracker Green (Invitrogen, cat#: M7514), Tetramethylrhodamine, Ethyl Ester,

Perchlorate/TMRE (Invitrogen, cat#: T669), and Rhod-2, AM (Invitrogen, cat#: R1244)
experiments, cells were stained with 1 μM of MitoSOX Deep Red, 100 nM of MitoTracker Green,
500 nM of TMRE, 10μM of DAPRed (stained for 30 minutes, and then gets replaced in regular
media) and 5μM of Rhod-2 for 30 min 37□°C, washed twice with warm PBS, and proceeded
with the same processing procedure as unstained cells. Fluorescence activated cell sorting
(FACS) was performed on Bigfoot Spectral Cell Sorter (Invitrogen), in which the fluorescence
intensity of minimum 10K individual cells was quantified in their specific channel. Data were
analyzed using FlowJo software (TreeStar).

Flow cytometry for mitochondria: HT1080 and SW480 cells were seeded at 1.5 million and 3
million in a 15cm plate to achieve final 70% confluency by the time of analysis. Upon 24 hr of
treatment with Fedratinib cells were stained with MitoTracker Deep Red or MitoSOX Deep Red.
Upon staining, 10% of whole cell fraction was set aside and then mitochondria were isolated
with the protocol mentioned. Briefly, cells were fixed at room temperature for 2 min with 2% PFA,
and then washed in mitochondria isolation buffer. FACS was performed on Bigfoot Spectral Cell
Sorter (Invitrogen), in which the fluorescence intensity of minimum 10K individual cells or 100K
mitochondria was quantified in their specific channel. Data were analyzed using FlowJo
software (TreeStar).

Fluorescence activated mitochondria sorting (FAMS): SW480 and HT1080
SPLICS/mitoTagRFP-expressing cells are plated at $3e^6$ and $6e^6$ respectively in a 150-mm plate
48 hours before sorting with doxycycline treatment to induce SPLICS expression. Cells were
treated with fedratinib or vehicle (DMSO) control overnight. The next morning, cells were
stained with MitoBrilliant 646 (Tocris #7700) at 200 nM for 1 hour prior to mitochondrial isolation.
Mitochondria isolations were conducted as previously indicated. FAMS was conducted on the
Bigfoot Spectral Cell Sorter (Invitrogen) with 70-micron nozzle. The fluidics were flushed with
deionized water for 5 minutes followed by equilibration with mitochondrial isolation buffer for
another 5 minutes. Polystyrene beads (Spherotech) with defined sizes (0.2 μm, 0.5 μm, 1 μm,
and 2 μm) from were used to determine gating size of mitochondria particles. After gating for
size, mitochondria populations were determined using mitoTagRFP and MitoBrilliant 646 double
positive gates. The populations then were analyzed based on GFP signal. SPLICS[Lo]
mitochondria are determined based on vehicle control, and SPLICS[Hi] mitochondria are
separated based on the formation of distinct GFP population in fedratinib-treated cells. For
proteomics, contain 60 million mitochondria were sorted in each condition and four replicates
were collected.

Live cell confocal imaging: 35mm glass bottom plates (Ibidi) were coated with 0.1mg/ml of poly-
D-Lysine (Sigma) for 1 hour at 37°C, washed once with sterile water, dry in the biosafety cabinet
for 2 hours, and then complete media was added on to equilibrate for 20 min in the incubator.
50K of HT1080 and 100K of SW480 cells were then plated to achieve final 60-70% confluency
by the time of imaging. Cells with HaloTag-Sec61 were stained with 200nM of JaneliaFluor 646
(JF646, Promega) for 30min at 37°C, and then washed once with warm PBS. Cells were
counterstained with 2000x dilution of Hoechst 33342 (Thermofisher) for 15min at 37°C.
FluoroBrite DMEM was used as the media for live cell imaging. The cells were imaged using
Zeiss LSM 980 Airyscan 2 microscope and detector equipped with a 63x oil objective with a 1.4
numerical aperture. CO_2 and humidity was equilibrated for at least 30 min before imaging. Post-
processing was done with Zen 3.4 (Blue edition).

Immunofluorescence (IF): Number 1.5 round 10-mm glass coverslips were coated with
0.1mg/ml of poly-D-Lysine (Sigma) for 1 hour at 37°C, washed once with sterile water, dry in the

biosafety cabinet for 2 hours, and then complete media was added on to equilibrate for 20 min
in the incubator. 50K of HT1080 and 100K of SW480 cells were then plated to achieve final 60-
70% confluency by the time of IF processing. Cells were washed with 1x PBS 3 times, then
fixed in 4% PFA for 10 min at room temperature in the dark. PFA was washed with 1x PBS for
another 3 times. Cells were then permeabilized with PBST (1x PBS with 0.25% Triton-X) for 10
min. Cells were then blocked with 5% bovine serum albumin (BSA) in 0.3% Triton-X PBST for
30 min. Cells were then processed for primary antibody staining at 4^{o}C for overnight with a
negative control without any primary antibody with indicated antibody dilution shown in the
methods section. Antibody staining and washing buffer is 1% BSA in 0.3% PBST. After primary
antibody staining, cells were washed for 3 times. Secondary antibody was stained at 1:400
dilution regardless of types of fluorophore conjugated secondary antibody for 1 hr at room
temperature along with 1:10,000 dilution of Hoechst 33342. After secondary staining, cells were
washed again for 3 times. Last, the coverslips were mounted using Vectashield Vibrance
Antifade Mounting Media (H-1700-10). After overnight curing in the dark at room temperature,
the cells were imaged using Zeiss LSM 980 Airyscan 2 microscope and detector equipped with
a 63x oil objective with a 1.4 numerical aperture. Post-processing was done with Zen 3.4 (Blue
edition).
754
Proximity ligation assay: Cover glasses (#1.5H Thickness Ø12, Thor Labs) were coated with
0.1mg/ml of poly-D-Lysine (Sigma) for 1 hour at 37^{o}C, washed once with sterile water, dry in the
biosafety cabinet for 2 hours, and then complete media was added on to equilibrate for 20 min
in the incubator. 25K of HT1080 and 50K of SW480 cells were then plated to achieve final 60-
70% confluency by the time of imaging. PLA procedure was conducted via suggested
manufacture using Duolink® In Situ Detection Reagents FarRed (Sigma-Aldrich,
cat#DUO92013) with primary antibody pairs TOM20 and IP3R-I/II/III.
762
Lattice Light sheet imaging: Same cell conditions were used as live cell confocal imaging. Zeiss
Lattice Light Sheet 7 were used and imaged with 20x objective with an NA of 1.0. Cells were
treated with Fedratinib and then monitored for 16 hr with images getting acquired every 5 min.
Initial post-processing was done with Zen 3.4 (Blue edition). Images were deconvolved and then
deskewed in the Zen 3.4. (Deskewing transforms the volume into traditional X, Y, Z coordinates).
Then, Arivis Vision 4D (ver. 4.1.0) was used for puncta and cell segmentation. Deconvolved and
deskewed datasets were analyzed with Arivis Vision 4D (ver. 4.1.0). For punctas, segmentation
was done via the Blob Finder tool. The resulting segments were then filtered by size to remove
false positives. For cells, maximum intensity projections of the deconvolved deskewed data
were created. Hand segmentations were used to train a machine learning segmentation model.
The model was then used to segment each time point in the maximum intensity projection time
series. To get the number of cells per time point, the total area of the segments was divided by
the average area of a cell. Puncta per cell was then further calculated.
776
Transmission electron microscopy: SW480 and HT1080 cells treated with vehicle or Fedratinib
for 24 hr were fixed in 3% glutaraldehyde and 3% paraformaldehyde in 0.1 M Cacodylate
buffer (CB; pH 7.2) overnight at 4 °C. Cells were then washed with PBS and centrifuged at
2000 rpm for 2 mins. Pre-warmed 2% agarose solution was then carefully added to islet pellets,
centrifuged and allowed to cool at 4 °C for 30 min. Samples were then subjected to
osmification in 1.5% K4Fe(CN)6 + 2% OsO4 in 0.1 CB for 1 h, dehydrated by serial washes
in EtOH (30%, 50%, 70%, 80%, 90%, 95% and 100%) and embedded in Spurr's resin by
polymerization at 60 °C for 24 h. Polymerized resins were then sectioned at 90 nm thickness
using a Leica EM UC7 ultramicrotome and imaged at 70 kV using a JEOL 1400 TEM equipped
with an AMT CMOS imaging system. Cristae classifications were done by blinding all images,
and then manually annotated with the four cristae classification: laminar, non-laminar,

compartmentalized, and loss-of-cristae. Mitochondrial structures (Aspect Ratio, Perimeter,
Circularity) were analyzed and quantified by Fiji/ImageJ. A minimum of 150 mitochondria are
analyzed from 25 independent cells.
791
<u>Correlative light and electron microscopy (CLEM)</u>: SW480 SPLICS reporter cells were induced
with doxycycline (100 ng/ml) for 24 hr, and then treated with Fedratinib for 24 hr. Cells were
high pressure frozen with Leica EM High Pressure Freezer, freeze substituted using 0.1%
Uranyl acetate in acetone, and finally infiltrated in Lowicryl HM20, following the protocol
published in Ronchi et al., 2021. 300nm sections were cut using a Leica UC7 ultramicrotome
and picked up on carbon coated 200 mesh grid**s**. Fluorescence microscopy imaging of the
sections was carried out as previously described (Kukulski et al., 2012) using a wide-field
Olympus IX81, equipped with an Olympus PlanApo 100X 1.4NA oil immersion objective. After
that, tilt series of the areas of interest containing fluorescent signal were acquired using a
TECNAI F30 (Thermo Fisher Scientific) and tomograms were reconstructed using IMOD
(Kremer et al, 1996). High precision overlays between fluorescence and electron microscopy
images were carried out using the plug in ec-CLEM (Paul-Gilloteaux et al., 2017) of the software
platform Icy (de Chaumont et al., 2012), by clicking on corresponding features in the two
imaging modalities. These articles were cited for reference[56–59].
806
<u>Segmentation for CLEM:</u> Automated detection of membranes in the tomogram was performed
using the TomoSegMemTV (April 2020 version) software package (PMID: 24625523). The
parameters used for membrane enhancement were as follows: scale_space -s 3; dtvoting -s 10
(default); surfaceness -m 0.45 -s 1.0 -p 1.0 (default); dtvoting -w -s 10 (default); surfaceness -S -
m 0.3 (default); thresholding -l 0.05; global analysis –3 100 (default). The output volumes were
imported into AMIRA 2022.2 (Thermo Fisher Scientific), and segmented membrane pixels were
manually annotated as specific organelle membranes using the 3D magic wand tool and saved
as different labels. Further, gaps in the segmentation were filled using the paint tool. The final
membrane surfaces were generated using the 'generate surface' module and smoothened
using the 'smooth surface' module with the following parameters: iterations 100; lambda 0.2.
817
<u>High pressure freezing, FIB-SEM, and data processing:</u> Cells were cultured on optically flat
sapphire disks (3mm diameter, 50-80um thick, Nanjing Co-Energy Optical Crystal Co., Ltd.).
Cell growth media was exchanged 3x for freezing media containing Fluorobrite media
(A1896702; ThermoFisher), 25% Dextran (Mr ~40,000, 31389–100G; Sigma), and 0.8pM
TetraSpeck microspheres (0.2μm diameter, T7280; Invitrogen). They were then immediately
subjected to High Pressure Freezing (HPF Compact 01; Wohlwend GmbH) according to the
manufacturer's instructions. Samples were then stored in liquid N_2. Samples were loaded onto a
custom CryoSIM microscope as described in the following citation at the end of the paragraph.
Epi-fluorescence montage images were taken of each coverslip to select cells for higher-
resolution structured illumination microscopy (SIM). 3D SIM images were acquired (5 phases, 3
angles) over a 7-10um z-depth, and were reconstructed as described in the following citation at
the end of the paragraph. Typical reconstruction parameters included 0.007 Wiener Filter, 0.7
gamma apodization, 15-pixel radii of the singularity suppression at the OTF origins. From SIM
images, candidate cells for FIB-SEM were selected. Samples were subjected to a freeze-
substitution/resin-embedding procedure in 2% OsO_4, 0.1% Uranyl Acetate, and 3% water in
acetone under liquid N_2 using a Leica AFS system. Samples were infiltrated and embedded
with Eponate 12 resin as described in in the following citation at the end of the paragraph. After
coverslips were removed from the resin block (leaving cells embedded), repeated
ultramicrotome trimming (Leica EM UC7) and micro X-ray CT imaging (Zeiss Xradia 510) was
used in conjunction with cryo-fluorescence image registration to locate the cells of interest in the
resin block. Samples were mounted on a copper stud, and sputter coated with 10nm gold and

100nm carbon (PECS 682; Gatan). Samples were loaded onto a custom FIB-SEM, consisting of
a Zeiss Gemini 450 Field Emission SEM and a Zeiss Capella FIB column oriented at 90
degrees to the SEM beam[60].Samples were imaged using 0.9kV landing energy and 0.25nA
electron dose, and images were acquired at 200kHz readout rate (5us dwell time), and 4nm
pixel size. Samples were milled using a 15nA gallium source at 30kV acceleration voltage.
Actual milling time and current per slice was adjusted to maintain average 4nm milling rate,
producing isotropic 4nm datasets. Images slices were registered with respect to each other
using a python implementation of the SITF algorithm[61] with RANSAC solver and a regularized
affine transformation model, available at https://github.com/gleb-shtengel/FIB-SEM. Images
were further de-noised using the noise 2 void 2 (N2V2) algorithm, available
at https://github.com/juglab/n2v. Overall, similar workflow was performed in these cited
publications[60,62,63].
851
Segmentation and analysis of FIBSEM: The ER and mitochondria were segmented using deep-
learning based segmentation software Empanada (EM Panoptic Any Dimension Annotation)
(PMID: 36657391). The pretrained MitoNet model was finetuned for instance-based
segmentation of mitochondria. A new model was trained for ER segmentation. Semantic and
instance-based segmentations were generated for ER and mitochondria, respectively. Instance-
based segmentation allowed assignment of unique identifier to each mitochondrion. The
mitochondrial segmentations were further improved by manually deleting the falsely segmented
objects. The ER segmentations were cleaned by removing unspecific objects less than 2000
pixel2. Following this, the segmentations were further improved by manually deleting larger
false positive objects. The segmentations were binned by a factor of 4 for analysis in Fiji (PMID:
22743772). Mitochondrial volumes and ERMCS distances were calculated using the 3D
manager plugin. For the quantification of ER-mitochondria contact surface area, single-voxel-
thick outer mitochondrial surfaces were generated. Then, the ER objects were expanded by a
distance of 64 nm (4 voxels). The overlapping voxels of the expanded ER objects and
mitochondria surface were marked as the contact surface. The percentage of mitochondrial
contact surface area was calculated by dividing the contact surface area by the total
mitochondrial surface area.
869
Sample preparation for LC-MS/MS metabolomics and lipidomics analysis: Samples for
metabolomics and lipidomics LC-MS/MS analyses were prepared by following the automated
dual-metabolite/lipid sample preparation workflow described in the Agilent application note
5994-5065EN. Briefly, cells (ca. 1M) were collected, washed with PBS and lysed with 1:1
trifluoroethanol/water at room temperature. Lysates were transferred to microcentrifuge tubes,
incubated for 10 minutes, quickly centrifuged at 250 xg for 30 seconds, and dried out by
centrifugation under reduced pressure with no heat. Samples were resuspended with 1:1
trifluoroethanol/water, transferred to a 96-well plate, and processed on a Bravo Metabolomics
sample preparation platform (Agilent Technologies, Inc.) with two separate VWorks protocols to
sequentially and selectively isolate cell metabolites and lipids as described (5994-5065EN).
880
LC-MS/MS metabolomics and lipidomics analysis: Sample were analyzed on an Agilent 1290
Infinity II Bio LC ultra-high performance liquid chromatography (UPLC) system consisting of a
high-pressure binary pump, multicolumn thermostat, and a temperature controlled multisampler.
The LC modules were setup with a standard configuration for omics workflows, which allowed
easy acquisition method interchange for polar metabolites and lipid analyses. Samples for both
targeted metabolomics and lipidomics analysis were analyzed in randomized order on an
Agilent 6495C triple quadrupole mass spectrometer equipped with an Agilent Jet Stream Dual
ESI ion source. For targeted metabolomics, isolated polar metabolites were analyzed with a
HILIC LC-MS/MS method as described in the Agilent application note 5994-5628EN. For

targeted lipidomics analysis, samples were analyzed with the reverse phase LC-MS/MS method
reported in the Agilent application note 5994-3747EN. After acquisition, both metabolomics and
lipidomics datasets were processed with MassHunter Quantitative analysis software and
subsequently imported into Mass Profiler Professional (MPP) for chemometrics analysis.
Metabolomics and lipidomics data were analyzed using MetaboAnalyst. In particular, statistical
analysis [one factor] was selected. Data table with metabolite peak intensities in rows and
samples in column was used. Data were filtered based on interquartile range (IQR) set filter out
25%, normalized based on protein quantitation, transformed with Log base 10, scaled by mean-
centering and dividing by the standard deviation of each variable. Data were visualized with
principal component analysis (PCA) and volcano plots.

Mitochondrial lipidomics: Mitochondria were isolated with the same method detailed prior. Lipid
extraction was performed as described by Bielawski[64] with slight modifications. Samples were
thawed on ice and then samples were extracted with 1.0 mL of IPA:Water:EtOAc (30:10:60,
v:v:v) and internal standard mixture of EquiSPLASH™ LIPIDOMIX® Quantitative Mass Spec
Internal Standard (Avanti Polar Lipids, Birmingham, AL). The extract was vortexed and
sonicated 2 minutes, followed by centrifugation for 10 minutes at 8000 x g at 4 °C. The organic
upper phase was transferred to a new tube. The pellet was re-extracted with an additional 0.5
mL of IPA:Water:EtOAc (30:10:60, v:v:v). The supernatants were combined and placed at -20
°C for 24 h. The supernatants were dried down using a speed vac. The dried sample was
reconstituted in 150 µL of the initial condition of the mobile phase. The suspension was
vortexed for 5 minutes and then centrifuged for 10 minutes at 17000 x g at 4 °C. The
supernatant was transferred to an auto-sampler vial for UHPLC-MS analysis.

Lipid profiling was conducted using a Vanquish UHPLC system with an Orbitrap Fusion Lumos
Tribrid™ mass spectrometer using a H-ESI™ ion source (all Thermo Fisher Scientific, Waltham,
MA) with a Waters ACQUITY UPLC CSH C18 column (150 mm x 1mm, 1.7 µm particle size,
Milford, MA). Solvent A was HPLC grade Water:acetonitrile (40: 60, v:v) with 0.1% formic acid
and 10 mM ammonium formate. Solvent B was HPLC grade isopropanol:acetonitrile (95:5, v:v)
with 0.1% formic acid and 10 mM ammonium formate. The column was maintained at 65 °C and
a flow rate was set at of 110 µL/min. The gradient of the solvent B is 15 % (B) at 0 min, 30 % (B)
at 2 min, 2–2.5 min 48% (B), 2.5–11 min 82 % (B), 11–11.01 min 99 % (B), 11.01–12.95 min 99
% (B), 12.95–13 min 15 % (B), and 13–15 min 15 % (B). Data acquisition was carried out in
positive and negative charge modes, with the ion source spray voltage configured to 4,000 V
and 3,000 V, respectively. The mass spectrometry analysis spanned a scan range of 200 to
1600 m/z for the full scan, and the MS1 resolution was established at 120K at m/z 200. The
AcquireX deep scan mode was employed for the MS2 acquisition, which was performed with a
stepped collision energy (SCE) of 30%, along with a 5% spread for the positive fragment ion
MS/MS scan and with a SCE of 40 % along with a 20 % spread for the negative fragmental
scan.

Raw data was converted into mzML format using Proteowizard mscovert software[65]. MS-DIAL
software (version v5.4.241004)[66] was used for general lipidomics data analysis including
compound identification with LipidBlast[67,68] which is default library in MS-DIAL. The data was
normalized by EquiSPLASH spiked-in internal standards.

Targeted CoQ Measurement by LC-MS/MS: Determination of the CoQ content and redox state
in mammalian cell culture and isolated mitochondria (with described isolation process) was
performed as previously described with modifications. In brief, frozen cell pellets or mitochondria
pellets were resuspended in 200 µL of PBS and added to ice cold extraction solution (200 µL
acidified methanol [0.1% HCl final], 300 µL hexane, with 0.1 µM CoQ_8 internal standard).

Samples were vortexed and centrifuged (5 min, 17000 x g, 4°C) and the top hexane layer was
transferred to a new tube. Extraction was repeated twice before the hexane layers were
combined and dried under argon gas at room temperature. Extracted dried lipids were
resuspended in methanol containing 2 mM ammonium formate and overlaid with argon.
945
LC-MS analysis was performed using a Thermo Vanquish Horizon UHPLC system coupled to a
Thermo Exploris 240 Orbitrap mass spectrometer. For LC separation, a Vanquish binary pump
system (Thermo Fisher Scientific) was used with a Waters Acquity CSH C18 column (100 mm ×
2.1 mm, 1.7 μm particle size) held at 35 °C under 300 μL/min flow rate. Mobile phase A
consisted of 5 mM ammonium acetate in acetonitrile:H_2O (70:30, v/v) with 125 μL/L acetic acid.
Mobile phase B consisted of 5 mM ammonium acetate in isopropanol:acetonitrile (90:10, v/v)
with the same additive. For each sample run, mobile phase B was initially held at 2% for 2 min
and then increased to 30% over 3 min. Mobile phase B was further increased to 50% over 1 min
and 85% over 14 min and then raised to 99% over 1 min and held for 4 min. The column was
re-equilibrated for 5 min at 2% B before the next injection. Five microliters of the sample were
injected by a Vanquish Split Sampler HT autosampler (Thermo Fisher Scientific), while the
autosampler temperature was kept at 4 °C. The samples were ionized by a heated ESI source
kept at a vaporizer temperature of 350 °C. Sheath gas was set to 50 units, auxiliary gas to 8
units, sweep gas to 1 unit, and the spray voltage was set to 3500 V for positive mode and 2500
V for negative mode. The inlet ion transfer tube temperature was kept at 325 °C with 70% RF
lens. For targeted analysis, the MS was operated in parallel reaction monitoring mode with
polarity switching acquiring scheduled, targeted scans to oxidized CoQ_{10} H+ adduct (m/z
863.6912), oxidized CoQ_{10} NH+ adduct (m/z 880.7177), reduced $CoQ_{10}H_2$ H+ adduct (m/z
865.7068), reduced $CoQ_{10}H_2$ NH+ adduct (m/z 882.7334), CoQ_8 H+ adduct (m/z 727.566) and
CoQ intermediates: DMQ_{10} H+ adduct (m/z 833.6806), and $PPHB_{10}$ H- adduct (m/z 817.6504).
MS acquisition parameters include resolution of 15,000, HCD collision energy (30% for positive
mode and stepped 20%, 40%, 60% for negative mode), and 3s dynamic exclusion. Automatic
gain control targets were set to standard mode. The resulting CoQ intermediate data were
processed using TraceFinder 5.1 (Thermo Fisher Scientific). Raw intensity values were
normalized to the CoQ_8 internal standard and protein content as determined by BCA[69].
971
<u>FACS sorted mitochondrial protein proteomics:</u> All samples were lysed in lysis buffer (8M Urea,
4-(2-Hydroxyethyl)-1-piperazinepropanesulfonic acid (EPPS)) with sonication. Samples were
precipitated with SP3 beads and labeled with iodoacetamide, washed and resuspended in
100mM EPPS buffer, pH 8.5 and digested at 37C with trypsin/LysC overnight. The samples
were labeled with TMT Pro and quenched with hydroxylamine. All the samples were combined
and desalted using a 50 mg Sep-Pak cartridge, followed by drying in a speedvac. Samples were
desalted via StageTip and dried with speedvac. Samples were resuspended in 5% formic acid,
and 5% acetonitrile for LC-MS/MS analysis.
980
981 Mass spectrometry data were collected using a Orbitrap Astral mass spectrometer (Thermo
982 Fisher Scientific, San Jose, CA) coupled with Neo Vanquish liquid chromatograph. Peptides
983 were separated on a 110 cm uPAC C18 column (Thermo Fisher Scientific). For each analysis,
984 we loaded ~0.5 μg onto the column. Peptides were separated using a 150 min gradient of 5 to
985 29% acetonitrile in 0.125% formic acid with a flow rate of 300 μL/min.
986
The scan sequence began with an Orbitrap MS^1 spectrum with the following parameters:
resolution 120,000, scan range 350–1350 Th, automatic gain control (AGC) target 200%,
maximum injection time 50ms, RF lens setting 50%, and centroid spectrum data type. FAIMS
was enabled with compensation voltages (CVs): -35V, -45V, -55V, -60V, and -70V using
TopSpeed setting of 1 sec for each CV. We selected the top twenty precursors for MS^2 analysis

which consisted of HCD high-energy collision dissociation with the following parameters: Astral
data acquisition (TMT off), AGC 200%, maximum injection time 15ms, isolation window 1.2 Th,
normalized collision energy (NCE) 36, and centroid spectrum data type. In addition, unassigned
and singly charged species were excluded from MS^2 analysis and dynamic exclusion was set to
90 s.
997
Whole cell proteomics: All samples were labeled with iodoacetamide and resuspended in 100
mM of 4-(2-Hydroxyethyl)-1-piperazinepropanesulfonic acid (EPPS) buffer, pH 8.5 and digested
at 37C with trypsin/LysC overnight. The samples were labeled with TMT Pro and quenched with
hydroxylamine. All the samples were combined and desalted using a 100mg Sep-Pak cartridge,
followed by drying in a speedvac. Samples were fractionated with basic pH reversed phase
(BPRP) high-performance liquid chromatography (HPLC) as described previously[70]. Samples
were desalted via StageTip and dried with a speedvac. Samples were resuspended in 5%
formic acid, and 5% acetonitrile for LC-MS/MS analysis. The mass spectrometry proteomics
data have been deposited to the ProteomeXchange Consortium via the PRIDE partner
repository with the dataset identifier PXD056917.
1008
Liquid chromatography and tandem mass spectrometry: Mass spectrometric data were
collected on an Exploris480 mass spectrometer coupled to a Proxeon NanoLC-1200 UHPLC.
The 100 μm capillary column was packed with 35 cm of Accucore 150 resin (2.6 μm, 150Å;
ThermoFisher Scientific) at a flow rate of 450 nL/min for 90min. The scan sequence began with
an MS1 spectrum (Orbitrap analysis, resolution 60,000, 350-1350 Th, automatic gain control
(AGC) target is set to "standard", maximum injection time set to "auto"). Data were acquired for
150 minutes per analysis. The hrMS2 stage consisted of fragmentation by higher energy
collisional dissociation (HCD, normalized collision energy 32%) and analysis using the Orbitrap
(AGC 300%, maximum injection time 96☐ms, isolation window 0.7 Th, resolution 30,000 with
TurboTMT activated). Data were acquired using the FAIMSpro interface the dispersion voltage
(DV) set to 5,000V with compensation voltages (CVs) set at -40V, -60V, and -80V. The
TopSpeed parameter was set at 1 sec per CV.
1021
Proteomics data analysis: Raw files were searched using the Comet algorithm with a custom
database search engine reported previously[71]. Database searching included human (Homo
Sapiens) entries from UniProt (http://www.uniprot.org, downloaded 2021) with the reversed
sequences, and common contaminants (i.e., keratins, trypsin). Peptides were searched using
the following parameters: 50 ppm precursor mass tolerance; up to 2 missed cleavages; variable
modifications: oxidation of methionine (+15.9949); static modifications: TMTpro (+304.2071) on
lysine and peptide N terminus, carboxyamidomethylation (+57.0215) on cysteines. The protein-
level FDR was determined using the ModScore algorithm where a score of 13 corresponds to
95% confidence in correct localization. TMT reporter ions were used for quantification of peptide
abundance. Isotopic impurities were corrected according to the manufacturer's specifications,
and signal-to-noise (S/N) was calculated. For Astral MS data, samples were processed by
removing peptides with summed S/N lower than 1000 across all channels or isolation specificity
lower than 0.7. Spectra with resolution <45,000 were discarded. For samples acquired with
Exploris 480, peptides with summed S/N lower than 100 across all channels or isolation
specificity lower than 0.7 were excluded. The high confidence peptides were then used to
quantify protein abundance by summing up S/N values for all peptides assigned to the same
protein, and only proteins in the linear quantification range of the instrument were included in
the analysis. The normalization for protein quantification was then performed by adjusting
protein loadings of total sum of S/N values to that of every TMT channel.
1041

<u>Chromatin immunoprecipitation (ChIP)</u> Chromatin immunoprecipitation (ChIP) experiments were
performed using the Ideal ChIP-seq kit for transcription factors (Diagenode) as per the
manufacturer's protocol. 4 million SW480 and HT1080 cells treated with vehicle (DMSO),
thapsigargin or fedratinib were used per ChIP reaction with 8ug of the BRD4 antibody
(Diagenode, Cat# C15410337). Briefly, cells were crosslinked for 10 minutes in a 1%
formaldehyde solution, followed by termination with 1/10th the volume of 1.25M glycine for 5
minutes at room temperature. Following this, the cells were lysed and sonicated (Bioruptor Pico,
Diagenode) to obtain desired chromatin fragments of about 200bp. Sheared chromatin was then
incubated with the BRD4 antibody overnight at 4□°C. Next day, ChIP-DNA was de-crosslinked
and purified using the Diagenode iPure Kit V3, following the manufacturer's protocol. Utilizing
the manufacturer's instructions (Illumina), purified DNA was prepared for sequencing. About 1-
10ng of ChIP DNA were converted to blunt-ended fragments using T4 DNA polymerase,
Escherichia coli DNA polymerase I large fragment (Klenow polymerase), and T4 polynucleotide
kinase (New England BioLabs (NEB)). Klenow fragment (3′ to 5′ exo minus; NEB) was used to
add a single adenine base to fragment ends, followed by ligation of Illumina adaptors (Quick
ligase, NEB). PCR enriched the adaptor-ligated DNA fragments using the Illumina Barcode
primers and Phusion DNA polymerase (NEB). PCR products were size-selected using 3%
NuSieve agarose gels (Lonza) followed by gel extraction using QIAEX II reagents (Qiagen).
Quantified libraries quality-checked using the Bioanalyzer 2100 (Agilent) and sequenced on the
Illumina HiSeq 2500 Sequencer (125-nucleotide read length). Reads were first processed using
Trimmomatic version 0.39 (settings TruSeq3-PE-2.fa:2:30:10, minlen 50) followed by alignment
with bwa ("bwa mem," options -5SP -T0, version 0.7.17-r1198-dirty) to hg38 (GRCh38)
reference[72,73]. After alignment the reads were filtered using Markduplicates from Picard and then
by quality score of >20 via samtools. MACS2 was used to call peaks, filtered using bedtools,
and converted to bigwigs with UCSC wigtoBigwig[74,75]. Cistrome overlap analysis was performed
in R (v3.6.0) using ChipSeekAnno (v3.0.0) and ChipSeeker (v1.29.1)[76,77]. Enrichment heatmaps
were generated using Deeptools.
1069
<u>RNA Extraction, Reverse Transcription, Real-time PCR and Bromouridine nascent mRNA-</u>
<u>Sequencing:</u> RNA was extracted from SW480 and HT1080 cells using the QIAGEN RNeasy
mini kit according to the manufacturer's instructions. 1 ug of RNA was reverse transcribed using
the Lunascript Supermix (NEB) according to the manufacturer's instructions. Quantitative real-
time PCR (qPCR) from cDNA was performed using SYBR green master mix and beta actin
(ACTB) was used as a control. qPCR from BRD4 ChIP-DNA was performed using 1% of input
genomic DNA with two sets of primers targeting MYC PVT enhancer locus, and a control primer
set targeting an upstream region that does not have BRD4 occupancy. The primer sequences
are listed in Supplementary Table 5. BrU-Seq method is described in these papers[78,79].
1079
<u>Reproducibility:</u> Each cell line experiment was performed in technical replicates for each
condition and repeated at least three times with biological triplicates to ensure reproducibility.
Figures show a representative biological replicate unless otherwise indicated. Blinding was
performed whenever appropriate. Sample description and identification was unavailable to the
core personnel during data collection and analysis. Statistical details of all experiments can be
found in the figure legends. The sample numbers are mentioned in each figure legend and
denote biological replicates. Statistical details are reported in figure legends. Results are
expressed as the mean plus or minus the standard error of the mean for all figures unless
otherwise noted. Significance between 2 groups was tested using a 2 tailed unpaired t test.
Significance among multiple groups was tested using a one-way ANOVA. GraphPad Prism 7.0
was used for the statistical analysis. Statistical significance is described in the figure legends as:
∗ $p < 0.05$, ∗∗ $p < 0.01$, ∗∗∗ $p < 0.005$, ∗∗∗∗ $p < 0.001$.
1092

<u>Data availability:</u> Images and details of the image-based drug repurposing screen were
deposited to Zenodo and are publicly available in the repository as part of this record: DOI:
10.5281/zenodo.13963526. Datasets generated are also available from the corresponding
author upon reasonable request. BrU-seq data has been deposited to NCBI GEO under the
accession number: GSE278077. ChIP-seq data has been deposited to NCBI GEO under the
accession number: GSE280900. The mass spectrometry proteomics data have been deposited
to the ProteomeXchange Consortium via the PRIDE partner repository with the dataset identifier
PXD056917. Due to the size of the FIBSEM data, it will be available upon request.
1101
1102 <u>Figure Illustrations:</u> Figure were created using Adobe Illustrator and Biorender.com.
1103
1104 <u>Drug treatment</u>: Small molecule inhibitors used in this studied are listed in the following table.
1105 Concentration of each compound is listed in figure legend other than Fedratinib. Fedratinib was
1106 used at 1 µM for all studies unless indicated differently.
1107

Compound	Company	Catalog number
Fedratinib	MedChem Express	HY-10309
Thapsigargin	Cayman Chemicals	10522
Ruxolitinib	Cayman Chemicals	11609
dBET6	Cayman Chemicals	35385
JQ1	Cayman Chemicals	11187
Actinomycin D	Cayman Chemicals	11421
AS-2863619	Cayman Chemicals	30976
NVP2	Cayman Chemicals	34725
Piericidin A	Cayman Chemicals	15379
TTFA	Cayman Chemicals	15517
Antimycin A	Cayman Chemicals	34799
Myxothiazol	Sigma Aldrich	T5580
Oligomycin	Cayman Chemicals	11341
CCCP	Cayman Chemicals	25458
FG4592	Cayman Chemicals	15294
MitoPQ	Cayman Chemicals	18808
S3QEL2	Cayman Chemicals	18556
Z-VAD-FMK	Cayman Chemicals	14463
PKR-IN-C16	MedChem Express	HY-13977A
Gamitrinib TPP hexafluorophosphate (GTPP)	MedChem Express	HY-102007A

1108
1109 **References:**
1110

1. Eisenberg-Bord, M., Shai, N., Schuldiner, M. & Bohnert, M. A Tether Is a Tether Is a Tether:
Tethering at Membrane Contact Sites. *Dev. Cell* **39**, 395–409 (2016).

2. Scorrano, L. *et al.* Coming together to define membrane contact sites. *Nat Commun* **10**, 1287
(2019).

3. Shiao, Y. J., Balcerzak, B. & Vance, J. E. A mitochondrial membrane protein is required for
translocation of phosphatidylserine from mitochondria-associated membranes to mitochondria.
Biochem. J. **331 (Pt 1)**, 217–23 (1998).

4. Hajnóczky, G., Robb-Gaspers, L. D., Seitz, M. B. & Thomas, A. P. Decoding of cytosolic
calcium oscillations in the mitochondria. *Cell* **82**, 415–424 (1995).

5. Friedman, J. R. *et al.* ER Tubules Mark Sites of Mitochondrial Division. *Science* **334**, 358–362
(2011).

6. Prudent, J. *et al.* MAPL SUMOylation of Drp1 Stabilizes an ER/Mitochondrial Platform
Required for Cell Death. *Mol. Cell* **59**, 941–955 (2015).

7. Lewis, S. C., Uchiyama, L. F. & Nunnari, J. ER-mitochondria contacts couple mtDNA
synthesis with mitochondrial division in human cells. *Science* **353**, aaf5549 (2016).

8. Hamasaki, M. *et al.* Autophagosomes form at ER–mitochondria contact sites. *Nature* **495**,
389–393 (2013).

9. Lee, K.-S. *et al.* Altered ER–mitochondria contact impacts mitochondria calcium homeostasis
and contributes to neurodegeneration in vivo in disease models. *Proc. Natl. Acad. Sci.* **115**,
E8844–E8853 (2018).

10. Arruda, A. P. *et al.* Chronic enrichment of hepatic endoplasmic reticulum–mitochondria
contact leads to mitochondrial dysfunction in obesity. *Nat Med* **20**, 1427–1435 (2014).

11. Çoku, J. *et al.* Reduced ER–mitochondria connectivity promotes neuroblastoma multidrug
resistance. *EMBO J.* **41**, e108272 (2022).

12. Thoudam, T. *et al.* PDK4 Augments ER–Mitochondria Contact to Dampen Skeletal Muscle
Insulin Signaling During Obesity. *Diabetes* **68**, 571–586 (2019).

13. Rodríguez, L. R. *et al.* Oxidative stress modulates rearrangement of endoplasmic reticulum-
mitochondria contacts and calcium dysregulation in a Friedreich's ataxia model. *Redox Biol.* **37**,
101762 (2020).

14. Vallese, F. *et al.* An expanded palette of improved SPLICS reporters detects multiple
organelle contacts in vitro and in vivo. *Nat Commun* **11**, 6069 (2020).

15. Cieri, D. *et al.* SPLICS: a split green fluorescent protein-based contact site sensor for narrow
and wide heterotypic organelle juxtaposition. *Cell Death Differ* **25**, 1131–1145 (2017).

16. Calì, T. *et al.* splitGFP Technology Reveals Dose-Dependent ER-Mitochondria Interface
Modulation by α-Synuclein A53T and A30P Mutants. *Cells* **8**, 1072 (2019).

17. Giamogante, F. *et al.* Stable Integration of Inducible SPLICS Reporters Enables Spatio-
Temporal Analysis of Multiple Organelle Contact Sites upon Modulation of Cholesterol Traffic.
Cells **11**, 1643 (2022).

18. Mishra, P., Carelli, V., Manfredi, G. & Chan, D. C. Proteolytic Cleavage of Opa1 Stimulates
Mitochondrial Inner Membrane Fusion and Couples Fusion to Oxidative Phosphorylation. *Cell*
Metab **19**, 630–641 (2014).

19. Bravo, R. *et al.* Increased ER–mitochondrial coupling promotes mitochondrial respiration
and bioenergetics during early phases of ER stress. *J Cell Sci* **124**, 2143–2152 (2011).

20. Calì, T. & Brini, M. Quantification of organelle contact sites by split-GFP-based contact site
sensors (SPLICS) in living cells. *Nat. Protoc.* **16**, 5287–5308 (2021).

21. Hertlein, V. *et al.* MERLIN: a novel BRET-based proximity biosensor for studying
mitochondria–ER contact sites. *Life Sci Alliance* **3**, e201900600 (2019).

22. Cole, N. B. & Lippincott-Schwartz, J. Organization of organelles and membrane traffic by
microtubules. *Curr. Opin. Cell Biol.* **7**, 55–64 (1995).

23. Talpaz, M. & Kiladjian, J.-J. Fedratinib, a newly approved treatment for patients with
myeloproliferative neoplasm-associated myelofibrosis. *Leukemia* **35**, 1–17 (2021).

24. Wernig, G. *et al.* Efficacy of TG101348, a Selective JAK2 Inhibitor, in Treatment of a Murine
Model of JAK2V617F-Induced Polycythemia Vera. *Cancer Cell* **13**, 311–320 (2008).

25. Hantschel, O. Unexpected Off-Targets and Paradoxical Pathway Activation by Kinase
Inhibitors. *ACS Chem. Biol.* **10**, 234–245 (2015).

26. Ciceri, P. *et al.* Dual kinase-bromodomain inhibitors for rationally designed
polypharmacology. *Nat Chem Biol* **10**, 305–312 (2014).

27. Kleppe, M. *et al.* Dual Targeting of Oncogenic Activation and Inflammatory Signaling
Increases Therapeutic Efficacy in Myeloproliferative Neoplasms. *Cancer Cell* **33**, 29-43.e7
(2018).

28. Sakamaki, J. *et al.* Bromodomain Protein BRD4 Is a Transcriptional Repressor of Autophagy
and Lysosomal Function. *Mol. Cell* **66**, 517-532.e9 (2017).

29. Hinton, A. *et al.* A Comprehensive Approach to Sample Preparation for Electron Microscopy
and the Assessment of Mitochondrial Morphology in Tissue and Cultured Cells. *Adv. Biol.* **7**,
e2200202 (2023).

30. Venkatraman, K. *et al.* Cristae formation is a mechanical buckling event controlled by the
inner mitochondrial membrane lipidome. *EMBO J.* e114054 (2023)
doi:10.15252/embj.2023114054.

31. MacDonald, J. A. *et al.* A nanoscale, multi-parametric flow cytometry-based platform to
study mitochondrial heterogeneity and mitochondrial DNA dynamics. *Commun. Biol.* **2**, 258
(2019).

32. Baik, A. H. & Jain, I. H. Turning the Oxygen Dial: Balancing the Highs and Lows. *Trends*
Cell Biol. **30**, 516–536 (2020).

33. Spinelli, J. B. *et al.* Fumarate is a terminal electron acceptor in the mammalian electron
transport chain. *Science* **374**, 1227–1237 (2021).

34. Geng, J. *et al.* Deregulation of ER-mitochondria contact formation and mitochondrial calcium
homeostasis mediated by VDAC in fragile X syndrome. *Dev. Cell* **58**, 597-615.e10 (2023).

35. Anastasia, I. *et al.* Mitochondria-rough-ER contacts in the liver regulate systemic lipid
homeostasis. *Cell Rep.* **34**, 108873 (2021).

36. Hofstadter, W. A. *et al.* Infection-induced peripheral mitochondria fission drives ER
encapsulations and inter-mitochondria contacts that rescue bioenergetics. *Nat. Commun.* **15**,
7352 (2024).

37. Poggio, E. *et al.* Perturbation of the host cell Ca2+ homeostasis and ER-mitochondria
contact sites by the SARS-CoV-2 structural proteins E and M. *Cell Death Dis.* **14**, 297 (2023).

38. Filippakopoulos, P. *et al.* Selective inhibition of BET bromodomains. *Nature* **468**, 1067–1073
(2010).

39. Barrow, J. J. *et al.* Bromodomain Inhibitors Correct Bioenergetic Deficiency Caused by
Mitochondrial Disease Complex I Mutations. *Mol. Cell* **64**, 163–175 (2016).

40. Costa-Mattioli, M. & Walter, P. The integrated stress response: From mechanism to disease.
Science **368**, (2020).

41. Kalkavan, H. *et al.* Sublethal cytochrome c release generates drug-tolerant persister cells.
Cell **185**, 3356-3374.e22 (2022).

42. Palchaudhuri, R. *et al.* A Small Molecule that Induces Intrinsic Pathway Apoptosis with
Unparalleled Speed. *Cell Rep.* **13**, 2027–2036 (2015).

43. Chipuk, J. E. *et al.* Sphingolipid Metabolism Cooperates with BAK and BAX to Promote the
Mitochondrial Pathway of Apoptosis. *Cell* **148**, 988–1000 (2012).

44. Verfaillie, T. *et al.* PERK is required at the ER-mitochondrial contact sites to convey
apoptosis after ROS-based ER stress. *Cell Death Differ* **19**, 1880–1891 (2012).

45. Csordás, G. *et al.* Structural and functional features and significance of the physical linkage
between ER and mitochondria. *J Cell Biology* **174**, 915–921 (2006).

46. Bassot, A. *et al.* The endoplasmic reticulum kinase PERK interacts with the oxidoreductase
ERO1 to metabolically adapt mitochondria. *Cell Rep.* **42**, 111899 (2023).

47. Carreras-Sureda, A. *et al.* Non-canonical function of IRE1α determines mitochondria-
associated endoplasmic reticulum composition to control calcium transfer and bioenergetics.
Nat Cell Biol **21**, 755–767 (2019).

48. Stefely, J. A. & Pagliarini, D. J. Biochemistry of Mitochondrial Coenzyme Q Biosynthesis.
Trends Biochem. Sci. **42**, 824–843 (2017).

49. Bersuker, K. *et al.* The CoQ oxidoreductase FSP1 acts parallel to GPX4 to inhibit ferroptosis.
Nature **575**, 688–692 (2019).

50. Doll, S. *et al.* FSP1 is a glutathione-independent ferroptosis suppressor. *Nature* **575**, 693–
698 (2019).

51. Jones, M. E. Pyrimidine Nucleotide Biosynthesis in Animals: Genes, Enzymes, and
Regulation of UMP Biosynthesis. *Annu. Rev. Biochem.* **49**, 253–279 (1980).

52. Kumar, R. *et al.* A redox cycle with complex II prioritizes sulfide quinone oxidoreductase-
dependent H2S oxidation. *J. Biol. Chem.* **298**, 101435 (2022).

53. Guerra, R. M. & Pagliarini, D. J. Coenzyme Q biochemistry and biosynthesis. *Trends*
Biochem Sci **48**, 463–476 (2023).

54. Williams, R. T. *et al.* ZBTB1 Regulates Asparagine Synthesis and Leukemia Cell Response
to L-Asparaginase. *Cell Metab.* **31**, 852-861.e6 (2020).

55. Chan, S. W. & Chen, J. Z. Measuring mtDNA damage using a supercoiling-sensitive qPCR
approach. *Methods Mol. Biol. (Clifton, NJ)* **554**, 183–97 (2009).

56. Kukulski, W. *et al.* Chapter 13 Precise, Correlated Fluorescence Microscopy and Electron
Tomography of Lowicryl Sections Using Fluorescent Fiducial Markers. *Methods Cell Biol.* **111**,
235–257 (2012).

57. Kremer, J. R., Mastronarde, D. N. & McIntosh, J. R. Computer Visualization of Three-
Dimensional Image Data Using IMOD. *J. Struct. Biol.* **116**, 71–76 (1996).

58. Paul-Gilloteaux, P. *et al.* eC-CLEM: flexible multidimensional registration software for
correlative microscopies. *Nat. Methods* **14**, 102–103 (2017).

59. Chaumont, F. de *et al.* Icy: an open bioimage informatics platform for extended reproducible
research. *Nat. Methods* **9**, 690–696 (2012).

60. Xu, C. S. *et al.* Enhanced FIB-SEM systems for large-volume 3D imaging. *Elife* **6**, e25916
(2017).

61. Lowe, D. G. Distinctive Image Features from Scale-Invariant Keypoints. *Int. J. Comput. Vis.*
60, 91–110 (2004).

62. Hoffman, D. P. *et al.* Correlative three-dimensional super-resolution and block-face electron
microscopy of whole vitreously frozen cells. *Science* **367**, (2020).

63. Bharathan, N. K. *et al.* Architecture and dynamics of a desmosome–endoplasmic reticulum
complex. *Nat. Cell Biol.* **25**, 823–835 (2023).

64. Bielawski, J. *et al.* Lipidomics, Volume 1: Methods and Protocols. *Methods Mol. Biol.* **579**,
443–467 (2009).

65. Chambers, M. C. *et al.* A cross-platform toolkit for mass spectrometry and proteomics. *Nat.*
Biotechnol. **30**, 918–920 (2012).

66. Tsugawa, H. *et al.* MS-DIAL: data-independent MS/MS deconvolution for comprehensive
metabolome analysis. *Nat. Methods* **12**, 523–526 (2015).

67. Kind, T. *et al.* LipidBlast in silico tandem mass spectrometry database for lipid identification.
Nat. Methods **10**, 755–758 (2013).

68. Tsugawa, H. *et al.* A lipidome atlas in MS-DIAL 4. *Nat. Biotechnol.* **38**, 1159–1163 (2020).

69. Burger, N. *et al.* A sensitive mass spectrometric assay for mitochondrial CoQ pool redox
state in vivo. *Free Radic. Biol. Med.* **147**, 37–47 (2020).

70. Liu, X., Gygi, S. P. & Paulo, J. A. A Semiautomated Paramagnetic Bead-Based Platform for
Isobaric Tag Sample Preparation. *J. Am. Soc. Mass Spectrom.* **32**, 1519–1529 (2021).

71. Huttlin, E. L. *et al.* A Tissue-Specific Atlas of Mouse Protein Phosphorylation and Expression.
Cell **143**, 1174–1189 (2010).

72. Bolger, A. M., Lohse, M. & Usadel, B. Trimmomatic: a flexible trimmer for Illumina sequence
data. *Bioinformatics* **30**, 2114–2120 (2014).

73. Li, H. & Durbin, R. Fast and accurate short read alignment with Burrows–Wheeler transform.
Bioinformatics **25**, 1754–1760 (2009).

74. Zhang, Y. *et al.* Model-based Analysis of ChIP-Seq (MACS). *Genome Biol.* **9**, R137 (2008).

75. Raney, B. J. *et al.* The UCSC Genome Browser database: 2024 update. *Nucleic Acids Res.*
52, D1082–D1088 (2023).

76. Zhu, L. J. *et al.* ChIPpeakAnno: a Bioconductor package to annotate ChIP-seq and ChIP-
chip data. *BMC Bioinform.* **11**, 237 (2010).

77. Yu, G., Wang, L.-G. & He, Q.-Y. ChIPseeker: an R/Bioconductor package for ChIP peak
annotation, comparison and visualization. *Bioinformatics* **31**, 2382–2383 (2015).

78. Paulsen, M. T. *et al.* Use of Bru-Seq and BruChase-Seq for genome-wide assessment of
the synthesis and stability of RNA. *Methods* **67**, 45–54 (2014).

79. McShane, A. *et al.* Characterizing nascent transcription patterns of PROMPTs, eRNAs, and
readthrough transcripts in the ENCODE4 deeply profiled cell lines. *bioRxiv* 2024.04.09.588612
(2024) doi:10.1101/2024.04.09.588612.

1280

1281

Acknowledgments: We thank all members of the Shah and Lyssiotis labs for their advice and
suggestions. In collaboration with this research, we acknowledge support from the University of
Michigan Biomedical Research Core Facilities (Flow Cytometry core, Advanced Genomics core,
Microscopy core), and Bru-seq Lab. FIB-SEM imaging were performed in collaboration with the
Advanced Imaging Center (AIC) at Janelia Research Campus, which is supported by the
Howard Hughes Medical Institute. The funders (Agilent Technologies) for the whole cell
metabolomics and lipidomics had no role in study design, data collection and analysis, or the
content and publication of this manuscript. We acknowledge the European Molecular Biology
Laboratory (EMBL) Electron Microscopy Core Facilities' assistance, data acquisition, and
analysis for CLEM expertise. B.C was supported by NCI F99CA284256-01. D.C.S. was
supported by NIGMS 5T32GM145304-03. Y.M.S. was supported by NCI R01CA148828,
R01CA245546, and NIDDK R01DK095201. C.A.L. was supported by NCI R37CA237421,
R01CA248160, and R01CA244931. S.M. was supported by NIGMS DP2GM150019. D.J.P. was
supported by NIDDK R01DK098672 and NIGMS R35GM131795. N.J.R. was supported by
NIGMS T32GM145470 and T32GM150581. R.B. was supported by NIGMS R35GM130183.
D.A.H. was supported by NIGMS F32GM140694. A.D.P. was supported by NIH S10OD021750,
USDA National Institute of Food and Federal Appropriations Project PEN04917, Accession
7006712 and Pennsylvania Department of Health using Tobacco CURE funds. T.C was
supported by Italian Ministry of University and Research PRIN2017, University of Padova
STARS Consolidator Grant 2019 and Progetto di Ateneo 2023 no. CALI_BIRD23_01, and
PNRR – CN3 National Center for Gene Therapy and Drugs based on RNA Technology n.
CN00000041 (2022-26).

1304
Author contributions: B.C., C.A.L., and Y.M.S. designed the study and wrote the manuscript.
B.C. designed experiments and collected data for the bulk of the experimental studies. B.C.,
D.C.S, P.J., T.M.L., B.S.H., N.J.R., N.S, M.S., J.A.P, M.C., M. P., S.K., I.K., S.E., P.R, P.M.,
D.A.H., J.L. R.M.G., and carried out experimental aspects of the project, and contributed to data
analysis. B.C., S.M., D.J.P., N.J.R., R.B., D.A.H., M. L., A.P., J.D.M., T.C., C.A.L., and Y.M.S.
provided resources, funding, and conceptual input for experiments and supervised the research.
All authors are involved in the throughout the research process, agreed amongst authors
regarding roles and responsibilities, and contributed to review, editing, and approval of the
manuscript.

1314
Conflicting interests: In the past three years, C.A.L. has consulted for Astellas
Pharmaceuticals, Odyssey Therapeutics, Third Rock Ventures, and T-Knife Therapeutics, and
is an inventor on patents pertaining to Kras regulated metabolic pathways, redox control
pathways in pancreatic cancer, and targeting the GOT1-ME1 pathway as a therapeutic
approach (US Patent No: 2015126580-A1, 05/07/2015; US Patent No: 20190136238,
05/09/2019; International Patent No: WO2013177426-A2, 04/23/2015). J.D.M. reports research
support from Novartis and Casma Therapeutics and has consulted for Third Rock Ventures and
Skyhawk Therapeutics, all unrelated to the submitted work.

1323
1324   **Materials & Correspondence:** Supplementary Information is available for this paper. All data
1325   generated or analyzed during this study are included in this published article and its
1326   supplementary information files. Correspondence and requests for materials should be
1327   addressed to Yatrik M. Shah (shahy@umich.edu) or Costas A. Lyssiotis (clyssiot@umich.edu).
1328

1329
Fig. 1: High throughput image-based chemical screen identifies novel regulators of
ERMCS.
a, Schematic of the SPLICS and mitochondria TagRFP (mitoTagRFP) reporter and analysis
pipeline for ERMCS. **b**, Relative fluorescence intensity of SPLICS, mitoTagRFP, and
SPLICS/mitoTagRFP ratio following treatment with vehicle or thapsigargin (50 nM) for 24 hr, or
c, 48 hr post transfection of ER-MT tether. **d**, D. Schematic representation of the bioassay
design for the ERMCS drug screen. Blue: Hoechst 33342/nuclei. Red: MitotagRFP/mitochondria.
Green: SPLICS/ERMCS. Clockwise: 1. Negative control demonstrating low levels of ERMCS; 2.
Cellpose nuclear segmentation using Hoechst counterstain; 3. Positive control demonstrating
induction of high levels of ERMCS. 4. GFP spot identification and feature extraction using
Cellprofiler for ERMCS. Machine learning was performed by training an XGBoost model against
per-plate controls. 5. Shapley additive explanations showing GFP features used to identify hits

in drug screening. Beeswarm plot showing SHAP values indicating feature importance for drug
effect scoring based on GFP features. 6. Drug screening scatter plot showing drug tested on the
x-axis and XGBoost score on the y-axis indicating ERMCS induction. Each color categories for
individual data points suggest different plates instead of categorization of drug pathways.
Fedratinib is highlighted in red on the scatter plot. Scale bar: 15 µm. **e**, Relative ERMCS in a
panel of cell lines treated with vehicle or fedratinib treatment for 24 hr. **f**, Relative ERMCS of
cells treated with vehicle or fedratinib for 24 hr, followed by PBS washout, and monitored for
additional 48 hr. **g**, Dose dependent increase of relative ERMCS of cells treated with vehicle or
fedratinib for 24 hr. **h,** Time-lapse imaging using Lattice light sheet microscopy to monitor
ERMCS puncta of cells treated with fedratinib for 16 hr with quantified puncta per cell.
Experiment (**b, c, e, f, h**) were performed three times, and experiment (**d**) was performed once,
and experiment (**g**) was performed twice. Significance was calculated with an unpaired two-
tailed t-test. *P ≤ 0.05, **P ≤ 0.01, ***P ≤ 0.005, ****P ≤ 0.001. Data are mean ± SD. Statistical
source data are provided in Source Data Fig. 1.
1356

Fig. 2: ERMCS induction requires a transcriptional response dependent on bromodomain
and extra-terminal protein (BET).
a, Mechanisms of action for fedratinib, ruxolitinib, and dBET6 targeting JAK2 and BRD4 (Left).
Relative ERMCS in cells treated with JAK2 and/or BRD4 inhibitors. **b**, Relative ERMCS and
BRD4 mRNA expression of cells following knockdown (KD) using scrambled or BRD4 targeting
siRNAs. **c**, Relative ERMCS in cells treated with inhibitors of transcription kinase CDK8 (AS-
2863619, 1 μM) or CDK9 (NVP2, 1 μM) for 6 hr, and then treated with vehicle or fedratinib for
24 hr. **d**, Representative BrU-seq transcript profile of SW480 cells treated with fedratinib or
vehicle for 30 min. **e**, Schematic of a fluorescence-activated cell sorting (FACS)-based CRISPR
screen using a nuclear genome targeted sgRNA library. **f**, Volcano plot of BrU-seq analysis after
SW480 cells treated with fedratinib for 2 hr (top panel). X-axis is the CRISPR gene scores (delta
Z score) with Y-axis being -Log$_{10}$ of false discovery rate (FDR). Highlighted red circles are
sgRNAs targeting the Mediator complex components. **g**, Histogram indicating the distribution of
sgRNA. Shown below is the relative enrichment of mediator complex genes. **h**, GSEA ridgeplot
displaying significantly enriched (FDR < 0.01) protein complexes based on differential CRISPR
screen scores.) BP: Biological processes. CC: Cellular components. **i**, Schematic of the
mediator complex and BRD4 in transcriptional regulation. Mediator components from the
CRISPR library, annotated in red demonstrate suppression of ERMCS. **j**, Relative ERMCS and
gene expression level of MED16 and MED24 mRNA transcripts upon CRISPRi mediated
knockdown of cells treated with vehicle or fedratinib for 24 hr. **k**, Relative MED16 and MED24

expression of sgMED16 and sgMED24 cells. Experiments. **l**, Relative ERMCS in cells treated
with inhibitors of transcription kinase CDK8 (AS-2863619, 1 µM) or CDK9 (NVP2, 1 µM) for 6 hr,
and then treated with vehicle or thapsigargin (50 nM) for 24 hr. (**b**, **c**, **j-l**) were performed three
times, experiment (**d**) was performed once with duplicates, and experiment (**e-h)** was conducted
once. Significance was calculated with an unpaired two-tailed t-test. n.s. = not significant, *P ≤
0.05, **P ≤ 0.01, ***P ≤ 0.005, ****P ≤ 0.001. Data are mean ± SD. Statistical source data are
provided in Source Data Fig. 2.
1386

1387
Fig. 3: Fedratinib induces ER-wrapping of mitochondria at ERMCS.
a, Workflow of our multi-modal electron microscopy imaging approach. **b**, Representative TEM
images (scale bar: 400 nm) of WT cells treated with vehicle or fedratinib for 24 hr (Mitochondria

green, ER: magenta (Above)). Distribution of cristae morphologies quantified from 25 cells as a
percentage of total mitochondria (n=161-299) (Below). Cristae structures are annotated in
corresponding colors in the stacked bars with percentage of each classification listed. Red:
Laminar cristae. Blue: Non-laminar cristae. Yellow: Compartmentalized cristae. Green: Loss of
cristae. Each example cristae structure is indicated with its corresponding color in the EM
images. **c**, Correlative light and electron microscopy (CLEM) images of SW480 cells treated
with fedratinib for 24 hr (Scale bar: 500 nm). White arrows indicate an example of SPLICS[Hi]
area. **d**, Segmentation of organelle membranes of SW480 cells treated with fedratinib for 24 hr
(M: mitochondria, ER: endoplasmic reticulum, EL: endolysosome, G: golgi apparatus; Scale bar:
500 nm). **e**, Representative slices through the tomograms of vehicle control (top left) and
fedratinib-treated (bottom left) HT1080 cells with an overlay of 3D segmentation. Inset highlights
a representative mitochondrion in each case. Scale bar, 200 nm and 50 nm for inset. 3D volume
reconstructions(right) displaying mitochondria. Scale bars, 1 μm. **f**, FIB-SEM images showing
instances of different morphologies of mitochondria wrapped around by ER membrane (red
arrowheads). Scale bar, 200 nm. **g**, Quantification of total mitochondrial volume. **h**, ER-
mitochondria contact site intermembrane distances. **i**, ER-mitochondria contact surface area. **j**,
Percentage contact surface area for individual mitochondria in vehicle-treated and fed-treated
cells. Experiments (**a-j)** were performed once. Red rectangle highlights area with cristae
collapse. Statistical source data are provided in Source Data Fig. 3.
1410
1411

Fig. 4: Metabolic and lipidomic changes associated with selective depolarization of ER-wrapped mitochondria.

a, Confocal images and, **b**, line intensity profile of representative SPLICS^{Hi} and SPLICS^{Lo} mitochondria of mitochondria membrane potential (MitoTracker Deep Red (MTDR)) and total mitochondrial mass (mitoTagRFP) in cells treated with fedratinib for 24 hr. **c**, Ratio metric analyses of mitochondria membrane potential from SPLICS^{Hi} and SPLICS^{Lo} mitochondria in the same cell. Cells treated with fedratinib for 24 hr (n = 16; Scale bar: 1 µm). Each dot indicates one cell that was imaged. **d**, Diagram describing the FACS analysis of mitochondria, and mitochondria flow cytometry performed on whole cell (n = 10,000) or isolated mitochondrial (n = 100,000) fraction from cells treated with vehicle or fedratinib for 24 hr. $\log_{10}$-scaled MTDR mean fluorescence intensity (MFI) displayed on x-axis and count on the y-axis. **e**, Principal component analysis (PCA) of metabolomic analysis of cells treated with vehicle or fedratinib for 24 hr. **f**, Volcano plot of mitochondrial lipidome profiles in SW480 and HT1080 cells treated with fedratinib for 24 hr. **g**, Volcano plot of mitochondrial PtdCho species highlighted in green in cells treated with fedratinib for 24 hr. Experiments (**a-d**) were repeated three times. Experiment (**e**) was conducted in biological triplicates. Experiments (**f** and **g**) were conducted once with four biological replicates. Two-tailed paired t-test were performed on (E). **P ≤ 0.01, ***P ≤ 0.005. Statistical source data are provided in Source Data Fig.4.

1432
1433
Fig. 5: Fedratinib induces subcellular proteomic changes specific to ER-enveloped
mitochondria.
a, Distribution of SPLICS[Hi] and SPLICS[Lo] mitochondria in fedratinib and vehicle treated cells. **b**,
Scatter plot of individual mitochondria from fedratinib and vehicle treated cells with x-axis as

SPLICS MFI and y-axis as forward scatter (FSC). **c**, Scatter plot of sorted mitochondria from
fedratinib and vehicle treated cells with x-axis as SPLICS MFI and y-axis as FSC. **d**, Schematic
of mitochondrial flow cytometry and sorting for proteomics. **e**, Histograms of pre- and post-
sorted mitochondria from fedratinib and vehicle treated cells. **f**, Distribution of SPLICS[Hi] and
SPLICS[Lo] mitochondria in pre- and post-sorted, fedratinib-treated cells. **g**, Subcellular proteomic
distribution of ER, mitochondria, and other compartmental proteins in vehicle-treated whole cell
and sorted mitochondria. **h**, Subcellular proteomic distribution of ER, mitochondria, and other
compartmental proteins in fedratinib-treated whole cell. **i**, Subcellular proteomic distribution of
ER, mitochondria, and other compartmental proteins in sorted vehicle-treated mitochondria,
SPLICS[Hi] mitochondria, and SPLICS[Lo] mitochondria. **j**, Subcellular proteomic distribution of ER,
mitochondria, and other compartmental proteins comparing SPLICS[Hi] to SPLICS[Lo] mitochondria.
k, GSEA analysis of SPLICS[Hi] mitochondria. **l**, Volcano plot showing enrichment of proteins
involved in: oligosaccharide transferase (OST) complex, fatty acid oxidation (FAO), voltage
dependent anion channel (VDAC), and reductive proline biosynthesis. Experiments (**a-c**, **e**, **f**)
were performed three times, experiments (**g-l**) were performed once with four biological
replicates. Statistical source data are provided in Source Data Fig. 5.
1454

Fig. 6: ETC Complex III function is necessary for ERMCS formation.
a, Relative ERMCS (right) in cells incubated in hypoxia (2% O_2) or treated with a PHD inhibitor
FG-4592 (50 µM) for 48 hr, and then treated with fedratinib for 24 hr. **b**, Diagram of
mitochondria electron transport chain (ETC) and specific mitochondrial complex inhibitors (CI:
piericidin A (20 nM), CII: TTFA (100 µM), CIII (antimycin A: 100 nM, myxothiazol: 50 nM), CIV
(Hypoxia: 2% O_2, H_2S: 250 ppm)). **c**, Relative ERMCS (right) in cells treated with specific ETC

inhibitors for 24 hrs, and then treated with vehicle or fedratinib for 24 hr. **d**, Diagram illustrating
the regulation of CoQ redox state following complex III inhibition, expression of alternative
oxidase (AOX) under complex III inhibition, and complex I/III dual inhibition. **e**, Volcano plot of
mitochondrial lipidome profiles comparing fedratinib (grey circle) and fedratinib plus antimycin A
combination treatment (yellow dot). **f**, Volcano plot of mitochondrial PtdCho species in fedratinib
(grey circle) and fedratinib plus antimycin A combination treatment (yellow dot). **g**, Relative
ERMCS in AOX expressed cell treated with complex III inhibition with myxothiazol (50 nM) or
antimycin A (100 nM) for 24 hr, and then treated with vehicle or fedratinib for 24 hr. AOX
expression was validated with immunoblotting (below). **h**, Relative ERMCS in cells treated with
complex I and III inhibitors Piericidin A (20 nM) and myxothiazol (50 nM), or antimycin A (100
nM) for 24 hr, and then treated with vehicle or fedratinib for 24 hr. Experiments (**a**, **c**, **g**, **h**) were
performed three times. Experiments (**e** and **f**) were performed once with four biological
replicates. Significance was calculated with an unpaired two-tailed t-test. n.s. = not significant,
**P ≤ 0.01, ***P ≤ 0.001, ****P ≤ 0.001. Data are mean ± SD. Statistical source data are
provided in Source Data Fig. 6. Immunoblot source data are provided in Supplemental Figure 1.
1478

1479
1480
Extended Data Figure 1: Control experiments for validating SPLICS reporter and
orthogonal validation.
a, Cell proliferation data of SPLICS reporter cells treated with doxycycline to induce reporter
expression. **b**, ECAR measurement in SPLICS cells treated with 24, 48, and 72 hr of

doxycycline to induce reporter expression. **c**, OCR measurement in SPLICS cells treated with
24, 48, and 72 hr of doxycycline to induce reporter expression. **d**, Schematic of control pairs for
SPLICS. 1. mKate2 tagged $beta_{11}$ + cytosolic GFP_{1-10} 2. mKate2 tagged $beta_{11}$ + OMM-GFP_{1-10}
3. ER-$beta_{11}$ + cytosolic GFP_{1-10} 4. ER-$beta_{11}$ + OMM-GFP_{1-10}. **e**, Relative fluorescence of
SPLICS and control pairs upon thapsigargin treatment. **f**, Relative fluorescence of SPLICS and
control pairs upon fedratinib treatment. **g**, Relative fluorescence of SPLICS and control pairs
upon Antimycin A treatment. **h**, Representative images of PLA experiment in HT1080 cells
treated with vehicle or fedratinib for 24 hr. Anti-TOM20 and Anti-IP3R were used as primary
antibodies in the assay (Left). Quantification of average number of PLA foci per cell (Right). 10
cells were quantified per condition per independent experiment. Experiments (**a-c**, **e**) were
performed 3 times, and experiment (**g**) was performed twice. n = 3. n.s. = not significant, *P ≤
0.05, **P ≤ 0.01, ***P ≤ 0.005, ****P ≤ 0.001. Data are mean ± SD. Statistical source data are
provided in Source Data Extended Data Fig. 2.
1498

Extended Data Figure 2: Transcriptional and epigenetic regulation of ERMCS.
a, **b** Immunoblot from cells treated with vehicle, fedratinib (Fed), ruxolitinib (Ruxo), or dBET6 for 24 hr. **c**, Immunoblot of JAK2 KD with CRISPRi in SW480 and HT1080 cells. **d**, Relative ERMCS of control and JAK2 CRISPRi-KD cells. **e**, Chromatin fractionation assay of cells treated with JQ1 (1 µM), fedratinib, and thapsigargin (50 nM) for 24 hr. **f**, Venn diagram showing overlaps of BRD4 ChIP-seq in thapsigargin (50nM) vs fedratinib-treated SW480 cells. **g**, ChIP-seq read density heatmaps of BRD4 at promoter and non-promoter sites in thapsigargin (50 nM) vs fedratinib-treated SW480 cells. **h**, Genomic locations of BRD4 bound sites defined from the overlap analysis in panel g. **i**, BRD4 ChIP-qPCR analysis of BRD4-bound PVT1 enhancer regions in HT1080 cells treated with fedratinib or thapsigargin (50 nM) for 4 hr. **j**, Relative ERMCS in cells treated with inhibitors of transcription kinase CDK8 (AS-2863619, 1 µM) or CDK9 (NVP2, 1 µM) for 6 hr, and then treated with vehicle or thapsigargin (50 nM) for 24 hr. **k**, Actinomycin D (10 µg/ml) pulse chase of mitoTagRFP and SPLICS fluorescence after vehicle, fedratinib, or thapsigargin (50nM) treatment. **l**, Relative ERMCS fluorescence from cells treated with actimonycin D (10 µg/ml) for 24 hr, and then treated with vehicle, fedratinib, or thapsigargin (50 nM) for 24 hr. **m**, Volcano plot of BrU-seq analysis after SW480 cells were treated with fedratinib for 30 min. **n**, GSEA of BrU-seq analysis after SW480 cells were treated with fedratinib for 30 min. Normalized enrichment score (NES) of top pathways altered. **o**, Relative ERMCS and gene expression level of MED16 and MED24 mRNA transcripts upon CRISPRi mediated knockdown of cells treated with vehicle or thapsigargin (50 nM) for 24 hr. Experiments (**a-f**, **j-l, o**) were performed three times, and experiment (**g-i**, **m**, **n**) was performed once with biological duplicates. Significance was calculated with an unpaired two-tailed t-test. n.s. = not significant, *P ≤ 0.05, **P ≤ 0.01, ***P ≤ 0.005, ****P ≤ 0.001. Data are mean ± SD. Statistical source data are provided in Source Data Extended Data Fig. 2. Immunoblot source data are provided in Supplemental Figure 2.

Extended Data Figure 3. Fedratinib induces ISR and PKR supports ERMCS induction.

a, BrU-Seq volcano plot of SW480 cells treated with fedratinib for 2 hr with ISR target genes
highlighted. **b**, GSEA analysis of BrU-seq analysis after SW480 cells were treated with
fedratinib for 2 hr with ISR target genes highlighted. Normalized enrichment score (NES) of top
pathways altered. **c**, Proteomic volcano plot of SW480 cells treated with fedratinib for 24 hr with
ISR target genes highlighted. **d**, Immunoblot of ISR targets from cells treated with fedratinib and
thapsigargin (50 nM) for 24 hr. **e**, Infographic of ISR delineating the ISR kinases and
phosphatases that mediate ATF4-dependent response. **f**, Relative ERMCS and ATF4
immunoblot of cells pretreated with ISRIB (1 µM) for 6 hr and then treated with fedratinib or
thapsigargin (50 nM) for another 18 hr. **g**, Relative ERMCS and ATF4 immunoblot of cells
treated with fedratinib and ISR inducers (CT020312, Histidinol, BEPP, BTdCPU, Salubrinal. **h**,
Relative ERMCS induction and **i**, immunoblot in cells pretreated with PKR inhibitor (PKR-IN-C16,
1 µM) and caspase inhibitor Z- VAD-FMK (10 µM) for 6 hr then treated with fedratinib or
thapsigargin (50 nM) for another 18 hr. **j**, Relative expression of UPRmito genes HSPD1 and
HSPE1 upon fedratinib or GTPP (10 µM). **k**, Comparison of ATF4 induction via immunoblot of
fedratinib to mitochondrial ETC inhibitors: Oligomycin (5 µM), Antimycin A (50 nM), Piericidin
(20 nM) and uncoupler CCCP (1 µM). Experiments (**b**, **d**, **f-k**) were performed three times,
experiments (**a**) was performed once with biological duplicates, and experiment (**c**) was
performed once with four biological replicates. Significance was calculated with an unpaired
two-tailed t-test. n.s. = not significant, *P ≤ 0.05, **P ≤ 0.01, ***P ≤ 0.005, ****P ≤ 0.001. Data
are mean ± SD. Statistical source data are provided in Source Data Extended Data Fig. 3.
Immunoblot source data are provided in Supplemental Figure 3.

Extended Data Figure 4: Fedratinib-dependent ERMCS formation is independent of its ability to induce autophagy
a, Immunoblot of delipidated (I) and lipidated (II) LC3B from fedratinib, chloroquine (CQ) and fedratinib/CQ double treated cells. **b**, Relative MFI of Lysotracker deep red in cells treated with dose dependent increase of fedratinib. **c**, Relative ERMCS from cells pretreated with CQ or bafilomycin A for 6 hr then 18 hr of fedratinib. **d**, Fluorescent images of cells treated with fedratinib with SPLICS, mitoTagRFP, and Lysotracker deep red. **e**, Fluorescent images of cells treate with fedratinib with SPLICS, mitoTagRFP, and DAPRed (autophagosome). Experiments (**a-c**) were performed three times, and experiment (**d**, **e**) were performed twice. Significance was calculated with an unpaired two-tailed t-test. n.s. = not significant, *P ≤ 0.05, **P ≤ 0.01, ***P ≤ 0.005, ****P ≤ 0.001. Data are mean ± SD. Statistical source data are provided in Source Data Extended Data Fig. 4. Immunoblot source data are provided in Supplemental Figure 4.

1563
Extended Data Figure 5: Fedratinib does not induce ER structural changes, UPR, and ER
stress.
a, **b**, **c**, Immunoblot of cells treated with fedratinib or thapsigargin (50 nM) for 24 hr for (**a**)
UPR^{ER} sensors and transcription targets, (**b**) BiP and CHOP, (**c**) ER sheet and tube shaping
proteins. **d**, Relative ERMCS in cells pre-treated with ER stress chaperone
tauroursodeoxycholic acid (TUDCA) (400 µM) for 6 hr, and then treated with fedratinib or
thapsigargin (50 nM) for another 18 hr. **e**, Microscopy images of SPLICS/mtRFP reporter cells
transfected with ER marker SEC61 treated with fedratinib or vehicle control for 24 hr.

Experiments (**a-d**) were performed three times, and experiment (**e**) was performed twice.
Significance was calculated with an unpaired two-tailed t-test. n.s. = not significant. Statistical
source data are provided in Source Data Extended Data Fig. 5. Immunoblot source data are
provided in Supplemental Figure 5.
1576

1577
1578
Extended Data Figure 6: Effects of fedratfnib on various mitochondrial function.
a, Immunoblot analyses of mitochondrial fusion (MFN2), fission (DRP1, DRP1^{pS637}, DRP1^{pS616}),
mitochondria contact site and cristae organization system (MICOS) components
(MIC25/CHCHD6, MIC60/IMMT) from cells treated with vehicle or fedratinib for 24 hr. **b**,
Immnuoblot analyses of mitochondria complex I (NDUFB8), complex II (SDHB), complex III

(UQCRC2), complex IV (COXII), and complex V (ATP5A) from cells treated with vehicle and
fedratinib for 24 hr. **c**, Relative mitochondrial content, **d**, mitochondrial DNA (MT-DNA) copy
number, **e**, mitochondrial membrane potential from cells treated with vehicle and fedratinib for
24 hr (Uncoupler CCCP (1 µM) was used as positive control). **f**, Mitochondrial ROS (mtROS)
was assessed using a ratio of MitoSOX Deep Red-to-MitoTracker Green. mito-paraquat
(mitoPQ, 10 µM), myxothiazol (25 nM), and piericidin A (20 nM) were used to induce mtROS. **g**,
General ROS MFI was measured with CellROX Deep Red with positive control tert butyl
hydrogen peroxide (TBHP, 200 µM), positive control Antimycin A (50 nM), and fedratinib for
24hr. **h**, Relative ERMCS from cells pretreated with ETC CIII ROS quencher S3QEL2 (10 µM)
for 3 hr then treated with Antimycin A (50 nM) for 3 hr, and then treated with fedratinib for 18 hr.
i, Relative ERMCS from cells treated with calcium chelation (BAPTA-AM; 5 µM) and **j**, MCU
inhibition (Ru360; 10 µM) for 24 hr, and then treated with vehicle or fedratinib for 24 hr following.
k, Mitochondria calcium was monitored with Rhod2 staining from cells treated with vehicle,
positive control NCLX inhibitor, CGP-37157 (50 µM), or fedratinib for 24 hr. Experiments (**a-l**)
were performed three times. Significance was calculated with an unpaired two-tailed t-test. n.s.
= not significant, *$P \leq 0.05$, **$P \leq 0.01$, ***$P \leq 0.005$, ****$P \leq 0.0001$. Data are mean ± SD.
Statistical source data are provided in Source Data Extended Data Fig. 6. Immunoblot source
data are provided in Supplemental Figure 6.
1602

1603
1604
Extended Data Figure 7: The effects of fedratinib on mitochondrial bioenergetics
a, Oxygen consumption (OCR), **b**, basal respiration, **c**, maximal respiration, **d**, ATP-linked
respiration, **e**, spare respiratory capacity, **f**, proton leak, **g**, extracellular acidification rate (ECAR)
and, **h**, basal ECAR of fedratinib or vehicle treated cells undergoing Seahorse MitoStress test.
Experiments (**a-g**) were performed three times. Significance was calculated with an unpaired
two-tailed t-test. n.s. = not significant, *P ≤ 0.05, **P ≤ 0.01, ***P ≤ 0.005, ****P ≤ 0.001. Data
are mean ± SD. Statistical source data are provided in Source Data Extended Data Fig. 7.

Extended Data Figure 8: Light and electron microscopy analyses of mitochondria in fedratinib-treated cells.

a, Confocal images of mtROS (MitoSOX Deep Red) and total mitochondrial mass (mitoTagRFP)
from cells treated with vehicle and fedratinib for 24 hr (Upper left). Line intensity profile of
representative SPLICS^{Hi} and SPLICS^{Lo} mitochondria (Lower left). Ratiometic analyses of
mitochondria membrane potential via MitoSOX/mitoTagRFP ratio were performed on SPLICS^{Hi}
and SPLICS^{Lo} mitochondria in cells following treatment with vehicle and fedratinib for 24 hr. Cell
number: n = 10 (Right). Scale bar: 1 μm. **b**, Representative confocal images of MitoTracker
Deep Red (MTDR) stained cells treated with vehicle and fedratinib for 24 hr. **c**, Representative
confocal images of MitoSOX Deep Red stained cells treated with vehicle and fedratinib for 24 hr.
d, Three representative TEM images in cells treated with vehicle or fedratinib for 24 hr. Cristae
structures are annotated in corresponding colors. Red: Laminar cristae. Blue: Non-laminar
cristae. Aspect ratio, circularity, and perimeter (μm) of mitochondria in cells treated with vehicle
or fedratinib for 24 hr (n=161-299Yellow: Compartmentalized cristae. Green: Loss of cristae.
Two-tailed paired t-test were performed on (A). n.s. = not significant. Data are median ±
interquartile range (**b-d**). Experiments (**a-c**) were performed twice, and experiment (**d**) was
performed once. n.s. = not significant, **P ≤ 0.01, ***P ≤ 0.001, ****P ≤ 0.0001. Statistical source
data are provided in Source Data Extended Data Fig. 8.
1631

1632
1633
Extended Data Figure 9: Whole cell metabolomics and lipidomics analyses of fedratinib-
treated cells.

a, Volcano plots of polar metabolites in cells treated with vehicle and fedratinib for 24 hr
(Log$_2$(Fold change) is plotted with fedratinib-over-vehicle control). **b**, Principal component
analysis (PCA) of the lipidome from cells treated with vehicle or fedratinib. **c**, Volcano plots of
non-polar lipids from vehicle or fedratinib-treated cells. Log2(FC) is plotted with fedratinib-over-
vehicle control. Experiment (**a**) was performed once with three biological replicates.
Experiments (**b** and **c**) were performed once with four biological replicates. Metabolomics and
Lipidomics raw data is included in Supplementary Table 5 and 6.
1643

Extended Data Figure 10: Regulation of ERMCS by mitochondrial ROS inducer and modulators of ETC flow.

a, Immunoblot of cells treated with PHD inhibitor (FG4592, 50 μM), hypoxia (2% oxygen), and/or fedratinib. **b**, Relative ERMCS in cells treated with mitochondrial-paraquat (mitoPQ, 10 μM) for 24 hr, and then with vehicle or fedratinib for another 24 hr. **c**, Relative ERMCS in AOX expressing cells. **d**, Relative ERMCS in AOX expressing cells treated with complex III inhibitor myxothiazol (50 nM) or antimycin A (100 nM) for 24 hr, and then treated with vehicle or thapsigargin (50 nM) for another 24 hr. **e**, Relative ERMCS in cells treated with complex I and III

inhibitor treatment, piericidin A (20 nM), myxothiazol (50 nM), or antimycin A (100 nM) for 24 hr,
and then treated with vehicle or thapsigargin (50 nM) for another 24 hr. **f**, Relative ERMCS level
in cells co-treated with complex I inhibitor piericidin A (20 nM) and H_2S or **g**, hypoxia. **h**, Mass
spectrometry measurement of relative CoQ redox state from cells treated with vehicle, fedratinib,
or positive control antimycin A (100 nM) for 24 hr. Mass spectrometry measurement of CoQ_{10}
levels in isolated mitochondria from SW480 cells treated with vehicle or fedratinib for 24 hr. **i**,
HPLC measurement of relative CoQ redox state and total CoQ concentrations in cells treated
with vehicle or fedratinib for 24 hr. **j**, Confocal imaging of COQ9 in SPLICS/mtRFP cells treated
with vehicle or fedratinib for 24 hr. Experiments (**a**-**g**) were performed three times, experiment (**h**,
i) was performed once with biological triplicates, and experiment (**j**) was performed twice,**.**
Significance was calculated with an unpaired two-tailed t-test. n.s. = not significant, *P ≤ 0.05,
**P ≤ 0.01, ***P ≤ 0.005, ****P ≤ 0.001. Data are mean ± SD. Statistical source data are
provided in Source Data Extended Data Fig. 9. Immunoblot source data are provided in
Supplemental Figure 7.
1668
1669

See accompanying Microsoft Excel document for Supplementary Tables and movies:

Supplementary Table 1. Primary results from image-based drug repurposing screen for ERMSC modulators.

Supplementary Table 2. Whole cell and FAMS-sorted mitochondrial proteomics results in vehicle (DMSO) treated cells, fedratinib-treated cells. SPLICS^{Lo} and SPLICS^{Hi} mitochondria from vehicle and fedratinib-treated cells.

Supplementary Table 3. BrU-seq data for SW480 cells treated with fedratinib for 30 min and 2 hr.

Supplementary Table 4. CRISPR screening results for regulators of ERMCS.

Supplementary Table 5. Metabolomics information for fedratinib-treated SW480 and HT1080 cells.

Supplementary Table 6. Lipidomic information for fedratinib-treated SW480 and HT1080 cells and mitochondria.

Supplementary Table 7. Nucleotide sequences used for primers and oligos.

Supplementary Movie 1: SW480 cells expressing SPLICS and mitoTagRFP being treated with fedratinib and imaged on Zeiss Lattice Light Sheet 7 for 16 hr.

Supplementary Movie 2: HT1080 cells expressing SPLICS and mitoTagRFP being treated with fedratinib and imaged on Zeiss Lattice Light Sheet 7 for 16 hr.

Supplementary Movie 3: CLEM Segmentation and EM overlay of SW480 cells treated with DMSO vehicle for 24 hr.

Supplementary Movie 4: CLEM Segmentation and EM overlay of SW480 cells treated with fedratinib for 24 hr.