# Nanoligomers targeting NF-κB and NLRP3 reduce neuroinflammation and improve cognitive function with aging and tauopathy

Devin Wahl[1,2], Sydney J. Risen[3,4], Shelby C. Osburn[1,2], Tobias Emge[1,2], Sadhana Sharma[5], Vincenzo S. Gilberto[5], Anushree Chatterjee[5], Prashant Nagpal[5], Julie A. Moreno[3,4], and Thomas J. LaRocca[1,2]*

[1] Department of Health and Exercise Science, Colorado State University, Fort Collins, CO
[2] Columbine Health Systems Center for Healthy Aging, Colorado State University, Fort Collins, CO
[3]Department of Environmental and Radiological Health Sciences, Colorado State University, Fort Collins, CO
[4]Brain Research Center, Colorado State University, Fort Collins, CO
[5]Sachi Bio, Colorado Technology Center, Louisville, Colorado

*Correspondence: Thomas LaRocca, PhD
Colorado State University
1582 Campus Delivery
Fort Collins, CO 80523
tom.larocca@colostate.edu

**SUMMARY**

Neuroinflammation contributes to impaired cognitive function in brain aging and neurodegenerative disorders like Alzheimer's disease, which is characterized by the aggregation of pathological tau. One major driver of both age- and tau-associated neuroinflammation is the NF-κB and NLRP3 signaling axis. However, current treatments targeting NF-κB or NLRP3 may have adverse/systemic effects, and most have not been clinically translatable. Here, we tested the efficacy of a novel, nucleic acid therapeutic (Nanoligomer) cocktail specifically targeting both NF-κB and NLRP3 in the brain for reducing neuroinflammation and improving cognitive function in old wildtype mice, and in a mouse model of tauopathy. We found that 4 weeks of NF-κB/NLRP3-targeting Nanoligomer treatment strongly reduced neuro-inflammatory cytokine profiles in the brain and improved cognitive-behavioral function in both old and tauopathy mice. These effects of NF-κB/NLRP3-targeting Nanoligomer treatment were associated with reduced glial cell activation in old wildtype mice, less pathology in tauopathy mice, favorable changes in transcriptome signatures of inflammation (reduced) and neuronal health (increased) in both mouse models, and positive systemic effects. Collectively, our results provide a basis for future translational studies targeting NF-κB/NLRP3 in the brain, perhaps using Nanoligomers, to inhibit neuroinflammation and improve cognitive function with aging and neurodegenerative disease.

## 1. INTRODUCTION

Aging is associated with declines in cognitive function and is the primary risk factor for most neurodegenerative diseases, including Alzheimer's disease (AD). These declines in cognitive function are closely linked with the accumulation of pathological tau, a main hallmark AD that also increases with aging [1, 2]. One key macro-mechanism that contributes to brain aging and tau pathology (tauopathy) is neuroinflammation, which is characterized by innate immune signaling, glial cell activation and the release of neurotoxic, pro-inflammatory cytokines [3]. In fact, evidence suggests that neuroinflammation may occur early in brain aging and both precede and be exacerbated by tauopathy [4]. As such, identifying strategies to reduce neuroinflammation in the context of aging and tauopathy is an important biomedical research goal [5].

The transcription factor Nuclear Factor Kappa B (NF-κB) and inflammasome member NLR Family Pyrin Domain Containing 3 (NLRP3) are key drivers of age- and tauopathy- associated neuroinflammation, and both play central roles in adverse processes that drive brain aging and AD (e.g., immune activation and pro-inflammatory signaling) [6, 7]. Recent studies have demonstrated that inhibition of either NF-κB or NLRP3 alone reduces immune activation/pro-inflammatory cytokine release in the brain and improves cognitive function [8-10]. However, current treatments meant to target NF-κB and NLRP3 may have off-target effects (e.g., by inhibiting activity of other enzymes) [11] and/or poor biodistribution, and many systemic anti-inflammatory treatments (e.g., immunosuppressive steroids) do not cross the blood-brain barrier. To address these concerns, our team recently developed a novel Nanoligomer (nucleic acid therapeutic) cocktail that targets both NF-κB and NLRP3 DNA, to inhibit transcription, and mRNA, to inhibit translation (**Figure 1A**). Nanoligomers comprise an antisense peptide nucleic acid (PNA) conjugated to a gold nanoparticle for improved delivery [12-17]. This Nanoligomer cocktail is highly specific, crosses the blood-brain barrier and inhibits neuroinflammation *in vitro* and *in vivo* [14, 15, 17], but its influence on cognitive function in the context of brain aging and tauopathy has not been explored.

In the present study, we evaluated the efficacy of targeting both NF-κB and NLRP3 in brain aging and tauopathy by treating both old wildtype mice and transgenic rTg4510 mice (a model of tauopathy commonly used in neurodegeneration research) with our Nanoligomer cocktail. We found that this treatment: 1) improved cognitive function and broadly reduced neuroinflammation in both models; 2) attenuated tau pathology and reduced age-associated glial cell activation; 3) reversed transcriptome changes common to brain aging and tauopathy; and 4) exerted favorable systemic effects. Our results show that selective targeting of both NF-κB and NLRP3 is a potent strategy for reducing neuroinflammation, and they provide a basis for similar studies in larger animal models and/or pilot clinical trials.

## 2. RESULTS

### 2.1. Short-term Nanoligomer treatment inhibits NF-κB and NLRP3 in the hippocampus.

NF-κB and NLRP3 are central mediators of age-related inflammation (i.e., "inflammaging"), neuroinflammation,

innate immune responses and cytokine release, including in the brain [6, 7, 18, 19]. For initial proof of concept and to provide insight into the ability of Nanoligomers to target NF-κB and NLRP3 in the brain, we first treated older (aged 19 months) male and female C57Bl/6J mice with a Nanoligomer cocktail targeting NF-κB/NLRP3 or saline control (1x intraperitoneal injection; 150 mg/kg body weight). Younger (aged 8 months) male and female mice, as a reference group, also received a saline injection. After 16 hours, mice were euthanized and immunoblots were performed on isolated whole hippocampus. We observed significant reductions in both NF-κB and NLRP3 in old treated vs. untreated mice (**Figure 1B,D**), and a modest increase in phosphorylated (activated) NF-κB in older vs. young control animals that was absent in older treated animals (**Figure 1C**). These effects of acute Nanoligomer administration on central mediators of neuroinflammation and innate immune responses show that the drug crosses the blood brain barrier, consistent with our previous reports [12, 13, 15, 17], and that it engages its targets, suggesting the potential for downstream effects on mediators of neuroinflammation (e.g., microglia, pro-inflammatory gene induction and cytokine release) [20].

**Figure 1. Short-term Nanoligomer treatment inhibits NF-κB and NLRP3 in the brain. (A)** NF-κB and NLRP3 interact to drive inflammation, Nanoligomers that block their translation may reduce age- and tau-associated neuroinflammation. **(B-D)** Western blot data showing protein levels of NF-κB, phosphorylated (p)-NF-κB and NLRP3 in the hippocampus of young (8 months) and older (19 months) wildtype mice, and older mice 16 h after treatment with a NF-κB/NLRP3-targeting Nanoligomer cocktail. Raw immunoblots shown in **(E)**. N = 4/group; *p < 0.05.

## 2.2. Long-term Nanoligomer treatment improves cognitive function in old wildtype and tauopathy mice.

Although NF-κB and NLRP3 play complex roles in brain health [21], most data suggest that inhibiting these proteins individually improves cognitive function [22, 23]. Therefore, given our finding that acute Nanoligomer treatment reduced NF-κB and NLRP3 in the hippocampus (a key area of the brain involved in memory and

cognitive function) and the central role of neuroinflammation in cognitive dysfunction with brain aging and AD [24], we next tested the hypothesis that long-term Nanoligomer treatment (150 mg/kg body weight; 3x week for one month) would improve cognitive function in old C57Bl/6J male and female mice. In addition, because tauopathy (a key hallmark of AD) is associated with NF-κB/NLRP3 activation [25] and cognitive dysfunction [26], we also treated male and female rTg4510 mice (aged 2 months) using the same approach. Importantly, we did not note any adverse effects with this treatment, and all body weights remained stable (**Supplementary Data**). After the treatment period, we measured short-term recognition memory via Novel Object Recognition testing and anxiety-like behavior via Elevated Plus Maze testing. In Novel Object Recognition testing, we found that treated older and tauopathy mice performed significantly better (as measured by recognition indices) than their untreated counterparts, and similarly to younger mice and littermate controls, respectively (**Figure 2A**). Also, in the Elevated Plus Maze test, there was a trend for treated older and tauopathy mice to spend more time in open maze arms than untreated mice, reflecting reduced anxiety (**Figure 2B**). Finally, we also measured grip strength because of its relationship to cognitive function and overall health [27], and we found significant increases in forelimb grip strength in old treated mice. Interestingly, however, these improvements were not noted in treated tauopathy mice (**Supplementary Figure S1**). Future, more comprehensive studies are needed to assess how NF-κB/NLRP3-targeting Nanoligomer treatment affects other peripheral functional outcomes (e.g., treadmill endurance, wire hang, rotarod performance). However, these results show that long-term Nanoligomer treatment improves several domains of cognitive-behavioral function in old, non-transgenic mice, and in tauopathy model mice.

**Figure 2. Long-term treatment with Nanoligomers targeting NF-κB and NLRP3 improves cognitive-behavioral function in old wildtype and tauopathy mice. (A)** Novel object recognition index (short-term recognition memory) quantified as time exploring the novel object versus total object exploration time in young, old and old Nanoligomer-treated wildtype mice, as well as littermate (LM) controls, rTg4510 tauopathy and rTg4510 Nanoligomer treated mice. **(B)** Time spent in open arms during Elevated Plus Maze testing, a reflection of anxiety-like behavior, in the same mice. N = 6-11/group; *p < 0.05.

## 2.3. Broad Nanoligomer-induced reductions in neuroinflammation correlate with cognitive function.

The activation of NF-κB and NLRP3 via aging or tauopathy induces the production of many pro-inflammatory

cytokines [28]. Therefore, we performed multiplex ELISAs on cortex tissue from the same mice described above to assess levels of key cytokines involved in neuroinflammation and innate immune responses. Principal component analyses (PCA) revealed that aging and tauopathy were associated with robust and broad increases in many cytokines, whereas cytokine levels in treated old and tauopathy animals were similar to those in young mice and littermate controls, respectively (**Figure 3A and 3B**, left panels). At the individual cytokine level, several important pro-inflammatory and immune-activating cytokines that increased with aging and tauopathy included Interleukin 5 (IL-5) and Macrophage-Inflammatory Protein 2 (MIP-2$\alpha$), but these effects were reduced with Nanoligomer treatment. Similar trends were noted for most Interleukins (ILs), Monocyte Chemoattractant Proteins (MCPs), and Interferons (IFNs) (**Supplementary Figure S2**). Additionally, we found that cytokine levels (global and individual) were significantly inversely correlated with short-term recognition memory (**Figure 3A and 3B,** center and right panels), which is consistent with existing evidence that increased cytokine levels in the brain are associated with cognitive decline [29]. Together, these data show that long-term NF-κB/NLRP3-targeting Nanoligomer treatment broadly reduces neuroinflammation, including many cytokines that are directly modulated by NF-κB and NLRP3 [19], and that these changes may contribute to improvements in cognitive function in old wildtype and tauopathy model mice.

**Figure 3. Broad reductions in neuroinflammation with Nanoligomer treatment are related to greater cognitive-behavioral function in old wildtype and tauopathy mice. (A)** Principal Component 1 (PC1) from PC analysis of 36 cytokines in hippocampus of young, old and old Nanoligomer-treated wildtype mice (left panel). Correlation of PC1 values with Novel Object Recognition Index scores in the same mice (middle panel) and heatmap showing magnitude of individual cytokine correlations with Recognition Index (right panel). **(B)** The same PC and cytokine analyses in littermate (LM) controls, rTg4510 tauopathy and rTg4510 Nanoligomer-treated mice. All analyses N = 6-11/group; *p < 0.05.

## 2.4. Reductions in phosphorylated tau with long-term Nanoligomer treatment.

Importantly, tau pathology can both activate and be exacerbated by NF-κB and NLRP3 [30-32], driving cycles of inflammation-mediated tau propagation and toxicity [33], and phosphorylated (activated) tau is closely associated with cognitive dysfunction [34]. To characterize the effects of Nanoligomer treatment on tau, we performed immunohistochemistry on hippocampus and frontal cortex sections from control and treated rTg4510 mice. We imaged and quantified two insoluble tau residues (phosphorylated at threonine-181 and threonine-231), which have been associated with AD pathology and progression [35, 36]. We found that these tau species increased in hippocampus and cortex of rTg4510 mice, similar to what others have reported in other tauopathy models [37, 38], and that this effect was attenuated with Nanoligomer treatment (**Figure 4**). Although preliminary in nature, these results suggest that, in addition to reducing neuroinflammation, NF-κB/NLRP3-targeting Nanoligomer treatment may suppress tau pathology, but future studies are needed to confirm this observation and clarify the order of events using additional animal/transgenic models.

**Figure 4. Reductions in phosphorylated tau in the hippocampus with Nanoligomer treatment.**
Quantification and representative immunohistochemistry images of phosphorylated tau (p-threonine-181 and p-threonine 231) in hippocampal and frontal cortex sections from littermate (LM) controls, rTg4510 tauopathy and rTg4510 Nanoligomer-treated mice. Insets in lower right corner of images are 2x magnification of representative cell bodies indicated in smaller boxes. Note reduction in p-tau staining density with Nanoligomer treatment. *p < 0.01.

## 2.5. Reversal of age-related glial cell activation with long-term Nanoligomer treatment.

Although aged wildtype mice do not develop tau pathology, they do develop gliosis like humans. Indeed, in human aging/AD, glial cells (astrocytes and microglia) become immune activated, secreting cytokines that impair neuronal and cognitive function [39]. This neuroinflammation is also associated with morphological changes [40, 41], including alterations in process length [42] and branching [43, 44], which have detrimental effects on neurons (e.g., synaptic dysfunction). To evaluate the effects of NF-κB/NLRP3-targeting Nanoligomer treatment on astrocyte and microglia activation/morphology with aging, we performed immunofluorescence staining on cortex tissue from old, old treated, and young mice. Consistent with previous findings showing that astrocyte activation increases with aging [40, 45], we found evidence of age-related increases in Glial Fibrillary Acidic Protein (GFAP) [46], but this effect was less pronounced in old treated mice (**Figure 5A**). Similar to previous studies [47, 48], we also found that aging was associated reduced astrocyte process length and branch number (**Figure 5B**). However, these effects were reversed with Nanoligomer treatment. We also found age-related increases in the activation of microglia (the resident immune cells of the brain), as reflected by increases in Allograft Inflammatory Factor 1 (IBA1), but these effects were less pronounced in old treated mice (**Figure 5C**). Furthermore, consistent with previous findings [49], we observed changes in microglial process length and branch number in old animals, and again, these effects of aging were attenuated with Nanoligomer treatment (**Figure 5D**). Collectively, these results suggest that the beneficial effects of Nanoligomers targeting NF-κB/NLRP3 in old mice may be due in part to reducing age-related changes in glial cells.

**Figure 5. Reversal of age-related cellular changes in astrocytes and microglia with Nanoligomer treatment.**
**(A)** Immunofluorescence images of Glial Fibrillary Acidic Protein (GFAP) staining in hippocampus sections from young, old and old Nanoligomer-treated wildtype mice. Note general increase in GFAP/altered astrocyte morphology in old mice that is reversed with Nanoligomer treatment. **(B)** Quantification of GFAP+ process length and branch number. **(C)** Immunofluorescence images of Ionized Calcium-Binding Adapter molecule 1 (IBA1) staining in the same mice. Note general increase in IBA1/altered microglial morphology in old mice that is reversed with treatment. **(D)** Quantification of IBA1+ process length and branch number. *p < 0.01 vs. young; #p < 0.01 vs. old control.

## 2.6. Transcriptome signatures of immune activation and reduced neuronal health in aging and tauopathy are reversed with long-term Nanoligomer treatment.

Perturbations in the transcriptome affect downstream biological processes that mediate brain aging/AD [50]. To provide broad insight into how Nanoligomer treatment affected genes/pathways most related to aging and tauopathy, we performed RNA-seq on hippocampus tissue from all groups of mice. We found that aging and tauopathy were associated with broad changes in gene/transcript expression (**Figure 6A**, top rows of heatmaps), and that many of these genes/transcripts were reversed in terms of relative expression with Nanoligomer treatment (**Figure 6A**, bottom rows of heatmaps). For example, one of the top transcripts that increased with aging but decreased with Nanoligomer treatment was S100 calcium-binding protein A8 (*S100A8*), which activates NF-κB and induces microglial activation [51]. One of the top transcripts that increased with tauopathy but decreased with Nanoligomer treatment was Interleukin 17 Receptor E (*IL17RE*), which is also associated with the activation of NF-κB [52] and cognitive dysfunction [53]. Given these observations and because aging and tau pathophysiology intersect [54], to understand how Nanoligomer treatment affected genes/pathways common to both aging and tauopathy, we next compared the top 1000 most increased/decreased genes/transcripts common to both aging and tauopathy, and those same genes/transcripts in old and tauopathy-treated mice (**Figure 6A,** top Venn diagrams). Gene ontology (GO) analyses revealed that increased genes common to both aging and tauopathy reflected pathways associated with inflammation/immune responses and cognitive dysfunction, including immune receptor activity, major histocompatibility complex (MHC) binding, and synapse pruning (**Figure 6B**, left). Decreased genes/transcripts common to both aging and tauopathy comprised pathways associated with neurotransmitter receptor activity and signal transduction (**Figure 6B**, right). However, these patterns were generally reversed with treatment. That is, among the transcripts that changed in both treated old and treated tauopathy mice (**Figure 6A,** bottom Venn diagrams), those that were increased reflected pathways involved in brain aging and neuronal/cognitive function including ion channel activity, neuronal signaling, and neurogenesis (**Figure 6C,** left), whereas common decreased transcripts with treatment in both aged and tauopathy mice reflected pathways associated with inflammation and immune responses, including cell adhesion and stress response processes (**Figure 6C,** right). Together, these results show that Nanoligomer treatment targeting NF-κB/NLRP3 favorably influences several important biological and molecular processes associated with brain aging and tauopathy, particularly those involved in neuroinflammation/immune activation and neuronal health (e.g., synaptic signaling and neurotransmitter activity).

**Figure 6. Transcriptome signatures of immune activation and reduced neuronal health with aging and tauopathy are reversed with Nanoligomer treatment. (A)** Heatmaps showing Log2Fold differences for the top 1000 increased and decreased genes/transcripts in old vs. young control mice and rTg4510 tauopathy mice vs. littermate (LM) controls (top heatmap rows), and Log2Fold differences for the same genes/transcripts in treated old and rTg4510 mice vs. their untreated counterparts (bottom heatmap rows). Venn diagrams in center show genes/transcripts that were increased or decreased in both old and rTg4510 mice (top), and genes/transcripts that were reversed in terms of relative expression in both treated old and treated rTg4510 mice. **(B)** Gene ontology analyses of genes/transcripts that were increased or decreased in both old and rTg4510 mice. **(C)** Gene ontology analyses of genes/transcripts that were increased or decreased in both treated old and treated rTg4510 mice.

## 2.7. Other favorable changes in response to Nanoligomer treatment.

Current therapies meant to target neuroinflammation in the brain often have off-target effects [11, 55], and drug-induced hepatotoxicity is common following exposure to many bioactive compounds [56]. The liver is a key metabolic organ that processes such compounds and therefore a relevant target for therapies meant to help the brain [57]. To understand how long-term Nanoligomer treatment affects the liver, we performed multiplex ELISAs on liver tissue from young, old, and old treated mice (**Figure 7A**). We found that several pro-inflammatory and immune-activating cytokines that increased with aging were reduced with Nanoligomer treatment, including IFN-ɣ inducible protein (IP-10) and chemokine ligand 1 (CXCL1). Additionally, the anti-inflammatory cytokine interleukin 10 (IL-10) was reduced with aging, but this effect was reversed with Nanoligomer treatment. We also found that aging was associated with evidence of hepatocyte polyploidy and polymorphonuclear leukocytes in the liver, a key source of tissue damage due to inflammation [58], and these effects were attenuated with Nanoligomer treatment (**Figure 7B,C**). Furthermore, in Nanoligomer-treated old mice, these favorable changes in the liver were associated with a preservation of circulating (plasma) levels of fibroblast growth factor 21 (FGF-21), a hormone synthesized by the liver that has been linked with longevity and neuroprotection [59, 60]. Collectively, these results suggest that Nanoligomers targeting NF-κB/NLRP3 are

not hepatotoxic and may even favorably modulate several markers of inflammation and liver/peripheral health in old mice.

**Figure 7. Favorable peripheral/systemic changes with long-term Nanoligomer treatment. (A)** Cytokine concentrations in liver of young, old and old Nanoligomer-treated wildtype mice. N = 4/group; *p < 0.05 vs. young; #p < 0.05 vs. old **(B)** Representative immuno-histochemistry images showing infiltrating immune cells at portal vein border (white arrows) and **(C)** quantification of immune cell infiltration in the same mice. *p < 0.01. **(D)** Plasma levels of fibroblast growth factor 21 (FGF-21) measured by ELISA in the same mice. N = 6-11/group; *p < 0.01.

## DISCUSSION

Neuroinflammation impairs cognitive function and is a key macro-mechanism uniting brain aging and AD [5], both of which are linked with tau pathology. As such, there is an urgent need to identify safe and well-tolerated treatments that target neuroinflammation, both in the context of brain aging and tauopathy [61]. Age- and tau-related neuroinflammation is characterized by immune activation, glial cell (microglia and astrocytes) reactivity, and the release of neurotoxic, pro-inflammatory cytokines [62, 63], and many of these events involve signaling via NF-κB and NLRP3. These two proteins are upstream mediators of immune activation and considered key components of a 'central axis' of neuroinflammation [7, 19, 64]. For example, NF-κB regulates many inflammatory processes, including the activation of pro-inflammatory cytokines and the transcription of neuroinflammatory genes [19], and NLRP3 is a major component of the inflammasome that directly regulates multiple immune-response pathways (e.g., complement/cytokine signaling) [65]. Our key finding is that simultaneous, targeted inhibition of both NF-κB and NLRP3 via Nanoligomers reduces neuroinflammation and

improves cognitive function in mouse models of both aging and tauopathy. This may be an important, clinically relevant finding, as these small molecule therapeutics are well positioned for human translation.

Several recent investigations have shown that separately targeting either NF-κB or NLRP3 reduces neuroinflammation and improves cognitive function in multiple mouse models [8, 32, 66, 67]. For example, an NF-κB inhibiting peptide was shown to reduce inflammatory cytokine expression and microglial activation in 5xFAD mice [10], and inhibition of NLRP3 with the small molecule MCC950 has been reported to reduce inflammation and improve cognitive function in APP/PS1 mice [8]. However, to the best of our knowledge, although NF-κB and NLRP3 signaling pathways are interconnected [18], no studies have investigated the effects of simultaneously targeting both NF-κB and NLRP3 in old and/or tauopathy model mice. Importantly, we found that this intervention was safe/well-tolerated, and elicited no adverse effects in the liver, a key metabolic organ. Our initial proof-of-concept study also showed that short-term treatment with Nanoligomers targeting NF-κB and NLRP3 caused reductions in target protein levels and activation in the hippocampus of old mice, consistent with previous studies showing that this same cocktail reduces key mediators of inflammation *in vitro* [14] and *in vivo* [17]. To provide translational relevance to this finding, we then performed a longer duration treatment (proportional to that which might be administered to older adults). We treated both old and tauopathy model mice with NF-κB/NLRP3-targeting Nanoligomers for one month, and we found that this treatment improved short-term recognition memory and reduced anxiety (which is oftentimes co-morbid with dementia) [68]. These results are important, because cognitive decline is a major risk factor for mild cognitive impairment and AD [69], and our findings are consistent with results from other studies focused on NF-κB and NLRP3 in mouse models of aging and AD [8, 10]. We note that intraperitoneal Nanoligomer delivery is a translational limitation, but our group is currently developing alternative approaches for administering Nanoligomers, including intranasal inhalation and oral delivery. Demonstrating efficacy of these approaches will be an important next step in this line of investigation.

Importantly, many pro-inflammatory/cytokine signaling pathways contribute to neuroinflammation that drives declines in cognitive function [5, 63, 70]. To address this in the present study, we probed the hippocampus for multiple cytokines that have been linked with cognitive dysfunction, brain aging, and tauopathy. We found that NF-κB/NLRP3-targeting Nanoligomer treatment reduced almost all of these cytokines, with PCA showing a marked reversal of overall cytokine patterns in both aged and tauopathy mice. Moreover, these cytokine profiles were highly (inversely) correlated with cognitive-behavioral function in these animals. Collectively, these results suggest that the mechanism by which Nanoligomer treatment improves cognitive function involves broadly reducing neuroinflammation. This idea is in agreement with our previous studies showing that Nanoligomer treatment reduces neuroinflammation and improves cognitive function in prion-disease mice (a model of severe neurodegeneration) [17], and with numerous other studies showing that targeting neuroinflammation improves cognitive function in mouse models of accelerated brain aging [71]. In future work, it will be important to identify the specific NF-κB/NLRP3-associated cytokine pathways that impact cognitive

function most directly, as some cytokines (e.g., IL-10) can have both protective [72] and deleterious effects in the brain [73].

Neuroinflammation is linked with protein pathology (including tau accumulation) and glial cell dysregulation in aging and AD [4, 74-76]. However, the order of these events is incompletely understood, and neuro-inflammatory signaling mediators like NF-κB and NLRP3 have been reported to both contribute to and be activated by pathology [32]. In our study, we observed several effects of NF-κB/NLRP3-targeting Nanoligomers that may provide insight in this context. First, we found that reduced neuroinflammation with NF-κB/NLRP3-targeting Nanoligomers was associated with reductions in phosphorylated tau in the rTg4510 mouse hippocampus. These insoluble tau residues (phosphorylated at threonine-181 and threonine-231) have been associated with cognitive dysfunction and AD pathology/progression [35, 36]. Because aggressive, transgene-driven expression of tau occurs in rTg4510 mice at young ages, our findings may suggest that the treatment protected against tau-induced NF-κB/NLRP3 activation that further promotes pathology in these animals. This idea would be in line with other studies showing that tau seeding can be reduced by small molecule inhibitors of NLRP3 [31], and that pharmacologically targeting neuroinflammation in general reduces the burden of phosphorylated tau [77, 78]. Second, in aging wildtype mice, we found that Nanoligomer-associated reductions in neuroinflammation were associated with reduced glial cell reactivity. In contrast to rTg4510 and other transgenic mouse models, aging wildtype mice do not develop tau or Aβ pathology—but they do develop glial cell-associated neuroinflammation. Activated microglia and astrocytes release pro-inflammatory, neurotoxic cytokines and are linked with both age-related cognitive dysfunction and tau propagation [2, 5, 39, 63, 79]. Here, similar to previous studies showing that astrocytes and microglia become reactive/pro-inflammatory with aging [40, 45], we found morphological and protein expression evidence of increased reactivity in both astrocytes and microglia of old mice—but NF-κB/NLRP3-targeting Nanoligomer treatment largely attenuated these effects. Together with our findings in tauopathy mice, these observations suggest NF-κB/NLRP3-associated neuroinflammation is a "proximal" target for enhancing cognitive function with aging and neurodegeneration. Future, more comprehensive studies using NF-κB/NLRP3-targeting Nanoligomers are needed to elucidate the order of these events, and/or to evaluate the treatment's effects on other neuroinflammation-associated pathological hallmarks of AD (e.g., Aβ aggregation), perhaps in additional transgenic and/or inducible mouse models (e.g., 5xFAD or APP/PS1).

Finally, given the protective effects of NF-κB/NLRP3-targeting Nanoligomers we observed in both old wildtype and tauopathy mice and evidence that aging and tauopathy may intersect [80], to provide insight into conserved mechanisms of action, we performed transcriptomics (RNA-seq) on hippocampus tissue. We found common transcriptome signatures of aging and tauopathy that reflected pathways associated with neuronal health, neuroinflammation and immune activation, similar to ours and others' observations in pre-clinical models of brain aging [81, 82] and in aging/AD in humans [83]. However, these age- and tauopathy-associated transcriptome effects were largely attenuated with Nanoligomer treatment, and even reversed in some cases. Again, these findings are consistent with the idea that neuroinflammation itself is the most proximal driver of

impaired cognitive function in both aging and tauopathy, which could explain the protective effects of NF-κB/NLRP3-targeting Nanoligomers we observed in both mouse models in the present study. Additional multi-omics studies could be performed in the future to evaluate up- and down-stream mechanisms of action for this novel therapeutic (e.g., epigenetic and/or proteomic changes underlying its efficacy).

## CONCLUSIONS AND FUTURE DIRECTIONS

Our findings demonstrate that Nanoligomers specifically formulated to target NF-κB and NLRP3 are safe and well-tolerated, engage their target proteins in the brain, reduce neuroinflammation and improve cognitive function with both aging and tauopathy. Our molecular/transcriptomic data suggest overlapping mechanisms of action in these contexts, supporting the idea that NF-κB/NLRP3-associated neuroinflammation is a key target in aging and neurodegeneration. Future studies are needed to determine if this approach will work in larger animal models and/or humans, and to disentangle differential effects in short-term vs long-term treatment periods with varying dosages).

## METHODS

### Nanoligomer design and synthesis

Nanoligomers were synthesized by Sachi Bioworks as previously reported [12-17]. Briefly, Nanoligomers comprise an antisense peptide nucleic acid (PNA) conjugated to a gold nanoparticle for improved delivery and membrane transport. The PNA sequences, provided in [13, 14], were previously screened for solubility, self-complementarity, and off-targeting in the human genome. The PNA portions of the Nanoligomers (NLRP3 Sequence: CTTCTACTGCTCACAGG, NFKB Sequence: AGTGGTACCGTCTGCTA) were synthesized on an Apex 396 peptide synthesizer (AAPPTec, LLC) with solid-phase Fmoc chemistry. Fmoc-PNA monomers were obtained from PolyOrg Inc., with A, C, and G monomers protected with Bhoc groups. Following synthesis, the peptides were conjugated with gold nanoparticles and purified via size-exclusion filtration. Conjugation and concentration of the purified solution was monitored through measurement of absorbance at 260 nm (for detection of PNA) and 400 nm (for quantification of nanoparticles). The sequence of the synthesized PNA was confirmed using LC-MS [12].

### Animal husbandry

Older and younger male and female C57Bl/6J mice (aged 19 months and 8 months, respectively) were purchased from the National Institute on Aging aged rodent colony. Male and female rTg4510 mice and appropriate littermate controls (2 months of age) were purchased from the Jackson Laboratory (strain #024854). For the first part of the study (proof of concept to show that the Nanoligomer cocktail reduced both NF-κB and NLRP3 in the brain), cohorts were treated with Nanoligomer acutely (1x intraperitoneal injection; 150 mg/kg body weight) and then studied 16 hours later. For long-term treatment, mice were treated for one month (150 mg/kg body weight; 3x week). Mice were housed throughout all experiments at ~18-23°C on a reverse 12 light/12 dark cycle. Fresh water and *ad-libitum* food (Tekland 2918; 18% protein) was routinely provided to all cages. Animals were consistently health checked by the veterinary staff at Colorado State University (CSU). This protocol was approved by CSU IACUC and Laboratory Animal Resources staff.

### Animal sacrifice and tissue collection

Mice were culled in a fed state in the late morning. After deep anesthetizing with isoflurane, ~1 ml of blood was removed via cardiac puncture followed by cervical dislocation. The left hippocampus and a piece of left cortex tissue were removed, flash frozen on dry ice, and stored at -80°C until further processing. The right hippocampus and right cortex were processed (parrafin-embedded) for immunostaining (details below).

## Immunoblotting

Frozen hippocampus samples were lysed in RIPA buffer containing 100 mM KCl, 40 mM Tris HCl, 10 mM Tris base, 5 mM $MgCl_2$, 1 mM EDTA, and 1 mM ATP (pH 7.5), phoSTOP phosphatase inhibitors and cOmplete protease inhibitors (ThermoFisher). Sample protein concentration was determined using a BCA kit (ThermoFisher/Pierce), and 20 µg of protein per sample was separated by electrophoresis (Bio-Rad Criterion system), transferred to nitrocellulose membranes (Trans-Blot Turbo, Bio-Rad), and blocked in TBS-T with 5% Bovine Serum Albumin (BSA). Primary antibodies were purchased from ABclonal and included phosphorylated nuclear factor kappa B (p-NF-κB, 1:1000 dilution); nuclear factor kappa B (NF-κB; 1:1000) and NLR Family Pyrin Domain Containing 3  (NLRP3; 1:1000). GAPDH (1:2000) was purchased from Novus Biologicals. Membranes were incubated in primary antibody (TBS-T in 5% BSA) for 24 hours at 4°C, followed by HRP-conjugated secondary antibodies (TST-T in 5% milk; 1:5000; Cell Signaling) for 1 h. Membranes were then incubated in ECL, imaged on a ProteinSimple Fluor Chem imaging system, and quantified with ImageJ.

## Cognitive-behavioral testing

Novel Object Recognition: For NOR testing, mice were assessed for short-term memory performance using no habituation phase [84]. Briefly, mice were placed in an opaque arena (50 cm x 50 cm) with two identical objects (tower of Legos) placed 8 cm from each side wall. After an inter-trial interval of two hours, mice were returned to the arena, but one of the old objects was replaced with a new object of similar shape and color (cell culture flask filled with multi-colored sand). Trials ended once 20 s total exploration time was reached, or 10 min had elapsed. If mice were unable to meet the exploration criteria of 20 s or 10 min, they were excluded from the analysis. Recognition index was calculated as novel object exploration time divided by 20 s [84].

Elevated Plus Maze: Elevated Plus Maze testing was performed as previously described [83, 85] and consisted of a cross with four arms (67 cm in length; 2 exposed arms, 2 enclosed arms). Mice were placed in the center of the maze facing an open arm and allowed to freely explore for 5 min. The amount of time in open arms (a measure of anxiety-like behavior) was recorded and analyzed with AnyMaze software.

Grip Strength: The grip strength test was performed as previously described [86]. Mice were tested using the DST-110 Digital Force Gauge (Maze Engineers). A total of 5 trials were performed per mouse and the highest and lowest values were discarded for analysis. The remaining 3 force values were averaged for each mouse. Data are presented as g force/g body weight.

## Multiplexed ELISAs

Brain homogenates from all mice were assessed with a 36-plex Procartaplex cytokine/chemokine panel (ThermoFisher) as previously reported [14, 15], which included the following cytokines and chemokines: ENA-78 (CXCL5), Eotaxin (CCL11), GRO-α (CXCL1), IP-10 (CXCL10), MCP-1 (CCL2), MIP-1α (CCL3), MIP-1β (CCL4), MIP-2α (CXCL2), RANTES (CCL5), G-CSF (CSF-3), GM-CSF, IFN-α, IFN-γ, IL-1α, IL-1β, IL-2, IL-3, IL-4, IL-5, IL-6, IL-9, IL-10, IL-12p70, IL-13, IL-15/IL-15R, IL-17A (CTLA-8), IL-18, IL-22, IL-23, IL-27, IL-28, IL-31, LIF, MCP-3 (CCL7), M-CSF, and TNF-α. For systemic effects (Figure 7), separate individual ELISAs were used. Briefly, 25uL of 10 mg/ml hippocampus, cortex, liver homogenate, or serum was processed using standard ELISA techniques and analyzed on a Luminex MAGPIX xMAP instrument. Standards for each cytokine/chemokine were used with 1:4 dilutions (8-fold dilutions) and background, controls and sample concentrations were determined from a standard curve using Five Parameter Logistic curve fit/quantification.

## Immunohistochemistry and immunofluorescence

Brain and liver tissues were fixed in 10% neutral buffered formalin at room temperature for at least 48 hours. Brain samples were processed on a Leica TP1020 Automatic Benchtop Tissue Processor and embedded in paraffin wax (Cancer Diagnostics). For immunohistochemistry analyses of phosphorylated tau, paraffin-embedded tissue was sectioned at 4 µm, deparaffinized, rehydrated, sodium citate treated, and blocked in Tris A/2% donkey serum (Jackson ImmunoResearch), then incubated overnight in the following antibodies: phospho-tau (Thr181; 1:400 dilution; Invitrogen) and phospho-tau (Thr231; 1:400; ABClonal). Wash steps were performed using 2% bovine serum albumin and 2% Triton-X in 1 M TBS, and an ABC HRP peroxidase detection kit (Vector Laboratories) and ImmPACT DAB Substrate Peroxidase (HRP) Kit (Vector Laboratories) were used for a chromogen. Slides were counterstained with hematoxylin (ThermoFisher), secured with a coverslip in mounting medium and stored at room temperature until imaging. Whole tissue images were taken on an Olympus BX53 microscope with an Olympus DP70 camera using an Olympus UPlanSApo 20x objective

(N.A. = 0.75). Representative images were taken using an Olympus BX53 microscope with an Olympus DP70 camera using an Olympus UPlanFL N 40x objective (N.A. = 0.75). To quantify and select the phosphorylation of tau positive stained cells the manual threshold on the Count and Measure function of Olympus CellSens software (v1.18) was used.  For immunofluorescence analyses of GFAP, IBA1 and glial cell process length and branching, sections were incubated overnight in primary antibodies: IBA1 (1:50 dilution; Abcam) and GFAP (1:100; Dako). Slides were washed with TBS and incubated for 1 h in the dark with AlexaFluor 555 or 647 (Invitrogen) secondary antibodies at 1:500 dilution and 2% normal donkey serum, then washed and incubated in Hoechst dye (1:2000; ThermoFisher) for 3 min. Slides were mounted with Prolong Gold Antifade mounting media (ThermoFisher) and stored in the dark at 4°C prior to imaging. Images were captured using an Olympus BX63 fluorescence microscope equipped with a motorized stage and Hamamatsu ORCA-flash 4.0 LT CCD camera using a 40x Olympus X-Apochromat air objective (N.A. = 0.80). For astrocyte/microglial process length and number analyses, four regions between the dentate gyrus and CA1-CA3 region of the hippocampus were imaged at 40x. To identify/quantify processes and branches, skeletonization of GFAP+ astrocytes and IBA1+ microglia was performed using IMARIS software as previously described [87] to trace and quantify stained filaments. For histology analyses of liver, tissues were sectioned on a ThermoFisher HM 325-2 microtome at 5µm thickness and mounted on positively charged glass slides (Superfrost Plus, Cancer Diagnostics). Liver sections were deparaffinized and stained with hematoxylin and eosin (Cancer Diagnostics), and whole tissue images were taken for analysis using an Olympus DP70 camera and an Olympus UPlanSApo 20x objective. Livers were scored for pathological infiltration of immune cells specifically around the portal triad, and for overall hepatocyte morphology by a trained animal pathologist.

## RNA sequencing and transcriptomic analyses

Transcriptome (RNA-seq) and gene expression analyses were performed on whole hippocampus via methods previously described **[82]**. Frozen samples (n=6/group; 3M and 3F) were transferred into Trizol (ThermoFisher) and homogenized on ice (SP Bel-Art ProCulture Micro-Tube Homogenizer System; VWR Scientific). RNA was isolated using an RNA-specific spin column kit (Direct-Zol, Zymo Research) and treated with DNase to remove genomic DNA. Poly(A)-selected libraries were generated using magnetic oligo dT beads (ThermoFisher) and Illumina TruSeq kits. Libraries were sequenced on an Illumina NovaSeq 6000 platform to yield >40 M 151-bp paired-end FASTQ reads/sample. Reads were trimmed and filtered with the fastp program **[88]**, then aligned to the mm10 *Mus musculus* genome using the STAR aligner **[89]**. Differential gene expression was analyzed with DESeq2 **[90]** and genes/transcripts with BaseMean read counts of <10 were filtered and removed for analyses **[91, 92]**. Differentially expressed genes were analyzed for gene ontology (GO) enrichment using the Metascape program **[93],** and heatmaps were constructed with GraphPad Prism software.

## Statistics

One-way ANOVA with Tukey's *post-hoc* testing was used to assess differences among groups in protein/cytokine levels and measures of cognitive function, as well as glial cell morphology, and simple linear regression (Pearson correlations) were used to relate cytokine levels to cognitive function data. All data were analyzed and presented using GraphPad Prism software. Differentially expressed genes were detected using DESeq2, and genes were sorted by p-value and Log2Fold change to construct volcano plots and heatmaps, respectively. The Metascape program was used for GO and TRRUST analyses [93].

## Data Availability
The RNA-seq data generated in this study are available on the Gene Expression Omnibus (GEO) website under accession number GSE251888. All additional data are included in the data supplement, and raw data will be made available upon request to the corresponding author (T.J.L).

## Funding
This project was supported in part by NIH awards AG078859, AG070562 and AG060302 (T.J.L.) and AG069361 (D.W.), NASA SBIR Awards 80NSSC22CA116 and 80NSSC23CA171 (P.N.), and by intramural (startup) funding from Colorado State University (T.J.L.).

## Declaration of competing interests
S.S, V.G, A.C., and P.N. work at Sachi Bio, a for-profit company, where the Nanoligomer technology was developed. A.C. and P.N. are the founders of Sachi Bio, and P.N. has filed a patent on this technology. All other authors declare no competing interest.

**Supplementary Figures**

**Figure S1. Nanoligomers targeting NF-κB and NLRP3 increase grip strength in old wildtype mice.** Forelimb grip strength in young, old and old Nanoligomer-treated wildtype mice, as well as littermate (LM) controls, rTg4510 tauopathy and rTg4510 Nanoligomer-treated mice. N = 6-11/group; *p < 0.05.

**Figure S2. Nanoligomers targeting NF-κB and NLRP3 reverse age- and tauopathy-related increases in many cytokines in the brain.** Multiplex ELISA analyses of prefrontal cortex tissue from young, old and old Nanoligomer-treated wildtype mice, as well as littermate (LM) controls, rTg4510 tauopathy and rTg4510 Nanoligomer-treated mice. N = 6-11/group; *p < 0.05 vs. young/LM; #p < 0.05 vs. old/rTg4510; red lines represent limit of detection.

# References

1. Xia, X., et al., *Aging and Alzheimer's disease: Comparison and associations from molecular to system level.* Aging Cell, 2018. **17**(5): p. e12802.
2. Wyss-Coray, T., *Ageing, neurodegeneration and brain rejuvenation.* Nature, 2016. **539**(7628): p. 180-186.
3. Heneka, M.T., M.P. Kummer, and E. Latz, *Innate immune activation in neurodegenerative disease.* Nat Rev Immunol, 2014. **14**(7): p. 463-77.
4. Metcalfe, M.J. and M.E. Figueiredo-Pereira, *Relationship between tau pathology and neuroinflammation in Alzheimer's disease.* Mt Sinai J Med, 2010. **77**(1): p. 50-8.
5. Andronie-Cioara, F.L., et al., *Molecular Mechanisms of Neuroinflammation in Aging and Alzheimer's Disease Progression.* Int J Mol Sci, 2023. **24**(3).
6. Youm, Y.H., et al., *Canonical Nlrp3 inflammasome links systemic low-grade inflammation to functional decline in aging.* Cell Metab, 2013. **18**(4): p. 519-32.
7. Walker, K.A., et al., *Connecting aging biology and inflammation in the omics era.* J Clin Invest, 2022. **132**(14).
8. Dempsey, C., et al., *Inhibiting the NLRP3 inflammasome with MCC950 promotes non-phlogistic clearance of amyloid-β and cognitive function in APP/PS1 mice.* Brain Behav Immun, 2017. **61**: p. 306-316.
9. Jha, N.K., et al., *Nuclear factor-kappa β as a therapeutic target for Alzheimer's disease.* J Neurochem, 2019. **150**(2): p. 113-137.
10. Lindsay, A., D. Hickman, and M. Srinivasan, *A nuclear factor-kappa B inhibiting peptide suppresses innate immune receptors and gliosis in a transgenic mouse model of Alzheimer's disease.* Biomedicine & Pharmacotherapy, 2021. **138**: p. 111405.
11. Kennedy, C.R., et al., *A Probe for NLRP3 Inflammasome Inhibitor MCC950 Identifies Carbonic Anhydrase 2 as a Novel Target.* ACS Chemical Biology, 2021. **16**(6): p. 982-990.
12. McCollum, C.R., et al., *Safety and Biodistribution of Nanoligomers Targeting the SARS-CoV-2 Genome for the Treatment of COVID-19.* ACS Biomaterials Science & Engineering, 2023. **9**(3): p. 1656-1671.
13. McCollum, C.R., et al., *Nanoligomers Targeting Human miRNA for the Treatment of Severe COVID-19 Are Safe and Nontoxic in Mice.* ACS Biomaterials Science & Engineering, 2022. **8**(7): p. 3087-3106.
14. Courtney, C.M., et al., *Reversing radiation-induced immunosuppression using a new therapeutic modality.* Life Sciences in Space Research, 2022. **35**: p. 127-139.
15. Sharma, S., et al., *Identifying an Optimal Neuroinflammation Treatment Using a Nanoligomer Discovery Engine.* ACS Chemical Neuroscience, 2022. **13**(23): p. 3247-3256.
16. McDonald, J.T., et al., *Role of miR-2392 in driving SARS-CoV-2 infection.* Cell Reports, 2021. **37**(3): p. 109839.
17. Risen, S.J., et al., *Targeting neuroinflammation by pharmacologic down-regulation of inflammatory pathways is neuroprotective in protein misfolding disorders.* ACS Chemical Neuroscience, 2024.
18. Bauernfeind, F.G., et al., *Cutting edge: NF-kappaB activating pattern recognition and cytokine receptors license NLRP3 inflammasome activation by regulating NLRP3 expression.* J Immunol, 2009. **183**(2): p. 787-91.
19. Liu, T., et al., *NF-κB signaling in inflammation.* Signal Transduct Target Ther, 2017. **2**: p. 17023-.
20. Shih, R.H., C.Y. Wang, and C.M. Yang, *NF-kappaB Signaling Pathways in Neurological Inflammation: A Mini Review.* Front Mol Neurosci, 2015. **8**: p. 77.
21. Mattson, M.P. and M.K. Meffert, *Roles for NF-κB in nerve cell survival, plasticity, and disease.* Cell Death & Differentiation, 2006. **13**(5): p. 852-860.
22. Datusalia, A.K. and S.S. Sharma, *NF-κB Inhibition Resolves Cognitive Deficits in Experimental Type 2 Diabetes Mellitus through CREB and Glutamate/GABA Neurotransmitters Pathway.* Curr Neurovasc Res, 2016. **13**(1): p. 22-32.
23. Saggu, R., et al., *Astroglial NF-kB contributes to white matter damage and cognitive impairment in a mouse model of vascular dementia.* Acta Neuropathologica Communications, 2016. **4**(1): p. 76.
24. Bettio, L.E.B., L. Rajendran, and J. Gil-Mohapel, *The effects of aging in the hippocampus and cognitive decline.* Neurosci Biobehav Rev, 2017. **79**: p. 66-86.
25. Jiang, S., et al., *Proteopathic tau primes and activates interleukin-1β via myeloid-cell-specific MyD88- and NLRP3-ASC-inflammasome pathway.* Cell Reports, 2021. **36**(12): p. 109720.

26. Di, J., et al., *Abnormal tau induces cognitive impairment through two different mechanisms: synaptic dysfunction and neuronal loss.* Scientific Reports, 2016. **6**(1): p. 20833.

27. Jiang, R., et al., *Associations between grip strength, brain structure, and mental health in > 40,000 participants from the UK Biobank.* BMC Medicine, 2022. **20**(1): p. 286.

28. Liu, T., et al., *NF-κB signaling in inflammation.* Signal Transduction and Targeted Therapy, 2017. **2**(1): p. 17023.

29. Flores-Aguilar, L., et al., *Cognitive and brain cytokine profile of non-demented individuals with cerebral amyloid-beta deposition.* Journal of Neuroinflammation, 2021. **18**(1): p. 147.

30. Bhaskar, K., et al., *Pathological tau activates nuclear factor-kappa B (NF-κB) and drives neuroinflammation.* Alzheimer's & Dementia, 2022. **18**(S4): p. e069386.

31. Stancu, I.C., et al., *Aggregated Tau activates NLRP3-ASC inflammasome exacerbating exogenously seeded and non-exogenously seeded Tau pathology in vivo.* Acta Neuropathol, 2019. **137**(4): p. 599-617.

32. Van Zeller, M., et al., *NLRP3 Inflammasome: A Starring Role in Amyloid-beta- and Tau-Driven Pathological Events in Alzheimer's Disease.* J Alzheimers Dis, 2021. **83**(3): p. 939-961.

33. Wang, C., et al., *Microglial NF-κB drives tau spreading and toxicity in a mouse model of tauopathy.* Nat Commun, 2022. **13**(1): p. 1969.

34. Sydow, A., et al., *Tau-induced defects in synaptic plasticity, learning, and memory are reversible in transgenic mice after switching off the toxic Tau mutant.* J Neurosci, 2011. **31**(7): p. 2511-25.

35. Suárez-Calvet, M., et al., *Novel tau biomarkers phosphorylated at T181, T217 or T231 rise in the initial stages of the preclinical Alzheimer's continuum when only subtle changes in Aβ pathology are detected.* EMBO Mol Med, 2020. **12**(12): p. e12921.

36. Lilek, J., et al., *Accumulation of pTau231 at the Postsynaptic Density in Early Alzheimer's Disease.* J Alzheimers Dis, 2023. **92**(1): p. 241-260.

37. Song, L., et al., *Analysis of tau post-translational modifications in rTg4510 mice, a model of tau pathology.* Mol Neurodegener, 2015. **10**: p. 14.

38. Bennett, R.E., et al., *Enhanced Tau Aggregation in the Presence of Amyloid β.* Am J Pathol, 2017. **187**(7): p. 1601-1612.

39. Yu, Y., et al., *The Role of Glial Cells in Synaptic Dysfunction: Insights into Alzheimer's Disease Mechanisms.* Aging Dis, 2023.

40. Deng, Q., et al., *Microglia and Astrocytes in Alzheimer's Disease: Significance and Summary of Recent Advances.* Aging Dis, 2023.

41. Niraula, A., J.F. Sheridan, and J.P. Godbout, *Microglia Priming with Aging and Stress.* Neuropsychopharmacology, 2017. **42**(1): p. 318-333.

42. Streit, W.J., S.A. Walter, and N.A. Pennell, *Reactive microgliosis.* Prog Neurobiol, 1999. **57**(6): p. 563-81.

43. Rodríguez-Callejas, J.D., E. Fuchs, and C. Perez-Cruz, *Atrophic astrocytes in aged marmosets present tau hyperphosphorylation, RNA oxidation, and DNA fragmentation.* Neurobiol Aging, 2023. **129**: p. 121-136.

44. Godeanu, S., et al., *Microglial morphology in the somatosensory cortex across lifespan. A quantitative study.* Dev Dyn, 2023. **252**(8): p. 1113-1129.

45. Clarke, L.E., et al., *Normal aging induces A1-like astrocyte reactivity.* Proceedings of the National Academy of Sciences, 2018. **115**(8): p. E1896-E1905.

46. Liddelow, S.A. and B.A. Barres, *Reactive Astrocytes: Production, Function, and Therapeutic Potential.* Immunity, 2017. **46**(6): p. 957-967.

47. Bondi, H., et al., *Complex and regional-specific changes in the morphological complexity of GFAP+ astrocytes in middle-aged mice.* Neurobiology of Aging, 2021. **100**: p. 59-71.

48. Popov, A., et al., *Astrocyte dystrophy in ageing brain parallels impaired synaptic plasticity.* Aging Cell, 2021. **20**(3): p. e13334.

49. Streit, W.J., et al., *Dystrophic microglia in the aging human brain.* Glia, 2004. **45**(2): p. 208-12.

50. Ham, S. and S.V. Lee, *Advances in transcriptome analysis of human brain aging.* Exp Mol Med, 2020. **52**(11): p. 1787-1797.

51. Wu, M., et al., *S100A8/A9 induces microglia activation and promotes the apoptosis of oligodendrocyte precursor cells by activating the NF-κB signaling pathway.* Brain Res Bull, 2018. **143**: p. 234-245.

52. Liu, Y., et al., *Activation of the IL-17/TRAF6/NF-κB pathway is implicated in Aβ-induced neurotoxicity.* BMC Neurosci, 2023. **24**(1): p. 14.

53. Brigas, H.C., et al., *IL-17 triggers the onset of cognitive and synaptic deficits in early stages of Alzheimer's disease.* Cell Rep, 2021. **36**(9): p. 109574.
54. Kovacs, G.G., *Chapter 25 - Tauopathies*, in *Handbook of Clinical Neurology*, G.G. Kovacs and I. Alafuzoff, Editors. 2018, Elsevier. p. 355-368.
55. Sivamaruthi, B.S., et al., *NF-κB Pathway and Its Inhibitors: A Promising Frontier in the Management of Alzheimer's Disease.* Biomedicines, 2023. **11**(9).
56. Fisher, K., R. Vuppalanchi, and R. Saxena, *Drug-Induced Liver Injury.* Arch Pathol Lab Med, 2015. **139**(7): p. 876-87.
57. Hunt, N.J., et al., *Targeting the liver in dementia and cognitive impairment: Dietary macronutrients and diabetic therapeutics.* Adv Drug Deliv Rev, 2022. **190**: p. 114537.
58. Baggiolini, M., et al., *The polymorphonuclear leukocyte.* Agents Actions, 1978. **8**(1-2): p. 3-10.
59. Hill, C.M., et al., *FGF21 is required for protein restriction to extend lifespan and improve metabolic health in male mice.* Nat Commun, 2022. **13**(1): p. 1897.
60. Zhang, Y., et al., *The starvation hormone, fibroblast growth factor-21, extends lifespan in mice.* Elife, 2012. **1**: p. e00065.
61. Wahl, D., A.N. Cavalier, and T.J. LaRocca, *Novel Strategies for Healthy Brain Aging.* Exerc Sport Sci Rev, 2021. **49**(2): p. 115-125.
62. Carson, M.J., J.C. Thrash, and B. Walter, *The cellular response in neuroinflammation: The role of leukocytes, microglia and astrocytes in neuronal death and survival.* Clin Neurosci Res, 2006. **6**(5): p. 237-245.
63. Heneka, M.T., et al., *Neuroinflammation in Alzheimer's disease.* Lancet Neurol, 2015. **14**(4): p. 388-405.
64. Afonina, I.S., et al., *Limiting inflammation-the negative regulation of NF-κB and the NLRP3 inflammasome.* Nat Immunol, 2017. **18**(8): p. 861-869.
65. Kelley, N., et al., *The NLRP3 Inflammasome: An Overview of Mechanisms of Activation and Regulation.* Int J Mol Sci, 2019. **20**(13).
66. Sun, E., et al., *The Pivotal Role of NF-kB in the Pathogenesis and Therapeutics of Alzheimer's Disease.* Int J Mol Sci, 2022. **23**(16).
67. Li, J., et al., *Protection of MCC950 against Alzheimer's disease via inhibiting neuronal pyroptosis in SAMP8 mice.* Exp Brain Res, 2020. **238**(11): p. 2603-2614.
68. Kwak, Y.T., Y. Yang, and M.S. Koo, *Anxiety in Dementia.* Dement Neurocogn Disord, 2017. **16**(2): p. 33-39.
69. Corey-Bloom, J., *The ABC of Alzheimer's disease: cognitive changes and their management in Alzheimer's disease and related dementias.* Int Psychogeriatr, 2002. **14 Suppl 1**: p. 51-75.
70. Kumar, A., *Editorial: Neuroinflammation and Cognition.* Front Aging Neurosci, 2018. **10**: p. 413.
71. Cheng, C., et al., *Targeting neuroinflammation as a preventive and therapeutic approach for perioperative neurocognitive disorders.* Journal of Neuroinflammation, 2022. **19**(1): p. 297.
72. Richwine, A.F., et al., *Cognitive deficits in interleukin-10-deficient mice after peripheral injection of lipopolysaccharide.* Brain Behav Immun, 2009. **23**(6): p. 794-802.
73. Sanchez-Molina, P., et al., *Chronic IL-10 overproduction disrupts microglia-neuron dialogue similar to aging, resulting in impaired hippocampal neurogenesis and spatial memory.* Brain Behav Immun, 2022. **101**: p. 231-245.
74. Chen, Y. and Y. Yu, *Tau and neuroinflammation in Alzheimer's disease: interplay mechanisms and clinical translation.* Journal of Neuroinflammation, 2023. **20**(1): p. 165.
75. Kwon, H.S. and S.H. Koh, *Neuroinflammation in neurodegenerative disorders: the roles of microglia and astrocytes.* Transl Neurodegener, 2020. **9**(1): p. 42.
76. Miao, J., et al., *Microglia in Alzheimer's disease: pathogenesis, mechanisms, and therapeutic potentials.* Front Aging Neurosci, 2023. **15**: p. 1201982.
77. Cai, Z., Y. Yan, and Y. Wang, *Minocycline alleviates beta-amyloid protein and tau pathology via restraining neuroinflammation induced by diabetic metabolic disorder.* Clinical Interventions in Aging, 2013. **8**(null): p. 1089-1095.
78. Yoshiyama, Y., et al., *Anti-Inflammatory Action of Donepezil Ameliorates Tau Pathology, Synaptic Loss, and Neurodegeneration in a Tauopathy Mouse Model.* Journal of Alzheimer's Disease, 2010. **22**: p. 295-306.
79. Pascoal, T.A., et al., *Microglial activation and tau propagate jointly across Braak stages.* Nature Medicine, 2021. **27**(9): p. 1592-1599.

80. Saha, P. and N. Sen, *Tauopathy: A common mechanism for neurodegeneration and brain aging.* Mech Ageing Dev, 2019. **178**: p. 72-79.

81. Cavalier, A.N., et al., *Protective Effects of Apigenin on the Brain Transcriptome with Aging.* Mech Ageing Dev, 2023: p. 111889.

82. Wahl, D., et al., *Nontransgenic Guinea Pig Strains Exhibit Hallmarks of Human Brain Aging and Alzheimer's Disease.* J Gerontol A Biol Sci Med Sci, 2022. **77**(9): p. 1766-1774.

83. Wahl, D., et al., *The reverse transcriptase inhibitor 3TC protects against age-related cognitive dysfunction.* Aging Cell, 2023. **22**(5): p. e13798.

84. Leger, M., et al., *Object recognition test in mice.* Nature Protocols, 2013. **8**(12): p. 2531-2537.

85. Okun, E., et al., *Toll-like receptor 3 inhibits memory retention and constrains adult hippocampal neurogenesis.* Proc Natl Acad Sci U S A, 2010. **107**(35): p. 15625-30.

86. Bernier, M., et al., *Age-dependent impact of two exercise training regimens on genomic and metabolic remodeling in skeletal muscle and liver of male mice.* npj Aging, 2022. **8**(1): p. 8.

87. Hay, A.J.D., et al., *Intranasally delivered mesenchymal stromal cells decrease glial inflammation early in prion disease.* Front Neurosci, 2023. **17**: p. 1158408.

88. Chen, S., et al., *fastp: an ultra-fast all-in-one FASTQ preprocessor.* Bioinformatics, 2018. **34**(17): p. i884-i890.

89. Dobin, A., et al., *STAR: ultrafast universal RNA-seq aligner.* Bioinformatics, 2013. **29**(1): p. 15-21.

90. Love, M.I., W. Huber, and S. Anders, *Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2.* Genome Biology, 2014. **15**(12): p. 550.

91. Bourgon, R., R. Gentleman, and W. Huber, *Independent filtering increases detection power for high-throughput experiments.* Proc Natl Acad Sci U S A, 2010. **107**(21): p. 9546-51.

92. Law, C.W., et al., *RNA-seq analysis is easy as 1-2-3 with limma, Glimma and edgeR.* F1000Res, 2016. **5**.

93. Zhou, Y., et al., *Metascape provides a biologist-oriented resource for the analysis of systems-level datasets.* Nat Commun, 2019. **10**(1): p. 1523.