Altered spawning seasons of Atlantic salmon broodstock genetically and epigenetically influence cell cycle and lipid-mediated regulations in their offspring

Takaya Saito[1], Marit Espe[1], Maren Mommens[2], Christoph Bock[3], Jorge M.O. Fernandes[4], Kaja H. Skjaerven[1*]

[1] Institute of Marine Research, IMR, Bergen, Norway

[2] AquaGen, Trondheim, Norway

[3] CeMM, Research Center for Molecular Medicine of the Austrian Academy of Sciences, 1090, Vienna, Austria

[4] Nord University, Bodø, Norway

[*] Corresponding author

E-mail address: Kaja.Skjaerven@hi.no

Abstract

Manipulating spawning seasons of Atlantic salmon (*Salmo salar*) is a common practice to facilitate year-round harvesting in salmon aquaculture. This process involves adjusting water temperature and light regime to control female broodstock maturation. However, recent studies have indicated that altered spawning seasons can significantly affect the nutritional status and growth performance of the offspring. Therefore, gaining a deeper understanding of

the biological regulations influenced by these alterations is crucial to enhance the growth

performance of fish over multiple generations.

In this study, we investigated four different spawning seasons achieved through

recirculating aquaculture systems (RAS) and sea-pen-based approaches. In addition to the

normal spawning season in November (sea-pen), three altered seasons were designated as

off-season (five-month advance, RAS), early season (two-month advance, sea-pen), and late

season (two-month delay, sea-pen). We conducted comprehensive gene expression and DNA

methylation analysis on liver samples collected from the start-feeding larvae of the next

generation.

Our results revealed distinct gene expression and DNA methylation patterns

associated with the altered spawning seasons. Offspring from RAS-based off-season

spawning exhibited lower larval weights and showed altered lipid-mediated regulations,

which may potentially hinder growth performance. In contrast, early and late seasons affected

cellular processes, particularly cell cycle regulation, with implications for potential negative

impacts on future growth potential. The implications of our findings are significant for

aquaculture and nutritional studies, providing valuable tools for assessing growth potential

and optimizing production strategies.

Keywords

Broodstodk, Spawning season, Intergenerational, Nutrition, Omics, DNA methylation, Cell

cycle, Lipid metabolism

Introduction

The Atlantic salmon (*Salmo salar*) follows an anadromous life cycle, beginning in freshwater, migrating to seawater for most of its life, and returning to freshwater to spawn. In salmon aquaculture, the timing of spawning is adjusted to ensure a year-round availability for harvesting. This manipulation involves controlling abiotic factors such as water temperature and photoperiod to influence the spawning timing [1, 2]. Additionally, apart from the traditional open sea-based cages, a land-based approach utilising recirculating aquaculture systems (RAS) is employed to replicate the salmon's natural lifecycle more accurately, encompassing both saltwater gonad maturation and final freshwater maturation [3]. Despite the ongoing and successful practices of spawning season alterations in salmon aquaculture, recent studies have reported that these practices can impact the nutritional status of broodstock, leading to negative effects on the growth performance in the next generation [4, 5].

Nutritional programming, in the context of fish nutrition, refers to strategically manipulating the feeding regime of fish during critical developmental stages to enhance their long-term growth performance [6-8]. It can be expanded into intergenerational programming by providing specific nutrients or dietary components to the parents before reproduction [8, 9]. Hence, spawning season alterations of salmon broodstock can be seen as an example of intergenerational nutritional programming even though the nutritional manipulation occurs indirectly through abiotic factors. The principal target of the programming is to ensure the production of healthy and robust offspring in this context.

While various nutrients influence the health condition and growth performance of fish, including Atlantic salmon [10-12], altered spawning seasons have appeared to affect the status of multiple metabolites and nutrients including micronutrients, amino acids, and lipid

classes [4, 5]. These nutrients are also closely associated with vital biological functions

essential for growth, such as one carbon (1C) metabolism, the citric acid cycle, and the Cahill

cycle. Key micronutrients like vitamin B12, folate, vitamin B6, and methionine are integrated

with the 1C metabolism [13, 14], which also regulates cellular methylation potential through

S-adenosylmethionine (SAM) and S-adenosylhomocysteine (SAH), critical for various

cellular processes [15]. The citric acid cycle enables the catabolism of carbohydrates, fats,

and amino acids, releasing stored energy crucial for growth and development [16], while the

Cahill cycle facilitates the transport of amino groups and carbons between muscle and liver

[17], both playing essential roles in various developmental processes. Amino acids are

essential for growth and tissue development, in addition to being the building blocks of

proteins [18], while lipids serve as major energy reserves and contribute to the formation of

cell membranes, also supporting various developmental processes [19].

Beyond the importance of nutrients, understanding genetic influences within the

context of nutritional programming provides invaluable insights from a molecular biology

perspective. Recent studies have revealed that alterations in micronutrient levels within the

feed can have genetic effects on various biological pathways in salmon, including lipid

metabolism, protein synthesis, and post-transcriptional regulation [20, 21]. Moreover, an

additional layer of information comes from the analysis of DNA methylation, which offers

valuable epigenetic perspectives. DNA methylation, being an epigenetic regulatory process,

responds to environmental stimuli, governing cell stability, differentiation, and development

[22, 23]. As DNA methylation is inherited during cell replication, it can lead to changes in

epigenetic regulation over the long term [22, 24].

The present study investigates the genetic and epigenetic effects resulting from four

different spawning seasons. These seasons include the natural sea-pen-based spawning in

November, an accelerated RAS-based spawning occurring five months earlier, and two

variations of sea-pen-based spawning: one with a two-month advance and another with a two-month delay relative to the natural spawning season. The main analysis of the study focuses on liver samples collected from start-feeding larvae at the end of the endogenous feeding phase. The liver serves as a key target organ for various biological reactions crucial to growth, and using start-feeding larvae helps eliminate the influences of exogenous feeding. To uncover both genetic and epigenetic impacts comprehensively, we employed RNA-sequencing (RNA-seq) for gene expression and reduced-representation bisulfite sequencing (RRBS) for DNA methylation analysis. Our findings provide valuable insights into the significance of the nutritional state in broodstock, as it can affect the growth of their offspring. Additionally, our study offers potential usages of gene expression and DNA methylation data as powerful tools for assessing current and future growth performance, which can significantly benefit a wide range of nutritional programming studies.

Methods and Materials

Ethical considerations

The experiment adhered to the ARRIVE guidelines for design and reporting. Broodstock and embryos used in the study were obtained from AquaGen's commercial production facility of Atlantic salmon at their breeding station in Kyrksæterøra, Norway.

The conditions and protocols applied in this experiment were identical to those used in commercial production, including certain confidential information protected under commercial interests. Throughout the sampling procedures for broodstock and larvae, anesthetics were administered by the supplier's instructions and compliance with both Norwegian and European legislation on animal research.

Formal approval from the Norwegian Animal Research Authority (NARA) was not

required for this experiment. This exemption is because the experimental conditions fall

under practices recognized as commercial animal husbandry, thereby exempting them from

the European Convention on the protection of animals used for scientific purposes

(2010/63/EU), specifically under Article 5d. Additionally, the Norwegian ethics board also

approved the experiment by the Norwegian regulation on animal experimentation, § 2, 5a, d,

for activities classified as "non-experimental husbandry (agriculture or aquaculture)" and

"procedures in normal/common breeding and husbandry".

Experimental design and sampling

As the details of experimental design and sampling methods have been described elsewhere

[4, 5], we only present summarized descriptions here. The study utilized broodstock and

embryos obtained from AquaGen's breeding station in Kyrksæterøra, Norway. The female

broodstock, sourced from the 15th generation, covered four spawning seasons: off-season,

early, normal, and late seasons.

The off-season broodstock were reared in a land-based recirculating aquaculture

system (RAS) in saltwater (12% salinity), while the rest of the broodstock were raised in sea-

based net pens. All broodstock were fed to satiation with broodstock feed (EWOS Breed

3500) until transferred to freshwater. The normal season broodstock matured naturally under

ambient photoperiod and temperatures, whereas artificial controls were applied to regulate

maturation for the other spawning seasons. Specifically, the off-season and early season

broodstock were subjected to five- and two-month advance maturation, respectively, while

the late season broodstock were subjected to a two-month delay. The fish received no further

feeding after the transfer to freshwater. The maturation process in freshwater lasted for 109,

163, 120, and 166 days for off-season, early, normal, and late season broodstock,

respectively, before fertilization to obtain the next generation (Supplementary Table S1).

From each spawning group, five females from separate tanks were sacrificed using

Benzoak (200 mg/L, ACD Pharmaceuticals AS). After stripping, each female's growth

measures, including body weights (n=5 per spawning season), were recorded. Liver samples

were dissected and flash frozen in liquid N2 for nutrient analysis. All oocytes were fertilised

with cryopreserved sperm from two pooled males. Newly fertilized eggs (2 d°) were flash-

frozen in liquid N2 for nutrient analysis (Supplementary Table S1).

At the end of the endogenous feeding phase (start-feeding larvae 979–994 d°;

Supplementary Table S1), 36 larvae from each broodstock were randomly selected,

euthanised with an overdose of buffered tricane methanesulfonate (MS222, Pharmaq,

Norway), and their growth measures, including body weights (n=180 per spawning season),

were recorded. Additionally, liver samples were dissected from larvae and flash frozen in

liquid N2 for RNA and DNA analysis.

Growth measure and nutrient analysis

We collected the data of body weights for broodstock and larvae from previous studies [4, 5].

In addition, we obtained the data on selected nutrients and metabolites from the same studies,

specifically vitamin B6 & B12, folate, S-adenosylmethionine (SAM), S-

adenosylhomocysteine (SAH), cholesterol, the sum of six different lipids, glutamate, L-

serine, L-lysine, L-alanine, L-glutamine, urea, and B-alanine [4, 5]. To analyse the data from

all four spawning seasons, we performed ANOVA followed by Tukey's HSD test.

DNA and RNA extraction

RNA extraction was carried out using the BioRobot EZ1 and EZ1 RNA Universal Tissue kit (Qiagen), followed by DNase treatment using the Ambion DNA-free DNA removal kit (Invitrogen, USA) according to their respective protocols. The RNA quantity was assessed using the NanoDrop ND-1000 Spectrophotometer (Nanodrop Technologies) and the Agilent 2100 Bioanalyzer with the RNA 6000 Nano LabChip kit (Agilent Technologies). For DNA isolation, the DNeasy Blood & Tissue Kit (Qiagen, Cat. No. #69506) was used as per the manufacturer's protocol. Liver samples underwent RNase A treatment (provided by the Qiagen kit, 50ng/µL, 10 min at room temperature) followed by proteinase K treatment (New England Biolabs, #8102S 20µg/µL, 1.5 h at 55°C). DNA was eluted in Milli Q water. DNA quantification was performed using the Qubit High Sensitivity Assay (Life Technologies #Q32854). Detailed methods for both DNA and RNA extraction were described elsewhere [20].

Library preparation of high throughput sequencing

Liver samples were sequenced at two different sequencing facilities for specific analyses. For RNA-seq, the liver samples were processed at the DeepSeq sequencing facility at Nord University, Bodø, Norway, and the library preparation was conducted using the NEBNext Ultra II Directional RNA Library Prep Kit for Illumina (New England Biolabs). Sequencing was performed on the NextSeq500 machine (Illumina). For RRBS, the liver samples were processed at the CeMM Biomedical Sequencing Facility, Vienna, Austria. The genomic DNA was extracted, digested by MspI, and bisulfite-converted before library preparation. The RRBS libraries were sequenced on Illumina HiSeq 3000/4000 instruments. Detailed methods of the library preparation for RNA-seq and RRBS were described elsewhere [20].

Atlantic salmon genome and genomic annotation

The reference genome (ICSASG version 2) and RefSeq data (version 100) for gene
annotation were downloaded from the NCBI web site
(https://www.ncbi.nlm.nih.gov/assembly/GCF_000233375.1). In cases where gene symbols
were either outdated or unavailable from RefSeq data (version 100), gene symbols from a
newer version of RefSeq data (version 102) and UniProt [25] were used. The genome regions
were annotated into intron, exon, three promoter regions, and flanking regions. Promoter
regions were divided into three categories based on their distance from transcription start
sites: P250 (1-250 bp), P1K (251-1000 bp), and P5K (1001-5000 bp). Flanking regions were
defined as 5000 upstream from P5K and 10000 downstream from the 5' end of the gene.

Pre-processing of high throughput sequencing

For both RNA-seq and RRBS data, the same pre-processing procedures were followed as
described in a previous study [20]. This involved trimming the reads using Cutadapt [26] and
Trim Galore! (Barbraham Institute), aligning the trimmed reads to the reference genome
using STAR [27] for RNA-seq and Bismark/Bowtie 1 [28, 29] for RRBS with their default
parameters. The mapped RNA-seq reads were quantified using featureCounts [30], while the
mapped RRBS reads were processed by Bismark [28] for methylation calling and CpG site
extraction. The samples were divided into four spawning season groups for further analysis:
off-season, early, normal, and late seasons, and clustering analysis was performed using the
factoextra package (https://CRAN.R-project.org/package=factoextra). RNA-seq counts were
subjected to a variance stabilizing transformation (VST) using DESeq2 [31] before principal
component analysis (PCA).

206   Raw reading data were uploaded to the repository of the Sequence Read Archive

207   (SRA) on the NCBI web site and available under the accession numbers PRJNA680425 and

208   PRJNA642998 for RNA-seq and RRBS, respectively.

Differential gene expression analysis

Differentially expressed genes (DEGs) were identified using DESeq2 [31] with an adjusted p-

value cut-off of less than 0.05. P-values were adjusted by the Benjamini-Hochberg procedure

[32]. From the candidate DEGs, those with absolute log fold changes (LFCs) greater than or

equal to 1.2 were selected as the final set of DEGs by using the lfcThreshold argument of the

results function of DESeq2 [31]. Shrunken LFCs values, calculated by the normal shrinkage

method provided by DESeq2 [31], were used for both heatmaps showing functional

annotation results and scatter plots showing merged results between DNA methylation and

gene expression differences.

Functional annotation with KEGG

219   Over-representation analysis (ORA) was performed on the Kyoto Encyclopedia of Genes and

220   Genomes (KEGG) database [33] using clusterProfiler [34]. Gene lists of DEGs identified

221   from three pair-wise comparisons were used as input, and enriched pathways were

222   determined based on adjusted p-values less than 0.05. P-values were adjusted by the

223   Benjamini-Hochberg procedure [32].

Methylation rate analysis

Mapped RRBS reads were filtered by methylKit [35], discarding reads with coverage less

than or equal to 10 and above the 99.9th percentile. Additional RRBS data from two other

studies were obtained for post-smolt and harvesting stages [20, 21] and underwent the same

pre-processing procedures to compare the distributions of methylation rates. Raw data of

these RRBS sequencing data can be found in SRA on the NCBI site under the accession

numbers of PRJNA680423 and PRJNA628740 for post-smolt and harvesting stages,

respectively. Ridge plots were generated using the ggridges R library (https://CRAN.R-

project.org/package=ggridges).

Differential methylation analysis

Differentially methylated CpG sites (DMCs) were identified by methylKit [35], and Q-values

were calculated using the sliding linear model (SLIM) method [36]. CpGs were identified as

DMCs when Q-values were less than 0.01 and methylation differences were greater than or

equal to 15%.

To filter out DMCs that were strongly affected by altered spawning seasons, we used

the counts of DMCs per gene in two promoter regions: P250 and P1K. Genes were sorted by

the count of DMCs in a descendant order, separately for P250 and P1K, and genes that

contained at least three DMCs were selected for further analysis.

Bioinformatics analysis

In-house R and Python scripts with Snakemake [37] were utilized for high-throughput

sequence analysis, basic statistical analysis, and figure generation.

Results

Altered spawning seasons impacted the weight and nutritional status of

offspring

The present study examines the effects on gene expression and DNA methylation patterns in

the liver of offspring hatched from four different spawning seasons: an off-season based on a

recirculating aquaculture system (RAS) and three sea-pen based seasons (early, normal, and

late) (Fig. 1a). Off-season spawning occurred in June, early season in September, and late

season in January, relative to the normal spawning months of November. (see detailed dates

and degree days in Supplementary Table S1). As the growth performance and nutritional

status of the broodstock and offspring have already been assessed and discussed elsewhere [4,

5], this section only provides a summary emphasizing the weight and the status of selected

nutrients and metabolites.

Fig. 1. Experimental design, body weights, and nutritional status across spawning seasons.

a) Schematic diagram illustrating four distinct spawning seasons: off-season (O), early (E), normal (N), and late
(L). Off-season was conducted in RAS, while others were in a sea-based environment. Broodstock were
transferred from seawater (blue bars) to freshwater (light blue bars) before spawning. Offspring were raised in
freshwater until sampling (green bars). Triangles indicate sampling points: broodstock (blue), eggs (red), and
larvae (green). **b)** Boxplots showing body weights of broodstock (n=5) and larvae (n=180). Red letters represent
Tukey's HSD compact letter display after ANOVA tests (p<0.05). **c)** Boxplots of 12 nutrients and metabolites in
egg samples. Nutrients and metabolites are categorised into four groups: 1C metabolism, lipid classes, citric acid
cycle, and Cahill cycle. Units: 1C metabolism group (Vitamin B6: mg/kg ww, Folate: μg/100 g ww, Vitamin
B12: μg/kg ww); remaining groups: lipid classes (mg/g ww), citric acid cycle (μmol/g ww), Cahill cycle
(μmol/g ww). Red letters represent Tukey's HSD compact letter display after ANOVA tests (p<0.05).

Broodstock weights showed no significant differences at spawning (upper part of Fig.

1b), but offspring weights at the larvae stage (979 and 994 degree days) showed noticeable

variation (lower part of Fig. 1b). Specifically, off-season and early season larvae had lower

weights compared to normal and late seasons. This trend was consistent with the observations

in egg sizes, where eggs from off-season and early-season exhibited smaller sizes compared

to the other two seasons (Supplementary Table S2).

Altered spawning seasons widely impacted the nutritional status of offspring eggs,

particularly regarding 1C metabolism, lipid classes, citric acid cycle, and Cahill cycle [4, 5],

but with inconsistent patterns across nutrients and metabolites (summarised in Fig. 1c and

Supplementary Tables S3-S6). Previous studies also reported that altered spawning seasons

affected the nutritional status of the broodstock [4, 5]. Thus, altered spawning seasons

significantly influenced the nutritional status of broodstock without impacting their weights,

subsequently affecting offspring nutritional status and growth potential of their offspring.

To investigate the underlying genomic and epigenetic regulations associated with the

observed differences in nutritional status and growth performance, we utilised RNA-seq for

gene expression analysis and RRBS for DNA methylation analysis on liver samples collected

from larvae at the end of the endogenous feeding phase.

288 Early and late spawning seasons exhibited similar gene expression patterns
289 when compared to the normal season

We performed gene expression analysis on 20 liver samples divided into four groups (n=5) to

investigate the impact of different spawning seasons to the offspring when compared to the

normal season (detailed statistics for each sample are provided in Supplementary Table S7).

Principal component analysis (PCA) differentiated the off-season from the normal

season when considering all mapped genes (left part of Fig. 2a). However, when focusing on

the top 500 genes with high variances, the early season moderately deviated from the normal

season (right part of Fig. 2a). In both cases, the late season fell between the normal and early

seasons.

Fig. 2. Gene expression analysis comparing altered spawning seasons to the normal season.

a) PCA plots displaying clustering patterns of four spawning seasons - off-season (red), normal (green), early
(blue), and late (purple) - using 20 RNA-seq samples. "All mapped" plot includes 45,153 genes, while "Top
500" includes top 500 genes with high variance. **b)** Volcano plots showing differential expression analysis
results for three pairwise comparisons: off-season vs. normal (Off-season), early vs. normal (Early), and late vs.
normal (Late). Dots represent down-regulated DEGs (blue), up-regulated DEGs (red), and non-DEGs (grey).
Counts of DEG and non-DEGs were provided in the legend. **c)** Venn diagrams illustrating overlaps of DEGs
identified between two comparisons, indicating common DEGs as well as those exclusive to a single
comparison.

In three pairwise comparisons for differential expression analysis (off-season vs.

normal, early vs. normal, and late vs. normal), we identified 173, 337, and 265 differentially

expressed genes (DEGs), respectively (Fig. 2b). All comparisons revealed a higher number of

down-regulated genes than up-regulated genes. Additionally, Venn diagrams indicated that

the DEGs from the off-season vs. normal comparison had very few overlapping genes (<1%)

with the other two comparisons (depicted in the first and second diagrams in Fig. 2c). On the

other hand, early vs. normal and late vs. normal comparisons shared approximately 25% of

the DEGs (depicted in the third diagram in Fig. 2c).

The results from PCA and differential expression analysis suggest that the expression

pattern of the off-season differed from both the early and late seasons when compared to the

normal season. Furthermore, the early and late seasons exhibited similar gene expression

patterns, with the late seasons showing more resemblance to the normal seasons.

Altered spawning seasons affected various biological pathways, including those related to metabolism, cellular processes, and organismal systems

To explore the impact of altered spawning seasons on biological pathways, we conducted

functional analysis on DEGs using the Kyoto Encyclopedia of Genes and Genomes (KEGG)

database. Over-representation analysis (ORA) identified a total of 11 enriched KEGG

pathways, which were classified into three functional categories: metabolism, cellular

processes, and organismal systems (Fig. 3).

Fig. 3. Enriched KEGG pathways for DEGs identified in three pair-wise comparisons.

Heatmaps showing log fold changes (LFCs) of genes associated with 11 KEGG pathways enriched through
over-representation analysis. LFCs underwent normal shrinkage transformation. Genes were included if
identified as DEGs in at least one of the three comparisons: off-season vs normal (Off), early vs normal (Early),
and late vs normal (Late). Number below each pathway represents DEG count. KEGG pathways are categorized
into metabolism, cellular processes, and organismal systems groups.
335

In the off-season vs. normal comparison, ORA identified only one enriched pathway,

carbon metabolism (Supplementary Table S8). All five DEGs associated with carbon

metabolism showed down-regulation (Fig. 3).

For the early vs. normal comparison, ORA revealed six enriched pathways: one in

metabolism, three in cellular processes, and two in organismal systems (Supplementary Table

S9). Metabolism-related pathways exhibited both up- and down-regulation, while most genes

in cellular processes showed significant down-regulation. In organismal systems, PPAR

signalling showed up-regulation, while progesterone-mediated oocyte maturation showed

down-regulation (Fig. 3).

The late vs. normal comparison identified nine enriched pathways: three in

metabolism, four in cellular processes, and two in organismal systems (Supplementary Table

S10). Remarkably, this comparison shared five enriched pathways with the early vs. normal

comparison, indicating similar impacts on biological pathways between them. Moreover,

these pathways displayed consistent expression patterns in terms of up- and down-regulation

between early and late seasons. Notably, genes associated with cellular processes exhibited

strong down-regulation for both seasons (Fig. 3).

Altered spawning seasons affected multiple pathways in metabolism, cellular

processes, and organismal systems. The off-season appeared to down-regulate carbon

metabolism, while the early and late seasons primarily had strong suppressive effects on

pathways associated with cellular processes. These findings may be linked to the reduced

weight of off-season and early season larvae, as well as variations in the nutritional status of

the eggs.

Altered spawning seasons had an impact on the expression of genes in a similar manner

To further investigate the genes commonly affected by multiple spawning seasons, we used

overlapping DEGs (represented as intersections of Venn diagrams in Fig. 2c) by applying two

filtering steps: selecting DEGs found in at least two comparisons and choosing the top five

DEGs from each comparison based on adjusted p-values (Supplementary Table S11). This

filtering approach revealed a total of 13 genes (Table 1). Among them, two genes (gene IDs:

106612264 and 106601246) lacked corresponding gene names and symbols. All genes,

except *pc* and *pitpnm2*, consistently displayed up- or down-regulation across all spawning

seasons, suggesting that all altered spawning seasons had a similar effect on this specific set

of genes (Table 1).

369
370 **Table 1**. List of top five DEGs identified in at least two pairwise comparisons.
371

Pattern[1]	Gene ID	Symbol	LFC[2]			Gene name[3]	Function
			Off-season	Early	Late		
All Down	106563634	*pcyt1b*	-1.9 *	-1.1	-1.6 *	choline-phosphate cytidylyltransferase B	Metabolism
	106612672	*bmp5*	-1.2 *	-1.2 *	-0.7	bone morphogenetic protein 5-like	Signalling
	106585803	*rhobtb4*	-0.3	-1.1 *	-0.9 *	Rho related BTB domain containing 4	Ubiquitin mediated proteolysis
	106608456	*prox3*	-0.2	-2.3 *	-1.7 *	prospero homeobox 3	Lymphangiogenesis
	106584736	*tcf12*	-0.7	-3.3 *	-2.3 *	transcription factor 12	Myogenic regulatory
Up & Down	106592454	*pc*	-1.7 *	1.5 *	0.9	pyruvate carboxylase, mitochondrial-like	Pyruvate metabolism, TCA cycle, redox
	106568312	*pitpnm2*	0.6	-2.1 *	-2.6 *	membrane-associated phosphatidylinositol transfer protein 2-like	Cell signalling
All Up	106612264	-	5.2 *	4.1	6.1 *	Uncharacterized	-
	106601246	-	1.8 *	2.2 *	1.5	Uncharacterized	-

106613828	crim1	0.3	3.0	*	3.0	*	cysteine-rich motor neuron 1 protein-like	CNS development,Bone morphogenic and growth factor
106566973	krt8	0.1	1.3	*	1.0	*	keratin, type II cytoskeletal 8	Liver development
106583637	klf15	2.3	4.5	*	3.8	*	Krueppel-like factor 15	Metabolism
106569008	cyp7a1	1.4	3.1	*	2.9	*	cytochrome P450 7A1	Metabolism

[1]All Down: All three comparisons showed down-regulation, Down+Up: Comparisons showed both down- and up-regulation. All Up: All three comparisons showed up-regulation. [2]Log fold changes of pairwise comparison against the normal season. *: significant LFC (adjusted p-value<0.05). [3]When gene symbols were unavailable, we utilised the gene symbols from either the latest NCBI version or UniProt as substitutes.

These genes can be classified into three categories by relevant biological functions: metabolism (*pcyt1b*, *pc*, *klf15*, and *cyp7a1*) [38-40], tissue development (*prox3*, *crim1*, and *krt8*) [41-44], and information processing, which can be further classified into three sub categories as signalling (*bmp5* and *pitpnm2*) [38, 45], transcription (*tcf12*) [46], and post-translation (*rhobtb4*) [38]. As these categories are typically associated with growth, all three altered spawning seasons affected a certain set of genes associated with growth regulation, despite PCA plots (Fig. 2a) and Venn diagrams (Fig. 2c) showing a distinct gene expression pattern of the off-season.

Overall methylation rates increased until the harvesting stage, except around transcription start sites

We conducted DNA methylation analysis on 20 liver samples from larvae across four different spawning seasons (n=5; detailed statistics in Supplementary Table S12) to investigate the epigenetic effects of altered spawning seasons on DNA methylation. To examine regional DNA methylation patterns, CpG sites were categorised into three genomic regions based on their positions relative to closely located genes: gene body (GB; including exons and introns), promoter (P; including P250, P1K, and P5K), and flanking regions (Fig. 4a; see Methods and Materials for definitions).

Fig. 4. Distribution of methylated CpG sites across genomic regions.

a) Genomic region diagram illustrating three regions: flanking regions (Flank, red), promoters (P, blue), and gene bodies (GB, green). Promoter region is divided into three subregions: P250, P1K, and P5K. Gene body includes exons and introns. **b)** Ridge density plots showing distribution patterns of methylated CpG sites in percentage across six genomic regions. Plot includes methylation data from the present study (larvae, green) and two other studies: post-smolt (light blue) and harvest (blue) for comparison. **c)** Ridge density plots showing distribution patterns of methylated CpG sites for four spawning seasons: off-season (red), early (green), normal (blue), and late (purple). Plot is divided into two ranges: 0-25% and 75%-100%.

Before analysing the effects of altered spawning seasons, we compared RRBS data from two other studies focusing on liver samples from different developmental stages: post-smolt (~31 weeks) and harvest (~45 weeks) [20, 21]. Distributions of methylation rates showed significant differences among life stages, except for P250 (Kolmogorov–Smirnov test; Supplementary Table S13). Ridge density plots displayed peak values ranging from 75% to 100%, excluding P250 and P1K (Fig. 4b). Notably, the peaks of the harvesting stage shifted closer to 100% compared to the larvae and post-smolt stages, supporting the trend of average methylation rates: larvae < post-smolt < harvest (average methylation rates are shown in text labels of Fig. 4b). This comparison suggests that methylation rates generally

414   increase as salmon grow, at least until the harvesting stage, while methylation rates near

415   transcription start sites (TSSs) remain lower and more stable than in other genomic regions.

Altered spawning seasons contributed to similar shifts in DNA methylation rates as observed in growth performance

We analysed distributions of methylation rates across spawning seasons and observed

significant differences among them, except for P250 and P1K (Kolmogorov–Smirnov test;

Supplementary Table S14 and Supplementary Fig. S1). To highlight peaks across seasons, we

divided the ridge density plot into 0-25% and 75-100% sections (Fig. 4c).

Even though distributions in the 0-25% range were similar, significant differences

were evident in multiple comparison to some degree (Kolmogorov–Smirnov test;

Supplementary Table S15). Notably, the comparison between early and late seasons showed

significant differences in all regions except for P250. Moreover, peaks in the early season

were further from 0% compared to other seasons, supporting the trend of average methylation

rates in the 0-25% range (average methylation rates are shown as text labels in the left side of

Fig. 4c).

In the 75-100% range, distributions significantly differed among spawning seasons,

excluding P250 and P1K (Kolmogorov–Smirnov test; Supplementary Table S16). Notably,

peaks in normal and late seasons shifted closer to 100% compared to the off-season and early

season, except for P250, supporting the trend of average methylation rates: early ≈ off-season

< normal < late (average methylation rates are shown as text labels in the right side of Fig.

4c). This order closely aligns with weight measurements observed at the larvae stage (Fig. 1b,

Supplementary Table S2). Considering both these comparisons and those across different life

stages, methylation rates are linked to growth with region-specific patterns of methylation

shifts.

Differential methylation patterns indicated higher dynamic regulation in promoter regions compared to other regions

We conducted six pairwise comparisons to identify CpG sites with differential methylation (Fig. 5a). Our objective was to capture a comprehensive pattern of methylation differences by considering all possible pair-wise combinations. Remarkably, all comparisons consistently demonstrated similar characteristics, for both the number of differentially methylated CpGs (DMCs) and the balanced ratio of hypo- and hyper-methylation (Fig. 5a).

Fig. 5. DNA methylation differences and correlations with gene expression.

a) Histograms displaying distributions of methylation rate differences in six pairwise comparisons. Hypo-methylated DMCs in blue, hyper-methylated DMCs in red, non-DMCs in grey. All CpG sites in plots showed significant methylation differences. **b)** Ridge density plots illustrating distribution patterns of methylated CpG sites across four spawning seasons: off-season (red), early (green), normal (blue), and late (purple). CpG sites were selected if at least one site identified as DMC in any of six comparisons. **c)** Scatter plots showing relationship between methylation rate differences and log fold changes (LFCs) of gene expression. DMCs were selected from three comparisons: off-season vs. normal (Off season), early vs. normal (Early), and late vs. normal (Late), and also selected based on two regions, P250 and P1K. Corresponding genes were linked to these DMCs chosen, and their LFCs were transformed by normal shrinkage. Data points are color-coded to indicate

Further investigation focused on a set of 10,308 common DMCs, identified in at least

one comparison. Distributions of methylation rates for these common DMCs exhibited a

distinct left-skewed pattern across the exon, intron, P5K, and flanking regions (Fig. 5b). This

observation strongly suggested that methylation differences resulting from altered spawning

seasons primarily occurred within regions that were already highly methylated. In contrast,

the P250 and P1K regions displayed nearly uniform methylation rates across a range of 0-

100%, indicating significant methylation differences occurred irrespective of the underlying

methylation rates. These findings imply a potentially greater level of dynamic regulation of

DNA methylation in P250 and P1K compared to the other regions.

Filtering multiple DMCs identified key genes affected by spawning season alterations

We examined genes with multiple DMCs in three pairwise comparisons against the normal

season. Since most genes had only a single DMC, especially in promoters (~75% in P250 and

~70% in P1K, Supplementary Fig. S2), we avoided identifying differentially methylated

regions (DMRs) using a sliding window approach. Instead, we filtered out genes strongly

supported by multiple DMCs based on specific criteria (see Methods and Materials), resulting

in a total of 11 genes (Table 2), including six genes for off-season vs. normal and three genes

each for the other two comparisons (Supplementary Table S17).

476
477   **Table 2**. List of genes with multiple DMCs in their promoter region.
478

Pattern[1]	Gene ID	Symbol	DMC[2]			Gene name	Function
			Off-season	Early	Late		
All hypo	106563027	*slc31a1*	Hypo (3) *	-	Hypo (3) *	solute carrier family 31 member 1	Transporter

Pattern	ID	Gene	Comp 1	Comp 2	Comp 3	Description	Function
Hyper & Hypo	106567512	stk32a	-	Hypo (2)	Hypo (2), * Hyper (1)	serine/threonine-protein kinase 32A	Signalling
	106607889	inf2	Hypo (1)	Hyper (3) *	-	inverted formin-2-like	Actin filament polymerization
	106584059	scp-2	Hyper (4) *	Hypo (1)	Hyper (1)	sterol carrier protein 2-like	Lipid transfer, transporter
	106579168	klhl11	Hypo (1), * Hyper (2)	-	-	kelch-like protein 11	Ubiquitination
All hyper	100196480	ctl2b	Hyper (3)	Hyper (7) *	Hyper (2)	CTLA-2-beta	Unknown
	106573279	-	-	Hyper (1)	Hyper (3) *	Uncharacterized	-
	106583651	helz2	Hyper (4) *	-	-	helicase with zinc finger domain 2-like	Transcriptional coactivator
	106581656	lrrfip2	Hyper (3) *	-	-	leucine-rich repeat flightless-interacting protein 2-like	Signalling
	106609256	prrx2	Hyper (3) *	-	-	paired mesoderm homeobox protein 2-like	Transcription
	106576160	ttll12	-	Hyper (3) *	-	tubulin tyrosine ligase-like family, member 12	Post-translational modification

[1]All hypo: All three comparisons showed hypo-methylated DMCs, Hyper & Hypo: Comparisons showed both hypo- and hyper-methylated DMCs. All hyper: All three comparisons showed hyper-methylated DMCs. [2]Either hyper or hypo-methylated DMC against the normal season. The number in brackets represent the number of DMCs. *: genes selected by filtering of multiple DMCs.

All 11 genes were identified by a single comparison except for *slc31a1* identified in two comparisons (off-season vs. normal and late vs. normal). Most genes (9 out of 11) exhibited either hypo-methylated or hyper-methylated DMCs, but not both, within a single comparison (Table 2). Furthermore, approximately 60% of the genes (7 out of 11) consistently displayed either hypo-methylation or hyper-methylation even across multiple comparisons ('All hypo' and 'All hyper' in the pattern column of Table 2, respectively).

Literature analysis associated these genes with two biological functions: development (*inf2*) and information processing, which can be further classified into four sub categories as transporter (*slc31a1* and *scp-2*) [47, 48], signalling (*stk32a* and *lrrfip2*) [25, 49], post-

494 translation (*klhl11* and *ttll12*) [25, 50], and transcription (*helz2* and *prrx2*) [51, 52]. However,

495 *ctl2b* and an uncharacterised gene (Gene ID: 106573279) had no clear associations with

496 specific biological functions. Although these genes were potentially linked to growth

497 performance, none of the 11 genes were identified as DEGs in our gene expression analysis.

498 Altered spawning seasons influenced gene expression and DNA methylation

499 of cell cycle-related genes

To investigate potential associations between DNA methylation and gene expression, we

focused on DMCs identified in three pairwise comparisons against the normal season (off-

season vs. normal, early vs. normal, and late vs. normal) in P250 and P1K. By correlating

their methylation differences with corresponding log-fold changes (LFCs), we found a total

of 11 DMCs: four DMCs in P250 and seven DMCs in P1K associated with DEGs (Fig. 5c,

Supplementary Table S18). Surprisingly, CpG sites with methylation differences greater than

30% lacked clear association with differentially expressed genes (light grey dots labelled as

non-DEGs in Fig. 5c), suggesting that differentially methylated sites showed inconsistent

links with differential gene expression.

These 11 DMCs were associated with seven genes that were DEGs in the

corresponding comparisons (Table 3). We expanded this list of seven genes with additional

information from all pair-wise comparisons (additional entries were marked with parentheses

in Table 3). Among them, three genes (*caprin-1*, *kifc1*, and *aurkb*) exhibited consistent

regulations of DMCs and DEGs compared to the normal season. Specifically, *caprin-1* and

kifc1 had hypo-methylated CpG sites, while *aurkb* had hyper-methylated CpG sites; all three

genes showed down-regulated gene expression (Table 3). Notably, all three genes (*caprin-1*,

kifc1, and *aurkb*) were associated with cell cycle regulation [53-55], while the remaining

genes were linked to metabolism (*cyp8b1*, *adrenodoxin*, and *lpin1*) [33, 56, 57] and transport

(*slc43a1a*) [58]. These findings indicate that altered spawning seasons had genetic and

epigenetic effects on several genes associated with cell cycle regulation.

Table 3. List of DEGs that contain DMCs in their promoter region.

Gene ID	Symbol	DMC[1]			DEG[2]			Gene name	Function
		Off-season	Early	Late	Off-season	Early	Late		
106562317	*caprin-1*	Hypo	(Hypo)	(Hypo)	Down	(Down)	(Down)	caprin-1	Cell cycle
106599887	*cyp8b1*	Hypo	-	-	Down	-	-	5-beta-cholestane-3-alpha,7-alpha-diol 12-alpha-hydroxylase	Metabolism, cytochrome P450
106588407	*kifc1*	-	Hypo	Hypo	-	Down	Down	carboxy-terminal kinesin 2	Cell cycle, Meiosis
106582038	*adrenodoxin*	-	Hypo		-	Down	-	adrenodoxin	Metabolism, cytochrome P450
106604665	*slc43a1a*	-	Hypo	-	-	Up	(Up)	solute carrier family 43 member 1a	Transporter
106561604	*aurkb*	(Hyper)	Hyper	Hyper	-	Down	Down	aurora kinase B	Cell cycle
106570052	*lpin1*	-	Hyper	-	-	Up	-	phosphatidate phosphatase LPIN1	Metabolism

[1]Either hypo- or hyper-methylated DMC against the normal season. Parentheses indicate that the site was identified as DMC, but the corresponding genes was not identified as stringent DEG. [2]Either up- or down-regulation against the normal season. Parentheses indicate that down- or up-regulation was determined by non-stringent DEGs.

Discussion

In the present study, we investigated the influence of altered spawning seasons on gene

expression and DNA methylation in offspring. Previous studies emphasized the significant

impacts of these changes on the nutritional status of both broodstock and their offspring, with

no observed effects on the body weights of broodstock [4, 5]. Our primary aim was to reveal

534   the genetic and epigenetic effects resulting from observed variations by utilizing liver

535   samples from larvae at the first feeding stage.

What insights do our omics analyses provide?

Gene expression analysis revealed distinct patterns associated with altered spawning seasons.

Specifically, early and late seasons exhibited similar gene expression patterns when

compared to the normal season. The differential expression of genes associated with

metabolism, cellular processes, and organismal systems indicates that spawning seasons can

have wide-ranging effects on biological pathways related to growth and development.

Furthermore, the analysis with overlapped DEGs revealed that all altered spawning seasons

affected genes associated with metabolism, tissue development, and information processing

similarly. Hence, while early and late seasons influenced a wide range of biological pathways

in a similar manner, all three seasons, including the off-season, also similarly affected a

certain set of genes, especially those strongly differentiated from the normal season.

DNA methylation analysis provided valuable insights into the epigenetic effects of

different spawning seasons. Altered spawning seasons led to increased methylation rates,

which appeared to be linked to higher body weights. Notably, these shifts primarily occurred

in regions that were already highly methylated, but the rates were altered independently of

the underlying methylation rates in promoter regions around transcription start sites. Our

findings imply that methylation rates could serve as a measure of the current growth levels,

even when comparing different treatment groups at the same developmental stage.

Several genes with multiple DMCs exhibited consistent hypo- or hyper-methylation

patterns across multiple spawning seasons in promoter regions, suggesting a common

epigenetic response to altered spawning seasons. These genes were associated with key

biological functions, such as development and information processing, further supporting

their role in influencing growth performance. Although both gene expression and DNA

methylation analyses resulted in similar biological functions impacted by altered spawning

seasons, they showed little overlap between DEGs and genes with DMCs. This could be

attributed to regulatory differences between gene expression and DNA methylation within the

same biological pathway, or it might be a limitation caused by the restriction enzyme used in

RRBS.

The observed correlations between DNA methylation and gene expression were

weaker than anticipated in our liver samples. Notably, CpG sites with methylation differences

exceeding 30% lacked a clear link to differential gene expression. Additionally, our findings

revealed inconsistent correspondence between hypo-methylation and active gene expression

and between hyper-methylation and gene suppression. These results suggest that the

regulation of DNA methylation is more intricate than commonly assumed as it is known that

high promoter CpG methylation reduces gene expression, while low promoter CpG

methylation allows for active gene expression [59]. Within promoter regions, several factors,

including underlying gene expression levels, transcription factor binding sites, and chromatin

remodelling, likely play a role in regulating actual gene expression.

What are the significant biological differences observed in offspring?

Off-season spawning in June, facilitated through a RAS, resulted in Atlantic salmon offspring

with lower larval weights compared to normal and late seasons. Offspring eggs also exhibited

reduced levels of vitamin B12 and lipid classes relative to normal and late seasons. Gene

expression analysis indicated significant down-regulation of genes associated with central

carbon metabolism (CCM). CCM involves several biological pathways, including the citric

acid cycle, and acts as a major source of energy for growth and development. Moreover,

lipids are interconnected with CCM through the utilisation of acetyl-CoA as a central

metabolite [60]. Among the six genes with multiple DMCs from the off-season vs. normal

comparison, sterol carrier protein2-like (*scp-2*) contained four hyper-methylated DMCs in its

promoter. The *scp-2* gene encodes a transfer protein that plays a key role in intracellular lipid

transport [48]. Overall, RAS-based off-season spawning appeared to impact lipid-mediated

regulations both genetically and epigenetically. This negative impact on lipid-related

mechanisms potentially accounted for impaired growth performance observed in body

weight.

Offspring resulting from early season spawning in September also exhibited lower

weights at the larval stage compared to those from normal and late seasons. However, the

nutritional status of offspring eggs displayed higher levels of SAM/SAH, lysine, glutamine,

and alanine. Gene expression analysis revealed significant down-regulation of genes

associated with cell cycle regulation, while genes involved in metabolism showed both down-

and up-regulation. Among the five genes that exhibited differential expression and had at

least one DMC in their promoters, two of them were associated with cell-cycle regulation.

Specifically, the carboxy-terminal kinesin 2 (*kifc1*) gene promotes mitotic spindle assembly

[54], and the aurora kinase B (*aurkb*) gene is involved in mitotic progression [55]. The *kifc1*

gene was hypomethylated, while the *aurkb* gene was hypermethylated. Consequently, the

early spawning season had an impact on cell-cycle regulation both genetically and

epigenetically. However, it is challenging to hypothesise that this negative impact on cell-

cycle regulation was linked to impaired growth performance because similar effects were

observed for the late season.

Late season spawning in January showed that offspring weights were almost as close

to, but slightly heavier than those from the normal season. The nutritional status of offspring

eggs also exhibited similar levels of various nutrients and metabolites to the normal season,

except for higher vitamin B12 and lower B-alanine. Gene expression patterns in the late

season were comparable to those observed in the early season, showing significant down-

regulation of genes related to cellular processes, especially cell cycle regulation. Two genes,

kifc1 and *aurkb*, showed differential expression and had at least one DMC in their promoters,

and both were associated with cell-cycle regulation. Hence, the late season showed similar

growth performance, nutritional status, and global DNA methylation pattern to the normal

season, while gene expression pattern was similar to the early season.

Although strong similarities were observed in gene expression patterns between early

and late spawning seasons, certain genes exhibited common regulation in both gene

expression and DNA methylation across all three spawning seasons. Among the four DEGs

identified in the off-season, choline-phosphate cytidylyltransferase B (*pcyt1b*) and bone

morphogenetic protein 5-like (*bmp5*) showed concordant down-regulation across all three

spawning seasons. The *pcyt1b* gene encodes an enzyme involved in phosphatidylcholine

biosynthesis, playing essential roles in multiple metabolic pathways [25], while the *bmp5*

gene is involved in signalling [33]. Additionally, all DMCs found in the promoter of the

cytotoxic T lymphocyte-associated protein 2 beta (*ctl2b*) gene were hyper-methylated across

all three spawning seasons. The function of *ctl2b* remains unknown, but it shares similarities

with genes encoding cysteine proteinase, an enzyme involved in protein breakdown within

cells [61]. Furthermore, among genes that were both differentially expressed and had at least

one DMC in their promoters, the *caprin-1* gene exhibited consistent regulation, with all

DMCs being hypo-methylated and its expression being suppressed. The *caprin-1* gene is

associated with cell cycle regulation, playing a crucial role in cellular activation or

proliferation [53]. This indicates that the off-season also influenced cell cycle regulation to a

certain degree.

Can we link our findings with future growth performance?

Although methylation levels roughly correspond to body weights, indicating current growth levels, we can still estimate future growth performance from all kinds of observed data. Offspring from the off-season exhibited lower lipid levels and suppressed lipid related mechanism, which could hinder their growth potential, while offspring from the early season showed suppressed cellular processes, which are also important for growth. However, the early season had the highest methylation potential (SAM/SAH ratios), suggesting superior growth potential due to methionine's role in vital growth-related biology [10, 62]. Therefore, even though fish from the off-season and early seasons are expected to underperform those from the normal season in later development stages, fish from the early season potentially outperforms those from the off-season at some point in later stages.

Late and normal seasons exhibited similar overall DNA methylation rates and nutritional status measures. Nonetheless, the late season displayed suppressed expression of genes involved in cellular processes, including cell cycle regulation. Furthermore, the late season exhibited the lowest SAM/SAH ratio among the four seasons. This suggests that fish from the late season may underperform those from the normal season in later development stages.

Hence, the projected future growth levels can be represented as: off-season < early < late < normal, in comparison to the current growth levels: early ≈ off-season < normal ≈ late, without considering compensation from exogenous feeding. To enhance the accuracy of these estimations, compiling tissue and developmental stage-specific DNA methylation maps would be valuable, as specific DNA methylation levels at certain loci are likely associated with the underlying growth potential.

653 **Why are our findings important in the context of aquaculture and**

654 **nutritional studies?**

Our study unveiled the significant effects of altered spawning season on offspring gene

expression and DNA methylation patterns in Atlantic salmon. These findings provide

valuable insights into assessing current and future growth potential, which can be challenging

to determine using only nutritional status and growth performance measures. Understanding

the genetic and epigenetic responses to altered spawning seasons will help develop strategies

to optimize growth and production in aquaculture practices. Additionally, our study can be

linked to the potential impact of increasing global ocean temperatures on the spawning of

aquatic vertebrates, as temperature is a key abiotic factor influencing spawning seasons.

Conclusions

The present study emphasises the substantial impact of altering spawning seasons on gene

expression and DNA methylation patterns in Atlantic salmon offspring. Our observations

indicate that RAS-based off-season spawning exerts an influence on lipid-mediated

regulations in offspring, while sea-pen based early and late seasons affect the regulation of

the offspring's cellular processes, especially cell cycle regulation. These effects potentially

play a crucial role in shaping both current and future growth performance.

These findings have important implications for aquaculture and nutritional studies, as

they provide valuable tools for assessing growth potential and optimizing production

strategies. Future research focusing on tissue and developmental stage-specific DNA

methylation maps will enhance the accuracy of growth performance estimation and provide

further insights into the genetic and epigenetic responses to altered spawning conditions.

Acknowledgements

678 We thank Amelie Nemc and Bekir Ergüner for advice on RRBS analysis, and Hui-Shan Tung

679 and Eva Mykkeltvedt for DNA and RNA extraction at the Institute of Marine Research

680 (Bergen, Norway). This work was supported by the Norwegian Research Council under

681 project no: 267787 (NutrEpi) and by the Institute of Marine Research under the Nutritional

682 programming project.

Author contributions

685 K.H.S conceived and designed the research. M.E. and M.M. conducted and designed the

686 feeding trial. J.M.O.F. and C.B. prepared RNA-seq and RRBS data. T.S., M.E., and K.H.S.

687 analysed and interpreted data. T.S. drafted the manuscript, and K.H.S revised it. The authors

688 read and approved the final manuscript.

Disclosure statement

691 No potential conflict of interest was reported by the authors.

References

1.   Pankhurst NW, King HR. Temperature and salmonid reproduction: implications for aquaculture. J Fish Biol. 2010;76(1):69-85. Epub 2010/08/27. doi: 10.1111/j.1095-8649.2009.02484.x. PubMed PMID: 20738700.

2.   Taranger GL, Carrillo M, Schulz RW, Fontaine P, Zanuy S, Felip A, et al. Control of puberty in farmed fish. Gen Comp Endocrinol. 2010;165(3):483-515. Epub 2009/05/16. doi: 10.1016/j.ygcen.2009.05.004. PubMed PMID: 19442666.

3.   Good C, Davidson J. A Review of Factors Influencing Maturation of Atlantic Salmon, Salmo salar, with Focus on Water Recirculation Aquaculture System Environments. Journal of the World Aquaculture Society. 2016;47(5):605-32. doi: https://doi.org/10.1111/jwas.12342.

4.   Skjaerven KH, Oveland E, Mommens M, Samori E, Saito T, Adam AC, et al. Out-of-season spawning affects the nutritional status and gene expression in both Atlantic salmon female broodstock and their offspring. Comp Biochem Physiol A Mol Integr Physiol. 2020;247:110717. Epub 2020/05/08. doi: 10.1016/j.cbpa.2020.110717. PubMed PMID: 32376458.

5.   Skjærven KH, Mommens M, Adam A-C, Saito T, Oveland E, Espe M. Earlier or delayed seasonal broodstock spawning changes nutritional status and metabolic programming of growth for next-generation Atlantic salmon. Aquaculture. 2022;554:738187. doi: https://doi.org/10.1016/j.aquaculture.2022.738187.

6.   Lucas A. Programming by early nutrition: an experimental approach. J Nutr. 1998;128(2 Suppl):401S-6S. Epub 1998/03/21. doi: 10.1093/jn/128.2.401S. PubMed PMID: 9478036.

7.   Agosti M, Tandoi F, Morlacchi L, Bossi A. Nutritional and metabolic programming during the first thousand days of life. Pediatr Med Chir. 2017;39(2):157. Epub 2017/07/05. doi: 10.4081/pmc.2017.157. PubMed PMID: 28673078.

8.   Skjærven KH, Adam A-C, Saito T, Waagbø R, Espe M. Epigenetics in Fish Nutritional Programming. Epigenetics in Aquaculture2023. p. 231-44.

9.   Heard E, Martienssen RA. Transgenerational epigenetic inheritance: myths and mechanisms. Cell. 2014;157(1):95-109. Epub 2014/04/01. doi: 10.1016/j.cell.2014.02.045. PubMed PMID: 24679529; PubMed Central PMCID: PMCPMC4020004.

10.   Council NR. Nutrient requirements of fish and shrimp. Washington, DC: National academies press; 2011.

11.   Hamre K, Sissener NH, Lock EJ, Olsvik PA, Espe M, Torstensen BE, et al. Antioxidant nutrition in Atlantic salmon (Salmo salar) parr and post-smolt, fed diets with high inclusion of plant ingredients and graded levels of micronutrients and selected amino acids. PeerJ. 2016;4:e2688. Epub 2016/11/16. doi: 10.7717/peerj.2688. PubMed PMID: 27843721; PubMed Central PMCID: PMCPMC5103829.

12.   Hemre GI, Lock EJ, Olsvik PA, Hamre K, Espe M, Torstensen BE, et al. Atlantic salmon (Salmo salar) require increased dietary levels of B-vitamins when fed diets with high inclusion of plant based ingredients. PeerJ. 2016;4:e2493. Epub 2016/10/06. doi: 10.7717/peerj.2493. PubMed PMID: 27703849; PubMed Central PMCID: PMCPMC5047143.

13.   Danchin A, Sekowska A, You C. One-carbon metabolism, folate, zinc and translation. Microb Biotechnol. 2020;13(4):899-925. Epub 2020/03/11. doi: 10.1111/1751-7915.13550. PubMed PMID: 32153134; PubMed Central PMCID: PMCPMC7264889.

14.   Lyon P, Strippoli V, Fang B, Cimmino L. B Vitamins and One-Carbon Metabolism: Implications in Human Health and Disease. Nutrients. 2020;12(9). Epub 2020/09/24. doi:

10.3390/nu12092867. PubMed PMID: 32961717; PubMed Central PMCID:
PMCPMC7551072.
15.   Ducker GS, Rabinowitz JD. One-Carbon Metabolism in Health and Disease. Cell
Metab. 2017;25(1):27-42. Epub 2016/09/20. doi: 10.1016/j.cmet.2016.08.009. PubMed
PMID: 27641100; PubMed Central PMCID: PMCPMC5353360.
16.   Akram M. Citric acid cycle and role of its intermediates in metabolism. Cell Biochem
Biophys. 2014;68(3):475-8. Epub 2013/09/27. doi: 10.1007/s12013-013-9750-1. PubMed
PMID: 24068518.
17.   Felig P. The glucose-alanine cycle. Metabolism. 1973;22(2):179-207. Epub
1973/02/01. doi: 10.1016/0026-0495(73)90269-2. PubMed PMID: 4567003.
18.   Wu G. Functional amino acids in growth, reproduction, and health. Adv Nutr.
2010;1(1):31-7. Epub 2011/11/02. doi: 10.3945/an.110.1008. PubMed PMID: 22043449;
PubMed Central PMCID: PMCPMC3042786.
19.   Fahy E, Cotter D, Sud M, Subramaniam S. Lipid classification, structures and tools.
Biochim Biophys Acta. 2011;1811(11):637-47. Epub 2011/06/28. doi:
10.1016/j.bbalip.2011.06.009. PubMed PMID: 21704189; PubMed Central PMCID:
PMCPMC3995129.
20.   Saito T, Whatmore P, Taylor JF, Fernandes JMO, Adam AC, Tocher DR, et al.
Micronutrient supplementation affects transcriptional and epigenetic regulation of lipid
metabolism in a dose-dependent manner. Epigenetics. 2021;16(11):1217-34. Epub
2020/12/15. doi: 10.1080/15592294.2020.1859867. PubMed PMID: 33315488; PubMed
Central PMCID: PMCPMC8813078.
21.   Saito T, Espe M, Vikeså V, Bock C, Thomsen TH, Adam A-C, et al. One-carbon
metabolism nutrients impact the interplay between DNA methylation and gene expression in
liver, enhancing protein synthesis in Atlantic Salmon. bioRxiv. 2023:2023.09.18.558254. doi:
10.1101/2023.09.18.558254.
22.   Moore LD, Le T, Fan G. DNA methylation and its basic function.
Neuropsychopharmacology. 2013;38(1):23-38. Epub 2012/07/12. doi: 10.1038/npp.2012.112.
PubMed PMID: 22781841; PubMed Central PMCID: PMCPMC3521964.
23.   Amenyah SD, Hughes CF, Ward M, Rosborough S, Deane J, Thursby SJ, et al.
Influence of nutrients involved in one-carbon metabolism on DNA methylation in adults-a
systematic review and meta-analysis. Nutr Rev. 2020;78(8):647-66. Epub 2020/01/25. doi:
10.1093/nutrit/nuz094. PubMed PMID: 31977026.
24.   Goll MG, Bestor TH. Eukaryotic cytosine methyltransferases. Annu Rev Biochem.
2005;74:481-514. Epub 2005/06/15. doi: 10.1146/annurev.biochem.74.010904.153721.
PubMed PMID: 15952895.
25.   Bateman A, Martin MJ, Orchard S, Magrane M, Ahmad S, Alpi E, et al. UniProt: the
Universal Protein Knowledgebase in 2023. Nucleic Acids Res. 2022. doi:
10.1093/nar/gkac1052. PubMed PMID: WOS:000893231800001.
26.   Martin M. Cutadapt removes adapter sequences from high-throughput sequencing
reads. 2011. 2011;17(1):3. Epub 2011-08-02. doi: 10.14806/ej.17.1.200.
27.   Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, et al. STAR:
ultrafast universal RNA-seq aligner. Bioinformatics. 2013;29(1):15-21. doi:
10.1093/bioinformatics/bts635. PubMed PMID: WOS:000312654600003.
28.   Krueger F, Andrews SR. Bismark: a flexible aligner and methylation caller for
Bisulfite-Seq applications. Bioinformatics. 2011;27(11):1571-2. doi:
10.1093/bioinformatics/btr167. PubMed PMID: WOS:000291062400018.
29.   Langmead B, Trapnell C, Pop M, Salzberg SL. Ultrafast and memory-efficient
alignment of short DNA sequences to the human genome. Genome Biol. 2009;10(3). doi:
ARTN R25

10.1186/gb-2009-10-3-r25. PubMed PMID: WOS:000266544500005.

30.   Liao Y, Smyth GK, Shi W. featureCounts: an efficient general purpose program for
assigning sequence reads to genomic features. Bioinformatics. 2014;30(7):923-30. doi:
10.1093/bioinformatics/btt656. PubMed PMID: WOS:000334078300005.

31.   Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion for
RNA-seq data with DESeq2. Genome Biol. 2014;15(12). doi: ARTN 550
10.1186/s13059-014-0550-8. PubMed PMID: WOS:000346609500022.

32.   Benjamini Y, Hochberg Y. Controlling the False Discovery Rate - a Practical and
Powerful Approach to Multiple Testing. J R Stat Soc B. 1995;57(1):289-300. doi: DOI
10.1111/j.2517-6161.1995.tb02031.x. PubMed PMID: WOS:A1995QE45300017.

33.   Kanehisa M, Goto S. KEGG: Kyoto Encyclopedia of Genes and Genomes. Nucleic
Acids Res. 2000;28(1):27-30. doi: DOI 10.1093/nar/28.1.27. PubMed PMID:
WOS:000084896300007.

34.   Yu GC, Wang LG, Han YY, He QY. clusterProfiler: an R Package for Comparing
Biological Themes Among Gene Clusters. Omics. 2012;16(5):284-7. doi:
10.1089/omi.2011.0118. PubMed PMID: WOS:000303653300007.

35.   Akalin A, Kormaksson M, Li S, Garrett-Bakelman FE, Figueroa ME, Melnick A, et
al. methylKit: a comprehensive R package for the analysis of genome-wide DNA methylation
profiles. Genome Biol. 2012;13(10). doi: ARTN R87
10.1186/gb-2012-13-10-R87. PubMed PMID: WOS:000313183900006.

36.   Wang HQ, Tuominen LK, Tsai CJ. SLIM: a sliding linear model for estimating the
proportion of true null hypotheses in datasets with dependence structures. Bioinformatics.
2011;27(2):225-31. doi: 10.1093/bioinformatics/btq650. PubMed PMID:
WOS:000286215200012.

37.   Koster J, Rahmann S. Snakemake-a scalable bioinformatics workflow engine.
Bioinformatics. 2012;28(19):2520-2. doi: 10.1093/bioinformatics/bts480. PubMed PMID:
WOS:000309687500018.

38.   Kanehisa M, Goto S. KEGG: kyoto encyclopedia of genes and genomes. Nucleic
Acids Res. 2000;28(1):27-30. Epub 1999/12/11. doi: 10.1093/nar/28.1.27. PubMed PMID:
10592173; PubMed Central PMCID: PMCPMC102409.

39.   Cappel DA, Deja S, Duarte JAG, Kucejova B, Inigo M, Fletcher JA, et al. Pyruvate-
Carboxylase-Mediated Anaplerosis Promotes Antioxidant Capacity by Sustaining TCA Cycle
and Redox Metabolism in Liver. Cell Metab. 2019;29(6):1291-305 e8. Epub 2019/04/23. doi:
10.1016/j.cmet.2019.03.014. PubMed PMID: 31006591; PubMed Central PMCID:
PMCPMC6585968.

40.   Takashima M, Ogawa W, Hayashi K, Inoue H, Kinoshita S, Okamoto Y, et al. Role of
KLF15 in regulation of hepatic gluconeogenesis and metformin action. Diabetes.
2010;59(7):1608-15. Epub 2010/04/16. doi: 10.2337/db09-1679. PubMed PMID: 20393151;
PubMed Central PMCID: PMCPMC2889759.

41.   Del Giacco L, Pistocchi A, Ghilardi A. prox1b Activity is essential in zebrafish
lymphangiogenesis. PLoS One. 2010;5(10):e13170. Epub 2010/10/27. doi:
10.1371/journal.pone.0013170. PubMed PMID: 20976189; PubMed Central PMCID:
PMCPMC2956630.

42.   Scott CL, Gil J, Hernando E, Teruya-Feldstein J, Narita M, Martinez D, et al. Role of
the chromobox protein CBX7 in lymphomagenesis. Proc Natl Acad Sci U S A.
2007;104(13):5389-94. Epub 2007/03/22. doi: 10.1073/pnas.0608721104. PubMed PMID:
17374722; PubMed Central PMCID: PMCPMC1828941.

43.   Kolle G, Georgas K, Holmes GP, Little MH, Yamada T. CRIM1, a novel gene
encoding a cysteine-rich repeat protein, is developmentally regulated and implicated in

vertebrate CNS development and organogenesis. Mech Dev. 2000;90(2):181-93. Epub
2000/01/21. doi: 10.1016/s0925-4773(99)00248-8. PubMed PMID: 10642437.
44. Ku NO, Strnad P, Zhong BH, Tao GZ, Omary MB. Keratins let liver live: Mutations
predispose to liver disease and crosslinking generates Mallory-Denk bodies. Hepatology.
2007;46(5):1639-49. Epub 2007/10/31. doi: 10.1002/hep.21976. PubMed PMID: 17969036.
45. Garner K, Hunt AN, Koster G, Somerharju P, Groves E, Li M, et al.
Phosphatidylinositol transfer protein, cytoplasmic 1 (PITPNC1) binds and transfers
phosphatidic acid. J Biol Chem. 2012;287(38):32263-76. Epub 2012/07/24. doi:
10.1074/jbc.M112.375840. PubMed PMID: 22822086; PubMed Central PMCID:
PMCPMC3442557.
46. Hu JS, Olson EN, Kingston RE. HEB, a helix-loop-helix protein related to E2A and
ITF2 that can modulate the DNA-binding ability of myogenic regulatory factors. Mol Cell
Biol. 1992;12(3):1031-42. Epub 1992/03/01. doi: 10.1128/mcb.12.3.1031-1042.1992.
PubMed PMID: 1312219; PubMed Central PMCID: PMCPMC369535.
47. Zhou B, Gitschier J. hCTR1: a human gene for copper uptake identified by
complementation in yeast. Proc Natl Acad Sci U S A. 1997;94(14):7481-6. Epub 1997/07/08.
doi: 10.1073/pnas.94.14.7481. PubMed PMID: 9207117; PubMed Central PMCID:
PMCPMC23847.
48. Stolowich NJ, Petrescu AD, Huang H, Martin GG, Scott AI, Schroeder F. Sterol
carrier protein-2: structure reveals function. Cell Mol Life Sci. 2002;59(2):193-212. Epub
2002/03/28. doi: 10.1007/s00018-002-8416-8. PubMed PMID: 11915938.
49. Alliance of Genome Resources C. Harmonizing model organism data in the Alliance
of Genome Resources. Genetics. 2022;220(4). Epub 2022/04/06. doi:
10.1093/genetics/iyac022. PubMed PMID: 35380658; PubMed Central PMCID:
PMCPMC8982023.
50. Brants J, Semenchenko K, Wasylyk C, Robert A, Carles A, Zambrano A, et al.
Tubulin tyrosine ligase like 12, a TTLL family member with SET- and TTL-like domains and
roles in histone and tubulin modifications and mitosis. PLoS One. 2012;7(12):e51258. Epub
2012/12/20. doi: 10.1371/journal.pone.0051258. PubMed PMID: 23251473; PubMed Central
PMCID: PMCPMC3520985.
51. Tomaru T, Satoh T, Yoshino S, Ishizuka T, Hashimoto K, Monden T, et al. Isolation
and characterization of a transcriptional cofactor and its novel isoform that bind the
deoxyribonucleic acid-binding domain of peroxisome proliferator-activated receptor-gamma.
Endocrinology. 2006;147(1):377-88. Epub 2005/10/22. doi: 10.1210/en.2005-0450. PubMed
PMID: 16239304.
52. Norris RA, Scott KK, Moore CS, Stetten G, Brown CR, Jabs EW, et al. Human
PRRX1 and PRRX2 genes: cloning, expression, genomic localization, and exclusion as
disease genes for Nager syndrome. Mamm Genome. 2000;11(11):1000-5. Epub 2000/11/04.
doi: 10.1007/s003350010193. PubMed PMID: 11063257.
53. Grill B, Wilson GM, Zhang KX, Wang B, Doyonnas R, Quadroni M, et al.
Activation/division of lymphocytes results in increased levels of cytoplasmic
activation/proliferation-associated protein-1: prototype of a new family of proteins. J
Immunol. 2004;172(4):2389-400. Epub 2004/02/07. doi: 10.4049/jimmunol.172.4.2389.
PubMed PMID: 14764709.
54. Walczak CE, Verma S, Mitchison TJ. XCTK2: a kinesin-related protein that promotes
mitotic spindle assembly in Xenopus laevis egg extracts. J Cell Biol. 1997;136(4):859-70.
Epub 1997/02/24. doi: 10.1083/jcb.136.4.859. PubMed PMID: 9049251; PubMed Central
PMCID: PMCPMC2132492.
55. Wheatley SP, Carvalho A, Vagnarelli P, Earnshaw WC. INCENP is required for
proper targeting of Survivin to the centromeres and the anaphase spindle during mitosis. Curr

Biol. 2001;11(11):886-90. Epub 2001/08/23. doi: 10.1016/s0960-9822(01)00238-x. PubMed
PMID: 11516652.
56.   Hansson R, Wikvall K. Hydroxylations in biosynthesis of bile acids. Cytochrome P-
450 LM4 and 12alpha-hydroxylation of 5beta-cholestane-3alpha, 7alpha-diol. Eur J
Biochem. 1982;125(2):423-9. Epub 1982/07/01. doi: 10.1111/j.1432-1033.1982.tb06700.x.
PubMed PMID: 6811268.
57.   Grinberg AV, Hannemann F, Schiffler B, Muller J, Heinemann U, Bernhardt R.
Adrenodoxin: structure, stability, and electron transfer properties. Proteins. 2000;40(4):590-
612. Epub 2000/07/19. doi: 10.1002/1097-0134(20000901)40:4<590::aid-prot50>3.0.co;2-p.
PubMed PMID: 10899784.
58.   Babu E, Kanai Y, Chairoungdua A, Kim DK, Iribe Y, Tangtrongsup S, et al.
Identification of a novel system L amino acid transporter structurally distinct from
heterodimeric amino acid transporters. J Biol Chem. 2003;278(44):43838-45. Epub
2003/08/22. doi: 10.1074/jbc.M305221200. PubMed PMID: 12930836.
59.   Lanata CM, Chung SA, Criswell LA. DNA methylation 101: what is important to
know about DNA methylation and its role in SLE risk and disease heterogeneity. Lupus Sci
Med. 2018;5(1):e000285. Epub 2018/08/11. doi: 10.1136/lupus-2018-000285. PubMed
PMID: 30094041; PubMed Central PMCID: PMCPMC6069928.
60.   Westfall CS, Levin PA. Comprehensive analysis of central carbon metabolism
illuminates connections between nutrient availability, growth rate, and cell morphology in
Escherichia coli. PLoS Genet. 2018;14(2):e1007205. Epub 2018/02/13. doi:
10.1371/journal.pgen.1007205. PubMed PMID: 29432413; PubMed Central PMCID:
PMCPMC5825171.
61.   Denizot F, Brunet JF, Roustan P, Harper K, Suzan M, Luciani MF, et al. Novel
structures CTLA-2 alpha and CTLA-2 beta expressed in mouse activated T cells and mast
cells and homologous to cysteine proteinase proregions. Eur J Immunol. 1989;19(4):631-5.
Epub 1989/04/01. doi: 10.1002/eji.1830190409. PubMed PMID: 2786470.
62.   Finkelstein JD. Methionine metabolism in mammals. J Nutr Biochem. 1990;1(5):228-
37. Epub 1990/05/01. doi: 10.1016/0955-2863(90)90070-2. PubMed PMID: 15539209.
919