TITLE

TMEM16 and TMEM63/OSCA proteins share a conserved potential to permeate ions and

phospholipids

AUTHORS

Augustus J. Lowry[1,†], Pengfei Liang[1,†], Y.C. Serena Wan[1], Zhen-Ming Pei[3], Huanghe Yang[1,2,*], Yang

Zhang[1,4,†,*]

[1]Department of Biochemistry, Duke University School of Medicine, Durham, NC 27710, USA

[2]Department of Neurobiology, Duke University School of Medicine, Durham, NC 27710, USA

[3]Department of Biology, Duke University, Durham, NC 27710, USA

[4]Current address: Shenzhen Bay Laboratory, Guangdong 518106, China

[†]Denotes equal contribution

[*]To whom correspondence should be addressed

ABSTRACT

The calcium-activated TMEM16 proteins and the mechanosensitive/osmolarity-activated

OSCA/TMEM63 proteins belong to the Transmembrane Channel/Scramblase (TCS) superfamily.

Within the superfamily, OSCA/TMEM63 proteins, as well as TMEM16A and TMEM16B, likely function

solely as ion channels. However, the remaining TMEM16 members, including TMEM16F, maintain an

additional function as scramblases, rapidly exchanging phospholipids between leaflets of the

membrane. Although recent studies have advanced our understanding of TCS structure-function

relationships, the molecular determinants of TCS ion and lipid permeation remain unclear. Here we

show that single lysine mutations in transmembrane helix (TM) 4 allow non-scrambling TCS members

to permeate phospholipids. This study highlights the key role of TM 4 in controlling TCS ion and lipid

permeation and offers novel insights into the evolution of the TCS superfamily, suggesting that, like

TMEM16s, the OSCA/TMEM63 family maintains a conserved potential to permeate ions and

phospholipids.

INTRODUCTION

Ion channels and phospholipid scramblases catalyze the passive flux of ions and phospholipids down their respective chemical gradients. Compared to ion channels, both our understanding of scramblases and the evolutionary origins of phospholipid scrambling are underdeveloped. Discoveries of two scramblase families—the TMEM16 calcium-activated phospholipid scramblases (CaPLSases) and XKR caspase-dependent phospholipid scramblases—have revealed roles for scramblases in processes such as blood coagulation[1], cell death signaling[2], phagocytosis[3], cell migration[4], membrane repair[5,6], microparticle release[7], cell-cell fusion[8–10], and viral infection[11,12]. Among identified scramblases, TMEM16 CaPLSases are the most extensively studied at the molecular level[13]. The TMEM16 family (Extended Data Fig. 1a) was originally classified as calcium-activated chloride channels (CaCCs) based on the first-discovered members, TMEM16A and TMEM16B[14–16]. The remaining members, however, exhibit CaPLSase activity, with TMEM16F representing the canonical CaPLSase[17] (Fig. 1a). Uniquely, CaPLSases also function as non-selective ion channels along with their non-specific permeability to phospholipids. Substrate permeation is facilitated by calcium-induced conformational changes causing the clamshell-like separation of TMs 4 and 6 to facilitate ion and phospholipid permeation[18–20] (Extended Data Fig. 1b-e). These conformational changes further catalyzed phospholipid permeation by thinning the membrane near the permeation pathway[21,22].

The TMEM16 family together with the TMEM63/OSCA family and TMC family[23–26] collectively form the Transmembrane Channel/Scramblase (TCS) superfamily[13]. Despite the conserved 10-TM architectural core[27–31], phospholipid permeability has not been experimentally demonstrated in mechanosensitive TMEM63/OSCA or TMC proteins, raising the intriguing question of how TCS proteins discriminate between ion and ion/phospholipid substrates. Interestingly, studies have shown that the TMEM16A CaCC can be genetically modified to enable phospholipid permeability, either by substituting TMEM16F domains as small as 15 amino acids[32], or through a single lysine mutation in TM 4 at the

hydrophobic gate[18]. These findings suggest a modest energetic barrier for scramblase activity and lead

us to hypothesize that evolutionary relatives of the TMEM16 family may maintain a conserved potential

for both ion and phospholipid permeation.

To test this hypothesis, we introduced single lysine mutations along TM 4 of TMEM16F,

TMEM16A, OSCA1.2, and TMEM63A at sites near the corresponding hydrophobic gate of TMEM16F[18].

TM 4 lysine mutations in TMEM16F and TMEM16A resulted in constitutive phospholipid scramblase

activity. Strikingly, the equivalent TM 4 lysine mutations of OSCA1.2 and TMEM63A also converted

these ion channels into phospholipid scramblases, which were either constitutively open or further

activated by osmotic stimulation. All mutants also resulted in commensurate gain-of-function ion

channel activity, suggesting that the charge in TM 4 disrupts the activation gate of these evolutionarily

related proteins. Together, our findings advance the mechanistic understanding of gating and substrate

permeation in the TMEM16 and OSCA/TMEM63 families, underscore a key design principle for

phospholipid scramblases, and support our hypothesis that the TCS superfamily maintains a conserved

potential for both ion and phospholipid permeation.

RESULTS

TM 4 lysine mutations result in a constitutively active TMEM16F scramblase

Our previous study identified the activation gate of TMEM16F, consisting of hydrophobic

residues from TMs 4, 5, and 6, which collectively govern ion and phospholipid permeation[18]. Introducing

polar or charged residues to the activation gate results in gain-of-function (GOF) CaPLSases, some of

which are constitutively activated without requiring calcium. Structural and computational studies on

the fungal NhTMEM16[33] and AfTMEM16[21] orthologs, as well as human TMEM16K[34] further support a

clamshell like gating model[18] whereby scrambling activity is promoted by separation of the TM 4/6

interface[13] (Extended Data Fig. 1b-d). The conformational transition involves the N-terminal half of TM

4, which bends away from TM 6, and the C-terminal half of TM 6, which collapses onto the calcium

binding sites formed together with TMs 7 and 8[13]. Given that the N-termini of TM 4 are largely

hydrophobic among TMEM16 CaPLSases (Extended Data Fig. 1f) and a lysine mutation at F518 leads

to a constitutively open TMEM16F activation gate[18], we hypothesized that introducing a charged side

chain along the TM 4/6 interface might disrupt the hydrophobic gate and result in TMEM16 scramblases

with GOF scramblase activity. To test this hypothesis, we overexpressed eGFP-tagged mutant

constructs in TMEM16F knockout (KO) HEK293T cells and used an established scramblase

assay[18,35,36] that detects phosphatidylserine (PS) exposure using fluorophore-tagged Annexin V (AnV)

as a reporter (Fig. 1b). In the absence of calcium stimulation, PS is predominantly in the inner leaflet of

the plasma membrane of the TMEM16F wildtype (WT) expressing cells and is therefore not labelled by

extracellular AnV. Similar to TMEM16F F518K[18], overexpressing the single lysine mutations TMEM16F

I521K and M522K led to spontaneous, global exposure of PS on the plasma membrane without

requiring calcium stimulation (Fig. 1b, c). We further tested whether I521K and M522K have GOF ion

channel activity using whole-cell patch clamp. In the absence of calcium, the mutants, but not WT

TMEM16F, showed robust depolarization-activated outward rectifying currents (Fig. 1d-f). Our imaging

and patch clamp experiments thus demonstrate that I521K and M522K are functionally expressed on

the plasma membrane and their GOF lipid scramblase activity results in spontaneous, global PS

exposure on the cell surface. I521 and M522 are approximately one helical turn below the hydrophobic

gate residue, F518,[18] (Fig. 1a) supporting the idea that a lysine mutation on TM 4 along the TM 4/6

interface promotes gate opening and substrate permeation (Fig. 1g).

TM 4 lysine mutations convert TMEM16A into a constitutively active scramblase

TMEM16A is a CaCC[14–16] without scramblase activity (Fig. 2a-c)[18,32]. We previously reported

that a single TM 4 lysine mutation (L543K) at the hydrophobic gate of TMEM16A (Fig. 2a) allows the

CaCC to permeate phospholipids spontaneously, analogous to the F518K mutation in TMEM16F[18].

Thus, we reasoned that lysine mutations equivalent to TMEM16F I521K and M522K might also result

in spontaneous phospholipid permeability. To test this hypothesis, we overexpressed eGFP-tagged TMEM16A I546K and I547K mutants in TMEM16F KO HEK293T cells, and assessed their ability to expose PS on the plasma membrane with confocal microscopy. Similar to TMEM16A L543K[18], I546K- and I547K-expressing cells exhibit spontaneous, global PS exposure (Fig. 2b, c), analogous to the equivalent TMEM16F mutations I521K and M522K (Extended Data Fig. 1f; Fig. 1b, c). Similarly, whole-cell patch clamp revealed GOF ion channel activity at depolarizing potentials, even in the absence of calcium stimulation (Fig. 2d-f). Together, these results suggest that lysine mutations in this region strongly destabilize the hydrophobic gate to endow the TMEM16A CaCC with constitutive phospholipid permeability and GOF ion channel activity (Fig. 2g).

TM 4 lysine mutation L438K converts OSCA1.2 into a constitutively active scramblase

We next applied our approach to a TMEM16 relative from the OSCA/TMEM63 family (Extended Data Fig. 1a), which was first discovered in plants as a family of mechanosensitive and osmolarity-activated cation non-selective ion channels[37] (Fig. 3a). We hypothesized that analogous lysine mutations on TM 4 along the TM 4/6 interface of OSCA/TMEM63 proteins would result in GOF channels that may also permeate phospholipids. Within the family, we chose OSCA1.2 from *Arabidopsis thaliana* due to previous structural[27] and biophysical[38] characterization demonstrating that the channel is activated directly by membrane tension. Overexpressing eGFP-tagged OSCA1.2 WT in TMEM16F KO HEK293T cells did not induce PS exposure, demonstrating that OSCA1.2 WT lacks spontaneous lipid scrambling activity (Fig. 3b). We then introduced TM 4 lysine mutations near the putative gating residues identified by structural (Fig. 3a) and sequence alignment (Extended Data Fig. 1e). Strikingly, a single point mutation, L438K, causes cells overexpressing the OSCA1.2 mutant to exhibit spontaneous and global PS exposure (Fig. 3b, c). This mirrors our results with both TMEM16F (Fig. 1) and TMEM16A (Fig. 2), suggesting that the L438K mutation allows the OSCA1.2 channel to permeate phospholipids. Next, we used inside-out patch clamp to examine if L438K also enhances OSCA1.2 channel activity. We found

that the mutant significantly left-shifts the conductance-voltage (*G-V*) relationship (Fig. 3e, f) and

accelerates channel activation kinetics (Fig. 3g) compared to WT. Under -50 mmHg of pressure, L438K

has half-maximal voltages ($V_{0.5}$) of 66.7±3.7 mV, while the WT $V_{0.5}$ is nearly 108.7±5.6 mV (Fig. 3f),

underscoring that this mutation also disrupts channel gating. Together, these experiments show that,

like TMEM16A, a single lysine mutation near the putative gate allows the mechanosensitive and

osmolarity-activated OSCA1.2 channel to become spontaneously permeable to phospholipids (Fig. 3h).

TM 4 lysine mutation A439K converts OSCA1.2 into an osmolarity-sensing scramblase

Interestingly, mutating the neighboring amino acid (Fig. 3a), A439, to lysine resulted in minimal

spontaneous PS exposure (Fig. 3b, c). We thus reasoned that A439K scramblase activity may require

additional stimulation. Given that OSCA1.2 is an osmolarity-sensitive ion channel[38] and reduction of

extracellular osmolarity does not induce PS exposure in OSCA1.2 WT expressing cells (Fig. 4a, c), we

acutely treated A439K-expressing cells with a hypotonic solution (120 mOsm/kg). Indeed, PS exposure

was robustly induced (Fig. 4b, c). Inside-out patch clamp further demonstrated that A439K enhances

OSCA1.2 ion channel activity as evidenced by the accelerated activation kinetics (Fig. 3g) and left-

shifted *G-V* relationship (Fig. 3f). Our experiments thus indicate that A439K disrupts OSCA1.2 gating

and converts the osmolarity-activated ion channel into an osmolarity-activated phospholipid scramblase

(Fig. 4d).

TM 4 lysine mutations convert TMEM63A into a constitutively active scramblase

We next turned our attention to TMEM63A to further investigate the evolutionary conservation

of our observation that TM 4 lysine mutations convert TMEM16 and OSCA members into scramblases.

TMEM63s represent the mammalian members of the OSCA/TMEM63 family with three members

present in humans (TMEM63A-C). Recent structural and functional characterizations indicate that

TMEM63s function as mechanosensitive ion channels gated by high-threshold membrane tension and,

in notable contrast to all other structurally resolved TCS members, they likely function as monomers[39–]

[41]. Given their structural homology to TMEM16s and OSCA1.2[39], we hypothesized that TM 4 mutations

in TMEM63A would also result in GOF activity. We again identified residues near the putative gate by

structural (Fig. 5a) and sequence alignment (Extended Data Fig. 1f), selecting W472 (equivalent to

F518 in TMEM16F and L543 in TMEM16A) and S475 (equivalent to L438 in OSCA1.2). Indeed,

overexpressing eGFP-tagged mouse TMEM63A with single lysine mutations at either W472 and S475

led to spontaneous PS exposure (Fig. 5b, c), though the AnV staining revealed punctate rather than

global patterns of PS exposure. To confirm membrane localization and further probe mutant effects,

we exploited TMEM63A ion channel function using cell-attached patch clamp. As TMEM63A exhibits

voltage-dependent activity under high pressures, we compared mutant and WT *I-V* relationships at -80

mmHg. W472K and S475K caused marked reductions in $V_{0.5}$ from 122.3±3.5 mV for WT to 95.9±6.3

mV and 92.1±8.6 mV, respectively (Fig. 5d-f). Together, these results are consistent with our

observations in TMEM16F, TMEM16A, and OSCA1.2, indicating that single lysine mutations along TM

4 of TMEM63A can facilitate both ion and phospholipid permeation (Fig. 5g).

DISCUSSION

Mechanistically, our study improves models of TMEM16 substrate permeation and gating. We

identified multiple lysine mutations in TMEM16F and TMEM16A that promote phospholipid permeation

and cause commensurate changes in ion channel activities. Our findings complement functional

characterizations of TMEM16F gating mutants where gate destabilization is inversely correlated with

side chain hydropathy[18,42]. Lysine is above only arginine on the hydropathy index and thus likely

explains why it readily destabilizes the gate. For instance, F518K exhibits spontaneous PS exposure,

even when the calcium binding site is destroyed[18], whereas F518H does not exhibit spontaneous PS

exposure[42]. Interestingly, the recent TMEM16F F518H structure (Extended Data Fig. 1d) shows local

membrane thinning due in part to unexpected conformational changes in TM 3[42]. Future structural

studies are needed to assess whether TMEM16A or OSCA/TMEM63 mutant scramblases also promote

membrane thinning and/or conformational rearrangements in TM 3. More broadly, our results highlight

an increasingly appreciated design principle of scramblases where polar and charged residues often

line a membrane-spanning groove. This observation has been noted for TMEM16[43,34,44], Xkr[45,46], and

opsin[47] scramblases and should be a key criterion for identifying and characterizing new scramblases.

Our findings also advance our understanding of evolutionary relatives of the TCS superfamily[13]

and help uncover their roles in human diseases. Although OSCA/TMEM63 proteins are not known to

scramble phospholipids, we show that TM 4 lysine mutations in OSCA1.2 and TMEM63A convert these

osmolarity-activated and mechanosensitive ion channels into phospholipid scramblases, analogous to

our findings with the equivalent TMEM16A mutations. We thus speculate that the conserved structural

architecture in the transmembrane region endows TCS proteins with a potential to scramble

phospholipids, though this capability may have been lost during evolution. It will be interesting to test

whether equivalent TM 4 lysine mutations can convert transmembrane channel-like (TMC) proteins—

the third TCS relative of TMEM16 and OSCA/TMEM63[13]—into phospholipid scramblases. TMC1 and

TMC2 are best known for their roles in auditory sensation, and thus far have mostly been characterized

in vivo due to expression difficulties in heterologous systems[48]. However, recent *in vivo* characterization

of mouse TMC1 M412K, known as the *Beethoven* mutation, provided an important insight[49]. The

deafness-associated mutation is located in TM 4 (Extended Data Fig. 1f; equivalent of M522 in

TMEM16F, I547 in TMEM16A, and A439 in OSCA1.2) and results in constitutive PS exposure when

expressed in the hair cell membranes of both heterozygous and homozygous mice[49]. This raises the

intriguing possibility that the *Beethoven* mutation may enable TMC1 to spontaneously permeate

phospholipids, leading to a loss of membrane homeostasis and ultimately, auditory sensation[50]. The

possibility of converting TMC proteins into phospholipid scramblases should be thoroughly investigated.

Additionally, disease-associated mutations in TMEM63 proteins are present along the TM 4/6 interface,

such as TMEM63B T481N[49]. We speculate that introducing more hydropathic side chains along this

interface may lead to spontaneous ion and or phospholipid permeability, perhaps contributing to

underlying pathophysiology.

METHODS

Cloning and mutagenesis

All constructs use a peGFP-N1 vector backbone. Wild type sequence and mutation numbers

correspond to NCBI: NP_780553.2 (*Mus musculus* TMEM16F) with a three amino acid (MQM) N-

terminal truncation, NCBI: NP_001229278 (*Mus musculus* TMEM16A), GenBank: AIU34614.1

(*Arabidopsis thaliana* OSCA1.2), and NCBI: NP_001404481.1 (*Mus musculus* TMEM63A). *At*OSCA1.2

and *Mm*TMEM63A cDNAs were subcloned using In-Fusion Snap Assembly (Takara, #638947). Point

mutants were generated by PCR site-directed mutagenesis with primers from IDT DNA Technologies.

Sequences were confirmed by Sanger sequencing (Azenta).

Bioinformatics

The following sequences were obtained from UniProt and aligned using Clustal Omega: Q8BHY3-2

(*Mm*TMEM16A), Q9NQ90 (*Hs*TMEM16B), Q9BYT9 (*Hs*TMEM16C), Q32M45 (*Hs*TMEM16D), Q75V66

(*Hs*TMEM16E), Q6P9J9 (*Mm*TMEM16F), Q6IWH7 (*Hs*TMEM16G), Q9HCE9 (*Hs*TMEM16H), A1A5B4

(*Hs*TMEM16J), Q9NW15 (*Hs*TMEM16K), C7Z7K1 (*Nh*TMEM16), Q4WA18 (*Af*TMEM16), Q9XEA1

(*At*OSCA1.1), Q5XEZ5 (*At*OSC1.2), B5TYT3 (*At*OSCA1.3), A0A097NUQ0 (*At*OSCA1.4),

A0A097NUS0 (*At*OSCA1.5), A0A097NUP1 (*At*OSCA1.6), A0A097NUP8 (*At*OSCA1.7), A0A097NUQ2

(*At*OSCA1.8), A0A097NUQ5 (*At*OSCA2.1), A0A097NUS5 (*At*OSCA2.2), A0A097NUP6 (*At*OSCA2.3),

A0A097NUQ3 (*At*OSCA2.4), A0A097NUQ7 (*At*OSCA2.5), Q9C8G5 (*At*OSCA3.1), A0A097NUT0

(*At*OSCA4.1), Q91YT8 (*Mm*TMEM63A), Q5T3F8 (*Hs*TMEM63B), and Q9P1W3 (*Hs*TMEM63C),

Q8R4P5 (*Mm*TMC1), Q8R4P4 (*Mm*TMC2), Q7TQ69 (*Mm*TMC3), Q7TQ65 (*Mm*TMC4), Q32NZ6

(*Mm*TMC5), Q7TN60 (*Mm*TMC6), Q8C428 (*Mm*TMC7), and Q7TN58 (*Mm*TMC8). A subset of the

alignment was selected for Extended Data Fig. 1. Structural models were obtained from the PDB,

aligned, and visualized using Pymol (Schrödinger).

Cell culture

The HEK293T cell line was authenticated by the Duke Cell Culture Facility. The TMEM16F KO

HEK293T cell line was generated by the Duke Functional Genomics Core and characterized in previous

studies[18,35]. All cells were cultured with DMEM (Gibco, #11995-065) supplemented with 10% fetal

bovine serum (FBS) (Sigma-Aldrich, # F2442) and 1% penicillin/streptomycin (Gibco, #15-140-122) at

37°C in 5% CO_2-95% air.

Transfection

Plasmids were transiently transfected into TMEM16F KO HEK 293T cells by using X-tremeGENE9 or

X-tremeGENE360 transfection reagents (MilliporeSigma). Media was replaced with calcium free DMEM

(Gibco, 21068-028) 3-4 hours after transfection. The transfected cells were imaged or patched 18-24

or 18-48 hours after transfection, respectively.

Fluorescence imaging of scramblase-mediated PS Exposure

A Zeiss 780 inverted confocal microscope was used to monitor scramblase activity in live cells using

the methods described in previous publications[9,18,35]. Briefly, 18-24 hours after transfection, the cells

were incubated in AnV buffer (1:175 dilution of the fluorescently tagged AnV (Biotium) in Hank's

balanced salt solution) before imaging. Spontaneous PS positive cells were labeled by fluorescently

tagged AnV and imaged. Results were quantified as a percentage of PS positive cells among all GFP

positive cells. For osmolarity activation, 2 mM $CaCl_2$ in ddH_2O was added to the AnV buffer at a 2:1

ratio. The final osmolarity was ~120 mOsm/kg as measured by a micro-osmometer (Advanced

Instrument). Cells overexpressing WT or mutant OSCA1.2 were treated with low osmolarity AnV buffer,

and the scramblase activity was measured by recording fluorescent AnV surface accumulation at 5

second intervals. A custom MATLAB code was used to quantify AnV signal and is available at Github (https://github.com/yanghuanghe/scrambling_activity).

Electrophysiology

All electrophysiology recordings were conducted using an Axopatch 200B amplifier with the signal digitally sampled at 10 kHz using an Axon Digidata 1550A (Molecular Devices, Inc.). All electrophysiology recordings were carried out at room temperature 18-48 hours after transfection. Glass pipettes were pulled from borosilicate capillaries (Sutter Instruments) and fire-polished using a microforge (Narishige). Pipettes had resistances of 2–3 MΩ in bath solution.

Inside-out patch clamp recordings. The pipette solution (external) contained 140 mM NaCl, 10 mM HEPES, 2 mM $MgCl_2$, adjusted to pH 7.3 (NaOH), and the bath solution (internal) contained 140 mM NaCl, 10 mM HEPES, 5 mM EGTA, adjusted to pH 7.3 (NaOH). OSCA1.2 WT and mutations were held at constant pressure administrated using a syringe calibrated with a manometer, similar to a previous study[54]. Patches were held at a membrane potential of -60 mV and at the indicated pressure, then stimulated using the indicated voltage protocol.

Cell-attached patch clamp recordings. For TMEM63 recordings, cell-attach mode was used. The bath solution contained (in mM): 140 KCl, 10 HEPES, 2 $MgCl_2$, 10 glucose, pH 7.3 adjusted with KOH. The pipette solution contained (in mM): 130 NaCl, 5 KCl, 10 HEPES, 10 TEA-Cl, 1 $CaCl_2$, 1 $MgCl_2$, pH 7.3 (with NaOH). The mechano-activated current was evoked with a 200 ms pressure pulse at -80 mmHg using a high-speed pressure clamp system (HSPC-1, ALA Scientific Instruments, Farmingdale, NY). The membrane potential inside the patch was held at -60 mV. The voltage pulse alone was run first followed by voltage pulse with pressure. The mechanosensitive current was obtained by subtracting the voltage pulse from the voltage with pressure pulse.

Whole cell patch clamp recordings. The pipette solution (internal) contained 140 mM CsCl, 10 mM HEPES, 5 mM EGTA, adjusted to pH 7.3 (CsOH), and the bath solution (external) contained

140 mM NaCl, 10 mM HEPES, 5 mM EGTA, adjusted to pH 7.3 (NaOH). Patches were held at a

membrane potential of -60 mV, then stimulated using the indicated voltage protocol.

Data analysis for electrophysiology

All data analysis was performed using Clampfit (Molecular Devices), Excel (Microsoft), MATLAB

(MathWorks), and Prism softwares (GraphPad). Individual *G-V* curves were fitted with a Boltzmann

function,

$$G = \frac{G_{max}}{1+e^{\frac{-ZF(V-V_{0.5})}{RT}}} \qquad (2)$$

where G_{max} denotes the fitted value for maximal conductance, $V_{0.5}$ denotes the voltage of half maximal

activation of conductance, Z denotes the net charge moved across the membrane during the transition

from the closed to the open state, and F denotes the Faraday constant.

REFERENCES

1. Suzuki, J., Umeda, M., Sims, P. J. & Nagata, S. Calcium-dependent phospholipid scrambling by TMEM16F. *Nature* **468**, 834–838 (2010).

2. Suzuki, J., Denning, D. P., Imanishi, E., Horvitz, H. R. & Nagata, S. Xk-related protein 8 and CED-8 promote phosphatidylserine exposure in apoptotic cells. *Science* **341**, 403–406 (2013).

3. Fadok, V. A. *et al.* Exposure of phosphatidylserine on the surface of apoptotic lymphocytes triggers specific recognition and removal by macrophages. *J. Immunol.* **148**, 2207–2216 (1992).

4. Chang, Z. *et al.* Anoctamin5 regulates cell migration and invasion in thyroid cancer. *Int. J. Oncol.* **51**, 1311–1319 (2017).

5. Griffin, D. A. *et al.* Defective membrane fusion and repair in Anoctamin5-deficient muscular dystrophy. *Hum. Mol. Genet.* **25**, 1900–1911 (2016).

6. Chandra, G. *et al.* Dysregulated calcium homeostasis prevents plasma membrane repair in Anoctamin 5/TMEM16E-deficient patient muscle cells. *Cell Death Discov.* **5**, 118 (2019).

7. Fujii, T., Sakata, A., Nishimura, S., Eto, K. & Nagata, S. TMEM16F is required for phosphatidylserine

exposure and microparticle release in activated mouse platelets. *Proc. Natl. Acad. Sci. U. S. A.* **112**,

12800–12805 (2015).

8. Whitlock, J. M., Yu, K., Cui, Y. Y. & Hartzell, H. C. Anoctamin 5/TMEM16E facilitates muscle

precursor cell fusion. *J. Gen. Physiol.* **150**, 1498–1509 (2018).

9. Zhang, Y. *et al.* TMEM16F phospholipid scramblase mediates trophoblast fusion and placental

development. *Sci. Adv.* **6**, eaba0310 (2020).

10.   Zhang, Y. *et al.* Functional coupling between TRPV4 channel and TMEM16F modulates human

trophoblast fusion. *eLife* **11**, e78840 (2022).

11.   Zaitseva, E. *et al.* Fusion Stage of HIV-1 Entry Depends on Virus-Induced Cell Surface Exposure

of Phosphatidylserine. *Cell Host Microbe* **22**, 99-110.e7 (2017).

12.   Nanbo, A. *et al.* Ebola virus requires a host scramblase for externalization of phosphatidylserine

on the surface of viral particles. *PLoS Pathog.* **14**, e1006848 (2018).

13.   Le, S. C., Liang, P., Lowry, A. J. & Yang, H. Gating and Regulatory Mechanisms of TMEM16 Ion

Channels and Scramblases. *Front. Physiol.* **12**, 787773 (2021).

14.   Caputo, A. *et al.* TMEM16A, a membrane protein associated with calcium-dependent chloride

channel activity. *Science* **322**, 590–594 (2008).

15.   Yang, Y. D. *et al.* TMEM16A confers receptor-activated calcium-dependent chloride

conductance. *Nature* **455**, 1210–1215 (2008).

16.   Schroeder, B. C., Cheng, T., Jan, Y. N. & Jan, L. Y. Expression cloning of TMEM16A as a

calcium-activated chloride channel subunit. *Cell* **134**, 1019–1029 (2008).

17.   Suzuki, J. *et al.* Calcium-dependent phospholipid scramblase activity of TMEM16 protein family

members. *J. Biol. Chem.* **288**, 13305–13316 (2013).

18.   Le, T. *et al.* An inner activation gate controls TMEM16F phospholipid scrambling. *Nat. Commun.*

10, 1846 (2019).

19.   Lam, A. K. M., Rheinberger, J., Paulino, C. & Dutzler, R. Gating the pore of the calcium-activated

chloride channel TMEM16A. *Nat. Commun.* **12**, 785 (2021).

20.   Lam, A. K. M. & Dutzler, R. Mechanism of pore opening in the calcium-activated chloride channel

TMEM16A. *Nat. Commun.* **12**, 786 (2021).

21.   Falzone, M. E. *et al.* Structural basis of Ca2+-dependent activation and lipid transport by a

TMEM16 scramblase. *eLife* **8**, e43229 (2019).

22.   Falzone, M. E. *et al.* TMEM16 scramblases thin the membrane to enable lipid scrambling. *Nat.*

Commun. **13**, 2604 (2022).

23.   Keresztes, G., Mutai, H. & Heller, S. TMC and EVER genes belong to a larger novel family, the

TMC gene family encoding transmembrane proteins. *BMC Genomics* **4**, 24 (2003).

24.   Hahn, Y., Kim, D. S., Pastan, I. H. & Lee, B. Anoctamin and transmembrane channel-like

proteins are evolutionarily related. *Int. J. Mol. Med.* **24**, 51–55 (2009).

25.   Medrano-Soto, A. *et al.* Bioinformatic characterization of the Anoctamin Superfamily of Ca2+-

activated ion channels and lipid scramblases. *PLOS ONE* **13**, e0192851 (2018).

26.   Ballesteros, A., Fenollar-Ferrer, C. & Swartz, K. J. Structural relationship between the putative

hair cell mechanotransduction channel TMC1 and TMEM16 proteins. *eLife* **7**, e38433 (2018).

27.   Jojoa-Cruz, S. *et al.* Cryo-EM structure of the mechanically activated ion channel OSCA1.2.

eLife **7**, e41845 (2018).

28.   Liu, X., Wang, J. & Sun, L. Structure of the hyperosmolality-gated calcium-permeable channel

OSCA1.2. *Nat. Commun.* **9**, 5060 (2018).

29.   Zhang, M. *et al.* Structure of the mechanosensitive OSCA channels. *Nat. Struct. Mol. Biol.* **25**,

850–858 (2018).

30.	Maity, K. *et al.* Cryo-EM structure of OSCA1.2 from Oryza sativa elucidates the mechanical basis of potential membrane hyperosmolality gating. *Proc. Natl. Acad. Sci. U. S. A.* **116**, 14309–14318 (2019).

31.	Jeong, H. *et al.* Structures of the TMC-1 complex illuminate mechanosensory transduction. *Nature* **610**, 796–803 (2022).

32.	Yu, K. *et al.* Identification of a lipid scrambling domain in ANO6/TMEM16F. *eLife* **4**, e06901 (2015).

33.	Kalienkova, V. *et al.* Stepwise activation mechanism of the scramblase nhTMEM16 revealed by cryo-EM. *eLife* **8**, e44364 (2019).

34.	Bushell, S. R. *et al.* The structural basis of lipid scrambling and inactivation in the endoplasmic reticulum scramblase TMEM16K. *Nat. Commun.* **10**, 3956 (2019).

35.	Le, T., Le, S. C. & Yang, H. Drosophila Subdued is a moonlighting transmembrane protein 16 (TMEM16) that transports ions and phospholipids. *J. Biol. Chem.* **294**, 4529–4537 (2019).

36.	Le, T., Le, S. C., Zhang, Y., Liang, P. & Yang, H. Evidence that polyphenols do not inhibit the phospholipid scramblase TMEM16F. *J. Biol. Chem.* **295**, 12537–12544 (2020).

37.	Yuan, F. *et al.* OSCA1 mediates osmotic-stress-evoked Ca2+ increases vital for osmosensing in Arabidopsis. *Nature* **514**, 367–371 (2014).

38.	Murthy, S. E. *et al.* OSCA/TMEM63 are an Evolutionarily Conserved Family of Mechanically Activated Ion Channels. *eLife* **7**, e41844 (2018).

39.	Zheng, W. *et al.* TMEM63 proteins function as monomeric high-threshold mechanosensitive ion channels. *Neuron* (2023) doi:10.1016/j.neuron.2023.07.006.

40.	Qin, Y. *et al.* Cryo-EM structure of TMEM63C suggests it functions as a monomer. *Nat. Commun.* **14**, 7265 (2023).

358   41.   Wu, X., Shang, T., Lü, X., Luo, D. & Yang, D. A monomeric structure of human TMEM63A

359   protein. *Proteins* (2024) doi:10.1002/prot.26660.

360   42.   Arndt, M. *et al.* Structural basis for the activation of the lipid scramblase TMEM16F. *Nat.*

361   *Commun.* **13**, 6692 (2022).

362   43.   Brunner, J. D., Lim, N. K., Schenck, S., Duerst, A. & Dutzler, R. X-ray structure of a calcium-

363   activated TMEM16 lipid scramblase. *Nature* **516**, 207–212 (2014).

364   44.   Alvadia, C. *et al.* Cryo-EM structures and functional characterization of the murine lipid

365   scramblase TMEM16F. *eLife* **8**, e44365 (2019).

366   45.   Straub, M. S., Alvadia, C., Sawicka, M. & Dutzler, R. Cryo-EM structures of the caspase-

367   activated protein XKR9 involved in apoptotic lipid scrambling. *eLife* **10**, e69800 (2021).

368   46.   Sakuragi, T. *et al.* The tertiary structure of the human Xkr8-Basigin complex that scrambles

369   phospholipids at plasma membranes. *Nat. Struct. Mol. Biol.* **28**, 825–834 (2021).

370   47.   Morra, G. *et al.* Mechanisms of Lipid Scrambling by the G Protein-Coupled Receptor Opsin.

371   *Struct. Lond. Engl. 1993* **26**, 356-367.e3 (2018).

372   48.   Pan, B. *et al.* TMC1 Forms the Pore of Mechanosensory Transduction Channels in Vertebrate

373   Inner Ear Hair Cells. *Neuron* **99**, 736-753.e6 (2018).

374   49.   Ballesteros, A. & Swartz, K. J. Regulation of membrane homeostasis by TMC1

375   mechanoelectrical transduction channels is essential for hearing. *Sci. Adv.* **8**, eabm5550 (2022).

376   50.   Sakuragi, T. & Nagata, S. Regulation of phospholipid distribution in the lipid bilayer by flippases

377   and scramblases. *Nat. Rev. Mol. Cell Biol.* 1–21 (2023) doi:10.1038/s41580-023-00604-z.

378   51.   Kumar, S., Stecher, G., Li, M., Knyaz, C. & Tamura, K. MEGA X: Molecular Evolutionary

379   Genetics Analysis across Computing Platforms. *Mol. Biol. Evol.* **35**, 1547–1549 (2018).

380   52.   Le, S. C., Jia, Z., Chen, J. & Yang, H. Molecular basis of PIP2-dependent regulation of the Ca2+-

381   activated chloride channel TMEM16A. *Nat. Commun.* **10**, 3769 (2019).

53.   Liang, P. & Yang, H. Molecular underpinning of intracellular pH regulation on TMEM16F. *J. Gen.*

Physiol. **153**, e202012704 (2021).

54.   Tsuchiya, M. *et al.* Cell surface flip-flop of phosphatidylserine is critical for PIEZO1-mediated

myotube formation. *Nat. Commun.* **9**, 2049 (2018).

ACKNOWLEDGEMENTS

This work was supported by the National Institute of Health (DP2GM126898 and R21GM146152 to

H.Y.) and the National Science Foundation Graduate Research Fellowship Program (DGE 2139754 to

A.J.L.). Any opinions, findings, and conclusions or recommendations expressed in this material are

those of the author(s) and do not necessarily reflect the views of the National Science Foundation.

AUTHOR CONTRIBUTIONS

H.Y. and Y.Z. conceived and H.Y. supervised the project. Y.Z. and A.J.L. imaging. P.L.

electrophysiology. A.J.L. sequence and structure alignments. A.J.L. and Y.C.S.W. cloning and

mutagenesis. Z.P. plasmids. Y.Z. MATLAB codes. A.J.L. and H.Y. drafted the manuscript with input

from all authors.

COMPETING INTERESTS

The authors declare no competing interests.

FIGURES

Figure 1: Lysine mutations along TM 4 enable TMEM16F channel and scramblase activities in the absence of calcium stimulation. (**a**) Top: TMEM16F is a calcium-activated phospholipid scramblase. Bottom: TM 4 mutant locations mapped on TMEM16F CaPLSase structure (PDB 6QPB). (**b**) Representative images of TMEM16F knockout (KO) HEK293T cells expressing eGFP-tagged

TMEM16F wildtype (WT), I521K, and M522K (center row). CF 594-conjugated Annexin V (AnV, bottom row) labelled PS exposing cells. BF denotes bright field images (top row). (**c**) Quantification of the percentage of cells with AnV labelling for TMEM16F WT (n=5), I521K (n=5), and M522K (n=5) transfected cells. Values were derived from images of biological replicates, with error bars representing standard error of the mean (SEM). Statistical comparisons were done using unpaired t-tests with Welch's correction (****: $p < 0.0001$). (**d**) Representative current recordings and (**e**) current-voltage (I-V) relationships of whole cell patches from TMEM16F KO HEK293T cells expressing eGFP-tagged TMEM16F WT (n=6), I521K (n=5), and M522K (n=5). Currents were elicited by the voltage protocol shown with the pipette solution containing 5 mM EGTA. Dotted line denotes zero current. (**f**) Quantification of current at +160 mV. Currents in (**e**) and (**f**) were normalized to cell capacitance with the mean ± SEM calculated from independent patches. Statistical comparisons were done using unpaired t-tests with Welch's correction (*: $p < 0.05$). (**g**) Lysine mutations along TM 4 in TMEM16F enable spontaneous phospholipid permeation.

Figure 2: Lysine mutations along TM 4 enable TMEM16A channel and scramblase activities in the absence of calcium stimulation. (**a**) Top: TMEM16A is a calcium-activated chloride channel Bottom: TM 4 mutant locations mapped on TMEM16A CaCC structure (PDB 5OYG). (**b**) Representative images of TMEM16F knockout (KO) HEK293T cells expressing eGFP-tagged TMEM16A wildtype (WT), I546K, and I547K (center rows). CF 594-conjugated annexin V (AnV, bottom row) labelled PS

exposing cells. BF denotes bright field images (top row). (**c**) Quantification of the percentage of cells

with AnV labelling for TMEM16A WT (n=4), I546K (n=4), and I547K (n=4) transfected cells. Values were

derived from images of biological replicates, with error bars representing the standard error of the mean

(SEM). Statistical comparisons were done using unpaired t-tests with Welch's correction (*: p<0.05, **:

p<0.01). (**d**) Representative whole-cell current recordings and (**e**) current-voltage (I-V) relationships of

whole cell patches from TMEM16F KO HEK293T cells expressing eGFP-tagged TMEM16A WT (n=14),

I546K (n=5), and I547K (n=5). Currents were elicited by the voltage protocol shown with the pipette

containing an EGTA solution. Dotted line denotes zero current. (**f**) Quantification of current at +160 mV.

Currents in (**e**) and (**f**) were normalized to cell capacitance with the mean ± SEM calculated from

independent patches. Statistical comparisons were done using unpaired t-tests with Welch's correction

(*: p<0.05, ****: p<0.0001). (**g**) Lysine mutations along TM 4 in TMEM16A enable spontaneous

phospholipid permeation.

Figure 3: Lysine mutations along TM 4 enable OSCA1.2 channel and scramblase activities. Top:

OSCA1.2 is a cation non-selective ion channel gated by membrane tension. Bottom: the TM 4/6

interface of OSCA1.2 (PDB 6MGV) with key residues shown as yellow sticks. (**b**) Representative

images of TMEM16F KO HEK293T cells expressing eGFP-tagged (middle row) OSCA1.2 WT (left

column), L438K (middle column), or A439K mutants (right column). CF 594-conjugated AnV (bottom

row) labelled PS exposing cells. BF denotes bright field images (top row). Asterisk highlights a PS

positive cell for the A439K mutant. (**c**) Quantification of the percentage of cells with AnV labelling for

OSCA1.2 WT (n = 4), L438K (n = 7), and A439K-transfected cells (n = 6). Statistical comparisons were

conducted with an unpaired t-test with Welch's correction (*: p<0.05, ****: p<0.0001). (**d**) Representative

current recordings and (**e**) normalized conductance-voltage (*G-V*) relationships of inside-out patches

from TMEM16F KO HEK293T cells expressing eGFP-tagged OSCA1.2 WT (n=8), L438K (n=8), and

A439K (n = 6). Currents were elicited by the voltage protocol shown next to the listed pressures. Dotted

line denotes zero current. (**f**) Quantification of half-maximal voltage at -50 mmHg for WT (109 mV),

L438K (67 mV), and A439K (63 mV). Error bars represent standard error of the mean (SEM) calculated

from independent patches. Statistical comparison was conducted with an unpaired t-tests with Welch's

correction (***: p<0.001, ****: p<0.0001). (**g**) Quantification of activation τ_{on} at -50 mmHg and 160 mV

for WT (41 ms), L438K (13 ms), and A439K (16 ms). Error bars represent standard error of the mean

(SEM) calculated from independent patches. Statistical comparison was conducted with an unpaired t-

tests with Welch's correction (***: p<0.001, ****: p<0.0001). (**h**) A lysine mutation along TM 4 converts

OSCA1.2 channel into a phospholipid scramblase with spontaneous phospholipid permeability.

Figure 4: OSCA1.2 A439K is an osmolarity-activated scramblase.

(**a-b**) Representative images of hypotonic osmolarity stimulation of TMEM16F KO HEK293T cells expressing eGFP-tagged OSCA1.2 (**a**) WT or (**b**) the A439K mutant (center rows). CF 594-conjugated AnV (bottom rows) labelled PS exposing cells. BF denotes bright field images (top rows). Each column of representative images corresponds to the indicated time after hypo-osmotic stimulation. (**c**) Quantification of AnV intensity for OSCA1.2 WT (n=5) and A439K (n=5) after hypo-osmotic stimulation. Statistical comparison was conducted with an unpaired t-tests with Welch's correction (**: p<0.01). (**d**) The A439K mutation converts OSCA1.2 to an osmolarity-activated phospholipid scramblase.

Figure 5: Lysine mutations along TM 4 enable TMEM63A channel and scramblase activities. Top:
TMEM63A is a cation non-selective ion channel gated by membrane tension. Bottom: the TM 4/6 interface of
*Hs*TMEM63A (PDB 8GRS) with key residues shown as yellow sticks using amino acid numbering corresponding
to the mouse ortholog. (**b**) Representative images of TMEM16F KO HEK293T cells expressing eGFP-tagged
(middle row) TMEM63A WT (left column), W472K (middle column), or S475K mutants (right column). CF 594-

conjugated AnV (bottom row) labelled PS exposing cells. BF denotes bright field images (top row). (**c**)

Quantification of the percentage of cells with AnV labelling for TMEM63A WT (n = 4), W472K (n = 4), and S475K-

transfected cells (n = 4). Statistical comparisons were conducted with unpaired t-tests with Welch's correction

(**: p<0.01, ****: p<0.0001). (**d**) Representative current recordings and (**e**) normalized conductance-voltage (*I-*

V) relationships of cell attached patches from TMEM16F KO HEK293T cells expressing eGFP-tagged TMEM63A

WT (n = 7), W472K (n = 7), and S475K (n = 5). Currents represent the subtraction of voltage alone from currents

elicited by the voltage and pressure protocols shown. Dotted line denotes zero current. (**f**) Quantification of half-

maximal voltage at -80 mmHg for WT (122 mV), W472K (96 mV), and S475K (92 mV). Error bars represent

standard error of the mean (SEM) calculated from independent patches. Statistical comparison was conducted

with an unpaired t-tests with Welch's correction (*: p<0.05, **: p<0.01). (**g**) Lysine mutations along TM 4 in

TMEM63A enable spontaneous phospholipid permeability.