Explicit and implicit locomotor learning in individuals with chronic hemiparetic stroke

Jonathan M. Wood[1,2], Elizabeth Thompson[1], Henry Wright[1], Liam Festa[1], Susanne M.

Morton[1,2], Darcy S. Reisman[1,2, †], and Hyosub E. Kim[1-4, †]

[†]These authors contributed equally to this work.

1. Department of Physical Therapy, University of Delaware, Newark, DE 19713, United

States

2. Biomechanics and Movement Sciences Program, University of Delaware, Newark, DE

19713, United States

3. School of Kinesiology, University of British Columbia, Vancouver, BC, Canada

4. Graduate Program in Neuroscience, University of British Columbia, Vancouver, BC,

Canada

Correspondence to:

Darcy S. Reisman

Address: 540 South College Ave, University of Delaware, Newark, DE 19713

Phone: 302-831-0508

Email: dreisman@udel.edu

Abstract

Motor learning involves both explicit and implicit processes that are fundamental for

acquiring and adapting complex motor skills. However, stroke may damage the neural

substrates underlying explicit and/or implicit learning, leading to deficits in overall motor

performance. While both learning processes are typically used in concert in daily life and

rehabilitation, no gait studies have determined how these processes function together after

stroke when tested during a task that elicits dissociable contributions from both. Here, we

compared explicit and implicit locomotor learning in individuals with chronic stroke to age-

and sex-matched neurologically intact controls. We assessed implicit learning using split-belt

adaptation (where two treadmill belts move at different speeds). We assessed explicit

learning (i.e., strategy-use) using visual feedback during split-belt walking to help individuals

explicitly correct for step length errors created by the split-belts. The removal of visual

feedback after the first 40 strides of split-belt walking, combined with task instructions,

minimized contributions from explicit learning for the remainder of the task. We utilized a

multi-rate state-space model to characterize individual explicit and implicit process

contributions to overall behavioral change. The computational and behavioral analyses

revealed that, compared to controls, individuals with chronic stroke demonstrated deficits in

both explicit and implicit contributions to locomotor learning, a result that runs counter to

prior work testing each process individually during gait. Since post-stroke locomotor

rehabilitation involves interventions that rely on both explicit and implicit motor learning,

future work should determine how locomotor rehabilitation interventions can be structured to

optimize overall motor learning.

New and noteworthy

Motor learning involves both implicit and explicit processes, the underlying neural substrates of which could be damaged by after stroke. While both learning processes are typically used in concert in daily life and rehabilitation, no gait studies have determined how these processes function together after stroke. Using a locomotor task that elicits dissociable contributions from both processes and computational modeling, we found evidence that chronic stroke causes deficits in both explicit and implicit locomotor learning.

Introduction

Introduction

Motor learning, the ability to acquire and maintain motor skills with practice (Schmidt

and Lee 2005), involves both explicit and implicit processes. Explicit learning, as in strategy

formation and use, is critical for skill acquisition because it provides the means for fast,

flexible changes in movements (Bond and Taylor 2015; Huberdeau et al. 2019; Roemmich

and Bastian 2018). Implicit processes keep movements finely calibrated in the face of

changes to the body or environment (Bond and Taylor 2015; Krakauer et al. 2019; Mazzoni

and Krakauer 2006; Roemmich and Bastian 2018). While both explicit and implicit processes

typically work in concert when learning complex motor skills, stroke may damage the neural

substrates underlying either component (or both), leading to deficits in overall motor

learning. Since motor learning is foundational to rehabilitation of individuals post-stroke, and

since explicit and implicit learning are often used concurrently (Roemmich et al. 2016;

Taylor and Ivry 2011, 2012), it is important to determine if individuals post-stroke have

impairments in one or both processes when assessed during the same motor task.

Explicit learning, sometimes referred to as "voluntary correction" (Roemmich et al.

2016) or "cognitive strategies" (Taylor and Ivry 2011), involves consciously directing

specific changes in movement patterns. For example, a patient may consciously increase their

step length in response to a clinician's verbal instructions. Explicit learning plays a critical

role in skill acquisition and motor memory for neurologically intact individuals, and is

hypothesized to be mediated primarily in prefrontal cortex (Krakauer et al. 2019; McDougle

et al. 2016; Taylor and Ivry 2014). We assume this process is driven by target error, the

difference between a movement outcome and the task goal (French et al. 2018; Mazzoni and

Krakauer 2006; Taylor and Ivry 2012). A key feature of this process is that it can be

volitionally "switched" on or off in response to context or instructions (Bond and Taylor

2015; French et al. 2018; Wood et al. 2020, 2021). Explicit learning can be used during gait

in both neurologically intact individuals and those post-stroke by providing visual feedback

and specific task instructions (French et al. 2021b, 2021a; Hussain et al. 2013). While our

prior work testing explicit learning post-stroke did not observe group-level impairments

(French et al. 2021b), we did find an effect of fluid cognition on explicit learning in a

separate study using the same paradigm (French et al. 2021a). This, in combination with the

use of a relatively small, visual perturbation in these studies suggests the potential to observe

explicit learning deficits under more challenging task conditions.

Sensorimotor adaptation is an implicit motor learning process that is essential for

maintaining well-calibrated movements in response to ever-changing environments and body

states. Sensorimotor adaptation, which we refer to here as "implicit adaptation", is driven by

sensory prediction error, the difference between the actual and expected sensory

consequences of a motor command, and mediated in large part by the cerebellum (Krakauer

et al. 2000; Morton and Bastian 2006; Shadmehr et al. 2010; Shadmehr and Mussa-Ivaldi

1994; Tseng et al. 2007). During gait, implicit adaptation can be elicited using a split-belt

treadmill where the belts under each limb move at different speeds (Dietz et al. 1994). This

perturbation initially produces asymmetric gait patterns (e.g., step length asymmetries) which

are slowly recalibrated back to baseline asymmetry levels (Malone et al. 2012; Reisman et al.

2005). The hallmark of implicit adaptation is the storage of the adapted stepping pattern when

the belts return to the same speed, termed an "implicit aftereffect" (Reisman et al. 2005).

When tested in isolation during locomotion, individuals with non-cerebellar stroke adapt their

step length asymmetry to a similar magnitude as neurologically intact participants by the end

of learning (Malone and Bastian 2014; Reisman et al. 2007; Savin et al. 2013; Tyrell et al.

2014) and demonstrate similar implicit aftereffects (Reisman et al. 2007; Savin et al. 2013;

c.f. Malone and Bastian 2014). However, individuals with stroke adapted at a slower rate

compared to controls (Malone and Bastian 2014; Reisman et al. 2007; Savin et al. 2013;

Tyrell et al. 2014).

While explicit learning and implicit adaptation are typically used simultaneously to

learn new skills in everyday life, including rehabilitation practice, they are mostly studied

individually during gait (French et al. 2018; Hussain et al. 2013; Reisman et al. 2005; Wood

et al. 2020, 2021). This may be because when they are studied within the same task, it can be

difficult to dissociate the individual contributions of each process to overall behavior

(Cherry-Allen et al. 2018; Long et al. 2016; Malone and Bastian 2010; Roemmich et al.

2016). A study in young neurotypical adults accomplished this using visual feedback to

induce explicit learning that helped correct the step length errors produced by the split-belt

treadmill (Roemmich et al. 2016). They found explicit learning improved performance during

split-belt walking compared to a group that did not receive feedback. However, the implicit

aftereffects (measured without visual feedback) were similar between groups, indicating

explicit learning did not impact the recalibration of motor commands (i.e., implicit

adaptation). Thus, the authors concluded that, within the same locomotor learning task, while

explicit learning improves overall performance, implicit adaptation proceeds despite

involvement from explicit learning in individuals with intact neurologic systems. Critically, it

is unclear to what degree explicit learning versus implicit adaptation is impaired in

individuals post-stroke when assessed in a task requiring dissociable contributions from both.

While explicit learning and implicit adaptation are broadly intact after stroke when

assessed individually during gait, it is unknown if this holds when they are combined in the

same task. Only two studies, both in reaching movements, have attempted to tackle this

question, but with mixed results (Binyamin-Netser et al. 2023; Taylor and Ivry 2014),

indicating more work is necessary to determine if there are deficits in these learning

processes when assessed within the same task. This is a critical gap because gait

rehabilitation post-stroke involves a combination of explicit learning and implicit adaptation

(e.g., a patient may explicitly try to increase step length based on their therapist's instructions

while simultaneously implicitly adapting to a more compliant walking surface). Therefore,

determining how each is impaired, when occurring together in the same task, has important

implications for how rehabilitation of locomotor tasks should be optimally structured.

The purpose of this study was to determine if individuals with chronic, hemiparetic

stroke demonstrate impaired explicit learning and/or implicit adaptation during a locomotor

task involving dissociable contributions from both processes. We accomplished this through a

combination of behavioral testing and computational modeling. Since cortical structures, and

in particular, prefrontal regions contribute to cognitive processes that are often impaired in

stroke (Barker-Collo and Feigin 2006), we hypothesized that individuals with stroke would

demonstrate impaired explicit learning compared to controls. Additionally, because the rate

of implicit adaptation on the split-belt treadmill is slow but the magnitude is intact in persons

post-stroke (Reisman et al. 2007; Savin et al. 2013; Tyrell et al. 2014) we hypothesized that

individuals with stroke would demonstrate similar levels of implicit adaptation as controls.

Materials and methods

<u>Participants</u>

We recruited 21 (10 Female) individuals with one prior unilateral, stroke to

participate in this study and 18 (9 Female) healthy age- and sex-matched control participants.

Individuals with stroke were included if they were between 18 and 85 years old, had a single

unilateral hemiparetic stroke (confirmed by an MRI or CT scan) more than 6 months prior,

and were able to walk without assistance from another person. Individuals with stroke were

excluded if they had evidence of cerebellar stroke, other neurologic diagnoses aside from

stroke, inability to walk outside of the home prior to stroke, pain limiting walking, neglect, or

significant aphasia. Control participants were excluded if they had any conditions that might

limit their walking or motor learning, any neurologic conditions, or uncorrected vision or

hearing loss. All individuals provided written informed consent prior to participating and the

study was approved by the University of Delaware Institutional Review Board.

Experimental design

To determine if individuals with stroke have impaired explicit learning or implicit

adaptation during a locomotor learning task that requires contributions from both processes,

we combined the split-belt adaptation paradigm with real-time visual feedback, similar to a

previous study (Figure 1A; Roemmich et al. 2016). Participants performed 4 phases of

treadmill walking: Baseline, Practice, Adaptation, and De-adaptation (Figure 1B). During

Baseline and Practice, both the treadmill belts moved at the same speed, set at half the speed

of the fast belt. During the Baseline phase, no visual feedback was provided on the screen and

individuals were told to "walk comfortably". The Practice phase served to introduce

participants to the visual feedback and ensure the individuals with stroke could respond to the

visual feedback by changing their step length. Step length targets first appeared at each

participant's baseline step length for 90 seconds, at which point, they were verbally oriented

to the feedback and instructed to practice changing their step lengths by stepping both above

and below the targets. For the next 30 seconds, the step length targets shifted 10 cm longer

for the limb taking the longer baseline step and 10 cm shorter for the limb taking the shorter

baseline step. The targets shifted back to the baseline step lengths for the final 60 seconds of

the Practice phase and individuals were asked to "walk comfortably". We confirmed that

participants in the stroke and control groups could change their step lengths from baseline in

response to the visual feedback (step length change from targets shift to walk comfortably,

stroke fast = 4.6 cm SD 3.6, slow = -3.0 SD 3.5; control fast = 7.8 SD 3.5, slow = -7.2 SD

3.2).

During the Adaptation phase (8 minutes), the fast belt speed was set at the fastest

overground gait speed (stroke group mean = 1.29 m/s SD 0.38; control = 1.81 SD 0.22), and

the slow belt moved at half the speed of the fast belt, producing a 2:1 speed ratio

(Charalambous et al. 2018; Tyrell et al. 2014). To ensure there were not large differences in

treadmill speeds between the stroke and control groups, we constrained the fast treadmill belt

speed between 0.6 and 1.0 m/s for all participants. This constraint resulted in both groups

walking at similar speeds (stroke group mean = 0.94 m/s SD 0.12; control = 1.0 SD 0). For

participants in both groups, the limb that took the longer step during the Baseline phase was

placed on the fast belt. This perturbation produces a large asymmetry of the left and right step

lengths (defined as the distance between two feet at heel strike), and is corrected on a stride-

by-stride basis through implicit adaptation (Charalambous et al. 2018; Reisman et al. 2005,

2007). Lastly, participants performed a De-adaptation phase (8 minutes) where they were

instructed to "walk comfortably", and both belts moved at the same speed as the Baseline

phase (i.e., the slow belt speed) so that we could measure the size of the implicit aftereffect,

our measure of the total magnitude of implicit adaptation.

To assess explicit learning, defined as the ability to consciously correct for errors

between a movement outcome and the task goal, we provided visual feedback of the left and

right step lengths during the first 40 strides of the Adaptation phase. The real-time visual

feedback was displayed on a vertically orientated LCD television screen placed 100 cm in

front of the treadmill (Figure 1A; Size: 123.3 x 71.1 cm; Sony Tokyo, Japan). The Motion

Monitor software (Innovative Sports Training Inc., Chicago, IL, USA) was used to display

the visual feedback during the experiment. The feedback consisted of a target grid of 12

possible step lengths, each 10 cm in height. This grid had a 1:1 correspondence with the

actual step length. The left and right step length feedback was displayed as a red and blue

foot, respectively. Each foot was presented in the center of the row corresponding to that step

length window, and appeared as soon as heel strike was detected, then disappeared once the

subsequent swing phase began. The target right and left step lengths during the Adaptation

phase were set at each participant's left and right baseline step lengths, denoted by

highlighting the corresponding row of the grid. Participants were instructed to "hit the

targets" when the feedback was visible. Therefore, because the targets were set at baseline

step length, the feedback guided participants to voluntarily correct the step length asymmetry

induced by the split-belt treadmill via explicit learning. We chose these targets to prevent any

ambiguity in the explicit task goal, and we reasoned that baseline step lengths were most

parsimonious targets to use.

The key manipulation that allowed us to assess the magnitude of explicit learning was

to turn off the feedback after the first 40 strides of Adaptation and instruct participants to

"walk comfortably". Here, we relied on the flexibility of explicit learning, assuming it could

be voluntarily "switched off", leaving no residual aftereffects due to implicit adaptation

(Bond and Taylor 2015; McDougle and Taylor 2019; Wood et al. 2020). Meaning, after the

instructions and removal of feedback, only implicit adaptation remained. Thus, our measure

of the total magnitude of explicit learning was the difference between the step length

behavior when the feedback was on and when it was first turned off (Roemmich et al. 2016).

Figure 1 Experimental Design. (A) Individuals walked on a split-belt treadmill with a vertically mounted television screen in front of them. The visual feedback was a grid of 12 different step lengths, each 10 cm in height. The step length feedback was represented on the screen as blue (left) and red (right) feet that appeared on the screen as soon as heel strike was detected and disappeared once toe off was detected. **(B)** All participants completed 4 walking phases: 1) A Baseline (Bsl) phase of normal walking where no feedback was on the screen; 2) A Practice phase where individuals were introduced to the visual feedback while walking (purple shading); 3) An Adaptation phase where the slow belt (dotted black line) moved at half the speed of the fast belt (solid black line), with feedback activated during only the first 40 strides (purple shading); 4) A De-adaptation phase where the belts returned to the same speed. The length of each phase (in minutes) is displayed in the grey shading at the bottom of the figure.

Data collection

During all phases of treadmill walking, individuals wore a ceiling mounted harness

(that did not provide body weight support) and held a handrail to prevent falls. Additionally,

we monitored heart rate (Polar, Kempele, Finland) for safety, and determined perceived

exertion using the Borg Rate of Perceived Exertion (RPE) scale after each walking phase. If

participants exceeded >80% their age-predicted max heart rate, the treadmill was stopped,

and the participant was provided with a seated rest break until their heart rate recovered.

However, no included participants needed the treadmill to stop during any of the walking

phases.

Participants walked on a dual belt treadmill that captured kinetic data through two

force plates, one under each belt, at 1000 Hz (Bertec, Columbus, OH, USA). Kinematic data

were captured and recorded at 100 Hz using a Vicon MX40, 8-camera motion capture

system, and time-synchronized with the kinetic data in Nexus software (v2.8.2, Vicon Motion

Systems, Inc., London, UK). We used a custom marker set with seven retroreflective

markers, one for each heel, lateral malleolus, and fifth metatarsal head, and the left medial

malleolus.

Data analysis

Step length, calculated in real-time using motion capture and the Motion Monitor, was

defined as the anterior-posterior distance between the two ankle markers at heel strike. Heel

strike was determined in real time using the following criteria: 1) The leading limb's heel

marker must be anterior with respect to the trailing limb's heel marker; 2) a ground reaction

force > one-third of the participants body weight detected through the treadmill force plate;

and 3) no ground reaction force > one-third of the participants body weight detected through

that same belt 40 ms prior to the detected step length.

The remainder of the data were analyzed with custom written MATLAB scripts

(vR2022a, Mathworks, Natick, MA, USA). Step lengths were used to calculate step length

asymmetry on each stride (s):

$step\ length\ asymmetry_{[s]} = \dfrac{fast\ step\ length_{[s]} - slow\ step\ length_{[s]}}{fast\ step\ length_{[s]} + slow\ step\ length_{[s]}} * 100\%$ (1)

Thus, values of 0 indicate perfect symmetry between the fast and slow step lengths, while

values further from 0 indicate greater asymmetry. We baseline corrected this measure by

subtracting the mean step length asymmetry during the Baseline phase from each stride in the

experiment for each individual. We removed outlier step length asymmetry strides, defined as

any step length asymmetry exceeding 3x the interquartile range of that participant's step

length asymmetry (mean percent removed [min max] stroke = 0.14% [0.0 0.39]; control =

0.92% [0.0 2.32]). This prevented strides outside the normal range of variability from

influencing the results. We also removed the first stride of each phase to account for treadmill

acceleration.

We used step length asymmetry data to calculate an Adaptation Index for each

individual for each stride (s; Charalambous et al. 2018; Mawase et al. 2014; Roemmich et al.

2016):

$Adaptation\ Index_{[s]} = \dfrac{step\ length\ asymmetry_{[s]} - condition * perturbation}{|perturbation|}$ (2a)

$condition = \begin{cases} 1, if\ belts\ are\ split \\ 0, if\ belts\ are\ tied \end{cases}$ (2b)

$perturbation = \begin{cases} min(step\ length\ asymmetry), if\ condition_{[s]} = 1 \\ max(step\ length\ asmmyetry), if\ condition_{[s]} = 0 \end{cases}$ (2c)

The min and max step length asymmetries used to determine the perturbation in equation 2c

were calculated only within the first 10 strides of each respective phase (Charalambous et al.

2018). Thus, during the Adaptation phase, an Adaptation Index of 0 represents the minimum

step length asymmetry (i.e., the max perturbation), and 1 indicates the perturbation has been

fully corrected. The reverse is true during the De-adaptation phase.

To test our hypotheses, we averaged Adaptation Index during 4 key timepoints of

interest: 1) Feedback On: the final five strides of the feedback being on during Adaptation, 2)

Feedback Off: the first five strides immediately after the feedback was turned off during

Adaptation, 3) End Adaptation: the last five strides of Adaptation, and 4) Implicit Aftereffect:

the first five strides of De-adaptation. Our hypothesis for explicit learning was tested by

comparing Feedback On (implicit adaptation plus explicit learning), to Feedback Off

(implicit adaptation only). Larger differences between Feedback On and Feedback Off

indicate greater explicit learning. We assessed the interaction between group (stroke vs

control) and time (Feedback On vs Feedback Off) as our primary behavioral measure testing

for impaired explicit learning in the stroke group. A secondary behavioral measure of explicit

learning was between group differences during Feedback On, as this reflected the ability of

individuals to use the visual feedback during implicit adaptation. Our primary behavioral

measure to assess implicit adaptation, defined as the recalibration of step lengths caused by

the split-belt treadmill, was comparing the Implicit Aftereffect between groups (Reisman et

al. 2005; Shadmehr and Mussa-Ivaldi 1994). A secondary behavioral measure of implicit

adaptation was a between groups comparison at End Adaptation.

<u>Computational modeling</u>

Since the behavioral analysis only provides a brief (i.e., 5 strides) and somewhat

arbitrary window into explicit learning and implicit adaptation, we used a computational

model to characterize each learning process. This approach allowed us to map the underlying

learning processes and their subcomponents onto each participant's behavior. Specifically,

we fit the model to individual data to obtain a unique set of parameter values for each

participant. Since these parameters represent specific aspects of explicit and implicit learning,

we can make inferences regarding the function of these underlying learning components

(Charalambous et al. 2018; Mawase et al. 2014; Roemmich et al. 2016). Then, we compared

the individual learning processes (i.e., model parameters) between the stroke and control

groups.

This "voluntary correction" model was previously used to capture explicit and

implicit learning in this paradigm (Roemmich et al. 2016), and the implicit adaptation

component of the model can successfully capture split-belt adaptation behavior in individuals

with stroke (Charalambous et al. 2018). The computational modeling used here followed that

of Roemmich and colleagues which defines the Adaptation Index (x) on each stride (s) as the

sum of both explicit learning ($x_{explicit}$) and implicit adaptation ($x_{implicit}$):

$$x_{[s]} = x_{explicit[s]} + x_{implicit[s]} \qquad\qquad (3)$$

Both processes correct for the same error ($error_{[s]} = perturbation_{[s]} - x_{implicit\ [s]}$), where

the perturbation = 1 during Adaptation and 0 during De-adaptation. Therefore, this model

assumes that the nervous system is correcting for this error using the implicit processes, and it

further reduces the error using the explicit process (Roemmich et al. 2016). Explicit learning

is only active when the feedback is on:

$$x_{explicit[s+1]} = \begin{cases} B_{explicit} * error_{[s]}, & if\ feedback\ is\ on \\ 0, & if\ feedback\ is\ off \end{cases} \qquad (4)$$

The free parameter, $B_{explicit}$, represents the learning rate for explicit learning as it is the

proportion of error that is explicitly corrected from one stride to the next (i.e., higher values

indicate faster learning). The implicit adaptation process has dual components, fast and slow,

and is active throughout the Adaptation and De-adaptation phases (Roemmich et al. 2016;

Smith et al. 2006):

$x_{implicit[s]} = x_{fast[s]} + x_{slow[s]}$ (5a)

$x_{fast[s+1]} = A_{fast}x_{fast[s]} + B_{fast}error_{[s]}$ (5b)

$x_{slow[s+1]} = A_{slow}x_{slow[s]} + B_{slow}error_{[s]}$ (5c)

Implicit learning has four free parameters. The learning rates, B_{fast} and B_{slow}, represent the

proportion of the error that is implicitly corrected from one stride to the next, and the

retention rates, A_{fast} and A_{slow}, represent the proportion of the current adapted state that is

retained. The fast process quickly learns from errors, but also quickly forgets, while the slow

process takes longer to learn from errors but retains longer (Smith et al. 2006).

<u>Model fitting and model comparison</u>

334   We fit the voluntary correction model to each participant's Adaptation Index data

during the Adaptation and De-adaptation phases using MATLAB's fmincon function, setting

the objective function as the sum of squared errors between the model output (x) and the data.

All parameters were constrained between 0 and 1. Additionally, we constrained the fast-

learning rate, B_{fast}, to be at least 5 times higher than the slow learning rate, B_{slow}; and the slow

retention rate, A_{slow}, was constrained to be greater than the fast retention rate, A_{fast}

(Roemmich et al. 2016; Smith et al. 2006). To ensure stable fits, we initialized the implicit

process parameters to the same values (Charalambous et al. 2018; Roemmich et al. 2016;

Smith et al. 2006) based on a prior locomotor adaptation study in individuals with stroke

(Charalambous et al. 2018): $A_{fast} = 0.92$, $B_{fast} = 0.03$, $A_{slow} = 0.996$, $B_{slow} = 0.004$. The

explicit parameter was initialized at uniformly random values between 0 and 1. To improve

the stability of the $B_{explicit}$ parameter, we performed 10 initializations for each participant. We

calculated individual fits (r^2) to this model, but due to the greater stride-to-stride variability

and idiosyncratic behavior of some individuals with stroke, the fits were worse for the stroke

group compared to controls. As we were primarily interested in capturing group-level

behaviors, we also performed a bootstrapping procedure to obtain a more consistent estimate

of both group's fits, while also quantifying our level of uncertainty around the sample means

(Charalambous et al. 2018; Kim et al. 2019; Smith et al. 2006; Taylor and Ivry 2011; Tsay et

al. 2021). We resampled with replacement the stride-by-stride data across participants 1000

times separately for each group, obtaining an average learning function for each bootstrap

sample. We then fit the model and calculated an r^2 value for each bootstrapped sample. We

report the mean and 95% confidence intervals of these r^2 values.

To ensure that we did not overfit the data with the five-parameter, voluntary

correction model, we also fit two simpler models to the data. For the single-rate model, the

motor output (x; i.e., Adaptation Index) is the result of a single process that has two

parameters B, the learning rate, and A the retention rate (Smith et al. 2006; Thoroughman and

Shadmehr 2000):

$x_{[s+1]} = A\, x_{[s]} + B\, error_{[s]}$ (6)

Additionally, we fit a four-parameter, dual-rate model to the data (Charalambous et al. 2018;

Roemmich et al. 2016; Smith et al. 2006). This dual-rate model represents motor output (x;

i.e., Adaptation Index) as the sum of a fast and slow process. This model is equivalent to the

implicit process in the voluntary correction model (equations 5a-c). Thus, neither alternative

model includes a voluntary correction, or explicit learning component. We compared fits of

the three different models using Akaike Information Criterion (AIC). To determine the

method of model comparison, we performed a model recovery analysis. We simulated the

experiment 100 times with each model, the subsequently fitted each model to the three

simulations. We calculated the number of times the model that simulated the data was best at

fitting its own simulation (which should be expected if the models can be distinguished;

Wilson and Collins 2019). We performed this procedure using both AIC and BIC and found

that AIC was better at distinguishing between the three models for the current experiment

(Wilson and Collins 2019).

Statistical analysis

For our Bayesian statistical analysis, we assumed the Adaptation Index data at each

timepoint of interest (Feedback On, Feedback Off, End Adaptation, Implicit Aftereffect) was

sampled from a normal distribution, with a mean which depended on both within subject (i.e.,

time) and between subject (i.e., group) parameters as well as an interaction parameter. We

confirmed a normal distribution was a reasonable assumption for our data by observing

favorable model comparison results against a student's t distribution. We estimated the

posterior distribution for these means (and all the statistical model parameters) using Bayes

rule, combining the evidence from our data with our prior assumptions about each parameter.

The prior distributions for the between and within subject effects were set as a wide normal

distribution centred on 0, and the prior for the standard deviation was set as a wide, positive-

valued uniform distribution. Thus, our prior assumptions only served to make our inferences

more conservative and did not bias the posterior. We estimated the joint posterior distribution

in Python v4.3.0 using the PyMc 4 library (Salvatier et al. 2016) and the bayes-toolbox

Python package (Kim 2023). We used Markov Chain Monte Carlo sampling to sample from

joint posterior distribution 10,000 times with 2,000 tuning samples. We performed posterior

predictive checks to ensure that the posterior samples accurately represented the data

(Kruschke 2014; McElreath 2016).

This procedure allowed us to report the full range of credible differences between the

groups along with the probability of a difference, given our data. To accomplish this, we

compared the posterior distributions of between group differences which are presented as

histograms representing the full distribution of possible differences based on the data we

collected. For each posterior distribution, we report the mean and 95% high density interval

(HDI), defined as the narrowest span of credible values that contain 95% of the distribution

(Kruschke 2014). The HDI can be interpreted as the true value falling between this range

with 95% certainty. Therefore, the 95% HDI provides an estimate of the size of the effect

with 95% certainty. We also report the probability of a difference as a percentage of posterior

distribution samples on one side of zero. Therefore, the $p_{difference}$ value provides an estimate of

the reliability of the effect. We do not provide a specific decision rule for the $p_{difference}$ values

as we would an alpha value. Rather, we made our inferences based on the actual degree of

certainty of the difference, with larger $p_{difference}$ values indicating greater certainty.

Results

Of the 21 individuals recruited to participate in the stroke group, we removed 4 from

the analysis because they either could not complete the task (1 due to double vision, and 2

due to fatigue that required stopping the treadmill) or they did not follow instructions (n=1).

Average participant characteristics for each group are displayed in Table 1. In Figure 2, we

display the mean, baseline-corrected step length asymmetry data during the Adaptation and

De-adaptation phases for both groups. For ease of group comparisons, we present our

primary analyses using the Adaptation Index. We note that similar results were obtained

when using step length asymmetry index, with no impact on any of our inferences.

	Stroke Group (n=17)	Control Group (n=18)
Age (years)	64.5 ± 10.2	64.8 ± 9.6
Sex	9M / 8F	9M / 9F
Time since stroke (months)	71.0 ± 49.1	
Side of brain lesion	7R/10L	
Self-selected (overground) walking speed (m/s)	0.92 ± 0.27	1.33 ± 0.28
Fastest (overground) walking speed	1.29 ± 0.38	1.81 ± 0.22
Fast treadmill belt speed	0.94 ± 0.12	1.00 ± 0
Lower Extremity Fugl Meyer	25.41 ± 6.27	

Table 1 Group characteristics. Demographic and clinical characteristics of participants. All
continuous variables are represented as mean ± 1 SD. (F = female, M = male, R = right, L =
left)

Figure 2 Step length asymmetry. Mean baseline-corrected step length asymmetry for each
group for the Adaptation and De-adaptation phases. Purple shading is the time when the
feedback was on. The vertical dashed line separates the Adaptation and De-adaptation phases.
Each phase was truncated to the participant with the shortest phase for visualization purposes.
Shading represents SEM.

In Figure 3, we display the mean Adaptation Index data and key timepoints of interest

for each group. First, we determined if individuals with stroke had impairments in explicit

learning (Figure 3B). Based on our instructions and previous work (Roemmich et al. 2016),

we assumed participants used explicit learning only while the feedback was on during the

Adaptation phase. Therefore, explicit learning magnitude was characterized as the difference

in Adaptation Index between Feedback On and Feedback Off. This difference was larger for

the control group (mean interaction effect [95% HDI] = 0.09 [-0.05 0.25], $p_{difference}$ = 88.2%),

providing evidence that the individuals with stroke had diminished explicit learning

compared to controls. Additionally, individuals in the stroke group were less able to use the

visual feedback during Adaptation compared to controls (Figure 3C), with much lower

Adaptation Index values during Feedback On (mean group difference = 0.23 [0.11 0.34],

$p_{difference}$ = 100.0%). Combined, these results point to impairments in explicit learning in

individuals with stroke compared to controls.

Next, we determined if individuals with stroke had impaired implicit adaptation by

comparing the size of the implicit aftereffect (Figure 3D). The control group demonstrated

larger implicit aftereffects compared to the stroke group (group difference = 0.10 [-0.02

0.21], $p_{difference}$ = 94.3%), providing evidence that individuals with stroke have impaired

implicit adaptation compared to controls. Additionally, we found large and reliable

differences between the groups at End Adaptation (group difference = 0.17 [0.07 0.28],

$p_{difference}$ = 99.9%). Overall, the behavioral results indicate that impairments may exist in both

explicit learning and implicit adaptation.

Figure 3 Adaptation Index. (**A**) Group averaged Adaptation Index data for the Adaptation and De-adaptation phases. The dashed line represents the walking period when the belts were split (i.e., the perturbation). Purple shading represents the time when the feedback was turned on. For visualization purposes, data for each phase were truncated to the individual with least number of strides. Solid lines represent group means, and shading represents SEM. (**B**) Group and individual data for the Feedback On and Feedback Off timepoints. Thick lines represent the group average slopes. (**C**) Group and individual data for the Feedback On timepoint. (**D**) Group and individual data for the Implicit Aftereffect timepoint. For panels C-D, bars represent group means, error bars represent 1 SD and smaller dots represent individuals. The insets display a histogram of the posterior distribution for the between group differences. The black vertical line in the histogram is there to aid visualization of the credibility of a between group difference (i.e., how much of the posterior probability distribution is on one side of zero). We report the 95% HDI regarding the range of credible effect sizes above the insets of the posterior distributions.

To provide a more complete picture of how each individual learning process contributed to overall adaptation, we applied a series of computational models to the data. The voluntary correction model, specifically, allowed us to map each individual's behavior to

explicit learning and implicit adaptation processes (Figure 4; see supplemental Figures S1-S2

for individual fits). This model fit the bootstrapped data well (stroke group mean r^2

[bootstrapped 95% CI] = 0.70 [0.11 0.89]; control group = 0.90 [0.81 0.95]). The model fits

to the individual stroke group's data were lower on average as compared to the control group

(mean [95% HDI] stroke = 0.31 [0.16 0.47]; control = 0.65 [0.53 0.77]), which is not

surprising given that stroke participants tend to be more heterogeneous than age-matched

controls. Nevertheless, based on the bootstrapping analysis (Fig. 4A and B), the voluntary

correction model reasonably approximated the systematic group-level behavior caused by the

experimental paradigm, as was our goal. Importantly, we confirmed that the voluntary

correction model had better (lower) AICs then both the single rate and the dual rate model for

both the stroke (single rate AIC difference mean [95% HDI] = 111 [54 167], $p_{difference}$ =

100.0%; dual rate AIC difference = 36 [-20 89], $p_{difference}$ = 89.9%) and control groups (single

rate AIC difference = 281 [226 336], $p_{difference}$ = 100.0%; dual rate AIC difference = 89 [35

142]; $p_{difference}$ = 99.9%), indicating that the voluntary correction model accurately

characterizes learning on this task without overfitting. As the single- and dual-rate state-space

models do not include a voluntary correction process, these results also support our

assumption that explicit learning contributed to behavioral change specifically when visual

feedback was on. Overall, this analysis provides further support for quantifying the level of

explicit learning impairment for the stroke group using the $B_{explicit}$ parameter

Comparing the individual parameters from the voluntary correction model allowed us

to determine the specific components of learning that were impaired. The learning rate

parameter for explicit learning, $B_{explicit}$, served as a measure of each individual's explicit

learning ability, with higher values indicating faster explicit learning (Figure 4C). The stroke

group had much smaller $B_{explicit}$ values compared to the control group (group difference =

0.23 [0.06 0.40], $p_{difference}$ = 99.5%), providing strong support for the hypothesis that explicit

learning is impaired in individuals with stroke compared to controls. Next, we examined the

four implicit adaptation process parameters (Figure 4F-G). While there was evidence of

differences between groups for most parameters, the magnitude of differences for three of the

four were near zero (group differences: A_{slow} = 0.00 [-0.00 0.01], $p_{difference}$ = 87.7%; B_{slow} =

0.00 [-0.00 0.00], $p_{difference}$ = 68.2%, B_{fast} = 0.01 [-0.01 0.03], $p_{difference}$ = 84.7%). In contrast,

there was a marked difference in the retention rate for the fast state (A_{fast} group difference =

0.09 [-0.04 0.24], $p_{difference}$ = 92.6%). Thus, it appears that individuals with stroke, as a group,

have a specific impairment in their ability to retain what was learned by the fast implicit

adaptation process. In sum, the results of our computational modeling provided strong

support for the hypothesis that explicit learning is impaired post-stroke and revealed that the

retention rate for the fast state could underlie slower implicit adaptation in stroke.

Figure 4 Computational model results. Mean model fits to bootstrapped samples plotted
against the bootstrapped stride-by-stride data for the **(A)** stroke group and **(B)** control group.
See supplemental Figures 1 and 2 for the model fits for each individual participant. Purple
shading represents the time when the feedback was turned on. For visualization purposes,
data for each phase were truncated to the individual with least number of strides. Shading
represents 1 SD of the bootstrapped samples (i.e., the standard deviation of sample means or
standard error of the mean) **(C-G)** Model parameter values for each group. Bars represent

group means and error bars represent 1 SD and smaller dots represent individuals. The insets
are histograms of the posterior of the between groups difference (contrast) in parameter
values. We report the 95% HDI regarding the range of credible effect sizes above the insets
of the posterior distributions. Note the scale of the x-axis varies for these inset plots.

Discussion

In the current study, we examined explicit learning and implicit adaptation within the

same locomotor learning task in individuals with chronic, hemiparetic stroke. We combined a

behavioral manipulation and computational modeling to determine the presence, and

potential degree, of impairment in both learning processes. The majority of work in

locomotor learning and stroke has primarily studied implicit adaptation (Betschart et al. 2017;

Charalambous et al. 2018; Reisman et al. 2007; Savin et al. 2013; Tyrell et al. 2014), with

less attention paid to explicit learning (French et al. 2021b, 2021a). While some gait studies

have examined both processes in the same task (Cherry-Allen et al. 2018; Day et al. 2019;

Moore et al. 2022; Mutha et al. 2011; Schaefer et al. 2009), only two studies in reaching have

attempted to discern their individual contributions to overall motor learning (Binyamin-

Netser et al. 2023; Taylor and Ivry 2014). To our knowledge, the current study is the first to

assess explicit and implicit motor learning within the same task in individuals with chronic

stroke using both behavioral manipulations and computational modeling. While our sample

size is relatively small, our results provide strong evidence that stroke impairs explicit

learning and the rate of implicit adaptation during a locomotor task that elicits dissociable

contributions from both processes. This has important implications for the design of

locomotor learning tasks in post-stroke rehabilitation because many interventions involve

both explicit and implicit components.

Explicit learning is impaired in chronic stroke

We found that individuals with chronic, hemiparetic stroke have impairments in explicit learning in a locomotor task involving both explicit learning and implicit adaptation. Individuals with stroke had a smaller change in behavior compared to controls after the visual feedback, intended to drive and support explicit learning, was removed. Additionally, the computational modeling revealed significantly slower explicit learning in individuals with stroke. To ensure these results were not due to motor control deficits that prevented participants with stroke from effectively implementing explicit strategies, we performed a series of Bayesian regressions to test the effect of lower extremity function (quantified by Lower Extremity Fugl-Meyer [LEFM] scores) on our learning measures. Our analyses showed that there was essentially no effect of LEFM on the $B_{explicit}$ parameter (beta coefficient = 0.005 [-0.01 0.02]), or on our behavioral measure of explicit learning (beta coefficient = 0.01 [-0.004 0.02]), providing further evidence that the behavioral differences observed were due to explicit learning deficits in participants with stroke.

While there are few prior studies assessing both explicit learning and implicit adaptation in the same task after stroke, one experiment in reaching showed that individuals with lateral prefrontal cortex (LPFC) lesions demonstrate impairments in explicit learning (Taylor and Ivry 2014). This and other work raises the possibility that cognitive processes such as working memory or general cognition in reaching studies (Anguera et al. 2011; Binyamin-Netser et al. 2023; Seidler et al. 2012; Vandevoorde and Orban de Xivry 2020), and fluid cognition in gait (French et al. 2021a) contribute to explicit motor learning, but more work is required to determine the specific contribution of cognition to explicit learning in stroke. Future work could use techniques such as lesion mapping along with computational modeling to provide a mechanistic explanation for the explicit learning impairments we observed.

Contrary to the current findings, prior work in reaching (Binyamin-Netser et al. 2023)

and gait (French et al. 2021b, 2021a) observed no differences in explicit learning in

individuals with stroke compared to controls. However, the studies in gait used a primarily

explicit task without a split-belt perturbation, likely making the task easier, which could

reduce the ability to detect explicit learning deficits in stroke. The reaching study dissociated

explicit learning and implicit adaptation using visual cues (the color and shape of a cursor). It

is possible that either this manner of distinguishing between explicit and implicit processes or

the broader inclusion criteria for their stroke group can account for the differences between

their findings and those of the current study. Similar to Taylor and Ivry (Taylor and Ivry

2014), we provided clear instructions and removed all visual feedback to ensure explicit

learning was minimized, and provided a narrower range of inclusion criteria, potentially

explaining why our results were more consistent with theirs. Still, it is critical to determine if

the manner of eliciting explicit learning (a specific type of cue or instruction) impacts the

ability to use this process in stroke given its ubiquity in rehabilitation settings.

<u>Slower implicit adaptation in stroke is due to worse retention of the fast process</u>

Contrary to our hypothesis, we found evidence that implicit adaptation is impaired

after stroke. The stroke group demonstrated smaller implicit aftereffects and a lower plateau

at the end of the Adaptation phase indicating a smaller overall magnitude of implicit

adaptation. Prior studies in locomotor adaptation after stroke indicate that the overall

magnitude of implicit adaptation is similar to controls, but the rate is slower (Malone and

Bastian 2014; Reisman et al. 2007; Savin et al. 2013; Tyrell et al. 2014). Therefore, the

slower rate of adaptation in stroke, combined with the relatively short Adaptation phase in the

current study (8 minutes compared to 10-15 minutes in the prior studies), could have

prevented us from observing asymptotic adaptation. While we assume that these impairments

in implicit adaptation during split-belt walking are related to learning from sensory prediction

error, it is also possible that split-belt adaptation is driven by energy optimization (Finley et

al. 2013; Sánchez et al. 2019, 2020; Sánchez and Finley 2018), a theory that could be tested

in individuals with stroke.

While it may seem that the visual feedback interfered with implicit adaptation for the

stroke group, prior work in young individuals with intact neurologic systems demonstrate that

visual feedback used to either help or hinder performance during split-belt walking does not

change the total magnitude of implicit adaptation (Long et al. 2016; Malone and Bastian

2010; Roemmich et al. 2016). Additionally, individuals with stroke can successfully adapt to

the split-belt treadmill while also explicitly learning to change a separate gait parameter (knee

flexion angle) using visual feedback (Cherry-Allen et al. 2018). Therefore, it is unlikely that

explicit learning itself hindered implicit adaptation in the current study since implicit

adaptation proceeds in spite of explicit learning, and even worsens performance in some

cases (Mazzoni and Krakauer 2006; Taylor and Ivry 2011), across reaching and walking

paradigms (Mazzoni and Krakauer 2006; Roemmich et al. 2016), including in stroke

(Binyamin-Netser et al. 2023; Cherry-Allen et al. 2018; Taylor and Ivry 2014).

The computational modeling utilized in this study provides insight into why the

learning rate of implicit adaptation was impaired in stroke in this task. The voluntary

correction model incorporates a dual-rate model of adaptation which frames implicit

adaptation as the combination of a fast state and a slow state (Smith et al. 2006). These states

represent updates to an internal model, a prediction of the sensory consequences of

movement, that could occur either in the cerebellum or motor cortex (Smith et al. 2006). One

theory suggests that the motor cortex is responsible for retention of the adapted state while

the cerebellum is responsible for learning (Galea et al. 2011; Li et al. 2001). Thus, damage to

motor cortices or possibly its outputs could explain poor retention of the fast process in

individuals with stroke. Alternatively, the fast process has been closely linked to explicit

learning during visuomotor rotation tasks (Krakauer et al. 2019; Taylor and Ivry 2014).

However, to date there is no evidence of contributions from explicit learning to standard

split-belt adaptation (i.e., without additional visual feedback; Charalambous et al. 2018;

Mawase et al. 2014; Roemmich et al. 2016). Another possibility is that the fast state

represents a reactive balance element that is sensitive to environmental changes (Mariscal et

al. 2023). Future studies are required to dissociate between these potential explanations.

Conclusion

Motor learning involves multiple processes, both explicit and implicit, that work

together to improve overall task performance. We found that individuals with chronic stroke

have impairments in explicit learning and implicit adaptation during a locomotor task that

elicits dissociable contributions from both. These findings are important because of the

potential application to post-stroke rehabilitation, which often combines different forms of

learning in a single task. To improve outcomes, future work should determine how locomotor

rehabilitation interventions can be structured to target these deficits and optimize overall

motor learning.

Supplemental Material

Supplemental Figures S1–S2: https://osf.io/pws2k/.

Data Availability

Source data for this study are openly available at https://osf.io/pws2k/

Acknowledgements

We would like to thank Saunders Penn for helping to create Figure 1A.

A preprint of this manuscript is available at https://doi.org/10.1101/2024.02.04.578807.

Grants

The authors acknowledge receipt of the following financial support for the research,

authorship, and/or publication of this article: This work was supported by the National

Institutes of Health to Darcy S. Reisman [NIH 2R01HD078330-05A1].

Disclosures

The Authors declare that there is no conflict of interest.

Author Contributions

JMW, DSR, HEK Conceived and designed research; HW, LF, JMW performed experiments;

JMW analyzed data; JMW, ET, SMM, DSR, HEK interpreted results of experiments; JMW

prepared figures; JMW drafted manuscript; All authors edited and revised manuscript; All

authors approved final version of manuscript.

References

Anguera JA, **Reuter-Lorenz PA**, **Willingham DT**, **Seidler RD**. Failure to Engage Spatial
Working Memory Contributes to Age-related Declines in Visuomotor Learning. *J Cogn*
Neurosci 23: 11–25, 2011.

Barker-Collo S, **Feigin V**. The impact of neuropsychological deficits on functional stroke
outcomes. *Neuropsychol Rev* 16: 53–64, 2006.

Betschart M, **Lauzière S**, **Miéville C**, **McFadyen BJ**, **Nadeau S**. Changes in lower limb
muscle activity after walking on a split-belt treadmill in individuals post-stroke. *J Electromyogr*
Kinesiol Off J Int Soc Electrophysiol Kinesiol 32: 93–100, 2017.

Binyamin-Netser R, **Goldhamer N**, **Avni I**, **Ressel Zviely A**, **Shmuelof L**. Cognitive
Impairments After Stroke Do Not Attenuate Explicit Visuomotor Adaptation in Reaching and
Savings With the Unaffected Arm. *Neurorehabil Neural Repair* 37: 444–457, 2023.

Bond KM, **Taylor JA**. Flexible explicit but rigid implicit learning in a visuomotor adaptation
task. *J Neurophysiol* 113: 3836–3849, 2015.

Charalambous CC, **Alcantara CC**, **French MA**, **Li X**, **Matt KS**, **Kim HE**, **Morton SM**,
Reisman DS. A single exercise bout and locomotor learning after stroke: physiological,
behavioural, and computational outcomes: Effect of exercise on locomotor learning in chronic
stroke survivors. *J Physiol* 596: 1999–2016, 2018.

Cherry-Allen KM, **Statton MA**, **Celnik PA**, **Bastian AJ**. A dual-learning paradigm
simultaneously improves multiple features of gait post-stroke. *Neurorehabil Neural Repair* 32:
810–820, 2018.

Day KA, **Cherry-Allen KM**, **Bastian AJ**. Individualized feedback to change multiple gait
deficits in chronic stroke. *J NeuroEngineering Rehabil* 16: 158, 2019.

Dietz V, **Zijlstra W**, **Duysens J**. Human neuronal interlimb coordination during split-belt
locomotion. *Exp Brain Res* 101, 1994.

Finley JM, **Bastian AJ**, **Gottschall JS**. Learning to be economical: the energy cost of walking
tracks motor adaptation: Split-belt adaptation reduces metabolic power. *J Physiol* 591: 1081–
1095, 2013.

French MA, **Cohen ML**, **Pohlig RT**, **Reisman DS**. Fluid Cognitive Abilities Are Important
for Learning and Retention of a New, Explicitly Learned Walking Pattern in Individuals After
Stroke. *Neurorehabil Neural Repair* 35: 419–430, 2021a.

French MA, **Morton SM**, **Charalambous CC**, **Reisman DS**. A locomotor learning paradigm
using distorted visual feedback elicits strategic learning. *J Neurophysiol* 120: 1923–1931,
2018.

French MA, **Morton SM**, **Reisman DS**. Use of explicit processes during a visually guided
locomotor learning task predicts 24-h retention after stroke. *J Neurophysiol* 125: 211–222,
2021b.

Galea JM, **Vazquez A**, **Pasricha N**, **de Xivry J-JO**, **Celnik P**. Dissociating the roles of the
cerebellum and motor cortex during adaptive learning: the motor cortex retains what the
cerebellum learns. *Cereb Cortex N Y N 1991* 21: 1761–1770, 2011.

Huberdeau DM, **Krakauer JW**, **Haith AM**. Practice induces a qualitative change in the
memory representation for visuomotor learning. *J Neurophysiol* 122: 1050–1059, 2019.

Hussain SJ, **Hanson AS**, **Tseng S-C**, **Morton SM**. A locomotor adaptation including explicit
knowledge and removal of postadaptation errors induces complete 24-hour retention. *J
Neurophysiol* 110: 916–925, 2013.

Kim HE. bayes-toolbox: A Python package for Bayesian statistics. *J Open Source Softw* 8:
5526, 2023.

Kim HE, **Parvin DE**, **Ivry RB**. The influence of task outcome on implicit motor learning.
eLife 8: e39882, 2019.

Krakauer JW, **Hadjiosif AM**, **Xu J**, **Wong AL**, **Haith AM**. Motor Learning. In:
Comprehensive Physiology. American Cancer Society, p. 613–663, 2019.

Krakauer JW, **Pine ZM**, **Ghilardi M-F**, **Ghez C**. Learning of visuomotor transformations
for vectorial planning of reaching trajectories. *J Neurosci* 20: 8916–8924, 2000.

Kruschke J. *Doing Bayesian Data Analysis: A Tutorial with R, JAGS, and Stan*. Academic
Press, 2014.

Li CS, **Padoa-Schioppa C**, **Bizzi E**. Neuronal correlates of motor performance and motor
learning in the primary motor cortex of monkeys adapting to an external force field. *Neuron*
30: 593–607, 2001.

Long AW, **Roemmich RT**, **Bastian AJ**. Blocking trial-by-trial error correction does not
interfere with motor learning in human walking. *J Neurophysiol* 115: 2341–2348, 2016.

Malone LA, **Bastian AJ**. Thinking about walking: effects of conscious correction versus
distraction on locomotor adaptation. *J Neurophysiol* 103: 1954–1962, 2010.

Malone LA, **Bastian AJ**. Spatial and Temporal Asymmetries in Gait Predict Split-Belt
Adaptation Behavior in Stroke. *Neurorehabil Neural Repair* 28: 230–240, 2014.

Malone LA, **Bastian AJ**, **Torres-Oviedo G**. How does the motor system correct for errors in
time and space during locomotor adaptation? *J Neurophysiol* 108: 672–683, 2012.

Mariscal DM, **Fjeld KL**, **Torres-Oviedo G**. Characterization of the adaptation and
generalization dynamics from high-dimensional neuromuscular data. In: *Advances in Motor
Learning and Motor Control (MLMC)*. Advances in motor learning and motor control.
Washington D.C.: 2023.

Mawase F, **Shmuelof L**, **Bar-Haim S**, **Karniel A**. Savings in locomotor adaptation explained
by changes in learning parameters following initial adaptation. *J Neurophysiol* 111: 1444–
1454, 2014.

Mazzoni P, **Krakauer JW**. An implicit plan overrides an explicit strategy during visuomotor
adaptation. *J Neurosci* 26: 3642–3645, 2006.

McDougle SD, **Ivry RB**, **Taylor JA**. Taking Aim at the Cognitive Side of Learning in
Sensorimotor Adaptation Tasks. *Trends Cogn Sci* 20: 535–544, 2016.

McDougle SD, **Taylor JA**. Dissociable cognitive strategies for sensorimotor learning. *Nat
Commun* 10: 40, 2019.

McElreath R. *Statistical Rethinking: A Bayesian Course with Examples in R and Stan*. CRC
Press, 2016.

Moore RT, **Piitz MA**, **Singh N**, **Dukelow SP**, **Cluff T**. Assessing Impairments in Visuomotor
Adaptation After Stroke. *Neurorehabil Neural Repair* 36: 415–425, 2022.

Morton SM, **Bastian AJ**. Cerebellar contributions to locomotor adaptations during splitbelt
treadmill walking. *J Neurosci* 26: 9107–9116, 2006.

Mutha PK, **Sainburg RL**, **Haaland KY**. Left Parietal Regions Are Critical for Adaptive
Visuomotor Control. *J Neurosci* 31: 6972–6981, 2011.

Reisman DS, **Block HJ**, **Bastian AJ**. Interlimb coordination during locomotion: what can be
adapted and stored? *J Neurophysiol* 94: 2403–2415, 2005.

Reisman DS, **Wityk R**, **Silver K**, **Bastian AJ**. Locomotor adaptation on a split-belt treadmill
can improve walking symmetry post-stroke. *Brain* 130: 1861–1872, 2007.

Roemmich RT, **Bastian AJ**. Closing the Loop: From Motor Neuroscience to
Neurorehabilitation. *Annu Rev Neurosci* 41: 415–429, 2018.

Roemmich RT, **Long AW**, **Bastian AJ**. Seeing the Errors You Feel Enhances Locomotor
Performance but Not Learning. *Curr Biol* 26: 2707–2716, 2016.

Salvatier J, **Wiecki TV**, **Fonnesbeck C**. Probabilistic programming in Python using PyMC3.
PeerJ Comput Sci 2: e55, 2016.

Sánchez N, **Finley JM**. Individual Differences in Locomotor Function Predict the Capacity to
Reduce Asymmetry and Modify the Energetic Cost of Walking Poststroke. *Neurorehabil
Neural Repair* 32: 701–713, 2018.

Sánchez N, **Simha SN**, **Donelan JM**, **Finley JM**. Taking advantage of external mechanical
work to reduce metabolic cost: the mechanics and energetics of split-belt treadmill walking. *J
Physiol* 597: 4053–4068, 2019.

Sánchez N, **Simha SN**, **Donelan JM**, **Finley JM**. Using asymmetry to your advantage:
learning to acquire and accept external assistance during prolonged split-belt walking. *J
Neurophysiol* 125: 344–357, 2020.

Savin DN, **Tseng S-C**, **Whitall J**, **Morton SM**. Poststroke Hemiparesis Impairs the Rate but
not Magnitude of Adaptation of Spatial and Temporal Locomotor Features. *Neurorehabil
Neural Repair* 27: 24–34, 2013.

Schaefer SY, **Haaland KY**, **Sainburg RL**. Dissociation of initial trajectory and final position
errors during visuomotor adaptation following unilateral stroke. *Brain Res* 1298: 78–91, 2009.

Schmidt RA, **Lee TD**. *Motor control and learning: A behavioral emphasis, 4th ed.*
Champaign, IL, US: Human Kinetics, 2005.

Seidler RD, **Bo J**, **Anguera JA**. Neurocognitive Contributions to Motor Skill Learning: The
Role of Working Memory. *J Mot Behav* 44: 445–453, 2012.

Shadmehr R, **Mussa-Ivaldi FA**. Adaptive representation of dynamics during learning of a
motor task. *J Neurosci* 14: 3208–3224, 1994.

Shadmehr R, **Smith MA**, **Krakauer JW**. Error Correction, Sensory Prediction, and
Adaptation in Motor Control. *Annu Rev Neurosci* 33: 89–108, 2010.

Smith MA, **Ghazizadeh A**, **Shadmehr R**. Interacting adaptive processes with different
timescales underlie short-term motor learning. *PLoS Biol* 4: e179, 2006.

Taylor JA, **Ivry RB**. Flexible cognitive strategies during motor learning. *PLoS Comput Biol*
7, 2011.

Taylor JA, **Ivry RB**. The role of strategies in motor learning. *Ann N Y Acad Sci* 1251: 1–12,
2012.

Taylor JA, **Ivry RB**. Cerebellar and prefrontal cortex contributions to adaptation, strategies,
and reinforcement learning. *Prog Brain Res* 210: 217–253, 2014.

Thoroughman KA, **Shadmehr R**. Learning of action through adaptive combination of motor
primitives. *Nature* 407: 742–747, 2000.

Tsay JS, **Lee A**, **Ivry RB**, **Avraham G**. Moving outside the lab: The viability of conducting
sensorimotor learning studies online. *Neurons Behav Data Anal Theory* 5: 1–22, 2021.

Tseng Y, **Diedrichsen J**, **Krakauer JW**, **Shadmehr R**, **Bastian AJ**. Sensory prediction errors
drive cerebellum-dependent adaptation of reaching. *J Neurophysiol* 98: 54–62, 2007.

Tyrell CM, **Helm E**, **Reisman DS**. Learning the spatial features of a locomotor task is slowed
after stroke. *J Neurophysiol* 112: 480–489, 2014.

Vandevoorde K, **Orban de Xivry J-J**. Why is the explicit component of motor adaptation
limited in elderly adults? *J Neurophysiol* 124: 152–167, 2020.

Wilson RC, **Collins AG**. Ten simple rules for the computational modeling of behavioral data.
eLife 8: e49547, 2019.

Wood JM, **Kim HE**, **French MA**, **Reisman DS**, **Morton SM**. Use-dependent plasticity
explains aftereffects in visually guided locomotor learning of a novel step length asymmetry.
J Neurophysiol 124: 32–39, 2020.

Wood JM, **Morton SM**, **Kim HE**. The Consistency of Prior Movements Shapes Locomotor
Use-Dependent Learning. *eNeuro* 8, 2021.

Supplemental material

Explicit and implicit locomotor learning in individuals with chronic hemiparetic stroke.

Authors: Jonathan M. Wood, Elizabeth Thompson, Henry Wright, Liam Festa, Susanne M.
Morton, Darcy S. Reisman, and Hyosub E. Kim
798

Supplemental Figures: Individual model fits

Supplemental Figure S1 Individual model fits for the control group. The voluntary
correction model was fit to each participant's Adaptation Index data in the control group
(n=18). Black dots represent the participant's Adaptation Index on that stride and the black
dashed line represents the perturbation (which equals one during the Adaptation phase and 0
during De-adaptation phase). The red function represents the model's motor output (x), the
light blue function (active only when the feedback is turned on – denoted by the gray shading)
represents the explicit component ($x_{explicit}$), and the yellow and purple functions represent the
fast (x_{fast}) and slow (x_{slow}) components of the implicit process, respectively.
810

Supplemental Figure S2 Individual model fits for the stroke group. The voluntary
correction model was fit to each participant's Adaptation Index data in the stroke group (n=17).
Black dots represent the participant's Adaptation Index on that stride and the black dashed line
represents the perturbation (which equals one during the Adaptation phase and 0 during De-
adaptation phase). The red function represents the model's motor output (x), the light blue
function (active only when the feedback is turned on – denoted by the gray shading) represents
the explicit component ($x_{explicit}$), and the yellow and purple functions represent the fast (x_{fast})
and slow (x_{slow}) components of the implicit process, respectively.

820