Chronospaces: an R package for the statistical exploration of divergence times

promotes the assessment of methodological sensitivity

ABSTRACT

[1] Much of our understanding of the history of life hinges upon time calibration, the process of

assigning absolute times to cladogenetic events. Bayesian approaches to time scaling phylogenetic trees

have dramatically grown in complexity, and depend today upon numerous methodological choices.

Arriving at objective justifications for all of these is difficult and time consuming. Thus, divergence times

are routinely inferred under only one or a handful of parametric conditions, often times chosen

arbitrarily. Progress towards building robust biological timescales necessitate the development of better

methods to visualize and quantify the sensitivity of results to these decisions.

[2] Here, we present an R package that assists in this endeavor through the use of chronospaces, i.e.,

graphical representations summarizing variation in the node ages contained in time-calibrated trees. We

further test this approach by estimating divergence times for three empirical datasets—spanning widely

differing evolutionary timeframes—using the software PhyloBayes.

[3] Our results reveal large differences in the impact of many common methodological decisions, with

the choice of clock (uncorrelated vs. autocorrelated) and loci having strong effects on inferred ages.

Other decisions have comparatively minor consequences, including the use of the computationally

intensive site-heterogeneous model CAT-GTR, whose effect might only be discernible for exceedingly old

divergences (e.g., the deepest eukaryote nodes).

[4] The package *chronospace* implements a range of graphical and analytical tools that assist in the

exploration of sensitivity and the prioritization of computational resources in the inference of divergence

times.

Keywords: chronospaces, divergence times, diversification, fossils, phylogenetic inference,

phylogenomics, sensitivity analysis, time calibration

INTRODUCTION

The discovery of the "molecular evolutionary clock" by Zuckerkandl and Pauling (1965), who first

described the relative constancy in the rates of change of molecular sequences both through time and

across clades, is a hallmark of the advent of molecular biology. By establishing a proportionality between

the amount of amino acid or nucleotide differences that can be directly observed and the time that must

have elapsed since those sequences last shared a common ancestor, the molecular clock greatly expands

the insights that can be drawn from phylogenetic trees. As such, scaling phylogenies to absolute (or

geologic) time is often the first step to explore macroevolutionary patterns and processes at deep

timescales, and one on which modern comparative biology relies on (Pennell & Harmon 2013).

Bayesian methods of divergence time estimation have become highly sophisticated (Kumar &

Hedges 2016), being able to accommodate among-lineage rate variation through the use of relaxed

clocks (Thorne *et al.* 1998; Drummond *et al.* 2006), exploit diverse sources of both character and

temporal information (Ronquist *et al.* 2012; O'Reilly *et al.* 2015), and use complex tree-building models

that account for fossilization and sampling rates (Heath *et al.* 2014) and accommodate gene tree

discordances (Rannala *et al.* 2020). While this increased complexity has allowed for an improved

modelling of diversification, it has also drastically increased the number of methodological choices that

the average scientist is confronted with. Dating software require users to specify which among several

competing models of diversification, trait evolution, and among-lineage rate variation—the three distinct

elements of the tripartite framework of Bayesian dating (Warnock & Wright 2020)—are enforced. Each

of these models includes a sizable number of parameters for which prior probability distributions need

to be specified. Furthermore, even though progress has been made regarding computational efficiency

(Dos Reis & Yang 2019; Douglas *et al.* 2022; Barba-Montoya *et al.* 2023), most methods remain limited in

terms of the size of the datasets that can be run within reasonable times, forcing further decisions

regarding taxon sampling and locus choice. The choice of number, location, and shape of fossil-derived

temporal constraints adds yet another layer of complexity to these analyses. These methodological

decisions can have a sizable effect on the inference of Bayesian time-calibrated trees (Warnock *et al.*

2011; Magallón *et al.* 2013; Eme *et al.* 2014; Mongiardino Koch *et al.* 2022; Sauquet *et al.* 2022)—also

known as chronograms—producing potentially incongruent reconstructions of biological diversification.

Two broad approaches have been explored to overcome this methodological uncertainty. First,

numerous methods have been developed to justify model selection, allowing the identification of a set

of parametric conditions considered optimal. These methods, however, have their own limitations:

model fit statistics do not reliably discriminate between partition and mixture models (Crotty & Holland

2022), while alternative tree priors might not be amenable to comparison using standard methods such

as Bayes factors (May & Rothfels 2023). General purpose approaches, such as cross-validation, are too

computationally consuming to be used exhaustively (Zhou *et al.* 2007; Lartillot 2023), while methods

designed to tackle specific aspects of the analysis can arrive at opposite conclusions depending on

unassessed decisions (e.g., detecting autocorrelated rates can depend on which model of substitution is

used to infer the tree being evaluated; Mongiardino Koch *et al.* 2022). The alternative to the use of

model selection algorithms involves sensitivity analyses, which explore the robustness of results to a

range of conditions (e.g., Sanderson & Doyle 2001; Britton *et al.* 2007; Warnock *et al.* 2011; Howard *et*

al. 2022; Mongiardino Koch *et al.* 2022). However, there is a dearth of tools that provide efficient ways of

summarizing multiple chronograms in either quantitative or visual ways, as well as a lack of criteria for

gauging the relative impact that different parameters have on the final results.

Here, we present *chronospace*, an R package (R Core Team 2022) for the statistical exploration of

time calibration. The package implements a set of visualization techniques that allow for a better

assessment of the sensitivity of divergence times to methodological decisions. It also provides new tools

to quantify the overall impact of each individual methodological choice, providing an estimate of its

effect size. We illustrate its functionalities by inferring divergence times for three empirical datasets

spanning widely differing evolutionary timescales. We do so using the program PhyloBayes v.4.1 (Lartillot

et al. 2013) which implements the complex site-heterogeneous model CAT-GTR (Lartillot & Philippe

2004).

MATERIALS AND METHODS

A) Chronospaces: the approach

Chronospaces are low-dimensional graphical representations that summarize variation in the

absolute node ages of populations of chronograms (Mongiardino Koch *et al.* 2022). They are similar to

treespaces (Hillis *et al.* 2005) in that they provide a visual summary of populations of phylogenetic trees,

but focus exclusively on temporal information. To do so, the chronograms analyzed must share the same

set of nodes, representing posterior distributions of Bayesian time-calibrated analyses run under a fixed

topology. While this limits their application, a two-step approach in which a preferred topology is first

inferred and then time scaled has become common in the phylogenomic literature due to the

computational burden of simultaneous inference.

Whereas treespaces are generated from distance matrices that capture topological similarity

(Smith 2022)—and are thus representations that are disconnected from the original variables—

chronospaces are built using ordination methods that reorganize variation, capturing major patterns in a

few synthetic axes. Here, we rely on between-group principal component analysis (bgPCA; Dolédec &

Chessel 1987; Dolédec & Chessel 1989; Yendle & MacFie 1989), an alternative to traditional PCA that

ordinates the covariance matrix of multivariate means for predefined groups of observations. Within the

context of a chronospace, posterior topologies constitute the 'observations', and the ages of each of

their internal nodes are the 'variables' (see also Fig. 1). When analyzing populations of topologies

inferred under different analytical conditions (e.g., alternative clock models), these constitute distinct

levels of a grouping variable. The estimated bgPCA axes will be those capturing the largest variation in

group (i.e., level) means, and onto which individual chronograms are projected. The result is an

ordination that maximizes the separation between groups through a rigid rotation of space, preserving

information regarding which nodes are more variable in age, and which factors this variation is

associated with. The approach will result in $g - 1$ bgPCA axes, where g is the number of groups, with

the remainder of multidimensional structure occurring on residual axes. Within the biological literature,

bgPCA has been mostly used to explore the discrimination of groups of observations using microarray

(Culhane *et al.* 2002) and geometric morphometric (Cardini *et al.* 2019) data. While bgPCA can introduce

spurious patterns when the number of variables exceeds the number of observations (Cardini *et al.*

2019), this does not represent an issue within the context of chronospaces as the number of nodes in a

phylogenetic tree (i.e., the number of variables) will typically be much lower than the number of trees

sampled in a Bayesian posterior distribution (i.e., the number of observations).

bgPCA axes capture the main directions of change in divergence times that are associated with a

given methodological decision. If the chronograms being analyzed are derived from a set of inferences

varying multiple parametric conditions, separate bgPCA rotations can be applied to the same

chronospace, deriving a set of eigenvectors and eigenvalues for each one of the factors being varied. The

proportion of the total variance accounted for by each factor, as well as the proportion that remains

unaccounted, is estimated both for the entire chronospace as well as for each individual bgPCA axis by

calculating the corresponding sums of squares for the balanced model. Chronospaces thus provide a

means of summarizing the entire temporal information of individual chronograms, and exploring the

effects (i.e., direction and magnitude of change) associated with alternative dating approaches. This

information can be used to aid in the interpretation of the overall sensitivity of divergence times, as well

as rank methodological choices from the most to the least impactful.

B) *chronospace:* the package

chronospace is an R package (R Core Team 2022) available for installation from the GitHub

repository (removed for anonymity). It relies on functions from phylogenetic packages *ape* (Paradis &

Schliep 2018), *phangorn* (Schliep 2011), and *phytools* (Revell 2012), visualization packages *deeptime*

(Gearty 2022), *ggpubr* (Kassambara 2020), *ggtree* (Yu *et al.* 2017), and *patchwork* (Pedersen 2020), as

well as data handling functionalities from the *tidyverse* (Wickham 2017).

The range of functions and capabilities implemented in the package are depicted in Figure 1 and

further documented on the GitHub repository. The typical workflow would begin with the use of

function `extract_ages` which gathers node ages from posterior distributions of Bayesian timetrees

inferred under a constrained topology. Typically, this would include trees from multiple runs (as most of

the methods implemented are designed to facilitate the comparison of results obtained under different

analytical conditions), but some analyses can be done with a single set of trees. Chronograms are read

from files in Newick format (one per run and/or chain). While not all phylogenetic software generate this

type of output, trees can be first converted and saved to file using functions from other R packages, such

as *treeio* (Wang *et al.* 2020). Function `extract_ages` can then import the trees, extract and format

the node ages contained in them, and annotate the data with a set of factors representing the

parameters changed between different runs. The constrained tree topology is also saved in order to

generate graphical output. This information (ages, factors, and topology) is stored as an object of class

`nodeAges`. Multiple graphical outputs can be directly obtained from this object: posterior distributions

of node ages can be plotted using both `sensitive_nodes`, which helps identify and visualize the

nodes whose inferred ages change the most between analyses, or `specified_node`, which plots ages

for a target node defined by the user; lineage-through-time plots can also be generated using function

`ltt_sensitivity` to depict the overall effect of methodological choices on the inferred history of

diversification (including different options for portraying uncertainty). Finally, bgPCA-rotated

chronospaces are generated using function `chronospace`, which creates an object of the same class,

and its associated printing and plotting functions summarize the results of the analysis in both graphical

and quantitative ways. The remainder of this article focuses on highlighting the use of the latter.

Figure 1: Structure of the R package *chronospace* and analytical workflow, represented using the data

that accompanies the package. Populations of posterior topologies from time calibration analyses

present in the working directory are imported, and their node ages annotated and stored as an object of

class `nodeAges` using function `extract_ages`. The package provides functions to visualize the effect

induced by analytical conditions on a node of interest (`specified_node`), and on the overall history

of diversification (`ltt_sensitivity`). It also detects the nodes whose ages change the most

between analyses, and plots their posterior distributions (`sensitive_nodes`). These plots can

optionally include a chronostratigraphic scale (shown here in the lineage-through-time plot). Finally, a

`chronospace` object can be built and used to summarize the overall effect of methodological

decisions on the results. These can be visualized as bgPCA-summarized chronospaces (which will be

univariate or multivariate, depending on the number of levels of each factor) and/or chronogram

warping plots. The latter depict the patterns of branch length contraction/expansion that are captured

by each bgPCA axis, and are akin to the landmark deformation plots that typically accompany the

morphospace axes in a geometric morphometric analysis (hence, their name). All plots are generated

with *ggplot2* (Wickham 2016) and several extensions, providing the user with control over basic

visualization aspects (object colors, sizes, transparencies, etc.). Chronospace summaries can also be

printed in the form of a table reporting the partitioning of variance.

C) **Sensitivity of divergence times across three empirical datasets**

Three genome-scale datasets of protein-coding loci were obtained from the literature, designed

to tackle the phylogenetic relationships and divergence times of Curculionoidea (65 terminals, 517 loci;

Shin *et al.* 2018), Decapoda (90 terminals, 410 loci; Wolfe *et al.* 2019), and Eukaryota (136 terminals, 313

loci; Strassert *et al.* 2021). Studies were selected so as to span widely differing evolutionary time frames

(Figures 2). For each of these, we gathered a phylogenomic dataset (including a concatenated matrix

coded as amino acids and an associated partition file), the rooted tree topology preferred by the

authors, and a list of node age calibrations derived from the fossil record. If studies included multiple

outgroups, only those forming part of the sister-group clade to the ingroup were retained. The eukaryote

dataset lacked outgroups altogether.

Divergence times were inferred using PhyloBayes v.4.1 (Lartillot *et al.* 2013) using a constrained

tree topology. Four different methodological decisions were assessed: the type of clock, the model of

molecular evolution, the prior distribution applied to calibrated nodes, and the approach used to

subsample loci. The first three of these are parameters specified when running the program. Divergence

times were obtained applying both an autocorrelated log-normal (`-ln`; Thorne *et al.* 1998) and an

uncorrelated gamma (`-ugam`; Drummond *et al.* 2006) clock; implementing unpartitioned site-

homogenous GTR + G (`-gtr`; Tavaré 1986) or site-heterogenous CAT-GTR + G (`-cat -gtr`; Lartillot &

Philippe 2004) models of molecular evolution; and using prior distributions of node ages that were

either uniformly distributed between stratigraphic bounds (`-ilb`), or implemented a truncated Cauchy

minimum bound (`-lb`; Inoue *et al.* 2010). Default parameter values were used for the latter ($p = 0.1$, $c =$

1), which results in the prior probability peaking at 110% the age of the minimum bound. The tree prior

was set to a birth-death process with free parameters and soft calibration bounds were activated (Yang

& Rannala 2006), allowing for 5% of prior probability to lie outside of the specified bounds for calibrated

nodes (`-bd -sb`). Default settings were used for all other parameters.

Regarding the fourth and last decision evaluated (the set of loci used), phylogenomic datasets

were subsampled using strategies intended to achieve four distinct goals: A) minimize topological

conflict; B) improve levels of phylogenetic usefulness; C) maximize the amount of information; and D)

enhance clock-like behavior. For the first of these, loci chosen corresponded to those displaying the

minimum Robinson-Foulds (RF) distance (Robinson & Foulds 1981) between gene trees and the

constrained species tree. For the second, loci were chosen using the multivariate subsampling strategy

genesortR (Mongiardino Koch 2021; Mongiardino Koch & Thompson 2021), which identifies loci that

simultaneously display high phylogenetic signal and low levels of potential sources of bias. Subsampled

matrices with maximal amount of information were built from loci with the highest levels of occupancy.

Finally, clock-likeness was estimated using the variance of root-to-tip distances displayed by gene trees.

In the presence of outgroups, gene trees were rooted ahead of estimation using the node corresponding

to the last common ancestor of all ingroup taxa; otherwise, gene trees were midpoint rooted along a

random internal edge. This is expected to cause few problems for the curculionoid and decapod datasets

for which outgroup occupancy was 98.6% and 81.0%, respectively, but could pose an issue for the

eukaryote dataset. To explore this caveat, we estimated the variance of root-to-tip distances for all

outgroup-containing gene trees using both outgroup and random midpoint rooting. Correlations

between the two were high (Peason's ρ = 0.83 for Curculionoidea and 0.92 for Decapoda, both p <

0.001). When used to select the top fifth most clock-like loci, both approaches agreed on 75.5%

(Curculionoidea) and 78.8% (Decapoda) of them, further reinforcing that loci displaying clock-like

evolution can be recognized in the absence of outgroups. Given the positive relationship between the

variance of root-to-tip distances and the overall rate of evolution of genes (Mongiardino Koch *et al.*

2022; see Fig. S1), loci were first sorted by rate (estimated using the overall tree length; Telford *et al.*

2014), and both the fastest- and slowest-evolving 15% of loci were excluded. Loci chosen corresponded

to those with minimum root-to-tip variances from among those showing relatively average evolutionary

rates. Gene trees necessary to measure these properties were inferred using ParGenes v.1.0.1 (Morel *et*

al. 2018), using models that minimized the Bayesian Information Criterion and estimating support with

100 bootstrap replicates. A fifth subsampled dataset was derived by selecting loci at random, providing a

means of estimating the overall effect of targeted subsampling schemes. To standardize these different

matrices in terms of the overall amount of data contained, the loci necessary to first exceed a total

length of 10,000 positions were concatenated.

Dating was performed under all combinations of these four methodological decisions, resulting

in 40 conditions per dataset, each of which was run using two independent chains for at least 20,000

generations. Posterior topologies were obtained using the verbose output option of the program

readdiv, setting a 50% burn-in and thinning chains to every two generations (`-x 10000 2 -v`), for a

total of 5,000 posterior trees per chain. Convergence was visually assessed using log-likelihood trace

plots (Fig. S2) and comparing median posterior estimates of node ages and model parameters between

replicate chains (Figs. S3-S6). After confirming convergence was attained, time-calibrated trees from

separate chains were combined into a single file. The remainder of analyses were performed using

package *chronospace*, summarizing each run using a random sample of 500 posterior trees. R code to

reproduce each of these steps is available as Supplementary Material S1.

RESULTS

Figure 2 summarizes the results obtained across the three empirical datasets used, showing the

median and 95% confidence interval (CI) of each inferred node age after combining the posterior

distributions of individual runs. Producing such overall summaries of temporal information across

populations of chronograms is straightforward once the data has been gathered and formatted using the

function `extract_ages`. For the ancient divergences estimated in the eukaryote dataset, pooled 95%

CI can span upwards of 1,000 Ma, as is the case for example of the last common ancestor of Amebozoa

(median: 1,291.2 Ma, 95% CI: 701.2-1,841.2 Ma). The temporal uncertainty associated with younger

divergences is much narrower in absolute terms, yet it can be even larger in relative terms (i.e., after

dividing the 95% CI width by its median value). For example, a handful of divergences between closely

related achelate decapods show 95% CI widths that are six to seven times the value of the median

estimate, with plausible dates spanning the last 142 Ma and falling anywhere between the Quaternary

and Early Cretaceous.

Figure 2: Uncertainty and sensitivity of node age estimates. Values shown are the median and width of pooled 95% CI per node across all analytical conditions. Fossil calibrations were placed on nodes highlighted with black margins. Insets show the distribution of the relative widths of pooled 95% CIs, obtained by dividing absolute widths by the median values. Only unconstrained nodes were tallied.

The spread of posterior dates shown in Figure 2 is a function of the width of the individual CI of

each run (uncertainty) and the degree of overlap of the CIs obtained under different conditions

(sensitivity). Chronospaces can help disentangle the two by visualizing and quantifying the contribution

of each individual methodological choice to overall levels of sensitivity (Figure 3). In the case of the

empirical datasets analyzed, this approach reveals that: A) there is a remarkable difference (over 1,000x)

in the overall impact of the parameters explored; and, B) the type of molecular clock and the subsample

of loci have larger effect sizes for the all three datasets. The implementation of the complex site-

heterogeneous CAT-GTR model seems to have a strong effect only for the eukaryote dataset. The most

sensitive nodes per dataset to each of the four factors evaluated are found in Figures S7-S9.

Figure 3: Sensitivity of divergence time estimates to methodological choices. **A.** bgPCA-summarized

chronospaces for each factor, from top to bottom: type of molecular clock, prior distribution on

constrained nodes, model of molecular evolution, and gene subsample. The proportion of total variance

explained by each factor is shown in grey boxes. For the loci subsample, only the first two axes (of four)

are shown. Ellipses include 95% of chronograms. **B.** Proportion of total variance explained by each factor,

divided by that of the least important factor for each dataset. Axis is log-scaled.

276   Regarding loci subsampling, chronograms built using high signal and high usefulness loci were

277   very similar across datasets, defining one extreme of the first (major) bgPCA axis (Fig. 4A). The other end

278   of this axis was always populated by chronograms derived from high occupancy loci, and occasionally

279   (for example, in the case of the eukaryote dataset) also by random and clock-like schemes. The branch-

280   length warping plots shown in Figure 4B reveal that the selection of high signal/usefulness loci results in

281   systematically longer internal branches and shorter terminal ones.

Change in branch length (Ma) relative to chronospace origin

Figure 4: Effect of locus choice on inferred ages. **A.** Chronospaces built discriminating by loci

subsampling strategy, as also shown in Fig. 3A. Circles depict the centroids of each distribution; the

thickness of the lines connecting them is scaled to the inverse of the Euclidean distances separating

them. Locations of centroids are shown on the first two bgPCA axes only, distances were calculated using

all dimensions. **B.** Graphical representation of chronograms representing negative (top) and positive

289   (bottom) extremes of bgPC1. The expansion/contraction of branches relative to a chronogram at the

290   origin of bgPC1 is color coded (red/blue, respectively). sd = standard deviation along bgPCA 1; values < 1

291   are automatically selected by the package in order to avoid generating negative branch lengths.

292   Chronogram warpings induced by other factors are shown in Figures S10-S12.

DISCUSSION

Understanding evolutionary processes across geologic timescales cannot be done without first

establishing an accurate timeline of biological diversification. While phylogenomic datasets, and

methods able to handle them, have allowed us to tackle phylogenetic questions that were once

intractable (e.g., Philippe *et al.* 1994; Poe & Chubb 2004), they have also introduced a plethora of

methodological steps and decisions with downstream effects that we are only beginning to explore (Tan

et al. 2015; Li *et al.* 2021; Mongiardino Koch 2021; Lozano-Fernandez 2022; Foster *et al.* 2023). In the

context of modern Bayesian relaxed-clock dating, these decisions relate to the selection of one among

several models that determine how the tree is built, how the characters evolve, and how a clock is

enforced—adding up to dozens of parameters that require user-specified prior probability distributions

(Álvarez-Carretero & dos Reis 2020; Warnock & Wright 2020). Further decisions need to be taken

regarding the character information that is employed, the taxa that are sampled, and the temporal

constraints that are enforced. These choices can have strong consequences: they can determine whether

we think clades survived mass extinctions (Budd & Mann 2023), emerged through sudden bursts of

innovation (Dos Reis *et al.* 2015), participated in events of co-diversification with mutualistic partners

(Barba-Montoya *et al.* 2018), and diverged—or not—as a direct consequence of continental drift (Tarver

et al. 2016).

Most of these methodological decisions are also hard to justify. Coupled with the high

computational cost of analyses, empirical evolutionary timescales are generally inferred using one (or a

few) among hundreds of possible parametric conditions, often times chosen arbitrarily. A literal reading

of the literature assessing the sensitivity of results to methodological factors suggests that every one of

these decisions can have large impacts on inferred dates (Hugall *et al.* 2007; Inoue *et al.* 2010; Clark et al.

2011; Duchêne *et al.* 2014; Hirt *et al.* 2017; Barba-Montoya *et al.* 2018; Dos Reis *et al.* 2018; Smith *et al.*

2018; Carruthers & Scotland 2021; Chen *et al.* 2021), providing little *a priori* guidance regarding which

should be prioritized given limited computational resources. These assessments are likely also

compounded by the choice of dating software, as estimates of sensitivity might be driven by the range of

options available, their implementation, or even their default parameterization (which is not always

tuned, as is the case of the present study). Progress in pinpointing the magnitude of these effects has

been hampered by a general focus on a handful of nodes of interest rather than on tree-wide effects, a

scarcity of methods of visualization and summary, as well as of metrics that quantify the effect size of

individual decisions in a manner that is comparable, and a lack of testing of whether the relevance of

these choices scales with temporal depth.

Chronospaces provide novel ways of visualizing, quantifying, and exploring the sensitivity of

divergence time estimates, contributing to the inference of more robust evolutionary timescales

(Mongiardino Koch *et al.* 2022; Song *et al.* 2023; Wolfe *et al.* 2023). By representing chronograms as

collections of node ages, standard multivariate statistical approaches can be readily employed on

populations of Bayesian posterior timetrees. This representation allows for more nuanced visualizations

that can effectively convey the overall impact of different choices (Fig. 3A), the relative similarity of trees

obtained across runs (Fig. 4A), and the patterns of branch length warping that are induced (Fig. 4B). At

the same time, partitioning the total variance into proportions explained by discrete factors using sums

of squares can result in tree-wide estimates of sensitivity with which to rank methodological choices by

their relative importance, helping prioritize computational resources. While the current implementation

is limited to analyses run under a constrained tree topology, chronospaces could potentially be built

using the subset of ages derived from nodes present in all (or most) of the posterior trees of a

simultaneous-inference analysis.

As shown in Fig. 3B, some decisions affect our reconstruction of diversification histories much

more than others, with effect sizes that vary by three to four orders of magnitude. This extreme

heterogeneity can help redirect the allocation of computational effort. For example, our results suggest

that empirical studies should either carefully validate their choice of evolutionary clock, or run

inferences that vary this component of the analysis, before deriving evolutionary insights (as shown

previously in both simulated and empirical studies; see Ho *et al.* 2005; Duchêne *et al.* 2014; Delsuc *et al.*

2018; Dos Reis *et al.* 2018; Mongiardino Koch *et al.* 2022; Song *et al.* 2023). Other factors can be

arbitrarily fixed without significant consequences, as is the case of the shape of prior distributions on

calibrated nodes (as noted previously by Betts *et al.* 2018, although see Clarke *et al.* 2011). We note that

the lack of an effect detected here applies exclusively to the use of the Cauchy lower bound (Inoue *et al.*

2010) as implemented in PhyloBayes under default settings, and does not mean that the fossil

calibration scheme is of little relevance—in fact, ample evidence points to the opposite (e.g., Near *et al.*

2005; Magallón *et al.* 2013; Montagna *et al.* 2019). The use of prior distributions with other alternative

shapes, the enforcement of hard or soft bounds, the choice of different calibration points (as well as

their overall number and relative depths), are all major components of dating analyses that were not

evaluated here. Similarly, the software of choice can impart idiosyncratic effects (Bapst *et al.* 2016),

meaning that all conclusions drawn in this study could also be software-specific. The main rationale for

choosing PhyloBayes is that it remains the only dating software implementing the site-heterogeneous

CAT-GTR model (Lartillot & Philippe 2004). Despite ongoing controversies, this model holds preeminence

in the inference of phylogenomic topologies (Whelan & Halanych 2017; Li et al. 2021; Bujaki & Rodrigue

2022; Al Jewari & Baldauf 2023; Szánthó *et al.* 2023), yet its impact on dating had so far remained

unexplored. Our results suggest that the CAT-GTR model induces relatively minor effects. Such

robustness of divergence times to the choice of substitution model has been reported before (Du *et al.*

2019; Tao *et al.* 2020; Mongiardino Koch *et al.* 2022), implying that the cost of using computationally

intensive models is not outweighed by the benefits. Notably, the relatively high effect of the choice of

loci and clock are not affected by the timeframe of the phylogenetic question, and affect equally nodes

of Cenozoic age as well as divergences spanning the Proterozoic (Fig. 3B). Only two differences were

observed as deeper timescales are probed: there is a decrease in the impact of the loci subsample

coupled with an increase in that of the clock model (with both remaining the two most important factors

overall); and the choice of substitution model becomes slightly more relevant for extremely ancient (e.g.,

Precambrian) divergences. The generality of these effects remains to be tested with a broader sample of

datasets.

Loci subsampling has become a standard approach in phylogenomic dating, yet again, the effects

introduced by the each of numerous available approaches have not been fully characterized. While some

authors have advocated for the selection of clock-like genes in the hopes that reduced lineage-specific

rate variation will improve model fit and diminish bias, others have favored loci with less topological

conflict, as these are expected to harbor higher phylogenetic signal and be less affected by incomplete

lineage sorting (ILS). Simulations have confirmed that both of these subsampling schemes result in more

accurate divergence times (Smith *et al.* 2018; Carruthers *et al.* 2020; Carruthers *et al.* 2022). Other

sensible options include selecting genes based on phylogenetic usefulness (as estimated using

genesortR; Mongiardino Koch 2021), which should balance both aforementioned strategies by

discovering loci with increased clock-likeness and decreased topological incongruence; and maximizing

occupancy, increasing the amount of information contained within a matrix of a given size.

Our results show that locus choice is a major factor determining the divergence times obtained,

and that alternative schemes impart characteristic signatures on inferred chronograms. Topologically

congruent loci (targeted by both the signal and usefulness schemes) produce shorter terminal branches

and longer internal ones (particularly those subtending species pairs; Fig. 4B). The opposite effect is

induced by maximizing occupancy, and to some extent also by favoring clock-like loci. These same

patterns of branch length warping were found in the simulation study of Carruthers *et al.* (2022), in

which the selection of congruent topologies was tied to an expansion of internal branch lengths and a

reduction of terminal ones. This is in fact the expected behavior if topological incongruence is driven by

biological factors such as ILS (Mendes & Hahn 2016). While our analyses were not designed to test this

phenomenon, the results are consistent with ILS driving topological incongruence even at exceedingly

ancient evolutionary timescales. Even if targeting clock-like sequences can improve the modelling of rate

variation, doing so in a manner that does not also account for topological incongruence can result in the

inference of distorted evolutionary timescales.

REFERENCES

Al Jewari, C. & Baldauf, S.L. (2023). Conflict over the eukaryote root resides in strong outliers, mosaics

and missing data sensitivity of site-specific (CAT) mixture models. *Systematic Biology*, **72**, 1-16.

Álvarez-Carretero, S. & dos Reis, M. (2020) Bayesian phylogenomic dating. In S. Y. W. Ho (Ed.), *The*

Molecular Evolutionary Clock: Theory and Practice (pp. 221-249). Springer.

Bapst, D.W., Wright, A.M., Matzke, N.J. & Lloyd, G.T. (2016). Topology, divergence dates, and

macroevolutionary inferences vary between different tip-dating approaches applied to fossil

theropods (Dinosauria). *Biology Letters*, **12**, 20160237.

Barba-Montoya, J., Sharma, S. & Kumar, S. (2023) Molecular timetrees using relaxed clocks and uncertain

phylogenies. *Frontiers in Bioinformatics,* **3**, 10.3389/fbinf.2023.1225807.

Barba-Montoya, J., Dos Reis, M., Schneider, H., Donoghue, P.C. & Yang, Z. (2018) Constraining uncertainty

in the timescale of angiosperm evolution and the veracity of a Cretaceous Terrestrial Revolution.

New Phytologist, **218,** 819-834.

Betts, H.C., Puttick, M.N., Clark, J.W., Williams, T.A., Donoghue, P.C. & Pisani, D. (2018) Integrated

genomic and fossil evidence illuminates life's early evolution and eukaryote origin. Nature

Ecology & Evolution, **2**, 1556-1562.

Britton, T., Anderson, C.L., Jacquet, D., Lundqvist, S. & Bremer, K. (2007). Estimating divergence times in

large phylogenetic trees. *Systematic biology*, **56**, 741-752.

Budd, G.E. & Mann, R.P. (2023) Two notorious nodes: a critical examination of relaxed molecular clock

age estimates of the bilaterian animals and placental mammals. *Systematic Biology*, syad057,

10.1093/sysbio/syad057.

Bujaki, T. & Rodrigue, N. (2022). Bayesian cross-validation comparison of amino acid replacement

models: contrasting profile mixtures, pairwise exchangeabilities, and gamma-distributed rates-

across-sites. *Journal of Molecular Evolution*, **90**, 468-475.

Cardini, A., O'Higgins, P. & Rohlf, F.J. (2019) Seeing distinct groups where there are none: spurious

patterns from between-group PCA. *Evolutionary Biology,* **46,** 303-316.

Carruthers, T., Sanderson, M.J. & Scotland, R.W. (2020) The implications of lineage-specific rates for

divergence time estimation. *Systematic Biology,* **69,** 660-670.

Carruthers, T. & Scotland, R.W. (2021) Uncertainty in divergence time estimation. *Systematic Biology*, **70**,

858-861.

Carruthers, T., Sun, M., Baker, W.J., Smith, S.A., de Vos, J.M. & Eiserhardt, W.L. (2022) The implications of

incongruence between gene tree and species tree topologies for divergence time estimation.

Systematic Biology, **71,** 1124-1146.

Chen, D., Hosner, P.A., Dittmann, D.L., O'Neill, J.P., Birks, S.M., Braun, E.L. & Kimball, R.T. (2021)

Divergence time estimation of Galliformes based on the best gene shopping scheme of

ultraconserved elements. *BMC Ecology and Evolution,* **21,** 209.

Clark, J.T., Warnock, R.C.M. & Donoghue, P.C.J. (2011) Establishing a time-scale for plant evolution. *New*

Phytologist, **192**, 266-301.

Crotty, S.M. & Holland, B.R. (2022) Comparing partitioned models to mixture models: Do information

criteria apply? *Systematic Biology,* **71,** 1541-1548.

Culhane, A.C., Perriere, G., Considine, E.C., Cotter, T.G. & Higgins, D.G. (2002) Between-group analysis of

microarray data. *Bioinformatics,* **18,** 1600-1608.

Delsuc, F., Philippe, H., Tsagkogeorga, G., Simion, P., Tilak, M.-K., Turon, X., Lopez-Legentil, S., Piette, J.,

Lemaire, P. & Douzery, E.J. (2018) A phylogenomic framework and timescale for comparative

studies of tunicates. *BMC Biology,* **16,** 39.

Dolédec, S. & Chessel, D. (1987) Rythmes saisonniers et composantes stationnelles en milieu aquatique.

I: Description d'un plan d'observation complet par projection de variables. *Acta Oecologica.*

Oecologia Generalis, **8,** 403-426.

Dolédec, S. & Chessel, D. (1989) Rythmes saisonniers et composantes stationnelles en milieu aquatique.

II: Prise en compte et élimination d'effets dans un tableau faunistique. *Acta Oecologica.*

Oecologia Generalis, **10,** 207-232.

Dos Reis, M., Gunnell, G.F., Barba-Montoya, J., Wilkins, A., Yang, Z. & Yoder, A.D. (2018) Using

phylogenomic data to explore the effects of relaxed clocks and calibration strategies on

divergence time estimation: primates as a test case. *Systematic Biology,* **67,** 594-615.

Dos Reis, M., Thawornwattana, Y., Angelis, K., Telford, M.J., Donoghue, P.C. & Yang, Z. (2015) Uncertainty

in the timing of origin of animals and the limits of precision in molecular timescales. *Current*

Biology, **25,** 2939-2950.

Dos Reis, M. & Yang, Z. (2019) Bayesian molecular clock dating using genome-scale datasets. In M.

Anisimova (Ed.), *Evolutionary genomics. Methods in molecular biology, vol. 1910* (pp. 309-330).

Humana.

Douglas, J., Jiménez-Silva, C.L. & Bouckaert, R. (2022) StarBeast3: adaptive parallelized Bayesian

inference under the multispecies coalescent. *Systematic Biology,* **71,** 901-916.

Drummond, A.J., Ho, S.Y., Phillips, M.J. & Rambaut, A. (2006) Relaxed phylogenetics and dating with

confidence. *PLoS Biology,* **4,** e88.

Du, Y., Wu, S., Edwards, S.V. & Liu, L. (2019) The effect of alignment uncertainty, substitution models and

priors in building and dating the mammal tree of life. *BMC Evolutionary Biology,* **19,** 203.

Duchêne, S., Lanfear, R. & Ho, S.Y. (2014) The impact of calibration and clock-model choice on molecular

estimates of divergence times. *Molecular Phylogenetics and Evolution,* **78,** 277-289.

Eme, L., Sharpe, S.C., Brown, M.W. & Roger, A.J. (2014) On the age of eukaryotes: evaluating evidence

from fossils and molecular clocks. *Cold Spring Harbor Perspectives in Biology,* **6,** a016139.

Foster, P.G., Schrempf, D., Szöllősi, G.J., Williams, T.A., Cox, C.J. & Embley, T.M. (2023) Recoding amino

acids to a reduced alphabet may increase or decrease phylogenetic accuracy. *Systematic Biology,*

72, 723-737.

Gearty, W. (2022) deeptime: plotting tools for anyone working in deep time. R package version 0.2.2,

https://CRAN.R-project.org/package=deeptime.

Heath, T.A., Huelsenbeck, J.P. & Stadler, T. (2014) The fossilized birth–death process for coherent

calibration of divergence-time estimates. *Proceedings of the National Academy of Sciences,* **111,**

E2957-E2966.

Hillis, D.M., Heath, T.A. & John, K.S. (2005) Analysis and visualization of tree space. *Systematic Biology,*

54, 471-482.

Hirt, M.V., Arratia, G., Chen, W.-J., Mayden, R.L., Tang, K.L., Wood, R.M. & Simons, A.M. (2017) Effects of

gene choice, base composition and rate heterogeneity on inference and estimates of divergence

times in cypriniform fishes. *Biological Journal of the Linnean Society,* **121,** 319-339.

Ho, S.Y., Phillips, M.J., Drummond, A.J. & Cooper, A. (2005) Accuracy of rate estimation using relaxed-

clock models with a critical focus on the early metazoan radiation. *Molecular Biology and*

Evolution, **22,** 1355-1363.

Howard, R.J., Giacomelli, M., Lozano-Fernandez, J., Edgecombe, G.D., Fleming, J.F., Kristensen, R.M., Ma,

X., Olesen, J., Sørensen, M.V. & Thomsen, P.F. (2022) The Ediacaran origin of Ecdysozoa:

integrating fossil and phylogenomic data. *Journal of the Geological Society,* **179,** jgs2021-107.

Hugall, A.F., Foster, R. & Lee, M.S. (2007) Calibration choice, rate smoothing, and the pattern of tetrapod

diversification according to the long nuclear gene RAG-1. *Systematic Biology,* **56,** 543-563.

Inoue, J., Donoghue, P.C. & Yang, Z. (2010) The impact of the representation of fossil calibrations on

Bayesian estimation of species divergence times. *Systematic Biology,* **59,** 74-89.

Kassambara, A. (2020) ggpubr: 'ggplot2' based publication ready plots. R package version 0.4.0,

https://CRAN.R-project.org/package=ggpubr.

Kumar, S. & Hedges, S.B. (2016) Advances in time estimation methods for molecular data. *Molecular*

Biology and Evolution, **33,** 863-869.

Lartillot, N. (2023) Identifying the best approximating model in Bayesian phylogenetics: Bayes factors,

cross-validation or wAIC? *Systematic Biology,* **72,** 616-638.

Lartillot, N. & Philippe, H. (2004) A Bayesian mixture model for across-site heterogeneities in the amino-

acid replacement process. *Molecular Biology and Evolution,* **21,** 1095-1109.

497 Lartillot, N., Rodrigue, N., Stubbs, D. & Richer, J. (2013) PhyloBayes MPI. Phylogenetic reconstruction

498   with infinite mixtures of profiles in a parallel environment. *Systematic Biology,* **62,** 611-615.

499 Li, Y., Shen, X.-X., Evans, B., Dunn, C.W. & Rokas, A. (2021) Rooting the animal tree of life. *Molecular*

500   *Biology and Evolution,* **38**, 4322-4333.

501 Lozano-Fernandez, J. (2022) A practical guide to design and assess a phylogenomic study. *Genome*

502   *Biology and Evolution,* **14,** evac129.

503 Magallón, S., Hilu, K.W. & Quandt, D. (2013). Land plant evolutionary timeline: gene effects are

504   secondary to fossil constraints in relaxed clock estimation of age and substitution rates.

505   *American Journal of Botany*, **100**, 556-573.

506 May, M.R. & Rothfels, C.J. (2023) Diversification models conflate likelihood and prior, and cannot be

507   compared using conventional model-comparison tools. *Systematic Biology,* **72,** 713-722.

508 Mendes, F.K. & Hahn, M.W. (2016) Gene tree discordance causes apparent substitution rate variation.

509   *Systematic Biology,* **65,** 711-721.

510 Mongiardino Koch, N. (2021) Phylogenomic subsampling and the search for phylogenetically reliable loci.

511   *Molecular Biology and Evolution,* **38**, 4025-4038.

512 Mongiardino Koch, N. & Thompson, J.R. (2021) A total-evidence dated phylogeny of Echinoidea

513   combining phylogenomic and paleontological data. *Systematic Biology,* **70,** 421-439.

514 Mongiardino Koch, N., Thompson, J.R., Hiley, A.S., McCowin, M.F., Armstrong, A.F., Coppard, S.E.,

515   Aguilera, F., Bronstein, O., Kroh, A. & Mooi, R. (2022) Phylogenomic analyses of echinoid

516   diversification prompt a re-evaluation of their fossil record. *eLife,* **11,** e72460.

518 Montagna, M., Tong, K.J., Magoga, G., Strada, L., Tintori, A., Ho, S.Y. & Lo, N. (2019). Recalibration of the

519   insect evolutionary time scale using Monte San Giorgio fossils suggests survival of key lineages

520   through the End-Permian Extinction. *Proceedings of the Royal Society B,* **286**, 20191854.

Morel, B., Kozlov, A.M. & Stamatakis, A. (2018) ParGenes: a tool for massively parallel model selection

and phylogenetic tree inference on thousands of genes. *Bioinformatics,* **35,** 1771-1773.

Near, T.J., Bolnick, D.I. & Wainwright, P.C. (2005). Fossil calibrations and molecular divergence time

estimates in centrarchid fishes (Teleostei: Centrarchidae). *Evolution*, **59**, 1768-1782.

O'Reilly, J.E., dos Reis, M. & Donoghue, P.C. (2015) Dating tips for divergence-time estimation. *TRENDS in*

Genetics, **31,** 637-650.

Paradis, E. & Schliep, K. (2018) ape 5.0: an environment for modern phylogenetics and evolutionary

analyses in R. *Bioinformatics,* **35,** 526-528.

Pedersen, T. (2020) patchwork: the composer of plots. R package version 1.1.1, https://CRAN.R-

project.org/package=patchwork.

Pennell, M.W. & Harmon, L.J. (2013) An integrative view of phylogenetic comparative methods:

connections to population genetics, community ecology, and paleobiology. *Annals of the New*

York Academy of Sciences, **1289,** 90-105.

Philippe, H., Chenuil, A. & Adoutte, A. (1994) Can the Cambrian explosion be inferred through molecular

phylogeny? *Development,* **1994,** 15-25.

Poe, S. & Chubb, A.L. (2004) Birds in a bush: five genes indicate explosive evolution of avian orders.

Evolution, **58,** 404-415.

R Core Team (2022) R: A Language and Environment for Statistical Computing. R Foundation for

Statistical Computing, Vienna, Austria.

Rannala, B., Leache, A., Edwards, S. & Yang, Z. (2020) The multispecies coalescent model and species

tree inference. In C. Scornavacca, F. Delsuc & N. Galtier (Eds.), *Phylogenetics in the genomic era*

(pp. 3:3:1-3:3:21). Self published.

Revell, L.J. (2012) phytools: an R package for phylogenetic comparative biology (and other things).

Methods in Ecology and Evolution, **3,** 217-223.

Robinson, D.F. & Foulds, L.R. (1981) Comparison of phylogenetic trees. *Mathematical Biosciences,* **53,**

131-147.

Ronquist, F., Klopfstein, S., Vilhelmsen, L., Schulmeister, S., Murray, D.L. & Rasnitsyn, A.P. (2012) A total-

evidence approach to dating with fossils, applied to the early radiation of the Hymenoptera.

Systematic Biology, **61,** 973-999.

Sanderson, M.J. & Doyle, J.A. (2001). Sources of error and confidence intervals in estimating the age of

angiosperms from rbcL and 18S rDNA data. *American Journal of Botany*, **88**, 1499-1516.

Sauquet, H., Ramírez-Barahona, S. & Magallón, S. (2022) What is the age of flowering plants? *Journal of*

Experimental Botany, **73**, 3840-3853.

Schliep, K.P. (2011) phangorn: phylogenetic analysis in R. *Bioinformatics,* **27,** 592-593.

Shin, S., Clarke, D.J., Lemmon, A.R., Moriarty Lemmon, E., Aitken, A.L., Haddad, S., Farrell, B.D., Marvaldi,

A.E., Oberprieler, R.G. & McKenna, D.D. (2018) Phylogenomic data yield new and robust insights

into the phylogeny and evolution of weevils. *Molecular Biology and Evolution,* **35,** 823-836.

Smith, M.R. (2022) Robust analysis of phylogenetic tree space. *Systematic Biology,* **71,** 1255-1270.

Smith, S.A., Brown, J.W. & Walker, J.F. (2018) So many genes, so little time: A practical approach to

divergence-time estimation in the genomic era. *PloS one,* **13,** e0197433.

Song, H., Wang, Y., Shao, H., Li, Z., Hu, P., Yap-Chiongco, M.K., Shi, P., Zhang, T., Li, C. & Wang, Y. (2023)

Scaphopoda is the sister taxon to Bivalvia: Evidence of ancient incomplete lineage sorting.

Proceedings of the National Academy of Sciences, **120,** e2302361120.

Strassert, J.F., Irisarri, I., Williams, T.A. & Burki, F. (2021) A molecular timescale for eukaryote evolution

with implications for the origin of red algal-derived plastids. *Nature Communications,* **12,** 1879.

Szánthó, L.L., Lartillot, N., Szöllősi, G.J. & Schrempf, D. (2023). Compositionally constrained sites drive

long-branch attraction. *Systematic Biology*, **72**, 767-780.

Tan, G., Muffato, M., Ledergerber, C., Herrero, J., Goldman, N., Gil, M. & Dessimoz, C. (2015) Current

methods for automated filtering of multiple sequence alignments frequently worsen single-gene

phylogenetic inference. *Systematic Biology, **64,** 778-791.

Tao, Q., Barba-Montoya, J., Huuki, L.A., Durnan, M.K. & Kumar, S. (2020) Relative efficiencies of simple

and complex substitution models in estimating divergence times in phylogenomics. *Molecular*

*Biology and Evolution, **37,** 1819-1831.

Tarver, J.E., Dos Reis, M., Mirarab, S., Moran, R.J., Parker, S., O'Reilly, J.E., King, B.L., O'Connell, M.J.,

Asher, R.J. & Warnow, T. (2016) The interrelationships of placental mammals and the limits of

phylogenetic inference. *Genome Biology and Evolution, **8,** 330-344.

Tavaré, S. (1986) Some probabilistic and statistical problems on the analysis of DNA sequence. *Lecture of*

*Mathematics for Life Science, **17,** 57.

Telford, M.J., Lowe, C.J., Cameron, C.B., Ortega-Martinez, O., Aronowicz, J., Oliveri, P. & Copley, R.R.

(2014) Phylogenomic analysis of echinoderm class relationships supports Asterozoa. *Proceedings*

*of the Royal Society B, **281,** 20140479.

Thorne, J.L., Kishino, H. & Painter, I.S. (1998) Estimating the rate of evolution of the rate of molecular

evolution. *Molecular Biology and Evolution, **15,** 1647-1657.

Wang, L.G., Lam, T.T.Y., Xu, S., Dai, Z., Zhou, L., Feng, T., Guo, P., Dunn, C.W., Jones, B.R., Bradley, T., Zhu,

H., Guan, Y., Jiang, Y. & Yu, G. (2020) *treeio*: an R package for phylogenetic tree input and output

with richly annotated and associated data. *Molecular Biology and Evolution*, **37,** 599-603.

Warnock, R.C. & Wright, A.M. (2020) *Understanding the tripartite approach to Bayesian divergence time*

estimation. Cambridge University Press.

Warnock, R.C., Yang, Z. & Donoghue, P.C. (2011) Exploring uncertainty in the calibration of the molecular

clock. *Biology Letters*, **8,** 156-159.

Whelan, N.V. & Halanych, K.M. (2017). Who let the CAT out of the bag? Accurately dealing with

substitutional heterogeneity in phylogenomic analyses. *Systematic biology*, **66**, 232-255.

Wickham, H. (2016) ggplot2: Elegant Graphics for Data Analysis. Springer-Verlag: New York.

Wickham, H. (2017) tidyverse: Easily install and load 'tidyverse' packages. R package version 1.2.1.

https://CRAN.R-project.org/package=tidyverse.

Wolfe, J.M., Ballou, L., Luque, J., Watson-Zink, V.M., Ahyong, S.T., Barido-Sottani, J., Chan, T.-Y., Chu, K.H.,

Crandall, K.A., Daniels, S.R. Felder, D.L., Mancke, H., Martin, J.W., Ng, P.K.L., Ortega-Hernández, J.,

Palacios Theil, E., Pentcheff, N.D., Robles, R., Thoma, B.P., Tsang, L.M., Wetzer, R., Windsor, A.M.

& Bracken-Grissom, H.D. (2023) Convergent adaptation of true crabs (Decapoda: Brachyura) to a

gradient of terrestrial environments. *Systematic Biology*, syad066, 10.1093/sysbio/syad066.

Wolfe, J.M., Breinholt, J.W., Crandall, K.A., Lemmon, A.R., Lemmon, E.M., Timm, L.E., Siddall, M.E. &

Bracken-Grissom, H.D. (2019) A phylogenomic framework, evolutionary timeline and genomic

resources for comparative studies of decapod crustaceans. *Proceedings of the Royal Society B*,

286, 20190079.

Yang, Z. & Rannala, B. (2006) Bayesian estimation of species divergence times under a molecular clock

using multiple fossil calibrations with soft bounds. *Molecular Biology and Evolution,* **23,** 212-226.

Yendle, P.W. & MacFie, H.J. (1989) Discriminant principal components analysis. *Journal of Chemometrics,*

3, 589-600.

Yu, G., Smith, D.K., Zhu, H., Guan, Y. & Lam, T.T.Y. (2017) ggtree: an R package for visualization and

annotation of phylogenetic trees with their covariates and other associated data. *Methods in*

Ecology and Evolution, **8,** 28-36.

Zhou, Y., Rodrigue, N., Lartillot, N. & Philippe, H. (2007) Evaluation of the models handling heterotachy in

phylogenetic inference. *BMC Evolutionary Biology,* **7,** 206.

Zuckerkandl, E. & Pauling, L. (1965) Evolutionary divergence and convergence in proteins. In V. Bryson &

615   H. J. Vogel (Eds.), *Evolving genes and proteins* (pp. 97-166). Academic Press.

SUPPLEMENTARY MATERIAL - Chronospaces: an R package for the statistical

exploration of divergence times promotes the assessment of methodological

sensitivity

Figure S1: Root-to-tip variance (a proxy for clock-likeness) is strongly determined by the evolutionary

rate of loci. Minimizing this variable, a common approach when subsampling matrices for time

calibration, also results in selecting genes that evolve at minimum rates. To avoid this confounding effect,

clock-like genes from these three matrices were selected after filtering the fastest and slowest evolving

15% of loci. Lines correspond to linear regressions; both axes are log-scaled.

Figure S2: Trace plots of the log-likelihood (logL) of chains. Results correspond to fifteen randomly

selected analyses. Dashed lines correspond to the burn-in threshold imposed (10,000 generations).

Convergence and stationarity vary between conditions and datasets, with mixing being more challenging

when using the site-heterogeneous model CAT + GTR + G, and convergence taking longer for the

eukaryote dataset. Nonetheless, both chains of each analysis seem to have converged to the same high

posterior region of parameter space by the time the burn-in generations have elapsed. Note that the

first few logL values were discarded in order to arrive at a meaningful scaling of the y-axis. The number

of excluded generations was 10 for analyses under the GTR + G model, and either 100 (Curculionoidea,

Decapoda) or 1,000 (Eukaryota) for analyses under the CAT + GTR + G model. Surprisingly, runs under

GTR + G converged within the first 50 generations (sometimes, as is the case of the second decapod plot,

within less than 10).

Figure S3: Median node ages for pairs of chains showing appropriate levels of convergence. Results

correspond to fifteen randomly selected analyses. Grey lines show 1-to-1 lines. The minimum Pearson

correlation coefficient (R) found across all analyses was 0.973.

645 **Figure S4:** Comparison of the posterior estimates of evolutionary rates across a random sample of 20

646 analyses per dataset (i.e., 60 analyses total, 2 chains each). Pairs of chains are shown next to each other.

647 Estimates correspond to posterior means ± 1 standard deviation.

Figure S5: Comparison of the posterior estimates of p1, one of the two parameters of the birth-death

650   tree prior, across a random sample of 20 analyses per dataset (i.e., 60 analyses total, 2 chains each). Pairs

651   of chains are shown next to each other. Estimates correspond to posterior means ± 1 standard deviation.

652   p1 is corresponds to the birth rate (λ) minus the death rate (μ), and is thus the net diversification rate.

653   Data is for the same analyses shown in Fig. S4.

Figure S6: Comparison of the posterior estimates of p2, one of the two parameters of the birth-death

tree prior, across a random sample of 20 analyses per dataset (i.e., 60 analyses total, 2 chains each). Pairs

of chains are shown next to each other. Estimates correspond to posterior means ± 1 standard deviation.

p2 is corresponds to the birth rate (λ) multiplied by the sampling fraction (ρ). Data is for the same anal-

yses shown in Fig. S4.

Figure S7: Posterior distributions of ages for the most sensitive nodes of the curculionid dataset. For each methodological choice being varied, the top five most sensitive nodes are depicted. Factors are ordered from the least impactful on the left, to the most impactful on the right (according to the proportions of explained variance, see Fig. 3). Nodes are identified using two randomly-selected terminals from each descendant clade.

Figure S8: Posterior distributions of ages for the most sensitive nodes of the decapod dataset. For each

methodological choice being varied, the top five most sensitive nodes are depicted. Factors are ordered

from the least impactful on the left, to the most impactful on the right (according to the proportions of

explained variance, see Fig. 3). Nodes are identified using two randomly-selected terminals from each

descendant clade.

Figure S9: Posterior distributions of ages for the most sensitive nodes of the eukaryote dataset. For each methodological choice being varied, the top five most sensitive nodes are depicted. Factors are ordered from the least impactful on the left, to the most impactful on the right (according to the proportions of explained variance, see Fig. 3). Nodes are identified using two randomly-selected terminals from each descendant clade.

Figure S10: Branch length warping along the bgPCA maximizing the discrimination of chronograms

inferred using different node age prior distributions (i.e., the least impactful decision across all datasets;

see Fig. 3). This decision seems to induce heterogeneous effects across subclades.

Figure S11: Branch length warping along the bgPCA maximizing the discrimination of chronograms

inferred using different models of molecular evolution (i.e., the second least impactful decision across all

datasets; see Fig. 3). This decision seems to induce heterogeneous effects across subclades.

Figure S12: Branch length warping along the bgPCA maximizing the discrimination of chronograms

inferred using different clock models (i.e., the first or second most impactful decision across all datasets;

see Fig. 3). This decision seems to induce effects that are strong, yet clade specific.