7Title: Comparison of six methods for stabilizing metapopulation dynamics and for their
robustness against census noise

Authors: Akanksha Singh[1,2], Sudipta Tung[2*]

Affiliations:

1. Indian Institute of Science Education and Research, Mohali, Punjab, India -140306
2. Integrated Genetics and Evolution Laboratory (IGEL), Department of Biology,
Ashoka University, Sonipat, Haryana, India, 131029.

1. akankshasingh8730@gmail.com
2. sudipta.tung@ashoka.edu.in

1. ORCID: 0000-0003-3403-8368
2. ORCID: 0000-0002-3201-3303

Running title: Comparing methods to stabilize metapopulation dynamics

* **Name and address of the corresponding author:**

Sudipta Tung

DBT/Wellcome Trust India Alliance Early Career Fellow

Department of Biology, Ashoka University

Plot No. 2, Rajiv Gandhi Education City,

Post Office Rai, Near Rai Police Station,

Sonipat, Rai, Haryana, India 131029

20
21 Tel: +91 9049131125

22 Email: sudipta.tung@ashoka.edu.in

Abstract

Natural populations face growing extinction risks due to mismatches between traits and ecological conditions, exacerbated by habitat degradation and climate change. While several population control methods have been proposed to protect vulnerable populations, their effectiveness in spatially structured populations (metapopulations) and resilience to census inaccuracies remain unexplored. This study uses simulations to evaluate six control methods, originally designed to stabilize isolated populations, in metapopulations. We compared these methods across twelve ecological conditions based on three stability metrics and implementation costs. Without interventions, the twelve metapopulation scenarios exhibited distinct dynamics and extinction susceptibilities. Comparing the control methods against these baseline dynamics, we found that Adaptive Limiter Control consistently reduced extinction probabilities, while Lower Limiter Control effectively minimized population fluctuations in most settings, followed by Adaptive and Both Limiter Controls. These performance rankings remained robust despite simulated census errors. Our findings provide actionable insights for conservation policies aimed at stabilizing vulnerable metapopulations and offer a general framework for studying metapopulation dynamics using external interventions.

1. Introduction

Complex temporal fluctuations in population size are a ubiquitous phenomenon in nature (Lundberg et al. 2000; Clark and Luis 2020). While external environmental factors are known to induce such fluctuations (Borowsky 1971), it's the intricate interplay between the life-history of resident populations and local environmental conditions that ultimately shapes the temporal dynamics of populations (Gamelon et al. 2017; Tung et al. 2019). Over time, some populations may exhibit diminishing magnitudes of fluctuation, harmonizing their life-histories with environmental cues, but not all populations demonstrate such resilience. If these fluctuations intensify or sustain at elevated levels, populations can undergo frequent declines, amplifying their risk of extinction (Escudero et al. 2004; Smith and Meerson 2016). Hence, devising strategies to stabilize such vulnerable populations has been at the forefront of research for ecologists, conservation biologists, and biological population managers for many years.

In the past three decades, theoretical propositions have introduced several methods to stabilize such inherently extinction-prone populations (McCallum 1992; Corron et al. 2000; Hilker and Westerhoff 2005; Dattani et al. 2011; Sah et al. 2013; Tung et al. 2014). These methods, commonly referred to as population control or stability methods, primarily operate by modulating population size — either by externally adding individuals or strategically removing them based on current census size. Their practical applicability is a key advantage, allowing deployment in actual biological populations without exhaustive knowledge of the underlying dynamics or system parameters, which is in contrast to the chaos-control methods proposed to stabilize chaotic non-linear dynamics by manipulating system parameters (Andrievskii and Fradkov 2003; Fradkov and Evans 2005). Despite these advantages, the widespread adoption of population control methods in conservation has been limited. One major factor for this reluctance stems from the observation that, with a couple of exceptions

(Dey and Joshi 2013; Sah et al. 2013), most methods have been explored solely in case of

single, isolated populations (Corron et al. 2000; Hilker and Westerho 2005; Dattani et al.

2011; Tung et al. 2014).

However, in natural settings, populations are seldom isolated. Organisms of a species

typically do not occupy a shared space or pool resources uniformly. Instead, more commonly,

groups of individuals of a species inhabit distinct, spatially dispersed habitats. Should these

habitats fall within an organism's range of movement, migration or dispersal (used

interchangeably here) becomes commonplace. These networks of spatially distributed

populations, interconnected by such migrations, are termed spatially structured populations or

'metapopulations'. Insights from studying population control methods in single isolated

populations prove inadequate for understanding metapopulation dynamics (Dey and Joshi

2007). This is primarily because, migration between subpopulations plays a major role in

shaping the dynamics and stability of metapopulations (Gyllenberg et al. 1993; Stacey et al.

1997; Hanski 1998; Dey and Joshi 2006).

Migration modalities, driven by factors such as food foraging, mate-seeking, or habitat

preference, and shaped by the connectivity and migratory tendencies of local populations,

play a crucial role in population dynamics. From a population dynamic perspective,

emigration can alleviate resource competition within a subpopulation, while immigration can

exacerbate it, increasing vulnerability to extinction. On the other hand, immigration can

rejuvenate a locally extinct population, eliminating the need for external intervention.

Therefore, the migration intricately influences local dynamics and, by extension, overall

metapopulation fluctuations (Quévreux et al. 2023; Suzuki and Economo 2024). Prior

research has shown that both low and high migration rates can differentially modulate

metapopulation stability, even under uniform environmental conditions across subpopulations

(Dey and Joshi 2006). Thus, extrapolating findings from spatially unstructured populations to

metapopulations becomes inadequate.

Our study introduces a simulation framework to explore performance of six population

control methods (mathematical formulations in Table 1) in metapopulations consisting of two

subpopulations connected via migration with varying characteristics. By introducing

combinations of intrinsic growth rate and carrying capacity parameters, we construct diverse

ecological scenarios (exact descriptions in Figure 1) to evaluate the efficacy of the population

control. For this comparison, we used an established statistic: composite scores (following

(Tung et al. 2014)) that combines measures of various aspects of population stability with the

costs associated with the implementation of the population control methods. Furthermore, we

assessed the relative performance of these population stability strategies in the presence of

three distinct census inaccuracies: unbiased noise or white noise, overestimation, and

underestimation—a pertinent aspect that has often been overlooked (Supplementary Figures

S8-S13).

2. Materials and Methods

2.1. Spatial structuring and population growth model

We adopted a metapopulation framework comprising two subpopulations interconnected

through migration. For our study, all metapopulations were treated as homogeneous, ensuring

identical environmental conditions across both subpopulations. Various migration rates and

environmental characteristics differentiated our treatment groups (Figure 1C). We used a

popular population growth model, the Ricker map (Ricker 1954), to model the dynamics of

each subpopulation. Mathematically this map is given as $N_{t+1} = N_t \exp[r(1-N_t/K)]$; where r, K

and N_t denote intrinsic growth rate, carrying capacity and population size at time t,

respectively. The popularity of this model stems from its intuitive formulation (May and

Oster 1976), and the fact that it can be derived from the first principles as long as the

individuals of the concerned population are distributed randomly over space and undergoing

scramble competition (Brännström and Sumpter 2005). Moreover, Ricker map has been

implemented to successfully capture empirical dynamics of species across diverse taxa,

including bacteria (Ponciano et al. 2005), fungi (Ives et al. 2004), ciliates (Fryxell et al.

2005), insects (Sheeba and Joshi 1998; Dey and Joshi 2006) and fishes (Ricker 1954; Denney

et al. 2002). Consequently, simulation outcomes using this model are expected to be broadly

generalizable.

2.2. Migration rates and environmental regimes

Migration rate is known to majorly influence metapopulation dynamics (Allen et al. 1993;

Gyllenberg et al. 1993; Hastings 1993; Ruxton 1997; Ranta et al. 1998). In order to capture a

diverse metapopulation dynamic scenarios in this comparative analysis, we have considered

metapopulations with bidirectionally symmetric migration with high ($m_1 = m_2 = 0.3$) and low

($m_1 = m_2 = 0.1$) level of migration rate (Dey and Joshi 2006), and also metapopulations with a

bidirectional asymmetric migration rate ($m_1 = 0.1$, $m_2 = 0.3$). We further incorporated different

values of the demographic parameters – intrinsic growth rate (r) and carrying capacity (K) to

capture contrasting environmental regimes of the subpopulations for each of the three types

of migration treatments. Since we aim to look at the *stabilising* effect of the different

population control methods, we deliberately chose two values of intrinsic growth rate r (low

$=2.8$, high $=4$) and carrying capacity K (low $=60$, high $=300$) that lead to extinctions and large

amplitude oscillations in single populations, following a previous such comparative effort for

spatially-unstructured populations (Tung et al. 2014). With this set of parameter values, we

capture diverse metapopulation phenomena such as local extinctions, and genetic bottlenecks.

Thus, we have 12 treatment regimes with various combinations of three migration modalities,

two intrinsic growth rates and two carrying capacity values (Figure 1).

2.3. Replications, initial conditions and reset

In order to obtain a generalizable output, we considered 70 replicates for each treatment

group, and thus all demographic or population dynamic matrices in the figures were

presented as average ($\pm$SEM) over all these replicates. At the beginning, time-series for all

replicates in all the treatment groups were initiated with 20 individuals within each

subpopulation. Whenever population size of the entire metapopulation became zero, we reset

population size of each of the subpopulations to eight (following (Dey and Joshi 2006)).

2.4. Biologically realistic assumptions

We incorporated a number of biologically realistic features in our simulations. Firstly, in

order to account for the fact that organisms come in whole numbers, we rounded off all

outputs of the Ricker model, the number of migrants and the number of individuals to be

introduced to or removed from the system as prescribed by the control methods, to their

nearest integer values as proposed by earlier work (Henson et al. 2001; Domokos and

Scheuring 2004). Secondly, empirically obtained timeseries are typically short and it is rarely

possible to exclude transients from them prior to analysis since natural populations are

situated in environments that are constantly undergoing change (Hastings 2004). In order to

keep a parity with that, we decided to include the transients in our study by restricting our

analysis for the first 50 generations of population growth time-series, instead of studying the

steady states of a system. Thirdly, stochastic noise is ubiquitous in any quantitative aspect of

biology and it is known that disregarding this noise may have significant impact on, *inter*

alia, the outcome of theoretical studies involving stabilizing populations through external

perturbations (Dey and Joshi 2007). To reflect this, we introduced noise in the growth rate

and carrying capacity values for each subpopulation at each iteration. A number was picked

using a uniform random number generator from U(-0.2, 0.2), and added to the regime-

specific value of r as mentioned in Figure 1. Similarly, the value of carrying capacity was

picked using the uniform random number generator from U(0.9×K, 1.1×K), where K is the

regime-specific value of carrying capacity. The values of the noise are at par with the

previous studies on extinction prone populations (Tung et al. 2014). Fourthly, whenever

population size becomes very low, the risk of extinction in the next generation increases

significantly due to various reasons including stochastic death of the breeding individuals, or

all breeding individuals being of same sex. We implemented this possibility of stochastic

extinction by assuming that there is 50% chance of extinction if subpopulation size goes

below 4 (following (Dey and Joshi 2006; Tung et al. 2014)).

2.5. Measures of stability and synchrony in unperturbed populations

Prior to comparing the control methods, we first studied the dynamics of the unperturbed

populations of all the twelve metapopulation regimes. For this, we measured two aspects of

demographic stability for each metapopulation in our analysis - Constancy and Persistence

Stability (Grimm and Wissel 1997). Constancy, as the name indicates, refers to how

unchanging or constant the size of a population is. In other words, the less the size of a

population fluctuates, the more stable it is in terms of constancy. We measured constancy

stability using Fluctuation Index (henceforth, FI, (Dey and Joshi 2006)), which is formulated

as $\sum |N_{t+1}-N_t| / (N \times T)$, where N_t is the population size after perturbation (if applicable) at

time t, $\bar{N}$ is the average population size over T generations. Thus, the lower the FI is, the

higher is the constancy stability.

In contrast, persistence can be said to be the resistance of a population to extinction or the

converse of the propensity of a population to go extinct. This is quantified by computing the

extinction probability (henceforth, EP) of the population as E/T, where E is the number of

extinction events in T number of generations.

Additionally, we measured genetic stability of the metapopulations by computing the

effective population size (EPS). This is computed as the harmonic mean (Allendorf et al.

2012) of post-perturbation (if applicable) population size, *i.e.* mathematically, EPS =

$T/\sum (1/N_t)$, where T is the length of the timeseries and N_t is breeding population size at the t^{th}

generation.

Synchrony between the subpopulations of a metapopulation is quantified by computing the

cross-correlation coefficient at lag zero of the first-differenced time series of log-transformed

values of the two subpopulation sizes (Bjørnstad et al. 1999; Dey and Joshi 2006).

2.6. Population control methods

For this comparative analysis, we considered six population control methods - Constant

Pinning (CP, (McCallum 1992; Parthasarathy and Sinha 1995; Solé et al. 1999)), Lower

Limit Control (LLC (Hilker and Westerho 2005)), Adaptive Limiter control (ALC, (Sah et

al. 2013)), Upper Limit Control (ULC, (Hilker and Westerho 2005)), Target Oriented

Control (TOC, (Dattani et al. 2011)), and Both Limit Control (BLC, (Tung et al. 2014)). A

description of each of these methods along with corresponding control parameter values can

be found in Table 1. These methods were chosen because they are implementable in real

biological settings (Gusset et al. 2009), well studied theoretically (McCallum 1992; Solé et al.

1999; Hilker and Westerho□ 2005; Dattani et al. 2011; Tung et al. 2014), and have been

validated empirically (Dey and Joshi 2007, 2013; Sah et al. 2013; Tung et al. 2016a,b).

Additionally, efficiency of these six methods has been compared in spatially-unstructured

populations (Tung et al. 2014), which can be contrasted qualitatively with the results we

obtain in the context of spatially-structured populations in this study. Outcomes of these two

studies together will provide a wholistic picture of stabilizing population dynamics. A more

detailed description of these methods can be found elsewhere (Tung et al. 2014).

2.7.The comparative framework

After implementing the methods on our spatially-structured simulation framework, we first

tested the performance of the methods by measuring two aspects of population stability –

constancy and persistence, over a range of control parameters (ranges can be found in Table

1). Promisingly, we observe that any level of constancy and persistence stability can be

achieved in case of two-patch metapopulations by varying the control parameters of *all* six

population control methods, similar to spatially-unstructured populations (Tung et al. 2014).

However, in order to reach a specific level of stability in one aspect of stability, the methods

varied substantially in their performance for the other aspects of stability and/or costs

incurred in implementing this level of stability.

So, in order to compare the effectiveness of these methods at a common level, we decided to

look at the performance of the methods in order to achieve 50% reduction of fluctuation

index (*i.e.,* improvement of constancy stability) and 50% reduction of extinction probability

(*i.e.,* improvement of persistence stability), separately. For each of these two scenarios, we

compared the methods using an established composite performance score (Tung et al. 2014)

and a novel robustness index of the performance of the methods against noise in census

count.

2.8. Composite Performance Score

Composite performance score (CPS) or composite index (following (Tung et al. 2014)), gives

equal weightage to the effort magnitude and effective population size and the other stability

value (*i.e.,* extinction probability when analysing for 50% reduction of fluctuation index, and

vice versa). Here, effort magnitude (EM) can be translated as the 'cost' of applying a certain

control method to the metapopulation and computed as $\sum |a_t\text{-}b_t| / (N\square \times T)$, where, b_t and a_t

denote population size before and after applying the control method, and $N\square$ denotes the

average population size over T generations.

The scales of the components of CPS, or "component indices" (CI), are different. This would

bias our comparison by giving more weightage to CIs with values a higher scale. To alleviate

this bias, the value for each of the CIs for each method was divided by the highest of that

index amongst all methods. This way, the scaled value of all the CIs remained between 0 to 1.

It is also noted that although a lower value of $CI_{FI/EP}$ (component index for FI or EP) and

CI_{EM} (component index for EM) are desirable for better performance of a method, for CI_{EPS}

(component index of effective population size), a *higher value* will denote better

performance. Thus, we computed composite performance index as, $CI_{FI/EP} + CI_{EM} + (1\text{-}$

$CI_{EPS})$, and overall, the lower the composite score for a method is, the better it works in

stabilising the regime in question.

For Lr populations, we found that the metapopulations were extremely persistent stable, i.e.,

they had negligibly low EP even without any external perturbations (refer to Section 3.1).

Therefore, we felt that including CI_{EP} in the calculations for composite performance score for

50% reduction in FI in Lr regimes would unnecessarily bias the results by reflecting a

method's ability of solving a problem (i.e., extinction) that was not there. Therefore, we

decided to not include CI_{EP} in the calculation of composite performance score for reducing FI

by 50% in Lr regimes.

2.9. Robustness Analysis

A notable challenge with external perturbation-based control methods is the need for precise

census counts in each generation when introducing or removing individuals, in all but the

Constant Pinning (CP) method. In natural settings, obtaining an exact animal count is nearly

impossible. Typically, only an estimate of the census size is achieved through methods like

transect walks, acoustic monitoring (Law et al. 2024), multiyear and multi-season

occupancies (Goldingay et al. 2022), bayesian occupancy models (Rodhouse et al. 2019) -

which can lead to overestimations or underestimations of the actual population size per

generation. The impact of such census errors on the efficacy of control methods remains

unexplored. To address this, we conducted a CPS-based comparison incorporating varying

levels of white noise or unbiased error in census values. We simulated random

overestimations or underestimations of the population size in each generation, based on the

level of noise. The perceived population size used for implementing control methods was

therefore *actual population size + Uniform()*, where *Uniform()* means that the error values

were picked from a uniform distribution. Subsequently, we calculated the "robustness index"

to evaluate the impact of the error. This index was computed as the *mean of*

(composite_score_with_error – composite_score_without_error)^2 for each noise level,

serving as a measure of dispersion around the no-error composite performance score. In

addition, factors like weather conditions, landscape characteristics, and vegetation density

can influence the accuracy of population estimates with specific biases. To capture this, we

considered two types of errors: positive noise (overestimating population) and negative noise

(underestimating population). For overestimations, error values were positive, while for

underestimations, they were negative. In the case of white noise, error values were randomly

drawn from a uniform distribution within the range [−noise level×actual population size,

+noise level×actual population size]. For each error type, noise level ranged from 0 to 0.5 in

increments of 0.05, enabling us to track the composite index's trajectory against error rates.

Although component indices were derived from the averages of 70 replicates, minor

differences in composite scores emerged in repeated runs of the same code. To address this,

we computed the composite scores 20 times, using the average of these scores to represent

the composite index in our graphical representation. The results for these can be found in

Figures 5-6, and Supplementary Figures S8 – S13.

3. Results and Discussion

3.1 Dynamics of the unperturbed regimes

First, we evaluated the dynamics of unperturbed metapopulation regimes, i.e., without subjecting the metapopulation to any population control methods, to establish a baseline for subsequently comparing the efficacy of various population stability strategies. For this purpose, we estimated two aspects of demographic stability – constancy and persistence – as well as genetic stability by computing effective population size for all 12 regimes (Figure 2). Apart from serving as a reference point for the effects of the different population control methods, this analysis reveals how these stability aspects are influenced by the complex interplay of intrinsic growth rate, carrying capacity of the constituent subpopulations, and migration rate between the subpopulations.

Fluctuation index and extinction probability values, as measures of inverse of constancy and persistence stability respectively of the unperturbed metapopulations, clearly suggested that the regimes with high intrinsic growth rate and low carrying capacity (i.e. HrLk regimes) exhibited the most unstable metapopulation dynamics (Figure 2a-2d). The instability in these dynamics stems from overcrowding due to high intrinsic growth rates, further aggravated by limited resource availability due to the low carrying capacities of these regimes. This leads to overutilization of resources, resulting in characteristic 'boom-bust dynamics' with significant population fluctuations and a heightened risk of extinction (Grimm and Wissel 2004; Tung et al. 2014; also see Figure S1). These regimes experienced severe population bottlenecks nearly every alternate generation (Figure S1), resulting in relatively poor genetic stability, as indicated by low effective population sizes (Figure 2e-1f). Interestingly, we also found that this pattern remained consistent regardless of the level of migration and the nature of

migration (i.e.,symmetric or asymmetric; comparing Figure 2a and Figure 2b; comparing

Figure 2c and Figure 2d).

Regimes characterized by high intrinsic growth rates and high carrying capacities, termed

HrHk regimes, emerged as the subsequent most unstable scenarios, as indicated by their high

fluctuation indexes, implying low constancy stability (Figure 2a, 2b). Notably, though these

regimes exhibited 'boom-bust dynamics' (Figure S1), their high carrying capacities resulted in

a significantly reduced frequency of population size reaching zero, as evidenced by their

relatively low extinction probabilities (Figure 2a, 2b; also see (Tung et al. 2014)). Despite

reaching higher population sizes due to their ample carrying capacities, these regimes did not

necessarily see an increase in effective population size (Figure 2e, 2f). This is attributed to

the substantial fluctuations in population size and that the calculation of effective population

size is more adversely impacted by smaller population values (Allendorf et al. 2012).

An intriguing characteristic of HrHk regimes with symmetric migration, is the observed

difference in metapopulation constancy stability between low and high migration levels

(Figure 2a). Metapopulations with low migration level exhibited substantially lower

fluctuation index i.e. greater constancy stability than those with high migration, although

constancy of the constituent subpopulations was comparable. This apparent discrepancy is

resolved when we look into the level of synchrony between the constituent subpopulations

(Figure 2g). We found that synchrony level was much lower in metapopulations with low

migration levels compared to metapopulations with high migration level (for example,

HrHkLm vs HrHkHm). This result capturing out-of-phase dynamics of coupled unstable

populations has been seen both theoretically (Gyllenberg et al. 1993; Doebeli 1995;

Amarasekare 1998; Kendall and Fox 1998; Ylikarjula et al. 2000; Briggs and Hoopes 2004;

Dey and Joshi 2006; Abbott 2011; Dey et al. 2014) and empirically (Lecomte et al. 2004;

Dey and Joshi 2006; Sah et al. 2013b; Mueller and Joshi 2020).

Similarly, in metapopulations with low intrinsic growth rates, we observed enhanced

constancy stability, which correlates with a lower synchrony level between subpopulation

dynamics, especially in scenarios of low symmetric migration (as seen in the comparison of

fluctuation index and synchrony plots for LrLkLm vs LrLkHm, and LrHkLm vs LrHkHm,

Figure 2a, 2g). This is in line with the observations in larger metacommunity networks

(Suzuki and Economo 2024) and fluctuating environments (Quévreux et al. 2023).

Comparing both constancy and persistence stability, metapopulations with low intrinsic

growth rate (Lr regimes) were found to be more stable demographically compared to the

regimes with high intrinsic growth rate (Hr regimes). This contrast is more prominent in

terms of persistent stability, as irrespective of the level and nature of migration,

metapopulations with Lr regimes rarely incurred extinction (Figure 2c-2d). However, it is

noteworthy that despite enhanced demographic stability, genetic stability of LrLk regimes is

not high. This is primarily because these regimes tend to maintain low population sizes due to

their lower carrying capacities and growth rates, making them more vulnerable to inbreeding

depression and loss of genetic diversity.

Following the same argument, LrHk regimes exhibited more stable dynamics with

significantly higher average population sizes, leading to considerably larger effective

population sizes (Figure 2e-2f). This divergence highlights the complex relationship between

demographic and genetic stability in metapopulations, influenced by intrinsic growth rates

and carrying capacities. We find that the regimes with high extinction probabilities were

always coupled with low genetic diversity (although not vice versa). This has also been seen

in empirical data (DiLeo et al. 2024).

Additionally, our study showed that the stability properties of metapopulations with

asymmetric migration lie between those of the two symmetric cases (Figure 2). This result is

in contrast to the notion that metapopulations with asymmetric migration rate are more stable

(Doebeli 1995; Ylikarjula et al. 2000). Instead, it aligns with more research suggesting that

the relative stability of metapopulations with asymmetric migration is contingent on specific

contextual factors (Dey et al. 2014).

Taken together, the 12 regimes considered in this study, each with their unique combinations

of demographic and genetic stability attributes, provide a robust framework for analysing the

effect of population control methods across a range of scenarios. This approach allows us to

compare the results across diverse scenarios with a consistent framework, thereby facilitating

the derivation of broader conclusions about population stability strategies in multiple

ecological contexts.

3.2 Comparing the control methods for inducing desired constancy stability

To assess the relative performance of the control methods on a common platform, we first

computed composite performance score (following (Tung et al. 2014)) for each of the

methods in each regime separately. This score integrated other stability measures, such as

extinction probability and effective population size, along with a metric for implementation

effort required to diminish population size fluctuation to 50% of the unperturbed scenario,

thereby improving constancy stability. As the extinction probabilities for Lr populations were

found to be negligible (Figure 2c-1d), we excluded the extinction probability component

from the calculation of composite scores in Lr regimes. This approach was adopted based on

previous findings that, while all methods can independently promote various stability aspects,

they often entail trade-offs in terms of other stability facets or require significant

implementation efforts (Tung et al. 2014, 2016a,b). For a comprehensive assessment, we

calculated these composite scores for each method across all 12 regimes. In this scoring

system, a lower composite score indicates a more favourable performance, reflecting a

method's efficacy in enhancing stability with minimal trade-offs and effort.

Our findings indicate that the effectiveness of control methods in metapopulations is context-

dependent, with no single method proving universally optimal across all regimes (Figure 3).

This aligns with previous observations in spatially unstructured populations (Tung et al.

2014). However, interestingly, our comparative analysis revealed a common trend for the

regimes with low intrinsic growth rate (i.e. Lr regimes). In these regimes, lower limiter

control (LLC), adaptive limiter control (ALC) and both limiter control (BLC) demonstrated

similar composite scores and consistently outperformed other methods (Figure 3a-3f). These

are closely followed by target-oriented control (TOC) and upper limiter control (ULC), in the

order of performance. Constant pinning (CP) method performed consistently worst in all six

scenarios. The reason behind this trend became clear when we checked the individual

components – effective population size and effort magnitude that constituted the composite

score in these scenarios (Figure S2-S4). LLC, ALC, and BLC achieved effective population

sizes that were comparable and only lower than those induced by the CP method. Although

CP had the potential to maximize effective population size across the regimes, its high

implementation costs, as evidenced by substantial effort magnitude, made it the least

favourable option. In contrast, LLC, ALC, and BLC struck a balance, attaining moderate

effective population sizes with reasonable level of implementation efforts to become the best

performing methods in these regimes.

In HrLk regimes, LLC stood out as the superior method compared to others (Figure 3g-3i). It

paralleled ALC in enhancing effective population size. However, LLC surpassed ALC in

reducing implementation effort and extinction probability, establishing itself as the most

effective method in these scenarios. While ULC and BLC proved more efficient in mitigating

extinction risks, their higher implementation costs and lower effective population sizes

negatively impacted their overall rankings according to the composite scores. On the other

hand, in HrHk regimes, BLC demonstrated superior efficacy, successfully eliminating

extinction risks while achieving an optimal balance between effort and effective population

size (Figure 3j-3l).

3.3 Comparing the control methods for inducing desired persistence stability

In our subsequent analysis, we focused on the performance of control methods in reducing

the extinction probability to 50% of the corresponding unperturbed scenario. Since Lr

regimes rarely experience extinction, this analysis was restricted to Hr regimes. Here, the

composite performance score was calculated from the fluctuation index, effective population

size, and the magnitude of implementation effort required for reducing the extinction

probability by 50% (following (Tung et al. 2014)).

We found that ALC was the most effective method across all Hr regimes (see Figure 4),

closely followed by LLC, BLC, and CP. In contrast, ULC and TOC were less effective in

addressing extinction risk. It appears that methods designed to prevent population size from

dropping below a certain threshold – whether constant (as in CP, LLC and BLC) or variable

(as in ALC) – are more successful in enhancing a population's resistance to extinction. In

contrast, the methods that involve culling individuals to counter overpopulation, i.e., TOC

and ULC, do not seem to make the metapopulation more persistent.

Interestingly, although we found that ALC performed the best in terms of inducing desired

persistence stability in all the regimes, analysis of the components of composite score

revealed that it becomes the best through different routes (Figure S5-S7). In case of the most

unstable HrLk regimes, when migration rate between the subpopulations was low (i.e. for

HrLkLm regime), CP induces the maximum effective population size followed by LLC, ALC

and BLC. But CP does that at the cost of a large effort magnitude leading to its poor

performance rank based on composite score. Among LLC, ALC and BLC, ALC performed

the best with respect to curbing population size fluctuation at the cost of minimum effort

magnitude, leading it to the best method to stabilize metapopulations in this regime. In

scenarios with high migration rate between the subpopulations i.e. in HrLkHm regime, effort

magnitude for LLC was minimum but this led to very low effective population size, thereby,

compromising its performance score. Other trends were similar, and overall ALC becomes

the best performing method. When migration between the subpopulations was asymmetric

i.e. in HrLkLmHm regime, effort magnitude incurred by ALC to reach the desired level of

persistence stability was actually the maximum, but as this method was excellent in

simultaneously improving constancy stability and genetic stability, it became the best

performing method with the lowest composite score. When carrying capacity was high, ALC

reduced population size fluctuations and increased the effective population size to the

maximum extent, incurring only moderate effort magnitude. While LLC and BLC also

performed well, they fell short in terms of one or more components of the composite

performance score.

To offer a comprehensive overview of the efficacy of six population control methods in

achieving target levels of constancy and persistence stability, we have compiled a summary

table that aggregates the rankings for each method across all evaluated regimes (Table S1).

Our consolidated findings indicate that the Lower Limiter Control (LLC) method delivers

consistent results in attaining the desired level of constancy stability, while the Adaptive

Limiter Control (ALC) method stands out as the most effective for achieving the desired level

of persistence stability. These results provide actionable insights for population ecologists

and conservation biology practitioners, without going into the details of the specific regime

conditions, aiding in the formulation of intervention strategies in case of spatially-structured

populations.

3.4 Comparing robustness of the control methods against census noise

To evaluate the robustness of population control methods against variations in census accuracy, we compared the composite performance scores of these methods across a range of white noise levels in population census. This approach addresses a significant concern in implementing external perturbation methods: the need for precise census counts of the target species, which may not always be easily available. We specifically analysed the impact of census noise on achieving the above-mentioned desired levels of constancy and persistence stability.

In scenarios focused on enhancing constancy stability, particularly within low intrinsic growth rate regimes (i.e., Lr regimes), the performance of most methods was comparable and robust, with the exception of ULC (Figure 5; Supplementary Figure S8). Combining their overall performance with resistance to census noise, LLC, ALC, and BLC emerged as the most promising methods in these regimes. Whereas, in HrLk regimes, LLC, ALC, and TOC demonstrated greater robustness to census noise. Consequently, LLC stood out as the most promising method in these settings. For HrHk regimes, LLC, TOC, BLC when migration rate is low between the subpopulations, ALC, TOC when migration rate is high between the subpopulations and LLC, ALC, TOC and BLC when migration rate is asymmetric between the subpopulations performed the better against census noise. Thus, combining performance of the methods and their robustness against census noise, BLC stood out to be promising in these regimes, although one needs to note that it is less robust to census noise when migration rate is high between the subpopulations.

When focusing on inducing persistence stability to a desired level, ULC performed poorly for HrLk regimes. Other methods performed similar as in without noise. In case of asymmetric migration between the subpopulations ALC performed the best, closely followed by TOC

(Figure 6; Supplementary Figure S9). Whereas, all methods performed well in HrHk regimes,

ALC was consistently good. Thus, considering both performance and robustness to census

noise, ALC was identified as the most effective method in these regimes.

In order to present a comprehensive evaluation of the resilience of the six control methods to

census inaccuracy in presence of white noise, we aggregated the rank of robustness indices of

these methods for inducing both constancy and persistence stability into a summarizing table

(Table S2). This analysis offers a broad view of each method's resilience to noise-induced

variability, independent of specific ecological regime parameters. When focusing on attaining

a target level of constancy stability, our results indicate a notable parity in performance

among all methods except for the Upper Limiter Control (ULC), which exhibits subpar

robustness. Conversely, in the pursuit of persistence stability, the Target-Oriented Control

(TOC), Adaptive Limiter Control (ALC), Lower Limiter Control (LLC), and Constant

Pinning (CP) methods show marginally superior resilience, in descending order of

effectiveness.

Performance of ALC was found to be rather resilient in the presence of a range of intensities

of positive (i.e. overestimation; Supplementary Figures S10-S11) and negative (i.e.

underestimation; Supplementary Figures S12-S13) noise to census. Composite score of ULC

methods was found to be the most variable across ranges of noise intensities. Notably, the CP

method, which is theoretically independent of population census data, was expected to excel

irrespective of census noise levels. However, contrary to our expectations, CP did not surpass

other methods in terms of composite score performance even in the presence of census noise.

3.5 Limitations and Future Scope

This study aims to prescribe population control methods to stabilize metapopulations, where

subpopulation dynamics are approximated using the two-parameter Ricker model. The

broader objective is to establish a general framework for studying ways to stabilise

metapopulations using population control strategies to implement conservation strategies

better. The proposed framework involves defining the metapopulation structure (e.g., two

subpopulations connected via bidirectional migration), describing their individual dynamics

(Ricker dynamics followed by migration), and applying different population control methods

to achieve a specific stability criterion (e.g., a 50% decrease in FI or EP). We then compare

various aspects of the controlled metapopulation (e.g., other stability measures,

implementation cost (EM), and genetic diversity (EPS)) using a composite performance score

(CPS). The CPS provides a guideline for selecting population control methods based on

different parameter values without census error. Additionally, we explore how CPSs for

control methods vary across different levels of census error.

This framework is customizable at each step based on the specific nature of metapopulation

and stability aspects of interest. For instance, if a different kind of growth model, such as

Beverton-Holt model (Beverton and Holt 2012), better describes a system, one can substitute

the subpopulation dynamics equations and evaluate control methods using composite scores.

The composite score definition can also be adjusted to fit the specific needs of the population

management. In this study, we did not include census effort as an explicit cost. However, this

cost can vary based on the species and population control strategy: CP requires no census

effort, LLC and ALC require minimal effort, while ULC, TOC, and BLC need sufficient

census to ensure population size is above a certain limit. We arbitrarily assigned equal weight

to stability (EP/FI), EM, and EPS for each control method, but this may not be useful in all

scenarios. For example, conserving a species might prioritize genetic diversity (EPS), or the

cost of adding/subtracting individuals (EM) might be critical. The weights of each composite

index in the CPS can be adjusted accordingly. This framework can also be extended to more

complex metapopulations with more than two subpopulations by incorporating corresponding

migration terms into the dynamical equations.

Conclusion

In conclusion, our comprehensive analysis of six methods for stabilizing metapopulation

dynamics, along with their robustness against census noise, reveals significant insights into

the complex interplay of demographic, genetic, and migration-related factors in population

stability. In comparing control methods, no single approach emerged as universally superior

across all regimes. However, in order to induce constancy stability to a desired level, lower

limiter control (LLC), adaptive limiter control (ALC), and both limiter control (BLC) proved

most effective in Lr regimes. LLC excelled in HrLk regimes by balancing efficacy and effort,

and BLC was the method of choice in HrHk regimes for its ability to eliminate extinction

risks effectively. Remarkably, ALC stood out for inducing persistence stability across all

regimes, achieving the best performance through distinct pathways in different migration

scenarios. Our findings underscored the importance of considering the specific aspects of

stability when selecting and implementing stabilization methods. Furthermore, our analysis of

robustness against census noise highlights the practicality of these methods in real-world

scenarios, with LLC, ALC, and BLC showing promising results in various regimes. This

study not only contributes to our understanding of metapopulation dynamics but also offers

practical guidelines for selecting appropriate population control methods in varied ecological

contexts. It highlights the need for a nuanced, context-specific approach when implementing

population control strategies for effective ecological management and conservation efforts. It

also provides a general framework to study the stability of metapopulations involving

population control strategies.

Acknowledgement
Akanksha Singh thanks Department of Science and Technology, Government of India, for

financial support through a KVPY fellowship. Sudipta Tung acknowledges the support of

DBT/Wellcome Trust India Alliance Early Career Fellowship (#IA/E/18/1/504347) and

Ashoka University.

Authors' contributions: Akanksha Singh and Sudipta Tung formulated the study. Akanksha
Singh carried out the simulations. Sudipta Tung and Akanksha Singh wrote the manuscript.
561

563 **Funding sources**

564
565 The research is funded by DBT/Wellcome Trust India Alliance Early Career Fellowship
566 (#IA/E/18/1/504347) to Sudipta Tung.

References

Abbott, K. C. 2011. A dispersal-induced paradox: synchrony and stability in stochastic
metapopulations. Ecology Letters 14:1158–1169.

Allen, J. C., W. M. Schaffer, and D. Rosko. 1993. Chaos reduces species extinction by
amplifying local population noise. Nature 364:229–232. Nature Publishing Group.

Allendorf, F. W., G. H. Luikart, and S. N. Aitken. 2012. Conservation and the Genetics of
Populations. John Wiley & Sons.

Amarasekare, P. 1998. Interactions between Local Dynamics and Dispersal: Insights from
Single Species Models. Theoretical Population Biology 53:44–59.

Andrievskii, B. R., and A. L. Fradkov. 2003. Control of Chaos: Methods and Applications. I.
Methods. Automation and Remote Control 64:673–713.

Beverton, R. J. H., and S. J. Holt. 2012. On the Dynamics of Exploited Fish Populations.
Springer Science & Business Media.

Bjørnstad, O. N., R. A. Ims, and X. Lambin. 1999. Spatial population dynamics: analyzing
patterns and processes of population synchrony. Trends Ecol Evol 14:427–432.

Borowsky, R. 1971. Principle of Competitive Exclusion and Drosophila. Nature 230:409–
410. Nature Publishing Group.

Brännström, Å., and D. j. t Sumpter. 2005. The role of competition and clustering in
population dynamics. Proceedings of the Royal Society B: Biological Sciences
272:2065–2072. Royal Society.

Briggs, C. J., and M. F. Hoopes. 2004. Stabilizing effects in spatial parasitoid–host and
predator–prey models: a review. Theoretical Population Biology 65:299–315.

Clark, T. J., and A. D. Luis. 2020. Nonlinear population dynamics are ubiquitous in animals.
Nat Ecol Evol 4:75–81. Nature Publishing Group.

Corron, N. J., S. D. Pethel, and B. A. Hopper. 2000. Controlling Chaos with Simple Limiters.

Phys. Rev. Lett. 84:3835–3838. American Physical Society.

Dattani, J., J. C. H. Blake, and F. M. Hilker. 2011. Target-oriented chaos control. Physics

Letters A 375:3986–3992.

Denney, N. H., S. Jennings, and J. D. Reynolds. 2002. Life–history correlates of maximum

population growth rates in marine fishes. Proceedings of the Royal Society of

London. Series B: Biological Sciences 269:2229–2237. Royal Society.

Dey, S., B. Goswami, and A. Joshi. 2014. Effects of symmetric and asymmetric dispersal on

the dynamics of heterogeneous metapopulations: Two-patch systems revisited.

Journal of Theoretical Biology 345:52–60.

Dey, S., and A. Joshi. 2013. Effects of constant immigration on the dynamics and persistence

of stable and unstable Drosophila populations. Sci Rep 3:1405. Nature Publishing

Group.

Dey, S., and A. Joshi. 2007. Local Perturbations Do Not Affect Stability of Laboratory

Fruitfly Metapopulations. PLOS ONE 2:e233. Public Library of Science.

Dey, S., and A. Joshi. 2006. Stability via Asynchrony in Drosophila Metapopulations with

Low Migration Rates. Science 312:434–436. American Association for the

Advancement of Science.

Doebeli, M. 1995. Dispersal and Dynamics. Theoretical Population Biology 47:82–106.

Domokos, G., and I. Scheuring. 2004. Discrete and continuous state population models in a

noisy world. Journal of Theoretical Biology 227:535–545.

Escudero, C., J. Buceta, F. J. de la Rubia, and K. Lindenberg. 2004. Extinction in population

dynamics. Phys. Rev. E 69:021908. American Physical Society.

Fradkov, A. L., and R. J. Evans. 2005. Control of chaos: Methods and applications in

engineering. Annual Reviews in Control 29:33–56.

Fryxell, J. M., I. M. Smith, and D. H. Lynn. 2005. Evaluation of alternate harvesting

strategies using experimental microcosms. Oikos 111:143–149.

Gamelon, M., V. Grøtan, A. L. K. Nilsson, S. Engen, J. W. Hurrell, K. Jerstad, A. S. Phillips,

O. W. Røstad, T. Slagsvold, B. Walseng, N. C. Stenseth, and B.-E. Sæther. 2017.

Interactions between demography and environmental effects are important

determinants of population dynamics. Science Advances 3:e1602298. American

Association for the Advancement of Science.

Grimm, V., and C. Wissel. 1997. Babel, or the Ecological Stability Discussions: An

Inventory and Analysis of Terminology and a Guide for Avoiding Confusion.

Oecologia 109:323–334. Springer.

Grimm, V., and C. Wissel. 2004. The intrinsic mean time to extinction: A unifying approach

to analysing persistence and viability of populations. Oikos 105:501–511.

Gusset, M., O. Jakoby, M. S. Müller, M. J. Somers, R. Slotow, and V. Grimm. 2009. Dogs on

the catwalk: Modelling re-introduction and translocation of endangered wild dogs in

South Africa. Biological Conservation 142:2774–2781.

Gyllenberg, M., G. Söderbacka, and S. Ericsson. 1993. Does migration stabilize local

population dynamics? analysis of a discrete metapopulation model. Mathematical

Biosciences 118:25–49.

Hanski, I. 1998. Metapopulation dynamics. Nature 396:41–49. Nature Publishing Group.

Hastings, A. 1993. Complex Interactions Between Dispersal and Dynamics: Lessons From

Coupled Logistic Equations. Ecology 74:1362–1372. Ecological Society of America.

Hastings, A. 2004. Transients: the key to long-term ecological understanding? Trends in

Ecology & Evolution 19:39–45.

Henson, S. M., R. F. Costantino, J. M. Cushing, R. A. Desharnais, B. Dennis, and A. A.

King. 2001. Lattice Effects Observed in Chaotic Dynamics of Experimental

Populations. Science 294:602–605. American Association for the Advancement of

Science.

Hilker, F. M., and F. H. Westerho□. 2005. Control of chaotic population dynamics:

Ecological and economic considerations. Beitraege des Instituts fuer Umwelt-

systemforschung (No 32).

Ives, A. R., S. T. Woody, E. V. Nordheim, C. Nelson, and J. H. Andrews. 2004. The

synergistic effects of stochasticity and dispersal on population densities. Am Nat

163:375–387.

Kendall, B. E., and G. A. Fox. 1998. Spatial structure, environmental heterogeneity, and

population dynamics: analysis of the coupled logistic map. Theor Popul Biol 54:11–

37.

Lecomte, J., K. Boudjemadi, F. Sarrazin, K. Cally, and J. Clobert. 2004. Connectivity and

homogenisation of population sizes: an experimental approach in Lacerta vivipara.

Journal of Animal Ecology 73:179–189.

Lundberg, P., E. Ranta, J. Ripa, and V. Kaitala. 2000. Population variability in space and

time. Trends in Ecology & Evolution 15:460–464.

May, R. M., and G. F. Oster. 1976. Bifurcations and Dynamic Complexity in Simple

Ecological Models. The American Naturalist 110:573–599. The University of

Chicago Press.

McCallum, H. I. 1992. Effects of immigration on chaotic population dynamics. Journal of

Theoretical Biology 154:277–284.

Mueller, L. D., and A. Joshi. 2020. Stability in Model Populations (MPB-31). P. *in* Stability

in Model Populations (MPB-31). Princeton University Press.

Parthasarathy, S., and S. Sinha. 1995. Controlling chaos in unidimensional maps using

constant feedback. Phys. Rev. E 51:6239–6242. American Physical Society.

Ponciano, J. M., F. P. J. Vandecasteele, T. F. Hess, L. J. Forney, R. L. Crawford, and P.

Joyce. 2005. Use of stochastic models to assess the effect of environmental factors on

microbial growth. Appl Environ Microbiol 71:2355–2364.

Ranta, E., V. Kaitala, and P. Lundberg. 1998. Population Variability in Space and Time: The

Dynamics of Synchronous Population Fluctuations. Oikos 83:376–382. [Nordic

Society Oikos, Wiley].

Ricker, W. E. 1954. Stock and Recruitment. J. Fish. Res. Bd. Can. 11:559–623. NRC

Research Press.

Ruxton, G. D. 1997. Low levels of immigration between chaotic populations can reduce

system extinctions by inducing asynchronous regular cycles. Proceedings of the Royal

Society of London. Series B: Biological Sciences 256:189–193. Royal Society.

Sah, P., J. Paul Salve, and S. Dey. 2013. Stabilizing biological populations and

metapopulations through Adaptive Limiter Control. Journal of Theoretical Biology

320:113–123.

Sheeba, V., and A. Joshi. 1998. A test of simple models of population growth using data from

very small populations of Drosophila melanogaster. Current Science 75:1406–1410.

Temporary Publisher.

Smith, N. R., and B. Meerson. 2016. Extinction of oscillating populations. Phys. Rev. E

93:032109. American Physical Society.

Solé, R. V., J. G. P. Gamarra, M. Ginovart, and D. López. 1999. Controlling Chaos in

Ecology: From Deterministic to Individual-based Models. Bulletin of Mathematical

Biology 61:1187–1207.

Stacey, P. B., M. L. Taper, and V. A. Johnson. 1997. 12 - Migration within Metapopulations:

The Impact upon Local Population Dynamics. Pp. 267–291 *in* I. Hanski and M. E.

Gilpin, eds. Metapopulation Biology. Academic Press, San Diego.

691 Tung, S., A. Mishra, and S. Dey. 2014. A comparison of six methods for stabilizing

692   population dynamics. J Theor Biol 356:163–173.

693 Tung, S., A. Mishra, and S. Dey. 2016a. Simultaneous enhancement of multiple stability

694   properties using two-parameter control methods in Drosophila melanogaster.

695   Ecological Complexity 26:128–136.

696 Tung, S., A. Mishra, and S. Dey. 2016b. Stabilizing the dynamics of laboratory populations

697   of Drosophila melanogaster through upper and lower limiter controls. Ecological

698   Complexity 25:18–25.

699 Tung, S., M. Rajamani, A. Joshi, and S. Dey. 2019. Complex interaction of resource

700   availability, life-history and demography determines the dynamics and stability of

701   stage-structured populations. Journal of Theoretical Biology 460:1–12.

702 Ylikarjula, J., S. Alaja, J. Laakso, and D. Tesar. 2000. Effects of patch number and dispersal

703   patterns on population dynamics and synchrony. J Theor Biol 207:377–387.

 Tables

Table 1: Mathematical formulation of the six control methods and their parameter ranges Here, b_t and a_t are population sizes before and after external perturbations at time t respectively. max() and min() indicate maximum and minimum function respectively.

Sl. No.	Control Method	Brief Description	Mathematical Expression	Control Parameters	Control Parameter Range	Step Size
1.	Constant Pinning (CP)	Add a constant number (p) of individuals at every time step	$a_t = b_t + p$	Pin (p)	[1, k - 1]	1
2.	Lower Limiter Control (LLC)	Increase population size to the lower threshold h whenever it falls below h	$a_t = \max(b_t, h)$	Lower limit (h)	[1, k – 1]	1
3.	Adaptive Limiter Control (ALC)	Similar to LLC, but with the lower threshold set as a fixed fraction (c) of the previous generation's breeding population size (a_{t-1})	$a_t = \max(b_t, c.a_{t-1})$	ALC intensity (c)	[0.05, 0.95]	0.05
4.	Upper Limiter Control (ULC)	Decrease population size to the upper threshold H whenever it goes above H	$a_t = \min(b_t, H)$	Upper limit (H)	[k+1, 3k]	1
5.	Target Oriented Control (TOC)	Limit the population size within a predefined c_d fraction of	$a_t = \max(0, b_t + c_d(T - b_t))$	Target (T)	k	

				the target threshold T			
				TOC intensity (c_d)	[0.05, 0.95]	0.05
6.	Both Limiter Control (BLC)	A combination of LLC and ULC	$a_t = \max(h, \min(b_t, H))$	Lower limit (h)	[1, k – 1]	1
				Upper limit (H)	[k+1, 3k]	1

Figure Captions:

Figure 1. Simulation schematics and parameters. This figure presents a schematic of the metapopulation framework (A), event timeline (B) and parameters used to define various regimes in this study (C). (A) The schematic diagram shows the two-patch metapopulation system used for this study, where local dynamics each patch is modelled using Ricker growth model. We incorporated noise in both intrinsic growth rate (r) and carrying capacity (K) parameters in each iteration of the simulations by drawing random numbers from a uniform distribution as indicated in the figure. (B) The event timeline indicates the sequence in which the population growth model, demographic stochasticity, migration and external perturbation were implemented in the simulations. The perturbation step is skipped for simulating the dynamics of the unperturbed regimes. (C) This table provides the intrinsic growth rate (r), carrying capacity (K) and migration (m1 and m2) parameter values used in the simulations to create contrasting metapopulation regimes.

Figure 2. Unperturbed Metapopulation Dynamics. This figure presents various indices of population dynamics across 12 distinct regimes. Indices for subpopulations and the overall metapopulation are displayed separately, with the left set of graphs (a, c, e, g) illustrating symmetric migration scenarios and the right set (b, d, f, h) depicting asymmetric migration. The indices detailed include: Fluctuation Index (FI) in (a) and (b), highlighting dynamic variability; Extinction Probability (EP) in (c) and (d), with Lr regimes displaying negligible to zero EP; Effective Population Size (EPS) in (e) and (f), representing genetic diversity; and Synchrony between subpopulations in (g) and (h). Standard error was used as error bars. For some cases, error bars are too small to be visible.

Figure 3. Composite Performance Scores for 50% Reduction in Fluctuation Index. Presented here are the composite performance scores for six population control methods evaluated across 12 distinct regimes, with the objective of achieving a 50% decrease in Fluctuation Index to promote constancy stability. Atop each bar, the numerical annotations represent the performance ranking of the method, with a lower composite score denoting a superior rank. Standard error is depicted as the error bars; however, in certain instances, the error bars are too small to be visible.

Figure 4. Composite Performance Scores for 50% Reduction in Extinction Probability. Presented here are the composite performance scores for six population control methods evaluated across six regimes, with high intrinsic growth rate (i.e. Hr regimes) while achieving a 50% decrease in Extinction Probability to promote persistence stability. Atop each bar, the numerical annotations represent the performance ranking of the method, with a lower composite score denoting a superior rank. Standard error is depicted as the error bars; however, in certain instances, the error bars are too small to be visible.

Figure 5. Robustness Indices for Population Stability Methods While Inducing Constancy Stability Under Varying White Noise Levels. This figure displays the robustness indices of six population stability methods evaluated across 12 distinct regimes tasked with inducing constancy stability, specifically a 50% reduction in the Fluctuation Index, in the presence of different intensities of white noise affecting population size estimates. The robustness index is determined by the average squared deviation between composite scores obtained under various white noise levels and that derived in the absence of noise. This metric evaluates the consistency of each method's performance in the face of census accuracy. To present the complete data range within one frame, a discontinuous Y-axis has been employed. Numerical annotations above each bar indicate the method's robustness rank, with lower scores corresponding to higher robustness against noise. Standard errors are represented by error bars. In certain instances, the error bars are too small to be visible.

Figure 6. Robustness Indices for Population Stability Methods While Inducing Persistence Stability Under Varying White Noise Levels. This figure displays the robustness indices of six population stability methods evaluated across six regimes with high intrinsic growth rate i.e. Hr regimes aimed to induce persistence stability, specifically a 50% reduction in the Extinction Probability, in the presence of different intensities of white noise affecting population size estimates. The robustness index is determined by the average squared deviation between composite scores obtained under various white noise levels and that derived in the absence of noise. This metric evaluates the consistency of each method's performance in the face of census accuracy. To present the complete data range within one frame, a discontinuous Y-axis has been employed. Numerical annotations above each bar indicate the method's robustness rank, with lower scores corresponding to higher robustness against noise. Standard errors are represented by error bars. In certain instances, the error bars are too small to be visible.

Unperturbed Regimes