Effects of excess phosphate on a coastal plankton community

Kristian Spilling[1,2], Mari Vanharanta[1,3], Mariano Santoro[4,5], Cristian Villena-Alemany[6,7], Matthias Labrenz[5], Hans-Peter Grossart[8, 9], Kasia Piwosz[10]

[1]Marine and Freshwater Solutions, Finnish Environment Institute, 00790 Helsinki, Finland

[2]Centre for Coastal Research, University of Agder, 4630 Kristiansand, Norway

[3]Tvärminne Zoological Station, University of Helsinki, 10900 Hanko, Finland

[4]Department of Biological Oceanography, Leibniz Institute for Baltic Sea Research Warnemünde - IOW, Rostock, Germany

[5]Department of Plant Physiology, Institute for Biosciences, University of Rostock, Rostock, Germany

[6]Laboratory of Anoxygenic Phototrophs, Institute of Microbiology, Czech Academy of Sciences, 379 01 Třeboň, Czech Republic

[7]Department of Ecosystem Biology, Faculty of Science, University of South Bohemia, České Budějovice, Czech Republic

[8]Leibniz Institute for Freshwater Ecology and Inland Fisheries, Germany

[9]Institute of Biology and Biochemistry, Potsdam University, Germany

[10]National Marine Fisheries Research Institute – NMFRI, Gdynia, Poland

Abstract

Eutrophication in the Baltic Sea has caused an imbalance in the inorganic nitrogen (N) to phosphorus (P) ratio, leaving excess phosphate (PO_4) remaining after the phytoplankton spring bloom that terminates after N depletion. Here we investigated the fate of such excess phosphate in a two-week mesocosm (1.2 m^3) experiment. The starting concentration of PO_4 was 0.66 µM, and treatments included a non-treated control (control), nitrate addition (N-add; 3.6 µM), glucose addition (C-add; 25 µM) and combined nitrate and glucose addition (N+C-add); all treatments done in triplicates. Addition of N (N-add and N+C-add) stimulated nano- and microphytoplankton, with the picophytoplankton abundance increasing after N depletion. Also the Copepod biomass was positively affected by the N-addition. N-fixing cyanobacteria were present but in low abundance. Carbon addition did not enhance heterotrophic bacterial uptake of PO_4, nor affecting the phyto- or zooplankton community composition. The PO_4 concentration was reduced to ~0.4 µM in the control and C-add treatments and to 0.16 µM in the two N-amended treatments, with an inorganic NP uptake ratio of 7.2. The results underscore the role of picophytoplankton in reducing the excess phosphate pool after the spring bloom, a function traditionally ascribed to bloom-forming diazotrophic cyanobacteria in the Baltic Sea.

Key words: biogeochemical cycles, coastal seas, eutrophication, nutrient stoichiometry, *Nodularia spumigena* and *Aphanizomenon flos-aquae*, Baltic Sea

Introduction

Loss of oxygen, i.e. deoxygenation, in the ocean is an increasing global problem (Keeling et al.

2010) that has implications not only directly for organisms using oxygen in their metabolism but

also effects biogeochemical cycles. There are two primary causes of the deoxygenation of the

ocean: global warming and human input of nutrients to marine ecosystems, i.e. eutrophication.

Warming of surface waters happens on a global scale and reduces gas solubility including oxygen.

Eutrophication is a local or regional environmental problem that causes algal blooms that typically

aggravate oxygen consumption in deeper water layers after the bloom, when the biomass is sinking

and decomposing.

In relatively shallow coastal seas, elevated oxygen consumption may cause hypoxic and anoxic

conditions on the seafloor, and sediment microbial processes typically reduce inorganic nitrogen

through denitrification and anammox (Hannig et al. 2007). At the same time, anoxia increases the

release of phosphate through the chemical reduction of iron-bound phosphate. The net result of

anoxic bottoms is a decreased bioavailable nitrogen and an increased phosphate concentration,

causing the inorganic NP ratio to decrease in several parts of the world (Conley et al. 2002, Hauss et

al. 2012, Kalvelage et al. 2013, Meyer et al. 2016).

The Baltic Sea has been affected by eutrophication for several decades, which has increased the

area of anoxic seafloor and the release of phosphate (Conley et al. 2002). In the Baltic Proper and

Gulf of Finland, the most productive period with accompanying flux of organic material to the

sediment is during the spring bloom, which terminates when dissolved inorganic Nitrogen (DIN)

has been depleted. There was a regime-like shift in the mid-1990s when the dissolved inorganic

phosphorus (DIP) pools started to be at elevated levels after the spring bloom; low phosphate

concentrations were generally detected during early summer before the mid-1990s, although N-

limitation of the spring bloom was demonstrated even then (Tamminen 1995).

Much of the Baltic Sea is nitrogen-limited and there has been a trend of decreasing inorganic NP

ratios in several sub-basins caused primarily by release of phosphate from the seafloor after turning

anoxic (Conley et al. 2002). This has led to a surplus of phosphate after the spring bloom that may

range from 0.1 up to 0.6 µmol PO_4 L^{-1} in the Gulf of Finland and Baltic Proper (Spilling et al.

2018). This excess phosphate is believed to directly benefit N-fixing cyanobacteria that often create

toxic or nuisance blooms during the summer (Larsson et al. 2001, Kangro et al. 2007).

Nodularia spumigena and *Aphanizomenon flos-aquae* are amongst the typical dominating

filamentous cyanobacteria, forming blooms in July and August in large parts of the Baltic Sea.

Although these species can take up and store P in the form of polyphosphate, they grow relatively

slowly in water temperatures below 15°C (although *A. flos-aquae* is less sensitive to cold water than

N. spumigena), and typically start growing after phosphate has been already depleted. An

alternative pathway of P is going through other primary producers or heterotrophic bacteria (Nausch

and Nausch 2004), reducing the available P for the filamentous cyanobacteria. However, the

prevailing N-limitation restricts the growth of other phytoplankton and consequently limits their

luxury uptake of P. Heterotrophic bacteria may also take up P, but these are typically limited by

carbon (Lignell et al. 1992). At present it remains unclear which organisms take up the excess

phosphate, as most of it seems to settle out of the euphotic zone before cyanobacterial blooms

develop (Nausch et al. 2008) and other P sources must exist for these cyanobacteria (Raateoja et al.

2011), most likely through upwelling (Wasmund et al. 2012).

Here we addressed the fate of this excess phosphate during an early summer experiment via

additions of bioavailable nitrogen (N), carbon (C) and both (N+C), to stimulate primary producers

and heterotrophic bacteria. We hypothesized that picophytoplankton would become relatively more

important after N depletion taking up part of the excess phosphate, but that heterotrophic bacteria

can also exploit the inorganic P pool when they have access to a labile carbon source.

Materials and methods

The experiment was set up in 1.2 m^3 mesocosm bags that were moored outside the Tvärminne

Zoological Station, Gulf of Finland (59° 50' 40" N, 23° 14' 57" E). The bags were placed in a row

next to a floating platform (Spilling et al. 2022a). In total, 14 bags were installed but the two

outermost bags were not used. These 'dummy' bags were used to create the same light conditions

for the outermost experimental units. All units were covered to prevent rain or bird droppings to fall

into the bags. The spring bloom in the area had peaked in April and the PO_4 concentration was

relatively low (0.013 μ mol L^{-1}), and we added PO_4 to all units to a final concentration of 0.66 μ

mol L^{-1}. The experimental setup consisted of a control, a treatment with added nitrate (N-add; 3.6

μmol N L^{-1}), a treatment with added carbon in the form of glucose (C-add, 36 μmol C L^{-1}), and a

treatment with both nitrate and glucose addition (N+C-add).

The bags were filled by gentle pumping in surface water on 7th June 2021, which was day -1 of the

experiment. This filling method had visually (microscope) been tested and found not to affect

zooplankton but prevents small fish from entering the mesocosms. The following day (day 0),

nutrients and carbon sources were added in the morning and the first full sampling took place right

after the addition. The temperature was approximately 18☐ and the salinity was 5.5 at the start of

the experiment (Supplementary Figs 1 and 2). Water samples were taken daily with a Limnos water

sampler (Hydro-Bios, Germany) from 1.5 m depth from the middle of the bags. CTD profiles were

taken daily from the bags and outside the experimental units, and we had temperature and dissolved

oxygen loggers (HOBO U26; Onset Inc. USA) mounted inside each mesocosm bag. In addition to

minor sampling every day (e.g. CTD profile), the regular sampling was carried out on days 0, 1, 3,

6, 8,10, 13 and 15.

Total chlorophyll a (Chla) fluorescence was determined from each mesocosm bag by daily use of a

handheld fluorometer (AquaPen, Photon Systems Instruments, Czech Republic), and direct Chla

measurements were carried out by filtration in duplicates on regular sampling days. For Chla, GF/F

filters (Whatman, UK) were used and the Chla was extracted in EtOH, stored at -20°C and in the

dark for up to two months, and measured with a spectrofluorometer (Varian Cary Eclipse, Agilent,

USA) with excitation and emission wavelengths of 430 and 670 nm, respectively (Jespersen and

Christoffersen 1987).

We used standard colorimetric methods (Grasshoff et al. 1999) using a photometric analyzer

(Aquakem 250, Thermo Scientific, USA) to measure dissolved inorganic nutrients: nitrite + nitrate

($NO_2 + NO_3$), phosphate (PO_4) and dissolved silicate (Dsi). Ammonium (NH_4) was measured

separately using a spectrophotometer (U-1100, Hitachi, Japan).

Particulate organic carbon (POC), nitrogen (PON) and phosphorus (POP) were determined from

duplicate water samples filtered onto acid-washed (2M HCl for 15 min, then rinsed carefully with

ultrapure water) and pre-combusted (450°C, 4 h) GF/F filters (Whatman, UK). POC and PON were

measured with an element analyzer coupled with a mass spectrometer (Europa Scientific ANCA-

MS 20-20 15N/13C, UK). POP was determined according to Solórzano and Sharp (1980) as

modified by Koistinen et al. (2017). Samples for biogenic silicate (Bsi) were filtered on

polycarbonate membrane filters (0.8 μm, GVS, Italy) and measured according to Koistinen et al.

(2018).

Flow cytometer counts were done daily with a Sysmex (Japan), Partec – Cube 8 equipped with two

lasers (488 and 561 nm), two scattering (forward and side) and three fluorescence detectors

(610/30; 661/16; and 670/40, corresponding to the detection of phycoerythrin, phycocyanin, and

Chla, respectively). The trigger was set on Chla fluorescence (670 nm). On each sampling day, we

made several measurement-runs with beads and blanks to ensure correct particle counts. Size

fractionations (0.8, 1, 2, 5, 10, 20 μm filters) were also used to identify the approximate size of

different phytoplankton groups. The scattering and fluorescence properties were used to gate

different phytoplankton groups using the FCS Express 6 software. Five groups were identified:

picoeucaryotes: size <2 µm with only Chla, *Synechococcus*-like cells <2 µm with phycocyanin

fluorescence, Cryptophyte-like >2 µm with Chla and phycoerythrin fluorescence,

nanophytoplankton size 2–20 µm and microphytoplankton: size >20µm with only Chla

fluorescence.

The abundance of heterotrophic bacteria were enumerated with another flow cytometer (LSR II, BD

Biosciences, USA equipped with a 488 nm laser) after the samples first had been fixed with 1%

paraformaldehyde (final concentration). Samples were stored for 15 min, flash frozen in liquid

nitrogen, and stored at -80□. Before measurements, samples thawed and stained with SYBRGreen I

(Molecular Probes, Eugene, OR, USA) and countBright beads (Molecular Probes) were added to

each sample to determine measured volume. Heterotrophic bacteria were detected according to their

green fluorescence and side scatter properties using FACSDiva Software (BD Biosciences). In

addition, aerobic anoxygenic phototrophic (AAP) bacteria were counted using the

bacteriochlorophyll-*a* fluorescence in epifluorescence microscope according to Villena-Alemany et

al. (2023).

Zooplankton was collected three times, at the start, middle and end of the experiment. Samples were

taken with a plankton net (25 cm diameter, 50 µm mesh size) carefully dragged from the bottom of

the mesocosm bags to the surface. Samples were immediately preserved with acid Lugol's solution

and stored at 4°C. Zooplankton was identified to genus level from images created using a flatbed

scanner and analyzed at 20 x magnification.

Data analysis

A one-way analysis of variance (ANOVA) was performed to test whether significant effects could

be revealed by the treatments. Tukey's Honest Significant Difference (HSD) post hoc test was

performed to compare different treatments and assess the significance of differences between them.

All statistical tests were performed using the SigmaPlot 15 software.

Results

There was a clear effect of N addition, but no effect of carbon addition, on the phosphate

concentration (Fig 1). The phosphate concentration decreased from ~0.66 µmol L^{-1} at the start of

the experiment to ~0.16 µmol L^{-1} in the two treatments with N added on Day 8, an inorganic NP

uptake ratio of 7.2. In the control and C-addition, the PO_4 concentration was 0.33 and 0.36 µmol

$PO_4 L^{-1}$, respectively, at the end of the experiment. There was residual ammonium with a

concentration in the range of 0.15 to 0.35 µmol L^{-1} throughout the entire experiment. Dissolved

silicate was constant in the control and C-add but decreased slightly towards the end of the

experiment in the N-add and N+C add treatments, with an uptake of approximately 1.9 µmol L^{-1}

(Fig 1).

The phytoplankton biomass was declining from the onset of the experiment in the control and C-

add treatments but with a delay of two days in the two treatments with added N (Fig 1). The initial

Chla concentration was 3.5 µg L^{-1} that decreased to 1–2 µg Chla L^{-1} and within two days in both

control and C-add treatments (Fig 2). After reaching a minimum of <1 µg L^{-1}, the Chla

concentration increased again at the end of the experiment to ~2 µg Chla L^{-1}. The nitrate added to

the N-add and N+C-add treatments was depleted already on day 3.

Heterotrophic bacteria had the same dynamics in all treatments (Fig 2); their abundance increased

after day 1, with a peak at day 8. The overall bacterial abundance was higher in the N-add treatment

compared to the control and C-add treatments (Tukey: p = 0.001 and p = 0.003 respectively), the

rest of the treatments were similar (p > 0.14). Aerobic anoxygenic phtotrophic (AAP) bacteria

followed similar abundance patterns that overall heterotrophic bacteria, increasing from the

beginning of the experiment in all conditions, but differed by showing higher abundance increase

than the overall heterotrophic bacteria (Supplementary Fig S3). This difference was more

conspicuous in N add treatment where AAP bacteria reached their maximum absolute abundance

(1.69×10^5 cells ml^{-1}) and the highest contribution to bacterial community (2.28%).

All particulate nutrients decreased in the experimental units during the first week, but the decrease

was more rapid in treatments without any nitrate addition (control and C-add), except for biogenic

silicate (BSi) where the decrease was similar in all treatments (Fig 3). Stoichiometric ratios were

relatively stable with an average C:N:P ratio of 140:14:1, and without any differences between

treatments (p >0.1).

The temperature and salinity throughout the experiment is presented in Supplementary Figs S1 and

S2. There was a drop in temperature after day 6 from ~18□ to ~14□ due to upwelling, but this

increased again towards the end of the experiment to ~20□. Salinity was constant in the bags and

there was no sign of mixing with the outside water.

Community

Nitrate additions benefited micro- and nanophytoplankton that increased in abundance in the N-add

and N+C add treatments (Fig 4), contributing to the high Chla concentration over the first three

days. This was also evident from the FlowCam images, where, *e.g.* the cell concentration and total

biovolume of dinoflagellates >10 µm increased in these two treatments until nitrate had been

depleted (Fig 5). The green algae *Monoraphidium* sp. (nanophytoplankton) increased also in the N-

amended treatments, but kept growing after N depletion, reaching a maximum biovolume at

experiment day 8 (Fig 5).

After nitrate depletion, picoeukaryotes started to increase in abundance, especially in the control

and N-add treatment, but also, to a lesser extent, in the N+C-add and C-add treatments (Fig 4).

Synechococcus-like cells decreased in all treatments after the start of the experiment and stayed at a

low abundance until the last few days when it started to increase in all treatments, with the highest

increase in the N-add treatment (Fig 3). *Planktothrix* sp., and *Pseudoanabaena* sp. were the main

component of colonial and filamentous cyanobacteria. They peaked in the N-add and N+C-add

treatments on day 3 and 6, respectively, with the highest peak in biovolume in the N+C-add

treatment (Fig 6). *Aphanizomenon flos-aquae* and *Nodularia spumigena* were present but was

combined less than 5% of the cyanobacterial biovolume.

Microzooplankton consisted of heterotrophic dinoflagellates and ciliates (Fig 5). Heterotrophic

dinoflagellates increased in the N-add and N+C-add treatments with a peak on day 8 before

declining to reach similar biovolume as in the control and the C-add treatments at the end of the

experiment (Fig 6). The biovolume of ciliates was highly variable between replicates and - although

the average biovolume was higher in the two N add treatments, there was no consistent difference

between treatments (ANOVA, $p = 0.767$).

The mesozooplankton community was dominated by copepods, cladocerans and pelagic stages of

Amphibalanus sp. (Fig 7). The number of copepods was highest in N-add and N+C-add treatments

towards the end of the experiment including both copepodite and nauplii stages ($p = 0.024$ and $p = $

0.017, respectively). Among cladocerans, the abundance of *Bosmina* sp. was positively affected by

the nitrate addition ($p = 0.04$), whereas *Podon* sp. (Fig 6) and *Evadne* sp. (data not shown) both

decreased over time and had completely disappeared in all treatments at the end of the experiment.

The difference in *Amphibalanus* nauplii abundance between treatments was not clear ($p = 0.06$), but

cypris larvae had only developed in the two N-amended treatments at the end of the experiment (Fig

7)

Discussion

Inorganic nutrients initially stimulated nano- and microphytoplankton. This was expected, as they

have a larger volume for internal storage, i.e. luxury uptake. Nonetheless, this effect only delayed

the gradual decline of phytoplankton biomass. The main driver for the differences between

treatments was the addition of nitrate, whereas the carbon addition did not affect the community of

the larger phytoplankton or zooplankton. After nitrate had been depleted, there was a clear increase

in picophytoplankton abundance, which was also observed in previous mesocosm experiments in

this area (Crawfurd et al 2017, Spilling et al. 2022b). Smaller cells have a larger surface-to-volume

area and are better adapted to low inorganic nutrient concentrations, which is the likely reason for

this group to be dominating during periods when easily available nutrient sources have been

depleted by larger cells. Often, a surplus of P would benefit nitrogen-fixing cyanobacteria,

especially if they have been able to increase their biomass during the nitrate depletion phase.

Although cyanobacteria were present in all treatments, there was no indication that the typical

bloom-forming diazotrophic species proliferated. Rather, the additional P uptake after N depletion

seems to have been driven by the increase in picoeukaryotes and possibly smaller filamentous

cyanobacterial (*Planktothrix* sp, and *Pseudoanabaena* sp) were taking up phosphate and storing part

of the excess phosphorus as polyphosphate (Jentsch et al. 2023). This was also seen under the

microscope, but we have no quantitative data on polyphosphate storage.

Interestingly, carbon addition had a negative effect on picoeukaryotes relative to the other

treatments. This could have been due to competition for nutrients with heterotrophic bacteria, which

likely got a boost due to the sudden, easily available carbon source that was added. This, however,

is not evidenced by the abundance of heterotrophic bacteria that remained similar between control

and C-add treatment. This could be different for particle-attached heterotrophic bacteria, but

unfortunately no data is available for this group. Alternatively, grazing pressure could have played a

role, but no noticeable difference in microzooplankton abundance was observed.

The addition of glucose did stimulate bacterial productivity and extracellular activity (Vanharanta et

al. 2024), but there was no indication of any bacterial drawdown of phosphate when compared to

the control. Also, in this respect, the combined N+C addition was similar to the N addition. This

further supports that the majority of phosphate uptake was due to picoeukaryotes with little to no

contribution of heterotrophic bacteria. The timing of the heterotrophic bacterial abundance peak is

typical following the spring bloom when there is a lot of organic matter available (Camarena-

Gómez at al. 2018), and the community is to some extent affected by the phytoplankton community

during the spring bloom (Camarena-Gómez et al. 2021). The absolute abundances and contribution

of AAP bacteria to bacterial community were consistent with previous findings in the Baltic Sea

(Mašín et al., 2006; Salka et al., 2008). Under all conditions AAP bacteria demonstrated higher

growth rates compared to the general heterotrophic bacteria, especially in the N-add treatment. The

phytoplankton spring bloom have been shown to enhance AAP bacteria proliferation in fresh waters

(Villena-Alemany et al., 2023), and our results support that AAP bacteria are impacted by the

phosphorous and nitrogen availability (Mašín et al., 2012) and has potential in phosphate removal.

In the N-add and N+C-add treatments, ~75% of the phosphate was removed, whereas in the control

and C-add treatments ~ 40% was removed within a week. Approximately 15% of the phosphate

pool, or 20–30% of the PO_4 taken up was released into the dissolved organic phosphorus (DOP)

pool, but DOP concentration decreased again during the last week of the experiment suggesting the

added DOP was relatively labile and microbially available. The NP uptake ratio of 7.2, less than

half the Redfield ratio of 16, suggest an active luxury uptake of P of the plankton community. The

decrease in the phosphate pool was not seen as an increase in the particulate phosphorus fraction

although the N-amended treatments had a higher concentration of particulate organic elements. This

was likely due to export of particulate organic matter to the bottom of the mesocosm bags.

The mesocosm bags prevent effective wind-driven mixing, although there was some temperature

driven advection. There was particulate organic matter sedimentation during the experiment, but

unfortunately, we could not sample the bottom part of the mesocosms to quantify it. This export

loss might have had some consequences for the phytoplankton community composition as non-

motile groups, such as diatoms, likely sedimented out quicker than motile flagellates. This could be

deduced from the development in BSi over time, and diatoms were a minor group in terms of

biovolume. The bags were not very deep and the available organic material in the bottom likely

supported the community of mesozooplankton that could easily move vertically between different

water depths, especially since higher predators were absent in the mesocosms. This could be a

likely reason for a treatment effect on mesozooplankton even though the available, suspended food

items were similar in all bags after the drawdown of nitrate.

In a recent study, using smaller 20 L tanks, we found that the excess phosphate had been depleted

completely under N-deficiency similar to this experiment, even without N addition (Vanharanta and

Spilling 2023). This previous experiment had a longer duration (35 d) than the one presented here,

with drawdown of phosphate in the control units of 20% to 50 % (from a starting concentration of

0.55 µM) after 15 days, depending on the temperature (Vanharanta and Spilling 2023), and it is

likely that we would have also seen a complete drawdown of phosphate, if the experiment would

have lasted a few weeks more.

One of the most prominent effects of eutrophication in the Baltic Sea is the intensifying occurrence

of diazotrophic cyanobacterial blooms during summer. The prevailing N-limitation leads to a low

inorganic N:P ratio, which favors the N-fixing cyanobacteria during warm, calm summer months

(Niemi 1979, Wasmund et al. 2005). These cyanobacterial blooms amplify the eutrophication

problem as the amount of cyanobacterial fixed N is comparable to the anthropogenic N-loading in

parts of the Baltic Sea (Savchuk 2005, Wasmund et al. 2005). The cyanobacterial input of N might

thus have a self-enforcing nature through a positive feedback loop: N-fixation increases the biomass

load reaching the sediment, increasing oxygen consumption, and adding more P to the ecosystem

(Vahtera et al. 2007a, Spilling et al 2018).

The two main cyanobacterial species: *Aphanizomenon flos-aquae* and *Nodularia spumigena* have

different P uptake strategies. *A. flos-aquae* requires inorganic phosphate and can take up and store

excess P for later growth, whereas *N. spumigena* has a higher affinity for phosphate and, thus, relies

more on recycled P (Hagström et al. 2001, Vahtera et al. 2007). Both species were present in the

experiment but at exceptionally low concentrations. It was other filamentous cyanobacteria

(*Planktothrix* sp, and *Pseudoanabaena* sp) that made up the bulk of the measured cyanobacterial

biovolume. On the one hand, this shift in cyanobacterial community composition could be due to an

enclosure effect, as there are examples of different primary producers being negatively affected in a

mesocosm setup. On the other hand, both. *A. flos-aquae* and *N. spumigena* have relatively low

growth rates, and it is likely that they did not grow fast enough in the experiment to increase much

biomass in the relatively short time span of the experiment.

A decreasing inorganic N:P ratio resulting from reduced O_2 concentrations can be observed in

different ecosystems. Short-term shifts are to some extent dependent on the plankton community

composition (Spilling et al. 2019, Bach et al. 2020). However, potential long-term consequences of

this phenomenon are little known. A decreasing NP ratio in marine ecosystems is not an universal

process, however, as there are examples of other processes that can increase this ratio, for example,

more effective P-removal from wastewater, increased use of N-rich fertilizer, or increased

atmospheric N-deposition from anthropogenic sources (McQuatters-Gollop et al. 2007, Kim et al.

2014).

In conclusion, we found that N-addition stimulated both nano- and microphytoplankton, and the

picophytoplankton abundance increased following nitrate depletion. Based on these results, we

conclude that it was primarily the picophytoplankton that incorporated substantial parts of the

excess P, confirming our initial hypothesis. Yet, it was unable to completely deplete the inorganic

P-pool, and most of the organic P formed was exported. At this point, we are missing direct

evidence and measurements of excess phosphate uptake, e.g. using radiolabeled P. Carbon addition

did stimulate heterotrophic bacteria production, but did not increase excess phosphate uptake,

contrary to our expectations. Heterotrophic bacterial uptake of excess phosphate could still take

place under other environmental conditions. The typical bloom-forming cyanobacteria in the Baltic

Sea, which are considered to greatly benefit from excess phosphate, were not abundant in our

mesocosms. The results here underscore the role of picophytoplankton in reducing the excess

phosphate pool after the spring bloom, a function traditionally ascribed to bloom-forming

diazotrophic cyanobacteria in the Baltic Sea.

Acknowledgements

We would like to thank the staff at Tvärminne Zoological Station for all their support, technical

assistance and nutrient measurements. In particular Joanna Norkko, Jaana Koistinen, Mervi

Sjöblom and field technicians Göran Lundberg and Jostein Solbakken.

This study used equipment part of the Finnish marine research infrastructure (FINMARI) program.

This work was supported by the Transnational Access program of the EU H2020-INFRAIA project

(No. 731065) AQUACOSM - Network of Leading European AQUAtic MesoCOSM Facilities

Connecting Mountains to Oceans from the Arctic to the Mediterranean - funded by the European

Commission. Additional funding came from Walter and Andree de Nottbeck foundation (KS and

MV). KP was supported by the National Science Centre, Poland under the Weave program (project

no 2021/03/Y/NZ8/00076). HPG was funded by the German Science Foundation (DFG) project

PycnoTrap (GR1540/37-1). MS was supported by Leibniz Science Campus Phosphorus Research

Rostock in the funding line strategic networks of the Leibniz Association.

References

Bach, L. T., A. J. Paul, T. Boxhammer, E. v. d. Esch, M. Graco, K. G. Schulz, E. Achterberg, P. Aguayo, J. Aristegui, P. Ayon, I. Banos, A. Bernales, A. S. Boegeholz, F. Chavez, S.-M. Chen, K. Doering, A. Filella, M. Fischer, P. Grasse, M. Haunost, J. Hennke, N. Hernandez-Hernandez, M. Hopwood, M. Igarza, V. Kalter, L. Kittu, P. Kohnert, J. Ledesma, C. Lieberum, S. Lischka, C. Loescher, A. Ludwig, U. Mendoza, J. Meyer, J. Meyer, F. Minutolo, J. O. Cortes, J. Piiparinen, C. Sforna, K. Spilling, S. Sanchez, C. Spisla, M. Sswat, M. Z. Moreira, and U. Riebesell. 2020. Factors controlling plankton productivity, particulate matter stoichiometry, and export flux in the coastal upwelling system off Peru. Biogeosciences Discussions.

Camarena-Gómez MT, Ruiz-Gonzalez C, Piiparinen J, Lipsewers T, Sobrino C, Logares R, Spilling K. 2021. Bacterioplankton dynamics driven by interannual and spatial variation in diatom and dinoflagellate spring bloom communities in the Baltic Sea. Limnol Oceanogr. 66: 255-271

Camarena-Gómez MT, Lipsewers T, Piiparinen J, Eronen-Rasimus E, Perez-Quemaliños D, Hoikkala L, Sobrino C, Spilling K. 2018 Shifts in phytoplankton community structure modify bacterial growth, production and composition. Aquat Microb Ecol. 81: 149-170.

Conley, D. J., C. Humborg, L. Rahm, O. P. Savchuk, and F. Wulff. 2002. Hypoxia in the Baltic Sea and basin-scale changes in phosphorus biogeochemistry. Environmental Science & Technology **36**:5315-5320.

Crawfurd, K. J., Alvarez-Fernandez, S., Mojica, K. D. A., Riebesell, U., and Brussaard, C. P. D.: Alterations in microbial community composition with increasing fCO$_2$: a mesocosm study in the eastern Baltic Sea, Biogeosciences, 14, 3831–3849, https://doi.org/10.5194/bg-14-3831-2017, 2017.

Grasshoff, K., K. Kremling, and M. Ehrhardt. 1999. Methods of seawater analysis 3 ed. Wiley-VCH, Weinheim.

Hagström, Å., F. Azam, J. Kuparinen, and U.-L. Zweifel. 2001. Pelagic plankton growth and resource limitations in the Baltic Sea. Pages 177-210 A systems analysis of the Baltic Sea. Springer.

Hannig, M., G. Lavik, M. Kuypers, D. Woebken, W. Martens-Habbena, and K. Jürgens. 2007. Shift from denitrification to anammox after inflow events in the central Baltic Sea. Limnology and Oceanography **52**:1336-1345.

Hauss, H., J. M. Franz, and U. Sommer. 2012. Changes in N: P stoichiometry influence taxonomic composition and nutritional quality of phytoplankton in the Peruvian upwelling. Journal of Sea Research **73**:74-85.

Jentzsch L, Grossart HP, Plewe S, Schulze-Makuch D, Goldhammer T. Response of cyanobacterial mats to ambient phosphate fluctuations: phosphorus cycling, polyphosphate accumulation and stoichiometric flexibility. ISME communications. 2023 Jan 25;3(1):6.

Jespersen, A. M., and K. Christoffersen. 1987. Measurements of chlorophyll-a from phytoplankton using ethanol as extraction solvent. Archiv fur Hydrobiologie **109**:445-454.

Kalvelage, T., G. Lavik, P. Lam, S. Contreras, L. Arteaga, C. R. Löscher, A. Oschlies, A. Paulmier, L. Stramma, and M. M. Kuypers. 2013. Nitrogen cycling driven by organic matter export in the South Pacific oxygen minimum zone. Nature Geoscience **6**:228-234.

Kangro, K., K. Olli, T. Tamminen, and R. Lignell. 2007. Species-specific responses of a cyanobacteria-dominated phytoplankton community to artificial nutrient limitation in the Baltic Sea. Marine Ecology Progress Series **336**:15-27.

Keeling, R. F., A. Körtzinger, and N. Gruber. 2010. Ocean deoxygenation in a warming world. Annual review of marine science **2**:199-229.

Kim, I.-N., K. Lee, N. Gruber, D. M. Karl, J. L. Bullister, S. Yang, and T.-W. Kim. 2014. Increasing anthropogenic nitrogen in the North Pacific Ocean. Science **346**:1102-1106.

Koistinen, J., M. Sjöblom, and K. Spilling. 2017. Determining inorganic and organic Phosphorus.

417   Koistinen, J., M. Sjöblom, and K. Spilling. 2018. Determining dissolved and biogenic Silica.

418   Larsson, U., S. Hajdu, J. Walve, and R. Elmgren. 2001. Baltic Sea nitrogen fixation estimated from
419   the summer increase in upper mixed layer total nitrogen. Limnology and Oceanography
420   **46**:811-820.

421   Lignell, R., S. Kaitala, and H. Kuosa. 1992. Factors controlling phyto- and bacterioplankton in late
422   spring on a salinity gradient in the northern Baltic. Marine Ecology Progress Series **84**:121-
423   131.

424   McQuatters-Gollop, A., D. E. Raitsos, M. Edwards, Y. Pradhan, L. D. Mee, S. J. Lavender, and M.
425   J. Attrill. 2007. A long‐term chlorophyll dataset reveals regime shift in North Sea
426   phytoplankton biomass unconnected to nutrient levels. Limnology and Oceanography
427   **52**:635-648.

428   Mašín, M., Čuperová, Z., Hojerová, E., Salka, I., Grossart, H., and Koblížek, M. (2012) Distribution
429   of aerobic anoxygenic phototrophic bacteria in glacial lakes of northern Europe. Aquat
430   Microb Ecol 66: 77–86.

431   Mašín, M., Zdun, A., Ston-Egiert, J., Nausch, M., Labrenz, M., Moulisová, V., and Koblízek, M.
432   (2006) Seasonal changes and diversity of aerobic anoxygenic phototrophs in the Baltic Sea.
433   Aquat Microb Ecol 45: 247–254.

434   Meyer, J., C. Löscher, S. Neulinger, A. Reichel, A. N. Loginova, C. Borchard, R. Schmitz, H.
435   Hauss, R. Kiko, and U. Riebesell. 2016. Changing nutrient stoichiometry affects
436   phytoplankton production, DOP accumulation and dinitrogen fixation–a mesocosm
437   experiment in the eastern tropical North Atlantic. Biogeosciences **13**:781-794.

438   Nausch, G., D. Nehring, and K. Nagel. 2008. Nutrient concentrations, trends and their relation to
439   eutrophication. State and Evolution of the Baltic Sea, 1952-2005: A Detailed 50-Year
440   Survey of Meteorology and Climate, Physics, Chemistry, Biology, and Marine
441   Environment:337-366.

442   Nausch, M., and G. Nausch. 2004. Bacterial utilization of phosphorus pools after nitrogen and
443   carbon amendment and its relation to alkaline phosphatase activity. Aquatic Microbial
444   Ecology **37**:237-245.

445   Niemi, Å. 1979. Blue-green algal blooms and N:P ratio in the Baltic Sea. Acta Botanica Fennica
446   **110**:57-61.

447   Raateoja, M., H. Kuosa, and S. Hällfors. 2011. Fate of excess phosphorus in the Baltic Sea: a real
448   driving force for cyanobacterial blooms? Journal of Sea Research **65**:315-321.

449   Salka, I., Moulisová, V., Koblíz☐ek, M., Jost, G., Ju☐rgens, K., and Labrenz, M. (2008)
450   Abundance, Depth Distribution, and Composition of Aerobic Bacteriochlorophyll a -
451   Producing Bacteria in Four Basins of the Central Baltic Sea. Appl Environ Microbiol 74:
452   4398–4404.

453   Savchuk, O. P. 2005. Resolving the Baltic Sea into seven subbasins: N and P budgets for 1991-
454   1999. Journal of Marine Systems **56**:1-15.

455   Solórzano, L., and J. H. Sharp. 1980. Determination of total dissolved phosphorus and particulate
456   phosphorus in natural waters. Limnology and Oceanography **25**:754-758.

457   Spilling, K., E. Asmala, N. Haavisto, L. Haraguchi, K. Kraft, A.-M. Lehto, A. Lewandowska, J.
458   Norkko, J. Piiparinen, and J. Seppälä. 2022a. Dataset from a mesocosm experiment on
459   brownification in the Baltic Sea. Data in Brief **45**:108755.

460   Spilling, K., E. Asmala, N. Haavisto, L. Haraguchi, K. Kraft, A.-M. Lehto, A. M. Lewandowska, J.
461   Norkko, J. Piiparinen, and J. Seppälä. 2022b. Brownification affects phytoplankton
462   community composition but not primary productivity in eutrophic coastal waters: A
463   mesocosm experiment in the Baltic Sea. Science of The Total Environment **841**:156510.

464   Spilling, K., M.-T. Camarena-Gómez, T. Lipsewers, A. Martinez-Varela, F. Díaz-Rosas, E. Eronen-
465   Rasimus, N. Silva, P. von Dassow, and V. Montecino. 2019. Impacts of reduced inorganic
466   N: P ratio on three distinct plankton communities in the Humboldt upwelling system.
467   Marine Biology **166**:114.

Spilling, K., K. Olli, J. Lehtoranta, A. Kremp, L. Tedesco, T. Tamelander, R. Klais, H. Peltonen,
and T. Tamminen. 2018. Shifting diatom-dinoflagellate dominance during spring bloom in
the Baltic Sea and its potential effects on biogeochemical cycling. Frontiers in Marine
Science **5**:327.
Tamminen, T. 1995. Nitrate and ammonium depletion rates and preferences during a Baltic spring
bloom. Marine Ecology Progress Series **120**:123-133.
Vahtera, E., M. Laamanen, and J. Rintala. 2007. Use of different phosphorus sources by the bloom-
forming cyanobacteria *Aphanizomenon flos-aquae* and *Nodularia spumigena*. Aquatic
Microbial Ecology **46**:225-237.
Vanharanta, M., and K. Spilling. 2023. Uptake of excess phosphate at low inorganic N: P ratio in a
coastal sea afflicted with eutrophication. Marine Ecology Progress Series **718**:23-37.
Vanharanta, M., Santoro, M., Villena-Alemany, C., Piiparinen, J., Piwosz, K., Grossart, H.P.,
Labrenz, M. and Spilling, K., 2024. Microbial remineralization processes during post-
spring-bloom excess phosphate in the northern Baltic Sea. bioRxiv, doi:
10.1101/2024.02.02.577174.
Villena-Alemany, C., Mujakic, I., Fecskeova, L.K., Woodhouse, J., Auladell, A., Dean, J.,
Hanusova, M., Socha, M., Gazulla, C.R., Ruscheweyh, H.-J., Sunagawa, S., Kavagutti, V.,
Andrei, A.-S., Grossart, H.-P., Ghai, R., Koblizek, M., and Piwosz, K. (2023) Phenology
and ecological role of Aerobic Anoxygenic Phototrophs in fresh waters. bioRxiv doi:
10.1101/2023.11.17.567504.
Villena☐Alemany, C., Mujakić, I., Porcal, P., Koblížek, M., and Piwosz, K. (2023) Diversity
dynamics of aerobic anoxygenic phototrophic bacteria in a freshwater lake. Environ
Microbiol Rep 15: 60–71.
Wasmund, N., G. Nausch, B. Schneider, K. Nagel, and M. Voss. 2005. Comparison of nitrogen
fixation rates determined with different methods: a study in the Baltic Proper. Marine
Ecology Progress Series **297**:23-31.
Wasmund, N., G. Nausch, and M. Voss. 2012. Upwelling events may cause cyanobacteria blooms
in the Baltic Sea. Journal of Marine Systems **90**:67-76.

Figure legend

Fig 1. The concentration of nitrite + nitrate (NO2 + NO3), phosphate (PO4), ammonium (NH4) and

dissolved silicate (DSi) plotted against time. The error bares represent the standard error (SE, n =

3).

Fig 2. The concentration of Chlorophyll a (Chla) and abundance of heterotrophic bacteria plotted

against time. The error bares represent the standard error (SE, n = 3).

Fig 3. The concentration of particulate organic carbon (POC), nitrogen (PON), phosphate (POP),

and biogenic silicate (BSi) plotted against time. The error bares represent the standard error (SE, n

= 3).

Fig 4. The abundance of *Synechoccoccus*-like cells, picophyto(?)eukaryotes, nano- and

microphytoplankton plotted against time. The error bares represent the standard error (SE, n = 3).

Fig 5. The sum of all biovolume measurements throughout the experiment of different nano- and

microplankton groups. The presented p-value is an ANOVA statistical test comparing the different

treatments. The temporal development in biovolume is presented in Fig 6.

Fig 6. The biovolume of different nano- and microplankton groups plotted against time. The error

bars represent the standard error (SE, n = 3).

Fig 7. The abundance of different mesozooplankton groups plotted against time. The error bares

represent the standard error (SE, n = 3).

Supplementary Figures

FigS1. The temperature in the surface water (10 cm depth) throughout the experiment.

Fig S2. Salinity measurements taken from individual mesocosms and outside the mesocosms.

532   Fig 1

534   Fig 2

539   FIG 3

FIG 4

FIG 5

Fig 6

FIG 7

Supplementary Fig S1. Temperature measured every 30 s with a temperature probe placed at 50 cm

depth right outside the mesocosm bags.

566   Supplementary Fig S2. Salinity measured at 50 cm depth in the 12 experimental bags, plus outside

567   the mesocosm bags (thick red line).

571 Suppmentary Fig S3. Average AAP bacteria abundance (A) and relative contribution of AAP
572 bacteria to bacterial community (B) during the experiment. Error bars represent standard error
573 (n=3).

574