Effects of excess phosphate on a coastal plankton community: a mesocosm experiment in the Baltic Sea

Kristian Spilling[1,2, *], Mari Vanharanta[1,3], Mariano Santoro[4,5], Cristian Villena-Alemany[6], Matthias Labrenz[4], Hans-Peter Grossart[9,10], Kasia Piwosz[11]

[1]Marine and Freshwater Solutions, Finnish Environment Institute, 00790 Helsinki, Finland

[2]Centre for Coastal Research, University of Agder, 4630 Kristiansand, Norway

[3]Tvärminne Zoological Station, University of Helsinki, 10900 Hanko, Finland

[4]Department of Biological Oceanography, Leibniz Institute for Baltic Sea Research Warnemünde - IOW, Rostock, Germany

[5]Department of Plant Physiology, Institute for Biosciences, University of Rostock, Rostock, Germany

[6]Laboratory of Anoxygenic Phototrophs, Institute of Microbiology, Czech Academy of Sciences, 379 01 Třeboň, Czech Republic

[9]Leibniz Institute for Freshwater Ecology and Inland Fisheries, Germany

[10]Institute of Biology and Biochemistry, Potsdam University, Germany

[11]National Marine Fisheries Research Institute – NMFRI, Gdynia, Poland

*Corresponding author: kristian.spilling@syke.fi

Key words: biogeochemical cycles, coastal seas, eutrophication, nutrient stoichiometry, filamentous cyanobacteria, ocean deoxygenation

Running head: Effects of excess phophate in the Baltic Sea

Abstract

Eutrophication in the Baltic Sea has caused an imbalance in the inorganic nitrogen (N) to

phosphorus (P) ratio, leaving excess phosphate (PO_4) after the phytoplankton spring bloom that

terminates after N-depletion. Using monitoring data, we demonstrated that the PO_4 concentration

has continued to increase in the outermost Gulf of Finland during past decades. We further

investigated the fate of such excess PO_4 in a two-week mesocosm (1.2 m^3) experiment. The starting

concentration of PO_4 was 0.66 µM, and treatments included a non-treated control (control), nitrate

addition (N-add; 3.6 µM), glucose addition (C-add; 25 µM) and combined nitrate and glucose

addition (N+C-add). The addition of N both in N-add and N+C-add treatments stimulated nano- and

microphytoplankton, while the picophytoplankton abundance increased only after N-depletion.

Also, the copepod biomass was positively affected by the N-addition. N_2-fixing cyanobacteria were

present but in low abundance. Carbon addition did not enhance heterotrophic bacterial uptake of

PO_4 contrary to our expectations, nor did it affect the phyto- or zooplankton community

composition. The PO_4 concentration was reduced to ~0.4 µM in the control and C-add treatments

and to 0.16 µM in the two N-amended treatments, with an inorganic N:P uptake ratio of 6.7. These

results underscore the role of picophytoplankton in reducing the excess PO_4 pool after the spring

bloom, a function traditionally ascribed to bloom-forming diazotrophic cyanobacteria in the Baltic

Sea.

Introduction

Deoxygenation, i.e. loss of oxygen, in the ocean is an increasing global problem that has direct implications not only for aerobic organisms but also for biogeochemical cycles (Keeling et al. 2010). There are two primary causes of the deoxygenation of the ocean: global warming, and human input of nutrients to marine ecosystems leading to eutrophication. Warming of surface waters happens on a global scale and reduces gas solubility, including oxygen. Eutrophication is a local or regional environmental problem that causes algal blooms that typically aggravate oxygen consumption in deeper water layers after the bloom, when the biomass is decomposing while sinking. In relatively shallow coastal seas, such elevated oxygen consumption may cause hypoxic and anoxic conditions on the seafloor, where sediment microbial processes typically decrease bioavailable nitrogen through denitrification and anammox (Hannig et al. 2007). At the same time, anoxia increases the release of phosphate from the sediment through the chemical reduction of iron-bound phosphate. The net result of anoxic bottoms is often a decrease in bioavailable nitrogen and an increase in phosphate concentration, causing the inorganic N:P ratio to decrease in several parts of the world (Conley et al. 2002, Hauss et al. 2012, Kalvelage et al. 2013, Meyer et al. 2016).

The Baltic Sea has been affected by eutrophication for several decades, which has increased the area of anoxic seafloor (Conley et al. 2002). In the Baltic Proper and the Gulf of Finland, the most productive period with accompanying flux of organic material to the sea floor is during the spring bloom, which in these subbasins terminate when dissolved inorganic nitrogen (DIN) has been depleted. There was a regime-like shift in the mid-1990s when the dissolved inorganic phosphorus (DIP) pools started to be at elevated levels after the spring bloom; low phosphate concentrations were generally detected during early summer before the mid-1990s, though N-limitation of the spring bloom could be already demonstrated at that time (Tamminen 1995). The decreasing inorganic N:P ratio is caused primarily by a release of phosphate from the seafloor after turning anoxic (Conley et al. 2002). At present the surplus of phosphate after the spring bloom typically range from 0.1 up to 0.6 μmol PO$_4$ L^{-1} in the Gulf of Finland and the Baltic Proper (Spilling et al.

2018). However, there is relatively little data published in recent years, making it challenging to

ascertain whether this trend persists.

The excess phosphate is believed to directly benefit N_2-fixing cyanobacteria that often create toxic

or nuisance blooms during the summer (Larsson et al. 2001, Kangro et al. 2007), supported by

model work (Munkes et al 2021). *Nodularia spumigena* and *Aphanizomenon flos-aquae* are

amongst the typical dominating filamentous cyanobacteria, forming blooms in July and August in

large parts of the Baltic Sea. These species are N_2-fixers and can take up PO_4 and store it in the

form of polyphosphate. However, they grow relatively slowly in water temperatures below 15°C

(although *A. flos-aquae* is less sensitive to cold water than *N. spumigena*), and typically start

forming blooms only after the excess phosphate pool has been depleted (Lehtimäki et al 1997,

Wasmund 1997).

An alternative pathway of the excess P pool is going through other primary producers or

heterotrophic bacteria (Nausch and Nausch 2004), thereby reducing the available P for the

filamentous cyanobacteria. However, the prevailing N-limitation restricts the growth of other

phytoplankton, whereas heterotrophic bacteria, that may also take up P, are typically limited by

carbon during late spring (Lignell et al. 1992). Organic carbon produced by the spring bloom

phytoplankton triggers a bacterial response, dependent on the composition of the phytoplankton

community, but this labile carbon is rapidly consumed (Camarena-Gómez et al 2018, 2019). After

this labile carbon is depleted, there remains a large (compared to other coastal ecosystems)

refractory pool of dissolved organic carbon due to the high freshwater influence in the Baltic Sea

bringing in e.g. humic substance (Hoikkala et al 2015).

Currently, it remains unclear which organisms take up the excess phosphate, most of which seems

to settle out of the euphotic zone before cyanobacterial blooms develop (Nausch et al. 2008).

Consequently, other P sources must exist for these cyanobacteria (Raateoja et al. 2011), most likely

through upwelling, turbulent mixing events, or regeneration of organic P (Raateoja et al 2011,

Wasmund et al. 2012).

Recently, we demonstrated in a small-scale (20 L) experiment that the post-spring-bloom plankton

community could remove excess phosphate, even without any N-addition, but with a clear effect of

temperature that affected both the rate of phosphate uptake and the plankton community

composition (Vanharanta and Spilling 2023). Picophytoplankton seems to play a major part in the

drawdown of phosphate, but an open question is the role of heterotrophic bacteria that can be

competing with phytoplankton for this resource (Lignell et al. 1992, Vanharanta and Spilling 2023).

Here, we present an updated dataset of phytoplankton pigments and inorganic nutrients from the

Finnish monitoring program taken in the Gulf of Finland, together with an experiment where we

addressed the potential of an early-summer plankton community to remove the excess phosphate.

The experiment was done in larger scale mesocosms closer to natural conditions than our previous

indoor tank experiment (Vanharanta and Spilling 2023). In addition to a non-treated control, we

added bioavailable nitrogen (N-add), carbon (C-add) and both (N+C-add), to stimulate primary

producers and heterotrophic bacteria. We hypothesized that picophytoplankton would become

relatively more important after N-depletion, taking up part of the excess phosphate, but that

heterotrophic bacteria can also exploit the inorganic P pool when they have access to a labile carbon

source.

Materials and methods

Field data

Field data was obtained from the Algaeline data constituting two commercial ships passing through

the Gulf of Finland (Helsinki - Travemünde, and Helsinki – Stockholm, Supplementary Figure S1).

The data consists of discrete samples automatically taken from a flow through system on the return

to Helsinki where they were collected and analyzed on the same day. The variables measured

include Chlorophyll a (Chla) concentration and inorganic nutrients. For Chla, GF/F filters

(Whatman, UK) were used and the Chla was extracted in EtOH, stored at -20°C and in the darkness

for up to two months, and measured with a spectrofluorometer (Varian Cary Eclipse, Agilent,

USA), calibrated against known standards (Merck, Sigma-Aldrich), with excitation and emission

wavelengths of 430 and 670 nm, respectively (Jespersen and Christoffersen 1987). Inorganic

nutrients were determined using standard colorimetric methods (Grasshoff et al. 1999). The ship-of-

opportunity monitoring data are available from 1998 until present. In addition, fluorescence sensor

data in the flow-through system exists from 2013.

Mesocosm experiment

The experiment was set up in 1.2 m^3 mesocosm bags (Ø = 90 cm, length = 190 cm) that were

moored outside the Tvärminne Zoological Station, Gulf of Finland (59° 50' 40" N, 23° 14' 57" E).

The bags were attached to a floating platform (Spilling et al. 2022a). In total, 14 bags were installed

but the two outermost bags were not used. These 'dummy' bags enabled to create the same light

conditions for the outermost experimental units. All units were covered to prevent rain or bird

droppings from falling into the bags. The spring bloom in the area had peaked in April and the PO_4

concentration was relatively low at the onset of the experiment (0.013 µ mol L^{-1}), thus we added

PO_4 to all units to a final concentration of 0.66 µ mol L^{-1}. The experimental setup consisted of an

unamended control, a treatment with added nitrate (N-add; 3.6 µmol N L^{-1}), a treatment with added

carbon in the form of glucose (C-add, 36 µmol C L^{-1}), and a treatment with both nitrate and glucose

addition (N+C-add). The rationale behind the treatments was to stimulate the phytoplankton with

nitrate and heterotrophic bacteria with glucose. Each treatment had three replicates, and in total 12

bags were used for the experiment.

The bags were filled with surface water using gentle pumping on 7th June 2021, which was

considered day -1 of the experiment. This filling method had visually (microscope) been tested and

found not to affect zooplankton but prevents small fish from entering the mesocosms. The

following day (day 0), nutrients and carbon sources were added in the morning and the first full

sampling took place right after the addition. The temperature was approximately 18°C and the salinity was 5.5 at the start of the experiment (Supplementary Figs S2 and S3). Water samples were taken daily with a Limnos water sampler (Hydro-Bios, Germany) from 0.5 m depth from the middle of the bags for fluorescence measurements and flow cytometer counts. CTD profiles were also taken daily from the bags and outside the experimental units, and we had dissolved oxygen loggers (HOBO U26; Onset Inc. USA) mounted inside each mesocosm bag. Rate of change in dissolved oxygen was calculated by linear regression during daytime (09 – 17) and nighttime (23 – 03). In addition to the daily sampling (e.g. CTD profile), larger volumes (5 L) were taken on regular sampling days: 0, 1, 3, 6, 8,10, 13 and 15. During these days, additional variables such as inorganic nutrients and particular organic carbon, nitrogen and phosphorus were measured.

Total Chla fluorescence was determined from each mesocosm bag by daily use of a handheld fluorometer (AquaPen, Photon Systems Instruments, Czech Republic), and direct Chla measurements were carried out by filtration in duplicates on regular sampling days, and analyzed as described above.

We used standard colorimetric methods (Grasshoff et al. 1999) using a photometric analyzer (Aquakem 250, Thermo Scientific, USA) to measure dissolved inorganic nutrients: nitrite + nitrate ($NO_2 + NO_3$), phosphate (PO_4) and dissolved silicate (Dsi). Ammonium (NH_4) was measured separately using a spectrophotometer (U-1100, Hitachi, Japan).

Particulate organic carbon (POC), nitrogen (PON) and phosphorus (POP) were determined from duplicate water samples filtered onto acid-washed (2M HCl for 15 min, then rinsed carefully with ultrapure water) and pre-combusted (450°C, 4 h) GF/F filters (Whatman, UK). POC and PON were measured with an element analyzer coupled with a mass spectrometer (Europa Scientific ANCA-MS 20-20 15N/13C, UK). POP was determined according to Solórzano and Sharp (1980) as modified by Koistinen et al. (2017). Samples for biogenic silicate (BSi) were filtered on

polycarbonate membrane filters (0.8 µm, GVS, Italy) and measured according to Koistinen et al.

(2018).

Abundance of nano- and microplankton (phytoplankton and microzooplankton) was determined

with the FlowCam (Fluid Imaging, Yokogawa) from preserved (acid Lugol's solution) samples.

The samples were stored in a fridge and counted within a year from the sampling. We used the 10x

magnification and 100 µm flowcell during the runs and the accompanying Visual Spreadsheet

software for automatic calculation of biovolume. Further identification and visual inspection were

done with an inverted microscope (Letiz Labovert) using a 40x objective.

Flow cytometer counts of live samples were done daily with a Sysmex (Japan), Partec – Cube 8

equipped with two lasers (488 and 561 nm), two scattering (forward and side) and three

fluorescence detectors (610/30; 661/16; and 670/40, corresponding to the detection of

phycoerythrin, phycocyanin, and Chla, respectively). The trigger was set on Chla fluorescence (670

nm). On each sampling day, we made several measurement-runs with beads and blanks to ensure

correct particle counts. Size fractionations (0.8, 1, 2, 5, 10, and 20 µm filters) were also used to

identify the approximate size of different phytoplankton groups. The scattering and fluorescence

properties were used to gate different phytoplankton groups using the FCS Express 6 software. Four

groups were identified: picoeucaryotes: size <2 µm with only Chla, Cryptophyte-like >2 µm with

Chla and phycoerythrin fluorescence, nanophytoplankton size 2–20 µm and microphytoplankton:

size >20µm with only Chla fluorescence.

Synechococcus-like cells, <2 µm with phycocyanin fluorescence, were counted at the same time as

heterotrophic bacteria, using a LSR II, BD flow cytometer (Biosciences, USA) equipped with a 488

nm laser. These samples were fixed with 1 % paraformaldehyde (final concentration) for 15 min in

darkness, flash frozen in liquid nitrogen, and stored at -80°C until analysis. Before measurements,

samples were thawed and stained with SYBRGreen I (Molecular Probes, Eugene, OR, USA) at a

10^{-4} (v/v) concentration and incubated for 15 min in the darkness before counting. We used

CountBright beads (Molecular Probes) in each sample to determine the measured volume.

FACSDiva Software (BD Biosciences) and Flowing Software version number 2.5.1. were used to

gate and obtain the cell counts.

Zooplankton was collected three times, at the start, middle and end of the experiment. Samples were

taken with a plankton net (25 cm diameter, 50 µm mesh size) carefully dragged from the bottom of

the mesocosm bags to the surface. Samples were immediately preserved with acid Lugol's solution

and stored at 4°C. Zooplankton was identified to genus level from images created using a flatbed

scanner and analyzed at 20 x magnification.

Data analysis

A one-way analysis of variance (ANOVA) was performed to test whether significant effects could

be revealed by the treatments. Tukey's Honest Significant Difference (HSD) *post hoc* test was

performed to compare different treatments and assess the significance of differences between them

after ensuring normality and homogeneity of the residuals. In case of the data following a non-

normal distribution, the ANOVA was performed on ranks followed by Dunn's *post hoc* test. All

statistical tests were performed using SigmaPlot 15 software (Systat Software).

Results

Field data

From the Gulf of Finland monitoring data, the Chla fluorescence had a clear peak in spring at

around April 20 (Day 110), followed by a clear water period after mid-May (Supplementary Fig

S4). Phycocyanin fluorescence was more variable, elevated levels of phycocyanin fluorescence

started to appear after mid-May (around May 19, Day 140) but the regression we used indicated a

peak around mid-July (July 14, Day 195), when the average temperature reached 15°C.

From the discrete samples, the Chla peak during spring was around 20 µg Chla L^{-1}, with some

observations of even higher concentrations, up to 33 µg Chla L^{-1} (Supplementary Fig S5). Nitrate

was rapidly depleted during spring, with very low concentrations after the Chla peak. Phosphate

was also reduced, but often not completely depleted, and there was more phosphate in the water

during summertime (Day 138-220) in the 10-year period 2014-2023 compared with the other

periods 1998-2003 and 2004-2013 (Dunn; $p < 0.002$). There was no difference in phosphate

concentration between the two earliest periods (Dunn; $p > 0.9$). The concentration of dissolved

silicate decreased during spring but never completely depleted, and the concentration was higher in

the past decade (2014 - 2023) compared to the previous one (2004 – 2013; Dunn $p < 0.001$).

Mesocosm experiment

In the experiment, there was a clear effect of N addition, but no effect of carbon addition, on the

phosphate concentration (Fig 1). The nitrate added to the N-add and N+C-add treatments was

depleted already on day 3. The phosphate concentration decreased from ~0.66 µmol L^{-1} at the start

of the experiment to ~0.16 µmol L^{-1} in the two treatments with N added on Day 8, resulting in an

inorganic N:P uptake ratio of 6.7. In the control and C-add treatments, the PO_4 concentration was

0.33 and 0.36 µmol PO_4 L^{-1}, respectively, at the end of the experiment. There was residual

ammonium with a concentration in the range of 0.15 to 0.35 µmol L^{-1} throughout the entire

experiment. Dissolved silicate was constant in the control and C-add but decreased slightly towards

the end of the experiment in the N-add and N+C-add treatments, with an uptake of approximately

1.9 µmol L^{-1} (Fig 1).

The phytoplankton biomass declined from the onset of the experiment in the control and C-add

treatments but with a delay of two days in the two N amended treatments (Fig 1). The initial Chla

concentration was 3.5 µg L^{-1} and it decreased to 1–2 µg Chla L^{-1} and within two days in the control

and C-add treatments (Fig 2). After reaching a minimum of <1 µg L^{-1}, the Chla concentration

increased again at the end of the experiment to ~2 µg Chla L^{-1}.

Oxygen dynamics indicated more O_2 was added during daytime in the N-add treatment compared

with the N+C add treatment during the first three days (Supplementary Fig S6). During nighttime,

the loss of O_2 was higher in the N-add and N+C add treatments compared to the control and C-add

treatments at the start of the experiment, but O_2 dynamic was similar in all treatments after day 7

(Supplementary Fig S6).

Heterotrophic bacteria showed the same dynamic in all treatments (Fig 2); their abundance

increased after day 1, with a peak at day 8. The overall bacterial abundance was higher in the N-add

treatment compared to the control and C-add treatments (Tukey: $p = 0.001$ and $p = 0.003$

respectively), the rest of the treatments were similar (Tukey; $p > 0.14$).

All particulate nutrients decreased in the experimental units during the first week, but the decrease

was more rapid in treatments without any nitrate addition (control and C-add), except for biogenic

silicate (BSi) where the decrease was similar in all treatments (Fig 3). Stoichiometric ratios were

relatively stable with an average C:N:P ratio of 140:14:1, and without any differences between

treatments (ANOVA; $p > 0.1$).

The temperature and salinity throughout the experiment are presented in Supplementary Figs S2

and S3. There was a drop in temperature after day 6 from ~18°C to ~14°C due to upwelling,

followed by increase towards the end of the experiment to ~20°C. Salinity varied outside the bags

but stayed constant within the bags with no sign of mixing with the outside water indicating that our

bags remained intact during the experiment.

Plankton community composition

Nitrate additions benefited micro- and nanophytoplankton that increased in abundance in both the

N-add and N+C-add treatments (Fig 4), contributing to a high Chla concentration over the first

three days. This was also evident from the FlowCam images, where, *e.g.,* the cell concentration and

total biovolume of dinoflagellates >10 µm increased in these two treatments until nitrate had been

depleted (supplementary Fig S7). The green algae *Monoraphidium* sp. (nanophytoplankton)

increased also in the N-amended treatments, but kept growing after N depletion, reaching a

maximum biovolume at day 8 (Fig 5).

After nitrate depletion, picoeukaryotes started to increase in abundance, especially in the control

and N-add treatment, but also, to a lesser extent, in the N+C-add and C-add treatments (Fig 4).

Synechococcus-like cells decreased in all treatments after the start of the experiment but increased

again after day 3 and more so in the control and N-add treatments (Fig 4), peaking on day 8 when

the abundance was clearly higher in the control and N-add treatments compared with the carbon

amended treatments (Tukey p <0.014). *Planktothrix* sp., and *Pseudoanabaena* sp. were the main

component of colonial and filamentous cyanobacteria. They peaked in the N-add and N+C-add

treatments on day 3 and 6, respectively, with the highest peak in biovolume in the N+C-add

treatment (Fig 5). *Aphanizomenon flos-aquae* and *Nodularia spumigena* were present but

contributed combined less than 5% to the total cyanobacterial biovolume.

Microzooplankton consisted of heterotrophic dinoflagellates and ciliates (Fig 5). Heterotrophic

dinoflagellates increased in the N-add and N+C-add treatments with a peak on day 8 before

declining to a similar biovolume as in the control and the C-add treatments at the end of the

experiment (Fig 5). The biovolume of ciliates was highly variable between replicates, and although

the average biovolume was higher in the two N-add treatments, there was no consistent difference

between the treatments (ANOVA, p = 0.767).

The mesozooplankton community was dominated by copepods, cladocerans and pelagic stages of

Amphibalanus sp. (Fig 6). The number of copepods was highest in both N-add and N+C-add

treatments towards the end of the experiment including both copepodite and nauplii stages

(ANOVA, p = 0.024 and p = 0.017, respectively). Among cladocerans, the abundance of *Bosmina*

sp. was positively affected by the nitrate addition (ANOVA, p = 0.04), whereas *Podon* sp. (Fig 6)

and *Evadne* sp. (data not shown) both decreased over time and had completely disappeared in all

treatments at the end of the experiment. The difference in *Amphibalanus* nauplii abundance between

treatments was not clear (ANOVA, p = 0.06), but cypris larvae had only developed in the two N-

amended treatments at the end of the experiment (Fig 6).

Discussion

Here we demonstrate that the phosphate concentration was higher in the past decade compared to

earlier periods in the Gulf of Finland, suggesting that, despite a considerable reduction in the

external nutrient load over past decades, it is still increasing. The reason for the high phosphate

concentration is the release from the sediment (Pitkänen et al 2001). The effect of short-term shifts

in the inorganic N:P ratio depends to some extent on the plankton community composition, but

tends to be reflected in the organic N:P ratio (Spilling et al. 2019, Bach et al. 2020). Yet, potential

long-term consequences of this phenomenon are still not known. A decreasing inorganic N:P ratio

during N-depletion, as we found here, is often caused by reduced O_2 concentrations in different

ecosystems. However, it is not a universal process, as there are examples of other processes that

increase N:P ratio, for example, by more effective P-removal from wastewater, increased use of N-

rich fertilizer, or increased atmospheric N-deposition (McQuatters-Gollop et al. 2007, Kim et al.

2014). Despite all the efforts to reduce external loading to the Baltic Sea, the increasing excess

phosphate concentration after N-depletion demonstrates that improving ecological state takes a long

time to achieve, and arguably longer than it takes to deteriorate the ecosystem.

We also found that dissolved silicate had increased during the active growth season for algae. This

is somewhat surprising as the DSi concentration has decreased long term in the Baltic Sea, linked to

a higher retention of silicate by damming of rivers and increased burial of biogenic silicate in the

sediment due to eutrophication (Conley et al 2008). Increases in dissolved silicate could therefore

be interpreted as a positive sign of slightly improving conditions (Wasmund et al 2017), but further

studies are needed to investigate whether this is a general trend in the Baltic Sea.

Temporal plankton community development in the experiment

Inorganic nutrients initially stimulated nano- and microphytoplankton, in particular the chlorophyte

Monoraphidium sp. and autotrophic dinoflagellates. This was expected, as they have a larger

volume for internal storage, i.e. luxury uptake of nutrients. In addition, *Monoraphidium* sp. is

known to have a relatively high growth rate (Seip and Reynolds 1995). Nonetheless, this nutrient

effect only delayed the gradual decline of phytoplankton biomass. After inorganic nutrients had

been depleted, there was a clear increase in picophytoplankton abundance, which has been observed

in previous mesocosm experiments in this area (Crawfurd et al 2017, Spilling et al. 2022b). Smaller

cells have a larger surface-to-volume area and are better adapted to low inorganic nutrient

concentrations, which is the likely reason for this group to dominate during periods when easily

available nutrient sources have been depleted by larger cells.

The mesocosm bags prevent effective wind-driven mixing, although there was some temperature

driven advection. There was particulate organic matter sedimentation during the experiment, but

unfortunately, we could not sample the bottom part of the mesocosms to quantify it. This export

loss might have had some consequences for the phytoplankton community composition as non-

motile groups, such as diatoms, likely sedimented out quicker than motile flagellates. This could be

deduced from the development in BSi over time, and diatoms were a minor group in terms of

biovolume.

As expected, mesozooplankton required a longer time to respond to the experiment, yet, an effect of

the nutrient addition on its biomass was measurable (Fig. 6). The bags were not very deep (1.9 m)

and the available organic material in the bottom likely supported the community of

mesozooplankton that could easily move vertically between different water depths, especially since

higher predators were absent. This is a likely reason for a treatment effect on copepods and

Bosmina sp, even though the available, suspended food items were similar in all bags after the

drawdown of nitrate. Some groups, such as the cladoceran *Evadne* sp. disappeared over time, which

could be due to an enclosure effect as this cladoceran often tend to disappear in mesocosm

experiments (e.g. Graneli and Turner 2002).

Treatment effects

The main driver for the observed differences between treatments was the addition of nitrate that

directly affected the phytoplankton biomass. The carbon addition, on the other hand, had no effect

on particular organic nutrients, nor on bacterial abundance, suggesting that the total amount of

added labile carbon was relatively low compared to the extra phytoplankton-derived carbon in the

N-amended treatments. Interestingly, however, carbon addition did have a negative effect on

picoeukaryotes and *Synechococcus*-like cells relative to the other treatments. This could have been

due to competition for nutrients with heterotrophic bacteria, which likely got a boost due to the

sudden, easily available carbon source that was added. This was, however, not evidenced by the

abundance of heterotrophic bacteria that remained similar between control and C-add treatments.

Increased grazing pressure could have played a role, but no noticeable differences in protozoan or

microzooplankton abundance were observed and grazing would be expected to affect autotrophic

and heterotrophic picoplankton equally (Kuosa and Marcussen 1988). Thus, the underlying reason

for the apparent carbon effect on picophytoplankton still remains unclear.

Carbon addition also affected the O_2 dynamics during the first three days of the experiment. There

was not as much increase in O_2 during the daytime in the N+C compared to the N-add treatments,

which likely was due to higher primary production in the latter treatment. The N amended

treatments revealed an increase in biomass and O_2 reduction rate during nighttime suggesting a

higher O_2 loss due to respiration than in the control and C-add treatments. Any treatment effects on

O_2 dynamics disappeared after the first week of incubation, indicating similar biological activity

between all treatments during the final week of the experiment. The rate of change in O_2 is not

directly transferable to biological activity, however, as temperature fluctuations affect gas

solubility, and there were occasions of both O_2 out-gassing and in-gassing due to changes in

temperature.

Phosphate removal

In the N-add and N+C-add treatments, ~75 % of the phosphate was removed, whereas in the control

and C-add treatments ~ 40 % was removed within a week. Approximately 15 % of the phosphate

pool, or 20–30 % of the PO_4 taken up was released into the dissolved organic phosphorus (DOP)

pool, but DOP concentration decreased again during the last week of the experiment (Vanharanta et

al 2024), suggesting the added DOP was relatively labile and microbially available. The N:P uptake

ratio of 6.7, less than half the Redfield ratio of 16, suggests an active luxury uptake of P of the

plankton community. The decrease in the phosphate pool was not seen as an increase in the

particulate phosphorus fraction although the N-amended treatments had a higher concentration of

particulate organic elements. This was likely due to the export of particulate organic matter to the

bottom of the mesocosm bags.

In a recent study, using smaller 20 L tanks, we found that the excess phosphate had been depleted

completely under N-deficiency, even without N addition (Vanharanta and Spilling 2023). This

previous experiment had a longer duration (35 d) than the one presented here, with a drawdown of

phosphate in the control units of 20 % to 50 % (from a starting concentration of 0.55 µM) after 15

days, depending on the temperature (Vanharanta and Spilling 2023). Thus, it is likely that we would

have also seen a complete drawdown of phosphate, if the current experiment would have lasted a

few weeks more. In addition, the decrease of temperature by several degrees in the middle of the

experiment likely decelerated microbial activities and consequently phosphate uptake rates.

Possibly, more of the excess phosphate would have been taken up if temperature had stayed stable

during the whole duration of the experiment.

The addition of glucose did stimulate bacterial productivity and extracellular enzyme activity

(Vanharanta et al. 2024a), but there was no indication of any additional drawdown of phosphate by

heterotrophic bacteria when stimulated by glucose compared to the control. Also, in this respect, the

combined N+C addition behaved like the N addition. This further supports that most phosphate

uptake was due to picophytoplankton with little contribution of heterotrophic bacteria. The

extracellular enzymatic activity of free-living bacteria likely served to acquire carbon from

polymers released by phytoplankton which was demonstrated by higher cell-specific activities of

proteolytic and glycolytic enzymes during the bacterioplankton peak in the N-addition when

compared to other treatments (Vanharanta et al. 2024a). This finding demonstrates the dependency

of heterotrophic bacteria on phytoplankton and could explain why carbon addition alone did not

result in significantly higher phosphate uptake compared to the control. The timing of the

heterotrophic bacterial abundance peak is typical following the spring bloom when there is lots of

organic matter available (Camarena-Gómez at al. 2018), and the bacterial community is to some

extent affected by the phytoplankton community during the spring bloom (Camarena-Gómez et al.

2021).

Link to cyanobacterial blooms in the Baltic Sea

One of the most prominent effects of eutrophication in the Baltic Sea is the intensifying occurrence

of diazotrophic cyanobacterial blooms during summer. The prevailing N-limitation leads to a low

inorganic N:P ratio, which favors N_2-fixing cyanobacteria during warm, calm summer months (Niemi

1979, Wasmund et al. 2005). These cyanobacterial blooms amplify the eutrophication problem, as

the amount of cyanobacterial fixed N is comparable to the anthropogenic N-loading in parts of the

Baltic Sea (Savchuk 2005, Wasmund et al. 2005). The cyanobacterial input of N might thus create a

self-reinforcing cycle through a positive feedback loop, i.e. N_2-fixation increases the biomass load

reaching the sediment, which in turn raises oxygen consumption and releases additional P into the

ecosystem (Vahtera et al. 2007a, Spilling et al 2018).

In our experiment we categorized the different cyanobacteria present in three functional groups. The

first group consists of small, single-celled cyanobacteria, here termed *Synechococcus*-like cells,

which do not fix nitrogen and are presumably functionally closer to picoeukaryotes than to the

larger N_2-fixing filamentous species. The second group includes the relatively small filamentous

cyanobacteria *Planktothrix* sp. and *Limnothrix* sp. plus the larger *Pseudoanabaena* sp. Common for

all of them is that they do not fix N_2. The third functional group comprises the relatively large N_2-

fixing filamentous cyanobacteria *Aphanizomenon flos-aquae* and *Nodularia spumigena*. A surplus

of P would benefit the latter, especially if they have been able to increase their biomass during the

nitrate depletion phase. Although these cyanobacteria were present in all treatments, there was no

indication that these typical bloom-forming diazotrophic species proliferated (staying below 5% of

the cyanobacterial biomass). Rather, the additional P uptake after N depletion seems to have been

driven by an increase in picoeukaryotes, *Synechococcus*-like cells and possibly the smaller (non-N_2-

fixing) filamentous cyanobacteria. These organisms can take up phosphate and store excess

phosphorus as polyphosphate (Jentsch et al. 2023). This was also observed under the microscope,

although we lack quantitative data on polyphosphate storage. *Planktothrix* sp., and *Pseudoanabaena*

sp. species are common in the area during summer, but do not form extensive blooms.

Aphanizomenon flos-aquae and *Nodularia spumigena* have different P uptake strategies. *A. flos-*

aquae can take up and store excess P for later growth and also take up organic forms of phosphorus,

whereas *N. spumigena* more commonly takes up phosphate and, to some extent, relies more on

recycled P after phosphate depletion (Hagström et al. 2001, Vahtera et al. 2007, Schoffelen et al

2022). Although both species were present in the experiment, they were at exceptionally low

concentrations. Other filamentous cyanobacteria (*Planktothrix* sp., and *Pseudoanabaena* sp.) made

up the bulk of the measured cyanobacterial biovolume. The lack of a typical cyanobacterial

community composition could be due to an enclosure effect, but both. *A. flos-aquae* and *N.*

spumigena have relatively low growth rates (Reynolds 1984, Vahtera et al. 2005), and it is likely

that they did not grow fast enough in the experiment to produce much biomass in the relatively

short time span of the experiment. This is similar to the typical natural development where the

excess phosphate typically has been depleted before the onset of filamentous cyanobacterial blooms

later in summer. Our experimental results support our hypothesis that picophytoplankton plays a

key role in the uptake of the excessive phosphate after the spring phytoplankton bloom. This would

indicate that in contrast to current models/assumptions, the excess phosphate remaining after the

spring bloom do not directly benefit the typical bloom forming cyanobacteria (*A. flos-aquae* and *N.*

spumigena), which would rather rely on recycled or upwelled sources of phosphate.

In conclusion, we found that N-addition stimulated both nano- and microphytoplankton, and the

picophytoplankton abundance increased following nitrate depletion. Based on these results, we

conclude that picophytoplankton incorporate substantial parts of the excess P, which strengthen our

initial hypothesis. Yet, picophytoplankton was unable to completely deplete the inorganic P-pool

during the relatively short experiment. Most of the organic P formed was likely exported by the

overall decrease in total phosphorus combined with a negligible change in the DOP pool between

the start and the end of the experiment. Carbon addition, in contrast, stimulated heterotrophic

bacteria production but did not increase bacterial uptake of excess phosphate, contrary to our

expectations. The typical bloom-forming cyanobacteria in the Baltic Sea, considered to greatly

benefit from excess phosphate, were present but not abundant in our mesocosms. Our results

underscore the key role of picophytoplankton in reducing the excess phosphate pool after the spring

bloom, a function traditionally ascribed to bloom-forming diazotrophic cyanobacteria in the Baltic

Sea.

Acknowledgements

We would like to thank the staff at Tvärminne Zoological Station for all their support, technical

assistance, and nutrient measurements. In particular Joanna Norkko, Jaana Koistinen, Mervi

Sjöblom, Kia Rautava and field technicians Göran Lundberg and Jostein Solbakken. We would also

like to thank Sebastian Ehrhart for help with the monitoring data.

This study used equipment part of the Finnish marine research infrastructure (FINMARI)

consortium.

This work was supported by the Transnational Access program of the EU H2020-INFRAIA project

(No. 731065) AQUACOSM - Network of Leading European AQUAtic MesoCOSM Facilities

Connecting Mountains to Oceans from the Arctic to the Mediterranean - funded by the European

Commission. Additional funding came from Walter and Andree de Nottbeck foundation (KS and

MV) and the Research Council of Finland (KS, decision no 354272). KP was supported by the

National Science Centre, Poland under the Weave program (project no 2021/03/Y/NZ8/00076).

HPG was funded by the German Science Foundation (DFG) project PycnoTrap (GR1540/37-1). MS

was supported by Leibniz Science Campus Phosphorus Research Rostock in the funding line

strategic networks of the Leibniz Association.

Data availability

Data from all measured variables during the mesocosm experiment are available on PANGAEA

(Vanharanta et al. 2024b).

References

Bach, L. T., and others. 2020. Factors controlling plankton productivity, particulate matter stoichiometry, and export flux in the coastal upwelling system off Peru. Biogeosciences 17, 4831–4852.

Camarena-Gómez M.T., Ruiz-Gonzalez C., Piiparinen J., Lipsewers T., Sobrino C., Logares R., Spilling K. 2021. Bacterioplankton dynamics driven by interannual and spatial variation in diatom and dinoflagellate spring bloom communities in the Baltic Sea. Limnol Oceanogr. 66: 255-271

Camarena-Gómez, M.T., Lipsewers T., Piiparinen J., Eronen-Rasimus E., Perez-Quemaliños D., Hoikkala L., Sobrino C., Spilling K. 2018 Shifts in phytoplankton community structure modify bacterial growth, production and composition. Aquat Microb Ecol. 81: 149-170.

Conley, D. J., C. Humborg, L. Rahm, O. P. Savchuk, and F. Wulff. 2002. Hypoxia in the Baltic Sea and basin-scale changes in phosphorus biogeochemistry. Env Sci Technol 36:5315-5320.

Conley, D.J., Humborg, C., Smedberg, E., Rahm, L., Papush, L., Danielsson, Å., Clarke, A., Pastuszak, M., Aigars, J., Ciuffa, D. and Mörth, C.M., 2008. Past, present and future state of the biogeochemical Si cycle in the Baltic Sea. J Mar Sys 73: 338-346.

Crawfurd, K. J., Alvarez-Fernandez, S., Mojica, K. D. A., Riebesell, U., Brussaard, C. P. D. 2017. Alterations in microbial community composition with increasing fCO$_2$: a mesocosm study in the eastern Baltic Sea, Biogeosciences, 14, 3831–3849

Díez, B., Van Nieuwerburgh, L. and Snoeijs, P., 2013. Water nutrient stoichiometry modifies the nutritional quality of phytoplankton and somatic growth of crustacean mesozooplankton. *Mar Ecol Prog Ser 489*, 93-105.

Goldenberg, S.U., Spisla, C., Sánchez, N., Taucher, J., Spilling, K., Sswat, M., Fiesinger, A., Fernández-Méndez, M., Krock, B., Hauss, H. and Haussmann, J., 2024. Diatom-mediated food web functioning under ocean artificial upwelling. *Scientific Reports*, *14*(1), p.3955.

Granéli, E. and Turner, J.T., 2002. Top-down regulation in ctenophore-copepod-ciliate-diatom-phytoflagellate communities in coastal waters: a mesocosm study. *Mar Ecol Prog Ser 239*, 57-68.

Grasshoff, K., K. Kremling, and M. Ehrhardt. 1999. Methods of seawater analysis 3 ed. Wiley-VCH, Weinheim.

Hagström, Å., F. Azam, J. Kuparinen, and U.-L. Zweifel. 2001. Pelagic plankton growth and resource limitations in the Baltic Sea. Pages 177-210 A systems analysis of the Baltic Sea. Springer.

Hannig, M., G. Lavik, M. Kuypers, D. Woebken, W. Martens-Habbena, and K. Jürgens. 2007. Shift from denitrification to anammox after inflow events in the central Baltic Sea. Limnology and Oceanography **52**:1336-1345.

Hauss, H., J. M. Franz, and U. Sommer. 2012. Changes in N: P stoichiometry influence taxonomic composition and nutritional quality of phytoplankton in the Peruvian upwelling. Journal of Sea Research **73**:74-85.

Jentzsch L, Grossart HP, Plewe S, Schulze-Makuch D, Goldhammer T. Response of cyanobacterial mats to ambient phosphate fluctuations: phosphorus cycling, polyphosphate accumulation and stoichiometric flexibility. ISME communications. 2023 Jan 25;3(1):6.

Jespersen, A. M., and K. Christoffersen. 1987. Measurements of chlorophyll-*a* from phytoplankton using ethanol as extraction solvent. Archiv fur Hydrobiologie **109**:445-454.

Kalvelage, T., G. Lavik, P. Lam, S. Contreras, L. Arteaga, C. R. Löscher, A. Oschlies, A. Paulmier, L. Stramma, and M. M. Kuypers. 2013. Nitrogen cycling driven by organic matter export in the South Pacific oxygen minimum zone. Nature Geoscience **6**:228-234.

Kangro, K., K. Olli, T. Tamminen, and R. Lignell. 2007. Species-specific responses of a cyanobacteria-dominated phytoplankton community to artificial nutrient limitation in the Baltic Sea. Marine Ecology Progress Series **336**:15-27.

Keeling, R. F., A. Körtzinger, and N. Gruber. 2010. Ocean deoxygenation in a warming world.
Annual review of marine science **2**:199-229.
Kim, I.-N., K. Lee, N. Gruber, D. M. Karl, J. L. Bullister, S. Yang, and T.-W. Kim. 2014.
Increasing anthropogenic nitrogen in the North Pacific Ocean. Science **346**:1102-1106.
Koistinen, J., M. Sjöblom, and K. Spilling. 2017. Determining inorganic and organic Phosphorus.
In: Spilling K (ed.). Biofuels from algae. New York: Human Press, 2017, 87–94.
Koistinen, J., M. Sjöblom, and K. Spilling. 2018. Determining dissolved and biogenic Silica. In:
Spilling K (ed) Methods in molecular biology. Humana Press, New York, NY, p 95−102.
Kuosa, H., Marcussen, B. 1988: Grazing of bacteria and phytoplankton by heterotrophic
nanoflagellates in a Baltic Sea sample. – Hydrobiologia 161: 211–216.
Larsson, U., S. Hajdu, J. Walve, and R. Elmgren. 2001. Baltic Sea nitrogen fixation estimated from
the summer increase in upper mixed layer total nitrogen. Limnology and Oceanography
46:811-820.
Lignell, R., S. Kaitala, and H. Kuosa. 1992. Factors controlling phyto- and bacterioplankton in late
spring on a salinity gradient in the northern Baltic. Marine Ecology Progress Series **84**:121-
131.
McQuatters-Gollop, A., D. E. Raitsos, M. Edwards, Y. Pradhan, L. D. Mee, S. J. Lavender, and M.
J. Attrill. 2007. A long-term chlorophyll dataset reveals regime shift in North Sea
phytoplankton biomass unconnected to nutrient levels. Limnology and Oceanography
52:635-648.
Meyer, J., C. Löscher, S. Neulinger, A. Reichel, A. N. Loginova, C. Borchard, R. Schmitz, H.
Hauss, R. Kiko, and U. Riebesell. 2016. Changing nutrient stoichiometry affects
phytoplankton production, DOP accumulation and dinitrogen fixation–a mesocosm
experiment in the eastern tropical North Atlantic. Biogeosciences **13**:781-794.
Munkes, B., Löptien, U. and Dietze, H., 2021. Cyanobacteria blooms in the Baltic Sea: a review of
models and facts. *Biogeosciences*, *18*(7), pp.2347-2378.
Nausch, G., D. Nehring, and K. Nagel. 2008. Nutrient concentrations, trends and their relation to
eutrophication. State and Evolution of the Baltic Sea, 1952-2005: A Detailed 50-Year
Survey of Meteorology and Climate, Physics, Chemistry, Biology, and Marine
Environment:337-366.
Nausch, M., and G. Nausch. 2004. Bacterial utilization of phosphorus pools after nitrogen and
carbon amendment and its relation to alkaline phosphatase activity. Aquatic Microbial
Ecology **37**:237-245.
Niemi, Å. 1979. Blue-green algal blooms and N:P ratio in the Baltic Sea. Acta Botanica Fennica
110:57-61.
Pitkänen, H., Lehtoranta, J. and Räike, A., 2001. Internal nutrient fluxes counteract decreases in
external load: the case of the estuarial eastern Gulf of Finland, Baltic Sea. *AMBIO 30:* 195-
201.
Raateoja, M., H. Kuosa, and S. Hällfors. 2011. Fate of excess phosphorus in the Baltic Sea: a real
driving force for cyanobacterial blooms? Journal of Sea Research **65**:315-321.
Reynolds, C.S., 1984. The Ecology of Freshwater Phytoplankton. Cambridge University Press,
Cambridge.
Schoffelen, N.J., Mohr, W., Ferdelman, T.G., Littmann, S., Duerschlag, J., Zubkov, M.V., Ploug,
H. and Kuypers, M.M., 2018. Single-cell imaging of phosphorus uptake shows that key
harmful algae rely on different phosphorus sources for growth. Scientific Reports **8**:
p.17182.
Savchuk, O. P. 2005. Resolving the Baltic Sea into seven subbasins: N and P budgets for 1991-
1999. Journal of Marine Systems **56**:1-15.
Seip, K.L. and Reynolds, C.S. 1995. Phytoplankton functional attributes along trophic gradient and
season. – Limnology and Oceanography 40: 589–597.
Solórzano, L., and J. H. Sharp. 1980. Determination of total dissolved phosphorus and particulate
phosphorus in natural waters. Limnology and Oceanography **25**:754-758.

Spilling, K., E. Asmala, N. Haavisto, L. Haraguchi, K. Kraft, A.-M. Lehto, A. Lewandowska, J.
Norkko, J. Piiparinen, and J. Seppälä. 2022a. Dataset from a mesocosm experiment on
brownification in the Baltic Sea. Data in Brief **45**:108755.
Spilling, K., E. Asmala, N. Haavisto, L. Haraguchi, K. Kraft, A.-M. Lehto, A. M. Lewandowska, J.
Norkko, J. Piiparinen, and J. Seppälä. 2022b. Brownification affects phytoplankton
community composition but not primary productivity in eutrophic coastal waters: A
mesocosm experiment in the Baltic Sea. Science of The Total Environment **841**:156510.
Spilling, K., M.-T. Camarena-Gómez, T. Lipsewers, A. Martinez-Varela, F. Díaz-Rosas, E. Eronen-
Rasimus, N. Silva, P. von Dassow, and V. Montecino. 2019. Impacts of reduced inorganic
N: P ratio on three distinct plankton communities in the Humboldt upwelling system.
Marine Biology **166**:114.
Spilling, K., K. Olli, J. Lehtoranta, A. Kremp, L. Tedesco, T. Tamelander, R. Klais, H. Peltonen,
and T. Tamminen. 2018. Shifting diatom-dinoflagellate dominance during spring bloom in
the Baltic Sea and its potential effects on biogeochemical cycling. Frontiers in Marine
Science **5**:327.
Tamminen, T. 1995. Nitrate and ammonium depletion rates and preferences during a Baltic spring
bloom. Marine Ecology Progress Series **120**:123-133.
Vahtera, E., M. Laamanen, and J. Rintala. 2007. Use of different phosphorus sources by the bloom-
forming cyanobacteria *Aphanizomenon flos-aquae* and *Nodularia spumigena*. Aquatic
Microbial Ecology **46**:225-237.
Vahtera, E., Laanemets, J., Pavelson, J., Huttunen, M., and Kononen, K. 2005. Effect of upwelling
on the pelagic environment and bloom-forming cyanobacteria in the western Gulf of
Finland, Baltic Sea. Journal of Marine Systems **58**: 67-82.
Vanharanta, M., and K. Spilling. 2023. Uptake of excess phosphate at low inorganic N: P ratio in a
coastal sea afflicted with eutrophication. Marine Ecology Progress Series **718**:23-37.
Vanharanta, M., Santoro, M., Villena-Alemany, C., Piiparinen, J., Piwosz, K., Grossart, H.P.,
Labrenz, M. and Spilling, K., 2024a. Microbial remineralization processes during post-
spring-bloom excess phosphate in the northern Baltic Sea. bioRxiv, doi:
10.1101/2024.02.02.577174.
Vanharanta, M., Santoro, M., Villena-Alemany, C., Piiparinen, J., Grossart, H.-P., Labrenz, M.,
Piwosz, K., Spilling, K. 2024b. Effect of decreasing inorganic N:P ratio on the plankton
community - INN:PP. PANGAEA https://doi.org/10.1594/PANGAEA.966040
Wasmund, N. 1997. Occurrence of cyanobacterial blooms in the Baltic Sea in relation to
environmental conditions. Int. Revue ges. Hydrobiol **2**:169-184.
Wasmund, N., G. Nausch, B. Schneider, K. Nagel, and M. Voss. 2005. Comparison of nitrogen
fixation rates determined with different methods: a study in the Baltic Proper. Marine
Ecology Progress Series **297**:23-31.
Wasmund, N., G. Nausch, and M. Voss. 2012. Upwelling events may cause cyanobacteria blooms
in the Baltic Sea. Journal of Marine Systems **90**:67-76.
Wasmund, N., Kownacka, J., Göbel, J., Jaanus, A., Johansen, M., Jurgensone, I., Lehtinen, S. and
Powilleit, M., 2017. The diatom/dinoflagellate index as an indicator of ecosystem changes in
the Baltic Sea 1. Principle and handling instruction. Frontiers in Marine Science. **4**: 22
https://doi.org/10.3389/fmars.2017.00022

.

Fig 1. The concentration of nitrite + nitrate ($NO_2 + NO_3$), phosphate (PO4), ammonium (NH_4) and

dissolved silicate (DSi) plotted against time. The error bares represent the standard error (SE, n =

3).

Fig 2. The concentration of Chlorophyll a (Chla) and abundance of heterotrophic bacteria plotted

against time. The error bares represent the standard error (SE, n = 3).

Fig 3. The concentration of particulate organic carbon (POC), nitrogen (PON), phosphate (POP),

and biogenic silicate (BSi) plotted against time. The error bares represent the standard error (SE, n

= 3).

Fig 4. The abundance of *Synechoccoccus*-like cells, picophyto(?)eukaryotes, nano- and

microphytoplankton plotted against time. The error bares represent the standard error (SE, n = 3).

Fig 5. The biovolume of different nano- and microplankton groups plotted against time. The error

bars represent the standard error (SE, n = 3).

Fig 6. The abundance of different mesozooplankton groups plotted against time. The error bars

represent the standard error (SE, n = 3).

Fig 1

Fig 2

665　Fig 3

667   Fig 4

669   Fig 5

671   Fig 6

Supplementary Fig S1. Map of the Gulf of Finland with an outline of where the field data was

gathered from ships-of-opportunity (red box), and the mesocosm experiment (blue dot).

Supplementary Fig S2. Temperature during the experimental setup. Temperature was measured

every 30 s with a temperature probe placed at 50 cm depth right outside the mesocosm bags.

Supplementary Fig S3. Salinity measured at 50 cm depth inside the 12 experimental bags, plus

outside the mesocosm bags (thick grey line).

Supplementary Fig S4. Chla and phycocyanin fluorescence (upper graph) and temperature (lower

graph) taken outside the Hanko peninsula Gulf of Finland, from a flow through system on two

different ships of opportunity. Each point represents an average from the area of one passing. Data

from 2013 to 2016. Solid lines represent the Gaussian regression line.

Supplementary Fig S5. Chlorophyll and inorganic nutrients (nitrate, phosphate and dissolved

silicate) taken from discrete sample taken outside the Hanko peninsula Gulf of Finland, in a flow

through system on two different ships of opportunity. Each point represents one bottle filled from

the transect. Colors represent three different periods, cyan square = 1998 – 2003, blue triangle =

2004 – 2013 and red circle 2014 – 2023.

700   Supplementary Fig S6. Rate of change in dissolved oxygen during day- and nighttime.

Supplementary Fig S7. The sum of all biovolume measurements throughout the experiment of different nano- and microplankton groups. The presented p-value is an ANOVA statistical test comparing the different treatments. The temporal development in biovolume is presented in Fig 5.