1

Multiplexed *in situ* hybridization reveals distinct lineage identities for major and minor vein initiation during maize leaf development

Chiara Perico[1], Maricris Zaidem[1], Olga Sedelnikova[2], Samik Bhattacharya[3], Christian Korfhage[3] & Jane A. Langdale[1*]

[1] Department of Biology, University of Oxford, South Parks Rd, Oxford, OX1 3RB, UK

[2] Syngenta Jealott's Hill International Research Centre, Bracknell, RG42 6EY, UK

[3] Resolve BioSciences GmbH, Alfred-Nobel-Straße 10, 40789 Monheim am Rhein, Germany

* Author for correspondence: jane.langdale@biology.ox.ac.uk

ABSTRACT

Leaves of flowering plants are characterised by diverse venation patterns. Patterning begins with the selection of vein-forming procambial initial cells from within the ground meristem of a developing leaf, a process which is considered to be auxin-dependent, and continues until veins are anatomically differentiated with functional xylem and phloem. At present, the mechanisms responsible for leaf venation patterning are primarily characterized in the model eudicot *Arabidopsis thaliana* which displays a reticulate venation network. However, evidence suggests that vein development may proceed via a different mechanism in monocot leaves where venation patterning is parallel. Here, we employed Molecular Cartography, a multiplexed *in situ* hybridization technique, to analyse the spatiotemporal localisation of a subset of auxin related genes and candidate regulators of vein patterning in maize leaves. We show how different combinations of auxin influx and efflux transporters are recruited during leaf and vein specification, and how major and minor vein ranks develop with distinct identities. The localisation of the procambial marker *PIN1a* and the spatial arrangement of procambial initial cells that give rise to major and minor vein ranks further suggests that vein spacing is pre-patterned across the medio-lateral leaf axis prior to accumulation of the PIN1a auxin transporter. In contrast, patterning in the adaxial-abaxial axis occurs progressively, with markers of xylem and phloem gradually becoming polarised as differentiation proceeds. Collectively our data suggest that both lineage- and position-based mechanisms may underpin vein patterning in maize leaves.

SIGNIFICANCE STATEMENT

During the development of multicellular organisms specialized cell-types differentiate from pluripotent stem cells, with cell identity acquired via lineage- or position-based mechanisms. In plants, most organs develop post-embryogenesis and as such developmental processes are influenced by the external environment. To adapt to different environmental contexts and yet still form recognizable structures, position-based differentiation mechanisms are deployed in which cells adopt a certain fate depending on the activity of neighbouring cells. Such is the prevalence of position-based mechanisms in plant development that a role for lineage is rarely contemplated. Here we show that stem cells which give rise to different vein types in maize leaves are transcriptionally distinct, possibly reflecting a role for lineage-based mechanisms in the differentiation of leaf veins.

INTRODUCTION

Within flowering plants distinct leaf venation patterns characterise the leaves of eudicotyledonous and monocotyledonous species, which display reticulate networks and parallel venation respectively (reviewed in (1)). The characteristics of parallel vein formation have been studied primarily in leaves of the monocot grass *Zea mays* (maize). As in all flowering plants, maize leaves emerge from the flanks of the shoot apical meristem (SAM) in a regular pattern. The SAM itself comprises two layers, an outer (epidermal) L1 layer and an inner L2 layer, both of which are recruited into new leaf primordia as they are formed (2). Leaves emerge sequentially on opposite sides of the SAM

(alternate phyllotaxis) and at regular time intervals called plastochrons (P) (3) The incipient leaf, not
yet distinguishable anatomically but for which the position is already defined is referred to as P0, the
youngest visible leaf primordium as P1, and older leaves with increasing plastochron numbers
(4). Primordia inception is marked by an auxin maximum and by down-regulation of the homeobox
gene *KNOTTED1* (*KN1*) which acts in the SAM to promote indeterminacy within both the central
zone (CZ) of stem cells and the flanking peripheral zone (PZ) that gives rise to leaves. Before
primordia develop, proximo-distal (base-tip), medio-lateral (centre-margin) and adaxial-abaxial
(upper-lower) leaf axes are established, each of which provides the spatial context in which veins
develop (reviewed in (5)). Of particular importance is the establishment of a central layer of ground
tissue in the adaxial-abaxial axis at P1, from within which all veins are subsequently initiated (6, 7).
Parallel venation in maize leaves thus results from the specification of veins at regular intervals
across the medio-lateral leaf axis, within a single layer of ground tissue in the adaxial-abaxial axis.

The maize leaf is characterised by several vein ranks which develop in a spatiotemporally regulated
manner at successive plastochrons (Fig. 1A-C). Regardless of rank, development begins with the
specification of a procambial initial cell (referred to as a pre-procambial cell in eudicots) within the
central layer of the ground meristem. Once specified, this cell divides transversely to establish and
extend the procambial strand in the proximo-distal leaf axis, and longitudinally/obliquely to form the
procambial centre from which tissues of the vein differentiate in the medio-lateral and adaxial-abaxial
leaf axes (reviewed in (8)). Major veins, which include the mid- and lateral veins, develop acropetally
(base-to-tip) along the proximo-distal axis with the midvein positioned in the middle of the developing
primordium by P1 and lateral veins specified either side of it between P2 and P3 (Fig. 1B-C). Minor
veins, which include rank-1 and rank-2 intermediates, develop basipetally (tip-to-base) from P3 (Fig.
1B-C) and are spaced regularly between the major veins such that each vein pair is separated by
four cells (two mesophyll and two bundle sheath) across the medio-lateral axis (9–11). During the
final stages of vein development, differentiation occurs in the adaxial-abaxial leaf axis, positioning
xylem adaxially, phloem abaxially and sclerenchyma between the vein and epidermis. The final vein
rank to form develops transversely, connecting the longitudinal veins across the medio-lateral leaf
axis in a striate pattern.

Vein patterning mechanisms have been primarily studied in the model eudicot *Arabidopsis thaliana*
where the auxin efflux carrier PIN1 is required for leaf initiation on the flanks of the SAM, specification
of pre-procambial cells within the leaf primordium and directional extension of the procambial strands
(12). In the SAM, the simultaneous action of *PIN1* and auxin influx carriers of the *AUX1/LAX* family
generates the auxin maxima that determine the position of new leaves (13–15) and in the leaf *PIN1*
activity induces the auxin activated transcription factors *MONOPTEROS* (*MP/ARF5*) and *ATHB8*,
which commit cells from the ground meristem to pre-procambial and then procambial fate. A
feedback loop between auxin transport, MP and ATHB8 then restricts the procambial strand to a
couple of cell files in width (reviewed in (16)). The reticulate vein network results from the sequential

specification of vein ranks with new veins developing towards pre-existing veins, which are used as positional landmarks towards which the auxin flux is directed (12, 17). The presence of parallel veins that are precisely spaced across the medio-lateral leaf axis and a considerable expansion in the PIN and LAX transporter gene families (18) suggest that mechanisms underpinning vein patterning in monocots could be different from those found in *Arabidopsis*.

Recent genetic and transcriptome analyses have identified candidate regulators of venation patterns in monocot leaves. For example, a role for the *AINTEGUMENTA* (*ANT*) family in vein formation is supported by the finding that maize *ANT4* (19) is differentially expressed in median ground meristem (mGM) cells when compared to pre-mesophyll cells, and mutations in the *Setaria viridis AINTEGUMENTA1* (*ANT1*) gene lead to smaller leaves with distorted Kranz anatomy (20). Double mutations in *SCARECROW1* (*SCR1*) and its homeolog *SCR1h* condition narrower spaces between veins (21) and quadruple mutants with *NAKED ENDOSPERM1* (*NKD1*) and *NKD2* further enhance the spacing defect (22). In contrast, mutations in *SHORTROOT1* (*SHR1*) and its paralog *SHR2* disrupt the specification and/or subsequent development of veins (23–25), and mutations in *TOO MANY LATERALS* (*TML1*) form lateral veins in the position where the rank 1 intermediate veins that extend from blade to sheath normally develop (26). Whereas each of the genes mentioned thus far encode transcription factors, leaf venation defects have also been observed in mutants that perturb auxin biosynthesis, and transcriptome analyses have predicted a role for auxin transporters. For example, in *SPARSE INFLORESCENCE1* (*SPI1*) and *VANISHING TASSEL2* (*VT2*) mutants, reduced levels of auxin are associated with the formation of more intermediate veins in wider than normal leaves (27). Since neither pharmacological nor genetic approaches have disrupted the correlation between vein number and leaf width in monocots, any change of vein patterning accompanied by a change in leaf width is unlikely to result from directly disrupting the patterning mechanism (28). However, expression of the *AtPIN1* ortholog *ZmPIN1a* in all vein ranks from early stages of vein and leaf development (27) and a similar spatial expression pattern of the *LAX1* (note named *LAX2* in (19)) and *LAX2* (29) influx carriers at later stages of vein development, strongly suggest that auxin flux plays a role in vein patterning in the maize leaf. Although these observations are too fragmented to reveal any form of gene regulatory network, consistent expression profiles and mutant phenotypes of orthologs in different grass species (e.g. *TML1*, *SHR1, PIN1a/PIN1d*) suggest that core components of the mechanism underpinning parallel venation patterns in monocots have been identified.

Here we employed the Molecular Cartography platform to investigate cellular relationships during early stages of maize leaf development. Molecular Cartography is a high-throughput quantitative *in situ* hybridization technology that combines single cell detection of transcripts with a visual interface to identify spatial localisation within a tissue (30–33). Focusing on candidate regulators of leaf initiation and vein patterning, we qualitatively and quantitatively defined spatial and temporal patterns of transcript accumulation across consecutive stages of development in maize shoot apices. These

138 analyses revealed developmental patterns not previously detected by either morphological or cell
139 lineage analyses, providing insight into the specification of procambial initial cells, the establishment
140 of procambial centres giving rise to different vein ranks, and the differentiation of tissues within each
141 vein.
142
RESULTS
Molecular Cartography reliably reports transcript accumulation patterns in maize shoot apices
We sought to generate a spatial expression atlas in maize shoot apices for genes in which an
aberrant leaf venation phenotype has been described previously in either monocots or dicots (Table
1). This group of genes included auxin efflux and influx transporters such as *PIN* and LAX family
members, auxin response transcription factors (ARFs) such as the *MP* co-orthologs *ARF4* and
ARF29, plus candidate regulators of Kranz anatomy and vein patterning such as *TML1*, *SHR* and
SCR. Hybridizations were carried out using transverse sections of wild-type B73 maize shoot apices,
comprising the SAM (M) and the first five leaf primordia (P1 to P5) (Fig.1A-B). These stages allowed
the spatial distribution of transcripts to be captured during the development of both major (midvein
and laterals) and minor (rank 1 and rank 2 intermediate) veins (Fig. 1C). Data were collected across
two experiments for a total of six sections. Calcofluor White (cell wall) and DAPI (nuclear DNA)
staining allowed for automatic cell segmentation with CellPose 2.0 (34) (Fig.1B) and enabled
qualitative and quantitative single-cell studies. A qualitative assessment showed consistency between
Molecular Cartography and traditional *in situ* hybridizations (Figure S1). For example, *KNOTTED 1*
(*KN1)* transcripts were detected primarily in the L2 layer of the meristem with very low or no
expression detected in the L1 layer (Fig. S1A), consistent with published reports (35–37). Similarly,
PIN1a (Fig. S1B) and *TML1* (Fig. S1C) transcripts were detected as previously reported, that is in all
veins at all stages of leaf development (*PIN1a*) (27) or in spaces between major veins where
intermediate veins are developing (*TML1*) (26). For additional validation, we also carried out a
traditional *in situ* hybridization with *LAX2*, an auxin influx transporter. With Molecular Cartography,
LAX2 transcripts were detected in both major and minor veins (Fig. S1D), with higher levels detected
in laterals (Fig. S1D, arrowheads) than intermediates (Fig. S1D, asterisks). These observations were
consistent with results obtained by traditional *in situ* hybridization (Fig. S1E,F) (29). We thus
concluded that Molecular Cartography can reliably detect transcripts in transverse sections of maize
shoot apices and that the single-cell information available following cell segmentation would enable a
quantitative spatial expression atlas to be generated.
170
Transcript clusters distinguish functional domains in the maize shoot apical meristem
To investigate cellular relationships in the SAM, we segmented M-P1 stages and carried out
qualitative and quantitative analyses of the candidate gene expression profiles within these tissues.
Cluster analysis on all six sections revealed five distinct cell identities across the M-P1 stage, which
includes both the SAM and P1 primordium (Fig. 2A). To spatially visualise the clusters, two
representative sections were analysed: the first (E1D2) cut across the shoot apex at the base of P1

(Fig. 2B-C) and the second (S1A1) cut across the shoot apex within the region of the P0 incipient leaf

and included the tip of the P1 primordium (Fig. 2B, D). Images of both sections show that one cluster

corresponds to the central zone of the SAM, with high levels of *KN1* detected (yellow, Fig. 2C-D, Fig.

S2). A second 'peripheral zone' cluster (green) is apparent either opposite a cluster corresponding to

the developing P1 primordium (red, Fig. 2C) or completely encircling the central zone except for a

few cells with leaf primordium identity (Fig. 2D, asterisks). Cells classified as dividing, due to the

expression of genes encoding cyclins, primarily localise within the developing P1 primordium (blue,

Fig. 2A-D). This observation is consistent with a rapid increase in cell number during early stages of

leaf development. Finally, a handful of cells mark the boundary between the central zone of the

meristem and the developing P1 primordium (magenta, Fig. 2A, C). Although the nature and function

of these boundary cells in not clear, the other four clusters clearly demarcate regions of the SAM with

distinct functions. Given that leaf primordia are initiated within the peripheral zone, we next sought to

find traces of P0 identity in the peripheral zone opposite the tip of the P1 primordium (Fig. 2B). In

Brachypodium and tomato, PIN1d generates the auxin maximum that determines the position of new

leaves on the flanks of the SAM (38, 39). In our dataset, cells with the highest combined expression

of both *PIN1d* (Fig. 2E) and *LAX5* (Fig. 2F) were positioned within the peripheral zone opposite the

P1 primordium tip (Fig. 2G, Table S1). We therefore speculate that high levels of *LAX5* and *PIN1d*

mark the position of P0 in the maize meristem.

Leaf initiation and midvein formation are marked by distinct auxin transporters

The role of auxin transporters in directing auxin towards sites of both leaf and midvein initiation is well

documented in eudicots, with PIN1 playing a dominant role. To investigate whether specific auxin

transporters are associated with leaf and/or midvein initiation in maize, we compared transcript

accumulation profiles across the SAM and P1. Figure 3 shows hybridization of *PIN1d*, *PIN1a*, *LAX3*

and *LAX5* to transverse sections of two regions of the SAM, along with a visual summary of our

observations in a deduced longitudinal view. To verify our predicted P0 site, we first examined *KN1*

transcript accumulation. Importantly, transcripts were detected in the inner layers of the shoot apical

meristem (Fig. 3D-F) with near absence of accumulation in our predicted P0 region (Fig. 3E, dotted

line). For the genes encoding efflux transporters, *PIN1d* transcripts were detected throughout the

meristem and P1 (Fig. 3G-I), with the highest signal intensity at P0 (Fig. 3H). By contrast, *PIN1a*

transcripts were localised to the central zone of the meristem and were restricted to the midvein

position in the P1 primordium (Fig. 3J-L, asterisks). Notably, *PIN1a* transcript accumulation appeared

to be contiguous between P0 and the centre of the meristem, suggesting the presence of an auxin

conduit between P0 and the vasculature of the stem. Similar to the efflux carriers, the two influx

transporters also showed distinct profiles. *LAX3* (Fig. 3M-O) and *LAX5* (Fig. 3P-R), which are co-

orthologues of Arabidopsis *AUX1* and *LAX1* (18), were both detected in the peripheral zone of the

meristem but not in the central zone or in the P1 primordium. However, *LAX5* transcripts were

detected specifically in the newly forming P0 primordium (Fig. 3Q) whereas *LAX3* was detected both

within and beyond the P0 region (Fig. 3N). Collectively these results reveal that *PIN1d* and *LAX5*

mark the position of the P0 primordium within the peripheral zone of the SAM whereas *PIN1a* marks

both the region in the SAM where the developing midvein will connect the new leaf to the vasculature

of the stem, and the region in the P1 primordium where the midvein develops.

A transcript cluster corresponding to major vein identity emerges at P2

Whereas the midvein position is determined by P1, lateral veins continue to be specified during P2

and thus to investigate cellular relationships at this later stage, we segmented P2 primordia and

quantified transcript levels in each cell. Cluster analyses revealed five cellular identities, which we

labelled based on the transcripts present and the spatial localisation within the primordium (Fig. 4A-

C, Fig. S4). Through a visual analysis of the cluster distributions within the primordium, three cellular

identities appeared to coincide with the epidermis, ground tissue and vascular tissue respectively.

Dividing cells (as defined by expression of genes encoding cyclins) were present in all tissue types,

although at variable levels. Intriguingly, the profile of cells in the epidermal cluster varied slightly

between experiments. In one experiment there was a clear subdivision into two clusters (Fig. 4A-B,

Fig. S2I-L): an abaxial epidermal cluster (grey) and an adaxial epidermal plus leaf margin cluster

(cyan). The abaxial epidermis was characterized by high transcript levels of an epidermal marker

(*MYB67*) and several ARFs (*ARF16, ARF34*, *ARF23*, Fig. 4D), whereas the adaxial epidermis and

leaf margin cluster displayed high levels of *ANT4*, *PIN1d*, *LAX3* and *HB52* transcripts. This

subdivision of the epidermal tissue was not as marked in a second experiment, where an epidermal

cluster was identified but there was no clear subdivision between abaxial and adaxial/margin

domains (Fig. S2G-H). Notably, differences between experiments were also observed in overall

PIN1d, *ARF16* and *ARF34* levels (Fig. S3), suggesting that the detection of these transcripts may be

more sensitive to subtle changes and variations between experiments, or to the precise cutting plane

of the sections or developmental stage of the plant. In both experiments, cells in the ground tissue

predominantly accumulated *ARF4* (*MP*), *ANT4* and *LAX1* transcripts, as well as low levels of *RVN1*,

SCR1/SCR1h and *TML1* (Fig. 4D).

Unlike the epidermis and ground tissue, which are derived from the SAM and are present at P0,

procambial initial cells that give rise to the leaf veins have to be specified from within the ground

tissue at successive stages of primordium development. Our data reveal that a cell cluster with

vascular identity emerges at P2 (orange, Fig. 4B-C) and that it includes both the midvein (Fig. 4C,

arrowhead) and lateral veins (Fig. 4C, asterisks). The cluster is characterized by high levels of *PIN1a*,

SHR1, *SHR2*, and *ANT4* transcripts and, to a lesser extent, *LAX1* and *LAX2*. Notably, the distance

between veins identified by this cluster is consistently four to five cells (Fig. 4C, bracket), which

suggests that venation patterns and the distance between veins are specified by P2. Also of note is

that the midvein and lateral veins form a single cluster despite representing two distinct vein ranks

and developing at different timepoints (Fig. 4C). Collectively, results show that with the gene panel

used in this study the P2 maize leaf primordium can be broadly subdivided into three major tissue

types: ground tissue, epidermis and vasculature, with vascular identity marked by the accumulation

of *PIN1a*, *SHR1*, *SHR2* and *ANT4* transcripts.

TML1 marks a distinct intermediate vein cluster at P3 and P4

Given that all major veins can be identified by a specific transcript cluster at P2, we next investigated

whether minor vein ranks can be identified by the same or a different gene set. For this purpose,

cluster analysis was carried out with data from P3 and P4 primordia, to capture the development of

rank-1 intermediate veins. Analysis of P3 and P4 primordia identified six and seven clusters

respectively (Fig. 5A-D). In the context of the epidermis, as for P2, discrepancies were seen between

experiments in terms of whether the epidermis was divided into distinct zones: one cluster associated

with the abaxial epidermis and another with both the adaxial epidermis and the leaf margin (Fig.

4A,B; Fig. 5A,B,E,F; Fig S2M-R – grey and cyan respectively). In both experiments, however, the

epidermal cluster was distinct from the ground tissue, which again was similar to that seen at P2 with

PIN1a, *LAX1*, *RVN1*, *ANT4*, *HB52*, *ARF4*, *ARF16* and *ARF34* transcripts detected (Fig. 4D, Fig.

5C,D). The striking difference between clusters identified at P2 versus those identified at later stages

was that two distinct vascular clusters were evident at P3 and P4. Notably these two vascular

clusters were also evident when data were integrated across all stages of development (M-P1 to P5)

(Fig. S5) as opposed to being analyzed within individual primordia.

The spatial localisation of the two vascular clusters and the spacing between them within the P3 and

P4 primordia suggest that they represent distinct lateral (Fig. 5G,H - orange) and intermediate (Fig.

5G,H - magenta) vein identities. Both clusters share a common set of transcripts which include

PIN1a, *LAX1*, *ANT4*, *PIN1d*, *HB52*, *ARF16* and *ARF24*. However, the intermediate vein cluster is

distinguished by high levels of *TML1* transcript (Fig. 5C,D), which is consistent with the observation

that the *tml1* mutation in maize specifically perturbs the development of rank-1 intermediate veins

(Fig. S1C) (26). The intermediate vein cluster also displays lower levels of *SHR1* and *SHR*2

transcripts than the lateral vein cluster (Fig. 5C,D). In conclusion, clustering of cells from stage P3

and P4 highlighted similarities and differences in the development of different vein ranks. Both lateral

and rank-1 intermediate veins accumulate a similar set of auxin transporters and procambial

markers, such as *PIN1a*, *LAX1*, *ANT4* and *HB52*. However, the clear difference between the lateral

and rank-1 intermediate vein clusters, and the specific accumulation of *TML1* transcripts in the

intermediate cluster, suggest that vein ranks are transcriptionally distinct after just a few divisions of

the procambial initial.

Vein formation across the medio-lateral leaf axis reveals a temporal hierarchy of PIN1a, SHR1 and

LAX2 accumulation during early procambium development

To determine the extent to which the spatial regulation of genes associated with vein formation is

accompanied by temporal regulation, we analysed the distribution of transcripts across the shoot

apex. For this analysis we focused on genes that are shared between lateral and intermediate veins

clusters, such as *PIN1a*, *SHR1*, *SHR2* and *ANT4,* in order to dissect events during the first few divisions of the procambial initial. We discovered that *PIN1a* was the earliest and most persistent marker of vein development, being detected at all stages of development from M-P1 to P5 (Fig. 6A). In M-P1, *PIN1a* transcripts marked both the position of the developing midvein (Fig. 6B, white ellipse, "m") and the central zone of the meristem, suggesting a role in connecting the newly developing midvein with the vasculature of the stem. From P2 onwards, transcripts were detected in all vein ranks of all leaf primordia, i.e. the midvein (Fig. 6C-D, white ellipse), laterals (Fig. 6D, empty arrowheads) and intermediates (Fig. 6D, filled arrowheads). Notably, *PIN1a* transcripts were detected before and after procambial centres became anatomically distinguishable from the surrounding ground tissue, with other procambial and vascular markers following the appearance of *PIN1a*. For example, we detected transcripts of the GRAS transcription factor *SHR1* (Fig. 6E, Fig. S6A-E) in all vein ranks (as indicated by *PIN1a* expression), but at a slightly later stage of development. These observations revealed not only that the expression of *SHR1* occurs temporally after that of *PIN1a*, but also that the development of veins across the medio-lateral axis follows a specific dynamic: veins closer to the midrib express *SHR1* earlier than those closer to the leaf margin. In a similar manner, we observed that *LAX2* transcripts accumulated at least one plastochron later than *SHR1* (Fig. 6F, Fig. S7A-E). By contrast, however, *ANT4* is present in all intermediate veins across the medio-lateral axis of the leaf (arrowheads, Figs. S6F, S7F). This suggests that the observed spatio-temporal dynamic is specific for *SHR1* and *LAX2*. Together, our data suggest that PIN1a is the earliest leaf procambial marker at the developmental stages considered in this study and that other procambial markers follow in a specific temporal and spatial hierarchy across the medio-lateral axis.

Vascular markers become polarized in the adaxial-abaxial axis at late stages of vein development

Having observed that at least a subset of venation markers are sequentially expressed during procambium initiation, we next investigated whether localisation could also vary between early and late stages of vein development. We took into consideration two vein ranks, the midvein and lateral veins, for which a trajectory from early specification (M-P1 and P2 stage) to phloem and xylem differentiation (P3 to P5) could be traced across the developmental stages in our study. Several markers for which transcripts are localised in all vein ranks were analyzed. A representative example is provided in Figure 7A, where *PIN1a* (red), *HB52* (yellow) and *LAX2* (blue) transcripts are superimposed, and all transcripts are detected in all vein ranks and at all leaf developmental stages. Regardless of vein rank, at stages prior to differentiation of phloem or xylem, all three markers were detected throughout the procambial centre (Fig. 7B). Once phloem and xylem were differentiated, however, *PIN1a*, *HB52* and *LAX2* transcripts were polarly localized (Fig. 7C,E). In both the mid- (Fig. 7C) and lateral (Fig. 7E) veins, *PIN1a* transcripts were detected on the adaxial side, *HB52* transcripts towards the phloem on the abaxial side, and *LAX2* juxtaposed between the two. Collectively, our data suggest that vascular-associated genes are tightly regulated temporally and spatially, both within and between different vein ranks, with changes in spatial localisation marking transitions between the

331 specification of procambial initial cells, the development and extension of procambial strands, and
332 differentiation of tissues within the vein.
333
DISCUSSION
Here we have used multiplexed *in situ* hybridization and a set of marker genes associated with auxin
transport and vein patterning to investigate cellular relationships during leaf specification and vein
formation in maize shoot apices (Fig. 1). On leaf initation, we reveal that it is possible to identify the
region corresponding to incipient P0 leaves within the peripheral zone of the SAM, based on high
density of *PIN1d* and *LAX5* transcripts (Fig. 2). *PIN1a* transcripts connect the site of the incipient leaf
with the central zone of the meristem, likely connecting the new leaf with the vasculature of the stem
and determining the position of the midvein at P1 (Fig. 3). From P2 to P4, the specification of both
major (midvein and laterals) and minor (intermediate) vein ranks can be traced, with intermediate
veins characterized by the presence of the transcription factor *TML1* (Fig. 4, 5). Strikingly, the
localisation of the procambial marker *PIN1a* and the spatial position of major and minor vein clusters
suggest that vein spacing could be pre-patterned prior to *PIN1a* accumulation. Once *PIN1a*
accumulation (and presumably auxin flux along the proximo-distal leaf axis) is initiated, a
spatiotemporal hierarchy is triggered in which vein markers accumulate in a sequential manner that is
position-dependent across the medio-lateral leaf axis (Fig. 6). During the final stages of vein
development, in which phloem and xylem are differentiated, vascular markers become spatially
restricted in the adaxial-abaxial leaf axis (Fig. 7). Collectively these data suggest progressive
restriction of gene expression in all three leaf axes as veins are specified, initiated, extended and
then differentiated.
353
The importance of understanding cell lineage relationships during development has long been
recognised, not least to distinguish mechanisms in which cell fate is inherently specified in
descendents following mitosis, from those in which fate specification relies on positional information
from neighbouring cells (reviewed in (40)). In maize, many aspects of leaf development have been
elucidated by exploiting genetic mosaics, including the presence of distinct central and marginal
lineages in the medio-lateral leaf axis (41, 42) and the presence of a lineage in the centre of the
adaxial-abaxial axis from which all veins are derived (7). With the advent of single cell sequencing,
spatial transcriptomics and quantitative *in situ* hybridizations, discussion of lineage relationships have
altered. Cells with shared fates are identified regardless of whether that fate is determined by
lineage-based or positional mechanisms (33, 43–45). Importantly, combining information from both
approaches allows classical lineage distinctions to be associated with specific gene expression
profiles and also facilitates more refinement of cellular relationships during later stages of
development. For example, we identified transcriptionally distinct populations of cells in the peripheral
zone of the SAM that will give rise to the P0 primordium, and identified two distinct populations of
vein-forming procambial cells in the central ground meristem layer of P3 and P4 leaf primordia.
Notably, some of the markers that defined developing procambial lineages, such as *PIN1a, TML1*

and *HB52*, could be detected in a dispersed pattern prior to vein specification. For example, *TML1* transcripts were detected at P2, prior to intermediate vein specification, but accumulation patterns were not restricted to the central ground meristem layer (Fig. S1C) (26). Similarly, *TML1* transcripts were detected primarily in procambial cells that were dividing to form intermediate veins in the middle of the P3 primordia (arrowheads, Fig. S1C), but were detected in a more dispersed pattern at the leaf margins where procambium was still being specified. Undoubtedly, better refinement of the developmental trajectories of cell lineages within and between leaf primordia could be achieved using unbiased approaches such as scRNAseq (44, 46) and/or 3D analyses (31) that would avoid variability between sections. However, even this limited gene set has enabled the identification of transcriptionally distinct cell populations within the vein-forming cell lineage (*sensu stricto*) in the middle of the adaxial-abaxial leaf axis (7) and has provided information about when cell fate is specified.

In the context of vein patterning, cellular relationships identified through transcript profiles have revealed distinctions that have not been made previously by either cell lineage or histological analyses. The set of genes used in this study distinguished two separate cell clusters between stages P2 and P4, corresponding to major (midvein and lateral) and intermediate veins (Figs. 4, 5). No substantial differences were detected between midvein and lateral vein clusters, which were grouped together from P2 onwards. This observation could reflect the fact that our dataset does not include genes which are specific for either midvein or lateral vein identity. Alternatively, it suggests that the midvein and lateral veins are not intrinsically different, a suggestion supported by the phenotype of midribless mutants (47, 48). By contrast, minor veins are transcriptionally distinct from major veins once procambium is specified, as evidenced by transcript localisation profiles (see example of single cells in Figure 5G) and the phenotype of loss of function *TML1* mutants (26). Although both minor and major veins are specified *de novo* from the middle layer of the ground meristem, this observation suggests that procambial initials are inherently distinct. This hypothesis is supported by the observation that intermediate veins start to be specified near the midvein during P3, when growth in the medio-lateral axis is ongoing and lateral veins are still being specified near the leaf margin (Fig. 5 G-H, orange). This leads to a gradient in the medio-lateral axis, with veins closer to the midrib being more developed than those closer to the leaf margin (11). If major and minor vein lineages were not inherently distinct, intermediate veins close to the midrib would be of similar size to lateral veins close to the leaf margin but they are not (Table S2). These findings suggest that procambial initial cells that give rise to major and minor vein ranks in the maize leaf are distinct at inception.

In the model eudicot *Arabidopsis thaliana,* auxin fluxes have been associated with vein patterning, with pre-existing vasculature used as a positional landmark towards which auxin flows and new procambium is initiated (12, 17). In contrast, the localisation of transcripts encoding auxin transporters in maize suggests that auxin fluxes associated with procambial initiation are pre-

patterned. Specifically, *PIN1a* transcripts are first detected in individual procambial initial cells that are already spaced appropriately across the medio-lateral axis. As such, *PIN1a* activity is not necessary to determine which cells in the ground tissue will commit to procambial fate. Notably, the DR5 transcriptional reporter is detected along the leaf margin in maize leaves, indicating that an auxin maximum is present at the margin (27, 49). Given that PIN1a does not overlap with the DR5 signal at the margin of P0 to P5 primordia (27), auxin flux from the margin is unlikely to be inducing procambial initial cell specification or procambial strand extension. Instead, an unidentified factor must be inducing PIN1a accumulation in the subset of regularly spaced ground tissue cells that become procambial initial cells. Once *PIN1a* is activated, other auxin transporters and vascular markers are detected in developing veins but temporal and/or spatial patterns of transcript accumulation again suggest progressive restriction of activity. For example, maize abaxial and adaxial markers *ARF3a* (*AUXIN RESPONSE FACTOR 3a*) and *RLD1* (*ROLLED LEAF1*) are localised throughout the adaxial (RLD1) or abaxial (ARF3a) leaf domains between P0 and P3, but by P4 they are primarily restricted to the vasculature and are associated with differentiated xylem (*RLD1*) and phloem (*ARF3a*) (49). Together, these findings suggest that the position of procambial initials in the medio-lateral axis of the maize leaf is pre-patterned prior to auxin flux through the procambial strand, and that the patterning of differentiated tissues within each vein is similarly pre-patterned in the adaxial-abaxial leaf axis.

The role of auxin in both leaf initation and vein patterning has been acknowledged for some time (1, 50) but the spatial transcriptomes reported here reveal new insights. In *Arabidopsis thaliana*, *PIN1* acts as a master regulator of both leaf positioning and vein specification (12, 51, 52). In non-Brassicaceae, however, the expansion of the PIN1 family and the emergence of a sister clade (SoPIN1/PIN1d) enabled sub-functionalisation (18, 38, 39, 53). Our analysis reveals that positioning of the P0 primordium in maize is associated with combined accumulation of transcripts encoding the auxin efflux and influx carriers *PIN1d* and *LAX5* (Fig. 3), whereas the specification of procambium is associated with the accumulation of *PIN1a*, *PIN1d* and *LAX1*, with *LAX2* likely being the dominant influx carrier later in vein development. Our results are consistent with previous findings where combined expression of *PIN1a*, *PIN1d* and *LAX1* (note *LAX2* in (19)) was detected from procambial initial cell selection through to later stages of procambium development ((19)). Detection of *PIN1d* but not *PIN1a* at the site of the P0 primordium suggests a functional subdivision within the maize PIN1 family as seen in tomato and *Brachypodium* (38, 52), with *PIN1d* (*SoPIN*) regulating phyllotaxy and *PIN1a* tracing the midvein and its connection to the subtending vasculature (Fig. 3) (49). We further speculate that *LAX5*, co-ortholog with *LAX3* of *Arabidopsis AUX1/LAX1*, supports the role of *PIN1d* in generating an auxin maximum in the L1 of the SAM, whereas *LAX2* acts with *PIN1a* to extend procambial strands in the proximo-distal leaf axis. Together, these findings suggest a subdivision of roles within both the *PIN1* and *LAX* gene families that allows different combinations of auxin influx and efflux transporters to be recruited in different developmental contexts.

CONCLUSIONS

This study used a quantitative multiplex *in situ* hybridization platform to identify populations of cells with shared identity, and to elucidate how those populations diverged during the course of leaf primordium development in maize. Regardless of whether they arise as the result of positional or lineage-based cues, cell fates can be associated with the spatial co-localisation of transcripts within a tissue. As such, we propose that major and minor vein ranks in the maize leaf have distinct identities at inception from the ground meristem. We further propose that veins in the maize leaf are pre-patterned in all three leaf axes, with auxin triggering downstream specification events through finely tuned spatio-temporal accumulation of influx and efflux transporters.

METHODS

Molecular Cartography probe design

Probes were designed using Resolve BioSciences' proprietary design algorithm and gene annotations from the *Zea Mays* RefGen V4. The final set of probes was selected as outlined in (30).

Plant growth and tissue preparation for Molecular Cartography

Samples were prepared, hybridized and imaged as in Vlad et al. (26). In total six sections were analyzed - E1B2, E1D2, S1B1, S1B2, S1A1 and S1A2. (30)Two independent experiments were carried out – the first with sections E1B2 and E1D2 and the second with samples S1A1, S1A2, S1B1 and S1B2.

Data analysis

All data analyses and plots were carried out with R Studio and the Molecular Cartography ImageJ plugin Polylux proprietary of Resolve Biosciences. Unprocessed single-cell data was extracted from the spatial images using the ImageJ plugin Polylux from Resolve BioSciences. Correlation plots were generated using corrplot (54) to evaluate the linear relationships between genes across experimental repeats (note that each experimental repeat was also a biological repeat as each section was taken from a different shoot apex). The R package Seurat (55) was used to carry out all clustering data analyses and plotting.. The Seurat clustering workflow was adapted with minor adjustments to fit the data frame from molecular cartography. Briefly, for each stage, a Seurat object was generated from unprocessed data with parameters: min.cells = 20 (P2) or 30 (M-P1, P3, P4, M to P5); min.features=3. Parameters were adjusted to account for the difference in the number of cells per primordia stage. Data was normalised using the "LogNormalize" method and a scale factor of 1. Scaling using ScaleData, to prevent overrepresentation of frequent signals in dimensional reductionwas then done using default parameters. A principal component analysis (PCA) was performed using standard parameters. After determining dimensionality of the dataset via the Seurat ElbowPlot function, 10 components were taken into consideration to run the Uniform Manifold Approximation and Projection (UMAP) dimensionality reduction to cluster the cells Clustering was

based on neighbours, found using the "FindNeighbours" command (dims = 1:10, dims = 1:13 for M to P5), followed by a default "FindClusters". FindClusters allowed cluster generation by constructing a Shared Nearest Neighbour (SNN) graph and applying the Louvain algorithm. Cluster resolution was set at either 0.3 (P4, M to P5) or 0.4 (M-P1, P2, P3). Dimension plots were generated using the "DimPlot" command and "umap" as the reduction method. Cluster location was visually checked using the Polylux plugin, and identity was assigned based on the cluster localisation within the sections.

Tissue fixation and preparation for traditional in situ hybridization

Traditional *in situ* hybridization was carried out using wax-embedded wild-type B73 maize shoot apices as previously described (25) with a 268 bp digoxygenin (DIG)-RNA labelled probe complementary to the 3'UTR sequence of the *LAX2* gene (starting at 126 bp downstream of the stop codon).

Vein width quantification

Wild-type B73 maize seeds (n = 5) were germinated and grown as for Molecular Cartography, using a 3:1 mixture of John Innes #3 and vermiculite. Following full blade expansion of leaves three and four, 5 mm segments spanning the whole leaf width were harvested from the widest point of the blade. Samples were then fixed in 3:1 EtOH/acetic acid for 30 min at room temperature, followed by a 30 min wash in 70% EtOH. After an overnight incubation in 70% EtOH at 4°C, leaf segments were embedded in blocks of 6% agar and sectioned using a vibratome. Leaf sections were imaged using a stereomicroscope and the width of lateral and intermediate veins outside of the midrib region were measured using ImageJ. Lateral veins were labelled with increasing numbers across the medio-lateral axis (L1 indicating the lateral vein closest to the midrib). Average intermediate veins number and width were measured between successive lateral veins, and between the last lateral vein and the leaf margin. All vein width quantifications were carried out on one half of the leaf segments.

Data access

All raw data, Calcofluor White and DAPI images and cell segmentations used in this study and in Vlad *et al* (26) are deposited on Zenodo (URL: https://doi.org/10.5281/zenodo.10605855). All unprocessed and normalised and scaled data are available in Data S1, S3 and S4, and cell identities following cluster analyses are reported in Data S2.

LIST OF SUPPLEMENTARY MATERIAL

Table S1. Gene expression in P0 cells identified from the 'peripheral zone' cluster.

Table S2. Lateral and intermediate vein width quantification across the medio-lateral axis of leaves three and four.

Data S1. Unprocessed and processed (normalised and scaled) transcript counts for stages M-P1, P2, P3, P4. Cells are labelled based on the section of origin (E1B2, E1D2, S1A1, S1A2, S1B1 or

S1B2), and the plastochron stage under analysis (M-P1, P2, P3, P4). These data were used to generate Figs. 2, 4 and 5.

Data S2. Molecular Cartography data clustering. Identities are labelled as outlined in Figs. 2A, 4A, 5A-B and S4. Cell labels as in Data S1.

Data S3. Unprocessed transcript counts for the analysis shown in Fig. S5.

Data S4. Normalised and scaled data obtained from Data S3. Refers to the analysis shown in Fig. S5.

Figure S1. Validation of spatial transcriptomics for localization of transcripts in maize shoot apices.

Figure S2. Visual representation of clusters at all stages of leaf development.

Figure S3. Correlation analysis of the Molecular Cartography datasets.

Figure S4. Representative images showing the localization of transcripts for all genes in the Molecular Cartography panel

Figure S5. Data clustering from stage M-P1 to P5 indicates separate identities for major and minor veins

Figure S6. *PIN1a* expression precedes that of *SHR1* across the medio-lateral axis.

Figure S7. *PIN1a* expression precedes that of *LAX2* across the medio-lateral axis.

AUTHOR CONTRIBUTIONS

CP and JAL conceived and designed the experiments, MZ and CP wrote the code and performed the cluster analysis, OS carried out the traditional *in situ* hybridization experiments, SB carried out the spatial transcriptomics hybridizations under the supervision of CK, CP performed all the sample preparation and data analysis; CP and JAL wrote the first draft of the manuscript and all authors contributed to the final version.

ACKNOWLEDGEMENTS

The authors thank all members of the Langdale lab for constructive discussions of the results and data analysis, and feedback on the manuscript. We also thank Thomas Hughes for helping with the sample preparation, and Scott Poethig, Tina Schreier, Hilde Nelissen, Jessica Joossens and Denia Herwegh for feedback on the manuscript.

COMPETING INTERESTS

SB and CK are employees of Resolve Biosciences which is the proprietor of Molecular Cartography technology. All other authors declare no competing interests.

FUNDING

This research was funded by the Bill and Melinda Gates Foundation C_4 Rice grant awarded to the University of Oxford 2019-2024 (INV-002970).

REFERENCES

1. C. Perico, S. Tan, J. A. Langdale, Developmental regulation of leaf venation patterns: monocot versus eudicots and the role of auxin. *New Phytologist* **234**, 783–803 (2022).

2. R. S. Poethig, I. M. Sussex, The cellular parameters of leaf development in tobacco: a clonal analysis. *Planta* **165**, 170–184 (1985).

3. R. O. Erickson, F. J. Michelini, The plastochron index. *Am J Bot* **44**, 297 (1957).

4. A. W. Sylvester, W. Z. Cande, M. Freeling, Division and differentiation during normal and liguleless-1 maize leaf development. *Development* **110**, 985–1000 (1990).

5. J. W. Satterlee, M. J. Scanlon, Coordination of leaf development across developmental axes. *Plants* **8** (2019).

6. K. Esau, Plant Anatomy. *John Wiley and Sons* (1965).

7. J. A. Langdale, B. Lane, M. Freeling, T. Nelson, Cell lineage analysis of maize bundle sheath and mesophyll cells. *Dev Biol* **133**, 128–139 (1989).

8. O. V. Sedelnikova, T. E. Hughes, J. A. Langdale, Understanding the genetic basis of C4 Kranz anatomy with a view to engineering C3 crops. *Ann Rew Genetics* **52**, 249–270 (2018).

9. S. H. Russell, R. F. Evert, Leaf vasculature in Zea mays L. *Planta* **164**, 448–458 (1985).

10. T. Nelson, N. Dengler, Leaf vascular pattern formation. *Plant Cell* **9**, 1121–1122 (1997).

11. B. C. Sharman, Developmental anatomy of the shoot of Zea mays L. *Ann Bot* **6**, 245–282 (1942).

12. E. Scarpella, D. Marcos, J. Friml, T. Berleth, Control of leaf vascular patterning by polar auxin transport. *Genes Dev* **20**, 1015 (2006).

13. K. Bainbridge, *et al.*, Auxin influx carriers stabilize phyllotactic patterning. *Genes Dev* **22**, 810–823 (2008).

14. D. Reinhardt, *et al.*, Regulation of phyllotaxis by polar auxin transport. *Nature* **426**, 255–260 (2003).

15. E. Benková, *et al.*, Local, efflux-dependent auxin gradients as a common module for plant organ formation. *Cell* **115**, 591–602 (2003).

16. M. Biedroń, A. Banasiak, Auxin-mediated regulation of vascular patterning in Arabidopsis thaliana leaves. *Plant Cell Reports* **37**, 1215–1229 (2018).

17. N. M. Linh, C. Verna, E. Scarpella, Coordination of cell polarity and the patterning of leaf vein networks. *Curr Opin Plant Biol* **41**, 116–124 (2018).

18. M. S. Matthes, *et al.*, Auxin EvoDevo: conservation and diversification of genes regulating auxin biosynthesis, transport, and signaling. *Mol Plant* **12**, 298–320 (2019).

19. W. Y. Liu, *et al.*, Regulators of early maize leaf development inferred from transcriptomes of laser capture microdissection (LCM)-isolated embryonic leaf cells. *Proc Natl Acad Sci U S A* **119** (2022).

20. W.-Y. Liu, *et al.*, Maize ANT1 modulates vascular development, chloroplast development, photosynthesis, and plant growth. *Proceedings of the National Academy of Sciences* **117**, 21747–21756 (2020).

21. T. E. Hughes, O. V. Sedelnikova, H. Wu, P. W. Becraft, J. A. Langdale, Redundant SCARECROW genes pattern distinct cell layers in roots and leaves of maize. *Development* **146**, dev177543 (2019).

22. T. E. Hughes, O. Sedelnikova, M. Thomas, J. A. Langdale, Mutations in NAKED-ENDOSPERM IDD genes reveal functional interactions with SCARECROW during leaf patterning in C$_4$ grasses. *PLoS Genet* **19**, e1010715 (2023).

23. Q. Liu, *et al.*, SHORT ROOT and INDETERMINATE DOMAIN family members govern PIN-FORMED expression to regulate minor vein differentiation in rice. *Plant Cell* **35**, 2848–2870 (2023).

24. T. L. Slewinski, *et al.*, Short-Root1 plays a role in the development of vascular tissue and Kranz anatomy in maize leaves. *Mol Plant* **7**, 1388-1392 (2014).

25. M. L. Schuler, O. V. Sedelnikova, B. J. Walker, P. Westhoff, J. A. Langdale, SHORTROOT-Mediated increase in stomatal density has no impact on photosynthetic efficiency. *Plant Physiol* **176**, 757–772 (2018).

26. D. Vlad, *et al.*, The WIP6 transcription factor TOO MANY LATERALS specifies vein type in C_4 and C_3 grass leaves. *Current Biology* **34**, 1–17 (2024).

27. J. M. Robil, P. McSteen, Hormonal control of medial–lateral growth and vein formation in the maize leaf. *New Phytologist* **238**, 125–141 (2023).

28. I. R. A. Smillie, K. A. Pyke, E. H. Murchie, Variation in vein density and mesophyll cell architecture in a rice deletion mutant population. *J Exp Bot* **63**, 4563–4570 (2012).

29. S. Leiboff, *et al.*, Genetic control of morphometric diversity in the maize shoot apical meristem. *Nature Communications* **6**, 1–10 (2015).

30. P. Glenn, *et al.*, Wheat bZIPC1 interacts with FT2 and contributes to the regulation of spikelet number per spike. *Theoretical and Applied Genetics* **136**, 1–20 (2023).

31. T. Nobori, M. Oliva, R. Lister, J. R. Ecker, Multiplexed single-cell 3D spatial gene expression analysis in plant tissue using PHYTOMap. *Nature Plants* **9**, 1026–1033 (2023).

32. X. Yang, *et al.*, Spatial transcriptomics of a lycophyte root sheds light on root evolution. *Current Biology* **33**, 4069-4084.e8 (2023).

33. R. Laureyns, *et al.*, An in situ sequencing approach maps PLASTOCHRON1 at the boundary between indeterminate and determinate cells. *Plant Physiol* **188**, 782–794 (2022).

34. M. Pachitariu, C. Stringer, Cellpose 2.0: how to train your own model. *Nature Methods* **19**, 1634–1641 (2022).

35. M. Kitagawa, X. Xu, D. Jackson, Trafficking and localization of KNOTTED1 related mRNAs in shoot meristems. *Commun Integr Biol* **15**, 158 (2022).

36. L. G. Smith, D. Jackson, S. Hake, Expression of knotted1 marks shoot meristem formation during maize embryogenesis. *Dev Genet* **16**, 344–348 (1995).

37. D. Jackson, B. Veit, S. Hake, Expression of maize KNOTTED1 related homeobox genes in the shoot apical meristem predicts patterns of morphogenesis in the vegetative shoot. *Development* **120**, 405–413 (1994).

38. C. C. Martinez, D. Koenig, D. H. Chitwood, N. R. Sinha, A sister of PIN1 gene in tomato (Solanum lycopersicum) defines leaf and flower organ initiation patterns by maintaining epidermal auxin flux. *Dev Biol* **419**, 85–98 (2016).

39. D. L. O'Connor, *et al.*, A division in PIN-mediated auxin patterning during organ initiation in grasses. *PLoS Comput Biol* **10**, e1003447 (2014).

40. R. S. Poethig, Clonal analysis of cell lineage patterns in plant development. *Am J Bot* **74**, 581 (1987).

41. M. J. Scanlon, K. D. Chen, C. C. McKnight IV, The narrow sheath duplicate genes: sectors of dual aneuploidy reveal ancestrally conserved gene functions during maize leaf development. *Genetics* **155**, 1379–1389 (2000).

42. M. J. Scanlon, M. Freeling, Clonal sectors reveal that a specific meristematic domain is not utilized in the maize mutant narrow sheath. *Dev Biol* **182**, 52–66 (1997).

43. I. Efroni, P. L. Ip, T. Nawy, A. Mello, K. D. Birnbaum, Quantification of cell identity from single-cell gene expression profiles. *Genome Biol* **16**, 1–12 (2015).

44. R. Shahan, *et al.*, A single-cell Arabidopsis root atlas reveals developmental trajectories in wild-type and cell identity mutants. *Dev Cell* **57**, 543–560 (2022).

45. C. Kidner, V. Sundaresan, K. Roberts, L. Dolan, Clonal analysis of the Arabidopsis
root confirms that position, not lineage, determines cell fate. *Planta* **211**, 191–199
(2000).
46. T. Denyer, *et al.*, Spatiotemporal developmental trajectories in the Arabidopsis root
revealed using high-throughput single-cell RNA sequencing. *Dev Cell* **48**, 840-
852.e5 (2019).
47. A. E. Richardson, A. Sluis, S. Hake, The Hoja loca1 mutant and AUX/IAA function
in phytomer development. *bioRxiv* 2020.03.27.012211 (2021).
48. J. Strable, *et al.*, Maize YABBY genes drooping leaf1 and drooping leaf2 regulate
plant architecture. *Plant Cell* **29**, 1622–1641 (2017).
49. R. Johnston, S. Leiboff, M. J. Scanlon, Ontogeny of the sheathing leaf base in
maize (Zea mays). *New Phytologist* **205**, 306–315 (2015).
50. Y. Xiong, Y. Jiao, The diverse roles of auxin in regulating leaf development. *Plants*
8 (2019).
51. N. Bhatia, *et al.*, Auxin acts through MONOPTEROS to regulate plant cell polarity
and pattern phyllotaxis. *Current Biology* **26**, 3202–3208 (2016).
52. D. Kierzkowski, M. Lenhard, R. Smith, C. Kuhlemeier, Interaction between
meristem tissue layers controls phyllotaxis. *Dev Cell* **26**, 616–628 (2013).
53. D. L. O'Connor, *et al.*, Cross-species functional diversity within the PIN auxin
efflux protein family. *Elife* **6** (2017).
54. Y. Hao, *et al.*, Integrated analysis of multimodal single-cell data. *Cell* **184**, 3573-
3587.e29 (2021).
55. T. Wei, V. Simko, R package "corrplot": visualization of a correlation matrix.
(Version 0.92). (2021).
694
695
696

 FIGURE LEGENDS
698

Figure. 1: Stages of leaf and vein development in maize shoot apices. A) Cartoon representation

of a longitudinal section across a maize shoot apex. SAM (black) indicates the Shoot Apical

Meristem; younger to older stages of leaf development (leaf primordia, P) are indicated using

increasing numbers and different colours. The black dotted line indicates the position of the

transverse section depicted in (B). **B)** Representative transverse section of a maize shoot stained

with Calcofluor and segmented with CellPose 2.0. SAM and developing leaf primordia are segmented

and colour-coded as in (A). **C)** Vein ranks appear sequentially during leaf primordium development.

P1: midvein only (M, cyan); P2: midvein and laterals (L, yellow); P3, P4 and P5: midvein, laterals and

intermediate veins (i, magenta). Scale bars = 100 µm.

Figure 2: *PIN1d* and *LAX5* transcripts mark the P0 primordium. A) Cluster analysis of cells in the

SAM and P1 primordium reveals five domains. **B)** Schematic of shoot apex showing the position of

transverse sections illustrated in (C) and (D). **C)** The section taken at the base of the P1 primordium

(E1D2) shows that the central zone (*KN1*-expressing) cluster localizes between the developing P1

primordium (red) and the peripheral zone (green). **D)** The section taken at the tip of the P1

primordium (S1A1) shows that the peripheral zone cluster (green) surrounds the central zone cluster

(yellow), and includes the putative incipient primordium P0 region. Asterisks highlight cells with

primordium identity flanking the P0 region. **E-F)** A subset (arrowhead) of cells from the 'peripheral

zone' cluster that co-express *PIN1d* (E) and *LAX5* (F) mark the incipient P0 primordium (G, dotted

ellipse). Transcript levels and cell labels are reported in Table S1. Scale bar = 30 µm.

Figure 3: Distinct auxin transporter profiles mark leaf initiation and midvein specification. **A-**

C) Schematics of sections in longitudinal (A) and transverse view (B, C). P0 = P0 primordium, P1 =

P1 primordium, SAM = shoot apical meristem, L1 and L2 define the two layers within the shoot apical

meristem. **D-F)** Schematic (D) and spatial transcriptomics (E, F) of *KN1*. Transcripts accumulate in

the L2 layer of the shoot apical meristem, with very low or no transcript levels detected in the L1 layer

or at the site of P0 initiation (dotted region, E). **G-I)** Schematic (G) and spatial transcriptomics (H, I) of

PIN1d. Transcripts accumulate predominantly in the initiating P0 primordium (P0) but also in the SAM

and the developing P1 primordium (H, I). **J-L)** Schematic (J) and spatial transcriptomics (K, L) of

PIN1a. Transcripts are detected at the site of the developing midvein in P1 (asterisks), in the initiating

P0 primordium and in the central zone of the SAM. **M-O)** Schematic (M) and spatial transcriptomics

(N, O) of *LAX3*. Transcripts are detected in the peripheral zone of the SAM, including the P0 initiation

site, but not in the developing P1 primordium. Lower transcript levels are detected in the P0

primordium (N) than in the rest of the peripheral zone. **P-R)** Schematic (P) and spatial transcriptomics

(Q, R) of *LAX5*. Transcripts are only detected in the initiating P0 primordium, primarily in the L1 layer.

Scale bars = 20 µm.

Figure 4: A transcript cluster with vascular identity emerges at P2. A) Cluster analysis and UMAP plot for the P2 leaf primordium reveals five domains. **B)** Representative image of cluster localisation in a P2 primordium, colour-coded to match (A). **C)** The vasculature cluster marks the midvein (arrowhead) and the developing lateral veins (asterisks). White bracket indicates spacing between veins. **D)** Dot-plot summarising transcript levels in each cluster. Average expression refers to the average transcript level in a cluster (cell type), calculated from the scaled data. Percent expressed refers to the percent of cells in a cluster in which transcripts (as indicated on the x-axis) were detected. Note high levels of *SHR1*, *SHR2*, *ANT4*, *PIN1a* and *LAX2* in the vasculature cluster. Scale bars = 50 μm.

Figure 5: Intermediate veins cluster separately from lateral veins at P3 and P4. A, B) Cluster analyses and UMAP plots of P3 (A) and P4 (B) leaf primordia. **C, D)** Dotplots summarising transcript levels for each gene in each cluster at P3 (C) and P4 (D). Average expression refers to the average transcript level for each gene in a cluster (cell type), calculated from the scaled data. Percent expressed refers to the percent of cells in a cluster in which transcripts (as indicated on the x-axis) were detected. **E, F)** Representative image of cluster localisations in a P3 (E) and P4 (F) primordium, colour-coded to match (A, B). **G, H)** Two distinct vein clusters are identified, lateral (orange) and intermediate (magenta), at both P3 (G) and P4 (H). Arrowhead in G) indicates a potential procambial initial. Scale bars = 50 μm.

Figure 6: *PIN1a* accumulation precedes the sequential accumulation of *SHR1* and *LAX2* transcripts in dividing procambial centres. **A)** Representative image of *PIN1a* transcript accumulation (red pixels) in the SAM and P1 to P5 leaf primordia. Transcripts accumulate in both the leaf sheath and leaf blade, at all stages of vein development and in all vein ranks. White boxes represent the position of enlargements shown in (B-D). **B, C)** *PIN1a* is detected at low levels in the future midvein at P1 (B - m, circled) and at higher levels at P2 (C- m, circled). **D)** *PIN1a* transcripts are detected in all developing veins within the P5 primordium (m, midvein; i, intermediate; L, lateral). **E)** Superimposed images of *PIN1a* (red) and *SHR1* (blue) transcripts reveal that *PIN1a* accumulation precedes that of *SHR1*. Arrowheads in the inset indicate three developing intermediate veins (i1, i2, i3), in order of their initiation across the medio-lateral leaf axis. *SHR1* transcripts are detected in i1 and i2, but not in the more recently developed i3. **F)** Superimposed images of *SHR1* (blue) and *LAX2* (yellow) transcripts reveal that *SHR1* accumulation precedes that of *LAX2*. Arrowheads in the inset indicate the same developing intermediate veins as in (E): *LAX2* transcripts are not detected in any of these veins but are found in neighbouring older veins. Scale bars = 100 μm.

Figure 7: Differentiation of phloem and xylem is accompanied by polar localization of vascular markers. A-E) Transcripts of vascular markers *HB52* (yellow), *LAX2* (blue) and *PIN1a* (red) accumulate in developed and developing veins in all leaf primordia (A). All markers co-localise in developing mid-(B) and lateral (D) veins, with very little adaxial (ad) or abaxial (ab) separation. In

775 differentiated mid-(C) and lateral (E) veins, transcripts preferentially accumulate on the phloem-side

776 (P, as in the case of *HB52*, yellow), xylem-side (X, as in the case of *PIN1a*, red), or juxtaposed (as in

777 the case of *LAX2*, blue). Scale bars = 100 μm (A), 20 μm (B-E).

 SUPPLEMENTARY FIGURE LEGENDS

Figure S1 Validation of Molecular Cartography methodology for localization of transcripts in

maize shoot apices. **A)** *KN1* transcripts localize in the shoot apical meristem, specifically the L2

layer as previously shown by in situ hybridization (37). **B)** *PIN1a* transcripts mark all developing veins

at all stages of leaf development. **C)** *TML1* transcripts are detected in developing intermediate veins.

D, E) Molecular Cartography (D) and *in situ* hybridization (E) of *LAX2*. In both cases, transcripts are

detected in developing vascular tissue with higher levels detected in major veins (inset, arrowhead)

compared to minor veins (inset, asterisks). **F)** *In situ* hybridization control with sense *LAX2* probe.

Scale bars = 50 µm.

Figure S2: Visual representation of clusters at all stages of leaf development. Sample numbers

are depicted across the top and plastochron stages at the left. Cluster colours are as depicted in the

main figures. Scale bars = 30 µm (A-F), 50 µm (G-R), 100 µm (S-X).

792
793

Figure S3: Correlation analysis of the Molecular Cartography datasets. **A)** Correlation plot for

comparison of different experimental datasets with the same tissue preparation method. **B)**

Correlation plot for comparison of datasets with manual (dataset 3) and automatic (dataset 2) tissue

preparation method within the same experiment. **C)** All datasets together. Brown colour indicates

negative correlations between rows and columns, blue indicates a positive correlation. The number in

each cell represents the correlation coefficient (confidence level = 0.95).

Figure S4: Representative images showing the localization of transcripts for all genes in the

Molecular Cartography panel. Images are chosen across all sections. Representative transcript

localization for *PIN1a*, *KN1*, *LAX2*, *SHR1*, *TML1* and *HB52* are provided in Fig. S1, Fig. 6 and Fig. 7.

Scale bars = 100 µm.

Figure S5: Data clustering from stage M-P1 to P5 indicates separate identities for major and

minor veins. A) Cluster analysis and UMAP plot of data collectively analysed for stage M-P1 to P5.

B) Dotplot summarising transcript levels for each gene in each cluster at M-P1 to P5. Average

expression refers to the average transcription level for each gene in a cluster (cell type), calculated

from the scaled data. Percent expressed refers to the percent of cells in a cluster in which transcripts

(as indicated on the x-axis) were detected. Blue and red boxes highlight the expression profiles of

clusters 6 and 0. **C-F)** Representative images of cluster localisation from M to P5, colour-code as in

panel A. Scale bars = 100 µm.

Figure S6: *PIN1a* **expression precedes that of** *SHR1* **across the medio-lateral axis**. **A-E)**

Transcript accumulation for *PIN1a* and *SHR1* is detected in both the sheath and blade and in both

major and minor veins. Intermediate veins closer to the midrib accumulate *SHR1* before those closer

to the leaf margin. In insets, intermediate veins are labelled with progressive numbers indicating the
degree of proximity to the midrib. **F)** Unlike *SHR1*, *ANT4* is detected in all intermediate veins across
the medio-lateral axis. Scale bars = 100 µm.

Figure S7: *PIN1a* expression precedes that of *LAX2* across the medio-lateral axis. **A-E)**
Transcript accumulation for *PIN1a* and *LAX2* is detected in both the sheath and blade and in both
major and minor veins. Intermediate veins closer to the midrib accumulate *LAX2* before those closer
to the leaf margin. In insets, intermediate veins are labelled with progressive numbers indicating the
degree of proximity to the midrib. **F)** Unlike *LAX2*, *ANT4* is detected in all intermediate veins across
the medio-lateral axis. Scale bars = 100 µm.

Figure. 1: Stages of leaf and vein development in maize shoot apices. A) Cartoon representation of a longitudinal section across a maize shoot apex. SAM (black) indicates the Shoot Apical Meristem; younger to older stages of leaf development (leaf primordia, P) are indicated using increasing numbers and different colours. The black dotted line indicates the position of the transverse section depicted in (B). **B)** Representative transverse section of a maize shoot stained with Calcofluor and segmented with CellPose. SAM and developing leaf primordia are segmented and colour-coded as in (A). **C)** Vein ranks appear sequentially during leaf primordium development. P1: midvein only (M, cyan); P2: midvein and laterals (L, yellow); P3, P4 and P5: midvein, laterals and intermediate veins (i, magenta). Scale bars = 100 µm.

Figure 2: *PIN1d* and *LAX5* transcripts mark the P0 primordium. A) Cluster analysis of cells in the SAM and P1 primordium reveals five domains. **B)** Schematic of shoot apex showing the position of transverse sections illustrated in (C) and (D). **C)** The section taken at the base of the P1 primordium (E1D2) shows that the central zone (*KN1*-expressing) cluster localizes between the developing P1 primordium (red) and the peripheral zone (green). **D)** The section taken at the tip of the P1 primordium (S1A1) shows that the peripheral zone cluster (green) surrounds the central zone cluster (yellow) and includes the putative incipient primordium P0 region. Asterisks highlight cells with primordium identity flanking the P0 region. **E-F)** A subset (arrowhead) of cells from the 'peripheral zone' cluster that co-express *PIN1d* (E) and *LAX5* (F) mark the incipient P0 primordium (G, dotted ellipse). Transcript levels and cell labels are reported in Table S1. Scale bar = 30 μm.

Figure 3: Distinct auxin transporter profiles mark leaf initiation and midvein specification. **A-C)** Schematics of sections in longitudinal (A) and transverse view (B, C). P0 = P0 primordium, P1 = P1 primordium, SAM = shoot apical meristem, L1 and L2 define the two layers within the shoot apical meristem. **D-F)** Schematic (D) and spatial transcriptomics (E, F) of *KN1*. Transcripts accumulate in the L2 layer of the shoot apical meristem, with very low or no transcript levels detected in the L1 layer or at the site of P0 initiation (dotted region, E). **G-I)** Schematic (G) and spatial transcriptomics (H, I) of *PIN1d*. Transcripts accumulate predominantly in the initiating P0 primordium (P0) but also in the SAM and the developing P1 primordium (H, I). **J-L)** Schematic (J) and spatial transcriptomics (K, L) of *PIN1a*. Transcripts are detected at the site of the developing midvein in P1 (asterisks), in the initiating P0 primordium and in the central zone of the SAM. **M-O)** Schematic (M) and spatial transcriptomics (N, O) of *LAX3*. Transcripts are detected in the peripheral zone of the SAM, including the P0 initiation site, but not in the developing P1 primordium. Lower transcript levels are detected in the P0 primordium (N) than in the rest of the peripheral zone. **P-R)** Schematic (P) and spatial transcriptomics (Q, R) of *LAX5*. Transcripts are only detected in the initiating P0 primordium, primarily in the L1 layer. Scale bars = 20 μm.

Figure 4: A transcript cluster with vascular identity emerges at P2. A) Cluster analysis and UMAP plot for the P2 leaf primordium reveals five domains. **B)** Representative image of cluster localisation in a P2 primordium, colour-coded to match (A). **C)** The vasculature cluster marks the midvein (arrowhead) and the developing lateral veins (asterisks). White bracket indicates spacing between veins. **D)** Dot-plot summarising transcript levels in each cluster. Average expression refers to the average transcript level in a cluster (cell type), calculated from the scaled data. Percent expressed refers to the percent of cells in a cluster in which transcripts (as indicated on the x-axis) were detected. Note high levels of *SHR1*, *SHR2*, *ANT4*, *PIN1a* and *LAX2* in the vasculature cluster. Scale bars = 50 μm.

Figure 5: Intermediate veins cluster separately from lateral veins at P3 and P4. A, B) Cluster analyses and UMAP plots of P3 (A) and P4 (B) leaf primordia. **C, D)** Dotplots summarising transcript levels for each gene in each cluster at P3 (C) and P4 (D). Average expression refers to the average transcript level for each gene in a cluster (cell type), calculated from the scaled data. Percent expressed refers to the percent of cells in a cluster in which transcripts (as indicated on the x-axis) were detected. **E, F)** Representative image of cluster localisations in a P3 (E) and P4 (F) primordium, colour-coded to match (A, B). **G, H)** Two distinct vein clusters are identified, lateral (orange) and intermediate (magenta), at both P3 (G) and P4 (H). Arrowhead in G) indicates a potential procambial initial. Scale bars = 50 μm.

Figure 6: *PIN1a* accumulation precedes the sequential accumulation of *SHR1* and *LAX2* transcripts in dividing procambial centres. **A)** Representative image of *PIN1a* transcript accumulation (red pixels) in the SAM and P1 to P5 leaf primordia. Transcripts accumulate in both the leaf sheath and leaf blade, at all stages of vein development and in all vein ranks. White boxes represent the position of enlargements shown in (B-D). **B, C)** *PIN1a* is detected at low levels in the future midvein at P1 (B - m, circled) and at higher levels at P2 (C- m, circled). **D)** *PIN1a* transcripts are detected in all developing veins within the P5 primordium (m, midvein; i, intermediate; L, lateral). **E)** Superimposed images of *PIN1a* (red) and *SHR1* (blue) transcripts reveal that *PIN1a* accumulation precedes that of *SHR1*. Arrowheads in the inset indicate three developing intermediate veins (i1, i2, i3), in order of their initiation across the medio-lateral leaf axis. *SHR1* transcripts are detected in i1 and i2, but not in the more recently developed i3. **F)** Superimposed images of *SHR1* (blue) and *LAX2* (yellow) transcripts reveal that *SHR1* accumulation precedes that of *LAX2*. Arrowheads in the inset indicate the same developing intermediate veins as in (E): *LAX2* transcripts are not detected in any of these veins but are found in neighbouring older veins. Scale bars = 100 μm.

Figure 7: Differentiation of phloem and xylem is accompanied by polar localization of vascular markers. A-E) Transcripts of vascular markers *HB52* (yellow), *LAX2* (blue) and *PIN1a* (red) accumulate in developed and developing veins in all leaf primordia (A). All markers co-localise in developing mid-(B) and lateral (D) veins, with very little adaxial (ad) or abaxial (ab) separation. In differentiated mid-(C) and lateral (E) veins, transcripts preferentially accumulate on the phloem-side (P, as in the case of *HB52*, yellow), xylem-side (X, as in the case of *PIN1a*, red), or juxtaposed (as in the case of *LAX2*, blue). Scale bars = 100 μm (A), 20 μm (B-E).

Gene_name	Gene_name Maize GDB	Gene ID (v3)	Gene ID (v4)	Gene ID (v5)	Origin of gene name (*supported by phylogeny)	Orthology	Variations
TML1	dot5	GRMZMG2150011	Zm00001d020037_T001	Zm00001eb309530	Vlad et al 2024*	AtWIP6	Previously referred to as Zmdot1 or Zmdot5
TML2	–	GRMZM2G134998	Zm00001d005566_T001	Zm00001eb097850	Vlad et al 2024*	AtWIP6	Previously referred to as Zmdot1h
SCR1	scro1	GRMZM2G131516	Zm00001d052380_T001	Zm00001eb195650	Hughes et al. 2019*	AtSCR	
SCR1h	Gras 37	GRMZM2G015080	Zm00001d005029_T001	Zm00001eb093670	Hughes et al. 2019*	AtSCR	
SHR1	Gras 58	GRMZM2G132794	Zm00001d029607_T001	Zm00001eb020840	Schuler et al. 2017*	AtSHR; OsSHR2	
SHR2	Gras 19	GRMZM2G172657	Zm00001d021973_T001	Zm00001eb326020	Schuler et al. 2017*	AtSHR; OsSHR1	
SHR2h	Gras 85	GRMZM2G019060	Zm00001d006721_T001	Zm00001eb108090	Schuler et al. 2017*	AtSHR; OsSHR1	
RVN1	IDD10	GRMZM2G143723	Zm00001d039467_T001	Zm00001eb121010	Sedelnikova et al. 2018	–	
RVN1h	IDD 18	GRMZM2G046290	Zm00001d008842_T001	Zm00001eb337700	Sedelnikova et al. 2018	–	
ANT4	EREB161	GRMZM2G021573	Zm00001d048004_T002	Zm00001eb399440	Liu et al. 2019*	ANT1-4 all co-orthologs of AtANT1	
PIN1a	pin1	GRMZM2G098643	Zm00001d044812_T001	Zm00001eb372180	Matthes et al. 2019*	Co-ortholog AtPIN1, ortholog OsPIN1a	
PIN1d	pin4	GRMZM2G171702	Zm00001d052442_T001	Zm00001eb196240	O'Connor et al 2014*	Ortholog of OsPIN1c & OsPIN1d	Also named ZmPIN1d in many papers
LAX1	aic3	GRMZM2G149481	Zm00001d028401_T005	Zm00001eb010460	Matthes et al. 2019*	Co-ortholog AtLAX2, OsLAX2 & OsLAX4	Named ZmLAX2 in Liu et al 2022
LAX2	aic1	GRMZM2G129413	Zm00001d030310_T001	Zm00001eb026490	Matthes et al. 2019*	Co-ortholog AtLAX2, ortholog of OsLAX4	
LAX3	atl1	GRMZM2G127949	Zm00001d042809_T003	Zm00001eb147480	Matthes et al. 2019*	Co-ortholog AtAUX1 & AtLAX1, ortholog of OsLAX1	
LAX5	aic4	GRMZM2G067022	Zm00001d038275_T005	Zm00001eb288800	Matthes et al. 2019*	Co-ortholog AtAUX1 & AtLAX1, ortholog of OsLAX3	
HB52	hb52	(AC187157.4_FG005)	Zm00001d008869_T001	Zm00001eb337970	TAIR	Ortholog of AtHB8 & AtHB15	
ARF4	arftf4	GRMZM2G034840	Zm00001d001945	Zm00001eb067270	Matthes et al.	Co-ortholog of AtMP,	

			_T010		2019*	OsARF11	
ARF29	arftf29	GRMZM2G086949	Zm00001d026540_T016	Zm00001eb433020	Matthes et al. 2019*	Co-ortholog of AtMP, OsARF11	
ARF16	arftf16	GRMZM2G028980	Zm00001d053819_T008	Zm00001eb207450	Matthes et al. 2019*	Co-ortholog of AtARF6, ortholog of OsARF6	
ARF34	arftf34	GRMZM2G081158	Zm00001d031064_T006	Zm00001eb031700	Matthes et al. 2019*	Co-ortholog of AtARF6, ortholog of OsARF17	
ARF23	arftf23	GRMZM2G441325	Zm00001d038698_T007	Zm00001eb292830	Matthes et al. 2019*	Co-ortholog of AtARF3 & AtARF4, ortholog of OsARF14	
KN1	kn1	GRMZM2G017087	Zm00001d033859_T002	Zm00001eb055920	Hake et al. 1989	Class I KNOX	
MYB67	myb67	GRMZM2G089244	Zm00001d053060_T001	Zm00001eb201150	Maize GDB	–	
CYC4	cyc4	GRMZM2G310115	Zm00001d012560_T002	Zm00001eb369340	Maize GDB	Ortholog of AtCYCB1 genes	cyc7 in raw data
CYC7	cyc7	GRMZM2G073003	Zm00001d043164_T001	Zm00001eb150450	Maize GDB	Ortholog of AtCYCB1 genes	previously cyc4b, cyc4 in raw data
CYC3	cyc3	GRMZM2G073671	Zm00001d036360_T002	Zm00001eb272010	Maize GDB	Ortholog AtCYCB2 genes	

TABLE 1. List of genes used for Molecular Cartography

Figure S1: Validation of Molecular Cartography methodology for localization of transcripts in maize shoot apices. **A)** *KN1* transcripts localize in the shoot apical meristem, specifically the L2 layer as previously shown by *in situ* hybridization (Jackson et al., 1994). **B)** *PIN1a* transcripts mark all developing veins at all stages of leaf development. **C)** *TML1* transcripts accumulation in developing intermediate veins from stage P3 (white arrowheads). **D, E)** Molecular Cartography (D) and *in situ* hybridization (E) of *LAX2*. In both cases, transcripts are detected in developing vascular tissue with higher levels detected in major veins (inset, arrowhead) compared to minor veins (inset, asterisks). **F)** *In situ* hybridization control with sense *LAX2* probe. Scale bars = 50 μm.

Figure S2: Visual representation of clusters at all stages of leaf development. Sample numbers are depicted across the top and plastochron stages at the left. Cluster colours are as depicted in the main figures: A-F refer to the colour scheme in Fig. 2, G-L to the colour scheme in Fig. 4 and M-X to the colour scheme in Fig. 5. Scale bars = 30 μm (A-F), 50 μm (G-R), 100 μm (S-X).

Figure S3: Correlation analysis of the Molecular Cartography datasets. **A)** Correlation plot for comparison of different experimental datasets with the same tissue preparation method. **B)** Correlation plot for comparison of datasets with manual (dataset 3) and automatic (dataset 2) tissue preparation method within the same experiment. **C)** All datasets together. Brown colour indicates negative correlations between rows and columns, blue indicates a positive correlation. The number in each cell represents the correlation coefficient (confidence level = 0.95).

Fig. S4: Representative images showing the localization of transcripts for all genes in the Molecular Cartography panel. Images are chosen across all sections. Representative transcript localization for *PIN1a*, *KN1*, *LAX2*, *SHR1*, *TML1* and *HB52* are provided in Fig. S1, Fig. 6 and Fig. 7. Scale bars = 100 µm.

Figure S5: Data clustering from stage M-P1 to P5 indicates separate identities for major and minor veins. A) Cluster analysis and UMAP plot of data collectively analysed for stage M-P1 to P5. **B)** Dotplot summarising transcript levels for each gene in each cluster at M-P1 to P5. Average expression refers to the average transcription level for each gene in a cluster (cell type), calculated from the scaled data. Percent expressed refers to the percent of cells in a cluster in which transcripts (as indicated on the x-axis) were detected. Blue and red boxes highlight the expression profiles of clusters 6 and 0. **C-F)** Representative images of cluster localisation from M to P5, colour-code as in panel A. Scale bars = 100 μm.

Figure S6: *PIN1a* expression precedes that of *SHR1* across the medio-lateral axis. **A-E)** Transcript accumulation for *PIN1a* and *SHR1* is detected in both the sheath and blade and in both major and minor veins. Intermediate veins closer to the midrib accumulate *SHR1* before those closer to the leaf margin. In insets, intermediate veins are labelled with progressive numbers indicating the degree of proximity to the midrib. **F)** Unlike *SHR1*, *ANT4* is detected in all intermediate veins across the medio-lateral axis. Scale bars = 100 μm.

Figure S7: *PIN1a* expression precedes that of *LAX2* across the medio-lateral axis. **A-E)** Transcript accumulation for *PIN1a* and *LAX2* is detected in both the sheath and blade and in both major and minor veins. Intermediate veins closer to the midrib accumulate *LAX2* before those closer to the leaf margin. In insets, intermediate veins are labelled with progressive numbers indicating the degree of proximity to the midrib. **F)** Unlike *LAX2*, *ANT4* is detected in all intermediate veins across the medio-lateral axis. Scale bars = 100 μm.

Gene.ID	E1B2_M_P1_23	E1B2_M_P1_121	E1B2_M_P1_191	S1A1_M_P1_39	S1A1_M_P1_49	S1A1_M_P1_82	S1A1_M_P1_83	S1A1_M_P1_88	S1A1_M_P1_95	S1A1_M_P1_100	S1A1_M_P1_124	S1A1_M_P1_127	S1A1_M_P1_150	S1A1_M_P1_160	S1A2_M_P1_16	S1A2_M_P1_72	S1A2_M_P1_81	S1A2_M_P1_234	S1B1_M_P1_183
SHR2	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
TML2	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
PIN1a	0	0	1	0	0	3	0	1	3	6	0	0	3	0	1	0	0	0	0
SCR1h	0	0	0	0	1	1	0	0	0	0	0	0	0	0	0	0	0	0	1
RVN1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
KN1	1	3	0	0	0	14	0	0	0	0	0	0	0	0	0	0	0	0	0
CYC3	1	0	0	0	0	0	0	0	1	2	0	1	0	0	0	0	0	0	0
CYC4	0	0	0	0	0	0	0	0	0	5	2	0	0	0	0	0	0	0	0
ANT4	0	0	0	2	0	0	3	0	0	5	1	0	4	6	0	0	2	0	0
PIN1d	0	0	0	0	2	0	7	6	5	16	2	3	10	4	0	2	1	3	1
LAX1	0	0	0	0	0	0	0	1	0	1	0	2	0	0	0	0	1	0	1
LAX3	0	0	1	1	0	8	2	7	1	9	2	3	8	0	0	1	2	2	0
LAX5	1	1	1	1	0	2	1	2	2	5	6	3	5	2	1	1	1	1	1
HB52	0	1	0	0	2	1	0	3	1	5	4	4	2	7	11	4	5	6	0
ARF4	1	1	3	0	0	7	4	1	2	0	5	4	0	0	3	1	1	6	0
ARF29	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
ARF16	0	0	0	0	3	8	2	5	4	2	5	6	1	6	0	0	0	5	1
ARF34	5	4	3	2	7	0	1	7	5	9	4	11	7	5	2	6	3	1	5
ARF23	0	0	0	0	6	1	3	3	10	21	12	14	13	3	1	0	4	2	1
MYB67	0	0	0	0	0	0	0	1	0	0	0	1	1	0	0	0	0	0	0
CYC7	1	0	1	1	7	0	0	3	15	21	27	14	2	3	9	0	1	0	0

Table S1: Gene expression in P0 cells identified from the 'peripheral zone' cluster.

LEAF 3

Plant ID	Blade width (mm)	Blade length (mm)	# Laterals - half leaf	Lateral veins width (μm) - half leaf				# Intermediates - half leaf				Avg. intermediate veins width (μm) - half leaf			
				L1	L2	L3	L4	L1-L2	L2-L3	L3-L4	L4-margin	L1-L2	L2-L3	L3-L4	L4-margin
1	14.49	330	4	112.89	106	88.24	89.64	15	14	8	5	48.9	45	45.3	44.4
2	16.56	360	4	118.44	121.82	97.7	83.81	16	16	8	4	49.75	49.36	42.67	41.21
3	16.8	365	4	150.73	116.4	106.87	76.53	12	14	13	11	52.42	48.28	46.69	49.73
4	17.99	300	4	138.18	131.45	125.36	83.93	13	14	12	6	62.4	53.77	43.29	43.53
5	14.79	310	4	110.92	105.63	95.02	83.19	14	15	10	4	53.06	42.79	35.22	44.29
Avg.	16.13	333.00	4.00	126.23	116.26	102.64	83.42	14.00	14.60	10.20	6.00	53.31	47.84	42.63	44.63
stdev	1.46	29.07	0.00	17.44	10.95	14.35	4.65	1.58	0.89	2.28	2.92	5.38	4.22	4.44	3.13

LEAF 4

Plant ID	Blade width (mm)	Blade length (mm)	# Laterals - half leaf	Lateral veins width (μm) - half leaf					# Intermediates - half leaf					Avg. intermediate veins width (μm) - half leaf				
				L1	L2	L3	L4	L5	L1-L2	L2-L3	L3-L4	L4-L5	L5-margin	L1-L2	L2-L3	L3-L4	L4-L5	L5-margin
1	20.33	460	5	112.3	140.05	124.7	107.75	87.82	14	16	14	8	9	49.4	43.89	48.77	57.43	51.09
2	24.1	460	5	147.23	153.06	137.04	120.53	82.18	14	15	13	8	4	53.11	49.98	51.27	48.67	56.02
3	23.19	470	5	137.34	122.93	125.94	103.3	83.93	15	15	14	11	9	47.86	51.97	52.78	48.59	44.51
4	20.89	440	5	118.7	128.56	153.13	115.5	89.7	15	17	17	8	5	48.58	57.1	54.66	45.65	55.88
5	22.33	420	5	120.1	127.34	130.81	98.77	101.37	10	12	14	10	7	65.58	56.91	55.98	58.68	45.7
Avg.	22.17	450.00	5.00	127.13	134.39	134.32	109.17	89.00	13.60	15.00	14.40	9.00	6.80	52.91	51.97	52.69	51.80	50.64
stdev	1.57	20.00	0.00	14.57	12.20	11.58	8.86	7.54	2.07	1.87	1.52	1.41	2.28	7.37	5.48	2.83	5.85	5.44

Table S2: Lateral and intermediate vein width quantification across the medio-lateral axis of leaves three and four.