OsNF-YB7 inactivates OsGLK1 to inhibit chlorophyll biosynthesis in rice embryo

Zongju Yang[1,2,†], Tianqi Bai[1,†], Zhiguo E[3,†], Baixiao Niu[1,2], Chen Chen[1,2*]

[1]Jiangsu Key Laboratory of Crop Genomics and Molecular Breeding/ Zhongshan Biological

Breeding Laboratory, Agricultural College of Yangzhou University, Yangzhou, China

[2]Jiangsu Co-Innovation Center for Modern Production Technology of Grain Crops/ Key

Laboratory of Plant Functional Genomics of the Ministry of Education, Agricultural College of

Yangzhou University, Yangzhou, China

[3] Key Laboratory of Rice Biology, China National Rice Research Institute, Hangzhou, China

[†] These authors contributed equally to this work

[*] Correspondence: chenchen@yzu.edu.cn

Abstract

As a master regulator of seed development, Leafy Cotyledon 1 (LEC1) promotes chlorophyll (Chl) biosynthesis in *Arabidopsis*, but the mechanism underlying this remains poorly understood. Here, we found that loss of function of *OsNF-YB7*, a *LEC1* homolog of rice, leads to chlorophyllous embryo, indicating that *OsNF-YB7* plays an opposite role in Chl biosynthesis in rice compared with that in *Arabidopsis*. OsNF-YB7 regulates the expression of a group of genes responsible for Chl biosynthesis and photosynthesis by directly binding to their promoters. In addition, OsNF-YB7 interacts with Golden 2-Like 1 (OsGLK1) to inhibit the transactivation activity of OsGLK1, a key regulator of Chl biosynthesis. Moreover, OsNF-YB7 can directly repress *OsGLK1* expression by recognizing its promoter *in vivo*, indicating the involvement of OsNF-YB7 in multiple regulatory layers of Chl biosynthesis in rice embryo. We propose that OsNF-YB7 functions as a transcriptional repressor to regulate Chl biosynthesis in rice embryo.

Introduction

Angiosperms can be divided into chloroembryophytes and leucoembryophytes, depending on the presence or absence of chlorophyll (Chl) in the embryo, respectively (Puthur et al., 2013; Smolikova and Medvedev, 2016). Some plant species such as *Arabidopsis* (*A. thaliana*) produce chloroembryos, which have photochemically active chloroplasts capable of producing assimilates that are further converted into reserve biopolymers (Simkin et al., 2020). Grass species, such as rice (*Oryza sativa*), cannot produce Chl when their embryos develop. To the best of our knowledge, it remains completely unclear what determines the ability to biosynthesize Chl in plant embryo. However, studies have identified several genes that contribute to Chl degradation for chloroembryos, the mutation of which can lead to a stay-green phenotype in mature seed like Mendel's green peas (Sato et al., 2007; Delmas et al., 2013; Wang et al., 2018; Li et al., 2017; Thomas and Ougham, 2014).

Golden 2-Like (*GLK*) genes encode GARP-type transcription factors (TFs), which are key components regulating the development of plant chloroplasts and Chl biosynthesis (Rossini et al., 2001; Fitter et al., 2002; Nakamura et al., 2009). GLK can recognize the CCAATC

cis-element of the Chl biosynthesis- and photosynthesis-associated nuclear genes to trigger

their expression (Waters et al., 2009). G-box (CACGTG) is also enriched in the GLK-targeted

genes in *Arabidopsis*, possibly due to GLK being able to interact with certain G-box binding

factors (Tamai et al., 2002). A genetic study showed that GLK activates Chl biosynthesis in

roots, in a manner dependent on the G-box binding TF Elongated Hypocotyl 5 (HY5)

(Kobayashi et al., 2012).

Like most plant species, rice has two *GLK* copies, designated *OsGLK1*, and *OsGLK2* (Fitter

et al., 2002). They redundantly regulate a set of genes, such as rice *Chlorophyllide A*

Oxygenase (*OsCAO*) and *Protochlorophyllide Oxidoreductase A* (*OsPORA*) responsible for

Chl biosynthesis, and *Light Harvesting Complex B1* (*OsLHCB1*) and *OsLHCB4* responsible for

photosynthesis (Nakamura et al., 2009; Wang et al., 2013; Sakuraba et al., 2017). *OsGLK1*

overexpression in rice leads to green calli and chloroplast development in the vascular

bundles (Nakamura et al., 2009; Wang et al., 2017). Organelle development in rice vascular

sheath cells is induced by ectopically expressed maize (*Zea mays*) *GLK* genes, mimicking a

key step in the evolutionary transition from C3 to C4 plants (Wang et al., 2017). In accordance

with this, the ectopic expression of maize *GLK*s in rice can boost biomass and grain yield by

facilitating chloroplast development and photosynthesis (Li et al., 2020; Yeh et al., 2022).

Moreover, the overexpression of maize *GLK*s in calli was shown to improve the ability of rice

and maize to regenerate (Luo et al., 2022). A recent study found that *OsGLK1* also participates

in tapetum plastid development and programmed cell death, consequently affecting pollen

fertility in rice (Zheng et al., 2022). The findings indicate that GLK plays multiple roles in

relation of plant development.

Leafy Cotyledon 1 (LEC1), a member of the nuclear factor Y (NF-Y) TF family, is a central

regulator controlling many aspects of seed development, such as Chl accumulation in embryo

(Meinke et al., 1994; Pelletier et al., 2017). LEC1 can also act as a pioneer TF to regulate

flowering by reprogramming the embryonic chromatin state (Tao et al., 2017). Previous studies

reported that *lec1* mutants have paler green embryos than wild-type (WT) at maturation in

Arabidopsis (Meinke et al., 1994; West et al., 1994). LEC1 transcriptionally regulates the

expression of genes that encode light-reaction components of photosystems I and II, as well

as the expression of genes involved in chloroplast biogenesis in *Arabidopsis* and soybean

embryos (Pelletier et al., 2017; Jo et al., 2020). These findings suggest that LEC1 is important

in photosynthesis and chloroplast development during seed development. However, the

molecular mechanisms underlying this remain largely unclear.

It is still a mysterious why plant species like rice cannot, whereas species like Arabidopsis

can, synthesize Chl in embryos. There are two LEC1 homologs, *OsNF-YB7* and *OsNF-YB9*,

encoded by the rice genome (E et al., 2018). *OsNF-YB7* is restricted to the embryo, and

defective *OsNF-YB7* may result in seed lethality (Niu et al., 2021b; Zhang and Xue, 2013).

Here, we found that OsNF-YB7 acts as a key inhibitor of Chl biosynthesis in rice embryo.

OsNF-YB7 inactivates the transactivation activity of OsGLK1, at multiple regulatory layers, to

inhibit Chl accumulation in the embryo of rice, explaining achlorophyllous embryo produced in

rice.

Results

Loss of function of *OsNF-YB7* leads to chloroembryos

By observing seeds produced by the loss-of-function mutant of *OsNF-YB7*, we surprisingly

found that the *osnf-yb7* embryo was greenish (**Fig. 1A and B**), suggesting that *OsNF-YB7*

plays a negative role in chloroplast biogenesis or Chl biosynthesis, or both, during

embryogenesis. We therefore examined the phenotype of green embryos at various seed

development stages. The results showed that WT had an achlorophyllous embryo throughout

embryonic development, whereas *osnf-yb7* embryo turned green at 5 days after fertilization

(DAF) and the chloroembryo remained green until maturity (**Fig. S1A—C**). Supporting this,

transmission electron microscopy (TEM) showed well-developed chloroplast in the scutellum

tip of the mutant, but this was not seen in the WT sections (**Fig. 1C**). By measuring Chl

contents in the developing and mature embryos, we found that total Chl content in *osnf-yb7*

was consistently higher than that in the WT, both at 10 DAF and at maturation (30 DAF) (**Fig.**

1D). As revealed by confocal laser scanning microscopy (CLSM), Chl autofluorescence was

detectable as early as 5 DAF in *osnf-yb7*, indicating the initiation of Chl accumulation at this

stage (**Fig. 1E**). This is consistent with our previous finding that *OsNF-YB7* was highly

activated at 5 DAF in rice embryo (Niu et al., 2021b).

Light is a critical signal triggering Chl biosynthesis (Wietrzynski and Engel, 2021). To

determine whether the Chl biosynthesis in *osnf-yb7* is induced by light, we investigated the

embryo phenotype of WT and *osnf-yb7* in the dark, using aluminum foil to cover rice panicles

prior to flowering. The result showed that, similar to WT, *osnf-yb7* embryo was achlorophyllous

in the dark, although the embryogenesis defects, such as degenerated epiblast and coleorhiza,

and maldeveloped coleoptile, were still observable (**Fig. S2A—H**). This indicated that Chl

biosynthesis in mutant is light dependent. Only a small amount of light can be perceived by

rice embryos because the external hulls block penetration of light (Simkin et al., 2020). We

removed the hulls to directly expose the embryo to light; however, Chl accumulation still failed

to occur in the WT embryo (**Fig. S2I—L**), suggesting that there are internal signals that repress

Chl biosynthesis in rice embryo. We suspected that *OsNF-YB7* mutation may attenuate the

activity of such inhibitors; alternatively, OsNF-YB7 itself could be an inhibitor.

OsNF-YB7 negatively regulates photosynthesis- and Chl biosynthesis-related genes

Using 5- and 10-DAF-old WT and *osnf-yb7* embryos, we previously performed deep

sequencing of the transcriptome (RNA-seq) to identify possible downstream genes of

OsNF-YB7 (Niu et al., 2021a). As revealed by Mapman analysis, photosynthesis-related

pathways, such as photosystem I, photosystem II, and the light reaction, were significantly

enriched for the differentially expressed genes (DEGs) in the mutant embryos (**Fig. 1F, Fig.**

S3A and B). Moreover, 96.9% (95/98) and 96.4% (106/110) photosynthesis-related DEGs that

identified from 5- and 10–DAF-old embryos, respectively, were upregulated in the mutant (**Fig.**

1G). To confirm this finding, we next examined the expression of *OsLHCA*s and *OsLHCB*s,

which are primarily associated with photosystems I and II, respectively, via quantitative

real-time PCR (RT-qPCR). It showed higher expression of all of the studied *OsLHCA*s and

*OsLHCB*s in the *osnf-yb7* embryo at 10 DAF (**Fig. S4A**). Likewise, many of the genes

participating in Chl biosynthesis, including rice *Genomes Uncoupled 4* (*OsGUN4*),

Mg-Chelatase H Subunit (*OsCHLH*), *OsCHLI*, *OsCHLD*, *Copper Response Defect 1*

(*OsCRD1*), *OsPORA*, *OsPORB*, and *Divinyl Reductase* (*OsDVR*), were significantly activated

in the mutant (**Fig. 1H and Fig. S4B**). The findings indicated that OsNF-YB7 might act as a

repressor of Chl biosynthesis and photosynthesis, which is the opposite of the role of its

homolog, LEC1, in *Arabidopsis.*

Because OsNF-YB7 is a TF, we assumed that it may directly regulate the expression of

genes related to Chl biosynthesis and photosynthesis, such as *OsPORA*, and *OsLHCB4*,

which were significantly activated in the *osnf-yb7* embryo (**Fig. 2A and B**). We first generated

transgenic lines that overexpressed *OsNF-YB7*, tagged with either green fluorescence protein

(*NF-YB7-GFP*) or 3× Flag (*NF-YB7-Flag*) in Zhonghua11 (ZH11, *O. sativa* ssp. *geng*/*japonica*)

or Kitaake (*O. sativa* ssp. *geng*/*japonica*) background, respectively. As expected, *OsPORA*

and *OsLHCB4* were significantly downregulated in leaves of the *OsNF-YB7* overexpressors

(**Fig. 2C and D, and Fig. S4C**). Similar to previously reported (Zhang and Xue, 2013; Ito et al.,

2011), the OsNF-YB7 overexpression lines displayed severe reproductive development

defects, preventing us to obtain enough embryo tissues for subsequent experiments. Instead,

using the *NF-YB7-Flag* seedling, we conducted a chromatin immunoprecipitation assay

coupled with quantitative PCR (ChIP-qPCR). The results showed that OsNF-YB7 was highly

enriched in the promoter regions of *OsPORA* and *OsLHCB4* harboring the G-box motif (**Fig.**

2E and F), a putative binding site of OsNF-YB7 (Guo et al., 2022). To confirm the ability of

OsNF-YB7 to bind to the *OsPORA* and *OsLHCB4* promoters, we next performed

electrophoretic mobility-shift assays (EMSAs), using recombinant OsNF-YB7-His protein, and

biotin-labeled subfragments of the *OsPORA* or *OsLHCB4* promoters containing G-boxes. The

results showed that OsNF-YB7-His was able to bind to the labeled probes, and the shifted

band signals were substantially weakened upon application of the unlabeled cold probes or

hot probes with a mutated G-box (**Fig. 2G and H**).

To confirm that the binding of OsNF-YB7 represses target gene expression, dual luciferase

(LUC) reporter (DLR) assays were performed. We first generated LUC reporters driven by the

OsPORA or *OsLHCB4* promoters (designated as *proOsPORA:LUC* and *proOsLHCB4:LUC*

hereafter) (**Fig. 2I**). When the reporters were coexpressed with OsNF-YB7, which acted as an

effector, in rice protoplast, we found that OsNF-YB7 significantly repressed the activity of both

proOsPORA:LUC and *proOsLHCB4:LUC* (**Fig. 2J**). Taken together, these results suggested

that OsNF-YB7 directly binds to the promoters of photosynthesis- and Chl biosynthesis-related

genes and represses their transcription.

OsNF-YB7 represses *OsGLK1* in the embryo

Several TFs that regulate Chl biosynthesis or chloroplast development have been identified in

plants (Jarvis and López-Juez, 2013). Some of these, such as rice *OsGLK1*, *HY5-like 1*

(*OsHY5L1*), *PIF-like 14* (*OsPIL14*), and *GATA Nitrate-inducible Carbon-metabolism-involved*

(*OsGNC*), were found to be upregulated in the embryo of *osnf-yb7* (**Fig. 3A**).

OsGLK1 was the most strikingly activated TF (**Fig. 3A and B**); meanwhile, it was

significantly repressed *in NF-YB7-GFP* (**Fig. 3C**). To test the idea that *OsGLK1* is a direct

downstream target of OsNF-YB7, a ChIP-qPCR assay was first carried out using *NF-YB7-Flag*

transgenic seedlings. The results showed that the promoter segments P3 and P4 were

significantly enriched in the immunoprecipitated chromatin (**Fig. 3D**), suggesting that

OsNF-YB7 was able to bind to the *OsGLK1* promoter *in vivo*. However, we failed to validate

the binding ability *in vitro* using the EMSA assay (**Fig. S5**), suggesting that OsNF-YB7 requires

other NF-Ys or TFs for forming a TF complex to recognize the promoter, like many NF-Y

members act for function (Laloum et al., 2013). A DLR assay was next performed to

investigate the negative regulation of OsNF-YB7 on *OsGLK1* transcription. The *OsGLK1*

promoter was inserted upstream of 5× upstream activating sequence (UAS) as a reporter;

OsNF-YB7 was C-terminally fused with the DNA binding domain (BD) of the yeast GAL4 and

the herpes virus VP16 transactivation domain (VP16:OsNF-YB7), as an effector (**Fig. 3E**).

Coexpression of VP16:OsNF-YB7 with the reporter in rice protoplast significantly decreased

VP16's transcriptional activity (**Fig. 3F**), indicating that OsNF-YB7 represses *OsGLK1*

promoter activity.

OsGLK1 is involved in OsNF-YB7-regulated Chl biosynthesis in embryo

To confirm the contribution of *OsGLK1* for the production of chloroembryo in *osnf-yb7*, we first

generated *OsGLK1*-overexpressing lines (Os*GLK1-OX*) driven by the rice ubiquitin promoter.

In association with over-accumulated Chl in the glume and seed coat of the transformant,

RT-qPCR and Western blot assays confirmed *OsGLK1* activation in *OsGLK1-OX* (**Fig.**

S6A—C). As observed in *osnf-yb7*, green embryos were produced (**Fig. 4A**). The Chl content

in *OsGLK-OX* were higher than that in WT (**Fig. 4B and C**), suggesting that *OsGLK1*

overexpression in rice embryo induces Chl biosynthesis.

Using a clustered regularly interspaced short palindromic repeats

(CRISPR)/CRISPR-associated 9 (Cas9) cassette including three tandemly arrayed guide

RNAs targeting *OsNF-YB7*, *OsGLK1,* and *OsGLK2*, respectively, we successfully obtained the

osnf-yb7;*osglk2* double mutant and the *osnf-yb7*;*osglk1*;*osglk2* triple mutant (**Fig. S7A and B**).

By analyzing the embryos that the mutants produced, we found that significantly less Chl

accumulated in *osnf-yb7*;*osglk1*;*osglk2* than in *osnf-yb7* and *osnf-yb7*;*osglk2* (**Fig. 4D–G and**

Fig. S8). In comparison to the achlorophyllous embryo of WT, the *osnf-yb7*;*osglk1*;*osglk2* triple

mutant still showed somewhat green coloration in the apical part of the embryos (**Fig. 4G**). We

hypothesized that this was at least partially due to that genes like *OsLHCB4* and *OsPORA* can

be induced by the mutation of *OsNF-YB7*, given the fact that OsNF-YB7 represses the genes'

expression independent of OsGLK1 (**Fig. 2J**).

OsNF-YB7 and OsGLK1 regulate a common set of genes in the embryo

We next performed RNA-seq analysis to explore the transcriptomic changes in the

chloroembryo of *OsGLK1-OX* at 10 DAF. More than 64.4% of the DEGs identified from

OsGLK1-OX were overlapped with the ones identified from *osnf-yb7*, vast majority of which

were either activated (60.6%) or repressed (27.6%) in both *OsGLK1-OX* and *osnf-yb7* (**Fig. 4H**

and I, and Table S1 and S2). As revealed by the Gene Ontology (GO) analysis, genes

involved in Chl biosynthesis and photosynthesis, such as *OsPORA* and *OsLHCB4*, were

markedly enriched among the common DEGs (**Fig. 4J and Fig. S9**), implying that OsNF-YB7

and OsGLK1 antagonically regulate a common set of genes for Chl biosynthesis and

photosynthesis.

The ChIP-qPCR results showed that OsGLK1 associated with the regions of the *OsPORA*

and *OsLHCB4* promoters to which OsNF-YB7 binds (**Fig. 5A and B**). In agreement with this,

the EMSA results suggested that OsGLK1 directly binds to the same DNA probes of *OsPORA*

and *OsLHCB4 in vitro* (**Fig. 5C and D**). In opposite to OsNF-YB7, when we coexpressed the

OsGLK1 effector vector with the reporter vector *proOsLHCB4:LUC* or *proOsPORA:LUC* in rice

protoplast, OsGLK1 showed significant transactivation activity on *OsPORA* and *OsLHCB4*

(**Fig. S10A and B**).

Recently, the putative binding sites of OsNF-YB7 and OsGLK1 were investigated at the

whole-genome scale (Guo et al., 2022; Tu et al., 2022). This allows testing of our hypothesis

that OsNF-YB7 and OsGLK1 can target a common set of genes involved in Chl biosynthesis

and photosynthesis. By reanalyzing the ChIP-seq data, we found that 91.4% (235/257) and

90.7% (167/184) of the OsGLK1-binding genes overlapped with the OsNF-YB7-binding genes

in two replications (**Fig. 5E and Table S3**), although the peak number of OsNF-YB7 was much

greater than that of OsGLK1 (Guo et al., 2022; Tu et al., 2022). A large number of the common

targets were activated in the embryos of *osnf-yb7* and *OsGLK1-OX* at 10 DAF (**Fig. S11A**).

GO analysis suggested that most of the common targets are genes involved in Chl

biosynthesis or photosynthesis (**Fig. S11B**). By retrieving the sequences of the OsNF-YB7 and

OsGLK1 binding peaks in the common targets for MEME analysis, we found that the TFs

probably recognize similar DNA motifs. A short sequence containing the G-box motif was the

most significantly enriched (**Fig. S12A and B**). There was also enrichment of another

sequence containing a CCAAT motif recognized by the NF-Y TF complexes and a CCAATC

motif recognized by GLKs, as previously reported (Waters et al., 2009; Pelletier et al., 2017)

(**Fig. S12A and B**). The results suggested that OsNF-YB7 and OsGLK1 can bind to the same

region of their common target. In agreement with this, approximately 75.0% and 76.5% of the

peak summits of OsGLK1 were located proximally to the summit of OsNF-YB7 in two

replications, within an adjacent region no more than 100 bp away (**Fig. 5F and Table S3**).

Consistent with our biochemical evidence, the ChIP-seq results showed that OsNF-YB7 does

bind to the same regions of the *OsPORA* and *OsLHCB4* promoters as OsGLK1 binds to (**Fig.**

S12C).

OsNF-YB7 physically interacts with OsGLK1 to inhibit its transcriptional activity

To resolve how OsNF-YB7 and OsGLK1 bind to the same regions to regulate their common

targets, we speculate that the TFs probably form a dimer in rice in order to exert their functions.

We therefore carried out a yeast two-hybrid (Y2H) assay by transforming yeast cells with a bait

construct expressing OsNF-YB7 fused with the GAL4 DNA binding domain (BD), together with

a prey construct expressing OsGLK1 fused with the yeast GAL4 activation domain (AD). The

results showed that OsNF-YB7 interacts with OsGLK1 in yeast (**Fig. 6A and B**). Furthermore,

to determine the functional domains required for the interaction, we generated two truncated

versions of OsNF-YB7, which contained the N- or C-terminal, and three truncated versions of

OsGLK1, which contained the N-terminal, DNA-BD, or GCT box domain, as previously

reported (Fitter et al., 2002; Zhang et al., 2021b) (**Fig. 6A**). We found that the C-terminal of

OsNF-YB7 was sufficient for the interaction (**Fig. 6B**). In addition, the GCT box domain of

OsGLK1 strongly interacted with the full length or C-terminal of OsNF-YB7, while the DNA-BD

of OsGLK1 showed a weak interaction (**Fig. 6B**).

A split complementary LUC assay further confirmed the interaction between OsNF-YB7 and

OsGLK1 in the epidermal cells of *Nicotiana benthamiana* (**Fig. 6C**). As suggested by the

bimolecular fluorescence complementation (BiFC) analysis, the interaction occurred

exclusively in the nucleus (**Fig. 6D**). Moreover, we transiently coexpressed OsNF-YB7 tagged

with GFP (OsNF-YB7-GFP) and OsGLK1 tagged with 3× Flag (OsGLK1-Flag) in rice

protoplast, and co-immunoprecipitation (Co-IP) analysis showed that OsGLK1-Flag could be

immunoprecipitated by the anti-GFP antibody (**Fig. 6E**), indicating that the interactions do

occur *in vivo*. In addition, both Y2H and split complementary LUC assays showed that

OsGLK2 could interact with OsNF-YB7 (**Fig. S13A and B**). These findings indicated that

OsNF-YB7 interacts with OsGLKs, explaining why OsNF-YB7 and OsGLKs share a common

set of targets in rice.

To explore the biological meaning of the interaction, we next performed DLR assays.

Transient expression of OsGLK1 in rice protoplast substantially activated the reporters, driven

by either the *OsPORA* or the Os*LHCB4* promoter; however, when we coexpressed OsNF-YB7

with OsGLK1, the transactivation ability of OsGLK1 was significantly repressed in rice

protoplasts (**Fig. 6F**). These findings suggested that OsNF-YB7/OsGLK1 dimerization reduces

the transactivation ability of OsGLK1 for fine-tuning the Chl biosynthetic and photosynthetic

genes, such as *OsPORA* and Os*LHCB4*. The EMSA assay showed that OsGLK1-MBP

recombinant proteins could bind to the promoter of *OsPORA*; however, when we incubated the

probe with OsGLK1-MBP and OsNF-YB7-His together, the binding ability of OsGLK1

substantially decreased (**Fig. 6G**). As a control, when the probe was incubated with

OsGLK1-MBP and GST-His, the binding ability of OsGLK1 remained unchanged (**Fig. 6G**).

These findings implied that the reduced transactivity of OsGLK1 is likely due to the formation

of OsNF-YB7/OsGLK1 heterodimers inhibiting the binding of OsGLK1 to its downstream

genes.

Discussion

By surveying 1,094 species from 666 genera and 182 families, Yakovlev and Zhukova (1980)

found that 428 angiosperms produce embryos with the presence of chlorophyll. Embryo

photosynthesis contributes a large amount of oxygen to fuel energy-generating pathways in

seed (Simkin et al., 2020). Some algae and gymnosperm species have evolved an ability to

synthesize Chl in darkness (Myers, 1940; Ranade et al., 2019; Bogorad, 1950). However, it

has remained largely unclear whether access to light accounts for the induction of

chloroembryos (Dahlgren, 1980; Periasamy and Vivekanandan, 1981; Liu et al., 2017). Here,

we found that removing the tissues covering an embryo failed to produce chlorophyllous

embryos in the WT; however, light avoidance inhibits Chl accumulation in *osnf-yb7* embryos

(**Fig. S2A—H**). These results suggest that light is necessary but insufficient to trigger

chloroplast biogenesis in rice. Because light itself does not induce Chl accumulation in rice

embryo (**Fig. S2I—L**), we inferred that there are intrinsic cues to repress chloroplast

development in rice embryo. However, to the best of our knowledge, the underlying

mechanism that determines an embryo's ability to synthesize Chl is completely unknown. Here,

we showed that OsNF-YB7, a LEC1 homolog of rice, acts as an inhibitor to repress Chl

accumulation in embryo. In line with this, a recent independent study also showed that

OsNF-YB7 null mutant accumulates Chl in embryo, although the cause of this remains

unresolved (Guo et al., 2022).

Several TFs involved in chloroplast development and photomorphogenesis in rice have

been identified (Nakamura et al., 2009; Li et al., 2019; Bai et al., 2019; Hudson et al., 2013).

Interestingly, we found that many of them were upregulated in the *osnf-yb7* embryos. For

example, *OsGLK1* was shown to be the most activated (**Fig. 3A and B**). Our genetic and

biochemical evidence suggests that *OsGLK1* is involved in the OsNF-YB7-mediated

repression of Chl biosynthesis in rice embryo. *OsGLK1* overexpression mimicked the

chloroembryo phenotype in WT, while knockout of *OsGLK1* and *OsGLK2*, simultaneously,

suppressed Chl accumulation in *osnf-yb7* embryo (**Fig. 4A—G**). The biochemical experiments

suggested that OsNF-YB7 associates with the promoter of *OsGLK1*, *in vivo*, to

transcriptionally inactivate *OsGLK1* (**Fig. 3D–F**), indicating that *OsGLK1* is a downstream

target of OsNF-YB7. However, OsNF-YB7 alone failed to bind to the promoter of *OsGLK1 in*

vitro (**Fig. S5**), presumably due to some as-yet-unidentified TFs being recruited by OsNF-YB7,

assisting in recognizing *OsGLK1* for transcriptional regulation. An NF-Y TF complex usually

consists of three subunits (NF-YA/B/C) in order to exert its function (Laloum et al., 2013).

NF-YA is responsible for DNA binding, while NF-YB and NF-YC are primarily responsible for

transactivation (Chaves-Sanjuan et al., 2021; Gnesutta et al., 2017). OsNF-YB7 possibly

forms a heterotrimer with NF-YA and NF-YC in the embryo to recognize *OsGLK1*. In addition,

previous studies showed that *Arabidopsis* LEC1 can interact with different TFs for

developmental regulation (Huang et al., 2015b; Yamamoto et al., 2009; Fatihi et al., 2016;

Boulard et al., 2018; Huang et al., 2015a). Identifying such TFs may reveal how OsNF-YB7

recognizes *OsGLK1* in order to exert its function.

OsNF-YB7 could recognize a common set of genes involved in Chl biosynthesis and

photosynthesis that are recognized by OsGLK1 (**Fig. 2E—H and Table S3**). The biochemical

results suggested that OsNF-YB7 can directly repress *OsPORA* and Os*LHCB4*, which are

activated by OsGLK1 (**Fig. 2J, and Fig. S10**). OsNF-YB7 probably hinders OsGLK1 to access

the target genes by forming an OsGLK1/OsNF-YB7 heterodimer (**Fig. 6A—E**), or by

occupying the motif OsGLK1 recognized in the promoter, given the binding site of OsGLK1

and OsNF-YB7 are likely overlapped (**Fig. 5F, Fig. S12 and Table S3**). The first of these

hypotheses is more plausible because, upon co-incubation of OsGLK1 and OsNF-YB7 with

the *OsPORA* promoter *in vitro*, the shifted band signal of OsGLK1 was substantially decreased

(**Fig. 6G**), indicating that the protein-protein interactions overwhelm the protein-DNA

interactions. Thus, when OsNF-YB7 is expressed, it interacts with OsGLK1, then OsGLK1 is

less available to activate downstream targets. The findings suggested that OsNF-YB7 plays a

dual role in regulating Chl biosynthesis in rice embryo: first, it represses the downstream

genes, achieved via its function as a transcriptional inactivator; second, OsNF-YB7 can

interact with OsGLKs to disturb their abilities to transactivate genes related to Chl biosynthesis

and photosynthesis (**Fig. S14**). In addition to OsNF-YB7, a recent study showed that Deep

Green Panicle 1, a plant-specific protein with a conserved TIGR01589 domain, can interact

with OsGLKs to suppress OsGLK-mediated transcription (Zhang et al., 2021a). The findings

suggested that, as the central regulator responsible for chloroplast development, GLK is tightly

regulated at the post-translational level to finetune Chl biosynthesis in plants.

Previous studies suggested that *Arabidopsis* LEC1 is a positive regulator of Chl biosynthesis

(Pelletier et al., 2017), given that the mature embryos of *lec1* were paler than the WT (Meinke,

1992; West et al., 1994). However, null mutation of *OsNF-YB7* activated Chl biosynthesis,

implying that the LEC1-type gene acts as a negative regulator in rice. By surveying the

literature, we noticed that Meinke reported that the cotyledons of *lec1* mutant remained green

unusually late in development (Meinke, 1992). Although there was no significant difference in

Chl content upon using whole seeds for quantification, Parcy *et al.* did observe that the tip of

lec1 cotyledons accumulates more Chl (Parcy et al., 1997). Moreover, the *lec1*;*abi3*

double-mutant embryos produced much more Chl than the *abi3* single mutant (Parcy et al.,

1997). Pelletier *et al.* recently reported that a cluster of LEC1 targets are enriched with

photosynthesis and chloroplast development related genes, and many of them are

down-regulated in the embryo of *lec1* (Pelletier et al., 2017); however, we noticed that in their

dataset, the photosynthesis and Chl biosynthesis related genes were more strikingly enriched

in the up-regulated genes of *lec1*, at either the mature green or postmature green stage.

These observations challenge the concept that LEC1 positively regulates Chl biosynthesis and

photosynthesis in *Arabidopsis*. Most studies on *Arabidopsis* have emphasized the importance

of *LEC1* in embryo development at the maturation stage. However, *LEC1* is activated within 24

h after fertilization (Lotan et al., 1998), but its role in the early embryo developmental stages for

Chl biosynthesis is still unknown.

In addition to producing chlorophyllous embryo, the *osnf-yb7* mutants display an array of

developmental defects, including abnormal embryogenesis, reduced dormancy, and

desiccation intolerance, similar to those found in *Arabidopsis lec1* mutants (Niu et al., 2021a).

Mutation of *osglk1* and *osglk2* in *osnf-yb7* could recover the chloroembyo phenotype; however,

the other embryo defects were not alleviated (**Fig. 4D—G**). The findings suggest that OsGLKs

specifically function in Chl biosynthesis, but OsNF-YB7 is responsible for many aspects of

embryo development. In agreement with this, the number of OsNF-YB7-targeted genes is far

greater than that of OsGLK-targeted genes, and the DEGs in *osnf-yb7* embryo is far greater

than that in *OsGLK1-OX* embryo (**Fig. 4H and Fig. 5E**). The mechanisms underlying the

OsNF-YB7-regulated multiple embryo developmental processes require further investigation.

Methods and Materials

Plant materials and growth conditions

The *osnf-yb7* mutant lines used in this study were previously generated in our laboratory (Niu

et al., 2021b). Rice cultivars Zhonghua11 (ZH11) and Kitaake (Kit) were used for gene

transformation. Rice plants were grown in a paddy field in Yangzhou, Jiangsu Province, China,

or in a climate-controlled room under long-day conditions with a photocycle of 14 h of light

(32°C) and 10 h of darkness (28°C), at 50% humidity. The *N. benthamiana* plants were grown

in a growth chamber at 22°C with long-day conditions (16 h light/8 h dark). To determine

whether the chloroembryo of *osnf-yb7* is light-dependent, the emerging panicles of the

mutants were covered with aluminum foil until seed maturation. The lemmas of the WT were

carefully removed with forceps at 1–2 DAF to expose the WT embryo to light in a

climate-controlled room.

Vector construction and plant transformation

To generate the overexpression plants, full-length coding sequences (CDSs) of *OsNF-YB7* or

OsGLK1 (the stop codon removed) were cloned into pCAMBIA1300-Flag or pUN1301-GFP

under the control of *ubiquitin* promoter, using the ClonExpress® II One Step Cloning Kit

(Vazyme). The constructs were transformed into rice calli through an *Agrobacterium*-mediated

strategy, as described previously (Chen et al., 2016). The higher-order mutants of *OsNF-YB7*,

OsGLK1 and *OsGLK2* mutants, were generated using a previously described method for

multiple gene editing in ZH11 (Cheng et al., 2021). The primers used for vector construction

are listed in **Table S4**.

RNA extraction and RT-qPCR

Embryos or seedlings were collected by flash freezing in liquid nitrogen and stored at −80 °C

until processing. Samples were finely ground using a mortar and pestle with liquid nitrogen.

Total RNA was isolated using the RNA-easy Isolation Reagent (Vazyme, R701-01). The

experiments were performed with at least three biological replicates. The relative expression

levels of the tested genes were normalized using the rice *Actin* gene and calculated by the $2^{\Delta Ct}$

method. The primers used for the RT-qPCR are listed in **Table S4**.

RNA-sequencing and data analysis

Ten-DAF-old embryos of the Wild-type (WT) and *OsGLK1-OX* were used for RNA-sequencing.

Two biological replicates were set. RNA extraction, library preparation, and high-throughput

sequencing of the collected samples were outsourced to BGI Genomics Co., Ltd., Shenzhen,

China. CLC Genomics Workbench 12.0 software was used for RNA-seq data analysis, as

previously reported (Xu et al., 2021). The thresholds of fold change > 2 and

Bonferroni-corrected *FDR* < 0.05 were used for defining a DEG. The previously generated

ChIP-seq data of OsNF-YB7 and OsGLK1 (Guo et al., 2022; Tu et al., 2022) were reanalyzed

using CLC Genomics Workbench 12.0 software for peak calling. Enriched motifs were

identified by the online tool MEME-ChIP (https://meme-suite.org/meme/tools/meme-chip) with

default parameter set. The software Mapman was used for pathway analysis (Usadel et al.,

2009). The online tool AgriGO 2.0 was used for GO analysis (Du et al., 2010). The Venn

diagrams   were   drawn   using   an   online   tool

(https://bioinfogp.cnb.csic.es/tools/venny/index.html).   TBtools   was   used   for   heat   map

generation (Chen et al., 2020).

Dual luciferase reporter (DLR) assays

The *OsGLK1*, *OsLHCB4* (2 kb upstream of translation start site), and *OsPORA* (1.5 kb

upstream of translation start site) promoter sequences were amplified from ZH11 genomic

DNA and cloned into the vector pGreenII 0800-LUC (Hellens et al., 2005), as reporters; the

35S::OsGLK1 and *35S::OsNF-YB7* constructs were used as effectors. The reporters and

effectors were transfected into rice etiolated protoplasts in different combinations and

incubated overnight. Firefly LUC and *Renilla* luciferase (REN) activities were measured using

the DLR Assay Kit (Vazyme), following the manufacturer's instructions, and the LUC:REN

ratios were calculated for analysis. The primers used for generating these constructs are listed

in **Table S4**.

Chromatin immunoprecipitation (ChIP) assays

OsGLK1-Flag and *OsNF-YB7-Flag* transgenic lines were used for the ChIP assays, in

accordance with a previously described method (Zhao et al., 2020). Briefly, 0.2 g of 14-day-old

seedlings were harvested and crosslinked with 1% formaldehyde for 15 min, followed by

neutralization using 0.125 M glycine for an additional 5 min. The seedlings were then ground

into powder in liquid nitrogen. The nuclei were isolated and lysed using Buffer S [50 mM

HEPES-KOH (pH 7.5), 150 mM NaCl, 1 mM ethylenediaminetetraacetic acid (EDTA), 1%

Triton X-100, 0.1% sodium deoxycholate, 1% SDS] and Buffer F [50 mM HEPES-KOH (pH

7.5), 150 mM NaCl, 1 mM EDTA, 1% Triton X-100, 0.1% sodium deoxycholate]. The chromatin

was then sonicated with the segment size ranging from 200 to 600 bp. The lysates were then

immunoprecipitated by anti-Flag antibody (Sigma no. F3165). Immunocomplexes were

washed with low-salt ChIP buffer (50 mM HEPES-KOH, 150 mM NaCl, 1 mM EDTA, 1% Triton

X-100, 0.1% sodium deoxycholate, 0.1% SDS), high-salt ChIP buffer (low-salt ChIP buffer but

replacing 150 mM NaCl with 350 mM NaCl), ChIP wash buffer (10 mM Tris-HCl pH 8.0, 250

mM LiCl, 0.5% NP-40, 1 mM EDTA, 0.1% sodium deoxycholate), and TE buffer (10 mM

Tris-HCl, pH 8.0, and 1 mM EDTA). The protein-DNA complexes were eluted from beads using

ChIP Elution buffer (50 mM Tris-HCl pH 7.5, 10 mM EDTA, 1% SDS) for 15 min at 65 °C and

the crosslinking was reversed by incubation overnight with proteinase K. The fragment DNA

was extracted with phenol:chloroform:isoamyl alcohol (25:24:1), precipitated with ethanol, and

resuspended in TE buffer. The immunoprecipitated DNA was used as a template for qPCR.

The primers used here are listed in **Table S4**.

Yeast two-hybrid assays

The CDSs of *OsGLK1/2* and *OsNF-YB7* were cloned into pGADT7 and pGBKT7, respectively.

The constructs were cotransformed into yeast strain AH109 using Frozen-EZ Yeast

Transformation II kit (Zymo), in accordance with the manufacturer's instructions. The empty

pGADT7 and pGBKT7 vectors were cotransformed in parallel as negative controls. The

transformants were first selected on synthetic dropout medium (SD/-Leu-Trp) plates. We

tested protein-protein interactions using selective SD/-Leu-Trp-His dropout medium.

Interactions were observed after 3 days of incubation at 28 °C. The primers used for

generating these constructs are listed in **Table S4**.

Split complementary LUC assays

Split complementary LUC assays were performed as previously described (Niu et al., 2020).

The CDSs of *OsGLK1/2* and *OsNF-YB7* were cloned into JW771 and JW772 vectors to

generate nLUC-OsGLK1/2 and cLUC-OsNF-YB7, respectively. The constructs were

introduced into *Agrobacterium tumefaciens* strain GV3101 and then co-infiltrated into *N.*

benthamiana leaves, and the LUC activities were analyzed after 48 h of infiltration using Tanon

Imaging System (5200 Multi; Tanon). The primers used for vector construction are shown in

Table S4.

Bimolecular fluorescence complementation assays

The CDSs of *OsNF*-YB7 and *OsGLK1* were cloned into the vector pSPYNE (nYFP) and

pSPYCE (cYFP), respectively. The prepared plasmids were transformed into *Agrobacterium*

strain GV3101, and the indicated transformant pairs were infiltrated into *N. benthamiana*

leaves. Forty-eight hours after infiltration, the fluorescence signal of yellow fluorescent protein

(YFP) was observed with a confocal laser scanning microscope (CLSM) (Carl Zeiss, LSM 710).

Images were captured at 514 nm laser excitation and 519–620 nm emission for YFP. The

primers used for vector construction are shown in **Table S4**.

Co-immunoprecipitation (Co-IP) assays

Co-IP assays were performed using rice protoplast as described previously (Zhang et al.,

2011). The CDSs (stop codons removed) of *OsGLK1* and *OsNF-YB7* were cloned into the

vectors pUC19-35S-FLAG-RBS and pJIT163-GFP driven by a 35S promoter, respectively. Ten

micrograms of plasmid DNA (OsGLK1-GFP, GFP, and OsNF-YB7-Flag) was transformed or

cotransformed into 200 µl of protoplasts and incubated in WI buffer (0.5 M mannitol, 20 mM

KCl, and 4 mM MES at pH 5.7) for 12 h. The protoplasts were collected and lysed in 500 µl of

lysis buffer [50 mM Tris-HCl, 150 mM NaCl, 5 mM EDTA (pH 8.0), 1% NP-40, 0.1 mM PMSF].

The extracts were incubated with GFP-Trap agarose beads at 4 °C for 3 h and washed three

times with washing buffer. Samples were boiled in SDS protein loading buffer. Immunoblots

were detected using corresponding primary antibodies (anti-GFP, ABclonal no. AE012;

anti-Flag, Sigma no. F3165). The primers used for vector construction are shown in **Table S4**.

Chl measurement and confocal imaging

Approximately one hundred micrograms of embryos of the indicated genotypes were extracted

in 1 ml of 100% dimethyl sulfoxide (DMSO) with incubation at 65 °C for 1 h. Then, the

absorbance values at wavelengths of 648.2 and 664.9 nm were measured by

spectrophotometry and total Chl content was calculated as reported previously(Barnes et al.,

1992). Chl autofluorescence signal was detected by CLSM (Carl Zeiss, LSM 710), with

excitation and emission wavelengths of 633 and 625–730 nm, respectively.

Electrophoretic mobility-shift assays (EMSAs)

OsNF-YB7 CDS was amplified by PCR and cloned into pET-28a vector to generate the

OsNF-YB7-His construct. The full-length CDS of *OsGLK1* was cloned into pMAL-c5X vector to

generate *OsGLK1-MBP* construct. All constructs were transformed into *E. coli* strain BL21 to

produce recombinant proteins. The promoter subfragments of *OsPORA* (42 bp, from -278 to

-237) and *OsLHCB4* (38 bp, from -259 to -222) were PCR amplified and labeled with biotin at

the 3' hydroxyl end of the double strands using EMSA Probe Biotin Labeling Kit (Beyotime,

GS008). EMSA was performed using EMSA/Gel-Shift kit (Beyotime, GS009), in accordance

with the manufacturer's instructions. The labeled probes were detected in accordance with the

instructions provided with the EMSA/Gel-Shift kit. All oligonucleotides used to generate the

biotin-labeled probes are listed in **Table S4**.

Transmission electron microscopy (TEM)

TEM analysis was performed as described previously (Cheng et al., 2021). Briefly, embryos of

WT and *osnf-yb7* were fixed overnight at 4°C in 2.5% glutaraldehyde and 0.1 M PBS. The

samples were subsequently washed three times with 0.1 M PBS and then fixed with 1% osmic

acid for 4 h. The samples were dehydrated in a series of ethanol and embedded in acrylic

resin at 37 °C for 12 h and at 60 °C for 48 h. The samples were sectioned at 100 nm and

observed by TEM (TECNAI 12).

Accession Numbers

Gene sequence data of this article can be found in the Rice Annotation Project Database

(RAP-DB) under the following accession numbers: *OsNF-YB7* (Os02g0725700), *OsGLK1*

(Os06g0348800), *OsGLK2* (Os01g0239000), *OsLHCA1* (Os06g0320500), *OsLHCA2*

(Os07g0577600), *OsLHCA3* (Os02g0197600), *OsLHCA4* (Os08g0435900), *OsLHCA5*

(Os02g0764500), *OsLHCA6* (Os09g0439500), *OsLHCB1.1* (Os01g0720500), *OsLHCB2*

(Os03g0592500), *OsLHCB3* (Os07g0562700), *OsLHCB4* (Os07g0558400), *OsLHCB5*

(Os11g0242800), *OsLHCB6* (Os04g0457000), *OsGUN4* (Os11g0267000), *OsCHLH*

(Os03g0323200), *OsCHLI* (Os03g0563300), *OsCHLD* (Os03g0811100), *OsCRD1*

(Os01g0363900), *OsDVR* (Os03g0351200), *OsPORA* (Os04g0678700), *OsGNC*

(Os06g0571800), *OsPIL14* (Os07g0143200).

Acknowledgements

We thank Prof. Hengxiu Yu and Dr. Chao Zhang of Yangzhou University for kindly providing

the positive and negative control vectors for dual luciferase reporter assays.

Funding

This research was supported by grants from the National Natural Science Foundation of China

(32170344, 32300689), the Project of Zhongshan Biological Breeding Laboratory

(BM2022008-02), the Jiangsu Province Government (JBGS001), the Natural Science

Foundation of the Higher Education Institutions of Jiangsu Province (21KJB21003), the

Independent Scientific Research Project Funds of the Jiangsu Key Laboratory of Crop

Genomics and Molecular Breeding (PLR202101), the Priority Academic Program

Development of Jiangsu Higher Education Institutions (PAPD).

Author contributions

C.C. conceived the project. C.C., Z.Y. and T.B. designed the research. T.B., Z.Y. and B.N

performed the experiments. Z.Y., T.B., Z.E. and C.C. analyzed the data. Z.Y. and C.C. wrote

the manuscript.

Declaration of interests

The authors declare no competing interests.

 Figure Legends

Figure 1. OsNF-YB7 negatively regulates Chlorophyll (Chl) biosynthesis in embryo.

A, B. Morphologies of wild-type (WT) and *osnf-yb7* detached embryos at 10 days after

fertilization (DAF) (**A**) and longitudinally dissected embryos at maturation (**B**). Scale bars = 200

μm.

C. Transmission electron microscopy images of embryos from WT and *osnf-yb7* at 10 DAF.

Scale bars = 5 μm.

D. Chl levels in WT and *osnf-yb7* embryos at 10 DAF and maturation. Data are means ± SD (n

= 3). **, $p < 0.01$; Student's *t*-test was used for statistical analysis.

E. Chl autofluorescence of WT and mutant embryos at 5 DAF. BF, bright field; Chl, Chl

autofluorescence. Scale bars = 100 μm.

F. Photosynthesis-related pathways enriched among the differentially expressed genes (DEGs)

identified from the 5- and 10-DAF-old *osnf-yb7* embryos compared to WT, respectively.

G. Most of the photosynthesis-related genes were upregulated in the mutant embryos.

H. A heat map shows the expression of the Chl biosynthesis-related genes in the WT and

osnf-yb7 embryos at 5 and 10 DAF. Reads per kilobase per million mapped reads (RPKM) is

used to indicate the expression level.

Figure 2. OsNF-YB7 binds to the promoters of *OsPORA* and *OsLHCB4* to regulate their

expression.

A, B. Quantitative real-time PCR (RT-qPCR) analysis of the transcription levels of *OsPORA* (**A**)

and *OsLHCB4* (**B**) in the embryos of WT and *osnf-yb7* at 10 DAF. Data are means ± SD (n = 3).

**, $p < 0.01$; Student's *t*-test was used for statistical analysis.

C, D. Expression of *OsPORA* (**C**) and *OsLHCB4* (**D**) in the leaves of WT and

OsNF-YB7-overexpressing transgenic plants (*NF-YB7-GFP*). Data are means ± SD (n = 3). **,

$p < 0.01$; Student's *t*-test was used for statistical analysis.

E, F. Chromatin immunoprecipitation assay coupled with quantitative PCR (ChIP-qPCR)

analyses showing the enrichment of OsNF-YB7 at the *OsPORA* (**E**) and *OsLHCB4* (**F**)

promoters in 14-day-old *OsNF-YB7-Flag* seedlings. Precipitated DNA was quantified by qPCR

and DNA enrichment is displayed as a percentage of input DNA. Data are means ± SD (n = 3).

*, $p < 0.05$; **, $p < 0.01$; Student's *t*-test was used for statistical analysis. *ACTIN* was used as a

nonspecific target gene. Diagrams in the upper panel showing the promoter structures of

OsPORA and *OsLHCB4*, and the PCR amplicons used for ChIP-qPCR.

G, H. Electrophoretic mobility-shift assays (EMSAs) showing that OsNF-YB7 directly binds to

the promoters of *OsPORA* (**I**) and *OsLHCB4* (**J**). Hot probes were biotin-labeled. The hot

mProbes contain mutant nucleic acid from CACATG to AAAAAA. The arrow heads indicate the

shift bands.

I. Schematic diagram displaying the constructs used in the dual luciferase reporter (DLR)

assays of **J**. LUC, firefly luciferase; REN, *Renilla* luciferase.

J. DLR assays showing that OsNF-YB7 directly represses the promoter activities of *OsPORA*

and *OsLHCB4*. Data are means ± SD (n = 3). *, $p < 0.05$; **, $p < 0.01$; Student's *t*-test was

used for statistical analysis.

Figure 3. OsNF-YB7 associates with the promoter of *OsGLK1* and represses its

expression.

A. A heat map showing the expression of transcription factors associated with Chl biosynthesis

and chloroplast development in the 5- and 10-DAF-old embryos of WT and *osnf-yb7*. The

colored dots indicate $\log_2$(RPKM mean) of the genes in three biological replicates.

B. RT-qPCR analysis of *OsGLK1*, *OsGNC*, and *OsPIL14* expression levels in the embryos

from WT and *osnf-yb7* at 5- and 10 DAF. Numbers represent fold changes of expression. Data

are means ± SD (n = 3). *, $p < 0.05$; **, $p < 0.01$; Student's *t*-test was used for statistical

analysis.

C. RT-qPCR analysis of *OsGLK1* expression levels in leaves from WT and *NF-YB7-GFP*. Data

are means ± SD (n = 3). **, $p < 0.01$; Student's *t*-test was used for statistical analysis.

D. ChIP-qPCR analysis showing the enrichment of OsNF-YB7 at the *OsGLK1* promoter in

14-day-old *OsNF-YB7-Flag* seedling. Chromatin of each sample was immunoprecipitated

using anti-Flag or IgG antibodies. Precipitated DNA was quantified by qPCR and DNA

enrichment is displayed as a percentage of input DNA. Data are means ± SD (n = 3). *, $p <$

0.05; **, $p < 0.01$; Student's *t*-test was used for statistical analysis. *ACTIN* was used as a

nonspecific target gene. The experiment was performed three times with similar results. The

diagram in the upper panel showing the promoter structure

of *OsGLK1* and PCR amplicons (P1, P2, P3, P4, and P5) used for ChIP-qPCR.

E. Schematic diagram displaying the constructs used in the DLR assays of **F**. LUC, firefly

luciferase; REN, *Renilla* luciferase; UAS, upstream activating sequence.

F. DLR assays showing that OsNF-YB7 represses the promoter activity of *OsGLK1*. Data are

means ± SD (n = 3). *, $p < 0.05$; Student's *t*-test was used for statistical analysis.

Figure 4. Chl biosynthesis in *osnf-yb7* embryo requires active *OsGLK*s.

A. Embryo morphologies of WT and Os*GLK1-OX* detached embryos at 10 DAF. Scale bars = 2

mm.

B. Chl autofluorescence of the WT and *GLK-OX* embryos at 10 DAF. BF, bright field; Chl, Chl

autofluorescence. Scale bars = 100 μm.

C. Chl levels in WT and *OsGLK1-OX* embryos at 10 DAF. Data are means ± SD (n = 3). **, $p <$

0.01; Student's *t*-test was used for statistical analysis.

D-G. Morphologies of the embryos produced by WT (**D**), *osnf-yb7* (**E**), *osnf-yb7*;*osglk2* double

mutant (**F**) and *osnf-yb7*;*osglk1*;*osglk2* triple mutant (**G**). Scale bars = 1 mm.

H. A Venn diagram showing overlaps of the DEGs identified from the embryos of *osnf-yb7* and

OsGLK1-OX at 10 DAF.

I. A pie chart showing similar transcriptional changes of the common DEGs identified from

osnf-yb7 and *OsGLK1-OX*.

J. Gene Ontology (GO) analysis of the common DEGs identified from *osnf-yb7* and

OsGLK1-OX.

Figure 5. OsNF-YB7 and OsGLK1 regulate a common set of genes in the embryo.

A, B. ChIP-qPCR analysis showing the enrichment of OsGLK1 on the *OsPORA* (**A**) and

OsLHCB4 (**B**) promoters. PCR amplicons used for ChIP-qPCR are indicated in the schematic

diagrams. OsGLK1-GFP was transiently expressed in protoplasts isolated from green tissues

of the 14-day-old WT seedling. Chromatin of each sample was immunoprecipitated using

anti-GFP or igG antibodies. Precipitated DNA was quantified by qPCR and DNA enrichment is

displayed as a percentage of input DNA. Data are means ± SD (n = 3). *, $p < 0.05$; **, $p < 0.01$;

Student's *t*-test was used for statistical analysis. *ACTIN* was used as a nonspecific target gene.

The experiment was performed three times with similar results.

C, D. EMSA assays showing that OsGLK1 directly binds to the promoters of *OsPORA* (**C**) and

OsLHCB4 (**D**). The arrow heads indicate the shift bands.

E. A Venn diagram showing overlaps of the target genes between OsGLK1 and OsNF-YB7.

F. Distribution of the distance between the OsNF-YB7 and OsGLK1 summits showing that

OsNF-YB7 and OsGLK1 bind to proximal regions of their common targets.

Figure 6. OsNF-YB7 interacts with OsGLK1 to regulate the expression of *OsPORA* and

OsLHCB4.

A. Schematic diagrams showing the protein structures of OsGLK1 and OsNF-YB7.

B. Yeast-two-hybrid (Y2H) assays showing the interaction between OsNF-YB7 and OsGLK1.

AD and BD indicate the activation domain and binding domain of GAL4, respectively. The full

length or truncated OsNF-YB7 and OsGLK1 were fused with BD and AD, respectively. The

indicated combinations of constructs were cotransformed into yeast cells and grown on the

nonselective medium SD/-L-W and selective medium SD/-L-W-H.

C. A split complementary luciferase (LUC) assay confirmed the interaction between OsNF-YB7

and OsGLK1. Coexpression of the fusion of OsGLK1 and the N-terminal half of LUC

(nLUC-OsGLK1) and the fusion of the C-terminal half of LUC and OsNF-YB7

(cLUC-OsNF-YB7) in the epidermal cells of *N. benthamiana* leaves induced LUC activities,

whereas the epidermal cells coexpressing nLUC-OsGLK1 and cLUC, nLUC and

cLUC-OsNF-YB7, or nLUC and cLUC did not show LUC activities.

D. Bimolecular fluorescence complementation (BiFC) assays showed interactions between

OsNF-YB7 and OsGLK1 in the nucleus. OsNF-YB7 was fused with the N-terminal of yellow

fluorescent protein (nYFP-OsNF-YB7); OsGLK1 was fused with the C-terminal of yellow

fluorescent protein (cYFP-OsGLK1). The indicated combinations of constructs were transiently

coexpressed in the leaf epidermal cells of *N. benthamiana.* Scale bar = 10 μm.

E. Co-immunoprecipitation (Co-IP) assays showing that OsNF-YB7 interacts with OsGLK1 *in*

vivo. 35S::OsNF-YB7:GFP (OsNF-YB7-GFP) and 35S::OsGLK1:3×Flag (OsGLK1-Flag) were

coexpressed in rice protoplasts and were immunoprecipitated with an anti-GFP antibody, and

the immunoblots were probed with anti-GFP and anti-Flag antibodies. 35S::GFP (GFP) was

used as a negative control.

F. DLR assays showing that OsNF-YB7 represses the transactivation activity of OsGLK1 on

OsPORA and *OsLHCB4*. Protoplasts isolated from etiolated seedlings were used for the

analyses. EV, empty vector. Data are means ± SD (n = 3). *, $p < 0.05$; **, $p < 0.01$; Student's

t-test was used for statistical analysis.

G. EMSA assay indicated that OsNF-YB7 inhibits the DNA binding of OsGLK1 to the promoter

of *OsPORA*. The black and white arrow heads indicate the OsGLK1- and OsNF-YB7-bound

probes, respectively; "+" and "++" indicate that 2- and 4 μM recombinant proteins were used

for the reactions. The GST-His was used as a negative control.

Supplementary Table 1. Differentially expressed genes (DEGs) in the 10-DAF-old embryos of

OsGLK1-OX.

Supplementary Table 2. Common DEGs identified from the *OsGLK1-OX* and *osnf-yb7*

embryos at 10 DAF.

Supplementary Table 3. Common targets of OsGLK1 and OsNF-YB7.

Supplementary Table 4. Primers used in this study.

Supplementary Figure 1. Mutation of *OsNF-YB7* leads to chloroembryo.

Supplementary Figure 2. Light is required but not sufficient for Chl biosynthesis in the

chloroembryo of *osnf-yb7*.

Supplementary Figure 3. Mutation of *OsNF-YB7* actives the expression of

photosynthesis-related genes.

Supplementary Figure 4. OsNF-YB7 negatively regulates the expression of light harvest and

Chl biosynthesis-associated genes.

Supplementary Figure 5. OsNF-YB7 does not directly binds to the promoter of *OsGLK1 in*

vitro.

Supplementary Figure 6. Overexpression of *OsGLK1* induces chloroembryo in rice.

Supplementary Figure 7. Generation of *osnf-yb7*, *osglk1* and *osglk2* high-order mutants.

Supplementary Figure 8. Chl levels in WT, *osnf-yb7* and *osnf-yb7*;*osglk1*;*osglk2* embryos at

10 DAF.

Supplementary Figure 9. Overexpression of *OsGLK1* activates the expression of Chl

biosynthesis- and photosynthesis-related genes.

Supplementary Figure 10. OsGLK1 significantly activates the promoter activities of *OsPORA*

and *OsLHCB4*.

Supplementary Figure 11. Many of the OsNF-YB7 and OsGLK1 common targets that

activated in *osnf-yb7* and *OsGLK1-OX* are involved in Chl biosynthesis and photosynthesis.

Supplementary Figure 12. OsNF-YB7 and OsGLK1 bind to proximal regions of their

common targets.

Supplementary Figure 13. OsNF-YB7 interacts with OsGLK2.

Supplementary Figure 14. A proposed model of the OsNF-YB7-mediated suppression of Chl

biosynthesis in rice embryo.

References

Bai, B., Lu, N., Li, Y., Guo, S., Yin, H., He, Y., Sun, W., Li, W., and Xie, X. (2019). OsBBX14

promotes photomorphogenesis in rice by activating OsHY5L1 expression under blue

light conditions. Plant Sci. **284**: 192–202.

Barnes, J.D., Balaguer, L., Manrique, E., Elvira, S., and Davison, A.W. (1992). A

reappraisal of the use of DMSO for the extraction and determination of chlorophylls a and

b in lichens and higher plants. Environ. Exp. Bot. **32**: 85–100.

Bogorad, L. (1950). Factors Associated with the Synthesis of Chlorophyll in the Dark in

Seedlings of Pinus jeffreyi. Bot. Gaz. **111**: 221–241.

Boulard, C., Thévenin, J., Tranquet, O., Laporte, V., Lepiniec, L., and Dubreucq, B.

(2018). LEC1 (NF-YB9) directly interacts with LEC2 to control gene expression in seed.

Biochim. Biophys. Acta - Gene Regul. Mech. **1861**: 443–450.

Chaves-Sanjuan, A., Gnesutta, N., Gobbini, A., Martignago, D., Bernardini, A., Fornara,

F., Mantovani, R., and Nardini, M. (2021). Structural determinants for NF-Y subunit

organization and NF-Y/DNA association in plants. Plant J. **105**: 49–61.

Chen, C., Begcy, K., Liu, K., Folsom, J.J., Wang, Z., Zhang, C., and Walia, H. (2016). Heat

stress yields a unique MADS box transcription factor in determining seed size and

thermal sensitivity. Plant Physiol. **171**: 606–622.

Chen, C., Chen, H., Zhang, Y., Thomas, H.R., Frank, M.H., He, Y., and Xia, R. (2020).

TBtools: An Integrative Toolkit Developed for Interactive Analyses of Big Biological Data.

Mol. Plant **13**: 1194–1202.

Cheng, X., Pan, M., E, Z., Zhou, Y., Niu, B., and Chen, C. (2021). The maternally expressed

polycomb group gene OsEMF2a is essential for endosperm cellularization and imprinting

in rice. Plant Commun. **2**: 100092.

Dahlgren, R. (1980). The taxonomic significance of chlorophyllous embryos in angiosperm

seeds. Bot. Not. **133**: 337–341.

Delmas, F., Sankaranarayanan, S., Deb, S., Widdup, E., Bournonville, C., Bollier, N.,

Northey, J.G.B., McCourt, P., and Samuel, M.A. (2013). ABI3 controls embryo

degreening through Mendel's i locus. Proc. Natl. Acad. Sci. U. S. A. **110**: 1–7.

Du, Z., Zhou, X., Ling, Y., Zhang, Z., and Su, Z. (2010). agriGO: a GO analysis toolkit for the

agricultural community. Nucleic Acids Res. **38**: W64–W70.

E, Z., Li, T., Zhang, H., Liu, Z., Deng, H., Sharma, S., Wei, X., Wang, L., Niu, B., and Chen,

C. (2018). A group of nuclear factor Y transcription factors are sub-functionalized during

endosperm development in monocots. J. Exp. Bot. **69**: 2495–2510.

Fatihi, A., Boulard, C., Bouyer, D., Baud, S., Dubreucq, B., and Lepiniec, L. (2016).

Deciphering and modifying LAFL transcriptional regulatory network in seed for improving

yield and quality of storage compounds. Plant Sci. **250**: 198–204.

Fitter, D.W., Martin, D.J., Copley, M.J., Scotland, R.W., and Langdale, J.A. (2002). GLK

gene pairs regulate chloroplast development in diverse plant species. Plant J. **31**: 713–

727.

Gnesutta, N., Saad, D., Chaves-Sanjuan, A., Mantovani, R., and Nardini, M. (2017).

Crystal Structure of the Arabidopsis thaliana L1L/NF-YC3 Histone-fold Dimer Reveals

Specificities of the LEC1 Family of NF-Y Subunits in Plants. Mol. Plant **10**: 645–648.

Guo, F. et al. (2022). Rice LEAFY COTYLEDON1 Hinders Embryo Greening During the Seed

Development. Front. Plant Sci. **13**: 1–18.

Hellens, R.P., Allan, A.C., Friel, E.N., Bolitho, K., Grafton, K., Templeton, M.D.,

Karunairetnam, S., Gleave, A.P., and Laing, W.A. (2005). Transient expression

vectors for functional genomics, quantification of promoter activity and RNA silencing in

plants. Plant Methods **1**: 13.

Huang, M., Hu, Y., Liu, X., Li, Y., and Hou, X. (2015a). Arabidopsis LEAFY COTYLEDON1

controls cell fate determination during post-embryonic development. Front. Plant Sci. **6**:

955.

Huang, M., Hu, Y., Liu, X., Li, Y., and Hou, X. (2015b). Arabidopsis LEAFY COTYLEDON1

mediates postembryonic development via interacting with

PHYTOCHROME-INTERACTING FACTOR4. Plant Cell **27**: 3099–3111.

Hudson, D., Guevara, D.R., Hand, A.J., Xu, Z., Hao, L., Chen, X., Zhu, T., Bi, Y.M., and

Rothstein, S.J. (2013). Rice cytokinin GATA transcription factor1 regulates chloroplast

development and plant architecture. Plant Physiol. **162**: 132–144.

Ito, Y., Thirumurugan, T., Serizawa, A., Hiratsu, K., Ohme-Takagi, M., and Kurata, N.

(2011). Aberrant vegetative and reproductive development by overexpression and

lethality by silencing of OsHAP3E in rice. Plant Sci. **181**: 105–110.

Jarvis, P. and López-Juez, E. (2013). Biogenesis and homeostasis of chloroplasts and other

plastids. Nat. Rev. Mol. Cell Biol. **14**: 787–802.

Jo, L., Pelletier, J.M., Hsu, S., Baden, R., Goldberg, R.B., and Harada, J.J. (2020).

Combinatorial interactions of the LEC1 transcription factor specify diverse developmental

programs during soybean seed development. Proc. Natl. Acad. Sci. **117**: 1223–1232.

Kobayashi, K., Baba, S., Obayashi, T., Sato, M., Toyooka, K., Keränen, M., Aro, E.M.,

Fukaki, H., Ohta, H., Sugimoto, K., and Masudaa, T. (2012). Regulation of root

greening by light and auxin/cytokinin signaling in Arabidopsis. Plant Cell **24**: 1081–1095.

Laloum, T., De Mita, S., Gamas, P., Baudin, M., and Niebel, A. (2013). CCAAT-box binding

transcription factors in plants: Y so many? Trends Plant Sci. **18**: 157–166.

Li, X., Wang, P., Li, J., Wei, S., Yan, Y., Yang, J., Zhao, M., Langdale, J.A., and Zhou, W.

(2020). Maize GOLDEN2-LIKE genes enhance biomass and grain yields in rice by

improving photosynthesis and reducing photoinhibition. Commun. Biol. **3**: 151.

Li, Z., Mo, W., Jia, L., Xu, Y.C., Tang, W., Yang, W., Guo, Y.L., and Lin, R. (2019). Rice

FLUORESCENT1 Is Involved in the Regulation of Chlorophyll. Plant Cell Physiol. **60**:

2307–2318.

Li, Z., Wu, S., Chen, J., Wang, X., Gao, J., Ren, G., and Kuai, B. (2017).

NYEs/SGRs-mediated chlorophyll degradation is critical for detoxification during seed

maturation in Arabidopsis. Plant J. **92**: 650–661.

Liu, H., Wang, X., Ren, K., Li, K., Wei, M., Wang, W., and Sheng, X. (2017). Light

deprivation-induced inhibition of chloroplast biogenesis does not arrest embryo

morphogenesis but strongly reduces the accumulation of storage reserves during

embryo maturation in arabidopsis. Front. Plant Sci. **8**: 1–14.

Lotan, T., Ohto, M., Yee, K.M., West, M.A.L., Lo, R., Kwong, R.W., Yamagishi, K., Fischer,

R.L., Goldberg, R.B., and Harada, J.J. (1998). Arabidopsis LEAFY COTYLEDON1 is

sufficient to induce embryo development in vegetative cells. Cell **93**: 1195–1205.

Luo, W., Tan, J., Li, T., Feng, Z., Ding, Z., Xie, X., Chen, Y., Chen, L., Liu, Y.-G., Zhu, Q.,

and Guo, J. (2022). Overexpression of maize GOLDEN2 in rice and maize calli improves

regeneration by activating chloroplast development. Sci. China Life Sci.

Meinke, D.W. (1992). A Homoeotic Mutant of Arabidopsis thaliana with Leafy Cotyledons.

Science **258**: 1647–50.

Meinke, D.W., Franzmann, L.H., Nickle, T.C., and Yeung, E.C. (1994). Leafy cotyledon

mutants of Arabidopsis. Plant Cell **6**: 1049–1064.

Myers, J. (1940). A Study Of The Pigments Produced In Darkness By Certain Green Algae.

Plant Physiol. **15**: 575–588.

Nakamura, H., Muramatsu, M., Hakata, M., Ueno, O., Nagamura, Y., Hirochika, H., Takano,

M., and Ichikawa, H. (2009). Ectopic overexpression of the transcription factor osglk1

induces chloroplast development in non-green rice cells. Plant Cell Physiol. **50**: 1933–

1949.

Niu, B., Deng, H., Li, T., Sharma, S., Yun, Q., Li, Q., E, Z., and Chen, C. (2020). OsbZIP76

interacts with OsNF‑YBs and regulates endosperm cellularization in rice (Oryza

sativa). J. Integr. Plant Biol. **62**: 1983–1996.

Niu, B., Yang, Z., Bai, T., E, Z., Xu, X., Yun, Q., Zhang, J., Zhang, Z., Lu, X., Li, Q., Liu,

Q.-Q., and Chen, C. (2021a). Rice LEAFY COTYLEDON1 regulates embryonic

envelope development and chlorophyll biogenesis in embryo.

doi.org/10.21203/rs.3.rs-888827/v1.

Niu, B., Zhang, Z., Zhang, J., Zhou, Y., and Chen, C. (2021b). The rice LEC1‑like

transcription factor OsNF‑YB9 interacts with SPK, an endosperm‑specific sucrose

synthase protein kinase, and functions in seed development. Plant J. **106**: 1233–1246.

Parcy, F., Valon, C., Kohara, A., Miséra, S., and Giraudat, J. (1997). The ABSCISIC

ACID-INSENSITIVE3, FUSCA3, and LEAFY COTYLEDON1 loci act in concert to control

multiple aspects of arabidopsis seed development. Plant Cell **9**: 1265–1277.

Pelletier, J.M., Kwong, R.W., Park, S., Le, B.H., Baden, R., Cagliari, A., Hashimoto, M.,

Munoz, M.D., Fischer, R.L., Goldberg, R.B., and Harada, J.J. (2017). LEC1

sequentially regulates the transcription of genes involved in diverse developmental

processes during seed development. Proc. Natl. Acad. Sci. **114**: E6710–E6719.

Periasamy, K. and Vivekanandan, M. (1981). Photosynthesis in the Chloroembryo of

Cyamopsis tetragonaloba Taub. Ann. Bot. **47**: 793–797.

Puthur, J.T., Shackira, A.M., Saradhi, P.P., and Bartels, D. (2013). Chloroembryos: A

unique photosynthesis system. J. Plant Physiol. **170**: 1131–1138.

Ranade, S.S., Delhomme, N., and García-Gil, M.R. (2019). Global gene expression analysis

in etiolated and de-etiolated seedlings in conifers. PLoS One **14**: e0219272.

Rossini, L., Cribb, L., Martin, D.J., and Langdale, J.A. (2001). The maize Golden2 gene

defines a novel class of transcriptional regulators in plants. Plant Cell **13**: 1231–1244.

Sakuraba, Y., Kim, E.Y., Han, S.H., Piao, W., An, G., Todaka, D., Yamaguchi-Shinozaki, K.,

and Paek, N.C. (2017). Rice Phytochrome-Interacting Factor-Like1 (OsPIL1) is involved

in the promotion of chlorophyll biosynthesis through feed-forward regulatory loops. J. Exp.

Bot. **68**: 4103–4114.

Sato, Y., Morita, R., Nishimura, M., Yamaguchi, H., and Kusaba, M. (2007). Mendel's green

cotyledon gene encodes a positive regulator of the chlorophyll-degrading pathway. Proc.

Natl. Acad. Sci. U. S. A. **104**: 14169–14174.

Simkin, A.J., Faralli, M., Ramamoorthy, S., and Lawson, T. (2020). Photosynthesis in

non‐foliar tissues: implications for yield. Plant J. **101**: 1001–1015.

Smolikova, G.N. and Medvedev, S.S. (2016). Photosynthesis in the seeds of

chloroembryophytes. Russ. J. Plant Physiol. **63**: 1–12.

Tamai, H., Iwabuchi, M., and Meshi, T. (2002). Arabidopsis GARP transcriptional activators

interact with the Pro-rich activation domain shared by G-box-binding bZIP factors. Plant

Cell Physiol. **43**: 99–107.

Tao, Z., Shen, L., Gu, X., Wang, Y., Yu, H., and He, Y. (2017). Embryonic epigenetic

reprogramming by a pioneer transcription factor in plants. Nature **551**: 124–128.

Thomas, H. and Ougham, H. (2014). The stay-green trait. J. Exp. Bot. **65**: 3889–3900.

Tu, X. et al. (2022). Limited conservation in cross-species comparison of GLK transcription

factor binding suggested wide-spread cistrome divergence. Nat. Commun. **13**: 7632.

Usadel, B., Poree, F., Nagel, A., Lohse, M., Czedik-Eysenberg, A., and Stitt, M. (2009). A

guide to using MapMan to visualize and compare Omics data in plants: A case study in

the crop species, Maize. Plant, Cell Environ. **32**: 1211–1229.

Wang, M. et al. (2018). Parallel selection on a dormancy gene during domestication of crops

from multiple families. Nat. Genet. **50**: 1435–1441.

Wang, P., Fouracre, J., Kelly, S., Karki, S., Gowik, U., Aubry, S., Shaw, M.K., Westhoff, P.,

Slamet-Loedin, I.H., Quick, W.P., Hibberd, J.M., and Langdale, J.A. (2013). Evolution

of GOLDEN2-LIKE gene function in C3 and C4 plants. Planta **237**: 481–495.

Wang, P., Khoshravesh, R., Karki, S., Tapia, R., Balahadia, C.P., Bandyopadhyay, A.,

Quick, W.P., Furbank, R., Sage, T.L., and Langdale, J.A. (2017). Re-creation of a Key

Step in the Evolutionary Switch from C3 to C4 Leaf Anatomy. Curr. Biol. **27**:

3278-3287.e6.

Waters, M.T., Wang, P., Korkaric, M., Capper, R.G., Saunders, N.J., and Langdale, J.A.

(2009). GLK transcription factors coordinate expression of the photosynthetic apparatus

in Arabidopsis. Plant Cell **21**: 1109–1128.

West, M.A.L., Yee, K.M., Danao, J., Zimmerman, J.L., Fischer, R.L., Goldberg, R.B., and

Harada, J.J. (1994). LEAFY COTYLEDON1 is an essential regulator of late

embryogenesis and cotyledon identity in Arabidopsis. Plant Cell **6**: 1731–1745.

Wietrzynski, W. and Engel, B.D. (2021). Chlorophyll biogenesis sees the light. Nat. Plants **7**:

380–381.

Xu, X., E, Z., Zhang, D., Yun, Q., Zhou, Y., Niu, B., and Chen, C. (2021). OsYUC11

-mediated auxin biosynthesis is essential for endosperm development of rice. Plant

Physiol. **185**: 934–950.

Yakovlev, M.S. and Zhukova, G.Y. (1980). Chhlorophyll in embryos of angiosperm seeds, a

review. Bot. Not. **133**: 323–336.

Yamamoto, A., Kagaya, Y., Toyoshima, R., Kagaya, M., Takeda, S., and Hattori, T. (2009).

Arabidopsis NF-YB subunits LEC1 and LEC1-LIKE activate transcription by interacting

with seed-specific ABRE-binding factors. Plant J. **58**: 843–856.

Yeh, S.Y. et al. (2022). Maize Golden2-like transcription factors boost rice chloroplast

development, photosynthesis, and grain yield. Plant Physiol. **188**: 442–459.

Zhang, C., Zhang, J., Tang, Y., Liu, K., Liu, Y., Tang, J., Zhang, T., and Yu, H. (2021a).

DEEP GREEN PANICLE1 suppresses GOLDEN2-LIKE activity to reduce chlorophyll

synthesis in rice glumes. Plant Physiol. **185**: 469–477.

Zhang, D., Tan, W., Yang, F., Han, Q., Deng, X., Guo, H., Liu, B., Yin, Y., and Lin, H.

(2021b). A BIN2-GLK1 Signaling Module Integrates Brassinosteroid and Light Signaling

to Repress Chloroplast Development in the Dark. Dev. Cell **56**: 310–324.

Zhang, J.-J. and Xue, H.-W. (2013). OsLEC1/OsHAP3E participates in the determination of

meristem identity in both vegetative and reproductive developments of rice. J. Integr.

Plant Biol. **55**: 232–49.

Zhang, Y., Su, J., Duan, S., Ao, Y., Dai, J., Liu, J., Wang, P., Li, Y., Liu, B., Feng, D., Wang,

J., and Wang, H. (2011). A highly efficient rice green tissue protoplast system for

transient gene expression and studying light/chloroplast-related processes. Plant

Methods **7**: 30.

Zhao, L. et al. (2020). Integrative analysis of reference epigenomes in 20 rice varieties. Nat.

Commun. **11**: 2658.

Zheng, S., Dong, J., Lu, J., Li, J., Jiang, D., Yu, H., Ye, S., Bu, W., Liu, Z., Zhou, H., Ding,

Y., and Zhuang, C. (2022). A cytosolic pentatricopeptide repeat protein is essential for

tapetal plastid development by regulating OsGLK1 transcript levels in rice. New Phytol.

234: 1678–1695.

Figure 1. OsNF-YB7 negatively regulates Chlorophyll (Chl) biogenesis in embryo.
A, B. Morphologies of wild-type (WT) and *osnf-yb7* detached embryos at 10 days after fertilization (DAF) (**A**) and longitudinally dissected embryos at maturation (**B**). Scale bars = 200 μm.
C. Transmission electron microscopy images of embryos from WT and *osnf-yb7* at 10 DAF. Scale bars = 5 μm.
D. Chl levels in WT and *osnf-yb7* embryos at 10 DAF and maturation. Data are means ± SD (n = 3). **, *p* < 0.01; Student's *t*-test was used for statistical analysis.
E. Chl autofluorescence of WT and mutant embryos at 5 DAF. BF, bright field; Chl, Chl autofluorescence. Scale bars = 100 μm.
F. Photosynthesis-related pathways enriched among the differentially expressed genes (DEGs) identified from the 5- and 10-DAF-old *osnf-yb7* embryos compared to WT, respectively.
G. Most of the photosynthesis-related genes were upregulated in the mutant embryos.
H. A heat map shows the expression of the Chl biosynthesis-related genes in the WT and *osnf-yb7* embryos at 5 and 10 DAF. Reads per kilobase per million mapped reads (RPKM) is used to indicate the expression level.

Figure 2. OsNF-YB7 directly binds to the promoters of *OsPORA* and *OsLHCB4* and regulates their expression.

A, B. Quantitative real-time PCR (RT-qPCR) analysis of the transcription levels of *OsPORA* (**A**) and *OsLHCB4* (**B**) in the embryos of WT and *osnf-yb7* at 10 DAF. Data are means $\pm$ SD (n = 3). **, $p < 0.01$; Student's *t*-test was used for statistical analysis.

C, D. Expression of *OsPORA* (**C**) and *OsLHCB4* (**D**) in the leaves of WT and *OsNF-YB7*-overexpressing transgenic plants (*NF-YB7-GFP*). Data are means $\pm$ SD (n = 3). **, $p < 0.01$; Student's *t*-test was used for statistical analysis.

E, F. Chromatin immunoprecipitation assay coupled with quantitative PCR (ChIP-qPCR) analyses showing the enrichment of OsNF-YB7 at the *OsPORA* (**E**) and *OsLHCB4* (**F**) promoters in 14-day-old *OsNF-YB7-Flag* seedlings. Precipitated DNA was quantified by qPCR and DNA enrichment is displayed as a percentage of input DNA. Data are means $\pm$ SD (n = 3). *, $p < 0.05$; **, $p < 0.01$; Student's *t*-test was used for statistical analysis. *ACTIN* was used as a nonspecific target gene. Diagrams in the upper panel showing the promoter structures of *OsPORA* and *OsLHCB4*, and the PCR amplicons used for ChIP-qPCR.

G, H. Electrophoretic mobility-shift assays (EMSAs) showing that OsNF-YB7 directly binds to the promoters of *OsPORA* (**I**) and *OsLHCB4* (**J**). Hot probes were biotin-labeled. The hot mProbes contain mutant nucleic acid from CACATG to AAAAAA. The arrow heads indicate the shift bands.

I. Schematic diagram displaying the constructs used in the dual luciferase reporter (DLR) assays of **J**. LUC, firefly luciferase; REN, *Renilla* luciferase.

J. DLR assays showing that OsNF-YB7 directly represses the promoter activities of *OsPORA* and *OsLHCB4*. Data are means $\pm$ SD (n = 3). *, $p < 0.05$; **, $p < 0.01$; Student's *t*-test was used for statistical analysis.

Figure 3. OsNF-YB7 associates with the promoter of *OsGLK1* and represses its expression.
A. A heat map showing the expression of transcription factors associated with Chl biosynthesis and chloroplast development in the 5- and 10-DAF-old embryos of WT and *osnf-yb7*. The colored dots indicate log$_2$(RPKM mean) of the genes in three biological replicates.
B. RT-qPCR analysis of *OsGLK1*, *OsGNC*, and *OsPIL14* expression levels in the embryos from WT and *osnf-yb7* at 5- and 10 DAF. Numbers represent fold changes of expression. Data are means $\pm$ SD (n = 3). *, $p < 0.05$; **, $p < 0.01$; Student's *t*-test was used for statistical analysis.
C. RT-qPCR analysis of *OsGLK1* expression levels in leaves from WT and *NF-YB7-GFP*. Data are means $\pm$ SD (n = 3). **, $p < 0.01$; Student's *t*-test was used for statistical analysis.
D. ChIP-qPCR analysis showing the enrichment of OsNF-YB7 at the *OsGLK1* promoter in 14-day-old *OsNF-YB7-Flag* seedling. Chromatin of each sample was immunoprecipitated using anti-Flag or IgG antibodies. Precipitated DNA was quantified by qPCR and DNA enrichment is displayed as a percentage of input DNA. Data are means $\pm$ SD (n = 3). *, $p < 0.05$; **, $p < 0.01$; Student's *t*-test was used for statistical analysis. *ACTIN* was used as a nonspecific target gene. The experiment was performed three times with similar results. The diagram in the upper panel showing the promoter structure of *OsGLK1* and PCR amplicons (P1, P2, P3, P4, and P5) used for ChIP-qPCR.
E. Schematic diagram displaying the constructs used in the DLR assays of **F**. LUC, firefly luciferase; REN, *Renilla* luciferase; UAS, upstream activating sequence.
F. DLR assays showing that OsNF-YB7 represses the promoter activity of *OsGLK1*. Data are means $\pm$ SD (n = 3). *, $p < 0.05$; Student's *t*-test was used for statistical analysis.

Figure 4. Chl biosynthesis in *osnf-yb7* embryo requires active *OsGLK*s.

A. Embryo morphologies of WT and Os*GLK1-OX* detached embryos at 10 DAF. Scale bars = 2 mm.

B. Chl autofluorescence of the WT and *GLK-OX* embryos at 10 DAF. BF, bright field; Chl, Chl autofluorescence. Scale bars = 100 μm.

C. Chl levels in WT and *OsGLK1-OX* embryos at 10 DAF. Data are means ± SD (n = 3). **, $p < 0.01$; Student's *t*-test was used for statistical analysis.

D-G. Morphologies of the embryos produced by WT (**D**), *osnf-yb7* (**E**), *osnf-yb7;osglk2* double mutant (**F**) and *osnf-yb7;osglk1;osglk2* triple mutant (**G**). Scale bars = 1 mm.

H. A Venn diagram showing overlaps of the DEGs identified from the embryos of *osnf-yb7* and *OsGLK1-OX* at 10 DAF.

I. A pie chart showing similar transcriptional changes of the common DEGs identified from *osnf-yb7* and *OsGLK1-OX*.

J. Gene Ontology (GO) analysis of the common DEGs identified from *osnf-yb7* and *OsGLK1-OX*.

Figure 5. OsNF-YB7 and OsGLK1 regulate a common set of genes in the embryo.
A, B. ChIP-qPCR analysis showing the enrichment of OsGLK1 on the *OsPORA* (**A**) and *OsLHCB4* (**B**) promoters. PCR amplicons used for ChIP-qPCR are indicated in the schematic diagrams. OsGLK1-GFP was transiently expressed in protoplasts isolated from 14-day-old WT seedling. Chromatin of each sample was immunoprecipitated using anti-GFP or igG antibodies. Precipitated DNA was quantified by qPCR and DNA enrichment is displayed as a percentage of input DNA. Data are means $\pm$ SD (n = 3). *, $p < 0.05$; **, $p < 0.01$; Student's *t*-test was used for statistical analysis. *ACTIN* was used as a nonspecific target gene. The experiment was performed three times with similar results.
C, D. EMSA assays showing that OsGLK1 directly binds to the promoters of *OsPORA* (**C**) and *OsLHCB4* (**D**). The arrow heads indicate the shift bands.
E. A Venn diagram showing overlaps of the target genes between OsGLK1 and OsNF-YB7.
F. Distribution of the distance between the OsNF-YB7 and OsGLK1 summits showing that OsNF-YB7 and OsGLK1 bind to proximal regions of their common targets.

Figure 6. OsNF-YB7 interacts with OsGLK1 to regulate the expression of *OsPORA* and *OsLHCB4*.

A. Schematic diagrams showing the protein structures of OsGLK1 and OsNF-YB7.

B. Yeast-two-hybrid (Y2H) assays showing the interaction between OsNF-YB7 and OsGLK1. AD and BD indicate the activation domain and binding domain of GAL4, respectively. The full length or truncated OsNF-YB7 and OsGLK1 were fused with BD and AD, respectively. The indicated combinations of constructs were cotransformed into yeast cells and grown on the nonselective medium SD/-L-W and selective medium SD/-L-W-H.

C. A split complementary luciferase (LUC) assay confirmed the interaction between OsNF-YB7 and OsGLK1. Coexpression of the fusion of OsGLK1 and the N-terminal half of LUC (nLUC-OsGLK1) and the fusion of the C-terminal half of LUC and OsNF-YB7 (cLUC-OsNF-YB7) in the epidermal cells of *Nicotiana benthamiana* leaves induced LUC activities, whereas the epidermal cells coexpressing nLUC-OsGLK1 and cLUC, nLUC and cLUC-OsNF-YB7, or nLUC and cLUC did not show LUC activities.

D. Bimolecular fluorescence complementation (BiFC) assays showed interactions between OsNF-YB7 and OsGLK1 in the nucleus. OsNF-YB7 was fused with the N-terminal of yellow fluorescent protein (nYFP-OsNF-YB7); OsGLK1 was fused with the C-terminal of yellow fluorescent protein (cYFP-OsGLK1). The recombinant proteins were transiently coexpressed in leaf epidermal cells of *N. benthamiana.* Fluorescence signals indicated that OsGLK1 interacted with OsNF-YB7 in the nucleus. Scale bar = 10 μm.

E. Co-immunoprecipitation (Co-IP) assays showing that OsNF-YB7 interacts with OsGLK1 *in vivo*. 35S::OsNF-YB7:GFP (OsNF-YB7-GFP) and 35S::OsGLK1:3×Flag (OsGLK1-Flag) were coexpressed in rice protoplasts and were immunoprecipitated with an anti-GFP antibody, and the immunoblots were probed with anti-GFP and anti-Flag antibodies. 35S::GFP (GFP) was used as a negative control.

F, G. DLR assays showing that OsNF-YB7 represses the transactivation activity of OsGLK1 on *OsPORA* (**F**) and *OsLHCB4* (**G**) using etiolated protoplasts. EV, empty vector. Data are means $\pm$ SD (n = 3). *, $p < 0.05$; **, $p < 0.01$; Student's *t*-test was used for statistical analysis.

H. EMSA assay indicated that OsNF-YB7 inhibits the DNA binding of OsGLK1 to the promoter of *OsPORA*. The black and white arrow heads indicate the OsGLK1- and OsNF-YB7-bound probes, respectively; "+" and "++" indicate that 2- and 4 μM recombinant proteins were used for the reactions. The GST-His was used as a negative control.