Fine-scale tracking reveals visual field use for predator detection and escape in collective foraging of pigeon flocks

Mathilde Delacoux* (1, 2, 3)

Fumihiro Kano* (1, 2)

*correspondence

mdelacoux@ab.mpg.de, mathilde.delacoux@uni-konstanz.de

fkano@ab.mpg.de, fumihiro.kano@uni-konstanz.de

1. Centre for the Advanced Study of Collective Behaviour, University of Konstanz, Germany

2. Max Planck Institute of Animal Behavior, Radolfzell, Germany

3. International Max Planck Research School for Quantitative Behavior, Ecology and Evolution, Radolfzell, Germany

Abstract

During collective vigilance, it is commonly assumed that individual animals compromise their feeding time to be vigilant against predators, benefiting the entire group. One notable issue with this assumption concerns the unclear nature of predator "detection", particularly in terms of vision. It remains uncertain how a vigilant individual utilizes its high-acuity vision (such as the fovea) to detect a predator cue and subsequently guide individual and collective escape responses. Using fine-scale motion capture technologies, we tracked the head and body orientations of pigeons (hence reconstructed their visual fields and foveal projections) foraging in a flock during simulated predator attacks. Pigeons used their fovea to inspect predator cues. Earlier foveation on a predator cue was linked to preceding behaviors related to vigilance and feeding, such as head-up or down positions, head-scanning, and food-pecking. Moreover, earlier foveation predicted earlier evasion flights at both the individual and collective levels. However, we also found that relatively long delay between their foveation and escape responses in individuals obscured the relationship between these two responses. While our results largely support the existing assumptions about vigilance, they also underscore the importance of considering vision and addressing the disparity between detection and escape responses in future research.

Keywords

Fovea, predator detection, pigeon *(Columba livia)*, collective vigilance, social contagion

Introduction

In everyday natural tasks, such as locating food, avoiding predators, and interacting with conspecifics, animals constantly face challenges in deciding when and how to adjust their behaviors to increase their chances of survival. One such behavior is the focusing of attention, or "looking" behavior, which has been relatively understudied in natural tasks due to the technical challenges involved in tracking an animal's gaze during natural activities. However, this is relatively well studied within the context of vigilance, a scenario where an animal's survival hinges on effective attentional allocation and where researchers can observe their scanning behavior even in field conditions. Vigilance, especially during foraging, generally involves the compromise between staying alert for threats and searching for food, as well as maintaining a balance between individual and collective vigilance (Beauchamp, 2015).

Common assumptions of vigilance research

Common assumptions about collective vigilance, first coined by Pulliam (Pulliam, 1973; Pulliam et al., 1982), posit that a vigilant individual, often identified by a "head-up" posture, is likely to have a higher probability of detecting an approaching predator compared to a feeding individual, typically in a "head-down" posture. Consequently, the vigilant individual can react and escape more swiftly, thereby reducing its predation risk (also see Beauchamp, 2015; Godin & Smith, 1988). In a group setting, non-vigilant members can gain benefits from the vigilant individual by following its lead when it exhibits escape behavior (Pulliam, 1973). This potential group advantage, known as "collective detection," might explain that individuals in a larger group tend to be less vigilant (and hence spend more time in foraging), suggesting a critical advantage of grouping (Pulliam, 1973; Pulliam et al., 1982).

Empirical challenges

While these assumptions underpin many influential theoretical models in behavioral ecology, they have not gone unchallenged, with numerous empirical studies highlighting potential inconsistencies. Firstly, the trade-off between vigilance and foraging may not be as pronounced as previously assumed. Even though several studies found a relationship between vigilance and escape latency

(Devereux et al., 2006; Hilton et al., 1999; Lima & Bednekoff, 1999), various studies have demonstrated that these two behaviors are not necessarily mutually exclusive (Bednekoff & Lima, 2005; Cresswell et al., 2003; Devereux et al., 2006; Kaby & Lind, 2003; Lima & Bednekoff, 1999). In several species, especially those with a broad visual field, individuals can simultaneously engage in foraging activities while remaining vigilant (Fernández-Juricic, 2012), likely using peripheral vision to detect approaching threats (Bednekoff & Lima, 2005; Cresswell et al., 2003; Kaby & Lind, 2003; Lima & Bednekoff, 1999). Relatedly, although vigilance and foraging have been defined in many previous studies respectively as head-up and head-down due to the limitation associated with direct observation, several studies have pointed out that vigilance triggered by the appearance of a predator is not necessarily related to the head-up postures but to the pattern of head movements (Fernández-Juricic, 2012; Jones et al., 2007, 2009). Similarly, the patterns of head-up (number of head-up bouts, their length and regularity), rather than the proportion of time being head-up versus head-down, are sometimes better predictors of predator detection (Beauchamp, 2015; Beauchamp & Ruxton, 2016; Bednekoff & Lima, 2002; Cresswell et al., 2003; Hart & Lendrem, 1984).

Second, it is challenging to empirically differentiate the effect of collective detection from other group size-related effects, such as confusion, dilution, and edge effect. Although there is evidence that non-detectors can benefit from detectors' escape (Davis, 1975; Lima, 1995b; Lima & Zollner, 1996), other studies favor alternative hypotheses based on their findings (Beauchamp & Ruxton, 2008; Inglis & Lazarus, 1981). It appears that multiple factors influence vigilance across species, including group size, density, and spatial configuration, as well as social contagion (Elgar, 1989; Roberts, 1996).

Third, non-vigilant animals do not always respond to behavioral cues of other members of the group. The upright-freezing alert posture, often one of the first behavioral indications of predator detection, tends to be rather inconspicuous, and flockmates usually do not seem to respond to it (Fernández-Juricic et al., 2009; Lima, 1995b). Additionally, birds do not necessarily differentiate between predator-based escapes and non-threat departures from a flock mate, suggesting that simultaneous departures of multiple birds are required to trigger contagious flights (Cresswell et al.,

2000; Lima, 1995b; Proctor et al., 2001). As a result, the extent to which group-living animals benefit

from collective detection remains open to questions (Bednekoff & Lima, 1998).

Finally, the definition of "detection" is generally uncertain in studies. Previous studies

typically used overt escape responses, such as flying, as a measure of detection (using escape as only

measure: Kenward, 1978; Lima & Zollner, 1996; Quinn & Cresswell, 2005; or partially using escape

as response: Cresswell et al., 2003; Lima, 1995a, 1995b; Tisdale & Fernández-Juricic, 2009;

Whittingham et al., 2004). However, a bird likely identifies a potential threat before escaping

(Barbosa & Castellanos, 2005; Fernández-Juricic, 2012; Lima & Dill, 1990). Thus, subtler behavioral

responses like freezing or alertness were also used (e.g., Fernández-Juricic et al., 2009; Kaby & Lind,

2003; Lima & Bednekoff, 1999; Rogers et al., 2004). These studies have found a period between these

subtle behavioral shifts and overt escape actions (response time delay), which is considered a crucial

risk-assessment phase (Cresswell et al., 2009). This phase has been found to depend on multiple

factors such as species, context (e.g., food availability, surrounding environment: Fernández-Juricic et

al., 2002), individual characteristics (Jones et al., 2009) and flock size (Boland, 2003). An issue

remains, however, that even these subtle behaviors might not accurately capture the time when an

animal detects an approaching threat (Barbosa & Castellanos, 2005; Fernández-Juricic, 2012; Lima &

Dill, 1990; Tätte et al., 2019), due to the lack of a more direct measure of visual attention.

Visual sensory ecology of birds

More recent research in visual sensory ecology has emphasized the role of vision in the context of

vigilance. Vision is crucial for gathering distant predator cues, and it is thus expected to play a

significant role in predator-prey interactions (Barbosa & Castellanos, 2005). Specifically, predation is

considered one of the primary drivers of the evolution of the visual system in birds, as well as

foraging (Martin, 2017). Birds' retinal specializations, such as the *area* (a region of the bird's retina

with a high density of photoreceptors) or the fovea (a pit-like area in the retina where visual acuity is

highest), are thought to be crucial for the detection, identification and tracking of predators

(Fernández-Juricic, 2012; Martin, 2017) as well as risk assessment and response selection (Cresswell,

1993; Cresswell et al., 2009; Fuchs et al., 2019; Veen et al., 2000). Experimental studies tracking the

foveal projections demonstrated that birds indeed use their foveas to inspect predator cues (Butler &
Fernández-Juricic, 2018; Tyrrell et al., 2014; Yorzinski & Platt, 2014). Given the diversity and
complexity of birds' visual field configurations, it is necessary to understand how different types of
visual information are acquired based on body posture and head orientation, rather than simply to
consider a head-up and head-down dichotomy (Fernández-Juricic et al., 2004). Notably, while it is a
prevalent notion that vigilant individuals are quicker to detect predators, this fundamental assumption
has seldom been directly scrutinized in the literature (Beauchamp, 2015).

Methodological issues

Despite the inherent technical difficulties in measuring visual perception during vigilance-foraging
tasks, a few studies have pioneered using innovative methodologies. Yorzinski & Platt (2014), for
instance, employed wearable eye-tracking technology on peacocks, showing that these birds actively
use their foveas for the detailed inspection of predators. Tyrrell et al. (2014) employed an eye tracker
with restrained smaller birds (starlings), showing that they also rely on their foveas for high-resolution
viewing. These studies highlight the critical role of foveal vision in predator detection. However,
while these studies were noteworthy in accurately capturing the birds' foveations, they did not capture
the collective vigilance of multiple birds within a flock during natural foraging.

Experimental rationales

To fill the gap in knowledge, this study focused on the "meso-scale" level of analysis concerning the
dynamics of collective antipredator behaviors. This contrasts with "macro-scale" analyses typically
conducted in the field, where observation and control are often challenging, or the "micro-scale"
analysis conducted under controlled but often artificial laboratory settings. To achieve this, we utilized
a state-of-the-art, non-invasive motion capture technique to monitor the head orientations of pigeons
moving freely within a flock (Nagy et al., 2023; Fig 1a and b). This approach enabled us to closely
analyze the birds' fine-scale looking behaviors while maintaining their natural foraging, vigilance, and
social interactions. By implementing this approach, our specific goal was to test a series of prevalent
assumptions within vigilance research, assumptions that have been challenged by multiple empirical
studies. In particular, we first examined the relationship between predator detection and foveation,

specifically testing whether birds employ their foveas in response to potential predatory threats.

Subsequently, we tested whether foveating on predator cues is associated with vigilance and foraging

behaviors and influences the likelihood and timing of escape. Finally, we assessed whether an

individual bird's visual and escape responses to predator cues affects the escape responses of the entire

flock.

Our methodology allowed us to reconstruct the visual fields of all birds in a flock during their

natural behavior. Although we did not record eye movements, a prior study (Kano et al., 2022)

established that pigeons utilize restricted regions of their visual fields, corresponding to the regions

where the foveas would project. This is consistent with the observation that eye movement in this

species is relatively limited, typically confined within a 5-degrees range (Wohlschläger et al., 1993).

In addition to head-tracking, we developed simple threshold-based algorithms to automatically

classify the behavior of individuals (e.g., head-up, grooming, feeding, running, flying) based on their

reconstructed postures. We examined the behavior of flocks of 10 pigeons simultaneously before,

during and after simulated predation events (Fig 1c). These events consisted of a 10-sec looming

stimulus (a predator shade) displayed on a monitor followed by a model predator appearing from

under a table covered with fabric below the monitor, and running on a wire through the room. We

used a looming predator silhouette as a warning cue simulating a predator approach before presenting

the model predator; the looming stimulus was chosen because it is generally perceived as threatening

across species (Evans et al., 2019), and is subtle enough so that pigeons require effort to detect it in

our experimental setup.

A prior study (Kano et al., 2022) established that when focusing on a distant visual target

(beyond their pecking distance), pigeons primarily use their foveas, rather than their frontal visual

field (including the area of binocular overlap and the projection of the visually sensitive "red fields"

(Wortel et al., 1984). Consequently, we focused on analyzing their foveas, not their frontal visual

field. In our observations, pigeons foveated on the predator cue typically before the offset of looming

stimulus (Fig 1d); we considered this foveation as a potential indicator of early detection (in a few

instances, pigeons focused on the model predator after the looming stimulus had disappeared, but

these cases were excluded from our analysis). To eliminate the possibility of random gaze crossing

over the predator cue, the first foveation was defined as the first instance lasting longer than 300 ms

and occurring at least 200 ms after the stimulus onset; this decision was based on the typical

intersaccadic intervals and reaction times observed in this species (Blough, 1977; Kano et al., 2022).

Pigeons initiated running and then took flight typically during a predator presentation event (Fig 1d,

see also Table S1). Pigeons responded either directly to the looming stimulus or after the appearance

of the model predator, by running away and/or flying. Therefore, our first measure of escape was the

presence of an escape response (either running or flying) prior to the looming stimulus offset (i.e.,

whether the pigeon runs or flies before the looming stimulus disappears). Our second measure of

evasion focused on flight responses (predominantly occurring after the predator appeared), given that

flight is the most prominent form of escape response in birds and is widely noted in the literature for

its contagious nature among flockmates (Davis, 1975; Lima, 1995b; Lima & Zollner, 1996).

Figure 1: a) Experimental setup. In a large-scale motion capture system (SMART-BARN), the feeding area, the monitors and the predator running wire were installed (a 3D view on the left, and a top view on the right). b) Still frames captured from the periods before (top) and during (bottom) the predator presentation, respectively. c) Sequence of an event presenting a model predator. Each predator presentation event started with the presentation of a looming stimulus (predator silhouette), which lasted for approximately 10 s, on either of the two monitors, followed by the model predator running on the wire across the room (lasting for approximately 3 s). The "before-cue" period is defined as the

191 2 minutes directly preceding the onset of the looming stimulus. d) Histograms of the pigeons'

192 responses. Top: latency to foveate (green) and latency to escape –either running or flying (purple);

193 bottom: latency to run (blue) and latency to fly (red) separately. The vertical line shows the looming

194 stimulus offset.

Experimental hypotheses

We subsequently developed a series of hypotheses outlined in Table 1. The initial two hypotheses

(Hypotheses 1 and 2) aim to examine whether the use of retinal specializations, such as foveas (also

known as the *area centralis*), correlates with predator detection. Hypothesis 1 posits that pigeons

employ their foveas to evaluate the predator (looming) cue. Hypothesis 2.1 suggests that the birds'

behavioral state observed at the onset of the looming stimulus (e.g., head-up, foraging) predicts the

latency to foveate on the cue. Additionally, we assessed spatial positioning—relative to the threat

(distance from the monitor) or to conspecifics (nearest neighbor)—as these factors are known to

influence vigilance and escape behaviors (Beauchamp, 2008; Beauchamp & Ruxton, 2008; Inglis &

Lazarus, 1981). A related hypothesis, Hypothesis 2.2, posits that various aspects of vigilance/foraging

behaviors observed prior to the presentation of the looming stimulus (e.g., head-up, pecking rate,

head-saccade rate, time spent foveating on predator-related objects or conspecifics) correlate with the

latency to foveate on the cue. As noted, these related yet distinct facets of visual/foraging behaviors

have been identified as differentially related to vigilance and escape (Cresswell et al., 2003; Jones et

al., 2007, 2009). From Hypothesis 3, we asked whether earlier foveation on the predator cue predicts

quicker escape responses by individual birds. While this is a common assumption in the literature,

several factors appear to affect the length of the response time delay (Cresswell et al., 2009; Jones et

al., 2009; Tätte et al., 2019). The final hypothesis (Hypotheses 4) focus on the collective dynamics of

detection and escape. Specifically, Hypothesis 4.1 posits that the first pigeon in a flock to foveate on

the looming stimulus facilitates the escape behaviors of other members. A related hypothesis,

Hypothesis 4.2, posits that the timing of escape initiations is socially contagious, specifically that the

escape responses within a predator presentation event are more clustered than would be expected by

chance.

Table 1. A set of hypotheses testing the assumptions about vigilance

	Hypothesis	Test	Response	Description of the test
1	Pigeons foveate on the predator cue (monitor or predator hiding table) during the 10-sec presentation period of the looming stimulus.	Linear Mixed Model: LMM	Proportion of time foveating on the objects of interest	Testing the effect of: Object type (the monitor presenting the looming stimulus, the monitor that did not present the looming stimulus, and the nearest neighbor pigeon)
2.1	Pigeons that are head-up (while not grooming nor courting or courted) foveate on the predator cue earlier than those that are feeding at the onset of the looming stimulus.	LMM	Latency to foveate on the predator cue.	Testing the effects of: Behavior exhibited at the onset of the looming stimulus (Head-up/Feeding), Distance from the monitor presenting the looming stimulus, and Distance from the nearest neighbor conspecific.
2.2	Pigeons' vigilance/feeding-related behaviors during the 2-min "before-cue" period predict their earlier foveation on the predator cue.	LMM	Latency to foveate on the predator cue.	Testing the effects of: Vigilance/feeding-related behaviors (proportion of time spent head-up, saccade rate, pecking rate, proportion of time foveating on any monitor, proportion of time foveating on any conspecifics).
3	Pigeons' first foveation on the predator cue predicts their escape.	LMM and Generalized Linear Mixed Model: GLMM	Probability of evasion before the offset of the looming stimulus, latency to fly.	Testing the effect of: Latency to foveate on the predator cue.
4.1	The first pigeon foveating on the predator (looming) within a flock facilitates the escape behaviors of the other members.	LMM and GLMM	Probability of escape before the offset of the looming stimulus and latency to fly in the other members (excluding the first pigeon).	Testing the effect of: Latency to foveate on the predator cue by the first pigeon.
4.2	Escape responses within a predator presentation event are more clustered than expected by chance.	Permutation test	Time interval between the successive escape (or flying) responses of pigeons within a given predator presentation event.	Comparing the mean interval observed in the data with the mean intervals expected by chance in a control event created by shuffling the latencies between event ID.

Results

Overview of observations

In our study, we tested 20 pigeons across 6 trials each, with every trial involving a flock of 10 pigeons and consisting of 2 predator presentation events. This approach resulted in a total of 240 observations, with each observation representing data from a single event for an individual pigeon. Each model incorporated this number of observations while excluding null cases in which pigeons displayed no response or cases where the system lost track of the pigeons (see Method for the details).

Within a few predator presentation events, our pigeons rapidly decreased their latencies to foveate, escape, and fly, indicating that learning occurred rapidly during these trials (Fig S1). At the very first predator presentation event, approximately half of the flock members did not foveate on the monitor nor escaped even after the appearance of the model predator. From the second predator presentation event, most pigeons made foveation, escape, and flight responses. Our pigeons maintained these responses throughout the trials, although in the last few trials pigeons decreased their flight responses, indicating that habituation was minimal. To ensure that these behavioral changes did not confound the effect of our test predictors, we controlled for the trial and event IDs, as well as other basic parameters (predator side, sex, and pigeon ID) in all models.

Test 1 (Hypothesis 1)

From the markers coordinates, we were able to reconstruct for each pigeon their head orientations simultaneously (Fig 2a-b). In the heatmaps (Fig 2c), we projected the 3D representations of the three objects of interest—the monitor displaying the looming stimulus, the monitor without the cue serving as a control, and the nearest conspecific (see supplementary for the 3D object definitions)—onto the pigeons' local head coordinate system. We sampled this data during the period of the looming stimulus presentation and before any escape response by the pigeon (normalized to unit sum). Visual inspection suggested that pigeons foveated on the monitor displaying the looming stimulus, more so than on the other monitor that was not presenting the looming stimulus or on the nearest conspecific. It should be

noted that the two monitors were placed on opposite sides of the room. Consequently, a pigeon

located in the center and foveating on one monitor would have the other monitor falling ~105° in

azimuth angle on the opposite side of its head. This is likely resulting in the spot next to the foveal

region for the monitor not displaying the looming stimulus (see Fig 2c).

Subsequently, we quantified the frequency (normalized to a unit sum) at which each object of

interest was observed within the defined foveal regions on the heatmap (illustrated by red rectangles;

corresponds to the fovea location, adjusted by ± 10 degrees in both elevation and azimuth). This

quantification was conducted for each pigeon at every predator presentation event. After logit-

transforming this response to improve the normality of residuals, we conducted a Linear Mixed Model

(LMM) with the object of interest as a within-subject categorical test variable, in addition to control

variables and random effects (see Methods and Table S7). We observed a significant effect of the

objects of interest ($\chi^2(1) = 108.06$, $p < 0.001$) (Fig 2d). Subsequent follow-up analyses revealed that

pigeons focused on the monitor displaying the looming stimulus for a longer period compared to the

other two objects (vs. monitor not presenting the looming stimulus, $\chi^2(1) = 62.44$, $p < 0.001$; vs.

nearest neighbor, $\chi^2(1) = 97.41$, $p < 0.001$).

Figure 2: Pigeons' foveation during the looming stimulus presentation. a) A pigeon equipped with motion capture markers on its head and back. b) Reconstruction of estimated foveal projections in the local coordinate system of the head (left) and in the global (motion-capture) coordinate system (right). c) Reprojection of each object of interest—the monitor presenting the looming stimulus, the monitor without the cue acting as a control, and the nearest conspecific—within the visual field of all pigeons across all trials. The red squares denote the designated region of estimated foveal projections in the visual field, defined as $75\pm10°$ in azimuth and $0\pm10°$ in elevation. The color scale illustrates the normalized frequency (normalized to a unit sum) at which each object of interest was observed within each $5\times5°$ cell of the heatmap. d) Normalized frequency (normalized to unit sum then logit-transformed) at which each object of interest was observed within the defined foveal regions on the heatmap. The dots represent single observations (jittered horizontally for visualization), the grey shade represents the violin plot of the distributions, and the boxplot's boxes represent the 0.25 and

277 0.75 quartiles (with the median represented as a line inside the box) and the whiskers the minimum

278 and maximum values within the lower/upper quartile ± 1.5 times the interquartile range.

Test 2 (Hypothesis 2)

To test Hypothesis 2.1, we conducted a LMM using the latency to foveate (log-transformed to

improve the normality of residuals) as the response variable. The pigeons' behavioral state (either

head-up or feeding) was included as a categorical test variable, while the distance from the monitor

and the distance from the nearest neighbor were included as continuous test variables (in addition to

the control variables and random effects; Table S7). The analysis revealed that the latency to foveate

was significantly influenced by the behavioral state ($\chi^2(1) = 13.78$, $p < 0.001$). However, it was not

significantly affected by the other spatial factors, such as the distance from the monitor ($\chi^2(1) = 1.20$,

$p = 0.27$) or the distance from the nearest neighbor ($\chi^2(1) = 0.05$, $p = 0.82$) (Fig 3a).

Figure 3: Results of the models from Test 2 and 3. a) Latency to foveate on the predator cue (log-

transformed) as a function of behavioral state (feeding or head-up) at the onset of the looming

stimulus. The dots represent single observations (jittered horizontally for visualization), the grey shade represents the violin plot of the distributions, and the boxplot's boxes represent the 0.25 and 0.75 quartiles (with the median represented as a line inside the box) and the whiskers the minimum and maximum values within the lower/upper quartile ± 1.5 times the interquartile range. b-c) Latency to foveate on the predator cue as a function of the proportion of time spent being head-up during the "before-cue" period (b) and as a function of the pecking rate during the same period (c). d-e) Probability of escaping before the looming stimulus offset (d), and latency to fly (e) as a function of the latency to foveate. (f) Histogram of the distribution of the response delay (time between the latency to foveate and the latency to fly). For all depicted results, regression lines were determined with other variables held constant, set to their mean values. A comprehensive table detailing the outcomes of the models can be found in Table S7.

To test Hypothesis 2.2, we ran 5 LMMs, each of which include the latency to foveate as a response variable and any of the 5 related behaviors (the proportion of time spent head-up, the saccade rate, the pecking rate, the proportion of time foveating on the monitor and the proportion of time foveating on any conspecifics) as a continuous test variable (in addition to the control variables and the random effects as defined in the Method). The latency to foveate was significantly predicted by most of the test variables, including the proportion of time spent being head-up ($\chi^2(1) = 15.90$, p < 0.0001), the pecking rate ($\chi^2(1) = 14.07$, p = 0.0002), the proportion of time foveating on a monitor ($\chi^2(1) = 12.04$, p = 0.0005), and the saccade rate ($\chi^2(1) = 9.85$, p = 0.0017). However, the proportion of time foveating on any conspecifics did not have a significant effect ($\chi^2(1) = 0.06$, p = 0.8116) (Fig 3b-c).

When comparing the Akaike Information Criterion (AIC) of all five models, the most effective model included the proportion of time spent being head-up (AIC = 609.99). This was followed by models including the pecking rate (AIC = 612.40), the proportion of time foveating on the monitor (AIC = 614.01), and the saccade rate (AIC = 616.00).

In further analyses outlined in the Supplementary Material, we examined the changes in these

vigilance- and feeding-related behaviors shortly after the predator disappeared (1 minute after the

predator's disappearance) as compared to the 1 min period preceding the stimulus onset. The results

highlighted a significant increase in vigilance and a significant decrease in feeding immediately

following the predator's disappearance across all variables (see Fig S5 and Table S5 for details).

Test 3 (Hypothesis 3)

We conducted a binomial Generalized Linear Mixed Model (GLMM) analysis using the probability of

escape before the looming stimulus offset as a binary response variable and the latency to foveate as a

continuous test variable (in addition to control variables and random effects). This model indicated

that the latency to foveate had no significant effect ($\chi^2(1) = 0.02$, p = 0.8716). Subsequently, we

analyzed a Linear Mixed Model (LMM) with the latency to fly as a continuous response variable,

with the same test and control variables. This model revealed a significant effect of the latency to

foveate, suggesting that earlier foveation predicted quicker flight responses ($\chi^2(1) = 6.49$, p = 0.0108)

(Fig 3d-e). We observed that the response time delay (the interval between the latency to foveate and

the latency to fly) was relatively lengthy and exhibited considerable variation (Fig 3f). During this

period, pigeons rarely returned to the feeding activity, as indicated by the low mean pecking rate (0.10

pecks/sec). In the Supplemental Material, we demonstrate that this variation can be partially attributed

to individual differences; a within-individual repeatability analysis revealed that one contributing

factor to this variation is inter-individual differences (R = 0.128, p = 0.0404).

Test 4 (Hypothesis 4)

To test Hypothesis 4.1, we included into GLMM the probability of escaping before the looming

stimulus offset as a binary response and the first pigeon's latency to foveate as a test variable (in

addition to control variables and random effects). We found that earlier foveation of the first pigeon

was related to earlier escape responses among the other flock members, although the effect was not

significant ($\chi^2(1)$ = 3.66, p = 0.0559). We then examined the same model with latency to fly as a

continuous response and found that the earlier foveation of the first pigeon significantly predicted

earlier flying responses among the other flock members ($\chi^2(1)$ = 17.32, p = 0.0003) (Fig 4a-b).

To test Hypothesis 4.2, we conducted a permutation test based on time gaps between the

latencies of individual pigeons to either escape or fly within a given event. This time gap was defined

as the latency of the focal individual minus the latency of the immediately preceding individual. For

the permuted data, we sampled one event from two different flocks of pigeons. For each sample, the

model predator was presented to both flocks from the same side (North or South) during the same

trial. We matched the trial and predator presentation side because, among the three control factors

(trial, predator presentation side, and event), these two significantly influenced individual latencies as

determined by a LMM (the event ID was not a significant factor in this confirmatory test). We

combined the latency data from both events, randomly redistributed this data between the two events

10,000 times, and then recalculated the time gaps each time (Fig 4c). The mean time gaps from the

observed data were then compared to those from the 10,000 permutations to determine a p-value. The

observed time gaps were significantly shorter than the permuted gaps for both escape (p = 0.00038)

and fly responses (p < 0.0001; Fig 4d).

We also considered the possibility that the individuals' distance from the monitor could

confound the observed effect. Specifically, if the two pigeon flocks occupied different locations across

the two events, and the distance from the monitor influenced the individuals' latency to escape or fly,

the results might reflect spatial rather than social influences. Upon inspecting the data, this effect

seemed plausible. To account for this potential confound, we determined the overlapping areas by

calculating the range between the minimum and maximum distances of each flock and subsequently

analyzed only the individuals within this overlapping area. Using these controlled datasets, we found

that the mean time gap was not significantly different for the escape latency (p = 0.206), but it was

significantly shorter for the fly latency (p = 0.0237) (Fig 4d).

Figure 4: Results of test 4. a-b) Latency to foveate of the first individual foveating on the monitor as a

function of the other individuals' probability of escape before the cue offset (a) and the other

individuals' latency to fly (b). All the regression lines were calculated with other variables constant

and equal to their mean value. A detailed table of the models' results is available in Table S7. c) An

overview of the permutation test procedure. d) Results of the permutation test. Histograms denote the

distributions of the average time gap in permutated data. The vertical dashed line indicates the mean

time gap from the observed data. The proportion of permutated data less than the observed data (on

the left of the dashed line) gives the p-value; escaping (top) and flying (bottom); without (left) or with

(right) distance control.

Discussion

This study leveraged fine-scale behavior tracking to examine key assumptions of vigilance research in pigeon flocks during free foraging, particularly focusing on the role of foveation. Using motion-capture technology, we tracked the individuals' visual fields and quantified their behaviors automatically during predator presentation events. Our findings supported the assumptions about vigilance in two aspects: foveation in pigeons is associated with predator inspection, and behaviors related to vigilance and feeding are reliable predictors of earlier foveation in pigeons. Our findings also largely supported the assumptions about collective vigilance in two aspects: earlier foveation is indicative of earlier flight responses at both individual and collective levels, and there is a social contagion effect in their evasion flight responses. However, these results are somewhat complicated by the individuals exhibiting long and variable response time delays—the interval between foveation and escape responses. Moreover, social contagion of evasion was only observed in flight responses following the appearance of the model predator (after most flock members had foveated on the predator cue), not in earlier responses. In summary, while our results largely affirm the prior assumptions about vigilance, we have identified several confounding factors, which will be discussed further below.

Use of foveas during predator cue inspection (related to Hypothesis 1).

Our pigeons primarily use their foveas to inspect the looming stimulus. This is consistent with previous studies showing that starlings and peacocks use their foveas for predator inspection (Butler & Fernández-Juricic, 2018; Tyrrell et al., 2014; Yorzinski & Platt, 2014). Our research extends these findings to situations where pigeons forage freely on the ground in a flock. It has been previously shown that pigeons use their frontal visual fields for searching and pecking at grain, and their foveas to inspect distantly presented objects (Kano et al., 2022). In line with this, our study found that pigeons did not employ their frontal visual fields to inspect the looming stimulus. Additionally, pigeons did not use their foveas to view conspecifics during the stimulus presentation. Collectively, our results indicate that, during the stimulus presentation, pigeons primarily relied on their foveas to

attend to the predator cue. This consolidates the idea that the avian fovea is crucial in predator

detection and inspection (Martin, 2017).

Testing the assumptions about vigilance (related to Hypothesis 2)

We found that vigilance-related behaviors, including keeping the head-up, engaging in more frequent

head movements (saccades), and earlier monitoring of potential threat locations (i.e., monitors),

indeed lead to earlier foveation on the predator cue. Conversely, feeding-related behaviors, such as

keeping the head-down and a higher pecking rate, delay the foveation on the cue. The spatial

configuration of the flock, such as the distance from the predator cue and foveation on conspecifics,

did not influence earlier foveation on the predator cue. Further analyses showed that these vigilance-

related behaviors significantly increased, and these feeding-related behaviors decreased after the

presentation of the predator (see Fig S5).

These findings indicate that all the measured vigilance- and feeding-related behaviors are

relevant to predator detection in our study system. This is in line with several studies (Devereux et al.,

2006; Hilton et al., 1999; Lima & Bednekoff, 1999), though not others (Cresswell et al., 2003; Jones

et al., 2007, 2009; Kaby & Lind, 2003). While prior research has suggested that species with a

relatively large visual field can maintain vigilance peripherally (Fernández-Juricic et al., 2004), our

data propose that behaviors such as head-up and scanning are still advantageous, and head-down

while feeding is still costly to predator detection, even in pigeons that possess a relatively large visual

field. Furthermore, previous studies have noted that, in blue tits, vigilance- and feeding-related

behaviors are linked to detection only when the feeding task is more demanding (Kaby & Lind, 2003).

This might be partly relevant to our results, as we used a moderately attention-demanding task where

pigeons searched for grain in a grain-grit mixture.

Most notably, our fine-scale tracking of pigeon behaviors may have refined our ability to

identify predator detection. Previous studies have inferred predator detection by using escape

responses as the sole measure (Kenward, 1978; Lima & Zollner, 1996; Quinn & Cresswell, 2005) or

by observing more subtle signs of alertness such as freezing, adopting a straight upright posture,

crouching, or ceasing to feed (e.g., Fernández-Juricic et al., 2009; Kaby & Lind, 2003; Lima &

Bednekoff, 1999; Rogers et al., 2004), or by combining these responses (Cresswell et al., 2003; Lima,

1995a, 1995b; Tisdale & Fernández-Juricic, 2009; Whittingham et al., 2004). However, as previously

claimed (Barbosa & Castellanos, 2005; Fernández-Juricic, 2012; Lima & Dill, 1990; Tätte et al.,

2019), predator detection might occur earlier without overt behavioral responses, which can be too

subtle to detect under standard observational conditions. In our study, we measured the birds' first

foveation on the predator cue while excluding instances of short and early foveation on the cue (see

Method). Although making definitive claims is technically challenging without assessing the internal

perceptual process, the clear relationships observed between foveation and vigilance- and feeding-

related behaviors in our results suggest that foveation is likely a reliable proxy for "detection" within

our study system.

Detection and escape (related to Hypothesis 3)

In our study, earlier foveation on the predator cue predicted earlier evasion responses from the model

predator. However, this relationship was confirmed only when we analyzed the flight responses made

after the predator's appearance, not when we included pigeons' running as the escape responses.

Moreover, pigeons exhibited relatively long and variable response times between foveation and the

flight response. This response time delay was, in part, related to consistent individual differences

across trials.

One interpretation of this result is that pigeons, after foveating on the predator cue, assessed

its potential risk. The influence of earlier detection on the decision to fly earlier likely stems from

increased sensitivity to risk following the appearance of the model predator. The consistent individual

differences in response time delay suggest variations in risk sensitivities among individuals. The

relatively long and variable response time delay observed is likely inherent to our study design, which

involved using a looming stimulus as a warning for the appearance of a model predator. Notably,

pigeons rapidly decreased their latency to foveate after just a few presentations (as shown in Fig S1),

indicating increased sensitivity rather than habituation to the stimuli across repeated presentations.

The low pecking rate during the response time delay (as detailed in the Results section) suggest that

pigeons were alert when viewing the looming stimulus. However, this alertness more likely led them

to assess the stimuli rather than to initiate an explicit escape response in most cases. These findings

are in line with the previously established idea that detection and escape represent two distinct stages

in antipredator responses (Barbosa & Castellanos, 2005; Fernández-Juricic, 2012; Lima & Dill, 1990),

and that response time delay is associated with risk assessment (Cresswell et al., 2009) and individual

differences (Jones & Godin, 2010).

Social contagion of escape responses (related to Hypothesis 4)

When the first pigeon in the flock foveated on the predator cue earlier, the remaining flock members

flew earlier. Permutation tests indicated that their evasion flight responses are socially contagious (no

clear evidence for earlier escape response involving running).

One interpretation of these results is that the decision to fly by the first pigeon that foveated

triggered a following response from the rest of the flock. This social contagion in flight responses has

been previously demonstrated in birds, including pigeons (Davis, 1975). Notably, our pigeons did not

directly foveate on other flock members (as illustrated in Fig 2), suggesting that direct foveation on

other pigeons is not necessary for the social contagion of flights. This finding is consistent with the

notion that birds only need to maintain peripheral visual contact with their flock members to observe

and respond to their departure flights (Lima & Zollner, 1996). Additionally, considering the findings

related to Hypotheses 3 and 4, our results indicate a nuanced form of collective detection: while

earlier detectors may trigger earlier flight responses in the flock via social contagion, non-detectors do

not necessarily gain an advantage from these detectors, as most flock members likely already detected

the predator cue.

Limitation

479 A potential limitation of our study is its limited generalizability. We focused on pigeons, which are

480 middle-sized birds, and their escape responses may differ from those of smaller birds commonly

481 studied in vigilance research; pigeons might, for instance, be more hesitant to take flight due to the

482 higher energy costs related to their weight. Furthermore, while our system precisely tracked freely

foraging birds, some elements of our experimental setup were artificial. The looming stimulus and the model predator were presented in a somewhat disjointed manner. Pigeons had to rapidly learn the association between these two stimuli over several trials, potentially increasing their reluctance to fly. This learning process and the disjointed stimulus presentation might have contributed to the long and variable response time delays observed in our study. While these delays provided a unique insight into collective escape, particularly in cases where early detection does not immediately result in early escape, it is possible that this scenario might be more likely with a more naturalistic predator stimulus. Future research could address these limitations by conducting similar studies in the field, outside of the motion-capture system, perhaps using the emerging technology of markerless 3D posture tracking in birds (Waldmann et al., 2022, 2023).

Conclusion

Our research largely supports the common assumptions in vigilance studies. Key among our findings is the relationship between foveation and predator inspection, along with vigilance- and feeding-related behaviors. This suggests that foveation can be a useful proxy for detection in bird vigilance studies. Additionally, we observed that earlier foveation led to earlier flight responses in the flock, facilitated by social contagion. Interestingly, this contagion occurred even though most flock members likely detected the predator cues, as suggested by the long and variable response time delay, which are likely associated with risk assessment. Therefore, our results imply that collective escape does not always involve non-detectors benefiting from detectors. Our study highlights the importance of considering the vision as well as the disparity between detection and escape responses in future vigilance research.

Methods

Subjects

Twenty pigeons (*Columba livia*, 13 females and 7 males) participated in this study. All pigeons originated from a breeder and were juveniles of the same age (1 year old). They were housed together

509 in an aviary (2w x 2d x 2h m; w-width, d-depth, h-height) with perching and nesting structures. The

510 pigeons were fed grains once a day. On experimental days, they were fed after all the experiments

511 were completed. Water and grit were available in the aviary *ad libidum.*

Ethics statement

All the experiments using animals in this study were performed under the license 35-9185.81/G-

19/107 awarded by the Regierungspräsidium Freiburg, Abteilung Landwirtschaft, Ländlicher Raum,

Veterinär- und Lebensmittelwesen, animal ethics authorities of Baden-Württemberg, Germany.

Handling was reduced to a minimum to avoid stress. No animal was hurt or killed during the predator

presentation tests, and they were transferred back to their home aviary directly after the tests. The

pigeons were provided with perches and nesting structures in their aviary, and their health states were

checked on a daily basis. After all experiments were performed, all the individuals were given to a

breeder to continue living as breeding pigeons.

Experimental setup

The experiment took place in SMART-BARN, the state-of-the-art animal tracking system build at the

Max Planck Institute of Animal Behavior (Fig 1a; Nagy et al., 2023). We used the motion-capture

feature of SMART-BARN, which was equipped with 32 motion-capture cameras in an area of 15w x

7d x 4h m (12 Vero v2.2 and 20 Vantage 5, VICON). At two opposite corners of the room, we placed

tables covered with fabric ("predator hiding place"; 105w x 75d x 75h cm), which hide from the

pigeons' view a model predator (plastic sparrow hawk; wingspan 60cm and beak-tail length 35cm). At

the center of the room, we placed jute fabric (4.2l x 3.6w m; l-length) where we scatted food to

encourage the pigeons to stay there during the experiments ("feeding area"). The model predator ran

on a thin wire across the room via a motored pulley mechanism (Wiral LITE kit, Wiral Technologies

AS.) until it disappeared into another hiding place set at the end of the room. On top of each predator

hiding table, we installed a monitor (61.5w x 37h cm, WQHD (2560 x 1440), 144Hz, G-MASTER

GB2760QSU, Iiyama), which displayed a looming stimulus before the model predator appears from

the hiding place. This looming stimulus was a silhouette of a predator mimicking the approach of a

predator toward the pigeon flock. The looming stimulus lasted for 10 seconds, starting from a still

small silhouette of a predator (20w x 4h cm) in the first 5 seconds and then expanding until the silhouette covers the screen entirely in the last 5 seconds. The looming stimulus was played back on a laptop PC (ThinkPad P17 Gen 2, Lenovo) connected to both monitors, and its onset was manually triggered by the experimenter. This onset time was recorded in the motion-capture system via analogue electric signal, specifically by the experimenter pressing a button at the onset of the looming stimulus on a custom Arduino device (Arduino UNO R3, Arduino) connected to the motion capture system. The same Arduino device also indicated the onset of the model predator to the experimenter via a small LED flash.

Experimental procedures

General experimental design

Each pigeon underwent maximum one trial per day and a total of 6 trials. A flock of 10 pigeons participated in each trial, yielding 2 separate flocks in each trial, and a total of 12 trials in the whole experiment. Each pigeon was pseudo-randomly assigned into either flock in each trial, in such a way that each pigeon was paired within a group with any other pigeon at least once across all 6 trials. Each trial composed of 2 predator presentation events, and each event presents a model predator from either the hiding place ("North" or "South" side), yielding 24 events in the whole experiments. The order of presenting the North and South side of hiding places was counterbalanced across groups within each trial.

Preparation for the experiment

Before the experiment, the pigeons were transported from the aviary using a pigeon carrier (80w x 40d x 24h cm). They were then equipped with 4 motion capture markers (6.4mm diameter, OptiTrack) glued on the head feathers and 4 motion capture markers (9mm diameter) on a small solid styrofoam plate (7l x 3.5w cm); this small plate was attached to a backpack worn by each pigeon with Velcro. The experimenter then held the pigeon's head briefly (less than a minute) in a custom 3D frame equipped with 4 web cameras and a triangular scale (26w x 29d x 27h cm) and then filmed the head with web cameras (C270 HD webcam, Logitech) synchronized by a commercial software (MultiCam

Capture, Pinnacle Studio) in order to reconstruct the eyes and beak positions relative to the markers

post hoc (see below for the reconstruction methods). Additionally, the motion capture system was

calibrated just before the experiment starts with an Active Wand (VICON) until all cameras have

recorded 2000 calibration frames.

Trial design

At the start of each trial, all 10 pigeons were released in the motion capture room. After 2 minutes of

acclimatization period, an experimenter scattered a grain-grit mix evenly in the feeding area (Fig 1a-

b). This grain-grit mix comprised of 200g of seeds with 400g of grit to make the foraging task

moderately challenging. The experimenter then hid behind a curtain, outside the motion capture room.

After the experimenter visually confirmed that minimally half of the pigeons (≥5 individuals) started

feeding (pecking grain; this always happened within a few seconds after the experimenter started

scattering the food), a free feeding period started. The duration of this period was randomized

between 2 and 5 minutes across trials to reduce predictability of the predator event. The last 2 minutes

of this feeding period, referred as the "before-cue" period, were used to extract pre-stimulus

behavioral measures. At the end of the feeding period, the experimenter triggered the looming

stimulus. After the offset of the looming stimulus (mean 10.83 ± SD 0.43 s after the onset), the

experimenter ran the model predator on the wire, and the predator took approximately 3 seconds

before it disappeared from the room (mean 2.83 ± SD 1.18 s; Fig 1c). After half of the pigeons resume

feeding (within 1-4 min after the predator disappears), and another free feeding period of 2-5 minutes,

a second predator presentation event occurred following the same procedures, except that the looming

stimulus and the model predator appeared from the other monitor and hiding place. Each trial lasted

for approximately 30 minutes. After the experiment, the motion-capture markers were detached from

each pigeon before being brought back to the aviary.

Data analysis

Reconstruction of the pigeons' head

A custom pipeline (see Data availability section) was used to reconstruct the relative 3D position of the center of the eyes, the beak-tip and the center of the markers from the 4 still images of the pigeon's head (see above). These key points were manually labelled by an experimenter on each picture (an updated algorithm is now available for automatic labelling using YOLO, also included in the deposit). A custom MATLAB program based on structure-from-motion reconstructed the 3D coordinates of all the keypoints with a less than 2 pixels mean reprojection error. These procedures were confirmed to yield accurate reconstruction of head orientations, less than a degree of rotational errors (Itahara & Kano, 2022).

Processing of the motion capture data

The motion capture data were exported as csv files from the motion-capture software (Nexus version 2.14, VICON). The csv files were then imported and processed in the custom codes written in MATLAB (provide in Kano et al., 2022, also provided in our deposit). The motion capture data consisted of a time series of 3D coordinates of the markers attached to the head and back of the pigeons (Fig 2a). From the reconstructed 3D positions of eye centers and beak tip, the local coordinate system of the pigeon's head (or the "visual field" of the pigeon) was defined (Fig 2b). In this local coordinate system, the horizon (the elevation of the local X and Y axes in the global coordinate system) was 30° above from the principal axis of the beak, and the X, Y, Z axis pointed to the right, front, and top of the pigeon's head, respectively. It should be noted that this angle of 30° was determined based on the typical standing postures of pigeons, as identified in Kano et al. (2022). While this previous study determined the angle on a trial-by-trial basis, our study employed a fixed angle following previous recommendations, and also due to the small variation of this angle across trials and individuals.

The data were filtered using the custom pipelines described in Fig S8 of Kano et al. (2022). Briefly, the raw motion-capture data were gap-filled in NEXUS, then smoothed using custom MATLAB codes. The translational and rotational movements of the reconstructed local coordinate

system were further smoothed, and any improbable movements were removed using the same

pipelines. As a result, the loss of head local coordinate system data amounted to 10.07±11.79% (mean

± SD) per individual in each trial, and the loss of back-marker centroids was 2.53±0.98% (mean ±

SD). Although the loss of head data was nontrivial, visual confirmation showed that most loss

occurred when pigeons were self-grooming (thus self-occluding their head markers from the motion

capture cameras); i.e., when they likely were not attending to any object of interest. Head saccades

were also removed from all foveation data due to the possibility that visual processing is inhibited

during saccades in birds (Brooks & Holden, 1973). It should be noted that one deviation from the

previous study in terms of filtering parameters was that the data smoothing was performed at 60Hz,

rather than 30Hz, following previous recommendations.

Reconstruction of gaze vectors

We then reconstructed the gaze (or "foveal") vectors of each pigeon based on the known projected

angles of the foveas (also called *area centralis*), 75° in azimuth and 0° in elevation in the head local

coordinate system (Nalbach et al., 1990). Pigeons use these gaze vectors primarily when attending to

an object/conspecific in the middle to far distance (roughly >50 cm from the head center; Kano et al.,

2022). Although pigeons have another sensitive region of retina, known as red field, this region was

not examined in this study because pigeons primarily use this retinal region to attend to objects in the

close distance (roughly <50 cm), such as pecking and searching for grain (Kano et al., 2022).

Behavioral classification

"Foveation" was defined when an object of interest (such as the monitor or another pigeon) fell within

±10° of the gaze vectors. This margin was estimated to accommodate well the eye movement, which

was typically within 5° in pigeons (Wohlschläger et al., 1993). Head-saccades were defined as any

head movement larger than 5°, that lasted for at least 50ms and faster than 60°/s, and fixations were

defined as inter-saccadic intervals, based on the previous study (Kano et al., 2022). Our objects of

interest included the monitors, the tables that hid the predator, and conspecifics. These objects were

defined as sphere encompassing them (see Supplementary Material and Fig S3). In our observations,

pigeons foveated on the predator cue (monitor or predator hiding table) typically before its offset (Fig

1d); in a few instances, pigeons foveated on the model predator after the looming stimulus had

disappeared, but these cases were excluded from our analysis. To ensure that the first foveation of a

pigeon on the predator cue was not a result of random gaze crossing, we excluded instances of

foveation that were shorter than 300 ms, based on the typical intersaccadic interval for a pigeon,

which typically ranges from 300 to 400 ms (Kano et al., 2022). Moreover, to remove cases where the

gaze vector was on the predator cue at the onset of the looming stimulus, we excluded from the

analysis the first 200 ms of looming stimulus period based on typical reaction times of a pigeon,

which is usually longer than 200 ms (Blough, 1977).

In addition to the foveation, the pigeons' behavior was classified automatically based on

simple thresholds using the 3D postural data (Table 2; for more detailed explanations, see Table S3);

those thresholds were verified by two human raters (see Supplementary Materials and Table S4).

Table 2 - definition of the automated classification of the behaviors. See Table S3

Behavior	Definition
Head-down	The head centroid is lower than the body centroid.
Head-up	The head centroid is higher than the body centroid, and the individual is not exhibiting feeding, grooming, or courting nor courted.
Pecking	The beak tip is close from the ground but not oriented towards the body
Feeding	A period encompassing consecutive instances of pecking
Grooming	The individual is preening its back or breast feathers or scratching its head with its paw; these behaviors are defined based on the head orientation and position relative to the body centroid
Courting	A period encompassing consecutive instances of head bowing (typical courtship movement) while close and oriented toward another pigeon (the courted pigeon was defined as the individual toward which the courting pigeon is oriented)
Running away	Individual is moving fast (>0.6m/s), away from the monitor, and is not

from monitor	courting/courted
Flying	Individual is moving even faster (>2m/s, speed impossible to reach by only running)

Statistical analysis

All the statistical analyses were performed in R (R Core Team, 2022). We mainly relied on linear mixed models or generalized linear mixed models (LMM or GLMM; lmer or glmer function from the lme4 package; Bates et al., 2015) unless otherwise mentioned, and the significance of the predictors was tested using likelihood ratio test. We verified the model assumptions by checking the distribution of the residuals in diagnostic plots (histogram of the residuals, qq-plot and plot of the residuals against fitted values). We transformed the response variable (logit- or log-transformation) when it improved the normality of the residual distribution. To test for collinearity, we also checked the variance inflation factor (VIF) of the predictors. In all models, the continuous predictors were normalized (z-transformed). For all models, in addition to the test predictors, we also included several control variables: the event number within a trial (1 or 2), the predator side ("North" or "South"), and the sex of the subject, as well as the pigeon ID and the trial ID as random effects (see Table S6 for the used R formulas). We report the effects of test variables in the Result section and report the effects of all test and control variables in Table S7. Detailed descriptions about the responses and test variables can be found in Table S2.

Acknowledgements

We thank the members of the Max Planck Institute of Animal Behavior (MPI-AB) as well as the Centre for the Advanced Study of Collective Behaviour (CASCB) for their valuable supports in conducting this study. Special thanks are extended to Drs. Mate Nagy, Dora Biro, Oliver Deussen, and Iain D. Couzin, along with the members of MPI-AB and CASCB, for their insightful comments on our study. We also thank Drs. Inge Müller and Daniel Zuniga, as well as the caretakers, for their

675   dedicated hosting and care of the pigeons. Finally, we thank Mathias Günther and Alex Chan for

676   technical support. This study was financially supported by MPI-AB, the DFG Cluster of Excellence

677   2117 CASCB (ID: 422037984), and the CASCB BigChunk projects (ID: L21-07).

Data availability

680   A OSF repository containing the code used for the processing and analysis can be found on the OSF

681   website (Delacoux & Kano, 2023). Sample data is also provided for the user (the whole raw dataset

682   could not be included for size reason). The repository contains the code for the automatic labelling of

683   the head calibration of the pigeons, the Matlab pipeline used for the motion capture data processing

684   and filtering, and the statistical analyses in R (along with the tables containing the processed data).

Reference

Barbosa, P., & Castellanos, I. (2005). *Ecology of Predator-Prey Interactions*. Oxford University Press,

USA.

Bates, D., Mächler, M., Bolker, B., & Walker, S. (2015). Fitting Linear Mixed-Effects Models Using

lme4. *Journal of Statistical Software*, *67*(1), 1–48. https://doi.org/10.18637/jss.v067.i01

Beauchamp, G. (2008). What is the magnitude of the group-size effect on vigilance? *Behavioral

Ecology*, *19*(6), 1361–1368. https://doi.org/10.1093/beheco/arn096

Beauchamp, G. (2015). *Animal Vigilance: Monitoring Predators and Competitors*. Academic Press.

Beauchamp, G., & Ruxton, G. D. (2008). Disentangling risk dilution and collective detection in the

antipredator vigilance of semipalmated sandpipers in flocks. *Animal Behaviour*, *75*(6), 1837–

1842. https://doi.org/10.1016/j.anbehav.2007.12.016

Beauchamp, G., & Ruxton, G. D. (2016). Modeling scan and interscan durations in antipredator

vigilance. *Journal of Theoretical Biology*, *390*, 86–96.

https://doi.org/10.1016/j.jtbi.2015.11.015

Bednekoff, P. A., & Lima, S. L. (1998). Re–examining safety in numbers: Interactions between risk dilution and collective detection depend upon predator targeting behaviour. *Proceedings of the Royal Society of London. Series B: Biological Sciences*, *265*(1409), 2021–2026. https://doi.org/10.1098/rspb.1998.0535

Bednekoff, P. A., & Lima, S. L. (2002). Why are scanning patterns so variable? An overlooked question in the study of anti-predator vigilance. *Journal of Avian Biology*, *33*(2), 143–149. https://doi.org/10.1034/j.1600-048X.2002.330204.x

Bednekoff, P. A., & Lima, S. L. (2005). Testing for peripheral vigilance: Do birds value what they see when not overtly vigilant? *Animal Behaviour*, *69*(5), 1165–1171. https://doi.org/10.1016/j.anbehav.2004.07.020

Blough, D. S. (1977). Visual Search in the Pigeon: Hunt and Peck Method. *Science*, *196*(4293), 1013–1014. https://doi.org/10.1126/science.860129

Boland, C. R. J. (2003). An Experimental Test of Predator Detection Rates Using Groups of Free-living Emus. *Ethology*, *109*(3), 209–222. https://doi.org/10.1046/j.1439-0310.2003.00860.x

Brooks, B., & Holden, A. L. (1973). Suppression of visual signals by rapid image displacement in the pigeon retina: A possible mechanism for "saccadic" suppression. *Vision Research*, *13*, 1387–1390. https:// doi. org/ 10. 1016/ 0042- 6989(73) 90216-2

Butler, S. R., & Fernández-Juricic, E. (2018). European starlings use their acute vision to check on feline predators but not on conspecifics. *PLOS ONE*, *13*(1), e0188857. https://doi.org/10.1371/journal.pone.0188857

Cresswell, W. (1993). Escape responses by redshanks, *Tringa totanus*, on attack by avian predators. *Animal Behaviour*, *46*(3), 609–611. https://doi.org/10.1006/anbe.1993.1231

Cresswell, W., Butler, S., Whittingham, M. J., & Quinn, J. L. (2009). Very Short Delays Prior to Escape from Potential Predators May Function Efficiently as Adaptive Risk-Assessment Periods. *Behaviour*, *146*(6), 795–813.

Cresswell, W., Hilton, G. M., & Ruxton, G. D. (2000). Evidence for a rule governing the avoidance of superfluous escape flights. *Proceedings of the Royal Society of London. Series B: Biological Sciences*, *267*(1444), 733–737. https://doi.org/10.1098/rspb.2000.1064

Cresswell, W., Quinn, J. L., Whittingham, M. J., & Butler, S. (2003). Good foragers can also be good at detecting predators. *Proceedings of the Royal Society of London. Series B: Biological Sciences*, *270*(1519), 1069–1076. https://doi.org/10.1098/rspb.2003.2353

Davis, J. M. (1975). Socially induced flight reactions in pigeons. *Animal Behaviour*, *23*, 597–601. https://doi.org/10.1016/0003-3472(75)90136-0

Delacoux, M., & Kano, F. (2023). *Fine-scale tracking reveals visual field use for predator detection and escape in collective foraging of pigeon flocks* [dataset]. OSF. osf.io/d682s

Devereux, C. L., Whittingham, M. J., Fernández-Juricic, E., Vickery, J. A., & Krebs, J. R. (2006). Predator detection and avoidance by starlings under differing scenarios of predation risk. *Behavioral Ecology*, *17*(2), 303–309. https://doi.org/10.1093/beheco/arj032

Elgar, M. A. (1989). Predator Vigilance and Group Size in Mammals and Birds: A Critical Review of the Empirical Evidence. *Biological Reviews*, *64*(1), 13–33. https://doi.org/10.1111/j.1469-185X.1989.tb00636.x

Evans, D. A., Stempel, A. V., Vale, R., & Branco, T. (2019). Cognitive Control of Escape Behaviour. *Trends in Cognitive Sciences*, *23*(4), 334–348. https://doi.org/10.1016/j.tics.2019.01.012

Fernández-Juricic, E. (2012). Sensory basis of vigilance behavior in birds: Synthesis and future prospects. *Behavioural Processes*, *89*(2), 143–152. https://doi.org/10.1016/j.beproc.2011.10.006

Fernández-Juricic, E., Delgado, J. A., Remacha, C., Jiménez, M. D., Garcia, V., & Hori, K. (2009). Can a solitary avian species use collective detection? An assay in semi-natural conditions. *Behavioural Processes*, *82*(1), 67–74. https://doi.org/10.1016/j.beproc.2009.05.002

Fernández-Juricic, E., Erichsen, J. T., & Kacelnik, A. (2004). Visual perception and social foraging in birds. *Trends in Ecology & Evolution*, *19*(1), 25–31. https://doi.org/10.1016/j.tree.2003.10.003

Fuchs, R., Veselý, P., & Nácarová, J. (2019). *Predator Recognition in Birds: The Use of Key Features*. https://doi.org/10.1007/978-3-030-12404-5

Godin, J.-G. J., & Smith, S. A. (1988). A fitness cost of foraging in the guppy. *Nature*, *333*(6168), 69–71. https://doi.org/10.1038/333069a0

Hart, A., & Lendrem, D. W. (1984). Vigilance and scanning patterns in birds. *Animal Behaviour*,

32(4), 1216–1224. https://doi.org/10.1016/S0003-3472(84)80239-0

Hilton, G. M., Cresswell, W., & Ruxton, G. D. (1999). Intraflock variation in the speed of escape-

flight response on attack by an avian predator. *Behavioral Ecology*, *10*(4), 391–395.

https://doi.org/10.1093/beheco/10.4.391

Inglis, I. R., & Lazarus, J. (1981). Vigilance and Flock Size in Brent Geese: The Edge Effect.

Zeitschrift Für Tierpsychologie, *57*(3–4), 193–200. https://doi.org/10.1111/j.1439-

0310.1981.tb01921.x

Itahara, A., & Kano, F. (2022). "Corvid Tracking Studio": A custom-built motion capture system to

track head movements of corvids. *Japanese Journal of Animal Psychology*, *72*(1), 1–16.

https://doi.org/10.2502/janip.72.1.1

Jones, K. A., & Godin, J.-G. J. (2010). Are fast explorers slow reactors? Linking personality type and

anti-predator behaviour. *Proceedings of the Royal Society B: Biological Sciences*, *277*(1681),

625–632. https://doi.org/10.1098/rspb.2009.1607

Jones, K. A., Krebs, J. R., & Whittingham, M. J. (2007). Vigilance in the third dimension: Head

movement not scan duration varies in response to different predator models. *Animal*

Behaviour, *74*(5), 1181–1187. https://doi.org/10.1016/j.anbehav.2006.09.029

Jones, K. A., Krebs, J. R., & Whittingham, M. J. (2009). Heavier birds react faster to predators:

Individual differences in the detection of stalking and ambush predators. *Behavioral Ecology*

and Sociobiology, *63*(9), 1319–1329. https://doi.org/10.1007/s00265-009-0778-6

Kaby, U., & Lind, J. (2003). What limits predator detection in blue tits *(Parus caeruleus)*: Posture,

task or orientation? *Behavioral Ecology and Sociobiology*, *54*(6), 534–538.

https://doi.org/10.1007/s00265-003-0665-5

Kano, F., Naik, H., Keskin, G., Couzin, I. D., & Nagy, M. (2022). Head-tracking of freely-behaving

pigeons in a motion-capture system reveals the selective use of visual field regions. *Scientific*

Reports, *12*(1), Article 1. https://doi.org/10.1038/s41598-022-21931-9

Kenward, R. E. (1978). Hawks and Doves: Factors Affecting Success and Selection in Goshawk

Attacks on Woodpigeons. *Journal of Animal Ecology*, *47*(2), 449–460.

https://doi.org/10.2307/3793

Lima, S. L. (1995a). Back to the basics of anti-predatory vigilance: The group-size effect. *Animal*

Behaviour, *49*(1), 11–20. https://doi.org/10.1016/0003-3472(95)80149-9

Lima, S. L. (1995b). Collective detection of predatory attack by social foragers: Fraught with

ambiguity? *Animal Behaviour*, *50*(4), 1097–1108. https://doi.org/10.1016/0003-

3472(95)80109-X

Lima, S. L., & Bednekoff, P. A. (1999). Back to the basics of antipredatory vigilance: Can nonvigilant

animals detect attack? *Animal Behaviour*, *58*(3), 537–543.

https://doi.org/10.1006/anbe.1999.1182

Lima, S. L., & Dill, L. M. (1990). Behavioral decisions made under the risk of predation: A review

and prospectus. *Canadian Journal of Zoology*, *68*(4), 619–640. https://doi.org/10.1139/z90-

092

Lima, S. L., & Zollner, P. A. (1996). Anti-predatory vigilance and the limits to collective detection:

Visual and spatial separation between foragers. *Behavioral Ecology and Sociobiology*, *38*(5),

355–363. https://doi.org/10.1007/s002650050252

Martin, G. R. (2017). What Drives Bird Vision? Bill Control and Predator Detection Overshadow

Flight. *Frontiers in Neuroscience*, *11*.

https://www.frontiersin.org/articles/10.3389/fnins.2017.00619

Nagy, M., Naik, H., Kano, F., Carlson, N. V., Koblitz, J. C., Wikelski, M., & Couzin, I. D. (2023).

SMART-BARN: Scalable multimodal arena for real-time tracking behavior of animals in

large numbers. *Science Advances*, *9*(35), eadf8068. https://doi.org/10.1126/sciadv.adf8068

Nalbach, H.-O., Wolf-Oberhollenzer, F., & Kirschfeld, K. (1990). The pigeon's eye viewed through an

ophthalmoscopic microscope: Orientation of retinal landmarks and significance of eye

movements. *Vision Research*, *30*(4), 529–540. https://doi.org/10.1016/0042-6989(90)90065-S

Proctor, C. J., Broom, M., & Ruxton, G. D. (2001). Modelling antipredator vigilance and flight response in group foragers when warning signals are ambiguous. *Journal of Theoretical Biology*, *211*(4), 409–417. https://doi.org/10.1006/jtbi.2001.2353

Pulliam, H. R. (1973). On the Advantages of Flocking. *Journal of Theoretical Biology*, *38*, 419–422.

Pulliam, H. R., Pyke, G. H., & Caraco, T. (1982). The scanning behavior of juncos: A game-theoretical approach. *Journal of Theoretical Biology*, *95*(1), 89–103. https://doi.org/10.1016/0022-5193(82)90289-2

Quinn, J. L., & Cresswell, W. (2005). Escape response delays in wintering redshank, *Tringa totanus*, flocks: Perceptual limits and economic decisions. *Animal Behaviour*, *69*(6), 1285–1292. https://doi.org/10.1016/j.anbehav.2004.10.007

R Core Team. (2022). *R: A Language and Environment for Statistical Computing* [Computer software]. R Foundation for Statistical Computing. https://www.R-project.org/

Roberts, G. (1996). Why individual vigilance declines as group size increases. *Animal Behaviour*, *51*(5), 1077–1086. https://doi.org/10.1006/anbe.1996.0109

Rogers, L. J., Zucca, P., & Vallortigara, G. (2004). Advantages of having a lateralized brain. *Proceedings of the Royal Society of London. Series B: Biological Sciences*, *271*(suppl_6), S420–S422. https://doi.org/10.1098/rsbl.2004.0200

Tätte, K., Ibáñez-Álamo, J. D., Markó, G., Mänd, R., & Møller, A. P. (2019). Antipredator function of vigilance re-examined: Vigilant birds delay escape. *Animal Behaviour*, *156*, 97–110. https://doi.org/10.1016/j.anbehav.2019.08.010

Tisdale, V., & Fernández-Juricic, E. (2009). Vigilance and predator detection vary between avian species with different visual acuity and coverage. *Behavioral Ecology*, *20*(5), 936–945. https://doi.org/10.1093/beheco/arp080

Tyrrell, L. P., Butler, S. R., Yorzinski, J. L., & Fernández-Juricic, E. (2014). A novel system for bi-ocular eye-tracking in vertebrates with laterally placed eyes. *Methods in Ecology and Evolution*, *5*(10), 1070–1077. https://doi.org/10.1111/2041-210X.12249

Veen, T., Richardson, D. S., Blaakmeer, K., & Komdeur, J. (2000). Experimental evidence for innate

predator recognition in the Seychelles warbler. *Proceedings of the Royal Society of London.*

Series B: Biological Sciences, *267*(1459), 2253–2258. https://doi.org/10.1098/rspb.2000.1276

Waldmann, U., Chan, A. H. H., Naik, H., Nagy, M., Couzin, I. D., Deussen, O., Goldluecke, B., &

Kano, F. (2023). *3D-MuPPET: 3D Multi-Pigeon Pose Estimation and Tracking*

(arXiv:2308.15316). arXiv. https://doi.org/10.48550/arXiv.2308.15316

Waldmann, U., Naik, H., Máté, N., Kano, F., Couzin, I. D., Deussen, O., & Goldlücke, B. (2022). I-

MuPPET: Interactive Multi-Pigeon Pose Estimation and Tracking. In B. Andres, F. Bernard,

D. Cremers, S. Frintrop, B. Goldlücke, & I. Ihrke (Eds.), *Pattern Recognition* (pp. 513–528).

Springer International Publishing. https://doi.org/10.1007/978-3-031-16788-1_31

Whittingham, M. J., Butler, S. J., Quinn, J. L., & Cresswell, W. (2004). The effect of limited visibility

on vigilance behaviour and speed of predator detection: Implications for the conservation of

granivorous passerines. *Oikos*, *106*(2), 377–385. https://doi.org/10.1111/j.0030-

1299.2004.13132.x

Wohlschläger, A., Jäger, R., & Delius, J. D. (1993). Head and eye movements in unrestrained pigeons

(Columba livia). *Journal of Comparative Psychology*, *107*(3), 313–319.

https://doi.org/10.1037/0735-7036.107.3.313

Wortel, J. F., Wubbels, R. J., & Nuboer, J. F. W. (1984). Photopic spectral sensitivities of the red and

the yellow field of the pigeon retina. *Vision Research*, *24*(9), 1107–1113.

https://doi.org/10.1016/0042-6989(84)90089-0

Yorzinski, J. L., & Platt, M. L. (2014). Selective attention in peacocks during predator detection.

Animal Cognition, *17*(3), 767–777. https://doi.org/10.1007/s10071-013-0708-x

Supplementary Materials:

Fine-scale tracking reveals visual field use for predator

detection and escape in collective foraging of pigeon flocks

862 Mathilde Delacoux, Fumihiro Kano

Overview of observations

864 Table S1: Total number of individuals that exhibited foveation, escape, and flight behavior over the
865 course of the trials and events. Each event was composed of a flock of ten pigeons.

Trial	Event	Group	# individuals foveating	# individuals escaping	# individuals flying
1	1	1	6	5	4
1	2	1	9	9	7
1	1	2	5	4	4
1	2	2	8	9	9
2	1	1	9	9	6
2	2	1	10	10	6
2	1	2	6	9	9
2	2	2	8	9	8
3	1	1	10	9	10
3	2	1	10	10	6
3	1	2	8	9	9
3	2	2	9	10	8
4	1	1	8	10	8
4	2	1	9	10	7
4	1	2	9	10	6
4	2	2	5	10	10
5	1	1	9	9	6
5	2	1	10	10	6
5	1	2	10	8	0
5	2	2	9	9	6
6	1	1	8	10	7
6	2	1	10	6	0
6	1	2	8	10	6
6	2	2	8	10	9

870 Figure S3: Response variables as a function of the trials and events. a) latency to foveate; b) latency to
871 escape; c) latency to fly. The distribution of the data is displayed as violin plots and the color
872 represents the side of the predator event.

874 Figure S4: Predictor variables as a function of the trials and events. The distributions of the different
875 variables are displayed as violin plots and the color represents the side of the predator event. a-e)

behavior during the "before cue" period (2 minutes period straight before the onset of the looming stimulus) as a function of the trial and event: proportion of time head-up (a), proportion of time foveating on either of the monitors (b), proportion of time foveating on one of the conspecifics (c), saccade rate (d), pecking rate (e). f-h) behavioral and spatial state measures at the onset of the looming stimulus as a function of the trial and event: distance to the monitor (f), distance to the nearest neighbor pigeon (g), the number of individuals within a flock exhibiting respective behaviors (feeding, head-up or other behavior) exhibited at the onset on the looming stimulus (h).

Definition of object shape

We approximated objects of interest (monitors, hiding places, and other pigeons) as 3D volumes, defining them as spheres or combinations of spheres.

The monitors were represented as a single sphere. The sphere's center was the monitor's centroid, and its diameter was set at 86.13 cm, which is the monitor's diagonal length of 71.77 cm plus a 20% margin. The predator hiding place was also defined as a single sphere, with its center located below the monitor's centroid at a height of 38 cm above the ground (along the z-axis of the global coordinate system, corresponding to half the height of the hiding place). Its diameter was 158.47 cm, based on the hiding place's diagonal length of 132.06 cm, plus a 20% margin.

The pigeons were represented by two spheres, denoting the head and the body. The head sphere's centroid was the midpoint between the eyes, with a diameter of 12 cm (calculated from the distance from the beak tip to the midpoint of the eyes, approximately 4 cm, plus a 2 cm margin for the radius). The body sphere's centroid was set 6 cm below the center point of the backpack markers (along the Z-axis of the global coordinate system, approximately at the body's midpoint directly above the pigeon's paws), and its diameter was 24 cm (considering the centroid was usually around 10 cm above the ground, with an added 20% margin for the radius) (Fig S3).

Figure S3: Illustration of the defined spherical representations for the pigeon's head and body.

Definition of responses

Table S2: Explanation of the different responses and variables used in our models.

Variable type	Test Set	Variable name	description
Responses	1	Proportion of time foveating	Normalized frequency (normalized to a unit sum) at which each object of interest was observed within the foveal region (in the reprojection of the pigeon's visual

			field), during the predator event.
	2.1-2.2	Latency to foveate (ms)	Latency of the first foveation on the looming stimulus or the predator hiding place after the onset of the looming stimulus and before the predator appears. To exclude the possibility of random gaze crossing, the first foveation was defined as instances lasting longer than 300 ms, which occurred at least 200 ms after the onset of the looming stimulus.
	3-4.1	Escaped before the looming stimulus offset or not (binary)	Whether the first overt escape response (either running away or flying) occurred before the looming stimulus offset (1) or not (0; i.e., either the pigeon responded after the looming stimulus offset or did not exhibit a detectable response).
	3-4.1	Latency to fly (ms)	Latency of the first flight after the onset of the looming stimulus
Predictors	1	Object type	Type of object: the monitor displaying the looming stimulus (relevant environmental cue), the monitor not displaying the looming stimulus (irrelevant environmental cue), or the nearest neighbor (conspecific)
	2.1	Behavior at the onset	Behavior at the onset of the looming stimulus (either head-up or feeding, other behaviors are not considered as they occurred much less often (20 occurrences out of 240 observations)
	2.1	Distance from the monitor (mm)	Distance to the monitor at the onset of the looming stimulus
	2.1	Distance to nearest neighbor (mm)	Distance to the nearest neighbor at the onset of the looming stimulus
	2.2	Head-up	Proportion of time the pigeon spent being head-up during the 2-min "before-cue" period
	2.2	Foveate on the monitor	Proportion time spent foveating on either monitor during the 2-min "before-cue" period
	2.2	Foveate on any pigeon	Proportion time spent foveating on any pigeon during the 2-min "before-cue" period
	2.2	Saccade rate (saccades/s)	Saccade rate when the pigeon is head-up during the 2-min "before-cue" period
	2.2	Pecking rate (pecks/s)	Pecking rate during the 2-min "before-cue" period
	3	Latency to foveate (ms)	See above
	4.1	First latency to foveate (ms)	The latency to foveate on the predator cue by the first pigeon during the looming stimulus presentation

Automated behavior classification and rating

Behaviors were automatically classified using postural definitions on the 3D data from the motion capture system (Table S3).

Table S3: Description of the behavior classification definitions. Terms definitions: state = behavior that lasts in time; event = punctual behavior. Body-head vector = the vector originating from the body center and projecting to the head centroid. Head direction vector = the vector projecting to the front of the head and towards the horizon when the head is still (corresponds to the y axis of the local coordinate system, projecting 30° above the beak-head center axis). Pitch = head rotation movement corresponding to "nodding" (pitch = 0° corresponds to the head direction vector pointing towards horizon, pitch < 0° corresponds to the head direction vector pointing down, pitch > 0° corresponds to the head direction vector pointing up). Roll = head rotation movement where the head is "tilted" (roll = 0° corresponds to a straight head and roll ≠ 0° corresponds to a head tilted to the right or to the left).

Behavior	Category	Definition
head-down	state	The centroid of the head (middle point between the eyes) is lower than the centroid of the body (6cm below the backpack markers center point) along the z axis of the global coordinate system (=vertically)
head-up	state	The pigeon is not head-down and not exhibiting another behavior (feeding, grooming or courting)
pecking	event	The beak tip is closer than 4cm from the ground and is not oriented towards the body (pitch > -100° from horizon). The pecking event is defined as the frame where the beak tip is the closest from the ground
feeding	state	Any time between 2 pecks spaced no more than 6 seconds apart

grooming	state	Preening the back feathers: This occurs when the head is pointing backward (defined as when the head direction vector and the body-head vector form an angle larger than 60°) and the beak tip is close to the body center (less than 10cm) OR when the head is pointing backward (defined as when the head direction vector and body-head vector form an angle larger than 40°) and the beak tip is close to the body centroid (less than 10cm), and the head centroid is above the body centroid. Preening the breast feathers: This behavior occurs when the head is strongly pointing down (pitch < -100°) OR when the head is pointing down (pitch < -80°) and the head direction vector and the body-head vector form an angle larger than 60°. Scratching the head with the paw: This behavior is exhibited when the roll of the head is large (> 50°) and the head is low (less than 3cm above the body center).
courting	state	The pigeon is bowing its head, which is a typical head movement in pigeon courtship, at a rate of at least 2 bows within an 8-second period, while being within a proximity of less than 60cm and oriented towards another pigeon within an angle of less than 120°. We filtered out any instances shorter than 1 second for robust detection.
running away from the monitor	state	When the body centroid of the pigeon moves at a speed faster than 0.6m/s, and the pigeon is not classified as courting or being courted, while also increasing its distance from the monitor at a speed of 0.2m/s, we filter out any events shorter than 70ms for robust detection.
flying	state	When the body centroid of the pigeon moves at a speed greater than 2m/s, we filter out any events shorter than 70ms for robust detection.

The reliability of the automated classification was verified by two human raters through manual coding. This process was conducted on a dataset different from the one used in this study. The raters assessed the presence or absence of specific behaviors (or the number of pecks) in 1-second sample segments (for coding peck counts, 3-second segments were used to increase variability in the number of pecks). We coded 50 segments for each behavior, with the exception of flying (for which only 30 segments were coded due to its scarcity in the dataset) and pecking (for which 60 segments were coded to enhance variation in peck counts). Behaviors such as head-down, head-up, and feeding were not rated, as they were determined based on a simple threshold definition and/or depended on rated behaviors. Inter-rater reliability was calculated using R (R Core Team, 2022) and the 'irr' package (Gamer et al., 2019). The pecking count IRR was computed using Intraclass Correlation Coefficient (ICC) for count data, while grooming, courting, and flying were analyzed using Cohen's kappa for binary, presence-absence data.

The head-up/down state was defined based on the relative position of the head and body. The head-up state was further refined to exclude periods of feeding, grooming, and courting, to avoid moments when pigeons were likely distracted by these activities. To verify the detection of pecks, two human raters counted the number of pecks in short data segments. The Intraclass Correlation Coefficient (ICC) confirmed a very high agreement between their coding and automated detection ($ICC\_min =$ 0.98; see Table S4). Feeding was then defined based on consecutive pecking instances. Grooming and courting were identified by a combination of several postural states and validated by the same two human raters (presence or absence of the behavior in short data segments; Cohen's $kappa\_min =$ 0.8; see Table S4). Running was defined based on a speed threshold, determined by inspecting locomotive speed histograms (Fig S4). This definition was limited to pigeons moving away from the predator hiding place and not engaged in courting activities, to identify 'running away from the monitor' as an evasion response. Flying was observable when a pigeon took off but was specifically defined by the locomotive speed exceeding 2 m/s, a threshold unattainable by running alone. This threshold was further validated by the human raters' coding (presence or absence of the behavior in short segments; Cohen's $kappa\_min =$ 0.93; see Table S4). All rated behaviors achieved Cohen's Kappa or Intraclass Coefficient (ICC) scores above 0.8 ("almost perfect"; Landis & Koch, 1977), indicating a strong agreement between human raters and the automated behavior coding.

Table S4: Inter-rater reliability for behavior classification. Pecking count was rated with ICC and other behaviors with Cohen's kappa. HR1 and HR2 represent the human raters and ACC the automated computer classification.

Behavior	Inter-rater reliability			Sample number	Metric	Comparison method
	HR1-ACC	HR2-ACC	HR1-HR2			
Pecking	0.983	0.984	0.987	60	count	ICC
Grooming	1	0.96	0.96	50	presence-absence	Cohen's Kappa
Courting	0.88	0.8	0.838	50	presence-absence	Cohen's Kappa
Flying	0.933	1	0.933	30	presence-absence	Cohen's Kappa

Figure S4: Histogram of the pigeons' speed for the whole dataset. Running unusually fast is defined as the body center moving with a speed larger than 0.6 (vertical blue line)

Behavioral changes after the predator presentation event

We investigated whether various behavioral and spatial measures were influenced by a predation event. We collected data for all measures over a one-minute period immediately preceding the onset of the looming stimulus and following the predator's disappearance (averaging the distances over the same period). Linear Mixed Models (LMMs) were used, with the behavior/distance as the response variable. The time period (before vs. after the event) served as the test variable, while event type, predator side, and sex were control variables. Pigeon ID and trial were included as random effects. For all tested behaviors, measurements taken after the predation event showed significant differences compared to those before (p < 0.001). The distances also exhibited significant changes (distance from the closest monitor: p = 0.0034; distance from the nearest neighbor: p = 0.0043; see Fig S5 for a graphical representation and Table S5 for detailed model results).

Figure S5: Behavioral changes between the 1 min period before the onset of the looming stimulus and
the 1 min period after the disappearance of the predator: proportion of time spent being head-up (a),
proportion of time spent foveating on one of the monitors (b), proportion of time spent foveating on
one of the conspecifics (c), saccade rate (d), pecking rate (e), distance to the closest monitor (f),
distance to the nearest neighbor (g). The dots represent single observations (jittered horizontally for
visualization), the grey shade represents the violin plot of the distributions, and the boxplot's boxes
represent the 0.25 and 0.75 quartiles (with the median represented as a line inside the box) and the
whiskers the minimum and maximum values within the lower/upper quartile ± 1.5 times the
interquartile range.

Table S5: Model results for behavioral comparison between the 1 min period before the onset of the
looming stimulus and the 1 min period after the disappearance of the predator. For all models, the
table details the beta estimates, as well as the Chi square value, degrees of freedom and p-values from
the likelihood ratio test. Event, predator side, sex, pigeon ID (random effect) and trial (random effect)
has been added as control variables.

Response	Type	Variable	# of obs.	Beta	Chi-sq	Df	p
Head-up	Test variable	Time period	477	0.27	237.71	1	<0.00001
	Control variable	Event	477	0.13	33.39	1	<0.00001
		Predator side (North)	477	0.00	0.05	1	0.832
		Sex (male)	477	-0.08	3.18	1	0.07433
Pecking rate (log-transformed)	Test variable	Time period	477	0.27	237.71	1	<0.00001
	Control variable	Event	477	0.13	33.39	1	<0.00001
		Predator side (North)	477	0.00	0.05	1	0.832
		Sex (male)	477	-0.08	3.18	1	0.07433
Saccade rate	Test variable	Time period	477	0.27	94.75	1	<0.00001
	Control variable	Event	477	0.09	5.49	1	0.01907
		Predator side (North)	477	-0.02	0.35	1	0.55565
		Sex (male)	477	-0.03	0.15	1	0.69753
Proportion of	Test variable	Time period	477	0.09	174.32	1	<0.00001

time looking at any monitor	Control variable	Event	477	0.02	6.09	1	0.01357
		Predator side (North)	477	0.00	0.01	1	0.93284
		Sex (male)	477	-0.04	2.47	1	0.11573
Proportion of time looking at any conspecific	Test variable	Time period	477	0.07	76.63	1	<0.00001
	Control variable	Event	477	0.00	0.09	1	0.77029
		Predator side (North)	477	-0.01	0.89	1	0.34477
		Sex (male)	477	0.04	2.48	1	0.11564
Distance to the closest monitor	Test variable	Time period	477	101.61	8.58	1	0.00339
	Control variable	Event	477	9.47	0.04	1	0.84643
		Predator side (North)	477	51.44	1.11	1	0.2922
		Sex (male)	477	8.38	0.01	1	0.91392
Distance to the nearest neighbor	Test variable	Time period	477	-31.11	8.15	1	0.0043
	Control variable	Event	477	6.07	0.16	1	0.69182
		Predator side (North)	477	26.76	3.03	1	0.08195
		Sex (male)	477	-50.13	1.84	1	0.17515

R formula

987 Table S6: R formulas used for the different models. Variables starting with "Logit" have been logit-
988 transformed, with "Log" have been log-transformed. As control variables, we have included in all
989 models the event number, the predator side, the sex of the individual, the pigeon ID (random effect)
990 and the trial ID (random effect).

Test	Formula
1	lmer(Logit_prop_in_foveal_region ~ Object_type + Event + Predator_side + Sex + (1|Pigeon_ID) + (1|Trial), Data)
2.1	lmer(Log_latency_foveate ~ scale(Distance_monintor) + scale(Distance_NN) + Behavior_at_onset + Event + Predator_side + Sex + (1|Pigeon_ID) + (1|Trial), Data)
2.2	lmer(Log_latency_foveate ~ scale(Prop_headup) + Event + Predator_side + Sex + (1|Pigeon_ID) + (1|Trial), Data) In other models, Prop_headup replaced with: Saccade_rate, Peck_rate, Prop_foveate_monitor or Prop_foveate_conspecific
3	glmer(Escape_before_cue_offset ~ scale(Latency_foveate) + Event + Predator_side + Sex + (1|Pigeon_ID) + (1|Trial), Data, family = binomial(link = "logit")) lmer(Latency_fly ~ scale(Latency_foveate) + Event + Predator_side + Sex + (1|Pigeon_ID) + (1|Trial), Data)
4.1	glmer(Escape_before_cue_offset ~ scale(First_latency_foveate) + Event + Predator_side + Sex + (1|Pigeon_ID) + (1|Trial), Data, family = binomial(link = "logit")) lmer(Latency_fly ~ scale(First_latency_foveate) + Event + Predator_side + Sex + (1|Pigeon_ID) + (1|Trial), Data)

Results of main models

Table S7: Results of all models related to our main hypotheses. This table presents detailed information for each predictor in every model used in this study, including the beta estimates from the model, as well as the Chi-square value, degrees of freedom, and p-values derived from the likelihood ratio test. Note: (a) was incorporated as a random effect in the model and, as such, does not have a beta estimate.

Test	Response	Type	Variable	# of obs.	Beta	Chi-sq	Df	p
1	Proportion of time foveating on the objects of interest	Test variable	Object type (monitor not presenting stimulus)	720	-0.93	108.07	2	<0.00001
			Object type (Nearest neighbor)	720	-1.07	108.07	2	<0.00001
		Control variable	Event	720	0.19	4.86	1	0.02755
			Predator side (North)	720	-0.06	0.55	1	0.46013
			Sex (male)	720	-0.02	0.03	1	0.85906
2.1	Latency to foveate	Test variable	Distance from the monitor	163	0.14	1.20	1	0.27397
			Distance from NN	163	-0.03	0.05	1	0.82479
			Behavior (head-up)	163	-0.75	13.78	1	0.00021
		Control variable	Event	163	-0.21	1.30	1	0.25377
			Predator side (North)	163	0.09	0.19	1	0.66482
			Sex (male)	163	0.30	2.54	1	0.1107
2.2	Latency to foveate	Test variable	Proportion of time head-up	195	-0.35	15.90	1	0.00007
		Control variable	Event	195	-0.03	0.02	1	0.89658
			Predator side (North)	195	-0.14	0.80	1	0.36962
			Sex (male)	195	0.17	1.04	1	0.30679
	Latency to foveate	Test variable	Pecking rate	195	0.32	14.07	1	0.0002
		Control variable	Event	195	-0.08	0.19	1	0.6649
			Predator side (North)	195	-0.13	0.74	1	0.3904
			Sex (male)	195	0.25	2.41	1	0.1202
	Latency to foveate	Test variable	Proportion of time foveating on the monitor	195	-0.30	12.04	1	0.00052
		Control variable	Event	195	-0.14	0.72	1	0.39587
			Predator side (North)	195	-0.11	0.55	1	0.45892
			Sex (male)	195	0.20	1.44	1	0.22948
	Latency to foveate	Test variable	Saccade rate	195	-0.26	9.85	1	0.0017
		Control variable	Event	195	-0.20	1.54	1	0.2139
			Predator side (North)	195	-0.15	0.97	1	0.32492

	Response	Type	Predictor	N	Estimate	Stat	df	p
			Sex (male)	195	0.25	2.37	1	0.12403
	Latency to foveate	Test variable	Proportion of time foveating on conspecifics	195	0.02	0.06	1	0.81159
		Control variable	Event	195	-0.30	3.42	1	0.06454
			Predator side (North)	195	-0.10	0.44	1	0.50919
			Sex (male)	195	0.29	2.90	1	0.08863
3	Latency to escape	Test variable	Latency to foveate	201	-0.03	0.03	1	0.8716
		Control variable	Event	201	-0.04	0.01	1	0.92304
			Predator side (North)	201	1.57	18.00	1	0.00002
			Sex (male)	201	-0.58	1.16	1	0.28183
	Latency to fly	Test variable	Latency to foveate	201	214.61	6.49	1	0.01084
		Control variable	Event	201	77.75	0.32	1	0.57313
			Predator side (North)	201	-320.20	5.01	1	0.02523
			Sex (male)	201	49.33	0.06	1	0.80554
4.1	Latency to escape	Test variable	First latency to foveate	210	-0.70	3.66	1	0.05586
		Control variable	Event	210	-0.07	0.03	1	0.86216
			Predator side (North)	210	1.49	14.93	1	0.00011
			Sex (male)	210	-0.47	0.78	1	0.37693
	Latency to fly	Test variable	First latency to foveate	210	340.20	17.32	1	0.00003
		Control variable	Event	210	205.62	2.66	1	0.10316
			Predator side (North)	210	-276.02	4.50	1	0.03396
			Sex (male)	210	-57.44	0.18	1	0.6706
4.2	Latency to run	Test variable	Trial	240	(a)	37.49	1	<0.00001
			Event	240	-260.26	1.37	1	0.24161
			Predator side (North)	240	-1243.29	30.25	1	<0.00001
			Sex (male)	240	247.14	0.35	1	0.55150
	Latency to fly	Test variable	Trial	240	(a)	40.00	1	<0.00001
			Event	240	20.48	0.03	1	0.86097
			Predator side (North)	240	-202.39	2.65	1	0.10364
			Sex (male)	240	83.62	0.26	1	0.60673

Repeatability analysis

999 We assessed whether the variables used in the models (both responses and predictors) exhibited
1000 significant inter-individual differences by evaluating their repeatability across trials and events. The
1001 rpt() function from the rptR package in R (Stoffel et al., 2017) was utilized for this analysis,
1002 incorporating the event, predator side, and the sex of the individual as control variables, and the trial

as an additional random effect in the repeatability model. Of the three response latencies examined, only the latency to escape demonstrated significant repeatability (R = 0.15, p < 0.0001), suggesting that certain individuals consistently escaped faster than others. In examining the response delay, we specifically analyzed the response time delay for flights (latency to fly - latency to foveate), which was found to be significantly repeatable. However, neither the latency to foveate nor the latency to fly were repeatable on their own. Due to the observation that in many trials, some pigeons did not fly up during the predator presentation, we also investigated whether the decision to fly was repeatable. Our analysis revealed that certain pigeons were more prone to flying than others (R = 0.22, p < 0.0001). As for the predictors tested, nearly all were found to be significantly repeatable (R ranging from 0.06 to 0.17, p values all below 0.0362), except for the distance from the monitor (p = 1). Detailed results of this analysis are presented in Table S8.

Table S8: Results of all repeatability analyses including the observed variable, the repeatability estimate and the p-value.

Variable	Repeatability estimate	p-value
Latency to foveate (log-transformed)	0.011	0.359
Latency to escape	0.145	<0.00001
Latency to fly	0.0525	0.062
Response time delay between foveation and flight	0.128	0.0404
Flew or not	0.219	<0.00001
Proportion of time head-up	0.166	<0.00001
Saccade rate	0.126	0.00014
Pecking rate (log-transformed)	0.0919	0.00507
Proportion foveate on the predator cue	0.144	0.00001
Proportion foveate on conspecifics	0.132	0.00007
Distance from the monitor	<0.0001	1
Distance from nearest neighbor	0.0644	0.0362

Reference in Supplemental Material

Gamer, M., Lemon, J., & Puspendra Singh, I. F. (2019). *irr: Various Coefficients of Interrater Reliability and Agreement* [Computer software]. https://CRAN.R-project.org/package=irr

Jones, K. A., & Godin, J.-G. J. (2010). Are fast explorers slow reactors? Linking personality type and anti-predator behaviour. *Proceedings of the Royal Society B: Biological Sciences*, *277*(1681), 625–632. https://doi.org/10.1098/rspb.2009.1607

Landis, J. R., & Koch, G. G. (1977). The Measurement of Observer Agreement for Categorical Data. *Biometrics*, *33*(1), 159–174. https://doi.org/10.2307/2529310

R Core Team. (2022). *R: A Language and Environment for Statistical Computing* [Computer software]. R Foundation for Statistical Computing. https://www.R-project.org/

1028   Stoffel, M. A., Nakagawa, S., & Schielzeth, H. (2017). rptR: Repeatability estimation and variance

1029   decomposition by generalized linear mixed-effects models. *Methods in Ecology and*

1030   *Evolution*, *8*(11), 1639–1644. https://doi.org/10.1111/2041-210X.12797

1031

1032