PI3K/HSCB axis facilitates FOG1 nuclear translocation to promote erythropoiesis and megakaryopoiesis

Gang Liu[a,1,*], Yunxuan Hou[a,1], Xin Jin[a], Yixue Zhang[a], Chaoyue Sun[a], Chengquan Huang[a], Yujie Ren[a], Jianmin Gao[c], Xiuli Wang[b] and Xiumei Jiang[c,*]

[a]Key Laboratory of Molecular Epigenetics of the Ministry of Education, Northeast Normal University, Changchun 130024, People's Republic of China

[b]School of Life Science, Northeast Normal University, Changchun 130024, People's Republic of China

[c]School of Chemistry, Northeast Normal University, Changchun 130024, People's Republic of China

*Corresponding authors: Gang Liu and Xiumei Jiang

Emails: liug959@nenu.edu.cn and jiangxm161@nenu.edu.cn

Condensed title: HSCB is key to erythropoiesis and megakaryopoiesis

Keywords: HSCB, FOG1 nuclear translocation, PI3K, erythropoiesis, megakaryopoiesis

---

[1] These authors contributed equally to this study.

Abstract

Erythropoiesis and megakaryopoiesis are stringently regulated by signaling pathways. However,

the precise molecular mechanisms through which signaling pathways regulate key transcription

factors controlling erythropoiesis and megakaryopoiesis remain partially understood. Herein, we

identified heat shock cognate B (HSCB), which is well known for its iron-sulfur cluster delivery

function, as an indispensable protein for friend of GATA 1 (FOG1) nuclear translocation during

erythropoiesis of K562 human erythroleukemia cells and cord-blood-derived human

CD34+CD90+ hematopoietic stem cells (HSCs), as well as during megakaryopoiesis of the

CD34+CD90+ HSCs. Mechanistically, HSCB could be phosphorylated by phosphoinositol-3-

kinase (PI3K) to bind with and mediate the proteasomal degradation of transforming acidic coiled-

coil containing protein 3 (TACC3), which otherwise detained FOG1 in the cytoplasm, thereby

facilitating FOG1 nuclear translocation. Given that PI3K is activated during both erythropoiesis

and megakaryopoiesis, and that FOG1 a key transcription factor for these processes, our findings

elucidate an important, previously unrecognized iron-sulfur cluster delivery independent function

of HSCB in erythropoiesis and megakaryopoiesis.

Introduction

Erythropoiesis and megakaryopoiesis respectively refer to the processes through which hematopoietic stem cells (HSCs) differentiate into mature red blood cells (RBCs) and megakaryocytes/platelets. Given the physiological significance of RBCs, megakaryocytes and platelets, it is not surprising that erythropoiesis and megakaryopoiesis are tightly regulated by a myriad of molecular mechanisms. Erythropoietin (EPO) is a fundamental cytokine promoting erythropoiesis via binding with erythropoietin receptor (EPOR) (Kuhrt and Wojchowski, 2015). Converging evidence indicates that the EPO/EPOR signaling is crucial for the proliferation and survival of colony-forming unit erythroid cells (CFU-Es) in vitro (Nandakumar et al., 2016). Although the EPO/EPOR signaling was dispensable for in vivo production of early committed erythroid progenitors, $Epo^{-/-}$ and $EpoR^{-/-}$ mice exhibited embryonic death due to deficient definitive erythropoiesis (Wu et al., 1995). These studies highlight the indispensability of EPO/EPOR signaling during erythropoiesis. Thrombopoietin (TPO) and its receptor MPL constitute the primary regulator of megakaryopoiesis (Hitchcock et al., 2021). Studies indicated that TPO alone could induce the production of megakaryocyte colony-forming units (CFU-MKs), while addition of other cytokines including stem cell factor (SCF), interleukin 3 (IL3) and IL11 markedly increased the size, but not the number, of these colonies (Kaushansky et al., 1995; Kaushansky et al., 1994). The biological significance of TPO/MPL signaling pathway was also evidenced by the development of thrombocytopenia in Mpl-deficient mice as well as in human subjects carrying *MPL* and *THPO* mutations (Gurney et al., 1994; Noris and Pecci, 2017). Both EPOR and MPL belong to the type-I family of cytokine receptors. They do not possess kinase activity *per se*, but can transphosphorylate Janus tyrosine kinase 2 (JAK2), which in turn activates several downstream signaling cascades, including the JAK2/signal transducers and activators of transcription (STAT), phosphoinositol-3-kinase (PI3K)/AKT and rat sarcoma (RAS)/RAF/mitogen-activated protein kinase (MAPK) pathways (Eggold and Rankin, 2019; Hitchcock et al., 2021), all of which contribute more or less to both erythropoiesis and megakaryopoiesis.

The essence of differentiation is the generation of cells with specific functions, which is realized by the differentiation stage-dependent transcriptional activation and suppression of innumerable genes. This is why there is a generally accepted notion that transcription factors (TFs) are the main drivers of differentiation (Lambert et al., 2018). Each of the differentiation processes of erythropoiesis and megakaryopoiesis is controlled by a combination of key TFs, including GATA1, friend of GATA1 (FOG1), KLF1, TAL1, MEIS1, etc., for the erythroid lineage (Dore and Crispino, 2011; Kwon et al., 2021), and GATA1, FOG1, TAL1, MEIS1, FLI1, etc., for the megakaryocytic lineage (Dore and Crispino, 2011; Kwon et al., 2021). Since cell signaling is the dominant way by which erythroid and megakaryocytic progenitors (or their common progenitors) sense

3

differentiation signals, it is crucial to understand how these key erythroid and megakaryocytic TFs

are regulated by signaling pathways, especially those downstream of EPO/EPOR and TPO/MPL

signaling. However, to date, research on this topic remains scarce. A study deepened our

understanding about this issue by demonstrating that the PI3K-AKT pathway can phosphorylate

GATA1 at serine 310 in erythroid cells, thereby augmenting its transcription activity (Zhao et al.,

2006). However, given that the proper functioning of GATA1 in regulating target gene expression

relies on its binding with a specific cofactor (Gutierrez et al., 2020), how EPO/EPOR and

TPO/MPL signaling pathways regulate the activity of GATA1 binding partners, such as FOG1,

warrants further investigation.

Heat shock cognate B (HSCB or HSC20) is widely known for its function during the delivery of

nascent iron-sulfur clusters (ISCs) to recipient proteins (Maio et al., 2014). The biogenesis of

ISCs starts with the assembly of nascent ISCs on a multi-protein complex composed of iron-sulfur

cluster assembly enzyme (ISCU), nitrogen fixation 1 homolog (NFS1), LYR motif containing 4

(LYRM4 or ISD11), and acyl carrier protein (ACP) (Maio et al., 2020). After the ISCs are

synthesized, ISCU binds with an ISC delivery chaperone/cochaperone complex comprising

HSCB and HSPA9 to transfer the nascent ISCs to target Fe-S proteins or secondary ISC carriers

(Maio et al., 2020). Recently, Crispin et al. described a female patient who suffered from

congenital sideroblastic anemia (CSA) due to heterozygous variations that resulted in decreased

expression of HSCB (Crispin et al., 2020). In addition to the anemic phenotype, the proband also

developed moderate thrombocytopenia and mild neutropenia. Consistent with the proband's

clinical presentations, *Hscb*[fl/–] mice with a *Vav1-Cre* transgene also exhibited reduced RBC and

platelet counts (Crispin et al., 2020). Based on these findings, we speculated that HSCB may be

implicated with hematopoiesis, especially erythropoiesis and megakaryopoiesis. To test this

hypothesis, herein we explored the function of HSCB during erythropoiesis and

megakaryopoiesis. Our findings indicated that HSCB can be phosphorylated by PI3K to promote

FOG1 nuclear translocation during human erythropoiesis and megakaryopoiesis. This study

describes a previously unrecognized iron-sulfur cluster delivery independent function of HSCB,

thereby providing new insights into human erythropoiesis and megakaryopoiesis.

Results

HSCB exerted an ISC delivery independent function during erythropoiesis of HSCs

First, we investigated if HSCB is implicated in erythropoiesis by inhibiting its expression in

CD34+CD90+ HSCs via a mixture of shRNAs. We also knocked down the expression of human

4

ISCU and *NFS1* genes in these cells to determine the impact of ISC biogenesis deficiency on

erythropoiesis. The high-purity CD34+CD90+ HSC population was first obtained (supplemental

Figures 1 and 2A) and divided into five groups of Control, shNC, shHSCB, shISCU and shNFS1.

HSCs in the Control group were untransfected, whereas those in the other four groups were

transfected with the corresponding shRNA plasmids. The HSCs in these groups induced for

erythropoiesis for 17 days to yield erythrocytes at day 17 of differentiation (Ery 17 days) and

collected for analyses (supplemental Figure 1). Figure 1A demonstrates the efficiency of shRNA

knockdown maintained throughout the experimental period. The activities of aconitase and

pyruvate dehydrogenase (PDH), which depend on intact ISC biogenesis (Cameron et al., 2011;

Castro et al., 2019), decreased significantly in the shHSCB, shISCU and shNFS1 groups

compared with the shNC group (all P values were less than 0.05 or 0.01 in Student's t test, Figure

1B-C), demonstrating that ISC biogenesis was strongly inhibited in these groups. Since heme

biosynthesis is regulated by ISC biogenesis at several nodes (Liu et al., 2020), we further

explored whether heme biosynthesis was also inhibited in the shHSCB, shISCU and shNFS1

groups. Indeed, these three groups exhibited significantly lower heme contents (all P values were

less than 0.05 or 0.01 in Student's t test, Figure 1D) and paler cell pellets (supplemental Figure

2B) than the shNC group. Surprisingly, among the groups subjected to shRNA plasmids

transfection, only the shHSCB group displayed significantly inhibited erythropoiesis, as evidenced

by the reduced proportion of CD71+CD235a+ progenitors (P<0.001, Student's t test, Figure 1E–F)

and the decreased mRNA expression levels of erythroid marker genes *HBB*, *ALAS2* and *GYPC*

(all P values were less than 0.01 or 0.001 in Student's t test, Figure 1G). We then purified the

CD71-CD235a- subpopulation from the shHSCB group of Ery 17 days via cell sorting and

characterized the differential status of these cells. As shown in Figure 1H, the majority of CD71-

CD235a- cells from the shHSCB group were CD34+CD38+CD45RA-IL3RA-, exhibiting a surface

marker expression pattern (CD34+CD38+CD45RA-IL3RA-CD71-CD235a-) resembling that

observed on CD34+CD90+ HSCs induced for erythropoiesis for 2 days (E 2-day HSCs)

(supplemental Figure 2C and Figure 5C). These findings revealed that HSCB knockdown

detained the erythropoiesis of HSCs at an early stage and that HSCB might be involved in key

events regulating early erythropoiesis at this specific stage. Therefore, E 2-day HSCs were

characterized in several following experiments to explore the molecular mechanisms through

which HSCB deficiency impeded HSCs erythropoiesis. Altogether, these data indicated that

HSCB knockdown impaired both heme biosynthesis and erythropoiesis in erythroid progenitors,

while knockdown of ISC assembly proteins markedly reduced heme production but did not

significantly affect erythropoiesis, suggesting an ISC delivery independent function of HSCB

during early erythropoiesis of CD34+CD90+ HSCs.

HSCB deficiency inhibited the erythropoiesis of K562 cells

K562 erythroleukemia cells are widely employed to study the molecular events implicated with

human erythropoiesis (Nandakumar et al., 2016). To verify the function of HSCB in erythropoiesis

and explore the underlying molecular mechanisms, we established *HSCB*-knockout K562 cells

(CRISPR HSCB) and induced erythroid differentiation of these cells with EPO (EPO+CRISPR

HSCB). After FBS deprivation treatment (see the Materials and Methods section for details),

untransfected K562 cells without EPO treatment, those only treated with EPO and those first

transfected with the negative control CRISPR plasmid and then treated with EPO respectively

constituted the Untreated, EPO and EPO+CRISPR NC groups. As revealed by Figure 2A–C,

compared with the Untreated group, the EPO and EPO+CRISPR NC groups exhibited

significantly higher levels of erythrocyte membrane proteins GYPA and SPTA1 (all P values were

less than 0.05 or 0.01 in Student's t test), significantly increased heme contents (all P values

were less than 0.01 or 0.001 in Student's t test) and elevated mRNA expression levels of

erythroid-specific genes *HBB*, *ALAS2* and *GYPC,* indicating that the EPO treatment successfully

induced the erythropoiesis of K562 cells. In the EPO+CRISPR HSCB group, however, the protein

levels of GYPA and SPTA1, heme content and mRNA levels of *HBB*, *ALAS2* and *GYPC* all

decreased markedly compared with those of the EPO+CRISPR NC group (all P values were less

than 0.01 or 0.001 in Student's t test), demonstrating that HSCB knockout hindered the

erythropoiesis of K562 cells induced by EPO. Erythropoiesis is underpinned by transcriptional

changes mainly driven by nuclear proteins (Dore and Crispino, 2011; Nandakumar et al., 2016).

To explore why HSCB silencing inhibited the erythropoiesis of K562 cells induced by EPO, we

compared the nuclear proteomic profiles of the EPO+CRISPR NC and EPO+CRISPR HSCB

groups via performing tandem mass tag-based mass spectrometry (TMT-MS) on the nuclear

fractions. As shown in Figure 2D, 9 downregulated and 2 upregulated differentially-expressed

nuclear proteins (DENPs, see the Materials and Methods section for the definition) were identified

in the EPO+CRISPR HSCB group. Among these 11 DENPs, only NAA30, LYAR and FOG1 were

bona fide nuclear proteins (Figure 2E). We further tested whether these 3 DENPs were indeed

differentially expressed in the nuclear fractions of K562 cells and E 2-day HSCs after HSCB

silencing. Western blotting (WB) analyses revealed that FOG1 was the only nuclear protein

expressed at significantly different levels in both K562 cells and E 2-day HSCs upon HSCB

deficiency (P values were less than 0.05 or 0.01 in Student's t test, Figure 2F–G). We also tested

the nuclear abundance of GATA1 in these cells but observed no significant changes (Figure 2F–

G). These data confirmed that the nuclear abundance of FOG1 was decreased in HSCB-deficient

EPO-induced K562 cells and E 2-day HSCs.

HSCB deficiency impaired the erythropoiesis of K562 cells and HSCs by prohibiting FOG1

6

nuclear translocation

Given the importance of FOG1 during human erythropoiesis, we determined whether the

decreased nuclear abundance of FOG1 was responsible for the inhibited erythropoiesis in HSCB-

deficient EPO-induced K562 cells and CD34+CD90+ HSCs. We overexpressed human FOG1

protein in the EPO+CRISPR HSCB K562 cells and the shHSCB CD34+CD90+ HSCs to

respectively establish the EPO+CRISPR HSCB+FOG1 OE and shHSCB+FOG1 OE groups. As

shown in Figure 3A–E, FOG1 overexpression counteracted the inhibitory effects of HSCB

silencing on the erythropoiesis of K562 cells and CD34+CD90+ HSCs because, except for the

heme contents, the EPO+CRISPR HSCB+FOG1 OE and shHSCB+FOG1 OE groups exhibited

comparable levels of all the parameters evaluated relative to their corresponding control groups.

Interestingly, the total protein level of GATA1 was not significantly affected by HSCB deficiency or

FOG1 overexpression in EPO-induced K562 cells and CD34+CD90+ HSCs (Figure 3A). These

data verified that HSCB deficiency hindered the erythropoiesis of K562 cells and CD34+CD90+

HSCs by reducing the nuclear abundance of FOG1. To gain more insights into the molecular

mechanisms by which HSCB deficiency decreased FOG1 nuclear abundance, we also measured

cytoplasmic FOG1 protein levels in EPO+CRISPR HSCB K562 cells and the shHSCB group of E

2-day HSCs. We observed that these cells expressed significantly higher cytoplasmic levels of

FOG1 than their corresponding controls (all P values were less than 0.01 or 0.001 in Student's t

test, Figure 3F), indicating that HSCB deficiency prohibited FOG1 nuclear translocation. These

findings were further confirmed by immunofluorescence assays on K562 cells, in which the red

fluorescence of FOG1 in the EPO+CRISPR NC group was mostly observed in the nuclei and that

in the EPO+CRISPR HSCB group was mainly found in the cytoplasm (Figure 3G). Collectively,

these observations indicated that HSCB deficiency impaired the erythropoiesis of K562 cells and

HSCs by prohibiting FOG1 nuclear translocation.

HSCB deficiency prohibited FOG1 nuclear translocation by enhancing its interaction with

TACC3

Nuclear translocation of a specific protein is ultimately regulated by protein-protein interactions

(Cautain et al., 2015; Wing et al., 2022). Therefore, to further explain how HSCB deficiency

prohibited FOG1 nuclear translocation, we next sought to identify differential binding proteins

(DBPs, see the Materials and Methods section for the definition) through affinity purification-MS

analyses. We identified two DBPs, namely MYO1E and TACC3 (Figure 4A). Their binding

affinities with FOG1 increased significantly after HSCB expression was silenced (P values were

less than 0.05 in Student's t test,). We focused only on TACC3 because MYO1E did not bind with

FOG1 in either the shNC or shHSCB group of E 2-day HSCs (supplemental Figure 3). We first
performed co-immunoprecipitation (co-IP) experiments to test whether FOG1 indeed exhibited
increased binding affinities with TACC3 in EPO-induced K562 cells and E 2-day HSCs after
HSCB silencing. As shown in Figure 4B–C, the interaction between FOG1 and TACC3 was only
detectable in HSCB-deficient cells, confirming that HSCB deficiency promoted the interaction
between FOG1 and TACC3. KHS101, a TACC3 inhibitor, can both reduce the activity of TACC3
and decrease its protein abundance (Polson et al., 2018). Therefore, we tested whether KHS101
treatment could decrease the protein level of TACC3 in these cells and thereby alleviate the
inhibitory effect of HSCB deficiency on their erythropoiesis. Indeed, as shown in Figure 4D–H,
KHS101 treatment enhanced FOG1 nuclear translocation [as reflected by the significantly
increased relative nuclear FOG1 (N-FOG1) to cytoplasmic FOG1 (C-FOG1) ratios in the KHS101
treatment groups (P values were less than 0.05 or 0.01 in Student's t test,)] and restored
erythropoiesis in EPO+CRISPR HSCB K562 cells and shHSCB HSCs. Altogether, these data
exhibited that HSCB deficiency prohibited FOG1 nuclear translocation by promoting its interaction
with TACC3, and that reducing TACC3 abundance by KHS101 treatment alleviated the inhibitory
effect of HSCB deficiency on the erythropoiesis of K562 cells and CD34+CD90+ HSCs.

HSCB facilitated FOG1 nuclear translocation by binding with and mediating the
proteasomal degradation of TACC3 upon activation of the EPO/EPOR signaling

To further understand how HSCB facilitated FOG1 nuclear translocation and why HSCB
deficiency enhanced the protein level of TACC3, we transfected an HSCB-overexpressing
plasmid into K562 cells and treated the transfected K562 cells with EPO to establish an
EPO+HSCB OE group. As shown in Figure 5A, EPO treatment significantly reduced the protein
level of TACC3 and promoted FOG1 nuclear translocation (all P values were less than 0.05 or
0.01 in Student's t test). HSCB overexpression further decreased the protein level of TACC3 and
enhanced FOG1 nuclear abundance. However, after the EPO+HSCB OE K562 cells were treated
with MG132, a proteasome inhibitor, the protein level of TACC3 and the nuclear abundance of
FOG1 returned to those in untreated K562 cells. Co-IP assays revealed that TACC3 interacted
with HSCB when the K562 cells were treated with EPO, whereas it interacted with FOG1 in the
untreated and EPO+HSCB OE+MG132 K562 cells (Figure 5B). These data indicated that, upon
EPO treatment, HSCB could bind with and mediate the proteasomal degradation of TACC3,
thereby preventing TACC3 from binding with FOG1 in K562 cells. We further characterized the
protein expression profiles of HSCB, cytoplasmic/nuclear FOG1 and TACC3, as well as their
physical interactions, at different stages during erythropoiesis of CD34+CD90+ HSCs. The cells
were induced for erythropoiesis for 0, 2, 4, 10, 17 and 24 days and harvested for analyses (Figure
5C). As shown in Figure 5D–F, the nuclear abundance of FOG1 and the interaction between

HSCB and TACC3 all gradually increased from day 0 to day 4, resembling the changing trend of
the number of EPOR+ cells. These findings were consistent with those observed in K562 cells
and further highlighted that the EPO/EPOR signaling was indispensable for HSCB to bind with
and mediate the proteasomal degradation of TACC3 to facilitate FOG1 nuclear translocation
(hereinafter defined as HSCB functionalization).

Phosphorylation of HSCB by PI3K was necessary for its functionalization during human
erythropoiesis

To determine which downstream cascade of EPO/EPOR signaling was important for HSCB
functionalization, we treated K562 cells with EPO plus ruxolitinib, wortmannin and lonafarnib to
respectively inhibit the activities of JAK2, PI3K and RAS upon activation of the EPO/EPOR
signaling. As shown in supplemental Figure 4, treatments with ruxolitinib, wortmannin and
lonafarnib respectively decreased the phosphorylation levels of STAT5, AKT and MEK1/2 in
EPO-induced K562 cells, confirming that these treatments were effective. Among the inhibitors,
ruxolitinib and wortmannin significantly ameliorated the protein level of TACC3 and diminished
the nuclear translocation of FOG1 (all P values were less than 0.05 or 0.001 in Student's t test,
Figure 6A), suggesting that JAK2 and PI3K activities are both necessary for HSCB
functionalization in EPO-induced K562 cells. Since PI3K functions downstream of JAK2 in the
EPO/EPOR signaling, we then focused only on PI3K to investigate its role in HSCB
functionalization. Consistent with the WB findings, our co-IP assays demonstrated that inhibition
of PI3K activity by wortmannin abolished the interaction between HSCB and TACC3 and thereby
allowed TACC3 to bind with FOG1 in EPO-induced K562 cells and E 2-day HSCs (Figure 6B).
Further analysis of protein phosphorylation status via co-IP assays revealed that EPO treatment
enhanced the overall phosphorylation level of cytosolic HSCB, but not that of TACC3, in K562
cells (Figure 6C). Additionally, wortmannin treatment reduced the overall phosphorylation level of
cytosolic HSCB, but not that of TACC3, in EPO-induced K562 cells and E 2-day HSCs (Figure
6C). We then identified the physical interaction between cytosolic HSCB and PIK3R1, which is
the P85-alpha subunit of PI3K, via co-IP experiments (Figure 6D). These data indicated that
HSCB phosphorylation by PI3K was indispensable for its functionalization by the EPO/EPOR
signaling. In line with these findings, wortmannin treatment significantly inhibited the
erythropoiesis of both K562 cells and CD34+CD90+ HSCs (all P values were less than 0.01 or
0.001 in Student's t test, Figure 6E–I). Collectively, our results clarified that phosphorylation of
HSCB by PI3K was necessary for its functionalization during human erythropoiesis.

Activation of the PI3K-HSCB axis was also necessary for the megakaryopoiesis of
CD34+CD90+ HSCs

Given that PI3K is also activated in the TPO/MPL signaling and that FOG1 is also crucial for

megakaryopoiesis, we finally tested whether the PI3K-HSCB signaling axis is necessary for the

megakaryopoiesis of CD34+CD90+ HSCs. As displayed in Figure 7A–C, the CD34+CD90+ HSCs

transfected with the shHSCB plasmid (shHSCB group) and those treated with wortmannin

(Wortmannin group) exhibited significantly reduced proportions of CD41a+CD42b+ cells and

lower mRNA expression levels of megakaryocyte-specific genes *VWF* and *GP1BA* (all P values

were less than 0.001 in Student's t test), while the megakaryopoiesis of CD34+CD90+ HSCs

transfected with the shHSCB plasmid and treated with KHS101 (shHSCB+KHS101) was

undisrupted. Since most cells from the shHSCB group were CD34+CD38+CD45RA-IL3RA-

(Figure 7D), an expression pattern most similar to that observed on CD34+CD90+ HSCs induced

for megakaryopoiesis for 3 days (MK 3-day HSCs) (supplemental Figure 5), MK 3-day HSCs

were utilized for the following analyses. As exhibited in Figure 7E, the shHSCB and Wortmannin

groups of MK 3-day HSCs had significantly higher levels of TACC3 and decreased relative N-

FOG1/C-FOG1 ratios (all P values were less than 0.05 in Student's t test,), while the

shHSCB+KHS101 group possessed a lower level of TACC3 and a slightly increased level of

FOG1 nuclear translocation. Additionally, wortmannin treatment reduced the overall

phosphorylation level of HSCB and decreased its interaction with TACC3 (Figure 7F). We also

determined the nuclear abundance of FOG1 on days 0, 3, 7 and 14 during CD34+CD90+ HSCs

megakaryopoiesis and found that the level of N-FOG1 peaked on day 3 and declined thereafter

(Figure 7G). Taken together, these data were consistent with those observed in erythroid

progenitors, indicating that the PI3K-HSCB signaling axis was also important for the normal

megakaryopoiesis of CD34+CD90+ HSCs.

323　**Discussion**

324
FOG1 is a binding partner of GATA1 exerting a vital function during erythropoiesis and

megakaryopoiesis, as FOG1-deficient mice displayed megakaryopoiesis failure and impaired

erythropoiesis (Tsang et al., 1998). A previous study suggested that FOG1 was needed for the

formation of megakaryocyte/erythroid progenitors (MEPs), while GATA1 was required for the

development of committed erythroid progenitors (Mancini et al., 2012). Given the biological

significance of FOG1, it is crucial to understand how FOG1 is regulated by erythroid and

megakaryocytic signals to ensure its timely activation or expression during erythropoiesis and

megakaryopoiesis. Herein, we have established a PI3K-HSCB signaling axis that facilitated

FOG1 nuclear translocation upon activation of the EPO/EPOR and TPO/MPL signaling pathways

during human erythropoiesis and megakaryopoiesis (Figure 7H).

Although heme biosynthesis depends on an intact ISC assembly machinery, deficiencies in

proteins of the ISC assembly complex, namely ISCU, NFS1, and LYRM4 have not been

associated with erythropoietic consequences in both human subjects and cell models (Ye and

Rouault, 2010). In contrast, however, deficiencies in proteins responsible for ISC transfer, such

as GLRX5, HSPA9 and HSCB, have been tightly linked with sideroblastic anemia phenotypes

(Ducamp and Fleming, 2019). This could be due to that ISC biogenesis deficiencies are tolerable

in erythrocytes, while the proteins responsible for ISC transfer possess moonlighting functions

during erythropoiesis. Our research was initially inspired by the findings that a CSA proband

carrying heterozygous HSCB mutations developed moderate thrombocytopenia and mild

neutropenia (Crispin et al., 2020). It is not surprising that mutations in proteins responsible for ISC

transfer can cause anemia because two of the eight enzymes catalyzing heme biosynthesis,

namely aminolevulinate dehydratase and ferrochelatase, were identified to be iron-sulfur proteins

(Dailey et al., 1994; Liu et al., 2020). However, it was not clear why the HSCB-deficient proband

also exhibited moderately reduced numbers of platelets. Therefore, we tested whether HSCB is

implicated with erythropoiesis and megakaryopoiesis of human CD34+CD90+ HSCs. Our results

indicated that HSCB knockdown detained the erythropoiesis and megakaryopoiesis of HSCs at

early stages because most of the resultant progenitors were CD34+CD38+CD45RA-IL3RA-

CD71-CD235a- or CD34+CD38+CD45RA-IL3RA-CD41a-CD42b-, corresponding to common

myeloid progenitors or MEPs as per previous studies (Edvardsson et al., 2006; Kwon et al., 2021).

Further analysis of the underlying mechanisms revealed that HSCB deficiency mainly impaired

FOG1 nuclear translocation. The only known function of HSCB is to deliver nascent ISCs to

recipient proteins. Interestingly, knockdown of proteins responsible for ISC assembly did not

significantly affect erythropoiesis of HSCs, thus highlighting an ISC delivery independent function

of HSCB during erythropoiesis and megakaryopoiesis. HSCB is a chaperone expressed in both

mitochondria and cytosol. In this study, we were able to detect two bands for HSCB with western

blotting, among which the upper and lower bands respectively represent cytosolic and

mitochondrial HSCB (Liu et al. 2020). As revealed by Figure 5D and Figure 6A, the sub-cellular

distribution of HSCB was neither significantly altered during erythropoiesis of CD34+CD90+

HSCs nor significantly affected by ruxolitinib and wortmannin treatments in K562 cells. Therefore,

it seems that the sub-cellular localization of HSCB does not change during erythropoiesis or

correlate with its phosphorylation status. However, whether the phosphorylation status of HSCB

affects its ISC delivery function warrants further investigation. Additionally, since HSCB can bind

with a wide array of intracellular proteins (Maio et al., 2017), we speculate that the protein may be

involved in many biological processes other than nascent ISC delivery. In this context, we hope

our study may inspire more research efforts on ISC delivery independent functions of HSCB.

TACC3 belongs to a family of TACC-domain-containing proteins associated with mitotic spindles

and centrosomes (Ding et al., 2017). The protein participates in cell division by regulating the

organization of spindle and microtubule nucleation during mitosis (Ding et al., 2017). Recently,

TACC3 has attracted increasing research interest because it was found aberrantly expressed in

cancers (Ding et al., 2017; Saatci et al., 2023). In addition to its association with tumorigenesis,

TACC3 has also been implicated with erythropoiesis. According to a previous study, TACC3

could inhibit FOG1 nuclear translocation to suppress the erythropoiesis of mouse MEL and G1ER

cells (Garriga-Canut and Orkin, 2004). Herein we found that TACC3 exerted a similar function in

human erythroid and megakaryocytic progenitor cells because, in the absence of HSCB, the

protein could bind with and prevent the nuclear translocation of FOG1. Then we observed that

treatment with KHS101, a small-molecule inhibitor that could decrease the activity and protein

abundance of TACC3, alleviated the inhibitory effect of TACC3 on erythropoiesis upon HSCB

deficiency. These data clarified the mechanism through which HSCB deficiency inhibited FOG1

nuclear translocation and consequently impaired erythropoiesis and megakaryopoiesis. In sight of

the functions of TACC3 during cell division and erythropoiesis/megakaryopoiesis, it is possible

that TACC3 plays a crucial role in maintaining the balance between cell proliferation and

differentiation, which of course warrants further investigation. Furthermore, given that KHS101

could decrease TACC3 protein level to facilitate FOG1 nuclear translocation, this small-molecule

reagent may have the potential to treat diseases caused by reduced activity or protein level of

FOG1.

The precise mechanisms through which EPO/EPOR and TPO/MPL signaling pathways regulate

lineage-specific transcription events during erythropoiesis and megakaryopoiesis remain

incompletely understood. In a previous research, expression of a constitutively activated Stat5a

mutant in Jak2$^{-/-}$ and EpoR$^{-/-}$ mouse fetal liver cells was found to rescue their proliferation

defects independently of EPO (Grebien et al., 2008). Additionally, the authors also observed that

transplantation of Jak2$^{-/-}$ fetal liver cells expressing the Stat5a mutant into irradiated mice was

able to maintain erythropoiesis and myelopoiesis of the mice (Grebien et al., 2008). These data

verified the important function of Stat5a as a downstream effector of the Epo/EpoR signaling

pathway during murine erythropoiesis and myelopoiesis. However, since STAT5 possesses a

variety of target genes, why the protein is specifically required for erythropoiesis and

myelopoiesis warrants further investigation. Another early study demonstrated that wortmannin

could impede EPO-induced K562 cell erythropoiesis but did not inhibit erythropoiesis of the cells

triggered by hemin (Kubota et al., 1996). Since wortmannin is an inhibitor for PI3K, this study

revealed that PI3K plays a large part in EPO-induced erythropoiesis of K562 cells. Later, it was

demonstrated that the PI3K-AKT signaling cascade could directly activate GATA1 to stimulate

erythropoiesis (Zhao et al., 2006). In this study, we confirmed the importance of PI3K activity

during erythropoiesis and megakaryopoiesis and established a novel molecular mechanism. Our

data indicated that PI3K could bind with and phosphorylate HSCB to promote its interaction with

TACC3 to induce TACC3 degradation, thereby facilitating FOG1 nuclear translocation during

erythropoiesis of K562 human erythroleukemia cells and HSCs, as well as during

megakaryopoiesis of HSCs. Interestingly, the nuclear and cytoplasmic GATA1 expression levels

were not regulated by this PI3K-HSCB axis, as revealed by the data presented in Figure 2F and

Figure 3A, as well as the TMT-MS counts provided in Supplemental Dataset 5. These new

findings explain how EPO/EPOR and TPO/MPL signaling pathways simultaneously activate

GATA1 and FOG1 to ensure that this duo can function in a concerted and timely manner during

human erythropoiesis and megakaryopoiesis. In addition to the EPO/EPOR and TPO/MPL

signaling pathways, PI3K is also activated by the SCF-c-KIT pathway (Lennartsson and

Ronnstrand, 2012). Since the c-KIT and MRL receptors are expressed early on HSCs

(Lennartsson and Ronnstrand, 2012), it is conceivable that the timely activation of GATA1 and

FOG1 during erythropoiesis and megakaryopoiesis also requires the participation of other

differentiation stage-specific mechanisms regulating the abundance of these proteins. This idea

can be supported by our findings on HSCs induced for erythropoiesis for different periods of time.

As shown in Figure 5D, while the levels of cytoplasmic TACC3 and FOG1 exhibited no significant

changes from Day 2 to Day 4 of erythropoiesis, the level of N-FOG1 increased by more than

three folds during the same period, suggesting that the functionalization of FOG1 is determined

not only by the activation of the PI3K-HSCB axis, but also by the overall expression level of this

protein. Additionally, although we failed to detect nuclear FOG1 in uninduced CD34+CD90+

HSCs (Figure 5D), we did detect a faint band of FOG1 in the cytoplasmic fraction. Given that

HSCB was already expressed in uninduced CD34+CD90+ HSCs, why the SCF-c-KIT-PI3K

pathway failed to promote FOG1 nuclear translocation in these cells warrants further investigation.

This study has some limitations that should be acknowledged. First, we utilized the K562 cell line

as one of the cell models in this study because it has the potential to differentiate into erythroid,

megakaryocyte and macrophage lineages and thus may represent a useful tool in studying the

molecular mechanisms underlying early-stage erythropoiesis and

megakaryopoiesis(Nandakumar et al., 2016; Sutherland et al., 1986). However, although

K562 cells have been widely utilized to study erythropoiesis (Andersson et al., 1979; Cao et al.,

2020), and the capability of EPO to induce K562 erythropoiesis has been demonstrated by this

study and previous research (Hoffman et al., 1979; Kubota et al., 1996), this human

erythroleukemia cell line might not faithfully represent the normal physiological processes of

erythropoiesis. Fortunately, all our major conclusions can be successfully recapitulated in primary

cells. A second limitation is that we did not focus on LYAR in this study because the difference in

nuclear abundance of this protein between the shNC and shHSCB groups of E 2-day HSCs did

not reach a statistically significant level (Figure 2F–G). However, given the established function of

LYAR during late erythropoiesis , it will be intriguing to explore whether this protein is also

implicated with HSCB functionalization during human erythropoiesis. Finally, since we did not

have access to biological samples from the patient harboring the *HSCB* mutation and that HSCB-

deficient mice had almost no erythroid cells in their bone marrow, we were not able to further

validate our findings in in vivo models.

In spite of these limitations, our study established a PI3K-HSCB axis that acted downstream of

the EPO/EPOR and TPO/MPL signaling pathways to induce proteasomal degradation of TACC3,

which otherwise detained FOG1 in the cytoplasm, in K562 cells and HSCs. This work elucidates

an essential iron-sulfur cluster delivery independent function of HSCB in translating differentiation

signals into downstream transcriptional events during human erythropoiesis and

megakaryopoiesis.

Materials and methods

Culturing and treatment of K562 cell line and HSCs

Human K562 erythroleukemia cells were purchased from Procell Life Science&Technology Co.,

Ltd. (China) and maintained in Iscove's modified Dulbecco medium (IMDM, PM150510, Procell,

China) supplemented with 10% fetal bovine serum (FBS, 085-150, Wisent Bioproducts, Canada),

100 U/mL penicillin and 100 μg/mL streptomycin (PB180120, Procell, China) in a 37°C, 5% CO_2

incubator. For induction of erythropoiesis, K562 cells were cultured in IMDM without FBS (FBS

deprivation treatment) for 24 hours and then treated with 5 U/mL human recombinant EPO

(GP20193, GlpBio, USA) for 48 hours. After the FBS deprivation treatment, K562 cells were

incubated for 48 hours with 300 nM ruxolitinib (GC14191), 200 nM wortmannin (GC12338), 2 μM

lonafarnib (GC10330) dissolved in dimethylsulfoxide (DMSO) and 10 μM MG132 (GC10383, all

from GlpBio, USA) dissolved in phosphate buffered saline (PBS) for inhibiting the activities of

JAK2, PI3K, RAS and proteasome, respectively, and with only DMSO for the "vehicle" treatment.

To reduce the protein level of transforming acidic coiled-coil containing protein 3 (TACC3), K562

cells were treated with 5 μM KHS101 (GC13582, GlpBio, USA) dissolved in PBS.

Cord-blood-derived CD34+ human HSCs were originally obtained from healthy female donors

aged between 24-27 years and were commercially available at Precision BioMedicals Co., Ltd.,

China (903410). The CD34+ HSCs population was first expanded by incubating the cells in

StemSpan™ Serum-Free Expansion Medium (SFEM) II (09655) plus StemSpan™ CD34+

Expansion Supplement that contains recombinant human fms-like tyrosine kinase 3 ligand, SCF,

IL3, IL6 and TPO (02691, all from STEMCELL Technologies, Canada) for 7 days in a 37°C, 5%

CO_2 incubator. After that, CD34+CD90+ HSCs were purified by cell sorting (described below).

The HSCs were cultured for 2, 4, 10, 17 or 24 days in StemSpan™ SFEM II plus StemSpan™

Erythroid Expansion Supplement that contains recombinant human SCF, IL3 and EPO (02692,

STEMCELL Technologies, Canada) for induction of erythropoiesis, or for 3 or 14 days in

StemSpan™ SFEM II plus StemSpan™ Megakaryocyte Expansion Supplement that contains

SCF, IL-6, IL-9 and TPO (02696, STEMCELL Technologies, Canada) for induction of

megakaryopoiesis. The concentration of the HSCs or the subsequent erythroid/megakaryocytic

progenitors was maintained below 1×10^5 cells/mL during all the aforementioned experiments. The

levels of PI3K activity and TACC3 protein expression were reduced by culturing the cells with 100

nM wortmannin (GC12338) and 4 µM KHS101 (GC13582, all from GlpBio, USA), respectively.

Gene knockdown, overexpression and knockout

Plasmids carrying shRNAs against human *ISCU* (sc-270108-SH), *NFS1* (sc-75911-SH) and

HSCB (sc-75306-SH) genes, as well as that carrying a negative control (NC) shRNA (sc-108060),

were obtained from Santa Cruz Biotechnology, China. Plasmids overexpressing human FOG1

(pcDNA3.1-FOG1 OE), FLAG/MYC (F/M)-tagged FOG1 (pCMV-F/M-FOG1 OE) and human

HSCB (pcDNA3.1-HSCB OE), and the corresponding empty vectors were synthesized and

provided by Fenghui Biotechnology Co., Ltd., China. The HSCB CRISPR/Cas9 KO Plasmid (h)

(sc-409377), HSCB HDR Plasmid (h) (sc-409377-HDR) and Control CRISPR/Cas9 Plasmid (sc-

418922) were procured from Santa Cruz Biotechnology, China. These plasmids were transfected

into CD34+CD90+ HSCs by utilizing the RFect^SP Plasmid DNA Transfection Reagent (21027,

Biodai, China). After transfection with shRNA plasmids, the HSCs were first cultured in

StemSpan™ Serum-Free Expansion Medium (SFEM) II (09655, STEMCELL Technologies,

Canada) supplemented with 2 µg/mL puromycin (GC16384, GlpBio, USA) for 72 h and then

induced for erythropoiesis or megakaryopoiesis. Stable transfection of shRNA plasmids was

maintained by culturing the cells with 4 µg/mL puromycin. For FOG1 overexpression,

CD34+CD90+ HSCs were transfected with a pcDNA3.1-FOG1-OE plasmid, cultured in

StemSpan™ SFEM II for 48 h and then induced for erythropoiesis. During the erythropoiesis

process, the cells were transfected with the pcDNA3.1-FOG1-OE plasmid once every 72 hours

(see supplemental Figure 1 for a schematic diagram).

For FOG1, FLAG/MYC-FOG1 and HSCB overexpression in K562 cells, the cells were transfected

with the corresponding plasmids, cultured for 48 hours, and harvested for analyses. For silencing

of HSC expression, K562 cells were co-transfected with an HSCB CRISPR/Cas9 KO Plasmid

15

512 and an HSCB HDR Plasmid with the RFect^{SP} Plasmid DNA Transfection Reagent. Following that,

513 stably transfected K562 cells were selected for 7 days by 5 µg/mL puromycin and then subjected

514 to different treatments.

Enzyme activity assays

The activities of aconitase and PDH were respectively analyzed with the Aconitase Activity Assay

Kit (ab109712, Abcam, UK) and the PDH Enzyme Activity Microplate Assay Kit (ab109902,

Abcam, UK). For aconitase activity assays, $1{\times}10^6$ Ery 17 days were collected by centrifugation at

1500 g for 5 min, washed with PBS for two times, and then resuspended in the Aconitase

preservation solution provided with the kit. The cell suspension was incubated in the presence of

the detergent provided with the kit on ice for 30 min and then centrifuged at 20,000 g for 10 min.

The supernatant was diluted into a concentration of 25 µg/50 µL with the Buffer provided with the

kit, mixed with 200 µL of Assay buffer, and measured for optical density at the wavelengh of 240

nm during a 30-min interval under room temperature. The whole-cell-lysate aconitase activity was

calculated as per the detailed instructions provided in the protocol of the kit. For PDH activity

assays, $2{\times}10^6$ cells were collected and washed as described above. The lysates were prepared

with the Detergent solution, diluted into a concentration of 25 µg/50 µL with a prepared 1× Buffer,

and measured for optical density at the wavelengh of 450 nm during a 30-min interval under room

temperature. The PDH activity was calculated as per the detailed instructions provided in the

protocol. All these colorimetric assays were carried out based on a Tecan Infinite M Nano

microplate reader (Switzerland).

Heme content assay

Intracellular heme contents were determined with a method described previously (Sassa, 1976).

Briefly, whole cell lysates were prepared with the RIPA Lysis Buffer (PC101, Epizyme, China).

Afterwards, the lystates were analyzed with the Omni-Easy™ Instant BCA Protein Assay Kit

(ZJ102, Epizyme, China) for protein concentration determination. Then 30-µg protein samples

were added into 500 µL of 2 M oxalic acid, mixed thoroughly, and divided into two aliquots. One

aliquot was heated at 100°C for 30 min, while the other was not heated (to correct for

endogenous porphyrins). After cooling, fluorescence was determined at excitation and emission

wavelengths of 400 nm and 662 nm, respectively, using the Tecan microplate reader. A standard

curve was established based on hemin solutions to allow absolute quantification. The data are

expressed as pmol heme per milligram (mg) of cell lysate.

Flow cytometry

Briefly, 2×10^5 cells were rinsed and evenly dispersed in ice-cold phosphate buffered saline (PBS)

containing 10% fetal bovine serum (FBS), incubated at 4°C for 1 h in dark with fluorescein

isothiocyanate (FITC)- or allophycocyanin (APC)-conjugated primary antibodies, washed three

times with the ice-cold PBS containing 10% FBS, and analyzed immediately with a CytoFLEX

flow cytometer (Beckman Coulter, USA). The data were analyzed with Treestar FlowJo v.10.6.2.

The gating strategies for cell populations were formulated based on flow cytometry findings with

single stain controls for each sample (please see supplemental Figure 2D for an example).

Information on the primary antibodies utilized for flow cytometry analyses is summarized in

supplemental Table 1.

Cell sorting

Positive selection of CD34+CD90+ HSCs and negative selection of CD71-CD235a- progenitors

were respectively carried out with the CD34 MicroBead Kit UltraPure (130-100-453) plus CD90

MicroBeads, human (130-096-253) and CD71 MicroBeads, human (130-046-201) plus CD235a

(Glycophorin A) MicroBeads, human (130-050-501) combinations based on the Mini &

MidiMACS™ Starting Kit (130-091-632) (all from Miltenyi Biotec, Germany) in accordance with

the manufacturer's protocols.

Quantitative real-time PCR (qRT-PCR)

Total RNA was isolated from erythroid/megakaryocytic progenitor or K562 cells using the RNA

Isolation Kit (Omn-02, Omiget, China) and reverse-transcribed into cDNA with the All-In-One 5X

RT MasterMix (G592, ABM, Canada). For qRT-PCR, the 20-μL reaction systems were

established with the cDNA samples and the gene-specific primers listed in supplemental Table 2

based on the BlasTaq™ 2X qPCR MasterMix (G891, ABM, Canada). All reactions were

performed on the QuantStudio™ 3 Real-Time PCR System (Applied Biosystems, USA). Human

GAPDH gene served as the internal reference gene, relative to which the mRNA expression

levels of our target genes were calculated with the $2^{-\Delta\Delta Ct}$ algorithm. Four replicates were set up for

each sample, and three biologically independent samples were analyzed.

Immunofluorescence assay

K562 cells were washed with PBS, incubated in the 4% Paraformaldehyde Fix Solution (P0099,

Beyotime, China) for 20 min, and washed thrice and resuspended with the Immunol Staining

Wash Buffer (P0106, Beyotime, China). The cell suspension was dropped onto a glass slide and

allowed to dry for 30 min. Then the slide was sequentially incubated in the Immunol Staining

Blocking Buffer (P0102, Beyotime, China) for 60 min under ambient temperature and then in a

ZFPM1 Rabbit pAb solution (1:100 diluted in the wash buffer, A18183, ABclonal, China).

Afterwards, the slide was washed three times with the wash buffer and stained sequentially with

the Immunol Fluorence Staining Kit with Alexa Fluor 647-Labeled Goat Anti-Rabbit IgG (P0180,

Beyotime, China) and 4',6-diamidino-2-phenylindole (DAPI, MBD0015, Sigma-Aldrich, USA) as

per the manufacturers' instructions. Finally, the slide was observed with a Carl Zeiss LSM 880

AxioObserver Confocal Laser Scanning Microscope equipped with an Airyscan detector under

room temperature (approximately 23°C) (Zeiss, Germany). The objective was Zeiss Plan-

Apochromat 63x/1.40 Oil DIC M27 with a numerical aperture of 1.4. The imaging medium was

ZEISS Immersol 518 F fluorescence free oil. The excitation wavelengths for Alexa Fluor 647 and

DAPI were respectively 633 nm and 405 nm. The images were acquired and analyzed with the

ZEN 2.6 (blue edition) software. For data processinig, we only added scale bars (10 µm) to the

original images using the ZEN 2.6 (blue edition) software.

Separation of cytosolic and mitochondrial fractions

589   Cytosolic and mitochondrial fractions of K562 cells were obtained using the Mitochondria Isolation

590   Kit (89874, Thermo Scientific™, USA).

Co-immunoprecipitation (co-IP) analysis

Physical interactions between proteins and the phosphorylation status of target proteins were

analyzed by the Classic Magnetic Protein A/G IP/Co-IP Kit (YJ201, Epizyme, China) in

conjunction with WB analysis. The inputs used for WB in co-IP assays were 10% whole-cell

lysates. For the negative control, bait proteins were captured with Normal Rabbit IgG (2729, Cell

Signalling Technology, USA) or Mouse Control IgG (AC011, ABclonal, China). Information on the

primary antibodies used to capture bait proteins for the co-IP assays is available in supplemental

Table 1.

Nuclear proteomics analysis

Briefly, 500 µg of nuclear extracts were subjected to tandem mass tag-based mass spectrometry

(TMT-MS) performed by CapitalBio Technology, China to characterize nuclear proteomic

changes. Nuclear extracts of equal protein amounts were first subjected to proteolysis with trypsin

(V5113, Promega) at 37°C for 16 hours. The resulting peptide fragments were then desalted with

C18 Cartridge columns (WAT023590, Waters), freeze-dried, and reconstituted in a 40-µL 0.1%

formic acid solution. Equal amounts of peptide fragments from each experimental group were

then labelled utilizing the TMTpro 16plex Isobaric Label Reagent (A44520, Thermo Fisher) as per

the manufacturer's instructions. The labelled fragments were fractionated through high

performance liquid chromatography (HPLC) with the C18 Cartridge columns and analyzed by

liquid chromatography-tandem mass spectrometry (LC-MS/MS) assays using a Q Exactive HF-X

mass spectrometer (Thermo Fisher). Each assay involved a full-spectrum scan (Resolution

120,000, scan range 350‑1500 m/z, AGC target 3e6, maximum IT 50 ms) and target MS/MS

scans (Resolution 45,000, AGC target 5e4, maximum IT 80 ms). The raw MS data were searched

against the Uniprot database (uniprot-UP000005640_9606.fasta) for annotation with Proteome

Discoverer Version 2.4 (Thermo Fisher) with the following parameters: Enzyme, trypsin; Max

missed cleavages, 2; Fixed modifications, carbamidomethyl (C), TMTpro (N-term), and TMTpro

(K); Variable modifications, oxidation (M) and acetyl (protein N-term); Peptide mass tolerance,

±10 ppm; Fragment mass tolerance, 0.02 Da; Peptide confidence, high. A peptide false discovery

rate of ≤0.01 was applied as the threshold for filtering the identified peptides based on their

precursor intensity fraction. The final MS counts for each protein in different experimental groups

were compared using unpaired Student's t tests. Nuclear proteins with a |$\log_2$(fold change)| value

of ≥0.415 and an unpaired t test P value of <0.05 were deemed differentially-expressed nuclear

proteins (DENPs).

Affinity purification-MS analysis

To explore the proteins exhibiting significantly different binding affinities with FOG1, the binding

partners of F/M-tagged FOG1 in the EPO+CRISPR NC and EPO+CRISPR HSCB groups of K562

cells were purified by the Anti-FLAG® M2 Magnetic Beads (M8823, Sigma-Aldrich, USA).

Subsequently, these binding partners were subjected to TMT-MS, and their counts in the analysis

were compared using unpaired t test. A P value of <0.05 was considered to indicate proteins

exhibiting significantly different binding affinities with FOG1 between the two groups of K562 cells

(DBPs).

Western blotting (WB)

Whole cell lysates or lysates of cytosolic and mitochondrial fractions were prepared with the RIPA

Lysis Buffer (PC101, Epizyme, China). Afterwards, the lysates were subjected to bicinchoninic

acid assay (BCA) utilizing the Omni-Easy™ Instant BCA Protein Assay Kit (ZJ102, Epizyme,

China) for determination of protein concentration. Equal amounts of protein samples (20-50 μg)

were then separated by sodium dodecyl sulfate-polyacrylamide gel electrophoresis gels prepared

with the PAGE Gel Fast Preparation Kit (PG112, Epizyme, China) and blotted onto 0.22-μm

polyvinylidene fluoride membranes. After electrotransfer, the membranes were first incubated in

5% skimmed milk dissolved in Tris-buffered saline with Tween-20 (TBST) for one hour at room

temperature, then in diluted primary antibodies (see supplemental Table 1 for details) for 2-3

hours at room temperature or overnight at 4°C, and finally in appropriate 1:5000 diluted

horseradish peroxidase-conjugated secondary antibodies (LF101 or LF102, Epizyme, China).

Target protein bands were finally visualized using the Tanon™ High-sig ECL Western Blotting

Substrate (180-501, Tanon, China). For band intensity analysis, ImageJ v. 1.53 was employed.

Statistical analysis

Statistical data obtained in this study were analyzed by two-sided unpaired Student's t-test with Graphpad Prism 9.3 (GraphPad Software, USA). All data represent mean ± standard deviation of three independent biological replicates. P values less than 0.05, 0.01 and 0.001 respectively signify significant (* or #), very significant (** or ##) and extremely significant (*** or ###) differences.

Online supplemental materials

Supplemental Table 1 displays the primary antibodies used for flow cytometry, Co-IP and WB experiments in this study. Supplemental Table 2 lists the primers used for qRT-PCR assays. Supplemental Figure 1 shows a schematic diagram for work flows of gene knockdown and overexpression experiments in CD34+CD90+ HSCs and the downstream progenitors. Supplemental Figure 2 exhibits the flow cytometric profiles and appearance of CD34+CD90+ HSCs before and after induction for erythropoiesis, as well as the gating strategy used for flow cytometric analysis of cell populations. Supplemental Figure 3 displays the Co-IP results for the interaction between FOG1 and MYO1E in the shNC and shHSCB groups of E 2-day HSCs. Supplemental Figure 4 exhibits the WB data reflecting the effects of ruxolitinib, wortmannin and lonafarnib treatments on EPO-induced K562 cells. Supplemental Figure 5 shows flow cytometry scatter plots for MK 3-day HSCs.

Acknowledgments

This work was financially supported by the Jilin Province Science and Technology Development Plan Project (Grant No. 20220508072RC) and the High-level Talent Recruitment Project of Northeast Normal University for G.L., as well as the Fundamental Research Funds for the Central Universities for G.L. and Xiumei Jiang.

Author contributions:

Gang Liu and Xiumei Jiang acquired funding for the project; Gang Liu, Yunxuan Hou, and Xiumei Jiang conceived the project; all the authors performed the experiments and analyzed the data; Xin Jin and Xiuli Wang provided technical support; Gang Liu, Yunxuan Hou, and Xiumei Jiang drafted and revised the manuscript.

Conflict of interest:

The authors declare no competing financial interests.

Data availability statement

The original nuclear proteomics data have been deposited in the ProteomeXchange platform
under the identifier number of PXD041272 and will be made publicly available upon acceptance
of this manuscript. The downregulated and upregulated DENPs are respectively summarized in
Supplemental Datasets 1 and 2. The full lists of FOG1 binding partners in the EPO+CRISPR NC
and EPO+CRISPR HSCB groups of K562 cells are respectively displayed in Supplemental
Datasets 3 and 4. The full list of proteins detected by the tandem mass tag-based mass
spectrometry analysis of nuclear fractions of the EPO+CRISPR NC and EPO+CRISPR HSCB
groups of K562 cells are displayed in Supplemental Dataset 5. Other original data and more
details about experimental protocols can be obtained from the corresponding authors upon
reasonable request.
688

689
690
References

Andersson, L.C., M. Jokinen, and C.G. Gahmberg. 1979. Induction of erythroid differentiation in
the human leukaemia cell line K562. *Nature*. 278:364-365.
Cameron, J.M., A. Janer, V. Levandovskiy, N. Mackay, T.A. Rouault, W.H. Tong, I. Ogilvie, E.A.
Shoubridge, and B.H. Robinson. 2011. Mutations in iron-sulfur cluster scaffold genes
NFU1 and BOLA3 cause a fatal deficiency of multiple respiratory chain and 2-oxoacid
dehydrogenase enzymes. *Am J Hum Genet*. 89:486-495.
Cao, J.Z., H. Liu, A. Wickrema, and L.A. Godley. 2020. HIF-1 directly induces TET3 expression to
enhance 5-hmC density and induce erythroid gene expression in hypoxia. *Blood Adv*.
4:3053-3062.
Castro, L., V. Tortora, S. Mansilla, and R. Radi. 2019. Aconitases: Non-redox Iron-Sulfur Proteins
Sensitive to Reactive Species. *Acc Chem Res*. 52:2609-2619.
Cautain, B., R. Hill, N. de Pedro, and W. Link. 2015. Components and regulation of nuclear
transport processes. *FEBS J*. 282:445-462.
Crispin, A., C. Guo, C. Chen, D.R. Campagna, P.J. Schmidt, D. Lichtenstein, C. Cao, A.K.
Sendamarai, G.J. Hildick-Smith, N.C. Huston, J. Boudreaux, S.S. Bottomley, M.M. Heeney,
B.H. Paw, M.D. Fleming, and S. Ducamp. 2020. Mutations in the iron-sulfur cluster
biogenesis protein HSCB cause congenital sideroblastic anemia. *J Clin Invest*. 130:5245-
5256.
Dailey, H.A., M.G. Finnegan, and M.K. Johnson. 1994. Human Ferrochelatase Is an Iron-Sulfur
Protein. *Biochemistry*. 33:403-407.
Ding, Z.M., C.J. Huang, X.F. Jiao, D. Wu, and L.J. Huo. 2017. The role of TACC3 in mitotic spindle
organization. *Cytoskeleton (Hoboken)*. 74:369-378.
Dore, L.C., and J.D. Crispino. 2011. Transcription factor networks in erythroid cell and
megakaryocyte development. *Blood*. 118:231-239.
Ducamp, S., and M.D. Fleming. 2019. The molecular genetics of sideroblastic anemia. *Blood*.
133:59-69.

Edvardsson, L., J. Dykes, and T. Olofsson. 2006. Isolation and characterization of human myeloid
progenitor populations--TpoR as discriminator between common myeloid and
megakaryocyte/erythroid progenitors. *Exp Hematol*. 34:599-609.
Eggold, J.T., and E.B. Rankin. 2019. Erythropoiesis, EPO, macrophages, and bone. *Bone*. 119:36-
41.
Garriga-Canut, M., and S.H. Orkin. 2004. Transforming acidic coiled-coil protein 3 (TACC3)
controls friend of GATA-1 (FOG-1) subcellular localization and regulates the association
between GATA-1 and FOG-1 during hematopoiesis. *J Biol Chem*. 279:23597-23605.
Grebien, F., M.A. Kerenyi, B. Kovacic, T. Kolbe, V. Becker, H. Dolznig, K. Pfeffer, U. Klingmuller,
M. Muller, H. Beug, E.W. Mullner, and R. Moriggl. 2008. Stat5 activation enables
erythropoiesis in the absence of EpoR and Jak2. *Blood*. 111:4511-4522.
Gurney, A.L., K. Carver-Moore, F.J. de Sauvage, and M.W. Moore. 1994. Thrombocytopenia in c-
mpl-deficient mice. *Science*. 265:1445-1447.
Gutierrez, L., N. Caballero, L. Fernandez-Calleja, E. Karkoulia, and J. Strouboulis. 2020. Regulation
of GATA1 levels in erythropoiesis. *IUBMB Life*. 72:89-105.
Hitchcock, I.S., M. Hafer, V. Sangkhae, and J.A. Tucker. 2021. The thrombopoietin receptor:
revisiting the master regulator of platelet production. *Platelets*. 32:770-778.
Hoffman, R., M.J. Murnane, E.J. Benz, Jr., R. Prohaska, V. Floyd, N. Dainiak, B.G. Forget, and H.
Furthmayr. 1979. Induction of erythropoietic colonies in a human chronic myelogenous
leukemia cell line. *Blood*. 54:1182-1187.
Kaushansky, K., V.C. Broudy, N. Lin, M.J. Jorgensen, J. McCarty, N. Fox, D. Zucker-Franklin, and C.
Lofton-Day. 1995. Thrombopoietin, the Mp1 ligand, is essential for full megakaryocyte
development. *Proc Natl Acad Sci U S A*. 92:3234-3238.
Kaushansky, K., S. Lok, R.D. Holly, V.C. Broudy, N. Lin, M.C. Bailey, J.W. Forstrom, M.M. Buddle,
P.J. Oort, F.S. Hagen, and et al. 1994. Promotion of megakaryocyte progenitor expansion
and differentiation by the c-Mpl ligand thrombopoietin. *Nature*. 369:568-571.
Kubota, Y., T. Tanaka, G. Yamaoka, M. Yamaguchi, H. Ohnishi, K. Kawanishi, J. Takahara, and S.
Irino. 1996. Wortmannin, a specific inhibitor of phosphatidylinositol-3-kinase, inhibits
erythropoietin-induced erythroid differentiation of K562 cells. *Leukemia*. 10:720-726.
Kuhrt, D., and D.M. Wojchowski. 2015. Emerging EPO and EPO receptor regulators and signal
transducers. *Blood*. 125:3536-3541.
Kwon, N., E.N. Thompson, M.Y. Mayday, V. Scanlon, Y.C. Lu, and D.S. Krause. 2021. Current
understanding of human megakaryocytic-erythroid progenitors and their fate
determinants. *Curr Opin Hematol*. 28:28-35.
Lambert, S.A., A. Jolma, L.F. Campitelli, P.K. Das, Y. Yin, M. Albu, X. Chen, J. Taipale, T.R. Hughes,
and M.T. Weirauch. 2018. The Human Transcription Factors. *Cell*. 175:598-599.
Lennartsson, J., and L. Ronnstrand. 2012. Stem cell factor receptor/c-Kit: from basic science to
clinical implications. *Physiol Rev*. 92:1619-1649.
Liu, G., D. Sil, N. Maio, W.H. Tong, J.M. Bollinger, Jr., C. Krebs, and T.A. Rouault. 2020. Heme
biosynthesis depends on previously unrecognized acquisition of iron-sulfur cofactors in
human amino-levulinic acid dehydratase. *Nat Commun*. 11:6310.
Maio, N., A. Jain, and T.A. Rouault. 2020. Mammalian iron-sulfur cluster biogenesis: Recent
insights into the roles of frataxin, acyl carrier protein and ATPase-mediated transfer to
recipient proteins. *Curr Opin Chem Biol*. 55:34-44.

Maio, N., K.S. Kim, A. Singh, and T.A. Rouault. 2017. A Single Adaptable Cochaperone-Scaffold
Complex Delivers Nascent Iron-Sulfur Clusters to Mammalian Respiratory Chain
Complexes I-III. *Cell Metab*. 25:945-953 e946.
Maio, N., A. Singh, H. Uhrigshardt, N. Saxena, W.H. Tong, and T.A. Rouault. 2014. Cochaperone
binding to LYR motifs confers specificity of iron sulfur cluster delivery. *Cell Metab*.
19:445-457.
Mancini, E., A. Sanjuan-Pla, L. Luciani, S. Moore, A. Grover, A. Zay, K.D. Rasmussen, S. Luc, D.
Bilbao, D. O'Carroll, S.E. Jacobsen, and C. Nerlov. 2012. FOG-1 and GATA-1 act
sequentially to specify definitive megakaryocytic and erythroid progenitors. *EMBO J*.
31:351-365.
Nandakumar, S.K., J.C. Ulirsch, and V.G. Sankaran. 2016. Advances in understanding
erythropoiesis: evolving perspectives. *Br J Haematol*. 173:206-218.
Noris, P., and A. Pecci. 2017. Hereditary thrombocytopenias: a growing list of disorders.
Hematology Am Soc Hematol Educ Program. 2017:385-399.
Polson, E.S., V.B. Kuchler, C. Abbosh, E.M. Ross, R.K. Mathew, H.A. Beard, B. da Silva, A.N.
Holding, S. Ballereau, E. Chuntharpursat-Bon, J. Williams, H.B.S. Griffiths, H. Shao, A.
Patel, A.J. Davies, A. Droop, P. Chumas, S.C. Short, M. Lorger, J.E. Gestwicki, L.D.
Roberts, R.S. Bon, S.J. Allison, S. Zhu, F. Markowetz, and H. Wurdak. 2018. KHS101
disrupts energy metabolism in human glioblastoma cells and reduces tumor growth in
mice. *Sci Transl Med*. 10.
Saatci, O., O. Akbulut, M. Cetin, V. Sikirzhytski, M. Uner, D. Lengerli, E.C. O'Quinn, M.J. Romeo,
B. Caliskan, E. Banoglu, S. Aksoy, A. Uner, and O. Sahin. 2023. Targeting TACC3
represents a novel vulnerability in highly aggressive breast cancers with centrosome
amplification. *Cell Death Differ*.
Sassa, S. 1976. Sequential induction of heme pathway enzymes during erythroid differentiation
of mouse Friend leukemia virus-infected cells. *J Exp Med*. 143:305-315.
Sutherland, J.A., A.R. Turner, P. Mannoni, L.E. McGann, and J.M. Turc. 1986. Differentiation of
K562 leukemia cells along erythroid, macrophage, and megakaryocyte lineages. *J Biol
Response Mod*. 5:250-262.
Tsang, A.P., Y. Fujiwara, D.B. Hom, and S.H. Orkin. 1998. Failure of megakaryopoiesis and
arrested erythropoiesis in mice lacking the GATA-1 transcriptional cofactor FOG. *Genes
Dev*. 12:1176-1188.
Wing, C.E., H.Y.J. Fung, and Y.M. Chook. 2022. Karyopherin-mediated nucleocytoplasmic
transport. *Nat Rev Mol Cell Biol*. 23:307-328.
Wu, H., X. Liu, R. Jaenisch, and H.F. Lodish. 1995. Generation of committed erythroid BFU-E and
CFU-E progenitors does not require erythropoietin or the erythropoietin receptor. *Cell*.
83:59-67.
Zhao, W., C. Kitidis, M.D. Fleming, H.F. Lodish, and S. Ghaffari. 2006. Erythropoietin stimulates
phosphorylation and activation of GATA-1 via the PI3-kinase/AKT signaling pathway.
Blood. 107:907-915.
802

Figure legends

Figure 1. Heat shock cognate B (HSCB) played an important iron sulfur cluster (ISC)
delivery independent function during erythropoiesis of CD34+CD90+ hematopoietic stem
cells (HSCs). (A) Western blotting analyses on Ery 17 days verified the efficacy of short hairpin
RNA (shRNA) knockdown maintained throughout the experimental period. (B) Aconitase activity,
(C) pyruvate dehydrogenase (PDH) activity and (D) heme content of the shHSCB, shISCU and
shNFS1 groups were significantly lower than those of the shNC group, indicating inhibited ISC
biogenesis in the shHSCB, shISCU and shNFS1 groups. (E) Representative flow cytometry
scatter plots and (F) statistics of three biologically independent flow cytometry assays displaying
the percentages of CD71+CD235a+ progenitors in the shNC, shHSCB, shISCU and shNFS1
groups. (G) Quantitative real-time PCR (qRT-PCR) analyses of the mRNA levels of *HBB*, *ALAS2*
and *GYPC* genes for the shNC, shHSCB, shISCU and shNFS1 groups. The flow cytometry and
qRT-PCR results respectively signified significantly decreased percentages of CD71+CD235a+
progenitors and significantly lower mRNA levels of the erythroid-specific genes in the shHSCB
group compared with the shNC group, revealing inhibited erythropoiesis of the shHSCB group of
Ery 17 days. (H) Flow cytometry scatter plots showing that the majority of CD71-CD235a- cells
from the shHSCB group were CD34+CD38+CD45RA-IL3RA-. Error bars denote standard
deviations; *, ** and *** respectively denote *P* values less than 0.05, 0.01 and 0.001 relative to the
shNC groups, two-sided Student's t test.

Figure 2. Knockout of HSCB with the CRISPR-Cas9 technique impaired erythropoietin
(EPO)-induced erythropoiesis of K562 cells. (A) Western blotting analyses exhibited
successful knockout of HSCB and decreased levels of erythrocyte membrane proteins (GYPA
and SPTA1) in the EPO+CRISPR HSCB K562 cells. Error bars denote standard deviations; * and
** respectively signify *P* values less than 0.05 and 0.01 compared with the Untreated group, while
$^{##}$ means *P* < 0.01 relative to the EPO+CRISPR NC group, two-sided Student's t test. (B) Heme
contents and (C) mRNA levels of the erythroid-specific genes were significantly lower in the
EPO+CRISPR HSCB group than in the EPO+CRISPR NC group. Error bars denote standard
deviations; ** and *** respectively represent *P* values less than 0.01 and 0.001 compared with the
Untreated group, while $^{###}$ means *P* < 0.001 relative to the EPO+CRISPR NC group, two-sided
Student's t test. (D) A volcano plot for the differentially-expressed nuclear proteins (DENPs)
detected by tandem mass tag-based mass spectrometry (TMT-MS) assays and (E) detailed
information of the *bona fide* DENPs between the EPO+CRISPR NC and EPO+CRISPR HSCB
groups. (F) Western blotting and (G) the corresponding band intensity analyses for FOG1, NAA30,
LYAR, and GATA1 proteins in the nuclear fractions of the EPO+CRISPR NC and EPO+CRISPR
HSCB K562 cells and shNC and shHSCB E 2-day HSCs. Friend of GATA1 (FOG1) was identified
as the only nuclear protein expressed at significantly different levels in both K562 cells and E 2-
day HSCs after HSCB knockout or knockdown. Error bars denote standard deviations; *
represents *P* < 0.05 relative to the EPO+CRISPR NC group, while $^{##}$ means *P* < 0.01 compared
with the shNC group, two-sided Student's t test.

Figure 3. Impaired FOG1 nuclear translocation was responsible for the inhibited
erythropoiesis of HSCB-deficient K562 cells and HSCs. (A) Western blotting analyses
indicated that FOG1 overexpression increased the levels of nuclear FOG1 in HSCB-deficient
EPO-induced K562 cells and E 2-day HSCs, as well as the levels of erythrocyte membrane
proteins in HSCB-deficient EPO-induced K562 cells without significantly affecting the expression
level of GATA1. Error bars denote standard deviations; * and ** respectively represent *P* values
less than 0.05 and 0.01 compared with the EPO+CRISPR NC or shNC group, while $^{#}$ denotes *P* <

24

0.05 relative to the EPO+CRISPR HSCB or shHSCB group, two-sided Student's t test. (B) Heme
contents in the EPO+CRISPR NC, EPO+CRISPR HSCB and EPO+CRISPR HSCB+FOG1 OE
groups of K562 cells, as well as in the shNC, shHSCB and shHSCB+FOG1 OE groups of Ery 17
days. Error bars denote standard deviations; ** denotes $P < 0.01$ relative to the EPO+CRISPR NC
group, while ## and ### respectively represent P values less than 0.01 and 0.001 compared with
the shNC group, two-sided Student's t test. (C) qRT-PCR analyses on HSCB-deficient EPO-
induced K562 cells and HSCB-deficient Ery 17 days demonstrated resumed mRNA expression
levels of erythroid-specific genes after FOG1 overexpression. (D) Representative flow cytometry
scatter plots and (E) statistics of three biologically independent flow cytometry assays showing
that the percentages of CD71+CD235a+ progenitors among HSCB-deficient Ery 17 days
recovered after FOG1 overexpression. Error bars denote standard deviations; *** signifies $P <$
0.001 relative to the shNC group, two-sided Student's t test. (F) Western blotting analyses of
cytoplasmic FOG1 and (G) immunofluorescence (IF) assays for FOG1 revealed inhibited FOG1
nuclear translocation in HSCB-deficient EPO-induced K562 cells and E 2-day HSCs (scale
bars=10 μm). Error bars denote standard deviations; ** and *** in panel (F) respectively represent
P values less than 0.01 and 0.001 compared with the EPO+CRISPR NC or shNC group, two-
sided Student's t test.

Figure 4. Deficiency of HSCB blocked FOG1 nuclear translocation by enhancing its
interaction with transforming acidic coiled-coil containing protein 3 (TACC3). (A) A
histogram showing the two differential binding proteins (DBPs) of FOG1 between the
EPO+CRISPR NC and EPO+CRISPR HSCB groups. Error bars denote standard deviations; **
signifies $P < 0.01$ compared with the EPO+CRISPR NC group, two-sided Student's t test. Co-
immunoprecipitation (Co-IP) assays in (B) K562 cells and (C) E 2-day HSCs indicated that HSCB
deficiency promoted the binding between FOG1 and TACC3. (D) Western blotting analyses on
K562 cells and E 2-day HSCs and (E) immunofluorescence (IF) assays on K562 cells revealed
that treatment with KHS101, a small-molecule inhibitor that could reduce the activity and protein
abundance of TACC3, decreased TACC3 protein levels and increased relative nuclear FOG1 (N-
FOG1) to cytoplasmic FOG1 (C-FOG1) ratios. Scale bars for IF assays indicate 10 μm. Error bars
denote standard deviations; * and ** in panel (D) respectively denote P values less than 0.05 and
0.01 compared with the EPO+CRISPR NC or shNC group, while #, ## and ### respectively mean P
< 0.05, $P < 0.01$ and $P < 0.001$ relative to the EPO+CRISPR HSCB or shHSCB group, two-sided
Student's t test. (F) qRT-PCR analyses on HSCB-deficient EPO-induced K562 cells and HSCB-
deficient Ery 17 days demonstrated recovered mRNA expression levels of erythroid-specific
genes after KHS101 treatment. (G) Representative flow cytometry scatter plots and (H) statistics
of three biologically independent flow cytometry assays showing that the percentages of
CD71+CD235a+ progenitors among HSCB-deficient Ery 17 days recovered after KHS101
treatment. Error bars denote standard deviations; *** means $P < 0.001$ compared with the shNC
group, two-sided Student's t test.

Figure 5. HSCB bound with and mediated the proteasomal degradation of TACC3 to
facilitate FOG1 nuclear translocation upon EPO/erythropoietin receptor (EPOR) signaling
activation. (A) Western blotting analyses indicated that EPO treatment decreased the level of
TACC3 and increased the relative N-FOG1/C-FOG1 ratio, HSCB overexpression augmented the
effects of EPO treatment on the protein level of TACC3 and FOG1 nuclear translocation, whereas
treatment with the proteasome inhibitor MG132 abrogated the effects of EPO treatment+HSCB

overexpression on the protein level of TACC3 and FOG1 nuclear translocation. Error bars denote
standard deviations; * and ** respectively represent P values less than 0.05 and 0.01 compared
with the Untreated group, while $^{#}$ and $^{##}$ respectively mean $P < 0.05$ and $P < 0.01$ relative to the
EPO+Empty Vector group, two-sided Student's t test. (B) Co-IP results showing that TACC3
interacted with HSCB when the K562 cells were treated with EPO, whereas it interacted with
FOG1 in the untreated and EPO+HSCB OE+MG132 K562 cells. (C) Flow cytometry scatter plots
exhibiting the levels of CD71 and CD235a on the cell surface of CD34+CD90+ HSCs induced for
erythropoiesis for 0, 2, 4, 10, 17 and 24 days. The flow cytometry analyses were repeated twice
with similar findings to ensure reproducibility. (D) Western blotting analyses of the levels of
cytoplasmic FOG1, TACC3 and nuclear FOG1 in CD34+CD90+ HSCs induced for erythropoiesis
for 0, 2, 4, 10, 17 and 24 days. (E) Co-IP analyses of the binding affinities of TACC3 with HSCB
and FOG1 in CD34+CD90+ HSCs induced for erythropoiesis for 0, 2, 4, 10, 17 and 24 days. (F)
Flow cytometry scatter plots showing the levels of CD71 and EPOR on the cell surface of
CD34+CD90+ HSCs induced for erythropoiesis for 0, 2, and 4 days. The western blotting, co-IP,
and flow cytometry data highlighted that the EPO/EPOR signaling was indispensable for the
functionalization of HSCB in CD34+CD90+ HSCs induced for erythropoiesis.

Figure 6. HSCB phosphorylation by phosphoinositol-3-kinase (PI3K) was necessary for its
functionalization during erythropoiesis of K562 cells and CD34+CD90+ HSCs. (A) Western
blotting analyses demonstrated that treatment with ruxolitinib and wortmannin, which are
respectively inhibitors against Janus tyrosine kinase 2 (JAK2) and PI3K, increased the protein
level of TACC3 and decreased FOG1 nuclear translocation. Error bars denote standard
deviations; $^{#}$, $^{##}$ and $^{###}$ respectively mean $P < 0.05$, $P < 0.01$ and $P < 0.001$ relative to the
EPO+Vehicle group, two-sided Student's t test. (B and C) Co-IP assays reflected that wortmannin
treatment impaired the interaction between TACC3 and HSCB, promoted the binding between
TACC3 and FOG1, and reduced the phosphorylation level of cytosolic HSCB in EPO-induced
K562 cells and E 2-day HSCs. (D) Co-IP experiments verified the physical interaction between
cytosolic HSCB and the P85-alpha subunit of PI3K (PIK3R1). (E) Western blotting, (F) heme
content, and (G) qRT-PCR assays confirmed that wortmannin treatment inhibited the erythroid
differentiation of EPO-induced K562 cells. Error bars denote standard deviations; *, ** and ***
respectively denote P values less than 0.05, 0.01 and 0.001 relative to the Untreated group, while
$^{##}$ and $^{###}$ respectively represent P values less than 0.01 and 0.001 compared with the
EPO+Vehicle group, two-sided Student's t test. (H) Representative flow cytometry scatter plots
and (I) statistics of three biologically independent flow cytometry assays exhibiting significantly
reduced percentages of CD71+ cells among E 4-day HSCs in the Wortmannin group. Error bars
denote standard deviations; ** means $P < 0.01$ compared with the Vehicle group, two-sided
Student's t test.

Figure 7. Megakaryopoiesis of CD34+CD90+ HSCs also required the activation of the PI3K-
HSCB signaling axis. (A) Representative flow cytometry scatter plots and (B) statistics of three
biologically independent flow cytometry assays revealing significantly reduced percentages of
CD41a+CD42b+ cells among the MK 14-day HSCs in the shHSC and Wortmannin groups. (C)
qRT-PCR assays indicated decreased mRNA expression levels of megakaryocyte-specific genes
VWF and *GP1BA* in the MK 14-day HSCs of the shHSC and Wortmannin groups. Error bars
denote standard deviations; *** and $^{###}$ respectively denote $P < 0.001$ relative to the shNC and
Vehicle groups, two-sided Student's t test. (D) Flow cytometry scatter plots showing that the

majority of cells from the shHSCB group of MK 14-day HSCs were CD34+CD38+CD45RA-

IL3RA-. (E) Western blotting analyses demonstrated that both knocking down HSCB and

wortmannin treatment could increase the protein level of TACC3 while reduce FOG1 nuclear

translocation, whereas treating HSCB-deficient HSCs with KHS101 decreased the protein level of

TACC3 while enhanced FOG1 nuclear translocation in MK 3-day HSCs. Error bars denote

standard deviations; ** and *** respectively represent P values less than 0.01 and 0.001 compared

with the shNC group, while # and ## respectively mean $P < 0.05$ and $P < 0.01$ relative to the

Vehicle group, two-sided Student's t test. (F) Co-IP analyses revealed decreased phosphorylation

level of cytosolic HSCB and impaired interaction between HSCB and TACC3 in MK 3-day HSCs

treated with wortmannin. (G) Western blotting analyses of nuclear FOG1 levels in CD34+CD90+

HSCs induced for megakaryopoiesis for 0, 3, 7, and 14 days. (H) A schematic diagram

summarizing the main findings of this study.

Figure 1

Figure 2

A K562 cells

HSCB — 25 kD
GYPA — 35 kD
SPTA1 — 250 kD
GAPDH — 35 kD

Untreated / EPO / EPO+CRISPR NC / EPO+CRISPR HSCB

Band intensity relative to GAPDH — GYPA, SPTA1

B K562 cells

Heme content (pmol/mg cell lysate)
Untreated, EPO, EPO+CRISPR NC, EPO+CRISPR HSCB

C K562 cells

Relative gene expression levels — HBB, ALAS2, GYPC
EPO / EPO+CRISPR NC / EPO+CRISPR HSCB

D K562 cells

EPO+CRISPR HSCB vs. EPO+CRISPR NC

Downregulated / Not changed / Upregulated

$-\log_{10}$(P value) vs $\log_2$(Fold Change)

E

Protein Name	Fold Change (EPO+CRISPR HSCB vs. EPO+ CRISPR NC)	P Value	Brief Description
NAA30	0.746	0.0201	Enables peptide alpha-N-acetyltransferase activity and is involved in N-terminal peptidyl-methionine acetylation
LYAR	0.680	0.0329	Possesses transcription factor binding, identical protein binding and transcription regulator inhibitor activities, and is involved in regulation of erythrocyte development and innate immune response
FOG1	0.610	0.0299	Possesses transcription factor binding and transcription corepressor activities, and is involved in platelet formation and definitive erythrocyte differentiation.

F K562 cells nuclear fraction

Replicate 1 Replicate 2 Replicate 3
EPO+CRISPR NC / EPO+CRISPR HSCB

FOG1 — 100 kD
NAA30 — 40 kD
LYAR — 40 kD
GATA1 — 50 kD
LMNB1 — 70 kD

E 2-day HSCs nuclear fraction

Replicate 1 Replicate 2 Replicate 3
shNC / shHSCB

FOG1 — 100 kD
NAA30 — 40 kD
LYAR — 40 kD
GATA1 — 50 kD
LMNB1 — 70 kD

G

Band intensity relative to LMNB1 — FOG1, NAA30, LYAR, GATA1

EPO+CRISPR NC / EPO+CRISPR HSCB / shNC / shHSCB

Figure 3

Figure 4

A K562 cells

Affinity purification-MS counts

☐ EPO+CRISPR NC
☐ EPO+CRISPR HSCB

MYO1E   TACC3

B K562 cells

EPO+CRISPR NC

Input (10% lysates) | F/M-FOG1 IPed with IgG | F/M-FOG1 IPed with anti-FLAG

Untransfected | Empty Vector | F/M-FOG1

FLAG/MYC-FOG1 FOG1 — 100 kD
TACC3 — 150 kD

EPO+CRISPR HSCB

Input (10% lysates) | F/M-FOG1 IPed with IgG | F/M-FOG1 IPed with anti-FLAG

Untransfected | Empty Vector | F/M-FOG1

— 100 kD
— 150 kD

C E 2-day HSCs

shNC

Input (10% lysate) | IPed with anti-TACC3 | IPed with IgG

TACC3 — 150 kD
FOG1 — 100 kD

shHSCB

Input (10% lysate) | IPed with anti-TACC3 | IPed with IgG

— 150 kD
— 100 kD

D K562 cells

Total lysate

EPO+CRISPR NC | EPO+CRISPR HSCB | EPO+CRISPR HSCB+KHS101

HSCB — 25 kD
TACC3 — 150 kD
FOG1 — 100 kD
GAPDH — 35 kD

☐ EPO+CRISPR NC
☐ EPO+CRISPR HSCB
☐ EPO+CRISPR HSCB+KHS101

Band intensity relative to GAPDH

TACC3   FOG1

Nuclear fraction

EPO+CRISPR NC | EPO+CRISPR HSCB | EPO+CRISPR HSCB+KHS101

FOG1 — 100 kD
LMNB1 — 70 kD

Cytoplasmic fraction

FOG1 — 100 kD
GAPDH — 35 kD

Relative N-FOG1/C-FOG1 ratios

EPO+CRISPR NC
EPO+CRISPR HSCB
EPO+CRISPR HSCB+KHS101

E 2-day HSCs

Total lysate

shNC | shHSCB | shHSCB+KHS101

HSCB — 25 kD
TACC3 — 150 kD
FOG1 — 100 kD
GAPDH — 35 kD

☐ shNC
☐ shHSCB
☐ shHSCB+KHS101

Band intensity relative to GAPDH

TACC3   FOG1

Nuclear fraction

shNC | shHSCB | shHSCB+KHS101

FOG1 — 100 kD
LMNB1 — 70 kD

Cytoplasmic fraction

FOG1 — 100 kD
GAPDH — 35 kD

Relative N-FOG1/C-FOG1 ratios

shNC
shHSCB
shHSCB+KHS101

E K562 cells

FOG1 | DAPI | Merged

EPO+CRISPR NC
EPO+CRISPR HSCB
EPO+CRISPR HSCB +KHS101

10 μm

F K562 cells

☐ EPO+CRISPR HSCB
☐ EPO+CRISPR HSCB+KHS101

Relative gene expression levels

HBB   ALAS2   GYPC

Ery 17 days

☐ shHSCB
☐ shHSCB+KHS101

Relative gene expression levels

HBB   ALAS2   GYPC

G Ery 17 days

shNC
Q1 12.2 | Q2 62.6 | Q4 21.6 | Q3 3.64

shHSCB
Q1 14.3 | Q2 14.4 | Q4 68.8 | Q3 2.53

shHSCB+KHS101
Q1 33.7 | Q2 50.5 | Q4 15.4 | Q3 0.37

APC-CD71
FITC-CD235a

H

Percentage of CD71+CD235a+ cells

shNC
shHSCB
shHSCB+KHS101

Figure 5

Figure 6

Figure 7

Figure S1

Figure S2

A

Q1 4.50   Q2 91.1

Q4 3.32   Q3 1.08

APC-CD90

FITC-CD34

B

Ery 17 days

shNC   shHSCB   shISCU   shNFS1

C   **E 2-day HSCs**

Q1 72.4   Q2 4.21

Q4 23.4   Q3 0

APC-CD38

FITC-CD45RA

Q1 83.5   Q2 5.72

Q4 10.8   Q3 0

FITC-CD34

APC-IL3RA

D   **Gating strategy**

APC Single Staining

FITC positive

FITC Single Staining

APC positive

Figure S3

Figure S4

K562 cells

Figure S5

MK 3-day HSCs