A disrupted compartment boundary underlies abnormal cardiac patterning and

congenital heart defects

Irfan S. Kathiriya[1,#], Martin H. Dominguez[2,3,4], Kavitha S. Rao[1,2], Jonathon M. Muncie-Vasic[2], W.

Patrick Devine[2,5], Kevin M. Hu[1,2,6], Swetansu K. Hota[2,7], Bayardo I. Garay[1,8], Diego Quintero[2,9],

Piyush Goyal[1,2,10], Megan N. Matthews[1], Reuben Thomas[2], Tatyana Sukonnik[2,15], Dario Miguel-

Perez[2], Sarah Winchester[2], Emily F. Brower[2], André Forjaz[11], Pei-Hsun Wu[11], Denis Wirtz[11],

Ashley L. Kiemen[11], Benoit G. Bruneau[2,12,13,14,#]

[#]corresponding authors

[1] Department of Anesthesia and Perioperative Care, University of California, San Francisco,

San Francisco, CA

[2] Gladstone Institutes, San Francisco, CA

[3] Department of Medicine, University of California, San Francisco, San Francisco, CA

[4] Current address: Department of Medicine (Cardiovascular Medicine), Perelman School of

Medicine, University of Pennsylvania, Philadelphia, PA

[5] Current address: Department of Pathology, University of California, San Francisco, San

Francisco, CA

[6] Current address: Creighton University School of Medicine, Omaha, NE

[7] Current address: Herman B. Wells Center for Pediatric Research, Indiana University School of

Medicine, Indianapolis, IN

[8] Current address: University of Minnesota Medical Scientist Training Program, Minneapolis, MN

[9] Current address: Department of Human Genetics, Emory University School of Medicine,

Atlanta, GA

[10] Current address: Touro University California, Vallejo, CA

[11] Department of Chemical and Biomolecular Engineering, Johns Hopkins University, Baltimore,

MD

[12] Roddenberry Center for Stem Cell Biology and Medicine, Gladstone Institutes, San Francisco,

CA

[13] Cardiovascular Research Institute, University of California, San Francisco, San Francisco, CA

[14] Department of Pediatrics, University of California, San Francisco, San Francisco, CA

[15] Deceased

Abstract.

Failure of septation of the interventricular septum (IVS) is the most common congenital heart

defect (CHD), but mechanisms for patterning the IVS are largely unknown. We show that a

Tbx5$^+$/Mef2cAHF$^+$ progenitor lineage forms a compartment boundary bisecting the IVS. This

coordinated population originates at a first- and second heart field interface, subsequently forming

a morphogenetic nexus. Ablation of *Tbx5$^+$/Mef2cAHF$^+$* progenitors cause IVS disorganization,

right ventricular hypoplasia and mixing of IVS lineages. Reduced dosage of the CHD transcription

factor TBX5 disrupts boundary position and integrity, resulting in ventricular septation defects

(VSDs) and patterning defects, including *Slit2* and *Ntn1* misexpression. Reducing NTN1 dosage

partly rescues cardiac defects in *Tbx5* mutant embryos. Loss of *Slit2* or *Ntn1* causes VSDs and

perturbed septal lineage distributions. Thus, we identify essential cues that direct progenitors to

pattern a compartment boundary for proper cardiac septation, revealing new mechanisms for

cardiac birth defects.

Main.

Organogenesis relies on fine tissue patterning, and disturbances cause organ

malformations that result in birth defects. Examples of prevalent birth defects resulting from

abnormal patterning include congenital limb abnormalities, neural tube defects or congenital heart

defects (CHDs).

CHDs are the most common birth defects and are a leading cause of morbidity and

mortality in childhood. CHDs are thought to result from alterations to the orchestrated patterning

of heart development. Nearly half of patients with CHDs have ventricular septal defects (VSDs)

or atrial septal defects (ASDs), either in isolation or combined with other anatomic defects,

including abnormal chamber formation. For example, atrioventricular canal (AVC) defects include

VSDs, ASDs and abnormal development of the atrioventricular (AV) valves, with severe cases

leading to chamber hypoplasia and functional single ventricle physiology. There is a large gap in

our understanding of how early developmental events pattern cardiac morphogenesis and

anatomy. Addressing this understudied need can inform diagnostic prognosis, family planning

and therapeutic approaches for CHDs.

Complete atrioventricular septation into four chambers allows for separation of systemic

and pulmonary circulations and enables higher levels of oxygen for transport in arterial blood.

Indeed, some progress has been made to determine the embryonic origins of the IVS [1,23]. A dye-

labelled approach in chick embryos revealed that labelling of the ventral surface of the linear heart

tube becomes positioned at the outer curvature of the linear heart tube [1,2]. When labelled at the

outer curvature upon rightward looping, the label was found at the apical border of the primary

interventricular foramen [1,2]. Discovery of the *Ganglion Nodosum* epitope from chick, which

demarcated the primary interventricular foramen and subsequently the IVS, AV junction and

ventricular conduction system in mouse, chick and humans, led to the description of a "primary

ring" for the ventricular septum [3–5].

Discrete early cardiac progenitors in mesoderm contribute to specific cardiac anatomy [6–

8]. The first heart field (FHF) gives rise primarily to the left ventricle (LV) and parts of the atria,

while the second heart field (SHF) contributes predominantly to the right ventricle (RV), outflow

tract (OFT) and portions of the atria [9]. The heart fields can be largely recapitulated by marking

lineage-labelled progenitors at gastrulation. Specifically, a lineage labelled by the CHD-linked

transcription factor *Tbx5* (*Tbx5*+-lineage) contributes to the LV and atria, while the anterior heart

field enhancer of *Mef2c* (*Mef2cAHF*+)-lineage largely contributes to the RV and OFT [6].

An aspect of proper tissue patterning entails the establishment of compartment

boundaries to separate neighboring fields of cells that exhibit discrete functions [10–12]. Few

compartment boundaries have been identified in mammals, including in the limb [13–15] and at the

midbrain-hindbrain border [16]. In the developing heart, both the LV and RV lineages supply

descendants to the interventricular septum (IVS) and establish a mutual border between the two

lineages [6,17]. Clonal analysis shows segregation at the IVS of the two lineages into non-

intermingling cellular compartments, on either the left or right sides of the IVS [6,17]. This lineage

configuration is reminiscent of a compartment boundary[10–12]. Using an intersectional-lineage

labelling approach, a subset of cardiac progenitor cells labeled by both *Tbx5*+/*Mef2cAHF*+ were

described with contributions to the left side of this apparent compartment boundary [6].

Understanding the role of the *Tbx5*+/*Mef2cAHF*+ lineage-labeled compartment boundary at the

IVS may reveal clues about IVS patterning with potential relevance for VSD etiologies. More

broadly, how disturbances to compartment boundaries contribute to birth defects is largely

unknown.

Here, we combined genetic lineage labeling with light sheet microscopy to follow

contributions of *Tbx5*+/*Mef2cAHF*+ precursors to a compartment boundary during cardiac

morphogenesis. We ablated *Tbx5*+/*Mef2cAHF*+ precursors to determine the essential roles of the

lineage-labeled compartment boundary for cardiac septation and chamber development. By

leveraging a genetic lesion of the CHD gene *Tbx5,* our work uncovers the genetic regulation of

the *Tbx5*+/*Mef2cAHF*+ lineage-labeled compartment boundary for proper IVS patterning. We

deployed single cell RNA sequencing (scRNAseq) to discover novel Tbx5-sensitive cues that are

essential for compartment boundary regulation and formation of the IVS during heart

development.

Results.

<u>Early cardiac progenitors for the IVS, IAS and AVC.</u> We used conditional dual lineage labelling

by *Tbx5*[CreERT2/+] or *Mef2cAHF-DreERT2* and marked cells during gastrulation at embryonic day

(E)6.5 by tamoxifen-induced recombination. We followed reporter-labeled lineages by

epifluorescence microscopy, histology or light sheet microscopy (Figure 1). At E14.5, the *Tbx5*[+]

lineage contributed to the left side of the IVS, often ending at a presumptive line marked at the

apex by the interventricular (IV) groove, consistent with previous reports suggesting a lineage

boundary [6,17]. *Tbx5*[+] and *Mef2cAHF*[+] lineages showed largely complementary patterns, with

notable overlap of *Tbx5*[+] and *Mef2cAHF*[+] lineages at the IVS, AVC and IAS (Figure 1B, D, F). To

better visualize the *Tbx5*[+]/*Mef2cAHF*[+] lineage, we used a conditional intersectional-lineage

labeling approach. Using lineage reporters responsive to both CreERT2 and DreERT2 [18], we

observed a striking spatial pattern of the *Tbx5*[+]/*Mef2cAHF*[+] lineage in the IVS, AVC and IAS. At

E14.5, the septal lineage extended to the apex of the heart via the IVS, to the base of the heart

at the AV canal, as well as to the IAS superiorly[15–18] (Figure 1H-J). These results expanded upon

previous findings in the developing IVS at E10.5 [6], by identifying contributions from

Tbx5[+]/*Mef2cAHF*[+] progenitors to additional anatomic sites during chamber formation and

septation. Moreover, lineage contributions to sites of cardiac septation were derived from

Tbx5[+]/*Mef2cAHF*[+] progenitors at E6.5 (Supp. Figure 1), but not later at E8.5 or E10.5, implicating

a narrow, early window for capturing *Tbx5*[+]/*Mef2cAHF*[+] septal progenitors during cardiac

mesoderm formation.

<u>*Tbx5*[+]/*Mef2cAHF*[+] septal lineage is prefigured in early mouse heart development.</u> To characterize

the lineage derived from *Tbx5*[+]/*Mef2cAHF*[+] septal progenitors during early heart development, we

examined the localization of the lineage prior to heart tube formation at E8.0 (Figure 2, Video 1).

By light sheet imaging, the pattern of the *Tbx5*[+]/*Mef2cAHF*[+] lineage was detected at the dorsal

aspect of the $Tbx5^+$ lineage, at an interface between the presumptive first and second heart fields

(Figure 2A-E"). Cell segmentation at this early stage revealed that the $Tbx5^+/Mef2cAHF^+$ lineage

resembled a nascent halo of cells that became readily apparent at later stages of heart

development (Figure 2F-I).

We further examined the lineage contributions of $Tbx5^+/Mef2cAHF^+$ septal progenitors at

subsequent timepoints during cardiac morphogenesis by light sheet microscopy. At the linear

heart tube stage at E8.25, the labeled $Tbx5^+/Mef2cAHF^+$ lineage appeared intricately arranged

into a ring of cells between the future left and right ventricles (Figure 2J, K). This circlet

configuration was maintained during rightward looping of the heart at E8.5 (Figure 2O-Q) and

subsequent chamber formation. At E10.5, we observed a band of lineage-labeled cells from the

IV groove at the outer curvature to the inner curvature near the AV groove that extended

posteriorly to the atria (Figure 2R-T), reminiscent of the "primary ring" [3–5]. Optical sections showed

the labeled lineage superiorly at a crossroads of the AVC, outflow tract and atria (Figure 2S). This

populates a morphogenetic nexus, where VSDs can occur due to abnormal connections between

the IVS and the AV cushions, outflow tract cushions, or the muscular septum itself. In the atria,

we noted lineage-labeled cells at the midline, superiorly and inferiorly, suggesting that the

Tbx5$^+$/Mef2cAHF$^+$ lineage potentially marks the presumptive location of the IAS for later stages

(Supp. Figure 2B-E). Consequently, this data is consistent with a notion that the $Tbx5^+/Mef2cAHF^+$

lineage is configured early in heart development, well before subsequent morphogenetic events

of chamber formation and cardiac septation.

IVS disorganization and ventricular hypoplasia from cell ablation of $Tbx5^+/Mef2cAHF^+$

progenitors. To determine a role for the $Tbx5^+/Mef2cAHF^+$ septal progenitors during heart

development, we conditionally ablated these cells at E6.5. To this end, we generated an

intersectional recombinase-responsive $DTA176$ knock-in mouse allele at the $Hip11$ safe-harbor

locus, by which cells would be killed where approximately 100-200 molecules of DTA176 were

expressed [19]. Upon tamoxifen administration to pregnant dams at E6.5, intersectional-*DTA* mutant

(*Tbx5^{CreERT2/+}; Mef2cAHF-DreERT2; Hip11^{Intersectional-DTA176/+}*) embryos displayed RV hypoplasia at

E9.5 (Figure 3A, B) and E12.5 (Figure 3C-J), together with IVS disorganization and non-

compaction with a blunted IV groove (Figure 3H, J). Further, there were defects of the AV

cushions and absence of the IAS (Figure 3H, J). A few *Tbx5^+/Mef2cAHF^+* lineage cells remained

in mutant embryos, likely reflecting some degree of inefficient dual recombination of both the

reporter allele and the intersectional-*DTA* transgene. Beyond E12.5, intersectional-*DTA* mutant

embryos were not recovered. Hence, these findings suggested that the *Tbx5^+/Mef2cAHF^+* septal

progenitors were not only important for IVS development and AV septation, but also for RV

chamber formation.

Lineage mixing from disruption of an IVS compartment boundary upon ablation of

Tbx5^+/Mef2cAHF^+ septal progenitors. We hypothesized that the *Tbx5^+/Mef2cAHF^+* septal

progenitors establish a compartment boundary at the IVS. Upon ablation of *Tbx5^+/Mef2cAHF^+*

septal progenitors, we predicted that if either lineage expanded into the other ventricular chamber,

then this would suggest lineage mixing from loss of compartment boundary integrity. To determine

if *Tbx5^+/Mef2cAHF^+* septal progenitors were essential for a compartment boundary at the IVS, we

tested the effects of ablating *Tbx5^+/Mef2cAHF^+* septal progenitors on lineage segregation. We

quantified the distributions of the *Mef2cAHF^+* or *Tbx5^+* lineages at E10.5 after tamoxifen

administration at E6.5, by quantifying fluorescence intensity as a linear profile across the

ventricular chambers by light sheet microscopy. In control (*Tbx5^{CreERT2/+};Mef2cAHF-*

DreERT2;Hip11^{+/+}) (n=2) embryos at E10.5, cells of the *Mef2cAHF^+* lineage were enriched in the

RV. Cells of the *Tbx5^+* lineage were highly enriched in the LV and rarely found in the RV, and

both lineages overlapped at the developing IVS. This data was consistent with observations of

lineage segregation at the IVS during later stages of heart development. However, in mutant

intersectional-*DTA* (*Tbx5^{CreERT2/+};Mef2cAHF-DreERT2;Hip11^{Intersectional-DTA176/+}*) (n=2) embryos at

E10.5, cells of the *Mef2cAHF*[+] lineage were less leftward in the corresponding region of targeted

cell ablation, suggesting a rightward retreat of the Mef2cAHF[+] lineage from ablation of

Tbx5[+]*/Mef2cAHF*[+] progenitors. In contrast, cells of the Tbx5[+] lineage in intersectional-*DTA*

mutants were observed more rightward, including in the RV. Therefore, we inferred that

Tbx5[+]*/Mef2cAHF*[+] progenitors were essential for preventing lineage mixing between the RV and

LV, by maintaining compartment boundary integrity at the developing IVS. This implies a

preferential regulation of first heart field derivatives over the adjacent second heart field.

<u>Heterozygous loss of *Tbx5* in the IVS leads to VSDs</u>. *TBX5* is a transcription factor gene that

causes VSDs in Holt-Oram Syndrome from haploinsufficiency in humans [20–22] and mice [23]. We

wondered if heterozygous loss of *Tbx5* in the IVS alone would cause VSDs. We used the

Mef2cAHF-Cre [24] in combination with a conditional deletion of *Tbx5* (*Tbx5*[flox]) [23], to conditionally

delete *Tbx5* heterozygously in a domain that overlaps with *Tbx5* expression at the IVS [25]. We

observed membranous VSDs in *Mef2cAHF-Cre*;*ROSA26*[mTmG/+];*Tbx5*[flox/+] mutant embryos (n=4/7)

compared to controls (n=0/3), at E14.5 (Supp. Figure 3A-C). This result contrasted with a previous

report that did not identify VSDs, albeit using a mixed genetic background [26]. This provided

evidence that appropriate *Tbx5* dosage in the IVS was essential for proper ventricular septation.

<u>Disturbed septal lineage contributions from reduced *Tbx5*.</u> As TBX5 dosage reduction globally or

only in the IVS results in VSDs, we reasoned that reducing TBX5 dosage may affect the regulation

of the *Tbx5*[+]*/Mef2cAHF*[+] septal progenitors and their subsequent lineage contributions. We

evaluated a reduction of *Tbx5* using a hypomorphic (*Tbx5*[CreERT2]) allele [22] in combination with a

conditional deletion of *Tbx5* (*Tbx5*[flox]) [23] at E6.5. This resulted in levels of *Tbx5* that are estimated

to be about 25% of wildtype in the *Tbx5*[+] lineage. In control (*Tbx5*[CreERT2/+];*Mef2cAHF-*

DreERT2;ROSA26[Ai66/Ai6]) embryos, the *Tbx5*[+]*/Mef2cAHF*[+] septal lineage overlapped at the

lineage front of the *Tbx5*[+] lineage in the IVS, from the base to the apex of the heart at the IV

groove, spanning from anterior to posterior of the heart in the IVS (Figure 4A-A", Supp. Figure

4A, B, Video 2). Among *Tbx5^CreERT2/flox* mutant (*Tbx5^CreERT2/flox;Mef2cAHF-*

DreERT2;ROSA26^Ai66/Ai6) embryos, we found that reduced TBX5 dosage at E6.5 caused a

spectrum of defects, frequently including VSDs, ASDs and AVC defects at E14.5 (Figure 4B-D'),

reminiscent of features of Holt-Oram Syndrome [20,21,27]. The normally organized distribution of the

tdTomato+ cells was highly irregular in the *Tbx5^CreERT2/flox* mutant hearts regardless of VSDs (Supp.

Figure 4A-H). Often, tdTomato+ cells were less apparent in the posterior IVS (Figure 4A', A", B',

B", C', C", D'), and tdTomato+ cells were nearly absent in a heart that displayed a severe AVC

defect and chamber hypoplasia (Figure 4D, D', Supp Figure 4G, H).

We used quantitative morphometry to assess the position and distribution of the

Tbx5+/Mef2cAHF+ septal lineage at the IVS. We tested this by quantifying linear profiles of the

labeled *Tbx5+/Mef2cAHF+* septal lineage in the heart. We found that both distribution and position

of the *Tbx5+/Mef2cAHF+* septal lineage was disturbed in *Tbx5^CreERT2/flox* mutant (n=3; control,

n=4)(Figure 4E-H, Supp. Figure 4I, J), including an increase of tdTomato+ cells leftward in the LV,

suggesting abnormal lineage position. As well, we observed a broader band of *Tbx5+/Mef2cAHF+*

septal lineage cells, suggesting that proper TBX5 dosage maintains integrity of the compartment

boundary marked by the *Tbx5+/Mef2cAHF+* lineage at the IVS (Figure 4G, H). We also observed

a reduction, or sometimes absence, of the labeled septal lineage among AVC cells, as well as

remnant atrial septal tissue in *Tbx5^CreERT2/flox* mutants (Figure 4D, D').

We further observed that cells of the *Tbx5+/Mef2cAHF+* lineage in the IVS were arranged

like a stack of coins, especially from the apex to the mid-septum (Figure 4A"). We wondered if

proper TBX5 dosage might be necessary for maintaining appropriate cell orientation in the IVS.

Therefore, cell alignment was determined and quantified by two metrics, average orientation

score or directional coherency. In *Tbx5^CreERT2/flox* mutant hearts, *Tbx5+/Mef2cAHF+* lineage

(tdTomato+) and *Tbx5+* lineage (ZsGreen+) cells were reduced in the dominant direction of cell

orientation and were more frequently in the orientation orthogonal to the dominant direction

(Figure 4S, Supp. Figure 4T). Further, both lineages demonstrated worse directional coherency

in $Tbx5^{CreERT2/flox}$ mutants (Figure 4T, Supp. Figure 4U), implicating TBX5 in proper cell orientation

in the IVS.

<u>*Tbx5*-sensitive genes encode guidance cues</u>. To find downstream effectors of Tbx5 that may

mediate the regulation of $Tbx5^+/Mef2cAHF^+$ lineage cells, we applied scRNAseq at E13.5, prior

to completion of ventricular septation. We micro-dissected the RV, LV and IVS+AVC in controls

and *Tbx5* mutants ($Tbx5^{CreERT2/flox}$*;Mef2cAHF-DreERT2;ROSA26*$^{Ai66/Ai6}$) (Figure 5A). These

samples were labeled at E6.5 for progenitors of the $Tbx5^+/Mef2cAHF^+$ lineage (*tdTomato*$^+$) and

Tbx5$^+$ lineage (*ZsGreen*$^+$). In samples from each cardiac tissue region, we detected clusters

enriched for *Tnnt2*$^+$ cardiomyocytes (CMs), *Postn*$^+$ fibroblasts, *Plvap*$^+$ endothelial cells,

Tbx18$^+$*/Wt1*$^+$ epicardial cells, *Hba-x*$^+$ red blood cells, and *C1qb*$^+$ white blood cells (Figure 5B,

Supp. Figure 5A-F).

In particular, *tdTomato*$^+$ cells were most abundant among clusters of *Tnnt2*$^+$

cardiomyocytes (CMs) of the IVS+AVC region (Figure 5C). However, we were unable to identify

a gene expression signature that was unique to the $Tbx5^+/Mef2cAHF^+$ lineage, beyond the lineage

marker itself, at this stage. Likewise, *ZsGreen*$^+$ cells were also enriched among clusters of *Tnnt2*$^+$

CMs in this region (Figure 5D), indicative of the *Tbx5*$^+$ lineage contribution to CMs in the region,

as well as a site of conditional reduction of *Tbx5* among *Tbx5* mutants.

We then focused our analysis on a subset of *Tnnt2*$^+$ enriched clusters from IVS+AV canal

regions. We identified genotype-enriched clusters that were abundant with cells from controls

(Cluster 5) or *Tbx5* mutants (Cluster 10) (Figure 5E, F). Control-enriched cluster 5 included genes

expressed in the AV canal region, encoding the Wnt co-receptor *Rspo3*, morphogen *Bmp2*, and

transcription factors *Tbx2* and *Tbx3*, suggesting that these genes are reduced in *Tbx5* mutants

(Figure 5G, H). As well, the SLIT-receptor *Robo1*, which is implicated in mouse and human VSDs

[28–30], and Netrin-receptor *Unc5b*, were also downregulated in *Tbx5* mutant cells (Figure 5G, H).

In the mutant-enriched cluster 10, genes typically expressed in ventricular trabeculae were

enriched (Figure 5G, H). These included natriuretic peptide *Nppa*, transcription factor *Cited1,* and

bone morphogenetic protein *Bmp10*, suggesting ectopic expression of these trabecular genes in

the IVS of *Tbx5* mutants. Intriguingly, guidance cues *Netrin1* (*Ntn1*) and *Slit2,* which are best

known for axonal development or vasculogenesis [31], were also dysregulated (Figure 5G, H).

We examined differential gene expression between controls or *Tbx5* mutants in *tdTomato*⁺

(Tbx5⁺/Mef2cAHF⁺ septal lineage) or *ZsGreen*⁺ (Tbx5⁺ lineage) CMs. We found many

differentially expressed genes that overlapped among comparisons of *tdTomato*⁺ or *ZsGreen*⁺

clusters, including *Nppa*, *Cited1,* and *Slit2*, consistent with findings from cluster-to-cluster

comparisons (Supp. Figure 5G, H).

We used orthogonal assays to validate candidate TBX5-sensitive genes. By fluorescence

in situ hybridization in formalin fixed paraffin embedded (FFPE) sections, we observed a reduction

of gene expression of *Bmp2, Robo1 and Unc5b* in the AV canal region of mutant hearts when

compared to controls. (Supp. Figure S6M-X'). As well, we detected expansion of *Nppa* and *Slit2*

into the IVS and compact layer of mutant hearts, both of which are normally expressed only in the

trabecular regions of both ventricles of control hearts (Figure 5I-L'). Moreover, we discovered

Ntn1 expression enriched in LV trabeculae and in a gradient across the control IVS (Figure 5M,

M'), consistent with immunostaining of NTN1 in the heart at E14.5 (Supp. Figure 6Y). Notably,

Ntn1 was enriched in the LV earlier at E11.5 during ventricular septation (Supp. Figure 6Z).

However, in *Tbx5* mutants, the *Ntn1* gradient was flattened and expanded in the IVS by reduced

Tbx5 dosage (Figure 5N, N'). Interestingly, we observed abnormal *Nppa*, *Slit2* and *Ntn1* in *Tbx5*

mutants with or without VSDs, suggesting that this dysregulated gene expression was not

secondary to VSDs (Supp Figure 6A-L'). In addition, analysis of ChIP-seq of TBX5 from

embryonic mouse hearts [32] showed TBX5 occupancy at promoters of *Nppa, Slit2*, and *Ntn1* (Supp

Figure 7A-C), suggesting these guidance genes are likely to be direct targets of TBX5 in the heart.

Tbx5-Slit2 and *Tbx5-Ntn1* genetic interactions. We evaluated if genetic interactions existed

between *Tbx5* and *Slit2* or *Ntn1*. As *Tbx5* heterozygous mutant (*Tbx5del/+*) [23] mouse embryos

display muscular or membranous VSDs, we wondered if heterozygous *Slit2* or *Ntn1* loss of

function (LOF) mutant embryos might alter *Tbx5*-dependent phenotypes. We mated *Tbx5del/+* mice

with either *Slit2* (*Slit2+/-*) or *Ntn1* (*Ntn1beta-actin-Cre/+* referred here subsequently as *Ntn1+/-* [33,34])

heterozygous LOF mice. In *Tbx5del/+* embryos at E14.5, we observed membranous (n=5/8) or

muscular (n=3/8) VSDs by histology (Supp. Figure 8B, E), consistent with previous reports [23].

Among *Slit2+/-* embryos at E14.5, we observed a membranous VSD (n=1/8) (Supp. Figure 8C, E).

This contrasted with a previous report that did not detect any VSDs in *Slit2+/-* albeit using a

different mutant *Slit2* allele [28]. In *Tbx5del/+*;*Slit2+/-* compound heterozygous embryos, we observed

an estimated decrease in the incidence of membranous VSDs (n=5/8) (Supp. Figure 8D, E), which

approached significance (log OR -2.5; $p < 0.09$ in a Generalized Linear Model).

We next tested if there was a genetic interaction between *Tbx5* and *Ntn1* (Supp. Figure

8F-J). In *Ntn1+/-* embryos, we observed a membranous (n=1/9) or a muscular (n=1/9) VSD. In

Tbx5del/+;*Ntn1+/-* compound heterozygous embryos, we observed a significant decrease of

membranous VSDs (n=1/8; log OR -5.3; $p < 1 \times 10^{-3}$ by a Generalized Linear Model), but not

significant changes to prevalence of muscular VSDs (n=4/8; log OR 0.12; $p < 0.93$), consistent with

a genetic interaction between *Tbx5* and *Ntn1* for ventricular septation.

We wondered if there were other quantitative histologic differences, in addition to the

changes in incidence of VSD types. To quantify morphometry from histology, we leveraged a

deep learning algorithm, known a CODA [35], to detect embryonic mouse heart components from

hematoxylin and eosin (H&E) sections. We generated three-dimensional (3-D) tissue

reconstructions, to visualize and quantify anatomic structures and CHDs, including membranous

and muscular VSDs, at cellular resolution (Figure 6I-K). Using this machine learning method, we

discovered that the cell density of the IVS, which we termed IVS fill, was increased in *Tbx5del/+*;

Ntn1+/- embryos compared to wildtype ($p < 0.05$ by Fisher's exact test), while classification of the

IVS as trabecular was significantly reduced in $Tbx5^{del/+}$;$Ntn1^{+/-}$ embryos (p<0.05 by Fisher's exact

test) (Supp. Figure 8M). In addition, we estimated the minimal area for each membranous or

muscular VSD, when present (Supp. Figure 8O, P). In $Tbx5^{del/+}$;$Slit2^{+/-}$ compound mutants, we

detected statistically significant levocardia (p<0.001 by Fisher's exact test), as determined by the

axis of the heart compared to the spine and sternum (Supp. Figure 8Q). This uncovered a genetic

interaction between $Tbx5$ and $Slit2$ related to heart position, demonstrating that machine learning-

based morphometry can quantify unexpected anatomic findings.

<u>$Slit2$ and $Ntn1$ for ventricular septation.</u> To determine if $Slit2$ or $Ntn1$ are essential for ventricular

septation, we evaluated homozygous LOF mouse mutants for $Slit2$ or $Ntn1$ for VSDs. In $Slit2^{-/-}$

embryos at E14.5, we observed that homozygous loss of $Slit2$ displayed complete penetrance for

VSDs, including membranous (n=4/4; log OR 5.0, p<0.00) or muscular VSD (n=1/4) (Figure 6A-

D), demonstrating a far higher incidence than a previous report using an alternative $Slit2$ LOF

allele [28]. Further, we detected a statistically significant non-compacted IVS as determined by

reduced IVS fill and thinning of the right ventricular compact layer (Supp. Figure 9E, H). Likewise,

$Ntn1^{-/-}$ embryos displayed membranous VSDs (n=6/7), an apparent thinning of the ventricular

compact layer and a non-compacted IVS, although these observations were not statistically

significant (Supp. Figure 9). Collectively, this morphologic evidence implicates $Slit2$ and $Ntn1$ in

proper ventricular septation during heart development.

As $Slit2$ and $Ntn1$ were necessary for ventricular septation, we asked if loss of $Slit2$ or $Ntn1$ might

disturb distribution of the $Tbx5^{+}$/$Mef2cAHF^{+}$ lineage in the IVS. In homozygous $Slit2$ LOF mutants

with the lineage reporter ($Tbx5^{CreERT2/+}$;$Mef2cAHF$-$DreERT2$;$ROSA26^{Ai66/+}$;$Slit2^{-/-}$) (n=4), we

observed a statistically-significant broadened distribution of the $Tbx5^{+}$/$Mef2cAHF^{+}$ lineage

compared to controls (n=2) (Figure 6N-Q), consistent with lineage mixing from boundary

disruption and implicating a role for SLIT2 in maintaining boundary integrity. Conversely,

homozygous *Ntn1* LOF mutants with the lineage reporter (*Tbx5^CreERT2/+^;Mef2cAHF-DreERT2*;

*ROSA26^Ai66/+^;Ntn1^-/-^*) (n=6) demonstrated leftward displacement of the Tbx5^+^/Mef2cAHF^+^ lineage

compared to controls (n=4) (Figure 6R-U), implying that NTN1 signaling precisely positions the

compartment boundary at the IVS. Taken together, this evidence suggested that SLIT2- and

NTN1-signaling, as part of a Tbx5-dependent pathway, regulates the position and integrity of the

compartment boundary labeled by the *Tbx5^+^/Mef2cAHF^+^* lineage.

343
Discussion.

We show that *Tbx5^+^/Mef2cAHF^+^* progenitors prefigure a compartment boundary that becomes

located at the junction between the left and right sides of the IVS, providing a cellular framework

during development for heart patterning. Our study underscores a fundamental principle that early

developmental events preconfigure structure and function of the heart and are susceptible to

genetic risks that cause CHDs. Further, our results demonstrate developmental regulation of a

compartment boundary by a disease-associated TF and that aberrant tissue patterning from

boundary disturbances may be an etiology of birth defects. This study reiterates the importance

of compartment boundaries in tissue patterning, provides evidence for compartment boundary

disruptions as an etiology of birth defects, and adds to the few examples of compartment

boundaries that exist in mammals.

We surmise that the rarity of compartment boundaries in mammals likely stems from a few

reasons. The discovery of a compartment boundary was first made by studying the fate of marked

mosaic cells and their clonal segregation in the wing disk of the fly embryo [36]. Subsequent

examples of compartment boundaries were identified by broad genetic screens in Drosophila [37].

In mammals, compartment boundaries were discovered by lineage analysis informed by another

organism [16] or serendipitously [13], as we have done here. Compartment boundaries have been

defined by several criteria. First, a compartment boundary segregates juxtaposed cell populations

such as distinct cell lineages to restrict cell intermingling, and ablation of the boundary leads to

cell mixing. Second, the expression domain of a selector gene corresponds with a compartment

domain, loss of the selector gene eliminates the identity in this territory, and ectopic expression

of the selector induces this identity. Third, a selector gene establishes a signaling center to

maintain the compartments and boundary. Our results satisfy all of these criteria: Ablation of the

$Tbx5^+/Mef2cAHF^+$ lineage disrupts segregation of $Tbx5^+$ lineage cells in the LV from the $Tbx5^-$

lineage in the RV of the developing heart. Reduced $Tbx5$ expression impairs the integrity of the

boundary and its patterning, and in previous work misexpression of TBX5 eliminates IVS

formation [25]. Third, TBX5 patterns signaling molecules that are important for morphologies that

impact the boundary. In this context, we postulate that $Tbx5$ may function as a candidate selector

gene. Further experiments will be needed to formally ascertain this.

In this context, we identify a $Tbx5$-dependent pathway that regulates $Slit2$ and $Ntn1$.

Normally, TBX5 represses their ectopic expression in the IVS and compact layer. Both SLIT2 and

NTN1 are guidance or adhesive cues in several developmental contexts. Here, we provide

evidence that $Slit2$ and $Ntn1$ are direct targets of TBX5, and that $Tbx5$ and $Slit2$ or $Ntn1$ genetically

interact. Further, we found that $Slit2$ and $Ntn1$ are essential for proper ventricular septation, as

well as compartment boundary integrity and positioning, respectively. Like TBX5, proper

ventricular septation may be sensitive to imbalanced dosage of SLIT2 or NTN1. As $Ntn1$

expression is enriched in the LV and left-side of the IVS in the $Tbx5^+$ compartment, we propose

a working hypothesis that NTN1 functions as a TBX5-dependent signaling center for compartment

boundary regulation.

The spatiotemporal position of the boundary is at the crux of cardiac septation and, when

disrupted, spans sites for congenital cardiac anomalies. The sensitivity of the $Tbx5^+/Mef2cAHF^+$

lineage-labeled boundary to genetic perturbations, specifically $Tbx5$ deficiency, is of significance

in the context of congenital heart disease. Such deficiency selectively compromises the

boundary's integrity and position, along with disturbances to cell orientation, leading to VSDs or

AVSDs. This observation suggests that genetic pathways for patterning, like the

Tbx5$^+$/Mef2cAHF$^+$ lineage-labeled boundary, can be exceptionally susceptible to genetic

disruptions. This provides a potential "beacon" for understanding the cellular basis of septation

defects. Whether additional genetic perturbations that cause VSDs or AVSDs disrupt the

compartment boundary remains to be determined. Likewise, it will be interesting to evaluate if the

Tbx5$^+$/Mef2cAHF$^+$ lineage-labeled boundary corresponds with abnormal shifts in IVS position, like

those observed in the endothelial loss of *Hand2*, which displays rightward shifted IVS and double

inlet left ventricle (DILV) or two IVS primordia [38,39].

Broadly, our research demonstrates the delicate interplay of progenitor cell behavior, gene

expression and compartment boundary formation that determines heart development and genetic

susceptibility that increases the risk of CHDs. This evidence further underscores the significance

of early developmental stages for the ontogeny of some CHDs. The implications of these

discoveries likely extend beyond cardiac morphogenesis, as these findings offer insights for the

roles of compartment boundaries during mammalian development and uncover a vulnerability for

organ patterning from genetic disturbances.

METHODS.

Mouse lines. All mouse protocols were approved by the Institutional Animal Care and Use

Committee at UCSF. Mice were housed in a barrier animal facility with standard husbandry

conditions at the Gladstone Institutes. Mice of *Tbx5^{CreERT2IRES2xFLAG}* (abbreviated here as

Tbx5^{CreERT2}) and *Mef2cAHF-DreERT2* [6], *Tbx5$^{del/+}$* and *Tbx5$^{flox/+}$* [23], *ROSA26^{Ai66}* and *ROSA26^{Ai6}* [18]

were described previously. *Mef2cAHF-Cre$^{/+}$* mice [24] were obtained from Brian Black. *Slit2$^{+/-}$* mice

(MMRC, Strain 065588-UCD, donated by Kent Lloyd) were generated by CRISPR/Cas9-

targeted constitutive deletion of exon 8 and flanking splicing regions of *Slit2*. *Ntn1$^{+/-}$* mice

were derived from matings of *Ntn1* floxed mice (*Ntn1$^{flox/+}$*; Jackson Laboratory #028038) [40]

to beta-actin-Cre [34], which were obtained from Gail Martin. All mouse strains were maintained

in the C57BL6/J background (Jackson Laboratory #664), except for *Tbx5$^{del/+}$*, which was

maintained in Black Swiss (Charles River, Strain Code 492), and *Slit2*[+/-] which was maintained in

C57BL6/N (Jackson Laboratory, #005304). Both male and female embryos were collected from

timed matings and used at random for experiments. Tamoxifen (Sigma-Adrich; T5648) is

suspended in sesame oil at 10mg/mL. The injected dose is 100ug/1 gram of body weight.

Cloning and generation of mouse lines. We generated an attenuated diptheria toxin (DTA176)

transgenic knock-in mouse under the control of the dual-recombinase intersectional cassette.

DTA176 encodes an attenuated form of toxic fragment A from diptheria toxin [19], which requires

~100-200 molecules for cell killing. This strategy was an effort to reduce potential problems of

leaky DTA expression prior to recombinase-mediated activation. Briefly, DTA176 [19] was cloned

downstream of ROX-STOP-ROX and LOX-STOP-LOX sites to create a Dre- and Cre-dependent

expression construct, which was derived from the Dre- and Cre-dependent tdTomato expression

cassette of the Ai66 (RCRL-tdT) targeting vector [18]. After the tdTomato cassette was removed

using MluI sites and replaced by DTA176, the construct was cloned into the TARGATT targeting

construct pBT346.3 (Applied Stem Cells) to target the transcriptionally inactive *Hip11* locus [41]

using PacI and SmaI restriction enzyme sites. The final construct was verified using restriction

digestion and Sanger sequencing. DNA was purified and injected along with mRNA for the Phi31o

transposase according to manufacturer's protocol to generate intersectional-DTA mutant

($Hip11^{Rox\text{-}STOP\text{-}Rox\text{-}Lox\text{-}STOP\text{-}Lox\text{-}DTA176}$) mice. To generate Ai66b mice ($ROSA26^{Ai66b}$), which is a Dre-

dependent tdTomato reporter, the LOX-STOP-LOX sites were removed from the Ai66 (RCRL-

tdT) targeting vector [18], and then the transgene cassette was inserted into endogenous genomic

loci via homologous recombination, as previously described [42].

Whole mount embryo and heart immunofluorescence labeling. Embryos were dissected from

timed pregnant dams, including removal of yolk sac for genotyping, as previously described [6].

Embryos up to E10.5 were fixed for 4 hours in 4% PFA at room temperature, then incubated with

gentle rocking in clearing solution (8% SDS in 0.2M borate buffer) at 37°C for 1-3 hours until clear.

E13.5 or E14.5 hearts were micro-dissected from freshly harvested embryos in PBS with heparin

(10 Units/mL) (Sigma #H3393) or incubated in 1X RBC lysis solution (Roche #11814389001) for

10 minutes at room temperature, fixed at room temperature in 4% PFA for one hour, and

incubated with gentle rocking in clearing solution at 37°C for 2-4 days until clear. Specimens were

permeabilized and blocked for two hours to overnight with gentle rocking at 37°C in blocking buffer

(PBS containing 5% normal donkey serum and 0.8% to 1.5% Triton X-100, dependent on

embryo/heart stage). Specimens were labeled with primary antibody (list below) in blocking buffer

with gentle rocking at 37°C, either overnight for embryos up to E10.5, or 5 days for E13.5 or E14.5

hearts. For long incubations, antibody solution was replaced halfway. Following three washes in

blocking buffer, each 45 minutes except the third overnight for E13.5 and E14.5 hearts, specimens

were incubated with secondary antibody (raised in donkey; AF488-, Dy405-, Cy3, or AF647-

conjugated; used at 1:750; Jackson ImmunoResearch) and DAPI in blocking buffer with gentle

rocking at 37°C, 2-3 hours for embryos up to E10.5, or 5 days for E13.5 or E14.5 hearts. Following

three washes in PBS with gentle rocking at 37°C, samples were stored (up to overnight) in PBS

with 0.2% sodium azide. Primary antibodies used were: tdTomato (rabbit, Rockland 600-401-379,

1:1000), MEF2c (sheep, R&D AF6786, 1:250), TNNT2 (mouse, Thermo MS-295-P, 1:500).

Strong ZsGreen fluorescence persisted after clearing and did not require immunolabeling.

Lightsheet Image Acquisition. Images were acquired by lightsheet microscopy, as described

previously [43]. Brielfy, specimens were warmed and transferred to 2% low melting point agarose

(Fisher BP165-25) in PBS at 37°C, then embedded in glass capillary tubes with paired pistons

(Sigma Z328510 paired with BR701938, or Sigma Z328502 paired with BR701934) for embryos,

or tip-truncated 1mL syringes (Becton Dickinson) for hearts. After the gel solidified, the capillaries

or syringes were suspended specimen-down from 14mL polystyrene tubes sealed with Parafilm.

Columns containing the specimen were partially extended into an ample volume optical clearing

solution (OCS, EasyIndex EI-Z1001, LifeCanvas) overnight. With immersion in OCS, specimens

were rotated and imaged from multi-view whole-volume approach on a Light Sheet Z.1

microscope (Carl Zeiss Imaging). One of two objective setups was used: EC Plan Neofluar

5X/0.16 with 5X/0.1 pair (hearts), or Clr Plan Neofluar 20X/1.0 paired with 10X/0.2 clearing pair

(embryos). Z-stacks were collected at each view angle at the optimal slice thickness

determined by Zeiss' Zen software, ranging from 1.42 to 4.95 µm.

Image Dataset Preprocessing. Imaging data was processed, as previously described [43]. All data

processing was performed on an 8-core x86-64 desktop PC with 64GB RAM, running Kubuntu

20.04 LTS, principally using Fiji software [44]. Acquired light sheet image stacks were subjected to

single-view deconvolution using regularized generalized Tikhonov filtering (Parallel Spectral

Deconvolution, https://imagej.net/plugins/parallel-spectral-deconvolution) following theoretical

PSF calculation using the intersection of Gaussian z-plane illumination with Gibson-Lanni

widefield epifluorescence detection patterns [45]. Deconvolved stacks were co-registered, and

image volumes were generated for each desired orientation, for each specimen, using Content

Based Fusion in BigStitcher [46]. TrackMate [47] and ImageJ 3-D viewer were used for segmentation

and visualization of fused volumes.

Image Volume Quantification. Extensive whole-volume quantifications of imaged E14.5 hearts

were performed using Fiji and associated plugins, with analysis conducted with R. For assessing

the location of fluorescence signal, volumes were downsampled, and 2-dimensional regions of

interest (ROIs) corresponding to anatomic structures were manually drawn on each Z-slice, using

the DAPI channel. Integrated intensity for each fluorescence channel was programatically

measured for each ROI, and counts were normalized to DAPI signal and pooled by anatomic

region. For whole-heart quantifications, counts were further normalized to combined Ai6 plus Ai66

signal, then averaged across each anatomic region. For orientation assessments, a multi-channel

panel of 174 full resolution four-chamber image excerpts, evenly distributed between apical-

posterior, apical-anterior, posterior-basal, and posterior-apical IVS, were split by channel

(tdTomato or ZsGreen) and blindly sampled. Using automated batch processing, these samples

were subjected to background filtering, contrast enhancement, and measurements of orientation

distribution and dominant direction (OrientationJ plugin, https://github.com/Biomedical-Imaging-

Group/OrientationJ), and of directionality (Directionality plugin,

https://github.com/fiji/Directionality). Orientation score distributions were rotated so that dominant

direction was set to 0°, with average score (across the panel of images) was plotted as a function

of direction, and analyzed by Watson's two-sample test. Individual scores surrounding 0° and 90°

were pooled and compared with the Wilcoxon signed-rank test. Directionality coherence scores

(across the panel of images) were plotted by genotype and compared with the Wilcoxon signed-

rank test.

505
Quantification of right-left position of lineages. For each E10.5 heart analyzed, five

representative coronal optical sections were selected, approximately 20-35 μm apart. For each

optical section, the DAPI channel was used to draw five linear profiles spanning the ventricular

chambers from right to left. Linear profiles were drawn such that the "0" position corresponded to

the midline, identified by the interventricular groove. Fluorescence intensity of each channel

(green=$Tbx5^+$ lineage, magenta=$Mef2cAHF$+ lineage) was measured along each linear profile,

and intensity values were normalized to the maximum value along each profile. Profiles for each

channel were grouped by experimental condition (WT or DTA), and then the average ± SEM

normalized fluorescence intensity profiles were calculated and plotted. Welch's two-sample t-test

was used at each position along the right-left axis to determine regions of significant difference in

fluorescence intensity of each channel between experimental groups, indicating a change in the

distribution of cells of that lineage.

For each E13.5-E14.0 heart analyzed, maximum z-projections were generated from

optical sections spanning every 50 µm from the anterior to posterior of the heart. For each

maximum z-projection, the DAPI and red or magenta (*Tbx5$^+$/Mef2cAHF$^+$* intersectional lineage)

channels were split and background subtraction was performed on the red or magenta channel

using a 50-pixel rolling ball radius. The DAPI channel was used to draw three linear profiles (basal,

middle, apical) spanning the ventricular chambers from right to left. Linear profiles were drawn

such that the "0" position corresponded to the middle of the IVS. Fluorescence intensity of the red

or magenta channel was measured along each linear profile, and intensity values were

normalized to the maximum value along each profile. Profiles were grouped by experimental

condition (*Tbx5$^{CreERT2/+}$* or *Tbx5$^{CreERT2/flox}$*) or (e.g. WT, *Ntn1$^{+/-}$*, *Ntn1$^{-/-}$*) and then the average ± SEM

normalized fluorescence intensity profiles were calculated and plotted. Two sample F-test and

Welch's t-test were used to analyze the distribution of *Tbx5$^+$/Mef2cAHF$^+$* intersectional lineage

cells.

Cell Harvesting for Single Cell RNA Sequencing. Samples for single cell RNA sequencing

were collected from three independent litters at E13.5. The heart was microdissected to obtain

the interventricular septum (IVS), left ventricular (LV) and right ventricular (RV) regions. Control

samples   were   *Tbx5$^{CreERT2/+}$;Mef2cAHF-DreERT2;ROSA26$^{Ai66/Ai6}$*   (n=3)   and

Tbx5$^{CreERT2/+}$;Mef2cAHF-DreERT2;ROSA26$^{Ai66/+}$ (n=1) for LV, RV and IVS each. *Tbx5* mutant

samples were *Tbx5$^{CreERT2/flox}$;Mef2cAHF-DreERT2;ROSA26$^{Ai66/Ai6}$* for RV and LV (n=3 each) and

IVS (n=2).

Each microdissected tissue was singularized with TrypLE Express (Life Technologies,

Cat# 12604-013) at 37C and quenched with 1% FBS in PBS. The single cell suspension was then

filtered through a cell strainer cap (Corning, Cat# 352235) and centrifuged at 300g for 5 minutes.

The pellet was resuspended in 1% FBS in PBS and counted using an automated cell counter. A

30µL aliquot of the cell suspension was used to generate single cell droplet libraries with the

Chromium Next GEM Single Cell 5' Library & Gel Bead Kit v1.1, according to manufacturer's

instructions (10X Genomics). After KAPA qPCR quantification, a shallow sequencing run was

performed on a NextSeq 500 (Illumina) prior to deep sequencing on a NovaSeq S4 (Illumina). For

control IVS+AVC, at E13.5, 3 samples were *Tbx5$^{CreERT2/+}$;Mef2cAHF-DreERT2;ROSA26$^{Ai66/Ai6}$*

Data Processing Using Cellranger. All datasets were processed using Cellranger 2.0.2. FASTQ

files were generated using the mkfastq function. Reads were aligned to a custom mm9 reference

(version 1.2.0) containing tdTomato and ZsGreen reporter genes. Cellranger aggr was used to

aggregate individual libraries after read depth normalization.

Seurat Analysis. Outputs from the Cellranger pipeline were analyzed using the Seurat v3 [48–50].

Datasets from the IVS, LV and RV were analyzed separately. Cells with 10,000-50,000 UMIs and

1500-7250 genes were retained. Data was normalized using the NormalizeData function and

scaled using ScaleData, while also regressing unwanted sources of variation, such as differences

in the number of UMIs, number of genes, percentage of mitochondrial reads and differences

between G2M and S phase scores. Principal component analysis (PCA) was performed using the

most highly variable genes. Cells were then clustered based on the top 30 principal components

and visualized on a Uniform Manifold Approximation and Projection (UMAP) [51]. A clustering

resolution that separated cells by major cell types was chosen. Next, *Tnnt2*$^+$ clusters from the IVS

dataset were extracted and re-clustered, similar to the parent dataset above. A clustering

resolution was chosen based on separation of genotypes. Differential gene expression tests

between clusters or between *ZsGreen*$^+$ and *ZsGreen*$^-$ or *tdTomato*$^+$ and *tdTomato*$^-$ cells were

performed using the FindMarkers function with default parameters. Selected differentially

expressed genes with an adjusted p-value less than 0.05 from the Wilcoxon rank sum test were

displayed using the Dotplot or FeaturePlot functions.

Fluorescent *In Situ* Hybridization. E11.5 or E14.5 hearts were fixed with 4% paraformaldehyde

or 10% formalin overnight at 4C, embedded in paraffin and then sectioned for a transverse or

four-chambered view on slides. *In situ* hybridization was performed on sections using the

RNAscope Multiplex Fluorescent v2 Assay kit (Advanced Cell Diagnostics, Cat# 323100). Briefly,

sections were deparaffinized in xylene and 100% ethanol, treated with hydrogen peroxide for 10

minutes and boiled in target retrieval buffer for 10 minutes. A hydrophobic barrier was drawn

around each section using an Immedge pen (Vector Laboratories, Cat# H-4000), and slides were

allowed to dry overnight. The following day, sections were treated with Protease Plus for 30

minutes, followed by hybridization with probes for 2 hours at 40C. Probes used were *Mm-Tnnt2-*

C4 (Cat# 418681-C4), *Mm-Ntn1* (Cat# 407621), *Mm-Slit2* (Cat# 449691), *Mm-Nppa-C3* (Cat#

418691-C3), *Mm-Bmp2-C2* (Cat# 406661-C2), *Mm-Robo1* (Cat# 475951) and *Mm-Unc5b* (Cat#

482481). Amplification steps were carried out according to manufacturer's instructions. Opal dyes

520, 570, 620 and 690 (Akoya Biosciences, Cat# FP1487001KT, FP1487001KT, FP1487001KT

and FP1487001KT) were used at 1:750 dilution. After staining with DAPI, 1-2 drops of ProLong

Gold Antifade Mountant (ThermoFisher Scientific, Cat# P36930) were placed on the slides and

mounted with a coverslip. Slides were stored overnight at 4C and were imaged at 10X

magnification on the Olympus FV3000RS. Multi-Area Time Lapse (MATL) images were captured

and then stitched together using the Olympus FV3000RS software. Stitched images were

analyzed using ImageJ OlympusViewer plugin.

Immunohistochemistry. Cryopreserved slides were thawed from -80C at RT and washed in

0.1% Triton X-100 in PBS (PBST) for 3 x 5 min. Antigen-retrieval was performed by boiling

slides in 10mM sodium citrate buffer (pH 6.0) for 10 minutes and then washed in H2O for 3 x 5

min. Blocking was performed using 5% donkey serum in 0.1% PBST at RT for 1-2hr. Anti-

Netrin-1 antibody 1:500 (R&D Systems AF1109) in 1% BSA + 0.1% PBST was incubated at 4C

overnight. After washing 3 x 5 min in 0.1% PBST, slides incubated in Alexa Fluor 594 (1:300) in

1% BSA + 0.1% PBST at RT for 1 hr. Slides were then washed 3 x 5 min in 0.1% PBST and

incubated in 1:1000 DAPI in 0.1% PBST at RT for 5-10 minutes. Finally, slides washed 3 x 5

min in 0.1% PBST, glass coverslips were mounted using Prolong Gold antifade mounting

medium and stored at 4C.

Machine learning-based 3-D reconstruction from histology and quantitative micrometry.

Tissues were formalin-fixed, paraffin embedded (FFPE) and serially sectioned at a standard

thickness of 5 μm to exhaustively collect the heart tissue of each fetal mouse. Serial sections

were stained with hematoxylin and eosin (H&E) and digitized at 20x magnification. CODA, a

technique to 3-D reconstruct serial tissue images, was used to map the microanatomy of the

hearts [35,52]. The various steps of CODA can be broken down into image registration, nuclear

detection, and tissue labelling. Nonlinear image registration was used to align the serial images.

Cellular coordinates were generated through color deconvolution and detection of two-

dimensional intensity peaks in the hematoxylin channel of the images. A deep learning semantic

segmentation algorithm was trained to label the anatomical structures of the mice at 1 μm

resolution. To train the model, manual annotations of six structures were generated in 55

histological images corresponding to the different large structures present in the tissue: bones,

spinal cord, lung, liver, stroma, heart, and non-tissue pixels in the images. A second deep learning

model was trained to subclassify the regions of interest within the heart: compact myocardium,

trabeculae, IVS, AV canal and atria. Fifty images from the manual annotation datasets were used

for model training, with five images held out for independent testing of model accuracy. The

models were deemed acceptable when they reached a minimum per-class precision and recall of

>85%. Using the trained model, independent images were segmented and aligned to generate

digital 3-D datasets.

The reconstructed volumes enabled microanatomical quantification of anatomical

properties. IVS fill, or the solidity of the muscle layer in the IVS, was calculated by dividing the

number of dark (<200 in RBG color space) pixels within the IVS, normalized by all pixels (including

empty space) within the IVS. Percent trabeculation within the IVS was calculated by first isolating

the tissue classified as trabeculae that was spatially attached to the IVS, then normalizing this

number by the combined volume of trabeculae attached to IVS and the volume of the IVS itself.

The compact myocardium thickness was calculated by measuring the thickness of the compact

layer at the bottom 20% of the ventricles. To measure the thickness on the left and right ventricles,

the two sides of the heart were manually segmented in 3-D space. VSDs were identified by

calculating regions where the empty space of the left ventricle contacted the empty space of the

right ventricle. These defects were evaluated and manually sorted into membranous or muscular

sub-types. The areas of each defect were then manually calculated through annotation on serial

histological images representing the shortest path to close the defect. These lines were summed

across all images containing the defect multiplied by the thickness of the histological slides to

determine the area of the hole.

Quantification and Statistical Analysis. For statistical analysis of genetic interactions, the

number of animals with a given defect (muscular VSD or membranous VSD) was modeled in a

Generalized Linear Model assuming a binomial probability distribution for the observed counts.

This model included main effects (in terms of log odds ratios) capturing the number of mutant

alleles for *Tbx5* (0 or 1, representing the WT or the Het genotype, respectively) and the number

of mutant alleles for *Slit2* or *Ntn1* (0,1 or 2: representing the WT, Het or Hom genotype,

respectively) and an interaction between these effects. Given the relatively small number of

animals in this experiment, bias-reduced estimates were made using the brglm2 package

{Kosmidis.2020} in R <https://www.R-project.org/>.

References.

1. de la Cruz, M. V., Sánchez-Gómez, C. & Palomino, M. A. The primitive cardiac regions in the
straight tube heart (Stage 9) and their anatomical expression in the mature heart: An
experimental study in the chick embryo. *J. Anat.* **165**, 121–31 (1989).

2. de la Cruz, M. V., Castillo, M. M., G., L. V., Valencia, A. & Moreno-Rodriguez, R. A. Primitive
interventricular septum, its primordium, and its contribution in the definitive interventricular
septum: In vivo labelling study in the chick embryo heart. *Anatomical Rec* **247**, 512–520 (1997).

3. Wessels, A. *et al.* Spatial distribution of "tissue-specific" antigens in the developing human
heart and skeletal muscle. I. An immunohistochemical analysis of creatine kinase isoenzyme
expression patterns. *Anatomical Rec* **228**, 163–176 (1990).

4. Wessels, A. *et al.* The Development of the Atrioventricular Junction in the Human Heart. *Circ
Res* **78**, 110–117 (1996).

5. Lamers, W. H. *et al.* New findings concerning ventricular septation in the human heart.
Implications for maldevelopment. *Circulation* **86**, 1194–1205 (2018).

6. Devine, W. P., Wythe, J. D., George, M., Koshiba-Takeuchi, K. & Bruneau, B. G. Early
patterning and specification of cardiac progenitors in gastrulating mesoderm. *eLife* **3**, 508
(2014).

7. Später, D. *et al.* A HCN4+ cardiomyogenic progenitor derived from the first heart field and
human pluripotent stem cells. *Nat Cell Biol* **15**, 1–10 (2013).

8. Lescroart, F. *et al.* Early lineage restriction in temporally distinct populations of Mesp1
progenitors during mammalian heart development. *Nat Cell Biol* **16**, 829–840 (2014).

9. Meilhac, S. M., Esner, M., Kelly, R. G., Nicolas, J.-F. & Buckingham, M. E. The clonal origin
of myocardial cells in different regions of the embryonic mouse heart. *Dev Cell* **6**, 685–698
(2004).

10. Kiecker, C. & Lumsden, A. Compartments and their boundaries in vertebrate brain
development. *Nat Rev Neurosci* **6**, 553–564 (2005).

11. Dahmann, C., Basler, K., Dahmann, C. & Basler, K. Compartment boundaries: at the edge
of development. *Trends Genet* **15**, 320–326 (1999).

12. Wang, J. & Dahmann, C. Establishing compartment boundaries in Drosophila wing imaginal
discs: An interplay between selector genes, signaling pathways and cell mechanics. *Semin Cell
Dev Biol* **107**, 161–169 (2020).

13. Kimmel, R. A. *et al.* Two lineage boundaries coordinate vertebrate apical ectodermal ridge
formation. *Genes Dev.* **14**, 1377–1389 (2000).

14. Arques, C. G., Doohan, R., Sharpe, J. & Torres, M. Cell tracing reveals a dorsoventral
lineage restriction plane in the mouse limb bud mesenchyme. *Development* **134**, 3713–3722
(2007).

15. Qiu, Q., Chen, H. & Johnson, R. L. Lmx1b-expressing cells in the mouse limb bud define a
dorsal mesenchymal lineage compartment. *Genesis* **47**, 224–233 (2009).

16. Zervas, M., Millet, S., Ahn, S. & Joyner, A. L. Cell Behaviors and Genetic Lineages of the
Mesencephalon and Rhombomere 1. *Neuron* **43**, 345–357 (2004).

17. Franco, D. *et al.* Left and right ventricular contributions to the formation of the
interventricular septum in the mouse heart. *Dev Biol* **294**, 366–375 (2006).

18. Madisen, L. *et al.* Transgenic Mice for Intersectional Targeting of Neural Sensors and
Effectors with High Specificity and Performance. *Neuron* **85**, 942–958 (2015).

19. Wu, S., Wu, Y. & Capecchi, M. R. Motoneurons and oligodendrocytes are sequentially
generated from neural stem cells but do not appear to share common lineage-restricted
progenitors in vivo. *Development* **133**, 581–590 (2006).

20. Basson, C. T. *et al.* Mutations in human TBX5 [corrected] cause limb and cardiac
malformation in Holt-Oram syndrome. *Nat Genet* **15**, 30–35 (1997).

21. Li, Q. Y. *et al.* Holt-Oram syndrome is caused by mutations in TBX5, a member of the
Brachyury (T) gene family. *Nat Genet* **15**, 21–29 (1997).

22. Kathiriya, I. S. *et al.* Modeling Human TBX5 Haploinsufficiency Predicts Regulatory
Networks for Congenital Heart Disease. *Dev Cell* **56**, 292-309.e9 (2021).

23. Bruneau, B. G. *et al.* A murine model of Holt-Oram syndrome defines roles of the T-box
transcription factor Tbx5 in cardiogenesis and disease. *Cell* **106**, 709–721 (2001).

24. Verzi, M. P., McCulley, D. J., Val, S. D., Dodou, E. & Black, B. L. The right ventricle, outflow
tract, and ventricular septum comprise a restricted expression domain within the
secondary/anterior heart field. *Dev Biol* **287**, 134–145 (2005).

25. Koshiba-Takeuchi, K. *et al.* Reptilian heart development and the molecular basis of cardiac
chamber evolution. *Nature* **461**, 95–98 (2009).

26. Xie, L. *et al.* Tbx5-Hedgehog Molecular Networks Are Essential in the Second Heart Field
for Atrial Septation. *Dev Cell* **23**, 280–291 (2012).

27. Mori, A. D. & Bruneau, B. G. TBX5 mutations and congenital heart disease: Holt-Oram
syndrome revealed. *Curr Opin Cardiol* **19**, 211–215 (2004).

28. Mommersteeg, M. T. M., Yeh, M. L., Parnavelas, J. G. & Andrews, W. D. Disrupted Slit-
Robo signalling results in membranous ventricular septum defects and bicuspid aortic valves.
Cardiovasc Res **106**, 55–66 (2015).

29. Li, Y. *et al.* Global genetic analysis in mice unveils central role for cilia in congenital heart
disease. *Nature* **521**, 1–16 (2015).

30. Kruszka, P. *et al.* Loss of function in ROBO1 is associated with tetralogy of Fallot and septal
defects. *J Med Genet* **54**, 825 (2017).

31. Larrieu-Lahargue, F., Thomas, K. R. & Li, D. Y. Netrin Ligands and Receptors: Lessons
From Neurons to the Endothelium. *Trends Cardiovas Med* **22**, 44–47 (2012).

32. Akerberg, B. N. *et al.* A reference map of murine cardiac transcription factor chromatin
occupancy identifies dynamic and conserved enhancers. *Nat Commun* **10**, 4907 (2019).

33. Bin, J. M. *et al.* Complete Loss of Netrin-1 Results in Embryonic Lethality and Severe Axon
Guidance Defects without Increased Neural Cell Death. *CellReports* **12**, 1099–1106 (2015).

34. Lewandoski, M., Meyers, E. N. & Martin, G. R. Analysis of Fgf8 gene function in vertebrate
development. *Cold Spring Harb Sym* **62**, 159–68 (1997).

35. Kiemen, A. L. *et al.* CODA: quantitative 3D reconstruction of large tissues at cellular
resolution. *Nat Methods* 1–10 (2022) doi:10.1038/s41592-022-01650-9.

36. Vincent, J. P. Compartment boundaries: where, why and how? *Int. J. Dev. Biol.* **42**, 311–5
(1998).

37. Lawrence, P. A. & Struhl, G. Morphogens, compartments, and pattern: lessons from
drosophila? *Cell* **85**, 951–961 (1996).

38. VanDusen, N. J. *et al.* Hand2 Is an Essential Regulator for Two Notch-Dependent Functions
within the Embryonic Endocardium. *Cell Rep.* **9**, 2071–2083 (2014).

39. George, R. M. *et al.* Single cell evaluation of endocardial HAND2 gene regulatory networks
reveals critical HAND2 dependent pathways impacting cardiac morphogenesis. *Development*
150, (2023).

40. Bin, J. M. *et al.* Complete Loss of Netrin-1 Results in Embryonic Lethality and Severe Axon
Guidance Defects without Increased Neural Cell Death. *Cell Reports* **12**, 1099–1106 (2015).

41. Tasic, B. *et al.* Site-specific integrase-mediated transgenesis in mice via pronuclear
injection. *Proc National Acad Sci* **108**, 7902–7907 (2011).

42. Madisen, L. *et al.* A robust and high-throughput Cre reporting and characterization system
for the whole mouse brain. *Nat Neurosci* **13**, 133–140 (2010).

43. Dominguez, M. H., Krup, A. L., Muncie, J. M. & Bruneau, B. G. Graded mesoderm assembly
governs cell fate and morphogenesis of the early mammalian heart. *Cell* **186**, 479-496.e23
(2023).

44. Schindelin, J. *et al.* Fiji: an open-source platform for biological-image analysis. *Nat. Methods*
9, 676–682 (2012).

45. Kirshner, H., Aguet, F., Sage, D. & Unser, M. 3-D PSF fitting for fluorescence microscopy:
implementation and localization application. *J. Microsc.* **249**, 13–25 (2013).

46. Hörl, D. *et al.* BigStitcher: reconstructing high-resolution image datasets of cleared and
expanded samples. *Nat Methods* **16**, 870–874 (2019).

47. Tinevez, J.-Y. *et al.* TrackMate: An open and extensible platform for single-particle tracking.
Methods **115**, 80–90 (2017).

48. Satija, R., Farrell, J. A., Gennert, D., Schier, A. F. & Regev, A. Spatial reconstruction of
single-cell gene expression data. *Nat Biotechnol* **33**, 495–502 (2015).

49. Butler, A., Hoffman, P., Smibert, P., Papalexi, E. & Satija, R. Integrating single-cell
transcriptomic data across different conditions, technologies, and species. *Nat Biotechnol* **48**,
1070–17 (2018).

50. Stuart, T. *et al.* Comprehensive Integration of Single-Cell Data. *Cell* **177**, 1–37 (2019).

51. Becht, E. *et al.* Dimensionality reduction for visualizing single-cell data using UMAP. *Nat*
Biotechnol (2018) doi:10.1038/nbt.4314.

52. Forjaz, A. *et al.* Three-dimensional assessments are necessary to determine the true spatial
tissue composition of diseased tissues. *bioRxiv* 2023.12.04.569986 (2023)
doi:10.1101/2023.12.04.569986.

Acknowledgements. We dedicate this manuscript in memory of Tatyana Sukonnik. We thank

Brian Black for mouse lines and members of the Bruneau lab for troubleshooting help, discussions

and comments. We also thank the Gladstone Bioinformatics, Genomics, and Histology and

Microscopy Cores, the UCSF Laboratory for Cell Analysis, and the UCSF Center for Advanced

Technology. This work was supported by grants from the National Institutes of Health (NHLBI

Bench to Bassinet Program UM1HL098179 to B.G.B.; R01HL114948 to B.G.B., R01HL155906

to B.G.B. and I.S.K., F32 HL162450 to J.M.M-V, U54CA268083 and UG3CA275681 to A.L.K.),

Additional Ventures to B.G.B., Society for Pediatric Anesthesia (Young Investigator Award to

I.S.K.), Hellman Family Fund (I.S.K.), UCSF REAC Grant (I.S.K.), UCSF Pediatric Heart Center

Catalyst Award to I.S.K. and UCSF Department of Anesthesia and Perioperative Care (Ongoing

Investigator Award to I.S.K.). This work was also supported by an NIH/NCRR grant (C06

RR018928) to the J. David Gladstone Institutes, and the Younger Family Fund (B.G.B.).

Author Contributions. W.P.D. and B.G.B. originally conceived, and I.S.K. and B.G.B. refined

and developed the study. W.P.D. made initial observations using genetic lineage tools that he

gathered or generated, such as $ROSA26^{Ai66b}$ and the intersectional-DTA. I.S.K., M.H.D., W.P.D.

and K.M.H. performed embryo dissections. K.M.H., B.I.G., P.G., M.N.M., T.S., D.M.P., S.W. and

E.F.B. managed animal husbandry and genotyping. I.S.K., M.H.D., W.P.D. and K.M.H.

performed whole mount or section immunostaining experiments and imaging. M.H.D., J.M.M.

and D.Q performed lightsheet imaging and processing, and M.H.D. and J.M.M. implemented

analyses with hypothesis-based input from I.S.K. I.S.K. designed the strategy to use 5'

sequencing kit and developed synthetic genome with Gladstone Bioinformatics core, micro-

dissected heart regions for scRNA-seq and processed samples with K.S.R. and P.G. for

sequencing library preparations. K.S.R. generated scRNA-seq libraries and performed scRNA-

seq analyses with hypothesis-based input from I.S.K. K.S.R., K.M.H., P.G. and M.N.M.

performed section *in situ* hybridization and imaging. K.S.R. generated ChIP-seq browser tracks

with input from S.K.H. I.S.K. assessed cardiac defects by histology and provided anatomical

knowledge to A.L.K. to apply a deep learning tool for cardiac morphometry. A.F. trained deep

learning models and was supervised by A.L.K. and advised by P-H.W. and D.W. B.G.B.

supervised and advised the study. I.S.K. and K.S.R. prepared figures. I.S.K. wrote the original

draft with contributions from co-authors. I.S.K. and B.G.B. reviewed and edited the manuscript

with input from co-authors.

Declaration of Interests: B.G.B. is a co-founder and shareholder of Tenaya Therapeutics.

B.G.B. is an advisor for Silver Creek Pharmaceuticals. None of the work presented here is

related to these commercial interests.

Figure 1

Figure 1. *Tbx5$^+$/Mef2cAHF$^+$* lineage marks a compartment boundary at the cardiac interventricular septum At E14.5, (A, B) clonal cell descendants of a *Tbx5$^+$* lineage (ZsGreen) labeled at E6.5 contributes to the LV, and (C, D) a *Mef2cAHF$^+$* lineage (tdTomato immunostaining) contributes to the RV, demonstrating largely complementary patterns. (F-J) These lineages overlap (*Tbx5$^+$/Mef2cAHF$^+$*) at the interventricular septum (IVS), as well as the atrioventricular canal (AVC) and interatrial septum (IAS). *Tbx5$^{CreERT2/+}$;Mef2cAHF-DreERT2;ROSA26$^{Ai6/Ai66b}$* hearts by epifluorescence microscopy (A-C) and cryosections (D-F). (G-J) Maximal projection images by light sheet microscopy of an intersectional reporter for the *Tbx5$^+$/Mef2cAHF$^+$* lineage (tdTomato immunostaining) in *Tbx5$^{CreERT2/+}$;Mef2cAHF-DreERT2;ROSA26$^{Ai66/+}$* embryos.

Figure S1

Supplemental Figure 1. (A-C) *Tbx5$^+$/Mef2cAHF$^+$* lineage contributions to the interventricular
septum were descendants from progenitors labeled at (A) E6.5, but not at (B) E8.5 or (C) E10.5
in *Tbx5$^{CreERT2/+}$;Mef2cAHF-DreERT2;ROSA26$^{Ai6/Ai66}$* embryos. Epifluorescence microscopy
images of the *Tbx5$^+$/Mef2cAHF$^+$* lineage (tdTomato immunostaining) labeled at E6.5 from (A)
E7.75, (B) E8.0 and (C) E8.5 displayed an apparent "salt and pepper" pattern in the early
developing heart.
820
821

Figure 2

Figure 2. *Tbx5$^+$/Mef2cAHF$^+$* lineage is prefigured in early mouse heart development. Frontal (A-E), transverse (A'-E') and magnified transverse (E'') views of maximum Z-projections of whole mount embryos by lightsheet imaging at E8.0 show the *Tbx5$^+$* lineage (ZsGreen), and immunostaining of tdTomato for the *Tbx5$^+$/Mef2cAHF$^+$* lineage, MEF2c and cTNT in *Tbx5$^{CreERT2/+}$;Mef2cAHF-DreERT2;ROSA26$^{Ai6/Ai66}$* embryos. A-P, anterior-posterior axis; V-D, ventral-dorsal axis. First (FHF) and second (SHF) heart fields. At E8.0 (approximately 3 somites), *Tbx5$^+$* lineage cells are restricted to a ventral domain corresponding to the FHF, whereas *Tbx5$^+$/Mef2cAHF$^+$* lineage cells originate along the FHF-SHF boundary. *Tbx5$^+$/Mef2cAHF$^+$* lineage cells lie in an apparent planar ringlet in the dorsal-ventral axis that expands as the heart tube grows. (J-N) At E8.25 (approximately 6 somites), with dorsal closure of the linear heart tube (LHT) underway, the *Tbx5$^+$/Mef2cAHF$^+$* lineage cells form a ring situated between the RV and LV primordia, which is observed from multiple perspectives including anterior, short-axis (SAx) and long-axis (LAx) views. (O-Q) At E8.5 (approximately 9 somites), *Tbx5$^+$/Mef2cAHF$^+$* lineage cells are observed in a portion of the AVC primoridium (pAVC and arrowheads). (R-T) At E10.5, a band of *Tbx5$^+$/Mef2cAHF$^+$* lineage cells extends from the interventricular groove at the outer curvature to the inner curvature of the heart. SAx and LAx show that *Tbx5$^+$/Mef2cAHF$^+$* lineage cells occupies a crossroads for heart morphogenesis, spanning the growing interventricular septum to the AVC, adjacent to the OFT.

Figure S2

842   **Supplemental Figure 2.** (A-E) In the developing atria, optical sections from ventral to dorsal show
843   lineage-labeled $Tbx5^+/Mef2cAHF^+$ lineage cells from the AV cushion (AVC) region to the roof of
844   the atrium, prior to the formation of the interatrial septum, in $Tbx5^{CreERT2/+}$;$Mef2cAHF$-
845     $DreERT2$;$ROSA26^{Ai6/Ai66}$ embryos.
846
847   **Video 1.** Related to Figure 2 and Supplemental Figure 2.
848
849

Figure 3

Figure 3. Cell ablation of *Tbx5⁺/Mef2cAHF⁺* progenitors causes RV hypoplasia, IVS disorganization, and lineage mixing. Misexpression of *diphtheria toxin* (*DTA*) in *Tbx5⁺/Mef2cAHF⁺* progenitors in *Tbx5$^{CreERT2/+}$;Mef2cAHF-DreERT2;ROSA26$^{Ai66/+}$;Hip11$^{intersectional-DTA/+}$* embryos (*DTA*), compared to *Tbx5$^{CreERT2/+}$;Mef2cAHF-DreERT2;ROSA26$^{Ai66/+}$;Hip11$^{+/+}$* (control), resulted in RV hypoplasia (arrows) at E9.5 (A, B), and E12.5 (C-J), along with disorganization and non-compaction of the interventricular septum (IVS) (asterisk) (G-J). (K-N) At E10.5, mixing of the *Mef2cAHF⁺* lineage and *Tbx5⁺* lineage (white arrow in N shows ectopic *Tbx5⁺* lineage ZsGreen cell in the right heart) was observed by lightsheet microscopy in *Tbx5$^{CreERT2/+}$;Mef2cAHF-DreERT2;ROSA26$^{Ai6/Ai66b}$;Hip11$^{intersectional-DTA/+}$* embryos (*DTA*) (n=2) compared to *Tbx5$^{CreERT2/+}$;Mef2cAHF-DreERT2;ROSA26$^{Ai6/Ai66b}$;Hip11$^{+/+}$* embryos (control) (n=2). (O-R) Linear profiles quantified fluorescence intensity above a threshold (dashed line in P,R) across the heart at the right or left (R/L) positions. *Mef2cAHF⁺* lineage retreated rightward (depiction in O, purple arrow in P) while *Tbx5⁺* lineage cells expanded rightward into the RV (depiction in Q, green arrow in R), consistent with a compartment boundary disruption. Yellow shading represents *Tbx5⁺/Mef2cAHF⁺* lineage cells and region of cell ablation. Statistical significance ($p<0.05$) between control and *DTA* mutants was determined by Welch's two-sample t-test at each position along the right-left axis.

Figure S3

869 **Supplemental Figure 3. Conditional haploinsufficiency of *Tbx5* in the interventricular**
870 **septum (IVS) leads to ventricular septal defects (VSDs).** (A-C) At E14.5, membranous VSDs
871 were observed in *Mef2cAHF-Cre*;*ROSA26$^{mTmG/+}$*;*Tbx5$^{flox/+}$* mutant embryos (n=4/7) compared to
872 *Mef2cAHF-Cre*;*ROSA26$^{mTmG/+}$*;*Tbx5$^{flox/+}$* controls (n=0/3).
873
874

Figure 4

Tbx5⁺ **Tbx5⁺/Mef2cAHF⁺**

A *Tbx5^{CreERT2/+}*   A'   A''
RV IVS
Tam E6.5
E14.5   LV

B *Tbx5^{CreERT2/flox}*   B'   B''
RV   LV

C *Tbx5^{CreERT2/flox}*   C' *
C''
RV   LV
200μm

D *Tbx5^{CreERT2/flox}*   D'
RA
LV   **
RV
166μm

E *Tbx5^{CreERT2/+}*
RV IVS   LV

F tdTomato⁺ linear profile plot
200
100
0   200 400 600 800 1000
0
toward RV ◄ microns ► toward LV

G

H tdTomato⁺ linear plot peak intensity
x̄: p<0.05
Tbx5^{CreERT2/+}   σ: p<0.01
Tbx5^{CreERT2/flox}
toward RV ◄ center ► toward LV

I J K L M   orientation by color
Tbx5^{CreERT2/+}   90°
0°
N O P Q R   -90°
Tbx5^{CreERT2/flox}

S tdTomato⁺ average orientation score
-90° -75° -60° -45° -30° -15° 0° 15° 30° 45° 60° 75° 90°
p < 0.001

0±5° p<0.00001   65±20° p<0.00001   90±5° p<0.00001
12   8   6
8   6   4
4   4   2
                    2
CreERT2/+ CreERT2/flox CreERT2/+ CreERT2/flox CreERT2/+ CreERT2/flox

T tdTomato⁺ directional coherency
p < 0.005
0.4
0.3
0.2
0.1
0.0
CreERT2/+ CreERT2/flox

Figure 4. Reduced Tbx5 dosage caused ventricular septal defects and perturbations to IVS
boundary position and integrity. (A) Mid-posterior optical sections from light sheet microscopy
of a control (*Tbx5$^{CreERT2/+}$;Mef2cAHF-DreERT2;ROSA26$^{Ai6/Ai66}$*) heart display *Tbx5$^+$* lineage
(ZsGreen) and *Tbx5$^+$/Mef2cAHF$^+$* lineage (tdTomato immunostaining) cells. *Tbx5$^+$/Mef2cAHF$^+$*
lineage cells in the interventricular septum (IVS) extended from the base (A') to the apex (A") of
the heart, where cells were highly organized. (B-D) A spectrum of phenotypes of *Tbx5$^{CreERT2/flox}$*
mutants (*Tbx5$^{CreERT2/flox}$;Mef2cAHF-DreERT2;ROSA26$^{Ai6/Ai66}$*) were observed, including intact IVS
(B-B"), membranous VSD (C-C", asterisk) and atrioventricular septal defect (double asterisk in
D'). (B', B", C', C", D') A reduction, maldistribution and disorientation of *Tbx5$^+$/Mef2cAHF$^+$* lineage
cells were observed in *Tbx5$^{CreERT2/flox}$* mutants. (E-H) Linear plot profiles at positions along the
anterior-posterior and apical-basal axis of control (n=4) and *Tbx5$^{CreERT2/flox}$* mutant (n=3) hearts for
Tbx5$^+$/Mef2cAHF$^+$ lineage cells showed a leftward shift of boundary positioning and broadening
of the *Tbx5$^+$/Mef2cAHF$^+$* lineage position, consistent with lineage mixing. (E,F) An example of a
linear profile plot is shown, and (G,H) aggregated data is depicted and quantified. (I-T) Orientation
scores for cells from each channel (tdTomato$^+$ or ZsGreen$^+$) was delineated for each genotype,
and distributions were plotted as a function of angle. *Tbx5$^{CreERT2/flox}$* mutant hearts scored worse
for orientation of *Tbx5$^+$/Mef2cAHF$^+$* lineage (tdTomato$^+$) cells in the dominant direction (range
from -5 to 5 degrees) and scored higher in the orientation orthogonal to the dominant direction
(range from 85 to 95 degrees and -85 to -95 degrees), as determined (S) by Watson's two-sample
test across all orientations or by Wilcoxon signed-rank test for selected orientations. (T)
Tbx5$^+$/Mef2cAHF$^+$ lineage (tdTomato$^+$) cells demonstrated worse directional coherency scoring in
Tbx5$^{CreERT2/flox}$ mutants by Wilcoxon signed-rank test.
897
898

Figure S4

Supplemental Figure 4. (A-H) Volume renderings from lightsheet microscopy of anterior and
posterior surfaces, as well as anterior and mid-anterior 4-chamber optical sections of $Tbx5^{CreERT2/+}$
control ($Tbx5^{CreERT2/+}$;*Mef2cAHF-DreERT2*;$ROSA26^{Ai6/Ai66}$) and $Tbx5^{CreERT2/flox}$ mutant
($Tbx5^{CreERT2/flox}$;*Mef2cAHF-DreERT2*;$ROSA26^{Ai6/Ai66}$) hearts at E14.5 display $Tbx5^+$ lineage
(ZsGreen) and $Tbx5^+$/*Mef2cAHF*$^+$ lineage (tdTomato immunostaining) cells. (I, J) Quantification
of the anatomical location of tdTomato$^+$ and ZsGreen$^+$ cells in $Tbx5^{CreERT2/flox}$ mutants (n=4) and
$Tbx5^{CreERT2/+}$ controls (n=4), by region as outlined in (I). Over 1200 3-dimensional regions of
interest were drawn blinded, and then integrated fluorescence signal was assessed. (J) Although
no regional differences in the intensities of ZsGreen$^+$ were detected, tdTomato$^+$ displayed higher
integrated fluorescence in the left ventricle (LV) and less signal in atrioventricular canal (AVC) by
Wilcoxon signed-rank test. (L-U) $Tbx5^{CreERT2/flox}$ mutant hearts scored worse for orientation scores
of $Tbx5^+$ lineage (ZsGreen$^+$) cells in the dominant direction (range from -5 to 5 degrees) and
scored higher in the orientation orthogonal to the dominant direction (range from 85 to 95 degrees
and -85 to -95 degrees), as determined (T) by Watson's two-sample test across all orientations
or by Wilcoxon signed-rank test for selected orientations. (U) $Tbx5^+$ lineage (ZsGreen$^+$) cells
demonstrated worse directional coherency scoring in $Tbx5^{CreERT2/flox}$ mutants by Wilcoxon signed-
rank test.
916
917   **Video 2.** Related to Figure 4 and Supplemental Figure 4.
918
919

Figure 5

Figure 5. *Slit2* and *Ntn1* are *Tbx5*-sensitive genes in the interventricular septum (IVS). (A) At E13.5, we micro-dissected the right ventricle (RV), left ventricle (LV) and IVS+atrioventricular canal (AVC) in $Tbx5^{CreERT2/+}$ controls (n=4) and $Tbx5^{CreERT/flox}$ mutants ($Tbx5^{CreERT2/flox}$;*Mef2cAHF-DreERT2*;$ROSA26^{Ai6/Ai66}$)(n=2) with the labeled $Tbx5^+$/$Mef2cAHF^+$ lineage ($tdTomato^+$) and $Tbx5^+$ lineage ($ZsGreen^+$) cells, after tamoxifen at E6.5. (B) UMAP visualization by cell type clusters and *Tbx5* genotype (inset). CMs, cardiomyocytes; WBCs, white blood cells; RBCs, red blood cells. (C) $ZsGreen^+$ and (D) $tdTomato^+$ cells are enriched among CMs. (E) UMAP shows a $Tnnt2^+$ CM subset by *Tbx5* genotype or (F) Louvain clusters. (G) Feature plots show that atrioventricular canal genes (*Rspo3, Bmp2, Robo1, Unc5b*) are enriched among control-enriched cluster 5, suggesting downregulation of these genes in *Tbx5* mutants. Trabecular genes (*Cited1, Nppa, Slit2, Ntn1*) were enriched in the *Tbx5* mutant-enriched cluster 10. (H) Dot plots of selected genes that are upregulated in cluster 10 (C10) or in cluster 5 (C5). Significance of adj p<0.05 was determined by Wilcoxan rank sum. (I-N) Fluorescent *in situ* hybridization of trabecular genes *Nppa, Slit2* and *Ntn1* in $Tbx5^{CreERT2/+}$ controls or $Tbx5^{CreERT2/flox}$ mutants at E14.5. (I, I', K, K') *Nppa* and *Slit2* are normally expressed in the ventricular trabecular layer and excluded from the core of the IVS. (J, J', L, L') In a $Tbx5^{CreERT2/flox}$ mutant with an intact IVS, *Nppa* and *Slit2* are misexpressed in the IVS (asterisks). (M, M') *Ntn1* is normally expressed in the trabecular layer and in a gradient across the IVS from left to right, demarcated by yellow dashed line. (N, N') In a $Tbx5^{CreERT2/flox}$ mutant with an intact IVS, *Ntn1* is expanded across the IVS, flattening its gradient.

Figure S5

Supplemental Figure 5. In scRNA-seq samples from E13.5 hearts (control, n=4; *Tbx5* mutant,
n=3), we display cells by region (LV, RV, IVS) (A) or by genotype (B). IVS+AVC region (control,
n=4, *Tbx5* mutant, n=2), we detected clusters enriched for (C) *Tnnt2*⁺ cardiomyocytes (CMs), (D)
Tbx18⁺/*Wt1*⁺ epicardial cells, (E) *Plvap*⁺ endothelial cells, (F) *Postn*⁺ fibroblasts, (G) *Hbb-b2*⁺ red
blood cells, and (H) *C1qb*⁺ white blood cells. Dot plots of selected differentially expressed genes
in (I) *tdTomato*⁺ or (J) *ZsGreen*⁺ cells from the IVS+AVC region. Significance of adj p<0.05 was
determined by Wilcoxan rank sum.
948
949

Figure S6

Supplemental Figure 6. Fluorescent *in situ* hybridization of *Tbx5*-dependent genes. (A-L')
Fluorescent *in situ* hybridization at E14.5 after tamoxifen at E6.5 of *Nppa*, *Slit2* and *Ntn1* are
expressed in the trabecular layer in *Tbx5$^{CreERT2/+}$* controls (*Tbx5$^{CreERT2/+}$;Mef2cAHF-
DreERT2;ROSA26$^{Ai6/Ai66}$*), and misexpressed in the IVS in *Tbx5$^{CreERT2/flox}$* mutants
(*Tbx5$^{CreERT2/flox}$;Mef2cAHF-DreERT2;ROSA26$^{Ai6/Ai66}$*) with a membranous VSD at E14.5. Yellow
dashed line demarcates gradient of *Ntn1* expression across the IVS from left to right. *Tnnt2* is
expressed in cardiomyocytes. (M-X') Expression of *Bmp2*, *Robo1* and *Unc5b* are expressed in
the atrioventricular canal (boxed region) in *Tbx5$^{CreERT2/+}$* controls, and expression of these genes
are reduced in *Tbx5$^{CreERT2/flox}$* mutants. (Y) Immunostaining of NTN1 at E14.5. (Z) Fluorescent *in
situ* hybridization of *Ntn1* at E11.5. Asterisks indicate ectopic expression. Arrows highlight
enriched expression.
961
962

Figure S7

963   **Supplemental Figure 7. TBX5 chromatin occupancy near *Slit2* and *Ntn1*.** (A-C) Browser
964   tracks of TBX5 ChIP-seq from E12.5 hearts [32] shows peaks of TBX5 occupancy (boxed in red)
965   near promoters of *Nppa*, *Slit2* and *Ntn1*.
966
967

Figure S8

Supplemental Figure 8. *Tbx5-Ntn1* and *Tbx5-Slit2* genetic interactions. Histology from E14.5
embryos from matings of (A-D) $Tbx5^{del/+}$ and $Slit2^{+/-}$ or (F-I) $Tbx5^{del/+}$ and $Ntn1^{+/-}$ by genotype.
Arrowheads depict muscular ventricular septal defects (VSDs). (E,J) Quantification of
membranous or muscular VSDs by genotype. $Tbx5^{del/+}$;$Slit2^{+/-}$compound heterozygous embryos
displayed an estimated decrease in the incidence of membranous VSDs, which approached
significance (log OR -2.5; p<0.09 in a Generalized Linear Model). $Tbx5^{del/+}$;$Ntn1^{+/-}$ compound
heterozygous embryos displayed a decrease of membranous VSDs (log OR -5.3; $p<1x10^{-3}$ by a
Generalized Linear Model), but not changes to prevalence of muscular VSDs (log OR 0.12;
p<0.93). Machine-learning based quantitative morphometry of metrics for (K-P) $Tbx5^{del/+}$ and
$Ntn1^{+/-}$ or (Q-V) $Tbx5^{del/+}$ and $Slit2^{+/-}$ by genotype. IVS, interventricular septum. Significance
determined by Fisher's exact test (*p<0.05, **p<0.01, ***p<0.001).
979
980

Figure 6

Figure 6. *Slit2* and *Ntn1* are essential for proper ventricular septation and compartment
boundary regulation. (A-C) Histology and (D) incidence of muscular or membranous
(arrowhead) ventricular septal defects (VSDs) of *Slit2* mutants are shown. Asterisk demarcates
non-compacted interventricular septum (IVS) (E-G) Histology and (H) incidence of muscular or
membranous ventricular septal defects (VSDs) of *Ntn1* mutants are shown. (I-K) Machine
learning-based 3-D reconstructions of the embryonic heart and other tissues from histology at
E14.5 were used for quantitative morphometric analysis of the embryonic heart (Supplemental
Figure 7, Supplemental Figure 8). Compact myo, compact myocardium; AV canal, atrioventricular
canal; IVS, interventricular septum. (L,M) A 3-D reconstruction of a membranous ventricular
septal defect (VSD) anteriorly and muscular VSD posteriorly from a $Tbx5^{dell/+}$ mutant heart. (N-P)
Lightsheet imaging of $Slit2^{-/-}$ mutant ($Tbx5^{CreERT2/+}$;*Mef2cAHF-DreERT2*;$ROSA26^{Ai66/+}$;$Slit2^{-/-}$)(n=4)
hearts at E14.5 showed a broadened distribution of the $Tbx5^+$/$Mef2cAHF^+$ lineage (tdTomato$^+$
immunostaining) compared to controls ($Tbx5^{CreERT2/+}$;*Mef2cAHF-*
DreERT2;$ROSA26^{Ai66/+}$;$Slit2^{+/+}$)(n=2), as (Q) quantified by linear profiles of fluorescence signals
above a threshold (dashed line). Gray shading represents control signal above threshold, and
periwinkle represents mutant signal. Significance of p<0.05 was determined by Welch's two-
sample t-test at each position along the right-left axis. (R-T) Lightsheet imaging of $Ntn1^{-/-}$ mutant
($Tbx5^{CreERT2/+}$;*Mef2cAHF-DreERT2*;$ROSA26^{Ai66/+}$;$Ntn1^{-/-}$)(n=6) hearts at E14.5 showed a leftward
shift in positioning of the $Tbx5^+$/$Mef2cAHF^+$ lineage compared to controls
($Tbx5^{CreERT2/+}$;*Mef2cAHF-DreERT2*;$ROSA26^{Ai66/+}$;$Ntn1^{+/+}$)(n=4), as (U) quantified by linear
profiles.
1002
1003

Figure S9

1004   **Supplemental Figure 9. Quantitative morphometry of *Slit2* and *Ntn1* mutant hearts.** (A-D)
1005   Additional histology by *Ntn1* genotype. Arrowhead, muscular ventricular septal defect (VSD).
1006   Asterisk, membranous VSD. (E-P) Machine learning-based quantitative cardiac morphometrics
1007   for (E-K) *Slit2* or (L-R) *Ntn1* mutants. Significance determined by Fisher's exact test (*p<0.05).
1008   (Q,R) Morphometry of the area of membranous VSDs by *Slit2* or *Ntn1* genotypes.