Considerations for reporting variants in novel candidate genes identified during clinical genomic testing

Jessica X. Chong[1,2,*], Seth I. Berger[3], Samantha Baxter[4], Erica Smith[5], Changrui Xiao[6], Daniel G. Calame[7], Megan H. Hawley[8], E. Andres Rivera-Munoz[9], Stephanie DiTroia[4], [Genomics Research to Elucidate the Genetics of Rare Diseases (GREGoR) Consortium], Michael J. Bamshad[1,2,10], Heidi L. Rehm[11,4]

[1]Department of Pediatrics, Division of Genetic Medicine, University of Washington, 1959 NE Pacific Street, Box 357371, Seattle, WA, 98195, USA

[2]Brotman-Baty Institute for Precision Medicine, 1959 NE Pacific Street, Box 357657, Seattle, WA, 98195, USA

[3]Center for Genetic Medicine Research, Children's National Research Institute, 111 Michigan Ave, NW, Washington, DC, 20010, USA

[4]Program in Medical and Population Genetics, Broad Institute of MIT and Harvard, 415 Main St, Cambridge, MA, 02141, USA

[5]Department of Clinical Diagnostics, Ambry Genetics, 15 Argonaut, Aliso Viejo, CA, 92656, USA

[6]Department of Neurology, University of California Irvine, 200 South Manchester Ave. St 206E, Orange, CA, 92868, USA

[7]Department of Pediatrics, Division of Pediatric Neurology and Developmental Neurosciences, Baylor College of Medicine, One Baylor Plaza, Houston, TX, 77030, USA

[8]Clinical Operations, Invitae, 485F US-1 Suite 110, Iselin, NJ, 08830, USA

[9]Department of Molecular and Human Genetics, Baylor College of Medicine, 1 Baylor Plaza T605, Houston, TX, 77030, USA

[10]Department of Pediatrics, Division of Genetic Medicine, Seattle Children's Hospital, Seattle, WA, 98195, USA

[11]Center for Genomic Medicine, Massachusetts General Hospital, 185 Cambridge St, Boston, MA, 02114, USA

* correspondence should be addressed to:

Jessica Chong, PhD

Department of Pediatrics

University of Washington School of Medicine

Box 357371

1959 NE Pacific Street, HSB I607I

Seattle, WA 98195

Phone: (206) 221-4075

FAX: (206) 221-3795

jxchong@uw.edu

Abstract

Since the first novel gene discovery for a Mendelian condition was made via exome sequencing (ES), the rapid increase in the number of genes known to underlie Mendelian conditions[1] coupled with the adoption of exome (and more recently, genome) sequencing by diagnostic testing labs has changed the landscape of genomic testing for rare disease. Specifically, many individuals suspected to have a Mendelian condition are now routinely offered clinical ES. This commonly results in a precise genetic diagnosis but frequently overlooks the identification of novel candidate genes. Such candidates are also less likely to be identified in the absence of large-scale gene discovery research programs. Accordingly, clinical laboratories have both the opportunity, and some might argue a responsibility, to contribute to novel gene discovery which should in turn increase the diagnostic yield for many conditions. However, clinical diagnostic laboratories must necessarily balance priorities for throughput, turnaround time, cost efficiency, clinician preferences, and regulatory constraints, and often do not have the infrastructure or resources to effectively participate in either clinical translational or basic genome science research efforts. For these and other reasons, many laboratories have historically refrained from broadly sharing potentially pathogenic variants in novel genes via networks like Matchmaker Exchange, much less reporting such results to ordering providers. Efforts to report such results are further complicated by a lack of guidelines for clinical reporting and interpretation of variants in novel candidate genes. Nevertheless, there are myriad benefits for many stakeholders, including patients/families, clinicians, researchers, if clinical laboratories systematically and routinely identify, share, and report novel candidate genes. To facilitate this change in practice, we developed criteria for triaging, sharing, and reporting novel candidate genes that are most likely to be promptly validated as underlying a Mendelian condition and translated to use in clinical settings.

Introduction

Over the past decade, phenotype-driven exome and genome sequencing (ES, GS), has demonstrated utility as a first-line test for clinical diagnostic testing in individuals with suspected Mendelian conditions.[2–6] Nevertheless, a large fraction (~30%-80%, depending on prior testing, categorical phenotype, analytical approach, and other factors) of persons tested by ES/GS remain without a precise genetic diagnosis,[4,7–12] and a substantial proportion of these individuals may harbor variants in novel candidate genes that explain their clinical findings. For example, re-analysis of existing ES/GS data moderately improves diagnostic yield, with novel gene discoveries—i.e. newly-reported gene-disease relationships, of which an estimated ~100-200 are published per year[13,14]— accounting for the majority of new precise genetic diagnoses.[9,15–19] These findings highlight the value of incorporating recently-established gene discoveries in the clinical analysis of ES/GS data, and suggest that even greater benefits may be leveraged by systematically sharing knowledge about candidate gene discoveries.

In contrast to single gene and multi-gene panel testing, ES/GS agnostically detects variants throughout the genome, including variants in genes that have been found to underlie a Mendelian condition (i.e., known gene) and genes that have not been implicated but for which varying sources of evidence may support a role in a Mendelian condition (i.e. novel candidate genes). As a result, if an individual's clinical findings remain partially or completely unexplained after assessment of variants in known genes, assessing variants in novel candidate genes could result in a precise genetic diagnosis.[9,15–18] Moreover, the increased cost (e.g., labor, time) to assess these variants is likely modest when added to an existing computational pipeline, suggesting a high return on the investment of generating the ES/WGS data.

While extensive guidelines exist for assessing and reporting the pathogenicity of variants in known genes, including variants for which the clinical effect includes degrees of uncertainty,[20–26] guidance is lacking for clinical interpretation and reporting of variants in novel

candidate genes. We describe criteria to select novel candidate genes supported by sufficient evidence as to be of high clinical value, yet small enough in number so that routine inclusion in clinical ES/GS reporting at this time is manageable. Further, we provide guidance for reporting and sharing variants in novel candidate genes.

Drafting recommendations

We formed a working group within the GREGoR Consortium comprising clinicians and research scientists with expertise in gene discovery, assessment of genotype-phenotype relationships, diagnostic genome analysis, variant classification, and clinical result reporting. A series of biweekly online calls were conducted over six months to draft, discuss, and refine the guidance. The group reviewed novel candidate variant/gene criteria used by several large-scale gene discovery/diagnostic efforts[27–30] as well as descriptions from diagnostic laboratories about their evaluation and clinical reporting practices for novel candidate genes.[31–35]

Terminology

The terms below are used as follows:

- *Clinical gene-disease validity* – an evidence-based assessment that evaluates the role of a disrupted or dysregulated gene in causing a particular Mendelian condition.[36] Evaluating the validity of a gene's asserted role in disease is not a binary assessment and instead, validity ranges in evidence strength. ClinGen developed a semi-quantitative framework[36] to define evidence levels and after a delphi-survey process from the Gene Curation Coalition (GenCC)[37], this framework was endorsed by all major gene-disease databases with harmonized assessments available in the GenCC Database (search.thegenecc.org), except for OMIM and Orphanet which use a binary inclusion or

exclusion approach. While seven levels of evidence are articulated in the framework (Definitive, Strong, Moderate, Limited, No Human Disease Evidence (+/- Non-human animal model), Disputed, and Refuted), ACMG standards indicate that the levels of Moderate to Definitive are sufficient to include in a panel clinical genetic test and that genes with Limited or less evidence should only be examined in genome-wide assessments such as ES and GS.[38]

- *Known gene* – a gene with at least a Moderate score of gene-disease validity according to the ClinGen framework (scored genes are available through GenCC[37] (https://search.thegencc.org)

- *Novel candidate gene* – a gene in a newly-proposed gene-disease relationship. The gene either has not yet been implicated in any Mendelian condition or, more broadly, has already been implicated in a Mendelian condition but is being proposed to underlie a novel phenotype, expand the known phenotype (phenotypic expansion), or lead to a disease through a novel mechanism (e.g. different than the known mode of inheritance). These genes are a type of Gene of Uncertain Significance (GUS), which is a broader category of genes that also includes published gene-disease relationships with Limited gene-disease validity and genes that have never been implicated in disease.

- *Novel gene discovery* - the process by which variant(s) in a novel candidate gene are first identified in an affected individual, subsequently confirmed by a combination of identification of additional unrelated affected individuals with variant(s) in the same novel candidate gene and experimental studies, and finally cumulative knowledge of the new gene-disease relationship is shared with the broader community, typically via peer-reviewed publication. Additional unrelated affected individuals are most commonly identified via "matchmaking" using data sharing and networks like the Matchmaker Exchange.[39,40]

- *Candidate variant* – a variant being evaluated in the context of a novel gene-disease relationship for a causal role in a Mendelian condition. Such variants typically have a higher level of supporting evidence for functional disruption or are more rare observations (e.g. *de novo*) than variants considered in known disease genes.

- *Variant of uncertain significance* - a broad category of variation that includes variants of uncertain clinical significance found in known disease genes as well as candidate variants in novel disease genes

- *Reporting novel candidate genes* – inclusion of candidate variants in a novel candidate gene in a clinical test report provided to a patient and their clinician

- *Sharing novel candidate genes* – submission of candidate variants and phenotype data for a novel candidate gene to a repository (e.g. ClinVar) or network (e.g., Matchmaker Exchange) for purposes including facilitating future identification or "matchmaking" of additional individuals with variants in the same novel candidate gene or connection to functional investigations of the gene's potential biological role in disease

Results

Criteria for novel candidate gene inclusion in clinical ES/GS

We first reviewed scoring and prioritization systems previously described by diagnostic laboratories and others[31–35,37,41] for use in assessing the strength and types of evidence for novel candidate genes and to determine whether to share and/or report variants in the corresponding genes with the patient/provider. From this review, it was clear that evaluation of the types and strengths of evidence for novel candidate genes is just as complex, if not more so, than evaluation of evidence of variant pathogenicity. Indeed, some laboratories have developed extensive scoring and weighting systems for the evaluation of novel candidate

genes,[31,34,35] while others have more streamlined criteria or lean more heavily on statistical analysis and/or internal and external matchmaking.[32,33]

To reduce barriers to more widespread implementation of novel candidate gene identification and reporting, we focused on considering a limited set of criteria, optimized for currently computable attributes, to identify candidate genes that are most likely to become established as known disease genes in the near future. For example, 98% (124 out of 127) of genes newly implicated in a Mendelian condition in 2021 according to our analysis[14] of OMIM satisfy at least one of these criteria (Supplementary Table 1). The three genes that did not satisfy any of these criteria underlie conditions that are adult-onset, have relatively mild health implications, exhibit wide variability in severity, and/or underlie an autosomal dominant condition caused only by dominant negative or biallelic variants (see footnote[a]). We recommend that variants in novel candidate genes meeting these criteria be shared with the community by submission to the Matchmaker Exchange (MME) and ClinVar as well as considered for appropriate reporting and return to patients/providers by way of inclusion on clinical reports, as summarized in Figure 1.

Recommended criteria to identify novel candidate genes to be assessed, reported, and shared. Some scenarios may not be possible to satisfy in a proband-only analysis:

1. *A gene in which predicted loss of function (pLOF) (nonsense, frameshift, essential splice site, whole or partial gene deletion or other structural variant that disrupts the coding region of the gene) or missense variant is observed to be de novo in a proband and the gene is strongly constrained for pLOF and/or missense variation. For proband-only analyses, strongly constrained genes with heterozygous pLOF variants absent from*

population databases[42] should be considered as well. See footnote[a] and Figure 1.

Overview of recommended venues through which laboratories can share and report novel candidate genes. Each venue (Matchmaker Exchange, ClinVar, and

---

[a] Gene constraint is a measure of the degree of deficit of a type of variant (e.g. pLOF, missense) among individuals in a population database, where a greater deficit is suggestive of a gene with little tolerance for the variant type. In gnomAD, the observed/expected ratios[42] for pLOF (LOEUF), missense (MOEUF), and synonymous variants are provided along with other measures of constraint (pLI,[64] Z scores). GeneBayes s_{het} (bioRxiv doi:10.1101/2023.05.19.541520) and UNEECON-g[65] are two other estimates of gene-level strength of selection against LOF variants and missense variants, respectively, that use machine learning to enable more sensitive predictions of constraint.[65] Fine-grained measures of constraint can detect subgenic/regional [Gnocchi[66], Constrained Coding Regions (CCR)[67]] or even single amino acid [MPC (bioRxiv doi:10.1101/148353), HMC (medRxiv doi:10.1101/2022.02.16.22271023)] constraint.

Constraint metrics are most sensitive for genes when the pathogenic mechanism is LOF, the condition is early-onset, and the condition has at least moderate effects on reproductive fitness. In contrast, these metrics have reduced sensitivity when the pathogenic mechanism is typically dominant negative (DN) or gain-of-function (GOF) effects,[68,69] which are often due to missense variants but can also be caused by variants that appear to be pLOF but escape nonsense-mediated decay.[70,71] Such genes (e.g. *HRAS, TRPC6, POLR3B, PRSS1, HSPB1*) often do not exhibit constraint for pLOF or missense variation according to some or even all currently-available metrics as listed in criteria 1, and GOF and DN variants are also often not predicted as damaging by computational variant effect predictors. pLOF variants in novel candidate genes should be examined carefully as such variants are more likely to not result in true LOF.[72] For example, the variant may be found in an exon that is alternatively transcribed and not required for function.[42,73] Genes that underlie conditions with modest or low effect on reproductive fitness (e.g. *C9ORF72* underlying amyotrophic lateral sclerosis and *TYR* underlying oculocutaneous albinism are not constrained for pLOF or missense variation) also exhibit less constraint.

clinical test reports) has a different design, intended audience, and purpose, so it is not redundant to share variants in a novel candidate gene across all three. In fact, sharing across all three venues is necessary to maximize the likelihood that a novel candidate gene will be confirmed quickly.

2. **Table *1*** for further discussion of constraint.
3. A gene in which biallelic variants were identified in an individual/family with a condition seemingly inherited in an autosomal recessive pattern, and there are few or no individuals in population databases with similarly rare, predicted deleterious variants *in trans* in the gene. One resource for assessing the frequency of such individuals is the "variant co-occurrence" table in gnomAD,[43,44] which uses in-silico phasing to predict total counts of individuals with pairs of rare missense or pLOF variants that are *in cis*, *in trans*, or with undetermined phase. Requiring at least one of the variants in the individual/family being tested to be pLOF would increase stringency of this criteria to further select only the most promising candidate genes.
4. *A gene in which a heterozygous pLOF (including structural variant that disrupts the* coding region) or missense variant was observed in a multiplex family with inheritance pattern consistent with a dominant inheritance, for which the number of segregations would reach a strong threshold of evidence,[45] and the gene exhibits extreme constraint and is supported by functional evidence (e.g. animal model or known biological function

consistent with phenotype).[52] See footnote[b] and Figure 1. **Overview of recommended venues through which laboratories can share and report novel candidate genes.** Each venue (Matchmaker Exchange, ClinVar, and clinical test reports) has a different

---

[b] Gene constraint is a measure of the degree of deficit of a type of variant (e.g. pLOF, missense) among individuals in a population database, where a greater deficit is suggestive of a gene with little tolerance for the variant type. In gnomAD, the observed/expected ratios[42] for pLOF (LOEUF), missense (MOEUF), and synonymous variants are provided along with other measures of constraint (pLI,[64] Z scores). GeneBayes s_{het} (bioRxiv doi:10.1101/2023.05.19.541520) and UNEECON-g[65] are two other estimates of gene-level strength of selection against LOF variants and missense variants, respectively, that use machine learning to enable more sensitive predictions of constraint. Fine-grained measures of constraint can detect subgenic/regional [Gnocchi[66], Constrained Coding Regions (CCR)[67]] or even single amino acid [MPC (bioRxiv doi:10.1101/148353), HMC (medRxiv doi:10.1101/2022.02.16.22271023)] constraint.

Constraint metrics are most sensitive for genes when the pathogenic mechanism is LOF, the condition is early-onset, and the condition has at least moderate effects on reproductive fitness. In contrast, these metrics have reduced sensitivity when the pathogenic mechanism is typically dominant negative (DN) or gain-of-function (GOF) effects,[68,69] which are often due to missense variants but can also be caused by variants that appear to be pLOF but escape nonsense-mediated decay.[70,71] Such genes (e.g. *HRAS, TRPC6, POLR3B, PRSS1, HSPB1*) often do not exhibit constraint for pLOF or missense variation according to some or even all currently-available metrics as listed in criteria 1, and GOF and DN variants are also often not predicted as damaging by computational variant effect predictors. pLOF variants in novel candidate genes should be examined carefully as such variants are more likely to not result in true LOF.[72] For example, the variant may be found in an exon that is alternatively transcribed and not required for function.[42,73] Genes that underlie conditions with modest or low effect on reproductive fitness (e.g. *C9ORF72* underlying amyotrophic lateral sclerosis and *TYR* underlying oculocutaneous albinism are not constrained for pLOF or missense variation) also exhibit less constraint.

design, intended audience, and purpose, so it is not redundant to share variants in a novel candidate gene across all three. In fact, sharing across all three venues is necessary to maximize the likelihood that a novel candidate gene will be confirmed quickly.

5. **Table *1*** for further discussion of constraint.

6. Variant(s) in a novel candidate gene for which evidence has been gathered through collection of other cases, with or without functional studies, but the data have not yet been published. Or variant(s) in a novel candidate gene for which a convincing animal model with matching zygosity exists.

7. Variants in a known gene for which the known disease phenotype does not overlap with a proband's clinical findings, but for which the variant-phenotype combination could represent a phenotypic expansion or novel gene-disease relationship.[46] We recognize that such a judgement is difficult in the absence of corroborating evidence (e.g., animal model with a similar phenotype), so we recommend that such genes be reported and shared if they meet at least one of criteria 1-4.

Rationale for sharing and reporting of variants in novel candidate genes

There are myriad benefits to reporting variants in novel candidate genes to clinicians and patients (and research participants) and to sharing these genes with the community despite the uncertainty of dealing with a novel candidate gene. Indeed, in terms of clinical decision-making, there may be little difference in the range of uncertainty between reporting a VUS in a known disease gene versus a candidate variant in a novel candidate gene. In both scenarios,

uncertainty exists and reporting the result enables the clinician and family to collect evidence for or against a causal relationship between the variant and the patient's phenotype. For example, reasons to routinely report VUS in known disease genes include the existence of established functional tests (e.g. metabolic assays) that can confirm or exclude the VUS as partially or fully explanatory of a patient's clinical findings or consideration of management approaches (e.g., drug therapies, tumor surveillance, etc.) that may be low risk but of potential impact even if the VUS turns out to be benign. It is also worth noting that both VUS[47,48] and variants in novel candidate genes identified in underserved populations[14] are less likely to have been previously reported let alone had evidence helpful for adjudication of pathogenicity generated. Failing to report these results risks further missed opportunities to build equity in rare disease diagnosis. Reporting a variant in a novel candidate gene also enables the provider and/or family to participate in research efforts to establish a novel gene-disease relationship through robust platforms like the Matchmaker Exchange (MME) and ClinVar that facilitate global data sharing and the ability to rapidly collect evidence in support of a candidate variant or gene via the process of "matchmaking" or "matching".[39,49,50]

The speed and frequency with which connections to gene discovery research efforts can be made and the potential clinical impact to individual families of being connected with such research efforts should not be underestimated. Members of our group have been involved in multiple cases in which entry of the gene into MME resulted in an immediate match with a cohort described in a draft manuscript describing a novel gene discovery. Such matches enable providers and families to take advantage of results prior to publication (see Box 1). These are not rare occurrences: recently, a research group described matching immediately upon submission to MME to an existing cohort with phenotypic and genetic similarities for almost ~10% of their novel candidate genes[51] and 15% of their submissions led to publication, with more in progress at the time of publication. Some families who received reports and were returned information about their novel candidate gene prior to publication have even launched

support groups, fundraising efforts, and research into therapeutic options concurrently with

researcher efforts to formally publish their novel gene discovery (e.g. *FAM177A1*

[https://www.fam177a1.org]; *MAST2* and *MAST4* [https://mastgenes.org]). Thus, the benefits of

identifying, reporting, and returning novel candidate variants and genes can be meaningful and

immediate.

Box 1. Examples in which sharing of variants in novel candidate genes enabled reporting and return of results.	
A family with pontocerebellar hypoplasia and microcephaly underwent exome sequencing which identified compound heterozygosity for a missense and pLOF variant in *PPIL1*, a novel candidate gene. The variants were prioritized because they were in *trans* and one was pLOF and the gene was prioritized because no individuals in gnomAD have rare biallelic variants with at least one being pLOF. The variants and gene were submitted to MME, yielding >5 matches, including connection to investigators with a manuscript asserting a novel gene-disease relationship already under review.[52] This quickly led to return of the novel candidate gene to the family.	A family with developmental delay and Kabuki syndrome-like facial features underwent exome sequencing which identified a *de novo* missense variant in *KDM1A*, a novel candidate gene. The *KDM1A* variant was prioritized because it was predicted pathogenic by multiple algorithms, the gene exhibits both missense and pLOF constraint, and because variants in other genes in the same pathway lead to similar phenotypes. Return of the candidate variant as a VUS to the family enabled them to create a website and engage in matchmaking themselves, followed by collaboration with researchers to publish the novel gene-disease relationship.[53,54]

In addition to benefiting individual families, efforts by laboratories to share and report

novel candidate genes facilitate the process of novel gene discovery by alerting the broader

community of the existence of an individual with a novel candidate gene. Such alerts are

important because current gene discovery approaches are highly dependent upon matchmaking

to build evidence for a candidate gene. Consequently, laboratories cannot rely upon reanalysis

of past cases to trigger identification of variants in newly-published known genes, because if

such genes and the variants they harbor are never shared, the gene-disease relationships may

never become known.

As the number of persons with suspected rare genetic diseases who undergo clinical

testing increasingly dwarfs the number of individuals sequenced through gene discovery

studies, there is a risk that the process of establishing new gene-disease relationships will slow unless novel candidate genes identified via clinical testing can be incorporated into research discovery efforts. Ultimately, facilitating gene discovery and the subsequent establishment of a new gene-disease relationship should improve diagnostic rates. For example, one clinical-research group recently reported[55] that among 33 candidate genes/variants identified by their program, 16 were established as pathogenic and explanatory in an average of under three years, 15/33 continued to be candidates, and only 2/33 were ultimately excluded from consideration, suggesting that the yield of "real" diagnoses from novel candidate genes is high. Finally, many large-scale clinical and hybrid clinical-research genomic testing efforts are already identifying, sharing, and reporting results in novel candidate genes and have described substantial benefits in identifying a genetic diagnosis for their patients.[28,31–33,51,56] Thus we strongly encourage clinical laboratories and research programs to proactively identify, share, and report variants in novel candidate genes.

Guidance for sharing and reporting of variants in novel candidate genes

We encourage laboratories to adopt a tiered approach to the reporting and sharing of novel candidate genes. While we recommend that all novel candidate genes meeting at least one of the criteria described above be shared and reported, laboratories and research groups with sufficient resources[51] and motivation may consider applying a more lenient threshold to identify and share additional variants in novel candidate genes via MME and ClinVar compared to what might be reported to an ordering provider. These genes could then be offered for return to families once further supporting evidence was available. Examples of novel candidate genes/variants that meet such a lower threshold include variants with low pathogenicity predictor scores[57] and/or variants in genes with moderate or low pLOF/missense constraint scores[58] in genes with known pathway/animal model evidence.

Multiple resources support sharing of novel candidate genes (Table 2). Laboratories should at minimum share all of the following information, where supported, via each resource to facilitate efficient communication and use of the shared data for future research: gene, variant(s) (transcript ID with cDNA and amino acid change and/or genomic coordinates with genome build), zygosity (when providing case-level data), medium-to-high level structured phenotype information, and suspected mode(s) of inheritance. These data are considered minimal risk for re-identification of individuals and do not require explicit patient consent for sharing.[59–61] It is especially critical to note that when submitting to Matchmaker Exchange, the amount of effort needed to manage matching correspondence balloons enormously for all parties involved if only a gene name is shared without these additional, critical details.[51]

Guidance for choosing which platform to share candidate genes
When considering which platform to use to share candidate gene-disease relationship, several factors should be considered, as outlined here:

Matchmaker Exchange (MME). The Matchmaker Exchange[39,40] is a federated network of sites, or so-called "nodes," run by research and clinical institutions with the goal of facilitating novel gene discovery through gene-level matching. A similar federated platform is also being developed for variant-level matching.[62] MME nodes each support "matching" between cases based on a shared candidate gene, at minimum. Some nodes are institution-specific or have formal criteria limiting who may join to deposit case data, while three nodes (i.e. GeneMatcher, MyGene2, PhenomeCentral) are open to all self-declared clinicians and researchers. Two of these encourage (MyGene2) or allow (GeneMatcher) individuals with rare diseases or their families to join as well. We strongly encourage laboratories not yet submitting candidate genes to MME to join one of these nodes, which all have API implementations available for authorized users to facilitate integration with laboratory management systems. Use of MME is non-

redundant with use of ClinVar as the users of MME are enriched for groups engaged in gene discovery and MME will alert them to the existence of each new case with a matching candidate gene.

There are two main limitations of the current MME network. First, submissions to most MME nodes are not publicly-accessible, with exceptions as described above, so other users of MME are unable to learn about interest in a candidate gene unless they have also specifically submitted that gene (i.e. if two independent submitters individually submit one of the two genes that encode binding partners in an essential protein complex, they will not be notified of the others' submission). Second, while matching and notification of submitters is automated, follow-up to evaluate the strength of matches is manual (i.e. via email) and can be time-consuming.[51] We provide some specific recommendations to decrease the amount of communication required for engaging productively with MME (Box 2).

Box 2. Recommendations for productive engagement with Matchmaker Exchange

1. Case submission:
    a. Laboratories should submit their case including all of the following data elements: gene, variant(s) (transcript ID with cDNA and amino acid change and/or genomic coordinates with genome build), zygosity, and suspected mode(s) of inheritance. It is essential that medium-to-high level structured phenotype information (e.g. "intellectual disability") is also included during submission as providing this information up front greatly reduces the number of follow-up emails that will be required to evaluate the validity of a match. Including non-identifying phenotype information in this manner does not compromise patient confidentiality[59,60]
    b. Use a case-level identifier that is not identifiable but will also be recognizable to the submitter (and ideally the ordering clinician) in the future
    c. Consider using a node such as DECIPHER or MyGene2 that allows submitters to make candidate genes and submission information public to increase visibility of the candidate gene-disease relationship. If submissions are not public, then the knowledge that the candidate gene has been identified will be unavailable to stakeholders not using MME to match on that exact gene (e.g. experimentalists studying the gene, geneticists with other cases due to disruptive variants in the same pathway/gene family). Other considerations may apply in choosing a node. For more guidance, see the MME website: https://www.matchmakerexchange.org/i_am_a_clinician_laboratory.html
2. Match management:
    a. Submit cases to MME using a single email address/account representing the entire entity, e.g. matchmaker@laboratory.com. This helps to ensure continuity as matches may not occur for years after the initial submitter may have moved on and to facilitate management of communications.
    b. If responses may be delayed or conditional, the laboratory's MME email address

should have an auto-reply that details expectations about response time and the company's policy/approach/preferences for handling inquiries about prospective matches (e.g. query submitters must provide phenotype/variant/mode of inheritance about their case up front to receive a response or query submitters must intend to participate in or collect a cohort of cases for publication, not just intend to adjudicate variants in a single case).

c. Consider notifying physicians of case submission to MME, of promising matches, and/or at minimum, note in laboratory documentation that a laboratory submits candidate genes to MME. If the node enables sharing a case with a clinician (e.g. MyGene2), doing so can circumvent having to establish a workflow for notifying clinicians.

d. Laboratories should consider modifying their test requisition forms to inform providers that if providers do not respond about promising matches within a reasonable time frame, the laboratory may reach out directly to the patient if contact info is available as facilitating establishment of their candidate gene in a novel gene-disease relationship is in the patient's best interests.

e. If neither the patient nor clinician responds about a match that could be described in a future publication, the laboratory may consider providing a limited amount of detail (i.e. similar granularity of information that the laboratory typically submits to ClinVar) to facilitate establishment of the novel gene-disease relationship via publication.

ClinVar. ClinVar[50] is a widely used public database managed by the NCBI that collects and reports variant and their phenotype associations. It is free, web accessible, and easily implemented as an attribute in variant annotation pipelines. Clinical laboratories, researchers, clinicians, and patients all use ClinVar to gather information about variants and genes, which makes it an ideal location to share variants in novel candidate genes. Its nearly ubiquitous use across clinical genomics ensures that anyone who has additional cases or information about a candidate gene would be able to make a connection or discover evidence, even if not using MME. Additionally, ClinVar features, such as providing contact information for submitters and the ability to "follow" a variant of interest further facilitating rapid data sharing and evidence building. It also has the advantage that information can more easily be exchanged without requiring additional communication through the submitter, such as occurs in MME, and therefore may be a preferred method of sharing data for low resourced laboratories.

All variants in candidate genes with limited or lower evidence should be classified and submitted to ClinVar as uncertain significance (VUS), which includes novel candidate genes,

while variants in genes with moderate evidence can be classified no higher than Likely

Pathogenic. To improve candidate gene variant submissions to ClinVar, we propose providing a

brief summary, as follows, stating that the gene is currently a candidate for the phenotype,

providing the relevant phenotypic features, noting any other relevant variants observed (e.g.

those in trans) and noting the zygosity of any variant observed in the case.

Example language for evidence summaries when sharing a novel candidate gene in ClinVar:

> The {zygosity} {AminoAcidChange} variant in {Gene} was identified by our
> study/laboratory in [#] individual(s) with {Phenotype description}. Evidence of variant
> disruption includes {X}. While this gene is still lacking sufficient evidence to establish a
> gene-disease relationship, evidence suggests this is a possible novel gene candidate for
> {Disease Name/Phenotype description}. Given the limited information about this gene-
> disease relationship, the significance of the {AminoAcidChange} variant is uncertain. If
> you have any additional variant evidence or have observed other individuals with this
> phenotype that also have variants in {Gene} we encourage you to reach out to us.

Laboratories are encouraged to be maximally inclusive towards submitting candidate

genes to MME and ClinVar to increase the chances of building more evidence supporting or

refuting a candidate gene, even if the gene is not included on the corresponding clinical report.

Test report. Generating a clinical report describing the identification of a novel candidate gene

ensures that a patient has the opportunity to benefit from the information and therefore reporting

the novel candidate gene empowers them and their clinical team. The report endures as part of

the medical record even if the laboratory closes, the laboratory does not proactively issue

updated reports, and/or the patient's record/data (e.g. in MME or ClinVar) are no longer

maintained by the laboratory. We suggest these findings be reported in the same location of the

report as other VUSs but be described with language that marks them as distinct from VUSs in

genes with known disease relationships per the example templates below.

Example language for including a novel candidate gene on a report:

- This variant was not identified in either of the biological parents of this individual and therefore occurred de novo. Other *de novo* variants in {Gene} have been identified in [#] individuals with {Disease Name/Phenotype description} identified through the Matchmaker Exchange (publication pending). However, this knowledgebase is still emerging and is insufficient to definitively establish the association between {Gene} and {Disease Name/Phenotype description}. In summary, additional evidence is needed to determine if {variant} may play a role in this individual's phenotypic presentation.
- This variant was not identified in either of the biological parents of this individual and therefore occurred *de novo*. The variant is predicted to result in loss-of-function (LOF) and was identified in {Gene}, which is highly constrained for loss-of-function variants (o/e=0.1 in gnomAD), suggesting that loss-of-function variants are likely to lead to a Mendelian condition. However, no other individuals have been publicly reported with LOF variants in this gene and therefore additional evidence is needed to establish the association between {Gene} and {Disease Name/Phenotype description}. In summary, this variant occurs in a candidate gene with an uncertain role in this individual's phenotypic presentation.

Discussion

To date, ~3,400 genes have at least one Moderate, Strong or Definitive gene-disease relationship in GenCC[37] with an additional ~1,400 listed in OMIM with at least one report of a gene-disease relationship. Additionally, twice as many (~10,500) genes have been predicted to be novel candidate genes but currently lack a publicly asserted gene-disease relationship.[14] However, the low prevalence of the vast majority of Mendelian conditions[63] and increasingly widespread availability of clinical genetic testing means that many, if not most, individuals with pathogenic variants in novel candidate genes are sequenced by diagnostic laboratories alone. Together, this underscores the imperative and opportunity for diagnostic laboratories to facilitate gene discovery by assessing, sharing, and reporting novel candidate genes to clinicians, families, and researchers.

The approaches we outlined for identifying and sharing novel candidate genes, and the variants within, are conservative as we recognize that handling novel candidate genes will be new terrain for many clinical laboratories. We envision these criteria will, and should, expand

over time to cover additional sets of genes and variants, as new tools and approaches for prioritizing novel candidate genes and non-coding regions and the variants within are developed. Moreover, we support laboratories with the capacity to implement more expansive criteria to do so now, and we also encourage all groups to continue to explore emerging[39,62] and as-yet-undeveloped approaches to efficiently share data to accelerate the process of implicating variants in new gene-disease relationships. Such engagement will increase the opportunities for even greater number of individuals with Mendelian condition to benefit from a precise genetic diagnosis.

Acknowledgements

The GREGoR Consortium is funded by the National Human Genome Research Institute of the National Institutes of Health, through the following grants: U01HG011758, U01HG011755, U01HG011745, U01HG011762, U01HG011744, and U24HG011746. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health.

Disclosures

All authors of this manuscript are funded by the NIH. ES is a current employee of Ambry Genetics and a stockholder of Invitae genetics. MHH is an employee of Invitae and own stock in the company. MJB is the chair of the Scientific Advisory Board of GeneDx. HLR has received rare disease research funding from Microsoft and Illumina and compensation as a past member of the scientific advisory board of Genome Medical.

Supplementary Materials

1 table

Figure Legend and Tables

Figure 1. Overview of recommended venues through which laboratories can share and report novel candidate genes. Each venue (Matchmaker Exchange, ClinVar, and clinical test reports) has a different design, intended audience, and purpose, so it is not redundant to share variants in a novel candidate gene across all three. In fact, sharing across all three venues is necessary to maximize the likelihood that a novel candidate gene will be confirmed quickly.

Table 1. Suggested constraint metrics cutoffs for selecting novel candidate genes for criteria 1 and 3. We recommend labs apply the 20% cutoff for criteria 1 to detect strongly constrained genes and the 10% cutoff for criteria 3 to detect extremely constrained genes. However, laboratories are encouraged to consider alternative cutoffs (e.g. 25% or 30%) that would identify more candidate genes as many known genes, and therefore many novel candidate genes, are expected to fall in these ranges. In addition, note that gnomAD constraint values for LOEUF and MOEUF are not comparable across different releases (i.e. gnomAD v2.1.1 vs v4.0). We provide only gnomAD v2.0 values as the interpretation of v4.0 values is still under evaluation. See footnote[a] for further details for interpreting constraint values.

metric	cutoff direction	cutoffs to identify the top X% most constrained genes			
		10%	**20%**	**25%**	**30%**
gnomAD v2.0 LOEUF	≤	0.26	0.41	0.48	0.55
GeneBayes s_{het}	≥	0.147	0.073	0.054	0.039
gnomAD v2.0 MOEUF	≤	0.58	0.70	0.73	0.77
UNEECON-g	≥	0.426	0.317	0.278	0.245

Table 2. Overview of platforms that facilitate sharing of novel candidate genes. MOI = mode of inheritance.

Resource	Description	Structured data supported for sharing					Data availability
		Gene	Variant	Zygosity	MOI	Phenotype	
ClinVar https://www.ncbi.nlm.nih.gov/clinvar/	Clinically-oriented database archiving reports of relationships between variants and human phenotypes.	Y	Y	if submitting case-level data	Y	Y	public
GenCC https://thegencc.org	Clinically-oriented database archiving reports of laboratory assessments of the strength and validity of gene-disease relationships	Y	N	N	Y	Y	public
Matchmaker Exchange (MME) https://www.matchmakerexchange.org	Research-oriented federated network of independently-run databases designed to facilitate gene discovery and genotype-phenotype relationship studies by matching submitters with the same gene/variant of interest	Y	Y	Y	Y	Y	mixed

References

1. Chong JX, Buckingham KJ, Jhangiani SN, et al. The Genetic Basis of Mendelian Phenotypes: Discoveries, Challenges, and Opportunities. *Am J Hum Genetics*. 2015;97(2):199-215. doi:10.1016/j.ajhg.2015.06.009

2. Srivastava S, Love-Nichols JA, Dies KA, et al. Meta-analysis and multidisciplinary consensus statement: exome sequencing is a first-tier clinical diagnostic test for individuals with neurodevelopmental disorders. *Genet Med*. 2019;21(11):2413-2421. doi:10.1038/s41436-019-0554-6

3. Brockman DG, Austin-Tse CA, Pelletier RC, et al. Randomized prospective evaluation of genome sequencing versus standard-of-care as a first molecular diagnostic test. *Genet Med*. 2021;23(9):1689-1696. doi:10.1038/s41436-021-01193-y

4. Chung CCY, Hue SPY, Ng NYT, et al. Meta-analysis of the diagnostic and clinical utility of exome and genome sequencing in pediatric and adult patients with rare diseases across diverse populations. *Genet Med*. 2023;25(9):100896. doi:10.1016/j.gim.2023.100896

5. Stark Z, Schofield D, Alam K, et al. Prospective comparison of the cost-effectiveness of clinical whole-exome sequencing with that of usual care overwhelmingly supports early use and reimbursement. *Genet Med*. 2017;19(8):867-874. doi:10.1038/gim.2016.221

6. Manickam K, McClain MR, Demmer LA, et al. Exome and genome sequencing for pediatric patients with congenital anomalies or intellectual disability: an evidence-based clinical guideline of the American College of Medical Genetics and Genomics (ACMG). *Genet Med*. 2021;23(11):2029-2037. doi:10.1038/s41436-021-01242-6

7. Shashi V, McConkie-Rosell A, Rosell B, et al. The utility of the traditional medical genetics diagnostic evaluation in the context of next-generation sequencing for undiagnosed genetic disorders. *Genet Med*. 2014;16(2):176-182. doi:10.1038/gim.2013.99

8. Splinter K, Adams DR, Bacino CA, et al. Effect of Genetic Diagnosis on Patients with Previously Undiagnosed Disease. *New Engl J Med*. Published online October 10, 2018:NEJMoa1714458-9. doi:10.1056/nejmoa1714458

9. Fung JLF, Yu MHC, Huang S, et al. A three-year follow-up study evaluating clinical utility of exome sequencing and diagnostic potential of reanalysis. *npj Genom Med*. 2020;5(1):37. doi:10.1038/s41525-020-00144-x

10. Yang Y, Muzny DM, Xia F, et al. Molecular Findings Among Patients Referred for Clinical Whole-Exome Sequencing. *Jama*. 2014;312(18):1870-10. doi:10.1001/jama.2014.14601

11. Posey JE, Rosenfeld JA, James RA, et al. Molecular diagnostic experience of whole-exome sequencing in adult patients. *Genet Med*. 2015;18(7). doi:10.1038/gim.2015.142

12. Retterer K, Juusola J, Cho MT, et al. Clinical application of whole-exome sequencing across clinical indications. *Genet Med*. 2015;18(7):696-704. doi:10.1038/gim.2015.148

13. Amberger JS, Bocchini CA, Schiettecatte F, Scott AF, Hamosh A. OMIM.org: Online Mendelian Inheritance in Man (OMIM(R)), an online catalog of human genes and genetic disorders. *Nucleic Acids Res*. 2015;43(Database issue):D789-98. doi:10.1093/nar/gku1205

14. Bamshad MJ, Nickerson DA, Chong JX. Mendelian Gene Discovery: Fast and Furious with No End in Sight. *Am J Hum Genetics*. 2019;105(3):448-455. doi:10.1016/j.ajhg.2019.07.011

15. Wenger AM, Guturu H, Bernstein JA, Bejerano G. Systematic reanalysis of clinical exome data yields additional diagnoses: implications for providers. *Genet Med*. 2016;19(2). doi:10.1038/gim.2016.88

16. Eldomery MK, Akdemir ZC, Harel T, et al. Lessons learned from additional research analyses of unsolved clinical exome cases. *Genome Med*. 2017;9(1):1-15. doi:10.1186/s13073-017-0412-6

17. Wright CF, McRae JF, Clayton S, et al. Making new genetic diagnoses with old data: iterative reanalysis and reporting from genome-wide data in 1,133 families with developmental disorders. *Genet Med*. 2018;20(10):1-8. doi:10.1038/gim.2017.246

18. Liu P, Meng L, Normand EA, et al. Reanalysis of Clinical Exome Sequencing Data. *N Engl J Med*. 2019;380(25):2478-2480. doi:10.1056/nejmc1812033

19. Dias KR, Shrestha R, Schofield D, et al. Narrowing the Diagnostic Gap: Genomes, Episignatures, Long-Read Sequencing and Health Economic Analyses in an Exome-Negative Intellectual Disability Cohort. *Genet Med*. Published online 2024:101076. doi:10.1016/j.gim.2024.101076

20. Tavtigian SV, Greenblatt MS, Harrison SM, et al. Modeling the ACMG/AMP variant classification guidelines as a Bayesian classification framework. *Genet Med*. 2018;20(9):1054-1060. doi:10.1038/gim.2017.210

21. Richards S, Aziz N, Bale S, et al. Standards and guidelines for the interpretation of sequence variants: a joint consensus recommendation of the American College of Medical Genetics and Genomics and the Association for Molecular Pathology. *Genet Med*. 2015;17(5):405-423. doi:10.1038/gim.2015.30

22. Walker LC, Hoya M de la, Wiggins GAR, et al. Using the ACMG/AMP framework to capture evidence related to predicted and observed impact on splicing: Recommendations from the ClinGen SVI Splicing Subgroup. *Am J Hum Genet*. 2023;110(7):1046-1067. doi:10.1016/j.ajhg.2023.06.002

23. Pejaver V, Byrne AB, Feng BJ, et al. Calibration of computational tools for missense variant pathogenicity classification and ClinGen recommendations for PP3/BP4 criteria. *Am J Hum Genet*. 2022;109(12):2163-2177. doi:10.1016/j.ajhg.2022.10.013

24. Amendola LM, Jarvik GP, Leo MC, et al. Performance of ACMG-AMP Variant-Interpretation Guidelines among Nine Laboratories in the Clinical Sequencing Exploratory Research Consortium. *Am J Hum Genetics*. 2016;98(6). doi:10.1016/j.ajhg.2016.03.024

25. Ellingford JM, Ahn JW, Bagnall RD, et al. Recommendations for clinical interpretation of variants found in non-coding regions of the genome. *Genome Med*. 2022;14(1):73. doi:10.1186/s13073-022-01073-3

26. Austin-Tse CA, Jobanputra V, Perry DL, et al. Best practices for the interpretation and reporting of clinical whole genome sequencing. *npj Genom Med*. 2022;7(1):27. doi:10.1038/s41525-022-00295-z

27. Baxter SM, Posey JE, Lake NJ, et al. Centers for Mendelian Genomics: A decade of facilitating gene discovery. *Genet Med*. 2022;24(4):784-797. doi:10.1016/j.gim.2021.12.005

28. Wright CF, Campbell P, Eberhardt RY, et al. Genomic Diagnosis of Rare Pediatric Disease in the United Kingdom and Ireland. *New Engl J Med*. Published online 2023. doi:10.1056/nejmoa2209046

29. Beaulieu CL, Majewski J, Schwartzentruber J, et al. FORGE Canada Consortium: Outcomes of a 2-Year National Rare-Disease Gene-Discovery Project. *Am J Hum Genetics*. 2014;94(6):809-817. doi:10.1016/j.ajhg.2014.05.003

30. Kobren SN, Baldridge D, Velinder M, et al. Commonalities across computational workflows for uncovering explanatory variants in undiagnosed cases. *Genet Med*. 2021;23(6):1075-1085. doi:10.1038/s41436-020-01084-8

31. Towne MC, Rossi M, Wayburn B, et al. Diagnostic testing laboratories are valuable partners for disease gene discovery: 5‐year experience with GeneMatcher. *Hum Mutat*. 2022;43(6):772-781. doi:10.1002/humu.24342

32. McWalter K, Torti E, Morrow M, Juusola J, Retterer K. Discovery of over 200 new and expanded genetic conditions using GeneMatcher. *Hum Mutat*. 2022;43(6):760-764. doi:10.1002/humu.24351

33. Taylor JP, Malhotra A, Burns NJ, et al. A clinical laboratory's experience using GeneMatcher—Building stronger gene–disease relationships. *Hum Mutat*. 2022;43(6):765-771. doi:10.1002/humu.24356

34. Hagman KDF, Shinde DN, Mroske C, et al. Candidate-gene criteria for clinical reporting: diagnostic exome sequencing identifies altered candidate genes among 8% of patients with undiagnosed diseases. *Genet Med*. 2016;19(2):224-235. doi:10.1038/gim.2016.95

35. Smith ED, Radtke K, Rossi M, et al. Classification of Genes: Standardized Clinical Validity Assessment of Gene–Disease Associations Aids Diagnostic Exome Analysis and Reclassifications. *Hum Mutat*. 2017;38(5):600-608. doi:10.1002/humu.23183

36. Strande NT, Riggs ER, Buchanan AH, et al. Evaluating the Clinical Validity of Gene-Disease Associations: An Evidence-Based Framework Developed by the Clinical Genome Resource. *Am J Hum Genetics*. 2017;100(6):895-906. doi:10.1016/j.ajhg.2017.04.015

37. DiStefano MT, Goehringer S, Babb L, et al. The Gene Curation Coalition: A global effort to harmonize gene–disease evidence resources. *Genet Med*. 2022;24(8):1732-1742. doi:10.1016/j.gim.2022.04.017

38. Bean L, Funke B, Carlston CM, et al. Diagnostic gene sequencing panels: from design to report—a technical standard of the American College of Medical Genetics and Genomics (ACMG). *Genet Med*. 2020;22(3):453-461. doi:10.1038/s41436-019-0666-z

39. Boycott KM, Azzariti DR, Hamosh A, Rehm HL. Seven years since the launch of the Matchmaker Exchange: The evolution of genomic matchmaking. *Hum Mutat*. 2022;43(6):659-667. doi:10.1002/humu.24373

40. Philippakis AA, Azzariti DR, Beltran S, et al. The Matchmaker Exchange: a platform for rare disease gene discovery. *Hum Mutat*. 2015;36(10):915-921. doi:10.1002/humu.22858

41. Santen GWE, Leitch HG, Cobben J. Gene–disease relationship evidence: A clinical perspective focusing on ultra☐rare diseases. *Hum Mutat*. Published online 2022. doi:10.1002/humu.24367

42. Karczewski KJ, Francioli LC, Tiao G, et al. The mutational constraint spectrum quantified from variation in 141,456 humans. *Nature*. 2020;581(7809):434-443. doi:10.1038/s41586-020-2308-7

43. Gudmundsson S, Singer☐Berk M, Watts NA, et al. Variant interpretation using population databases: Lessons from gnomAD. *Hum Mutat*. 2022;43(8):1012-1030. doi:10.1002/humu.24309

44. Guo MH, Francioli LC, Stenton SL, et al. Inferring compound heterozygosity from large-scale exome sequencing data. *Nat Genet*. 2024;56(1):152-161. doi:10.1038/s41588-023-01608-3

45. Biesecker LG, Byrne AB, Harrison SM, et al. ClinGen guidance for use of the PP1/BS4 co-segregation and PP4 phenotype specificity criteria for sequence variant pathogenicity classification. *Am J Hum Genet*. Published online 2023. doi:10.1016/j.ajhg.2023.11.009

46. Wilkie AOM. Many faces of SMCHD1. *Nat Genet*. 2017;49(2):176-178. doi:10.1038/ng.3776

47. Appelbaum PS, Burke W, Parens E, et al. Is there a way to reduce the inequity in variant interpretation on the basis of ancestry? *Am J Hum Genetics*. 2022;109(6):981-988. doi:10.1016/j.ajhg.2022.04.012

48. Chen E, Facio FM, Aradhya KW, et al. Rates and Classification of Variants of Uncertain Significance in Hereditary Disease Genetic Testing. *JAMA Netw Open*. 2023;6(10):e2339571. doi:10.1001/jamanetworkopen.2023.39571

49. Hartley T, Lemire G, Kernohan KD, Howley HE, Adams DR, Boycott KM. New Diagnostic Approaches for Undiagnosed Rare Genetic Diseases. *Annu Rev Genom Hum G*. 2020;21(1):1-22. doi:10.1146/annurev-genom-083118-015345

50. Landrum MJ, Lee JM, Benson M, et al. ClinVar: improving access to variant interpretations and supporting evidence. *Nucleic Acids Res*. 2017;46(Database issue):gkx1153-. doi:10.1093/nar/gkx1153

51. Osmond M, Hartley T, Dyment DA, et al. Outcome of over 1500 matches through the Matchmaker Exchange for rare disease gene discovery: The 2-year experience of Care4Rare Canada. *Genet Med*. 2022;24(1):100-108. doi:10.1016/j.gim.2021.08.014

52. Chai G, Webb A, Li C, et al. Mutations in Spliceosomal Genes PPIL1 and PRP17 Cause Neurodegenerative Pontocerebellar Hypoplasia with Microcephaly. *Neuron*. 2021;109(2):241-256.e9. doi:10.1016/j.neuron.2020.10.035

53. Tunovic S, Barkovich J, Sherr EH, Slavotinek AM. De novo ANKRD11 and KDM1A gene mutations in a male with features of KBG syndrome and Kabuki syndrome. *Am J Med Genet A*. 2014;164A(7):1744-1749. doi:10.1002/ajmg.a.36450

54. Chong JX, Yu JH, Lorentzen P, et al. Gene discovery for Mendelian conditions via social networking: de novo variants in KDM1A cause developmental delay and distinctive facial features. *Genet Med*. 2016;18(8):788-795. doi:10.1038/gim.2015.161

55. Hills S, Li Q, Madden JA, et al. High number of candidate gene variants are identified as disease causing in a period of 4 years. *Am J Méd Genet Part A*. Published online 2023. doi:10.1002/ajmg.a.63509

56. Greene D, Consortium GER, Pirri D, et al. Genetic association analysis of 77,539 genomes reveals rare disease etiologies. *Nat Med*. 2023;29(3):679-688. doi:10.1038/s41591-023-02211-z

57. Mao D, Reuter CM, Ruzhnikov MRZ, et al. De novo EIF2AK1 and EIF2AK2 Variants Are Associated with Developmental Delay, Leukoencephalopathy, and Neurologic Decompensation. *Am J Hum Genet*. 2020;106(4):570-583. doi:10.1016/j.ajhg.2020.02.016

58. Chong JX, Talbot JC, Teets EM, et al. Mutations in MYLPF Cause a Novel Segmental Amyoplasia that Manifests as Distal Arthrogryposis. *Am J Hum Genetics*. 2020;107(2):293-310. doi:10.1016/j.ajhg.2020.06.014

59. Azzariti DR, Riggs ER, Niehaus A, et al. Points to consider for sharing variant-level information from clinical genetic testing with ClinVar. *Mol Case Stud*. 2018;4(1):a002345. doi:10.1101/mcs.a002345

60. Dyke SOM, Knoppers BM, Hamosh A, et al. "Matching" consent to purpose: The example of the Matchmaker Exchange. *Hum Mutat*. 2017;38(10):1281-1285. doi:10.1002/humu.23278

61. Wright CF, Ware JS, Lucassen AM, et al. Genomic variant sharing: a position statement. *Wellcome Open Res*. 2019;4:22. doi:10.12688/wellcomeopenres.15090.2

62. Rodrigues E da S, Griffith S, Martin R, et al. Variant☐level matching for diagnosis and discovery: Challenges and opportunities. *Hum Mutat*. 2022;43(6):782-790. doi:10.1002/humu.24359

63. Wakap SN, Lambert DM, Olry A, et al. Estimating cumulative point prevalence of rare diseases: analysis of the Orphanet database. *Eur J Hum Genet*. 2019;28(2):165-173. doi:10.1038/s41431-019-0508-0

64. Lek M, Karczewski KJ, Minikel EV, et al. Analysis of protein-coding genetic variation in 60,706 humans. *Nature*. 2016;536(7616):285-291. doi:10.1038/nature19057

65. Huang YF. Unified inference of missense variant effects and gene constraints in the human genome. *Plos Genet*. 2020;16(7):e1008922. doi:10.1371/journal.pgen.1008922

66. Chen S, Francioli LC, Goodrich JK, et al. A genomic mutational constraint map using variation in 76,156 human genomes. *Nature*. 2024;625(7993):92-100. doi:10.1038/s41586-023-06045-0

67. Havrilla JM, Pedersen BS, Layer RM, Quinlan AR. A map of constrained coding regions in the human genome. *Nat Genet*. 2018;75(1):1-12. doi:10.1038/s41588-018-0294-6

68. Gerasimavicius L, Livesey BJ, Marsh JA. Loss-of-function, gain-of-function and dominant-negative mutations have profoundly different effects on protein structure. *Nat Commun*. 2022;13(1):3895. doi:10.1038/s41467-022-31686-6

69. Backwell L, Marsh JA. Diverse Molecular Mechanisms Underlying Pathogenic Protein Mutations: Beyond the Loss-of-Function Paradigm. *Annu Rev Genom Hum Genet*. 2022;23(1):475-498. doi:10.1146/annurev-genom-111221-103208

70. Akdemir ZC, White JJ, Song X, et al. Identifying Genes Whose Mutant Transcripts Cause Dominant Disease Traits by Potential Gain-of-Function Alleles. *Am J Hum Genetics*. 2018;103(2):171-187. doi:10.1016/j.ajhg.2018.06.009

71. Torene RI, Sacoto MJG, Millan F, et al. Systematic analysis of variants escaping nonsense-mediated decay uncovers candidate Mendelian diseases. *Am J Hum Genet*. Published online 2023. doi:10.1016/j.ajhg.2023.11.007

72. Singer-Berk M, Gudmundsson S, Baxter S, et al. Advanced variant classification framework reduces the false positive rate of predicted loss-of-function variants in population sequencing data. *Am J Hum Genet*. 2023;110(9):1496-1508. doi:10.1016/j.ajhg.2023.08.005

73. Cummings BB, Karczewski KJ, Kosmicki JA, et al. Transcript expression-aware annotation improves rare variant interpretation. *Nature*. 2020;581(7809):452-458. doi:10.1038/s41586-020-2329-2

Members of the [Genomics Research to Elucidate the Genetics of Rare Diseases (GREGoR)](Consortium):

Siwaar Abouhala[1], Jessica Albert[2], Miguel Almalvez[3], Raquel Alvarez[4], Mutaz Amin[1], Peter Anderson[5], Swaroop Aradhya[6], Euan Ashley[4], Themistocles Assimes[4], Light Auriga[6], Christina Austin-Tse[1], Mike Bamshad[5], Hayk Barseghyan[2], Samantha Baxter[1], Sairam Behera[7], Shaghayegh Beheshti[7], Gill Bejerano[4], Seth Berger[2], Jon Bernstein[4], Sabrina Best[5], Benjamin Blankenmeister[1], Elizabeth Blue[5], Eric Boerwinkle[8], Emily Bonkowski[5], Devon Bonner[4], Philip Boone[9], Miriam Bornhorst[2], Harrison Brand[1], Kati Buckingham[5], Daniel Calame[7], Jennefer Carter[4], Silvia Casadei[5], Lisa Chadwick[10], Clarisa Chavez[4], Ziwei Chen[4], Ivan Chinn[7], Jessica Chong[5], Zeynep Coban-Akdemir[8], Andrea J. Cohen[2], Sarah Conner[5], Matthew Conomos[5], Karen Coveler[7], Ya Allen Cui[3], Sara Currin[10], Robert Daber[6], Zain Dardas[7], Colleen Davis[5], Moez Dawood[7], Ivan de Dios[3], Celine de Esch[9], Meghan Delaney[2], Emmanuele Delot[2], Stephanie DiTroia[1], Harsha Doddapaneni[7], Haowei Du[7], Ruizhi Duan[7], Shannon Dugan-Perez[7], Nhat Duong[2], Michael Duyzend[11], Evan Eichler[5], Sara Emami[4], Jamie Fraser[2], Vincent Fusaro[3], Miranda Galey[5], Vijay Ganesh[1], Brandon Garcia[7], Kiran Garimella[1], Richard Gibbs[7], Casey Gifford[4], Amy Ginsburg[5], Page Goddard[4], Stephanie Gogarten[5], Nikhita Gogate[7], William Gordon[5], John E. Gorzynski[4], William Greenleaf[4], Christopher Grochowski[7], Emily Groopman[12], Rodrigo Guarischi Sousa[4], Sanna Gudmundsson[1], Ashima Gulati[2], Stacey Hall[1], William Harvey[5], Megan Hawley[6], Ben Heavner[5], Martha Horike-Pyne[5], Jianhong Hu[7], Yongqing Huang[1], James Hwang[7], Gail Jarvik[5], Tanner Jensen[4], Shalini Jhangiani[7], David Jimenez-Morales[4], Christopher Jin[4], Ahmed K. Saad[7], Amanda Kahn-Kirby[3], Jessica Kain[4], Parneet Kaur[7], Laura Keehan[4], Susan Knoblach[2], Arthur Ko[2], Anshul Kundaje[4], Soumya Kundu[4], Samuel M. Lancaster[4], Katie Larsson[1], Arthur Lee[1], Gabrielle Lemire[1], Richard Lewis[7], Wei Li[3], Yidan Li[7], Pengfei Liu[7], Jonathan LoTempio[3], James (Jim) Lupski[7], Jialan Ma[1], Daniel MacArthur[1], Medhat Mahmoud[7], Nirav Malani[6], Brian Mangilog[1], Dana Marafi[13], Sofia Marmolejos[2], Daniel Marten[1], Eva Martinez[1], Colby Marvin[5], Shruti Marwaha[4], Francesco Kumara Mastrorosa[5], Dena Matalon[4], Susanne May[5], Sean McGee[5], Lauren Meador[4], Heather Mefford[14], Hector Rodrigo Mendez[4], Alexander Miller[4], Danny E. Miller[5], Tadahiro Mitani[15], Stephen Montgomery[4], Mariana Moyses[9], Chloe Munderloh[7], Donna Muzny[7], Sarah Nelson[5], Thuy-mi P. Nguyen[4], Jonathan Nguyen[4], Robert Nussbaum[6], Keith Nykamp[6], William O'Callaghan[6], Emily O'Heir[1], Melanie O'Leary[1], Jeren Olsen[4], Ikeoluwa Osei-Owusu[1], Anne

O'Donnell-Luria[1], Evin Padhi[4], Lynn Pais[1], Miao Pan[2], Piyush Panchal[7], Karynne Patterson[5], Sheryl Payne[5], Davut Pehlivan[16], Paul Petrowski[4], Alicia Pham[1], Georgia Pitsava[2], Astaria/Sara Podesta[4], Sarah Ponce[7], Elizabeth Porter[4], Jennifer Posey[7], Jaime Prosser[5], Thomas Quertermous[4], Archana Rai[7], Arun Ramani[6], Heidi Rehm[1], Chloe Reuter[4], Jason Reuter[6], Matthew Richardson[5], Andres Rivera-Munoz[7], Oriane Rubio[4], Aniko Sabo[7], Monica Salani[1], Kaitlin Samocha[1], Alba Sanchis-Juan[1], Sarah Savage[6], Evette Scott[7], Stuart Scott[4], Fritz Sedlazeck[7], Gulalai Shah[1], Ali Shojaie[5], Mugdha Singh[1], Kevin Smith[4], Josh Smith[5], Hana Snow[1], Michael Snyder[4], Kayla Socarras[1], Lea Starita[5], Brigitte Stark[4], Sarah Stenton[12], Andrew Stergachis[5], Adrienne Stilp[5], V. Reid Sutton[16], Jui-Cheng Tai[9], Michael (Mike) Talkowski[1], Christina Tise[4], Catherine (Cat) Tong[5], Philip Tsao[4], Rachel Ungar[4], Grace VanNoy[1], Eric Vilain[3], Isabella Voutos[4], Kim Walker[7], Chia-Lin Wei[5], Ben Weisburd[1], Jeff Weiss[5], Chris Wellington[10], Ziming Weng[4], Emily Westheimer[6], Marsha Wheeler[5], Matthew Wheeler[4], Laurens Wiel[4], Michael Wilson[1], Monica Wojcik[1], Quenna Wong[5], Changrui Xiao[3], Rachita Yadav[1], Qian Yi[5], Bo Yuan[7], Jianhua Zhao[6], Jimmy Zhen[4], Harry Zhou[6]

[1]Broad Institute, [2]Children's National, [3]University of California Irvine, [4]Stanford University, [5]University of Washington, [6]Invitae, [7]Baylor College of Medicine, [8]University of Texas Health Science Center, [9]Massachusetts General Hospital; Broad Institute, [10]National Institutes of Health (NIH), [11]Broad Institute; Massachusetts General Hospital; Boston Children's Hospital, [12]Broad Institute; Boston Children's Hospital, [13]Kuwait University, [14]St. Jude Children's Research Hospital, [15]Jichi Medical University, [16]Baylor College of Medicine; Texas Children's Hospital

Clinical Testing ▸ **Variant Reporting** ▸ **Immediate Impact** ▸ **Long Term Impact**

Research

Clinical

Matchmaker Exchange (MME)

- enables cohort building or functional modeling among engaged submitters with same candidate gene
- increased certainty in patient's diagnosis if matched with existing cohort

- establish new gene-disease relationship via gene discovery publication
- define spectrum of phenotype and variants through case aggregation

Clinical laboratory identifies potentially causative genetic variants in novel candidate genes

ClinVar

- variant and reported information can be combined with newly generated evidence
- variant and information globally accessible and archivable

- catalyze other labs to submit entries building on initial evidence
- catalog variation in gene

Clinician orders genetic testing on patient with a phenotype

Clinical Laboratory Report

- variant enters medical record
- clinical reverse phenotyping possible
- patient able to participate in research and advocacy

- clinical variant reclassification available to patient for diagnosis and treatment