Evolution of a Plasmid Regulatory Circuit Ameliorates Plasmid Fitness Cost

Clinton A. Elg[1,2,3], Erin Mack[3], Michael Rolfsmeier[3], Thomas C. McLean[4],

Olivia Kosterlitz[2,3,5], Elizabeth Soderling[3], Solana Narum[1,2,3], Paul A. Rowley[3],

Christopher M. Thomas[6], Eva M. Top[1,2,3]

[1] Bioinformatics and Computational Biology Program, University of Idaho, Moscow, Idaho,

USA.

[2] Institute for Interdisciplinary Data Sciences, University of Idaho, Moscow, Idaho, USA.

[3] Department of Biological Sciences, University of Idaho, Moscow, Idaho, USA.

[4] Department of Molecular Microbiology, John Innes Centre, Norwich, UK.

[5] Biology Department, University of Washington, Seattle, Washington, USA.

[6] School of Biosciences, University of Birmingham, Edgbaston, Birmingham, UK.

Corresponding Author: Eva M. Top

evatop@uidaho.edu

ABSTRACT

Plasmids play a major role in rapid adaptation of bacteria by facilitating horizontal transfer of diverse genes, most notably those conferring antibiotic resistance. While most plasmids that replicate in a broad range of bacteria also persist well in diverse hosts, there are exceptions that are poorly understood. We investigated why a broad-host range plasmid, pBP136, originally found in clinical *Bordetella pertussis* isolates, quickly became extinct in laboratory *Escherichia coli* populations. Through experimental evolution we found that inactivation of a previously uncharacterized plasmid gene, *upf31*, drastically improved plasmid maintenance in *E. coli*. This gene inactivation resulted in decreased transcription of the global plasmid regulators (*korA*, *korB*, and *korC)* and numerous genes in their regulons. It also caused transcriptional changes in many chromosomal genes primarily related to metabolism. *In silico* analyses suggested that the change in plasmid transcriptome may be initiated by Upf31 interacting with the plasmid regulator KorB. Expression of *upf31 in trans* negatively affected persistence of pBP136Δ*upf31* as well as the closely related archetypal IncP-1β plasmid R751, which is stable in *E. coli* and natively encodes a truncated *upf31* allele. Our results demonstrate that while the *upf31* allele in pBP136 might advantageously modulate gene expression in its original host, *B. pertussis*, it has harmful effects in *E. coli*. Thus, evolution of a single plasmid gene can change the range of hosts in which that plasmid persists, due to effects on the regulation of plasmid gene transcription.

INTRODUCTION

Bacterial evolution and rapid adaptation are frequently shaped by the horizontal transfer of genes, including those transferred by extra-chromosomal DNA elements known as plasmids (Norman et al. 2009; Soucy et al. 2015). One example is the plasmid-encoded spread of antibiotic resistance, which enables bacterial pathogens to resist traditional therapeutic antibiotic treatments, leading to the rise of highly resistant "superbugs" that increasingly threaten human health (Murray et al., 2022; San Millan, 2018). Understanding how plasmids evolve to successfully transfer and maintain themselves in bacterial populations and communities is thus a critical step towards limiting the spread of antibiotic resistance.

A central metric for assessing the degree to which a plasmid remains in a bacterial population over time is termed 'plasmid persistence'. Plasmid-host pairs often differ in plasmid persistence, and even the same plasmid has been shown to persist differently in closely related species (De Gelder et al. 2007; Kottara et al. 2018). Instances of poor plasmid persistence can often be attributed to decreased fitness of the plasmid-containing bacterium relative to its plasmid-free counterpart. This fitness cost of plasmid carriage imposed on the bacterial host is due to several factors, including metabolic costs and molecular conflicts between plasmid and host machinery (Modi and Adams, 1991). Several studies have shown compensatory mutations that arise during serial batch cultivation of plasmid-host pairs; these mutations ameliorate plasmid fitness cost and thereby restore plasmid persistence. Some of the important early studies showed that such cost amelioration could occur through genetic changes on the plasmid but were unable to explore the underlying molecular mechanisms (Bouma and Lenski, 1988; Dahlberg and Chao, 2003). With the advent of next generation sequencing, compensatory mutations have been identified in specific plasmid genes, including those encoding accessory functions (Bottery

et al. 2017; Stalder et al. 2017), replication initiation (Sota et al. 2010; Hughes et al. 2012; Yano

et al. 2016), and conjugation machinery (De Gelder et al. 2008; Jordt et al. 2020; Porse et al.

2016; Yang et al. 2023). Additionally, chromosomal evolution can stabilize host-plasmid pairs

through amelioration of plasmid cost via mutations targeting global regulators (Harrison et al.

2015; Stalder et al. 2017) and putative helicases (San Millan et al. 2014; Loftie-Eaton et al.

2017). Thus, compensatory mutations that increase plasmid persistence can occur on the

plasmid, chromosome, or both (Hall et al. 2021).

Some plasmid types, such as those of the incompatibility group IncP-1 used in this study,

are known to be highly persistent across many Proteobacteria (Schmidhauser and Helinski, 1985:

Shintani et al. 2010; Yano et al. 2012, 2013; Jain and Srivastava, 2013; Klümper et al. 2015;),

with few exceptions (De Gelder et al. 2007; Kottara et al. 2018). This remarkable persistence is

aided by several conserved regions (Thorsted et al. 1998; Thomas, 2000; Sen et al. 2013) of

backbone genes, which ensure the fidelity of critical functions like replication, stable inheritance

and control, mating pair formation and conjugative DNA transfer. These backbone genes are

controlled by a complex genetic regulatory circuit with four repressors located chiefly within the

stable inheritance and control region (Bingle et al. 2005; Rajasekar et al. 2016). The regulators

are part of a negative feedback loop referred to as autogenous control (Bingle and Thomas, 2001)

that limits their own expression and that of the operons they control. This in turn allows fine-

tuned and temporal plasmid gene expression to minimize cost of plasmid carriage to the host and

prevent potential conflicts with host cellular machinery. Collectively, this regulatory network

enables IncP-1 plasmids to replicate, transfer, and be stably maintained in multiple classes of the

Proteobacteria (Schmidhauser and Helinski, 1985; Shintani et al. 2010; Yano et al. 2012, 2013);

Jain and Srivastava, 2013; Klümper et al. 2015).

Given the known broad host range of IncP-1 plasmids and their role in the spread of

antibiotic resistance (Popowska and Krawczyk-Balska, 2013) we sought to examine the

previously reported but unexplained poor persistence of subgroup IncP-1β plasmid pBP136Km

in several *Escherichia coli* strains (Sota and Top, 2008). This plasmid was previously obtained

by inserting a kanamycin resistance gene in pBP136 (Sota et al. 2007), which was found in a

Bordetella pertussis strain isolated from a lethal infection in an infant in Japan (Kamachi et al.

2006).

We report that a single genetic change in pBP136Km that occurred within 50 generations

of evolution in *E. coli* in the absence of antibiotic selection drastically increased the plasmid's

persistence. This improved persistence was explained by a decrease in the plasmid fitness cost

due to partial deletion and presumed inactivation of a plasmid-encoded gene with unknown

protein function, *upf31*. This gene was previously thought to encode a putative adenine

methylase that is associated with an intriguing set of clustered repeats specific to the IncP-1β

plasmids (Thorsted et al. 1998). The high cost of plasmid carriage caused by this *upf31* allele

was shown to extend to the archetypal IncP-1β plasmid R751 (Thorsted et al. 1998), which is

otherwise stable in *E. coli*. In pBP136Km the presence of Upf31 was associated with changes in

expression of plasmid backbone regulatory genes and chromosomal genes, which we propose

may be due to its physical interaction with plasmid regulator KorB. Our study highlights that

evolution of the intricate gene regulatory systems of self-transmissible plasmids can improve

plasmid-bacteria pairings by amelioration of plasmid fitness costs.

RESULTS

Plasmid-encoded *upf31* results in poor plasmid persistence in *E. coli* hosts.

We first sought to confirm a previous finding (Sota and Top, 2008) that plasmid pBP136Km shows poor persistence in various *E. coli* hosts. Using *E. coli* K-12 MG1655 (hereafter K-12), we performed a plasmid persistence assay entailing daily passage of triplicate cultures in non-selective conditions (*i.e.* without antibiotics). This initial assay with the ancestral host-plasmid pair showed poor plasmid persistence (Figure 1a in blue). In line with the previous study, we randomly isolated one plasmid-bearing clone from each of the triplicate populations on Day 5 and tested plasmid persistence in these clones. The persistence of the plasmid in these clones was markedly improved (clones A, B, and C in Figure 1a in red, green, magenta, respectively).

Figure 1: Plasmid evolution rapidly improved pBP136Km persistence. (a) Persistence of the ancestral plasmid pBP136Km (blue) and three populations (red, green, magenta) inoculated with clones that were isolated on Day 5 of the ancestral plasmid assay (see dotted ellipse and arrow). Clones isolated on Day 5 had greatly improved persistence compared to the ancestor. Lighter shades indicate 95% confidence interval. **(b)** Genomic map of pBP136Km in evolved Day 5 clones revealed deletions in the accessory region (green) between the *trb-tra* operons (light and dark blue, respectively).

To identify the genetic changes that may explain the observed change in plasmid

persistence, we completely sequenced the three isolated clones (Supplemental Table S1). We

found deletions within pBP136Km consistently targeting the region often associated with

accessory genes between the *tra* and *trb* operons, and clones B and C had identical plasmids

(Figure 1b). Additionally, no genetic changes were observed in the chromosomes of these three

clones. As the plasmid deletions in all three clones at least included *upf30.5* and *upf31.0*, we

suspected the presence of one or both genes to be responsible for the poor plasmid persistence in

E. coli.

We also routinely sequenced clones recovered after transferring the ancestral plasmid

between strains of *E. coli* by conjugation (*i.e.* matings) or electroporation. Among these clones

were several with deletions either internal to *upf31* or in its putative promoter (Supplemental

Table S1)*.* To test if deletion of *upf31* alone could explain improved plasmid persistence, we

compared the persistence of one of these recovered variants, named pBP136KmΔ*upf31,* to that

of the ancestral plasmid pBP136Km in the same host, *E. coli* K-12 (Figure 2a).

This selected plasmid contained a 168-base pair (bp) deletion internal to the open reading

frame of *upf31* whereas *upf30.5* was intact (Figure 1b and Supplemental Table S1). Given that

the plasmid with inactivated *upf31* was persistent for at least 10 days, this gene of unknown

function must be responsible for the very poor persistence of pBP136 (compare red versus blue

squares of Figure 2a). The improvement in plasmid persistence after inactivation of *upf31* was

also seen in other well-studied strains of *E. coli* including DH5α and JM109 (Figure 2a).

Importantly, the pBP136KmΔ*upf31* containing clone was isolated from an overnight mating

followed by plating for transconjugants on selective media (Supplemental Table S1). This

provides an example of very rapid plasmid evolution that results in remarkably improved

plasmid persistence (*i.e.* < 48 hours, see Hall et al. 2020).

Figure 2: *upf31* inactivation alone explained improved persistence in *E. coli* hosts. (a) The poor persistence of plasmid pBP136Km (blue lines) in *E. coli* strains K-12 MG1655, DH5α, and JM109 (square, circle, and diamond respectively) was markedly improved after an internal 168-bp deletion in plasmid gene *upf31* (red lines). Lighter shades indicate 95% confidence interval. **(b)** A map of pBP136KmΔ*upf31* showing the 168-bp deletion (red) within the coding sequence of *upf31*.

Host fitness was lowered in the presence of upf31 and pBP136Km.

To investigate if the improved plasmid persistence associated with *upf31* inactivation was due to amelioration of plasmid fitness costs, we employed two approaches: *1)* a comparison of the growth dynamics between the two plasmid-host pairs, with a focus on the maximum growth rate and carrying capacity, and *2)* competition assays involving plasmid-bearing and plasmid-free strains. First, we observed that K-12 containing pBP136KmΔ*upf31* had a significantly higher maximum growth rate and reached a higher final density in comparison to K-12 containing ancestral pBP136Km (Figure 3a). These results demonstrate that the presence of *upf31* encoded on pBP136Km imposes a substantial fitness cost on its host.

Figure 1: The presence of *upf31* encoded on pBP136Km lowered host fitness. **(a)** Bacterial growth in batch culture was slower (blue) when carrying pBP136Km with ancestral *upf31* compared to the evolved plasmid pBP136KmΔ*upf31* with inactivated *upf31* (red). Lighter shades indicate 95% confidence interval. (Two sample independent T-test of max growth rates, $n=10$, p-value= 2.35×10^{-3}) **(b)** With the fitness of plasmid-free K-12 normalized to 1 (dashed line), the relative fitness (w) of *E. coli* with ancestral pBP136Km (in blue) was 0.51, *i.e.,* a ~49% cost (Two sample independent T-test, $n=5$, p-value=1.70×10^{-6}). The relative fitness of evolved genotype K-12 (pBP136KmΔ*upf31*) to plasmid-free K-12 was statistically indistinguishable (Two sample independent T-test, $n=5$, p-value=0.40). Box is interquartile range, green triangle is mean, orange line is median, ***=p $\leq$ 0.001.

Next, we confirmed the results of the growth rate observations by estimating the cost of

the ancestral and evolved plasmid in competition assays. This was done by individually

competing each plasmid-bearing strain against the plasmid-free K-12. We note that direct

competition assays between identical hosts with and without a highly transmissible conjugative

plasmid such as pBP136Km can be confounded by plasmid transfer during the assays. While

IncP-1 plasmids are less efficient at transferring in liquid than on surfaces, they have been shown

to transfer at detectable rates (Zhong et al. 2010). To overcome this limitation, we developed a

novel low-density competition assay which allows competing two strains at densities too low for

appreciable plasmid transfer to occur (see Methods). Indeed, as conjugation requires cell contact,

decreasing initial densities results in decreasing cell collisions and thus plasmid transfer events

(Kosterlitz et al. 2022). The results (Fig 3b) show a fitness of K-12 (pBP136Km) relative to K-12

of only 0.51, suggesting the plasmid caused a 49% reduction in fitness. By contrast, the fitness of

K-12 (pBP136KmΔ*upf31*) was statistically indistinguishable from that of K-12. We conclude

that the persistence differences in Figure 2 can be attributed to a very large fitness cost conferred

by pBP136Km, which was alleviated by the inactivation of one plasmid gene, *upf31*.

The fitness cost imposed by *upf31* requires the presence of pBP136Km.

We next sought to understand if the expression of *upf31* directly imposed a fitness cost

on K-12 in the absence of plasmid pBP136Km. This was done by expressing *upf31* within K-12

in trans from expression vector pCW-LIC-*upf31*. Production of protein Upf31 was verified via

induction of pCW-LIC-*upf31* and observation of the associated 25.4 kDa product (Supplemental

Figure S1). As a negative control we used the non-modified vector pCW-LIC-*sacB*. The

maximum growth rates of K-12 with or without the Upf31 protein were not significantly

different (Fig 4a). This suggests that Upf31 does not impose a significant fitness cost on K-12 in

the absence of pBP136Km where it is naturally encoded. In contrast, when *upf31* was expressed

in trans in the presence of pBP136KmΔ*upf31,* the maximum growth rate was significantly

reduced compared to the same strain with the control vector (Fig 4b). Thus, we demonstrated

that Upf31 did not directly impose a fitness cost on K-12 but rather the interaction of Upf31 with

pBP136Km was responsible for the host fitness cost.

Gene *upf31* also effects persistence and fitness cost of archetype Inc-1β

plasmid R751 when complemented *in trans*.

We examined if the high cost of plasmid carriage observed with *upf31* extends to other

IncP-1β plasmids. The archetype IncP-1β plasmid R751(Thorsted et al. 1998) encodes a *upf31*

Figure 2: Upf31 requires the presence of pBP136Km to significantly reduce K-12 growth rate.
(a) In the absence of plasmid pBP136Km there was no significant difference in maximum growth
rate between *E. coli* K-12 expressing control *sacB* (purple) and *upf31* (brown) (Two sample
independent T-test, *n*=10, p-value=0.46). **(b)** K-12 (pBP136KmΔ*upf31*) with *upf31* expressed *in
trans* (orange) showed a lower maximum growth rate than K-12 (pBP136KmΔ*upf31*) with control
sacB expressed *in trans* (green) (Two sample independent T-test*, n*=10, p-value=4.23x10^{-3}). Box is
interquartile range, green triangle is mean, orange line is median, *n.s.* = p > 0.05, **=p ≤ 0.01.

homolog with a 66 base pair deletion at the 3' end compared to pBP136Km. The high

persistence of R751 in *E. coli* suggests that this allele does not negatively affect *E. coli* fitness

(Supplemental Figure S2). To test if *upf31* from pBP136Km affects K-12 (R751) when

expressed *in trans*, we compared K-12 (R751) (pCW-LIC-*upf31*) to control K-12 (R751) (pCW-

LIC-*sacB*) in terms of plasmid persistence and growth rate. Plasmid R751 showed lower

persistence and resulted in a statistically significant lower host growth rate (Supplemental Figure

S2) when pBP136Km's *upf31* is present. This suggests that the longer *upf31* allele of pBP136

has a negative effect on R751 cost and persistence.

Forty-nine IncP-1β plasmids were identified in NCBI that contain a gene with full-length

homology to the ancestral *upf31* and associated promoter in pBP136, although none have an

identical protein sequence (Supplemental Data S1). We tested the persistence of two of these

plasmids, pAKD1, isolated from forest soil (Drønen et al. 1998), and the wastewater associated

pALTS29 (Law et al. 2021). Whereas plasmid pAKD1 showed poor persistence, similar to

pBP136Km, pALTS29 demonstrated high persistence (Supplemental Figure S3). The *upf31*

gene product of plasmid pAKD1 differs from that of pBP136Km in ten amino acids, with three

of these being non-conservative substitutions (D76A, A91V, R124G – see Supplemental Table

S2). Upf31 from pALTS29 differed from that of pBP136Km in twelve amino acids, including

the same three non-conservative substitutions found in pAKD1. When comparing the Upf31

amino acid sequence between all three plasmids, pAKD1 had a unique R60Q change and

pALTS29 had three unique differences in L78R, Q82P, S86T. Plasmid pALTS29 may be worth

further analysis, especially since the wastewater environment in which it was found might have

selected plasmid variants that persist better in a range of *Enterobacteriaceae*.

The computationally predicted methylase function of Upf31 was not supported by experimental results.

To provide further insight into the possible function(s) of Upf31 we performed a bioinformatic analysis using Phyre2. The output strongly predicted Upf31 to be a Dam methylase homolog with 70% of the amino acids modeling with 100% confidence (Supplemental Data S2). This is consistent with the pBP136 reference genome including a computationally automated annotation of *upf31* as encoding a DNA methylase based on protein similarity as noted in the original annotation of archetypal IncP-1β plasmid R751 (Thorsted et al. 1998). To test if Upf31 had Dam activity, we digested genomic DNA extracted from *dam⁻* strain *E. coli* JM110 with and without pBP136Km and *dam⁺* K-12, using restriction enzymes that digest only methylated or non-methylated 'GATC' DNA sequences. The results suggest that Upf31 is not a functional Dam homolog as the restriction profiles of DNA from JM110 (pBP136Km) were opposite those of the *dam+* strain K-12 (Supplemental Figure S4).

Next, to determine if Upf31 more broadly encoded a methyltransferase, we used the base pair modification detection available via Single Molecule Real Time (SMRT) sequencing. This method has the benefit of not requiring *a priori* knowledge of the base pair modification chemistry or target DNA sequence of the putative methylase. Briefly, the methylation-free *E. coli* ER2796 (Anton et al. 2015) was used with *upf31* expressed from vector pCW-LIC-*upf31* or pCW-LIC-*sacB* as the vector control. Genomic DNA extracts of both strains were analyzed using the base modification analysis on a Pacbio Sequel II, which measures the kinetic incorporation of base pairs to detect patterns distinctive from non-methylated base pairs. The output showed no discernible kinetic patterns associated with methylated base pair incorporation,

228   indicating that Upf31 does not function as a methylase under our experimental conditions, in

229   contrast to the computational predictions.

The presence of *upf31* alters the expression of global plasmid regulators and

associated operons.

To understand how *upf31* and pBP136Km interact to influence *E. coli* fitness, we next

performed RNA-seq on K-12 containing pBP136Km and K-12 containing pBP136KmΔ*upf31*.

We considered K-12 (pBP136Km) to be the treatment (presence of the intact gene *upf31*), and K-

12 (pBP136KmΔ*upf31*) with the internal 168-bp deletion to be the control (absence of a

functional *upf31*). The presence of *upf31* led to greater than two-fold expression changes in 654

of the ~4,400 chromosomal genes, with 412 of those upregulated and 242 downregulated

(adjusted P-value < 0.05, Supplemental Data S3). The gene ontology tool DAVID (Dennis et al.

2003) predicted downregulation of cellular pathways related to nitrogen and sugar metabolisms,

while biosynthesis of siderophores, ABC transporters, and sulfur metabolism pathways were all

upregulated (Supplemental Data S4).

Most interestingly, K-12 (pBP136Km) showed greater than two-fold expression increase

in 11 of its 46 plasmid genes relative to the *upf31* deletion variant (adjusted P-value < 0.05,

Figure 5a and Supplemental Data S3). These genes were primarily part of the stable inheritance

and control region made up of regulons controlled by proteins KorA, KorB and KorC (Figure 5a

in yellow). Strikingly, K-12 (pBP136Km) showed a relative decrease in expression in only one

gene, *upf31* itself. This relative comparison of gene expression was made possible by

pBP136KmΔ*upf31* not having a complete *upf31* deletion but rather an internal 168-bp deletion

that left an inactivated 507 bp *upf31* product to be expressed (Figure 5b).

Figure 3: Expression of global plasmid regulators and their associated operons was increased in the presence of ancestral *upf31* compared to the deletion mutant. (a) Genetic map of pBP136Km with operons and operator sites annotated for global regulators KorA, KorB, KorC, and TrbA. Asterisks denote differentially expressed plasmid genes in the presence of *upf31*. **(b)** Table of plasmid-encoded genes with at least 2-fold differential expression (black asterisks in panel a) in the presence of *upf31*. Note expression of the ancestral *upf31* is ~16-fold lower than that of the truncated *upf31* knockout.

The effect of Upf31 to cause at least two-fold increased expression of all four of the

canonical IncP-1 regulatory genes (Adamczyk and Jagura-Burdzy, 2003)– *korA, korB, korC,* and

trbA – along with the other genes in the operons they regulate (Figure 5a in bold) is notable.

These regulators are known to autogenously repress their own expression (Bingle and Thomas,

2001) by binding cognate operators near their respective promoters (orange, red, green, and

black dots in Figure 5a). They are also known to repress promoters for the other operons in their

regulons (yellow in Figure 5a). The increased repressor gene expression should therefore

decrease rather than increase expression of all other regulated operons as the expression of the

repressor genes increases. This suggests that *upf31* has a general effect of diminishing the

repressive ability of *kor* regulatory genes, both in terms of their autogenous control and

repression of their respective regulons.

Upf31 represses its own transcription.

A possible mechanistic explanation for the decrease in *upf31* transcription in the presence

of Upf31 is that Upf31 represses its own transcription, like the previously IncP-1 plasmid

regulators KorA, KorB, KorC and TrbA studied (Pansegrau et al. 1994). Testing this hypothesis

requires first verifying the putative promoter upstream of upf31(*i.e.* upf31p) and then measuring

expression from this promoter with and without Upf31 present in the cells. To test the

functionality of the putative *upf31* promoter upf31p, a 68-bp region upstream of *upf31* was

inserted into *xylE* reporter plasmid pGCMT1 to create pGCMTupf31p (Supplemental Figures S5

and S6). The *xylE* assay confirmed that this region possesses significant promoter activity with

7.4 XylE units compared to the 0.09 units for the empty vector (Supplemental Table 3).

Since the Thomas lab had previously failed to demonstrate strong repressor activity from

the 3'-end truncated *upf31* in R751(Akhtar, 2002), we tested if adding the C-terminus of

pBP136's Upf31 to R751's Upf31 restored the protein into a functional repressor. To do this, we

constructed an R751/pBP136 hybrid *upf31* gene (aa 1 to 172 from R751 and aa 173 to 225 from

pBP136) and inserted this hybrid gene into pBR322 to create pBR*upf31*. Expression from the

upf31 promoter on pGCMTupf31p was then measured in the presence and absence of pBR*upf31*.

The full-length hybrid Upf31 expressed from pBR*upf31* caused repression of upf31p, resulting in

an average of 0.09 XylE units compared to 4.76 XylE units with pBR322 as negative control

(Supplemental Table 3). This confirmed that *upf31* expression is likely to be autoregulated via

Upf31 binding near upf31p. The most obvious operator-like sequence in the region of this

promoter consists of inverted 5'-CAGCATCG-3' repeats, which belongs to the family of

inverted repeats (IR) that occur in four groups across the IncP-1β genome (Thorsted et al. 1998).

The presence of these strikingly conserved sequences of these IR across IncP-1β plasmids

suggests that Upf31 may bind to similar sequences across IncP-1β plasmids. In conclusion, these

assays confirmed the transcriptome results that suggested Upf31 represses itself and may have

additional regulatory functions.

Upf31 is computationally predicted to bind KorB via a C-terminal

dimerization domain.

A hypothesis to explain our transcriptome results emerges from the fact that the high

level of repression associated with the stable inheritance and control region depends on the

cooperative interaction of KorA (and TrbA) with KorB (Zatyka et al. 1997; Kostelidou et al.

1999; Bingle et al. 2008). If Upf31 were able to disrupt this cooperativity then it might decrease

repression while allowing the concentration of the repressors to rise. To explore the hypothesis

that Upf31 disrupts the cooperative interaction of KorA (and TrbA) with KorB we used

Alphafold2 (Jumper et al. 2021) and ColabFold (Mirdita et al. 2022) to predict the structures of

every individual pBP136 coding region in complex with Upf31. We subsequently ranked these

interactions using a self-assessment ranking score (ranking_confidence: 0.2pTM + 0.8ipTM, see

Supplemental Data S5). The structure prediction for Upf31 is that it possesses a globular DNA

binding domain (residues 1 to 174) and a C-terminal dimerization domain (CTD) which is

consistent with its proposed autogenous regulation of its own expression. However, the most

exciting result of this analysis is the prediction that the CTD mediates interaction with KorB

(Figure 6a) in a way that is very similar to KorA. Modeling of Upf31 with KorB gave a robust

Figure 4: AlphaFold2 modeling reveals the potential interaction between pBP136 Upf31 and KorB is mediated by the C-terminal dimerization domain. (a) AlphaFold2 modelling of the pBP136 Upf31 homodimer (light and dark blue) with the pBP136 KorB homodimer (red and orange). **(b)** Superimposed AlphaFold2 models of pBP136 Upf31 (dark blue) and R751 Upf31 (magenta) highlighting the additional C-terminal dimerization domain of pBP136 Upf31. **(c)** Overlay of the AlphaFold2 model of pBP136 Upf31 homodimer (dark blue) with the crystal structure of the RK2 KorA homodimer (PDB: 5CM3, DNA duplex removed). Basic residues in the dimerization domain are highlighted in magenta or yellow. KorA Y84 is highlighted in deep red, Upf31 W198 is highlighted in cyan.

ranking_confidence score of 0.64. Removal of the CTD from Upf31 drastically reduced the

predicted interaction between KorB and Upf31 to 0.274. When pBP136 KorB was modelled with

R751Upf31, which natively lacks a CTD, a similarly low score of 0.252 was predicted. Given

the globular domain sequence similarity between the Upf31 of pBP136Km and R751 these

results are highly suggestive of pBP136Km Upf31 interacting with KorB via its C-terminus CTD

(Figure 6b). Although the Upf31 CTD does not line up with KorA CTD in the way that the CTD

of TrbA does, one can detect similarities like the Tyrosine at position 180 and the patch of

predominantly basic amino acids. This predicted interaction might allow Upf31 to reduce

available KorB either when bound to DNA or free, since we previously observed that KorA and

KorB could be copurified in the absence of DNA (McLean et al. 2024). This would in turn

explain the increased transcription of the IncP-1 regulatory genes as well as the operons they

regulate.

Additional experiments were performed to test the computational prediction of Upf31

requiring a CTD for functionality. Plasmid R751 encodes a truncated Upf31 allele lacking a

CTD and persisted very well in *E.* coli but was rapidly lost in the presence of Upf31/pBP136Km

expressed *in trans* (Supplemental Figure S2 and Supplemental Data S6). This suggests the CTD

of Upf31 is critical to Upf31's negative effect on plasmid persistence and growth of *E. coli.* To

test further whether it is the CTD of pBP136's Upf31 that is critical for poor plasmid persistence

in *E. coli*, rather than other amino acid differences between the two alleles, we tested the effect

of the hybrid Upf31 (described earlier) on R751. This hybrid Upf31 combines R751's truncated

Upf31 with the C-terminus of Upf31/pBP136Km and was expressed *in trans* with vector

pBR*upf31*. Strikingly, transformation of pBR*upf31* into competent *E. coli* C600 (R751) gave

hundreds of pBR*upf31* transformants on L-agar with ampicillin but less than 10 transformants on

L-agar with ampicillin and trimethoprim (selecting for both pBR*upf31* and R751). Together our

computational and experimental findings strongly suggest that the presence of the CTD of Upf31

from pBP136 is necessary for the poor plasmid persistence phenotype, and likely because of its

physical interaction with KorB.

DISCUSSION

This work sheds new light on the biology and evolution of the largest sampled subgroup

of the most intensively studied conjugative plasmids. We demonstrate that the poor persistence

in *E. coli* of a conjugative broad-host-range IncP-1β plasmid from *Bordetella pertussis* rapidly

improved, even in the absence of selection for the plasmid. The associated drastic amelioration

of the plasmid fitness cost to *E. coli* was due to the inactivation of a previously uncharacterized

plasmid gene, *upf31,* which affected the expression of the plasmid backbone regulatory circuit

including all four major regulators as well as hundreds of chromosomal genes. This change in

transcriptome may have been initiated by Upf31 interacting with one of these major plasmid

regulators, KorB. Even though plasmids like pBP136 of the incompatibility group IncP-1 are

notably persistent in many Proteobacteria (Schmidhauser and Helinski, 1985; De Gelder et al.

2007; Shintani et al. 2010; Yano et al. 2012, 2013, Jain and Srivastava, 2013; Klümper et al.

2015; Kottara et al. 2018;), we show that different alleles of plasmid genes like *upf31* can cause

striking differences in plasmid persistence in particular hosts. Understanding how this gene acts

and why it may be an advantage in some hosts will expand our understanding of the genetic

toolbox plasmids can exploit for their success.

The last decade has seen a flood of seminal studies regarding compensatory evolution of

plasmid fitness cost, including multiple studies showing that changes in chromosomal regulatory

systems can improve plasmid-host relations. One example is mutations of the *fur* transcriptional

regulator involved in iron uptake (Hassan and Troxell, 2013; Stalder et al. 2017), while evolution

of the chromosomal *gacA/S* regulators (Harrison et al. 2015) was observed to lower the fitness

cost of plasmid carriage within 48 hours of plating on agar (Hall et al. 2020). A few studies have

found compensatory evolution via putative chromosomally-encoded helicases (Loftie-Eaton et

al. 2017), in one case largely restoring gene expression levels of the plasmid-host pair to that of

the plasmid-free bacteria (San Millan et al. 2015). Amid these discoveries, the complex and

tightly regulated plasmid transcriptional regulatory systems have perhaps been overlooked as a

key factor in plasmid-host relations. One exception was a study which found that a plasmid

improved persistence via differential expression of a *parAB* operon, with backbone genes and

plasmid regulators otherwise remaining unchanged (Hall et al. 2021). Our work shows that the

plasmid encoded Upf31 inactivated during evolution in *E. coli* regulates its own transcription as

well as that of the primary plasmid regulators associated with the stable inheritance and control

region. Thus, the regulatory systems of *both* plasmids and chromosomes should be included in

the growing list of evolutionary pathways that can decrease the fitness cost of a plasmid.

The exact mechanism of action of Upf31 remains experimentally unvalidated but we

hypothesize that its C-terminal dimerization domain interacts with plasmid regulator KorB. This

was suggested by our use of AlphaFold2 to predict structures and interactions between Upf31

and all other pBP136-encoded proteins, which identified KorB as a highly likely target for Upf31

(Supplemental Data S5). Furthermore, the CTD of Upf31 modeled very similar to the CTD of

KorA, which was previously shown to interact with KorB (Bingle et al. 2008). KorB is known to

interact co-operatively with the other plasmid backbone repressors (Kostelidou et al. 1999;

Kostelidou and Thomas, 2000; Zatyka et al. 2001). Therefore, a KorB-Upf31 interaction that

reduces KorB's ability to interact with other repressors could explain the observed weakening of

KorB autogenous control as well as weaker operon repression. These changes in plasmid

transcriptional regulation were associated with greatly reduced *E. coli* host fitness and the

associated lowered plasmid persistence. It would also align with our result that plasmid R751

with a *upf31* allele that natively lacks a CTD is highly persistent in *E. coli*. In sum, our study

suggests that just like KorA, Upf31 may also belong to the family of proteins that can interact

with KorB, and that its interaction may override the other interactions with concomitant

derepression of gene expression.

The Upf31-associated derepression of global plasmid repressors triggers a secondary

effect of large-scale differential gene expression. Thus, the next question becomes which of these

differentially expressed genes explains poor *E. coli* fitness. As this negative fitness effect of

upf31 was only observed in the presence of IncP-1β plasmids, we focus here on changes in the

plasmid transcriptome. First, of the three plasmids in this study that contained full-length *upf31*

homologs, only pALTS29 showed high persistence in *E. coli,* which may suggest low fitness

cost. Notably, this plasmid lacked a toxin-antitoxin (TA) system found in nearly all IncP-1β

plasmids. This suggests that overexpression of TA genes may explain poor *E. coli* fitness.

However, the higher persistence of pALTS29 could equally be explained by differences in Upf31

amino acid sequences (Supplemental Table S2). Given our hybrid Upf31 results, there are only

two amino acid differences between the Upf31/R751/pBP136Km hybrid and Upf31/pALTS29

that could explain differences in plasmid persistence, i.e. at positions 82 and 92. Another

possible explanation for the negative effect of Upf31 on *E. coli* fitness is a set of four genes in

the control region that were upregulated in the presence of Upf31 (*kleA, kleE, klcA, klcB,* or '*kil'*

genes). These genes have been shown to have harmful effects on the growth of *E.coli* strains

when not repressed by the global Kor regulators (Figurski et al. 1982; Kornacki et al. 1993;

Larsen and Figurski, 1994), with *klcB* found harmful only in the presence of an IncP-1 plasmid

(Bhattacharyya and Figurski, 2001). Both above hypotheses assume that the observed mis-

regulation of *non*-regulatory plasmid genes cause physiological change within the bacterial host

which would explain the responsive shift in chromosomal gene expression levels. An alternative

model is that the observed mis-regulation of the plasmid repressors (KorA/B/C) also directly

changes expression of chromosomal genes in an example of plasmid-chromosome 'cross-talk' (

Vial and Hommais, 2020; Hall et al. 2021; Marincola et al. 2021; Thompson et al. 2023). Future

studies will have to identify the specific overexpressed plasmid genes that have such a negative

effect on *E. coli* fitness and by what mechanism, and if the differentially transcribed

chromosomal genes are at all responsible. It also remains to be determined whether *upf31*

provides an advantage in the original *B. pertussis* host.

Upf31 might be an important clue to understanding the four groups of conserved inverted

repeats (IR) that were previously described (Thorsted et al. 1998) in IncP-1β plasmids but

without any explanation of purpose or function. The *upf31* gene itself is embedded in one of the

four IR regions and appears to autoregulate its own expression by binding to one of the IR

regions. It is therefore likely that Upf31 binds to all the IR sequences although we currently

have no indication of purpose. However, it is noteworthy that *klcB* is among the highest

overexpressed genes in the presence of Upf31 (Figure 6) and includes several IR within its

coding sequence which may correlate with being available to compete with KorB for repression.

We next wish to report two important lessons we learned during this study. First, we

frequently observed rapid loss of *upf31* during routine laboratory cultivation due to various

deletions, prompting us to always sequence the region of *upf31* to confirm its presence. This

suggests that rapid evolution of costly plasmid genes may often go unrecognized during

cultivation and plasmid transfer between strains by matings and electroporation. Our work aligns

with previous findings of rapid (<48 hours) compensatory evolution between plasmid-host pairs,

even without known selection (Hall et al. 2020). Second, our work showed that Upf31 is likely

not a functional methylase in contrast to automated annotations describing the gene as such

(without experimental data) in many plasmid genomes. This highlights the argument that

plasmid genes *without* experimental data on their function should be assigned locus tags rather

than putative function-related gene names (Thomas et al. 2017).

In conclusion, despite decades of great progress in plasmid biology and the pivotal role of

conjugative plasmids in antibiotic resistance dissemination, a substantial portion of plasmids

harbor multiple genes with unknown functions. Our study highlights the importance of one such

gene, which when inactivated led to a lasting host-plasmid pair, effectively rescuing a plasmid

from potential extinction (Gomulkiewicz and Holt, 1995) in a bacterial population. The

investigation of this specific gene revealed that this evolutionary rescue is closely linked to

alterations in the plasmids regulatory circuit. These findings underscore the significance of

identifying and experimentally validating uncharacterized plasmid genes to understand how

evolution underwrites the spread and persistence of plasmids in bacterial communities.

MATERIALS AND METHODS

Bacterial strains, plasmids, and media

IncP-1β plasmid pBP136 was discovered in a strain of *Bordetella pertussis* isolated from

a lethal case of infant whooping cough[35]. A kanamycin gene was later inserted to generate

pBP136Km (NCBI accession number NZ_OR146256.1), which provided a selectable marker in

the originally cryptic plasmid[56]. The expression vector pCW-LIC-*upf31* was derived from pCW-

LIC-*sacB* as described below in the cloning section.

Escherichia coli K-12 MG1655 was derived[57] from an isolate within the stool of a

diphtheria patient in 1925, and *E. coli* JM109 and DH5α[58] are later derivatives of this strain with

useful attributes relevant to cloning. We generated rifampicin and nalidixic resistant clones of *E.*

coli K-12 MG1655 by recovering resistant mutants after plating on agar with the respective

antibiotic. New constructs and initial, intermediate, and final populations of the plasmid

persistence assays were all archived at -70 °C in 30% glycerol.

All bacteria in this study were grown at 37°C in lysogeny broth (LB) shaken at 200 RPM

or on LB agar (LBA) plates. Strains were grown with 50 µg ml^{-1} kanamycin (km) or 100 µg ml^{-1}

ampicillin (amp) when appropriate for plasmid maintenance. Expression vectors were induced

with 500 µM isopropyl β- d-1-thiogalactopyranoside (IPTG) when induction is mentioned in the

text.

Plasmid Persistence Assay

All plasmid persistence assays were performed in triplicate and first grown overnight

(O/N) with kanamycin to select for initial plasmid maintenance (time T0). Daily serial transfers

of 4.9 µL of overnight culture into 5 mL of sterile broth *without* antibiotics were then performed

for ten more days (times T1-T10). Each day the cultures were diluted 10^{-6} in 1x phosphate

buffered saline (PBS) and 100 uL was spread onto dilution plates. The daily plasmid-containing

fraction of the population was determined by replica plating 52 randomly chosen colonies from

the dilution plates onto LB plates with and without kanamycin.

Growth Assays

We performed growth assays in batch culture to measure the effect of Upf31 on growth

rates and final densities. Strains were grown from freezer archives for 24 hours and diluted

1:100, then grown another 24 hours and finally diluted 1:1000 in sterile media. From these ten

biological replicates each were loaded at a volume of 200uL into the respective wells of a 96-

well plate. A SPECTROstar Nano plate reader measured optical density at 600nm every ten

minutes for 23 hours with incubation at 37C and 500 RPM orbital shaking. The R package

GrowthRates (Hall et al. 2014) v. 0.8.4 was used to calculate maximum growth rate using default

parameters.

Low-Density Competition Assays

The low-density competition assay was developed in this study to avoid plasmid transfer

during the assay. It contains two distinct steps: (*i*) determining the likely "conjugation-free" time

window for specific low donor and recipient densities, and (*ii*) performing a traditional

competition assay at low densities within the established time window.

To find the likely window of time before density is high enough for conjugation to occur,

we used K-12 MG1655 strains[34] with spontaneous mutations conferring resistance to nalidixic

acid (Nal) and rifampicin (Rif) (donor K-12N(pBP136Km) and recipient K-12R, respectively).

Donor and recipient were mixed1:1 at densities of 10^2/ml into 2.5 mL LB shaken 200 RPM at

37°C. To select for possible transconjugants, the cultures were mixed hourly with another 2.5

mL of LB with Km and Rif to create a 5mL mixture with 50 μg ml^{-1} Km and 50 μg ml^{-1} Rif.

These were grown 24 hours after which the earliest timepoint showing turbidity was associated

with the emergence of transconjugants K-12R(pBP136Km). The first turbid culture emerged

after six hours, and therefore a five-hour window of likely conjugation-free growth was chosen

for the subsequent competition assays.

Competition assays were then performed using K-12 vs. K-12(pBP136Km) and K-12 vs.

K-12(pBP136KmΔ*upf31*). Each experiment began with 1:1 mixtures at densities of 10^2/ml in 5

LB without antibiotics in shaken test tubes. Plating onto LBA occurred at the start time and after

five hours (after appropriate dilutions), and the relative increase in cell numbers (cfu/ml) was

used as a proxy for relative fitness as follows: (Wiser and Lenski, 2015)

490
$$w = \frac{ln\left(\frac{A_f}{A_i}\right)}{ln\left(\frac{B_f}{B_i}\right)}$$

491   where w is the Malthusian relative fitness term, A_I and A_F are the initial and final plasmid

492   containing K-12 populations, respectively, and B_I and B_F are the initial and final plasmid-free K-

493   12 populations, respectively.

Methylation Detection

To test if *upf1* encodes a methyltransferase as computationally predicted we utilized the

base modification analysis provided by the Pacbio Sequel II DNA sequencer. The methylase-free

E. coli strain ER2796[43] with expression vector pCW-LIC-*upf31* was tested against ER2796

(pCW-LIC-*sacB*) for differential base modification[36] on a Pacbio Sequel II at the Arizona

Genomic Institute in Tucson, AZ, USA. Both strains were grown with antibiotic selection and

500 μM IPTG induction before genomic extraction using the Sigma GeneElute Bacterial kit.

Base Modification Analysis was run on SMRT Link v. 10.2.0.133434 with a reference made by

merging the *E. coli* assembly with the respective plasmids and enabling "Find Modified Base

Motifs" and "Consolidate Mapped BAMs for IGV" options. All the other options were left as

default.

RNA-Seq

To understand how *upf31* and pBP136Km interact to influence *E. coli* fitness, we

performed RNA-seq. Strains K-12WT (pBP136Km) and K-12WT (pBP136KmΔ*upf31)* were

grown overnight in 5 ml LB with Km and centrifuged. The cell pellets were sent to Zymo

Research for Total RNA-Seq Service. Sequencing libraries were constructed from total RNA

samples and were prepared using the Zymo-Seq RiboFree Total RNA Library Prep Kit (Cat #

R3000) according to the manufacturer's instructions

(https://www.zymoresearch.com/products/zymo-seq-ribofree-total-rna-library-kit). RNA-Seq

libraries were sequenced on an Illumina NovaSeq to a sequencing depth of at least 30 million

read pairs (150 base paired-end sequencing) per sample.

A custom bioinformatics pipeline began with TrimGalore! v. 0.67 to remove adapters and

low-quality reads. Bowtie2 (Langmead and Salzberg, 2012) v. 2.4.5 then mapped reads to

merged chromosomal and plasmid references. These were counted using featureCounts(Liao et

al. 2014) v. 2.0.1, the output of which was input unto DESeq2 (Love et al. 2014) v. 1.38.1, for

final calculations of relative expression.

DNA Sequencing

Whole genome sequencing was performed by SeqCenter, LLC in Pittsburg, PA, USA.

Genomic DNA extractions were performed using Sigma GeneElute Bacterial kit. The libraries

were prepared using the Illumina DNA Prep kit with IDT 10bp UDI indices and sequenced on an

Illumina NextSeq 2000 with 2x151bp reads. Demultiplexing, quality control and adapter

trimming was initially performed using Illumina's bcl-convert v. 3.9.3. TrimGalore! v. 0.6 was

additionally used to remove remaining adapters and low-quality reads. Breseq (Barrick et al.

2014) v. 0.36.0 was used to identify mutations between ancestral and evolved strains.

Cloning

pCW-LIC (hereafter pCW-LIC-*sacB*) was a gift from Cheryl Arrowsmith (Addgene

plasmid #26098; http://n2t.net/addgene:26098; RRID:Addgene_26098). Plasmid pCW-LIC-*sacB*

was double cut at the *NdeI* and *HindIII* restrictions sites, which created a 4,962 base pair (bp)

linearized backbone and a 2,302 bp linear fragment containing *sacB* and its promoter. These

were separated and the linearized backbone recovered from an agarose gel using a Thermo

Scientific (TS) GeneJET Gel Extraction Kit. Next, gene *upf31* was amplified from pBP136Km

miniprep using upstream primer 5'-*GGTGGT*CATATGTCCAGGAAGAAGGCCATGAG-3'

and downstream primer 5'- *GGTGG*TAAGCTTCTACTCGGCCGCCTCTAG-3' (flanking

sequences for restriction enzymes in italics, *NheI* and *HindIII* sites underlined, respectively). The

ends of the *upf31* amplicon were then double digested at the *NdeI* and *HindIII* sites, and the

small digested ends removed using a TS GeneJET PCR Purification Kit. The *upf31* segment and

linearized pCW-LIC backbone were joined using T4 DNA ligase and the now-circularized pCW-

LIC-*upf31* was electroporated into DH5α using standard methods(Sambrook and Russell, 2001).

To construct the upf31p-*xylE* reporter gene fusion complementary 68-nt oligomers were

designed to create a double stranded fragment with *Xba*I and *Eco*RI sticky ends for insertion into

xylE reporter plasmid pGCMT1 to create pGCMT-upf31p as shown in Supplementary Figure S6.

The 68 bp region upstream of *upf31* was from plasmid R751.

To construct the R751-pBP136 hybrid *upf31*, translational start plus codons 1 to 173 were

amplified by PCR with primers 5'-

TGCAAGCTTTAATGCGGTAGCCAAGTCCCGATTTACTCCAG-3' and 5'-

CCGGCTCGGGGTAGTTCATC-3' from R751 template. To create codons 174 to 224 from

pBP136 PCR-driven overlapping(Heckman and Pease, 2007) of synthetic oligomers (104 nt and

117 nt) were utilized with pBP136 template. The whole segment was amplified with primers 5'-

TGCAAGCTTTAATGCGGTAGCCAAG-3' and 5'-TCGGTCGACGCAGGCGTGAC-3' and

after cutting with restriction enzymes inserted into both pBR322 (Accession J01749.1) and

pACYC184 (Accession X06403.1) between their HindIII and SalI sites so that *upf31* was

transcribed from the *tetA* promoter.

xylE reporter assay

Overnight LB cultures were inoculated from single colonies of strains with reporter

plasmid pGCMT1 or its derivative pGCMTupf31p alone (with just kanamycin selection) or with

a second plasmid vector (pBR322) or vector plus the hybrid *upf31* gene (with both kanamycin

and ampicillin) and incubated for 16 h. Bacteria from 1 ml aliquots were pelleted, resuspended in

0.5 ml sonication buffer, sonicated with three bursts of 5 seconds, the cell debris cleared with 10

min centrifugation at maximum speed in a microfuge at 4°C and then measured quantitatively for

XylE activity and protein concentration as described previously (Zukowski et al. 1983).

Upf31 protein expression

Induction cultures of pCW-LIC-upf31 in *E. coli* K-12 M1655 were started from an

overnight liquid culture of LB (Lennox) with 100 µg/ml ampicillin (LB Amp). Twenty-five

milliliter LB Amp cultures were initiated at OD600 0.05 using the liquid overnight culture to

serve as the induction cultures. The 25 milliliter cultures were grown at 37° C while shaking at

200 rpm. When the 25 milliliter cultures reached OD600 > 1.0, one milliliter of culture was

removed, and the cells were harvested by centrifugation for four minutes at 16,000 x g at room

temperature in an Eppendorf 5425 centrifuge. This material served as the pre-induction (or 0

hour) time point for each culture. ITPG was then added to each induction culture to a final

concentration of 0.1 mM, 0.5 mM, or 1 mM and growth was monitored at OD600. At the time

points indicated in Figure S2, one milliliter of culture was removed from each culture and cells

were isolated as above. Cells were resuspended in 1x Tris-Glycine running buffer (25 mM Tris-

Cl, pH 8.3, 192 mM glycine, and 0.1% SDS) at 100 microliters per $1.0 = OD600$ cells. Laemmli

buffer from Bio-Rad was added to the resuspended cell pellets to a final concentration of 1x and

heated 10 minutes at $> 98°$ C prior to gel loading.

Upf31 SDS-Page Gel Electrophoresis

Heated, denatured cellular extract (see previous section) samples were loaded on a Tris-

glycine gel with a 12% acrylamide resolving gel and a 4% acrylamide stacking gel. Each sample

lane contained 0.1 OD_{600} of cells. The molecular weight standards used were unstained broad

range protein standards (NEB) and then the gel was run at 100 volts until the dye front was 1 cm

from the bottom of the gel and then stained one hour in Coomassie blue stain (50% methanol,

10% acetic acid, 0.25% Coomassie brilliant blue R-250) followed by destaining (5% methanol,

7% acetic acid) overnight.

Computational Modeling of Interactions between Upf31 and pBP136-encoded Proteins

An 'in silico pulldown' of Upf31 was carried out using the LazyAF pipeline(McLean,

2024). In short, the pBP136 genome (AB237782) was retrieved from NCBI GenBank as a FASTA

protein coding sequence. The Upf31 protein coding sequence was used to generate individual

concatenated FASTA files with Upf31 and each pBP136 coding sequence. ColabFold v1.5.5:

AlphaFold2 w/ MMseqs2 BATCH was then run on Google Colaboratory using the High-RAM

A100 GPU with the following settings: msa_mode: MMseqs2 (UniRef+Environmental),

num_models: 5, num_recycles: 3, stop_at_score: 100. Subsequently the JSON files were analyzed

to retrieve the pTM and ipTM scores for each top ranked model and calculated the

ranking_confidence score (0.2 pTM + 0.8 ipTM).

ACKNOWLEDGEMENTS

We acknowledge Tùng Lê at the John Innes Centre (UK) for his work on the *in silico*

Upf31 pulldown. Dr. Ben Kerr at the University of Washington helped O.K design the low-

density competition assay. We are indebted for the laboratory efforts of undergraduate

researchers Luke Hoover, Morgan Sower, Sue Winger, Katlyn Schafer, Courtney Stattner, Mattie

Hagestad, Audrey Dingel, and Alexandra Gal and research technician Jack Millstein at the

University of Idaho. We also thank Salvador ("Chava") Castaneda Barba for improving this

manuscript. The project was made possible with the skilled sequencing efforts of IIDS Genomics

and Bioinformatics Resources Core (GBRC) at the University of Idaho as well as the Arizona

Genomics Institute at the University of Arizona.

FUNDING RESOURCES

E.M.T.and C.E. received partial support for this work from the National Institute of

Allergy and Infectious Diseases Extramural Activities grant no. R01AI084918 from the National

Institutes of Health. C.A.E. was also supported by the National Science Foundation Graduate

Research Fellowship grant no. DGE-2019265372 as well as the Bioinformatics and

Computational Biology (BCB) Fellowship and the Paul Joyce Memorial BCB Fellowship

Endowment at the University of Idaho. O.K. was supported by the National Science Foundation

Graduate Research Fellowship grant no. DGE-1762114. This work is supported by the Wellcome

Trust Investigator grant 221776/Z/2/Z to Tùng Lê that supported T.C.M, and by the BBSRC

funded Institute Strategic Program Harnessing Biosynthesis for Sustainable Food and Health

(HBio) (BB/X01097X/1). The funders had no role in study design, data collection and analysis,

decision to publish, or preparation of the manuscript.

REFERENCES

Adamczyk, M., & Jagura-Burdzy, G. (2003). Spread and survival of promiscuous IncP-1 plasmids. *Acta*

Biochimica Polonica, *50*(2), 425–453. https://doi.org/10.18388/abp.2003_3696

Akhtar, P. (2002). *Phylogeny and genomic motifs of the beta sub-family of IncP-1 plasmids.* PhD Thesis.

University of Birmingham.

Anton, B. P., Mongodin, E. F., Agrawal, S., Fomenkov, A., Byrd, D. R., Roberts, R. J., & Raleigh, E. A.

(2015). Complete genome sequence of ER2796, a DNA methyltransferase-deficient strain of

Escherichia coli K-12. *PLoS ONE*, *10*(5). https://doi.org/10.1371/journal.pone.0127446

Barrick, J. E., Colburn, G., Deatherage, D. E., Traverse, C. C., Strand, M. D., Borges, J. J., Knoester, D.

B., Reba, A., & Meyer, A. G. (2014). Identifying structural variation in haploid microbial

genomes from short-read resequencing data using breseq. *BMC Genomics*, *15*(1), 1039.

https://doi.org/10.1186/1471-2164-15-1039

Bhattacharyya, A., & Figurski, D. H. (2001). A small protein-protein interaction domain common to KlcB

and global regulators KorA and TrbA of promiscuous IncP plasmids11Edited by M. Belfort.

Journal of Molecular Biology, *310*(1), 51–67. https://doi.org/10.1006/jmbi.2001.4729

Bingle, L. E. H., Macartney, D. P., Fantozzi, A., Manzoor, S. E., Thomas, C. M., & Karn, J. (2005).

Flexibility in repression and cooperativity by KorB of broad host range IncP-1 plasmid RK2.

Journal of Molecular Biology, *349*(2), 302–316. https://doi.org/10.1016/j.jmb.2005.03.062

Bingle, L. E. H., Rajasekar, K. V., Muntaha, S. tul, Nadella, V., Hyde, E. I., & Thomas, C. M. (2008). A

single aromatic residue in transcriptional repressor protein KorA is critical for cooperativity with

its co-regulator KorB. *Molecular Microbiology*, *70*(6), 1502–1514.

https://doi.org/10.1111/j.1365-2958.2008.06498.x

Bingle, L. E. H., & Thomas, C. M. (2001). Regulatory circuits for plasmid survival. *Current Opinion in*

Microbiology, *4*(2), 194–200. https://doi.org/10.1016/S1369-5274(00)00188-0

Bottery, M. J., Wood, A. J., & Brockhurst, M. A. (2017). Adaptive modulation of antibiotic resistance

through intragenomic coevolution. *Nature Ecology & Evolution*, *1*(9), Article 9.

https://doi.org/10.1038/s41559-017-0242-3

Bouma, J. E., & Lenski, R. E. (1988). Evolution of a bacteria/plasmid association. *Nature*, *335*(6188),

Article 6188. https://doi.org/10.1038/335351a0

Dahlberg, C., & Chao, L. (2003). Amelioration of the Cost of Conjugative Plasmid Carriage in

Eschericha coli K12. *Genetics*, *165*(4), 1641–1649.

De Gelder, L., Ponciano, J. M., Joyce, P., & Top, E. M. (2007). Stability of a promiscuous plasmid in

different hosts: No guarantee for a long-term relationship. *Microbiology*, *153*(2), 452–463.

https://doi.org/10.1099/mic.0.2006/001784-0

De Gelder, L., Williams, J. J., Ponciano, J. M., Sota, M., & Top, E. M. (2008). Adaptive Plasmid

Evolution Results in Host-Range Expansion of a Broad-Host-Range Plasmid. *Genetics*, *178*(4),

2179–2190. https://doi.org/10.1534/genetics.107.084475

Dennis, G., Sherman, B. T., Hosack, D. A., Yang, J., Gao, W., Lane, H. C., & Lempicki, R. A. (2003).

DAVID: Database for Annotation, Visualization, and Integrated Discovery. *Genome Biology*,

4(9), R60. https://doi.org/10.1186/gb-2003-4-9-r60

Drønen, A. K., Torsvik, V., Goksøyr, J., & Top, E. M. (1998). Effect of mercury addition on plasmid

incidence and gene mobilizing capacity in bulk soil. *FEMS Microbiology Ecology*, *27*(4), 381–

394. https://doi.org/10.1016/S0168-6496(98)00085-3

Figurski, D. H., Pohlman, R. F., Bechhofer, D. H., Prince, A. S., & Kelton, C. A. (1982). Broad host

range plasmid RK2 encodes multiple kil genes potentially lethal to Escherichia coli host cells.

Proceedings of the National Academy of Sciences, *79*(6), 1935–1939.

https://doi.org/10.1073/pnas.79.6.1935

Gomulkiewicz, R., & Holt, R. D. (1995). When does Evolution by Natural Selection Prevent Extinction?

Evolution, *49*(1), 201–207. https://doi.org/10.2307/2410305

Hall, B. G., Acar, H., Nandipati, A., & Barlow, M. (2014). Growth Rates Made Easy. *Molecular Biology*

and Evolution, *31*(1), 232–238. https://doi.org/10.1093/molbev/mst187

Hall, J. P. J., Wright, R. C. T., Guymer, D., Harrison, E., & Brockhurst, M. A. (2020). Extremely fast

amelioration of plasmid fitness costs by multiple functionally diverse pathways. *Microbiology*,

166(1), 56–62. https://doi.org/10.1099/mic.0.000862

Hall, J. P. J., Wright, R. C. T., Harrison, E., Muddiman, K. J., Wood, A. J., Paterson, S., & Brockhurst,

M. A. (2021). Plasmid fitness costs are caused by specific genetic conflicts enabling resolution by

compensatory mutation. *PLOS Biology*, *19*(10), e3001225.

https://doi.org/10.1371/journal.pbio.3001225

Harrison, E., Guymer, D., Spiers, A. J., Paterson, S., & Brockhurst, M. A. (2015). Parallel Compensatory

Evolution Stabilizes Plasmids across the Parasitism-Mutualism Continuum. *Current Biology*,

25(15), 2034–2039. https://doi.org/10.1016/j.cub.2015.06.024

Hassan, H., & Troxell, B. (2013). Transcriptional regulation by Ferric Uptake Regulator (Fur) in

pathogenic bacteria. *Frontiers in Cellular and Infection Microbiology*, *3*.

https://www.frontiersin.org/articles/10.3389/fcimb.2013.00059

Heckman, K. L., & Pease, L. R. (2007). Gene splicing and mutagenesis by PCR-driven overlap extension.

Nature Protocols, *2*(4), 924–932. https://doi.org/10.1038/nprot.2007.132

Hughes, J. M., Lohman, B. K., Deckert, G. E., Nichols, E. P., Settles, M., Abdo, Z., & Top, E. M. (2012).

The role of clonal interference in the evolutionary dynamics of plasmid-host adaptation. *mBio*,

3(4), e00077-12. https://doi.org/10.1128/mBio.00077-12

Jain, A., & Srivastava, P. (2013). Broad host range plasmids. *FEMS Microbiology Letters*, *348*(2), 87–96.

https://doi.org/10.1111/1574-6968.12241

Jordt, H., Stalder, T., Kosterlitz, O., Ponciano, J. M., Top, E. M., & Kerr, B. (2020). Coevolution of host–

plasmid pairs facilitates the emergence of novel multidrug resistance. *Nature Ecology &*

Evolution, *4*(6), 863–869. https://doi.org/10.1038/s41559-020-1170-1

Jumper, J., Evans, R., Pritzel, A., Green, T., Figurnov, M., Ronneberger, O., Tunyasuvunakool, K., Bates,

R., Žídek, A., Potapenko, A., Bridgland, A., Meyer, C., Kohl, S. A. A., Ballard, A. J., Cowie, A.,

Romera-Paredes, B., Nikolov, S., Jain, R., Adler, J., … Hassabis, D. (2021). Highly accurate

protein structure prediction with AlphaFold. *Nature*, *596*(7873), Article 7873.

https://doi.org/10.1038/s41586-021-03819-2

Kamachi, K., Sota, M., Tamai, Y., Nagata, N., Konda, T., Inoue, T., Top, E. M., & Arakawa, Y. (2006).

Plasmid pBP136 from Bordetella pertussis represents an ancestral form of IncP-1β plasmids

without accessory mobile elements. *Microbiology*, *152*(12), 3477–3484.

https://doi.org/10.1099/mic.0.29056-0

Klümper, U., Riber, L., Dechesne, A., Sannazzarro, A., Hansen, L. H., Sørensen, S. J., & Smets, B. F.

(2015). Broad host range plasmids can invade an unexpectedly diverse fraction of a soil bacterial

community. *The ISME Journal*, *9*(4), Article 4. https://doi.org/10.1038/ismej.2014.191

Kornacki, J. A., Chang, C. H., & Figurski, D. H. (1993). kil-kor Regulon of promiscuous plasmid RK2:

Structure, products, and regulation of two operons that constitute the kilE locus. *Journal of*

Bacteriology, *175*(16), 5078–5090. https://doi.org/10.1128/jb.175.16.5078-5090.1993

Kostelidou, K., Jones, A. C., & Thomas, C. M. (1999). Conserved C-terminal region of global repressor

KorA of broad-host-range plasmid RK2 is required for co-operativity between KorA and a

second RK2 global regulator, KorB11Edited by J. Karn. *Journal of Molecular Biology*, *289*(2),

211–221. https://doi.org/10.1006/jmbi.1999.2761

Kostelidou, K., & Thomas, C. M. (2000). The hierarchy of KorB binding at its 12 binding sites on the

broad-host-range plasmid RK2 and modulation of this binding by IncC1 protein11Edited by J.

Karn. *Journal of Molecular Biology*, *295*(3), 411–422. https://doi.org/10.1006/jmbi.1999.3359

Kosterlitz, O., Muñiz Tirado, A., Wate, C., Elg, C., Bozic, I., Top, E. M., & Kerr, B. (2022). Estimating

the transfer rates of bacterial plasmids with an adapted Luria–Delbrück fluctuation analysis. *PLoS*

Biology, *20*(7), e3001732. https://doi.org/10.1371/journal.pbio.3001732

Kottara, A., Hall, J. P. J., Harrison, E., & Brockhurst, M. A. (2018). Variable plasmid fitness effects and

mobile genetic element dynamics across Pseudomonas species. *FEMS Microbiology Ecology*,

94(1). https://doi.org/10.1093/femsec/fix172

Langmead, B., & Salzberg, S. L. (2012). Fast gapped-read alignment with Bowtie 2. *Nature Methods*,

9(4), 357–359. https://doi.org/10.1038/nmeth.1923

Larsen, M. H., & Figurski, D. H. (1994). Structure, expression, and regulation of the kilC operon of

promiscuous IncP alpha plasmids. *Journal of Bacteriology*, *176*(16), 5022–5032.

Law, A., Solano, O., Brown, C. J., Hunter, S. S., Fagnan, M., Top, E. M., & Stalder, T. (2021). Biosolids

as a Source of Antibiotic Resistance Plasmids for Commensal and Pathogenic Bacteria. *Frontiers*

in Microbiology, *12*, 606409. https://doi.org/10.3389/fmicb.2021.606409

Liao, Y., Smyth, G. K., & Shi, W. (2014). featureCounts: An efficient general purpose program for

assigning sequence reads to genomic features. *Bioinformatics*, *30*(7), 923–930.

https://doi.org/10.1093/bioinformatics/btt656

Loftie-Eaton, W., Bashford, K., Quinn, H., Dong, K., Millstein, J., Hunter, S., Thomason, M. K.,

Merrikh, H., Ponciano, J. M., & Top, E. M. (2017). Compensatory mutations improve general

permissiveness to antibiotic resistance plasmids. *Nature Ecology and Evolution*, *1*(9), 1354–1363.

https://doi.org/10.1038/s41559-017-0243-2

Love, M. I., Huber, W., & Anders, S. (2014). Moderated estimation of fold change and dispersion for

RNA-seq data with DESeq2. *Genome Biology*, *15*(12), 550. https://doi.org/10.1186/s13059-014-

0550-8

Marincola, G., Jaschkowitz, G., Kieninger, A.-K., Wencker, F. D. R., Feßler, A. T., Schwarz, S., &

Ziebuhr, W. (2021). Plasmid-Chromosome Crosstalk in Staphylococcus aureus: A Horizontally

Acquired Transcription Regulator Controls Polysaccharide Intercellular Adhesin-Mediated

Biofilm Formation. *Frontiers in Cellular and Infection Microbiology*, *11*.

https://doi.org/10.3389/fcimb.2021.660702

McLean, T. C. (2024). *LazyAF, a pipeline for accessible medium-scale in silico prediction of protein-*

protein interactions (p. 2024.01.29.577767). bioRxiv. https://doi.org/10.1101/2024.01.29.577767

McLean, T. C., Balaguer-Pérez, F., Chandanani, J., Thomas, C. M., Aicart-Ramos, C., Burick, S.,

Olinares, P. D. B., Gobbato, G., Mundy, J. E. A., Chait, B. T., Lawson, D. M., Darst, S. A.,

Campbell, E. A., Moreno-Herrero, F., & Le, T. B. K. (2024). *Molecular switching of a DNA-*

sliding clamp to a repressor mediates long-range gene silencing (p. 2024.02.16.579611).

bioRxiv. https://doi.org/10.1101/2024.02.16.579611

Mirdita, M., Schütze, K., Moriwaki, Y., Heo, L., Ovchinnikov, S., & Steinegger, M. (2022). ColabFold:

Making protein folding accessible to all. *Nature Methods*, *19*(6), Article 6.

https://doi.org/10.1038/s41592-022-01488-1

Modi, R. I., & Adams, J. (1991). Coevolution in bacterial-plasmid populations. *Evolution*, *45*(3), 656–

667.

Murray, C. J. L., Ikuta, K. S., Sharara, F., Swetschinski, L., Aguilar, G. R., Gray, A., Han, C., Bisignano,

C., Rao, P., Wool, E., Johnson, S. C., Browne, A. J., Chipeta, M. G., Fell, F., Hackett, S., Haines-

Woodhouse, G., Hamadani, B. H. K., Kumaran, E. A. P., McManigal, B., … Naghavi, M. (2022).

Global burden of bacterial antimicrobial resistance in 2019: A systematic analysis. *The Lancet*,

399(10325), 629–655. https://doi.org/10.1016/S0140-6736(21)02724-0

Norman, A., Hansen, L. H., & Sørensen, S. J. (2009). Conjugative plasmids: Vessels of the communal

gene pool. *Philosophical Transactions of the Royal Society of London. Series B, Biological*

Sciences, *364*(1527), 2275–2289. https://doi.org/10.1098/rstb.2009.0037

Pansegrau, W., Lanka, E., Barth, P. T., Figurski, D. H., Guiney, D. G., Haas, D., Helinski, D. R., Schwab,

H., Stanisich, V. A., & Thomas, C. M. (1994). Complete nucleotide sequence of Birmingham

IncP alpha plasmids. Compilation and comparative analysis. *Journal of Molecular Biology*,

239(5), 623–663. https://doi.org/10.1006/jmbi.1994.1404

Popowska, M., & Krawczyk-Balska, A. (2013). Broad-host-range IncP-1 plasmids and their resistance

potential. *Frontiers in Microbiology*, *4*(MAR), 1–8. https://doi.org/10.3389/fmicb.2013.00044

Porse, A., Schønning, K., Munck, C., & Sommer, M. O. A. (2016). Survival and Evolution of a Large

Multidrug Resistance Plasmid in New Clinical Bacterial Hosts. *Molecular Biology and Evolution*,

33(11), 2860–2873. https://doi.org/10.1093/molbev/msw163

Rajasekar, K. V., Lovering, A. L., Dancea, F., Scott, D. J., Harris, S. A., Bingle, L. E. H., Roessle, M.,

Thomas, C. M., Hyde, E. I., & White, S. A. (2016). Flexibility of KorA, a plasmid-encoded,

global transcription regulator, in the presence and the absence of its operator. *Nucleic Acids*

Research, *44*(10), 4947–4956. https://doi.org/10.1093/nar/gkw191

Sambrook, Joseph., & Russell, D. W. (David W., 1957-. (2001). *Molecular cloning: A Laboratory*

Manual (3rd ed.). Cold Spring Harbor Laboratory Press; WorldCat.org.

San Millan, A. (2018). Evolution of Plasmid-Mediated Antibiotic Resistance in the Clinical Context.

Trends in Microbiology, *26*(12), 978–985. https://doi.org/10.1016/j.tim.2018.06.007

San Millan, A., Peña-Miller, R., Toll-Riera, M., Halbert, Z. V., McLean, A. R., Cooper, B. S., &

Maclean, R. C. (2014). Positive selection and compensatory adaptation interact to stabilize non-

transmissible plasmids. *Nature Communications*, *5*, 5208. https://doi.org/10.1038/ncomms6208

San Millan, A., Toll-Riera, M., Qi, Q., & MacLean, R. C. (2015). Interactions between horizontally

acquired genes create a fitness cost in Pseudomonas aeruginosa. *Nature Communications*, *6*,

6845. https://doi.org/10.1038/ncomms7845

Schmidhauser, T. J., & Helinski, D. R. (1985). Regions of broad-host-range plasmid RK2 involved in

replication and stable maintenance in nine species of gram-negative bacteria. *Journal of*

Bacteriology, *164*(1), 446–455.

Sen, D., Brown, C. J., Top, E. M., & Sullivan, J. (2013). Inferring the evolutionary history of IncP-1

plasmids despite incongruence among backbone gene trees. *Molecular Biology and Evolution*,

30(1), 154–166. https://doi.org/10.1093/molbev/mss210

Shintani, M., Takahashi, Y., Yamane, H., & Nojiri, H. (2010). The behavior and significance of

degradative plasmids belonging to Inc groups in Pseudomonas within natural environments and

microcosms. *Microbes and Environments*, *25*(4), 253–265.

https://doi.org/10.1264/jsme2.me10155

Sota, M., & Top, E. M. (2008). Host-specific factors determine the persistence of IncP-1 plasmids. *World*

Journal of Microbiology and Biotechnology, *24*(9), 1951–1954. https://doi.org/10.1007/s11274-

008-9653-2

Sota, M., Tsuda, M., Yano, H., Suzuki, H., Forney, L. J., & Top, E. M. (2007). Region-specific insertion

of transposons in combination with selection for high plasmid transferability and stability

accounts for the structural similarity of IncP-1 plasmids. *Journal of Bacteriology*, *189*(8), 3091–

3098. https://doi.org/10.1128/JB.01906-06

Sota, M., Yano, H., M Hughes, J., Daughdrill, G. W., Abdo, Z., Forney, L. J., & Top, E. M. (2010). Shifts

in the host range of a promiscuous plasmid through parallel evolution of its replication initiation

protein. *ISME Journal*, *4*(12), 1568–1580. https://doi.org/10.1038/ismej.2010.72

Soucy, S. M., Huang, J., & Gogarten, J. P. (2015). Horizontal gene transfer: Building the web of life.

Nature Reviews Genetics, *16*(8), 472–482. https://doi.org/10.1038/nrg3962

Stalder, T., Rogers, L. M., Renfrow, C., Yano, H., Smith, Z., & Top, E. M. (2017). Emerging patterns of

plasmid-host coevolution that stabilize antibiotic resistance. *Scientific Reports*, *7*(1), 1–10.

https://doi.org/10.1038/s41598-017-04662-0

Thomas, C. M. (2000). Paradigms of plasmid organization. *Molecular Microbiology*, *37*(3), 485–491.

https://doi.org/10.1046/j.1365-2958.2000.02006.x

Thomas, C. M., Thomson, N. R., Cerdeño-Tárraga, A. M., Brown, C. J., Top, E. M., & Frost, L. S. (2017). Annotation of plasmid genes. *Plasmid*, *91*(December 2016), 61–67. https://doi.org/10.1016/j.plasmid.2017.03.006

Thompson, C. M. A., Hall, J. P. J., Chandra, G., Martins, C., Saalbach, G., Panturat, S., Bird, S. M., Ford, S., Little, R. H., Piazza, A., Harrison, E., Jackson, R. W., Brockhurst, M. A., & Malone, J. G. (2023). Plasmids manipulate bacterial behaviour through translational regulatory crosstalk. *PLOS Biology*, *21*(2), e3001988. https://doi.org/10.1371/journal.pbio.3001988

Thorsted, P. B., MacArtney, D. P., Akhtar, P., Haines, A. S., Ali, N., Davidson, P., Stafford, T., Pocklington, M. J., Pansegrau, W., Wilkins, B. M., Lanka, E., & Thomas, C. M. (1998). Complete sequence of the IncPβ plasmid R751: Implications for evolution and organisation of the IncP backbone. *Journal of Molecular Biology*, *282*(5), 969–990. https://doi.org/10.1006/jmbi.1998.2060

Vial, L., & Hommais, F. (2020). Plasmid-chromosome cross-talks. *Environmental Microbiology*, *22*(2), 540–556. https://doi.org/10.1111/1462-2920.14880

Wiser, M. J., & Lenski, R. E. (2015). A Comparison of Methods to Measure Fitness in Escherichia coli. *PLOS ONE*, *10*(5), e0126210. https://doi.org/10.1371/journal.pone.0126210

Yang, J., Wu, R., Xia, Q., Yu, J., Yi, L.-X., Huang, Y., Deng, M., He, W.-Y., Bai, Y., Lv, L., Burrus, V., Wang, C., & Liu, J.-H. (2023). The evolution of infectious transmission promotes the persistence of mcr-1 plasmids. *mBio*, *14*(4), e00442-23. https://doi.org/10.1128/mbio.00442-23

Yano, H., Deckert, G. E., Rogers, L. M., & Top, E. M. (2012). Roles of long and short replication initiation proteins in the fate of IncP-1 Plasmids. *Journal of Bacteriology*, *194*(6), 1533–1543. https://doi.org/10.1128/JB.06395-11

Yano, H., Rogers, L. M., Knox, M. G., Heuer, H., Smalla, K., Brown, C. J., & Top, E. M. (2013). Host range diversification within the IncP-1 plasmid group. *Microbiology (United Kingdom)*, *159*(PART11), 2303–2315. https://doi.org/10.1099/mic.0.068387-0

841   Yano, H., Wegrzyn, K., Loftie-Eaton, W., Johnson, J., Deckert, G. E., Rogers, L. M., Konieczny, I., &

842   Top, E. M. (2016). Evolved plasmid-host interactions reduce plasmid interference cost.

843   *Molecular Microbiology*, *101*(5), 743–756. https://doi.org/10.1111/mmi.13407

844   Zatyka, M., Bingle, L., Jones, A. C., & Thomas, C. M. (2001). Cooperativity between KorB and TrbA

845   repressors of broad-host-range plasmid RK2. *Journal of Bacteriology*, *183*(3), 1022–1031.

846   https://doi.org/10.1128/JB.183.3.1022-1031.2001

847   Zatyka, M., Jagura-Burdzy, G., & Thomas, C. M. (1997). Transcriptional and translational control of the

848   genes for the mating pair formation apparatus of promiscuous IncP plasmids. *Journal of*

849   *Bacteriology*, *179*(23), 7201–7209. https://doi.org/10.1128/jb.179.23.7201-7209.1997

850   Zhong, X., Kro ̇l, JarosŁaw E., Top, E. M., & Krone, S. M. (2010). Accounting for mating pair formation

851   in plasmid population dynamics. *Journal of Theoretical Biology*, *262*(4), 711–719.

852   https://doi.org/10.1016/j.jtbi.2009.10.013

853   Zukowski, M. M., Gaffney, D. F., Speck, D., Kauffmann, M., Findeli, A., Wisecup, A., & Lecocq, J. P.

854   (1983). Chromogenic identification of genetic regulatory signals in Bacillus subtilis based on

855   expression of a cloned Pseudomonas gene. *Proceedings of the National Academy of Sciences*,

856   *80*(4), 1101–1105. https://doi.org/10.1073/pnas.80.4.1101

857