Rtt107 cooperates with Rad55 or Slx4 to maintain genome stability in *Saccharomyces cerevisiae*

Joshua A. R. Brown[1] and Michael S. Kobor[2*]

[1]Centre for Molecular Medicine and Therapeutics, BC Children's Hospital Research Institute,

Department of Medical Genetics, University of British Columbia, Vancouver, British Columbia,

Canada, V5Z 4H4

[2]Centre for Molecular Medicine and Therapeutics, Edwin S.H. Leong Centre for Healthy Aging,

Department of Medical Genetics, University of British Columbia, Vancouver, British Columbia,

Canada, V5Z 4H4

*Corresponding author: Michael S. Kobor

msk@cmmt.ubc.ca

Short title: Rtt107, Rad55, and Genome Maintenance

Abstract

A range of genome maintenance factors respond to endogenous and exogenous DNA damage to prevent mutations and cell death. The scaffold protein, Rtt107, is important for the growth of cells exposed to DNA-damaging agents in the budding yeast *Saccharomyces cerevisiae*. Rtt107 binds to a diverse array of partner proteins, such as Slx4, and responds to DNA damage by localizing to phosphorylated histone H2A. Rad55–Rad57, a heterodimer involved in DNA repair, also binds to Rtt107, but the function of the Rtt107–Rad55–Rad57 complex remains unclear. In addition to their sensitivity to DNA-damaging agents, *rtt107Δ* mutants exhibit spontaneous genome instability phenotypes, including spontaneous loss of heterozygosity (LOH) caused by crossovers and other genetic events. However, the binding partners with which Rtt107 interacts to prevent spontaneous genome instability have yet to be elucidated. Here, we showed that Rtt107 acts in the same pathway as Rad55 to limit LOH, specifically by preventing crossover events. A *rad55-S404A* phosphorylation site mutation largely disrupted the interaction between Rtt107 and Rad55–Rad57, resulting in increased LOH and crossover rates, consistent with the contribution of Rtt107–Rad55–Rad57 interaction to genome stability. Strikingly, an *rtt107-K887M* mutation that reduces Rtt107 recruitment to H2A did not result in an LOH phenotype, suggesting that the role of Rtt107 in preventing LOH is distinct from its function as an H2A-binding scaffold. Rtt107 did not function primarily in the same pathway as Rad55 to limit recombination at the sensitive ribosomal DNA (rDNA) locus, but instead acted with Slx4 to maintain rDNA stability, suggesting that interactions of Rtt107 with different partners prevented distinct types of instability. Taken together, our observations suggested that Rtt107 limits spontaneous LOH and crossover events in part by binding to Rad55 in a manner dependent on Rad55-S404.

Author Summary

Numerous proteins are involved in the repair of damaged DNA and prevention of genome instability in cells, which would otherwise result in persistent changes to DNA. Genome maintenance pathways are evolutionarily conserved, and the budding yeast, *Saccharomyces cerevisiae*, is a powerful model organism for investigating the maintenance of genome integrity. Rtt107 is a scaffold protein expressed in yeast, containing conserved protein domains that are important for the function of genome maintenance proteins. Although the functions of Rtt107 in cells treated with DNA-damaging agents have been characterized in some detail, it remains unclear how Rtt107 prevents spontaneous genome instability in cells growing under normal conditions. Here, we found that Rtt107 prevents specific types of spontaneous genome instability and acts in the same pathway as the DNA repair protein, Rad55, to which it binds. Mutation of a possible Rtt107 binding site on Rad55 showed that Rtt107 indeed limited genome instability in part by binding to Rad55. Strikingly, Rtt107 also showed Rad55-independent roles in preventing ribosomal DNA (rDNA) instability, and in this context Rtt107 cooperated in part with its binding partner Slx4. Taken together, our results revealed the pathways by which the evolutionarily conserved protein Rtt107 limits spontaneous genome instability.

1   Introduction

The genome is constantly under threat by a variety of endogenous and exogenous factors, which are counteracted by the DNA damage response (DDR) consisting of interconnected pathways that detect, signal, and repair DNA damage [1,2]. These DDR proteins play crucial roles in ultimately preventing cell death when DNA damage occurs. Genome instability, encompassing both mutations and genomic rearrangements, can also occur as a byproduct of DNA damage and repair. Cells therefore rely on a variety of genome maintenance factors, with functions in the DDR and in processes such as DNA replication, to both survive DNA damage and limit genome instability.

DNA double-strand breaks (DSBs) are among the most deleterious types of DNA damage, and can cause cell death if left unrepaired [3]. DSBs and DSB repair can result in a range of types of genome instability, including deletions, gross chromosomal rearrangements (GCRs), and loss of heterozygosity (LOH) [2,3]. DSBs are repaired by a number of different pathways depending on the stage of the cell cycle at which they occur and the nature of specific lesions. In the budding yeast, *Saccharomyces cerevisiae*, DSB repair most frequently involves the homologous recombination (HR) pathway, while non-homologous end-joining (NHEJ) of DSB ends is specifically upregulated in G_1 phase haploid yeast and is the predominant pathway in mammals [4]. HR occurs after resection, the $5'{\rightarrow}3'$ nucleolytic digestion of DSB ends that exposes single-stranded DNA (ssDNA) and leads to formation of a Rad51–ssDNA filament, and can generate reciprocal crossovers (RCOs) [3]. Alongside specific repair processes, DNA damage prompts a broader DDR that involves checkpoint signaling, which arrests the cell cycle while DNA damage is repaired [5]. Here, histone H2A-S129 phosphorylation (γH2A) is placed by the Mec1 and Tel1 sensor kinases around lesions, and forms a platform for the recruitment of

other DDR factors, including checkpoint proteins [6]. Another histone mark, H4T80

phosphorylation, is placed near sites of DNA damage by Cla4, although the functional

consequences of this chromatin mark in the DDR have yet to be fully elucidated [7].

Rtt107 is an evolutionarily conserved protein that plays multiple roles in the DDR, and is

required for normal cell growth during chronic exposure to DNA-damaging agents [8]. At the

molecular level, Rtt107 localizes to stalled replication forks, DSBs, and other DNA damage

sites, where it acts as a scaffold to recruit other proteins to these lesions [9–14]. Rtt107 promotes

recovery from treatments that damage DNA and the restart of DNA replication after damage

repair, at least in part by limiting DNA damage and preventing checkpoint hyperactivation

[8,15–17]. Rtt107 contains six BRCA1 C-terminal (BRCT) domains, which generally mediate

both phosphorylation-dependent and -independent protein–protein interactions [8,18], and

localizes near DSBs by binding γH2A via BRCT5/6 [10,19]. Rtt107 is also recruited near DSBs

by binding phosphorylated H4T80 via BRCT1–4, with this representing the first identified

molecular interaction partner of phosphorylated H4T80 [7]. The interaction of Rtt107 with γH2A

is mediated by Rtt107-K887 and is required for its localization near DSBs [10,19], whereas

interaction with phosphorylated H4T80 is mediated by Rtt107-K426 and partially contributes to

its localization near DSBs [7]. Both *rtt107-K426A* and *rtt107-K887M* mutants exhibit

hypersensitivity to chronic treatment with DNA-damaging agents, suggesting that Rtt107 confers

DNA damage resistance, at least in part, by localizing to histone marks [10]. Rtt107–BRCT1–4

forms a tetrahedral structure that binds a variety of complexes, and in turn Rtt107 may act as a

scaffold by recruiting these partners to γH2A-containing regions, to which Rtt107 itself is

tethered via BRCT5/6 [9,14,15,20].

Rtt107 has been studied most extensively in the context of its localization to γH2A and

its constitutive interaction with the Slx4 endonuclease [10,14,15]. DNA damage induces

interaction of the Rtt107–Slx4 complex with the replication factor, Dpb11 [21]. In turn, this

Rtt107– Slx4–Dpb11 interaction can competitively reduce the interaction between Dpb11 and

the checkpoint factor, Rad9, while increased Rad9–Dpb11 interaction in the absence of the

Rtt107–Slx4 complex can limit DSB resection and may contribute to checkpoint hyperactivation

[16,17,22–24]. Slx4 is also required for phosphorylation of the SQ/TQ motifs of Rtt107 by Mec1

after treatment with the DNA-damaging agent, methyl methanesulfonate (MMS) [8,15].

However, it is unclear which functions of Rtt107 are dependent on its phosphorylation [8,21,25].

Rtt107 interacts with the Rad55–Rad57 heterodimer via Rtt107–BRCT1–4, and Rad55

contains an Rtt107 interaction motif that was shown to interact with Rtt107–BRCT1–4 *in vitro*

[14,20]. However, the molecular function of the Rtt107–Rad55–Rad57 complex has yet to be

determined. In general, Rad55–Rad57 acts as a recombination mediator, which promotes Rad51–

ssDNA filament assembly for HR [26–30]. In contrast to the *rtt107Δ* mutant [8], both *rad55Δ*

and *rad57Δ* mutants are sensitive to treatment with ionizing radiation [27,31], which induces

DSBs. These findings suggest that Rtt107 may be less important for successful DSB repair than

Rad55–Rad57, or alternatively that the importance of Rtt107 is limited to specific contexts. In

addition to promoting DSB repair as a recombination mediator, Rad55–Rad57 also plays a role

in DNA damage tolerance pathways that allow DNA lesions to be bypassed during replication

and gap filling after replication [32]. Here, Rad55–Rad57 promotes the error-free postreplication

repair (PRR) tolerance pathway, also known as template switching, by bridging Rad51 to the Shu

complex [33–36]. Recent work also suggested that Rad55–Rad57 may downregulate the

translesion synthesis (TLS) pathway [37]. Although *rtt107Δ* mutants exhibit spontaneous Rad52

foci [38], a marker of DNA damage and recombination, it has not been determined whether

Rtt107 is important for any DNA damage repair function of Rad55–Rad57.

The function of Rtt107 in the context of exposure to DNA-damaging agents has been

documented in detail, while the role of Rtt107 in preventing spontaneous genome instability is

only beginning to be elucidated. Originally, the *rtt107Δ* mutant was identified in screens for

mutants with elevated levels of spontaneous LOH and GCRs, chromosomal instability, and

phenotypes reflecting spontaneous DNA damage foci and DDR signaling [38–42]. The *rtt107Δ*

mutant also exhibits hyperrecombination at the ribosomal DNA (rDNA) locus [38], a highly

transcribed series of tandem repeats spanning ~10% of the *S. cerevisiae* genome that represents a

unique challenge to the maintenance of genome stability [43]. Taken together, these phenotypes

indicate that Rtt107 plays an important role in preventing several types of spontaneous genome

instability. However, it remains to be explored whether the role of Rtt107 in preventing

spontaneous instability is related to its established functions in the response to exogenous DNA-

damaging agents. Rtt107 binds to a variety of DDR complexes in addition to Slx4 and Rad55–

Rad57, and it is possible that Rtt107 cooperates with one or several of its documented binding

partners to prevent spontaneous genome instability. These partners include the SMC5/6

(Structural Maintenance of Chromosomes) complex, Rtt101–Mms1–Mms22, and Mus81–

Mms4–Cdc7–Dbf4–Cdc5 [9,12,21,44–46]. Further, Rtt107 was shown to interact with the Sgs1

helicase in a yeast two-hybrid screen [20].

Genetic interactions underlying genome instability phenotypes represent powerful tools

for investigating gene function, and epistatic interactions can often be interpreted in the

framework of classical biochemical pathway models [2]. Here, we made use of a comprehensive

suite of genetic analyses to uncover the specific pathways in which Rtt107 prevents specific

types of spontaneous genome instability. First, we established that the function of Rtt107 in

preventing spontaneous LOH was distinct from its role as a γH2A-binding scaffold and from the

role of Rtt107 in the context of chronic exposure to DNA-damaging agents, but that Rtt107–

BRCT5/6 was still important for preventing spontaneous LOH. Next, we found that Rtt107

cooperated specifically with Rad55 to limit LOH as well as mitotic RCO events. A *rad55-S404A*

mutant showed reduced interaction between Rtt107 and Rad55 and increased rates of mitotic

RCOs and LOH, although the *rtt107Δ* and *rad55-S404A* mutations also made independent

contributions to the LOH rate. However, rather than functioning with Rad55 to limit spontaneous

recombination at the rDNA locus, Rtt107 cooperated with Slx4 to maintain rDNA stability.

Taken together, these results demonstrated that the role of Rtt107 in preventing LOH was largely

independent of its previously identified function as a γH2A-binding scaffold, and indicated that

Rtt107 cooperated with Rad55 or Slx4 to prevent different types of instability at distinct genetic

loci.

2   Materials and Methods

2.1   Yeast strains and plasmids

Yeast strains used in this study are listed in Table S1, made using standard genetic

techniques. The primers used to generate the Rad57-HA construct also encoded a

GCAGCAGCAGCAGCAGCAGCA linker between Rad57 and the HA tag (primers available

upon request).

The plasmids used in this study are listed in Table S2, and were produced using standard

genetic techniques. Plasmids pMK429 and pMK430 were generous gifts from Dr. Jocelyn Krebs

(University of Alaska Anchorage), and were used in the experiments presented in Figure 2C to

examine the *H2A-S129A* allele.

2.2 Genome instability assays

Several assays for genome instability phenotypes were applied in this study under normal growth conditions. Each assay is briefly summarized below and outlined schematically in Fig. 1.

2.2.1 Quantitative bimater assay

The procedure for the quantitative bimater assay was adapted from a previous report [25]. An overnight culture of a *MAT*a mating tester strain was grown in YPD at 30°C, diluted, and plated onto solid medium containing no amino acids (−AA plates). Overnight cultures of the test strain were grown at 30°C and diluted. Cultures were then plated at different dilutions onto −AA plates to count mating events with the *MAT*a tester and onto YPD plates to score total counts. Plates were grown at 30°C for 2–3 days before counting. Mating rates and 95% confidence intervals were calculated with the maximum likelihood estimate method, implemented using the `newton.LD.plating` function in the `rSalvador` R package [47]. Statistical significance was assessed by comparing the overlap of confidence intervals [48], and comparisons between strains were considered significant if the 95% confidence intervals did not overlap. This approach is more stringent than a conventional p-value cutoff of 0.05 [48].

The bimater assay shown in Figure 2B was adapted to maintain plasmid selection. Strains were grown in SC-Leu medium, and mated to a *leu2Δ* mating tester strain.

2.2.2 Reciprocal crossover (RCO) assay

RCO events were examined using *can1-100* and *SUP4-o* alleles on chromosome V, as described previously [49]. Independent colonies were grown on YPD plates for 2 days and resuspended in water. Cells were plated at different dilutions onto solid SC-Arg medium with or

without 250 μg/mL of canavanine sulfate (Sigma). Plates were grown for 4 days at 30°C and

stored at 4°C for 1 day to allow colony color development. Red/white sectored colonies on plates

containing canavanine were scored as representative of RCO events in the interval between

CEN5 and *CAN1* loci. Counts were compared to colony counts on plates lacking canavanine, and

the frequency of RCOs was calculated. Statistical significance was assessed using Student's *t*

test.

2.2.3   Quantitative A-like faker (ALF) assay

A quantitative version of the A-like faker (ALF) assay was performed using the same

method as the bimater assay, except that *MATα* haploid strains were examined for their ability to

mate with a *MATα* tester strain. Mating rates and statistical significance were calculated using

the same methods as described above.

2.2.4   rDNA recombination assay

An rDNA recombination assay was adapted from a previous report to allow for

quantification by fluctuation analysis [50]. Strains were plated onto solid medium lacking

adenine and grown for 2 days at 30°C. Independent cultures were grown in liquid medium at

30°C for 2 days. Diluted cultures were plated onto solid medium lacking arginine with or

without 120 μg/mL canavanine (Sigma). Plates with canavanine were incubated for 2 days,

replica plated onto media with or without adenine, and incubated for 2 days. Canavanine-

resistant colonies that did not grow on plates lacking adenine were scored as representative of

rDNA recombination. Scored colonies were compared to colony counts from plates lacking

canavanine. The rDNA recombination rates and statistical significance were calculated by the

method described for mating rates in the bimater assay.

2.2.5   Diploid instability assay

The assay used to examine various types of genome instability in diploids shown in

Figure 6C was adapted from a previous report [49]. Colonies were grown, diluted, and plated as

described for the RCO assay. Colonies were plated at separate dilutions to examine Class 5/6

events due to their lower rates. After 4 days of growth at 30°C, colonies were replica plated onto

several types of solid media to score growth phenotypes, as described previously [49]. Plates

were grown at 30°C for 2 days and stored at 4°C for 2 days to allow colony color development.

Non-sectored colonies were scored as representative of Class 1–6 events according to their

respective growth phenotypes. Mutation rates and statistical significance were calculated as

described for mating rates in the bimater assay. A total of 28 Class 3 mutants were examined by

PCR as described previously [49] to discriminate between local gene conversion and *de novo*

mutation events at the *SUP4-o* locus.

2.3   DNA damage sensitivity assays

Overnight cultures grown in YPD at 30°C were diluted to 0.5 OD_{600}. Cells were then 10-

fold serially diluted, and plated on YPD medium containing the indicated concentration of MMS,

HU, or CPT (Sigma). Plates were incubated for 2 days at 30°C.

2.4   TCA extraction

Overnight cultures were diluted to 0.3 OD_{600} in YPD medium and grown to mid-log

phase. Whole-cell extracts were prepared by glass bead lysis in the presence of trichloroacetic

acid (TCA). Samples were boiled for 5 min and resolved by 8% SDS-PAGE. Western blotting

analysis was performed using anti-FLAG M2 (Sigma) and anti-PGK1 (Thermo Fisher)

antibodies.

2.5 Analytical-scale protein–protein interaction assay

Overnight cultures were diluted to 0.3 OD_{600} in 4 L of YPD medium, and grown to mid-log phase. The procedure for co-immunoprecipitation was adapted from a previous report [51]. Briefly, cells were disrupted with a coffee grinder in the presence of dry ice pellets and resuspended in TAP-IP buffer (50 mM Tris, pH 7.8, 150 mM NaCl, 1.5 mM MgAc, 0.15% Nonidet P-40, 1 mM DTT, 10 mM NaPPi, 5 mM EGTA, 5 mM EDTA, 0.1 mM Na_3VO_4, 5 mM NaF, and Roche cOmplete™ Protease Inhibitor Mix). Crude extracts were prepared by centrifugation, and incubated for 2 hours at 4°C with anti-FLAG M2 antibody (Sigma)-coupled Dynabeads (Invitrogen). Beads were washed with TAP-IP buffer. Captured protein was detected by immunoblotting with anti-FLAG M2 antibody (Sigma) and visualized using an Odyssey Infrared Imaging System (Licor). Co-purifying proteins were imaged with anti-HA (Sigma) and anti-VSV (Bethyl Laboratories) antibodies. Anti-PGK1 (Thermo Fisher) and anti-VSV (Bethyl Laboratories) antibodies were applied to the input fraction as loading controls.

3 Results

3.1 Rtt107–BRCT1–4 and BRCT5/6 were important for limiting spontaneous LOH

Despite the identification of a number of Rtt107 binding partners and of the molecular functions of Rtt107 in cells exposed to DNA-damaging agents, it remains an important question how Rtt107 operates to prevent a variety of types of spontaneous genome instability. As Rtt107 acts as a scaffold by binding to different partners via Rtt107–BRCT1–4 and BRCT5/6, we first examined which structural features of Rtt107 were required for limiting spontaneous genome instability. For this purpose, we examined a spontaneous LOH phenotype that was previously identified for the *rtt107Δ* mutant using the bimater assay (Fig. 1) [25,39]; this assay determines the rate of LOH at the mating type locus in diploid yeast and therefore captures a range of

underlying mechanisms, encompassing mitotic recombination, gene conversion, chromosome

loss, and chromosomal rearrangements including deletions. We targeted previously studied

Rtt107 features with point mutations and a C-terminal truncation (Fig. 2A), and examined the

impacts of these mutations on spontaneous LOH rates. The interaction of Rtt107 with γH2A is

largely disrupted by the *rtt107-K887M* mutation [10,19], and C-terminal truncation of Rtt107

after aa820 (*rtt107ΔBRCT5/6*) was used to disrupt any other protein–protein interactions that

may require BRCT5/6 [10]. We used the *rtt107-K426M* mutation within BRCT1–4 to disrupt the

binding of Rtt107 to phosphorylated H4T80 [7], although it should be noted that Rtt107–

BRCT1–4 also has additional binding partners that could be affected by this specific point

mutation [9,14,15,20]. We examined other regions of Rtt107 using nonphosphorylatable

mutations of each SQ/TQ motif in the SQ/TQ cluster of Rtt107 (*rtt107-4A*) [8], and a mutation

designed to disrupt the functions of Rtt107– BRCT1/2 (*rtt107-K192M*) [10]. First, we confirmed

that plasmid-borne *RTT107* reduced the rate of LOH in the *rtt107Δ* mutant to close to that seen

in wild-type cells (Fig. 2B) consistent with previous reports of a bimater phenotype for the

rtt107Δ mutant [24,28]. Note that in these experiments, comparisons were considered

statistically significant if the 95% confidence intervals did not overlap (see *Materials and*

Methods). The *rtt107-K887M* mutation did not result in an increase in rate of LOH, suggesting

that the function of Rtt107 in this context may be independent of its localization to γH2A. In

contrast, the *rtt107ΔBRCT5/6* mutation resulted in an elevated LOH rate, similar to complete

loss of Rtt107. This suggested that BRCT5/6 played a γH2A-independent role in genome

maintenance. The *rtt107-K426M* mutation increased the rate of LOH in this assay, indicating that

BRCT1–4 also contributed to LOH prevention. In contrast, the *rtt107-K192M* and *rtt107-4A*

mutations did not affect rescue of the bimater phenotype. This was perhaps unsurprising in the

case of the *rtt107-K192M* mutation, which has been shown not to disrupt Rtt107 function in

other contexts [10]. Taken together, these data indicated that both Rtt107–BRCT1–4 and Rtt107–

BRCT5/6 contributed to LOH prevention.

We next confirmed that the function of Rtt107 in this context was indeed independent of

its localization to γH2A, as our data from the *rtt107-K887M* mutant suggested. For this purpose,

we examined whether a nonphosphorylatable *H2A-S129A* mutation affected the ability of Rtt107

to limit spontaneous LOH. The results indicated that incorporation of the *H2A-S129A* allele

increased the rate of LOH in the bimater assay, consistent with the function of γH2A in the DDR

(Fig. 2C) [52]. The *rtt107Δ* mutation increased the rate of LOH of the *H2A-S129A* mutant

indicating that Rtt107 and γH2A had independent functions in this context, and that Rtt107 did

not prevent spontaneous LOH by acting as a scaffold localized to γH2A.

3.2   Rtt107 acted in the same pathway as Rad55 to limit spontaneous LOH

Having found that multiple domains of Rtt107 contributed to its function in maintaining

genome stability but that Rtt107 localization to γH2A was not required, we next examined

whether Rtt107 limited genome instability in concert with its known physical interaction

partners. We first tested whether the role of Rtt107 in limiting spontaneous genome instability

was related to its well-characterized function in downregulating the Rad9–Dpb11 interaction,

which has been examined previously in the context of treatment with DNA-damaging agents

[16]. To determine whether the genome instability of the *rtt107Δ* mutant was due to increased

Rad9–Dpb11 interaction, we utilized a *rad9-ST462,474AA* allele (hereafter referred to as *rad9-*

2A), which disrupts the Rad9–Dpb11 interaction and partially suppresses the MMS sensitivity of

the *rtt107Δ* mutant [17,53]. We compared the *rtt107Δ*, *rad9-2A*, and *rtt107Δ rad9-2A* mutant

phenotypes in the bimater assay, and found that the *rad9-2A* allele modestly increased the LOH

rate compared to the wild-type control (Fig. S1). Introduction of the *rad9-2A* allele into the

rtt107Δ mutant background increased the rate of LOH to approximately double that of the

rtt107Δ single mutant, suggesting that spontaneous LOH in the *rtt107Δ* mutant was not caused

by Rad9–Dpb11 interaction. It should be noted that strains carrying genomic *H2A/H2B* gene

derivatives were used in this assay and subsequent experiments, and the *rtt107Δ* mutation caused

a somewhat stronger phenotype in this context compared to strains with plasmid-borne *H2A/H2B*

genes (Fig. 2C).

As Rtt107 did not prevent spontaneous LOH by regulating the Rad9–Dpb11 interaction

or localizing to γH2A, we next examined whether Rtt107 limits genome instability by

cooperating with its other known or putative physical interaction partners, specifically Slx4,

SMC5/6, Mms22, Mms4, Sgs1, and Rad55 [9,12,21,44–46]. For this purpose, we compared the

bimater phenotype of the *rtt107Δ* mutant to strains with mutations of Rtt107 physical interaction

partners as well as corresponding double mutants (Fig. 3A–F). Specifically, we sought to identify

a mutation in an Rtt107 interaction partner that had a similar spontaneous LOH phenotype to the

rtt107Δ mutant, and that exhibited epistasis with the *rtt107Δ* mutation. We used a hypomorphic

smc6-9 allele to examine the essential SMC5/6 complex (Fig. 3B) [54], as the *smc6-9* mutant

shows a defect similar to the *rtt107Δ* mutant in DSB relocalization to the nuclear periphery [55].

We used an *mms4-8A* allele to reduce the interaction between Rtt107 and Mus81–Mms4 (Fig.

3D) [45]. Of the mutations examined, only the *rtt107Δ rad55Δ* double mutant had a phenotype

similar to those of the *rtt107Δ* and *rad55Δ* single mutants, indicating an epistatic interaction

between the two mutations (Fig. 3F). This pattern was not observed for mutants of other Rtt107

binding partners, and was consistent with an overlap in function between Rtt107 and Rad55 in

preventing spontaneous LOH.

3.3   Rtt107 acted in the same pathway as Rad55 to limit spontaneous mitotic RCOs

As the *rtt107Δ*, *rad55Δ*, and *rtt107Δ rad55Δ* mutants all exhibited a similar LOH

phenotype, we next interrogated the relation between Rtt107 and Rad55 in greater detail. Given

the complexities and interrelations between genes involved in the DDR, we first tested whether

the relation observed between the *rtt107Δ* and *rad55Δ* mutations was dependent on strain

background. Specifically, a *rad5-535* allele is present in the canonical W303 yeast strain

background used here, and this can possibly affect findings of DDR studies [56]. To test whether

this mutation might affect our results, we repeated the bimater assay in a *RAD5*-corrected W303

background. Reassuringly, we observed a similar pattern of LOH phenotypes (Fig. 4A), although

perhaps not surprisingly the LOH rates for each strain were much lower than those in the *rad5-*

535 background (Fig. 3F). This indicated that the epistatic interaction between the *rtt107Δ* and

rad55Δ mutations was not dependent on the presence of the *rad5-535* allele, although LOH rates

may be mediated in some way by Rad5, and again suggested that Rtt107 may act in the same

pathway as Rad55 to limit spontaneous LOH

As a shared function for Rtt107 and Rad55 has not yet been identified, we sought to

characterize the degree of cooperation between Rtt107 and Rad55 in limiting spontaneous

genome instability in more detail. LOH in the bimater assay is predominantly mediated by

crossover events produced by mitotic recombination in wild-type yeast cells [39]. Given that the

Rad55–Rad57 heterodimer acts as a mediator of homologous recombination, which can produce

mitotic RCO, we hypothesized that spontaneous RCO may occur in the *rtt107Δ* and *rad55Δ*

mutants. Therefore, we tested whether Rtt107 and Rad55 limited the rate of spontaneous mitotic

RCO, and whether they did so by acting within the same pathway. We examined the *rtt107Δ* and

rad55Δ mutants by sectoring assay for spontaneous mitotic RCOs within a 120-kb interval on

chromosome V in diploid cells, as described previously [49]. It should be noted that this and all

subsequent experiments regarding the relation between *RTT107* and *RAD55* were performed in

the *RAD5*-corrected W303 background, as described above. The *rtt107Δ* and *rad55Δ* mutants

showed higher frequencies of spontaneous RCOs than the wild-type control (Fig. 4B). The

rtt107Δ single mutant and *rtt107Δ rad55Δ* double mutant showed similar increases in mitotic

RCO frequency, suggesting that Rtt107 and Rad55 limited RCOs by acting within the same

pathway.

Next, we tested whether Rtt107 and Rad55 cooperated to limit other genomic events in

addition to RCOs that may have caused LOH in the diploid bimater assay, such as gene

conversion, chromosomal rearrangements, and chromosome loss. Each of these events, other

than RCOs, can be captured by the ALF assay that measures loss of mating locus function in

haploid cells [39]. Consistent with an earlier screen, quantitative ALF assay showed that the

rtt107Δ and *rad55Δ* mutants both exhibited higher mating rates than the wild-type (Fig. 4C) [39].

The *rtt107Δ rad55Δ* double mutant phenotype trended toward a slightly higher mating rate than

either of the single mutants, although the difference was not statistically significant. These

observations suggested that Rtt107 and Rad55 may partly act within the same pathway to limit

the types of genome instability that are captured by the ALF assay. However, as it remained

formally possible that Rtt107 and Rad55 had independent functions in this context, we focused

on exploring other genome instability phenotypes.

The observations outlined above suggested that Rtt107 cooperated with Rad55 to prevent

specific types of genome instability at the mating locus and on chromosome V in diploid cells.

Next, to test whether Rtt107 prevented various types of spontaneous instability more broadly at

different genetic loci, we examined spontaneous recombination at the sensitive rDNA locus [43].

Specifically, we used a quantitative version of an assay for rDNA recombination based on the

loss of an *ADE2-CAN1* marker in the rDNA of haploid yeast to monitor rDNA instability [50].

The *rtt107Δ* mutant exhibited rDNA hyperrecombination, as reported previously (Fig. 4D) [38].

The *rad55Δ* mutant exhibited a modest (~2-fold) increase in spontaneous rDNA recombination,

which was significantly less than that in the *rtt107Δ* mutant (~5-fold), suggesting that Rtt107 did

not primarily maintain rDNA stability by cooperating with Rad55. Consistent with largely

separate roles of Rtt107 and Rad55 in rDNA maintenance, the *rtt107Δ rad55Δ* double mutant

showed a trend toward greater instability than the *rtt107Δ* single mutant, although this increase

was not statistically significant. We considered that Rtt107 may instead maintain rDNA stability

with its major binding partner, Slx4, which maintains rDNA structure during DNA replication

[57]. Both the *slx4Δ* single mutant and *rtt107Δ slx4Δ* double mutant exhibited rDNA

hyperrecombination at rates similar to that of the *rtt107Δ* mutant (Fig. S2), which suggested that

Rtt107 maintained rDNA stability by cooperating with Slx4 rather than Rad55.

Our results up to this point supported substantial overlap in function between Rtt107 and

Rad55 in preventing specific types of spontaneous genome instability. We next examined the

degree of cooperation between Rtt107 and Rad55 during chronic treatment with DNA-damaging

agents to clarify the degree of cooperation between Rtt107 and Rad55 in the DDR. Focusing on

agents previously shown to cause sensitivity to strains lacking Rtt107, we examined whether

Rtt107 and Rad55 acted in the same DDR pathway to confer resistance to the DNA-damaging

agents MMS, hydroxyurea (HU), or camptothecin (CPT) in a growth assay (Fig. 4E). Given the

extreme sensitivity of the *rad55Δ* mutant to these agents, phenotypic comparison of the *rad55Δ*

mutant and *rtt107Δ rad55Δ* double mutant required their use at low concentrations. However, it

should be noted that the reported sensitivity of the *rtt107Δ* mutant to these agents was not readily

apparent at the low concentrations used in this study [8,58]. We first confirmed that the *rtt107Δ*

rad55Δ mutant exhibited enhanced MMS sensitivity compared to single mutants, as reported

previously [20]. Next, we found that the *rad55Δ* mutant exhibited much greater sensitivity to HU

and CPT than the *rtt107Δ* mutant. The *rtt107Δ rad55Δ* double mutant also showed similar HU

sensitivity and a slight but reproducible increase in CPT sensitivity compared to the *rad55Δ*

single mutant (Fig. 4E). Taken together, these results suggested that Rtt107 was not required for

the function of Rad55 in the context of resistance to chronic DNA damage.

The observations outlined above suggested that Rtt107 cooperated with Rad55 to limit

LOH and mitotic RCOs, and we therefore hypothesized that Rtt107 may limit genome instability

through a DDR pathway that is known to be regulated by Rad55. Given that Rtt07 may not be

required for Rad55's canonical DSB repair function [20], we considered that Rtt107 may

cooperate with Rad55 in regulating DNA damage tolerance, as mutants deficient in

recombination can rely on tolerance to limit spontaneous LOH [59]. To test our hypothesis, we

compared the spontaneous LOH phenotype of the *rtt107Δ* mutant with those of a *rev3Δ* mutant

deficient in TLS, a *ubc13Δ* mutant deficient in error-free PRR, and a *rad18Δ* mutant deficient in

both pathways [60,61]. Consistent with previous work, the *rad18Δ* mutant exhibited a modest

(~2-fold) but significant increase in spontaneous LOH [39] (Fig. S3). The *ubc13Δ* mutation did

not affect the rate of spontaneous LOH in wild-type or *rtt107Δ* mutant, indicating that Rtt107 did

not primarily limit LOH by promoting error-free PRR. In addition, the *rev3Δ* mutation increased

the LOH phenotype of the *rtt107Δ* mutant, although this effect was modest (~1.5-fold). The

findings outlined above indicated that Rtt107 acted in the same pathway as Rad55 to limit

spontaneous LOH and mitotic RCOs, but that Rtt107 did not cooperate with Rad55 in all

contexts.

3.4   The *rad55-S404A* mutation largely disrupted the Rtt107–Rad55 interaction

To further delineate the mechanisms underlying the genetic relationship between *RTT107*

and *RAD55*, we tested whether the physical interaction between Rtt107 and Rad55 was important

for preventing genome instability. We considered that the Rtt107–Rad55 interaction may be

mediated by Rad55-S404 phosphorylation, which does not require DNA damage induction [62],

as the Rtt107–Rad55 interaction occurs under normal growth conditions [20]. To test this

hypothesis, we generated a *rad55-S404A* mutation and established that it did not affect Rad55

protein level (Fig. S4). To determine whether the *rad55-S404A* mutation affected the Rtt107–

Rad55 interaction, we performed a small-scale interaction assay using FLAG-Rad55 from log-

phase cultures (Fig. 5A). We observed copurification of Rad57-HA and Rtt107-VSV with

Rad55, which was consistent with a previous report [20]. The *rad57Δ* mutation did not eliminate

the copurification of Rtt107 with Rad55, indicating that the Rtt107–Rad55 interaction was

independent of Rad57. As Rtt107 was also found in Rad57 immunoprecipitations [20], these

results were consistent with a model in which Rtt107 interacts with Rad55 to form an Rtt107–

Rad55–Rad57 trimeric complex. Of note, the *rad55-S404A* mutation substantially reduced the

amount of Rtt107, but not Rad57, that copurified with Rad55, suggesting that the Rad55-S404

residue was indeed important for the interaction with Rtt107, while having no appreciable effect

on the Rad55–Rad57 interaction. For our purposes, we therefore concluded that this mutation

was appropriate for investigating the importance of the Rtt107–Rad55 interaction in functional

assays.

A previous report suggested that mutation of the Rad55-S404 site does not affect the

ability of cells to grow during chronic treatment with MMS [62]. As the phenotypes of DNA

repair mutants can be highly dependent on the type of DNA-damaging agent used as well as the

strain background, we examined whether the *rad55-S404A* mutation affected the ability of our

strains to grow during chronic exposure to MMS, HU, or CPT (Fig. 5B). We found that the

rad55-S404A mutation did not cause sensitivity to any of these agents at concentrations that

severely inhibited growth of the *rad55Δ* mutant. Taken together, these observations suggested

that the *rad55-S404A* mutation left Rad55 function in this context largely intact and that the

interaction between Rtt107 and Rad55 was not crucial for growth during chronic exposure to

DNA-damaging agents.

3.5   The *rad55-S404A* mutation increased spontaneous LOH and RCO rates

As the *rad55-S404A* mutation largely disrupted the physical Rtt107–Rad55 interaction,

we next used it to examine whether the interaction between Rtt107 and Rad55 contributed to

limiting spontaneous genome instability. First, we again employed the bimater assay to examine

LOH induced by a broad range of events. The *rad55-S404A* mutation caused a modest (~2-fold)

increase in spontaneous LOH in comparison to the wild-type strain (Fig. 6A), consistent with a

model in which Rtt107 and Rad55 prevented spontaneous RCOs in part by binding to each other.

However, we noted that the *rad55Δ* mutation caused a much greater increase in LOH rate in the

bimater assay (~10-fold) (Fig. 4A). Further, the LOH phenotype of the *rtt107Δ* mutant was also

greater than that of the *rad55-S404A* point mutant, and the *rtt107Δ rad55-S404A* double mutant

exhibited a higher rate of LOH than either single mutant alone, suggesting that the function of

Rtt107 in preventing LOH may not be solely dependent on the Rad55-S404 site and the Rtt107–

Rad55 interaction

To more granularly determine the types of spontaneous genome instability that may be

prevented by Rtt107–Rad55 interaction, we again used the sectoring assay to specifically

examine mitotic RCO events. The *rad55-S404A* mutation increased the frequency of mitotic

RCOs, indicating that the Rad55-S404 site contributed to the limitation of spontaneous RCOs,

and suggesting that the interaction between Rtt107 and Rad55 may be important in this context

(Fig. 6B). The *rtt107Δ rad55-S404A* double mutant phenotype was similar to that of the *rtt107Δ*

single mutant, and this pattern was broadly similar to the epistasis observed between the *rtt107Δ*

and *rad55Δ* mutations in the context of the RCO phenotype (Fig. 4B), consistent with a shared

role for Rtt107 and the Rad55-S404 site in limiting mitotic RCOs.

To further delineate the functional relevance of the Rtt107–Rad55 interaction, we

considered that the spontaneous LOH and RCO phenotypes of the *rad55-S404A* mutant may be

indicative of a defect in a specific DNA repair pathway, possibly the crossover-generating

homologous recombination pathway. We therefore examined whether the *rad55-S404A* and

rtt107Δ mutants exhibited various types of spontaneous genome instability related to

recombinational repair using an established assay that discriminates between several distinct

types of recombination and mutation in diploid cells (Figure 1) (see [49] for details). In this

assay, Class 1/2 events primarily represent RCOs and break-induced replication (BIR) events.

Class 3/4 events primarily represent local gene conversions that occur without crossovers, or *de*

novo mutations occurring within the markers. Class 5/6 events primarily represent loss of one

copy of chromosome V by nondisjunction. Using this approach, we found that the *rtt107Δ*

mutant exhibited higher rates of Class 1 and Class 3/4 events but wild-type rates of Class 5/6

events (Fig. 6C). In the case of Class 2 events, the *rtt107Δ* mutant showed a trend toward a

higher spontaneous mutation rate than wild-type, but this increase was not statistically

significant. We further examined the Class 3 events in the *rtt107Δ* mutant by colony PCR and

found that 24 of 28 events (86%) represented gene conversion rather than *de novo* mutation

within the marker (see *Materials and Methods*). These observations suggested that *rtt107Δ*

mutants have elevated rates of gene conversion, RCOs, and possibly BIR, but not chromosome

loss. The *rad55-S404A* single mutant did not exhibit significantly increased rates in any of these

classes, and the *rtt107Δ rad55-S404A* double mutant resembled the *rtt107Δ* single mutant. In

contrast to the importance of the Rad55-S404 phosphorylation site for limiting LOH and RCOs,

these results indicated that Rad55-S404 was not essential to limit multiple other types of

spontaneous genome instability that are common in *rtt107Δ* mutants.

3.6   *RTT107* and *RAD55/RAD57* acted in similar genetic pathways under normal growth

conditions

Our findings suggested that Rtt107 cooperated with Rad55 to limit spontaneous LOH and

RCOs, and that this function was at least partially distinct from its role in the response to chronic

DNA-damaging agent exposure. Therefore, we hypothesized that the genetic interaction

spectrum of *RTT107* drawn from studies that did not apply DNA-damaging agents may provide

relevant insights regarding the relations between *RTT107* and *RAD55*, and potentially also

RAD57. To test this hypothesis, we compared published genetic interaction data on *RTT107*,

RAD55, and *RAD57* from several high-throughput screens performed under normal growth

conditions, and determined which genetic interactions were shared between at least two of these

genes (Table 1) [63–71]. Of the 99 genetic interactions shared between *RAD55* and *RAD57*, we

found that 40 were also shared with *RTT107*. This represented significant overlap between genes

that interacted with *RTT107* and genes that interacted with *RAD55* + *RAD57* (Fisher's exact test,

two sided, $p < 2.2 \times 10^{-16}$). Fewer interactions (8) were shared between either *RTT107* + *RAD55*

or *RTT107* + *RAD57* only. While further analyses will be required to fully establish the

relationships between *RTT107*, *RAD55*, and *RAD57*, the greater than expected by chance overlap

of shared genetic interactors suggested that these genes may act, at least in part, in the same

pathways under normal growth conditions.

     Table 1. Shared and distinct genetic interaction profiles between *RTT107*, *RAD55,* and

     RAD57

Shared genetic interaction partners	Genes with shared interactions		Total shared interactions
	Negative interactions	**Positive interactions**	
RAD55 + RAD57 (only)	*BCS1, BIM1, BRE1, CIK1, CSM3, DBF2, DEP1, DFR1, GET1, HOM6, HSL1, HYR1, IMG2, IRC19, LGE1, LRS4, LSM6, LSM7, MCM1, MDM12, MET7, MMS22, MRC1, NAT1, NPT1, NUP133, POL3, POL31, POL32, PRI1, PRI2, RAD5, REV1, REV3, REV7, RFA2, RFC2, RFC4, RFC5, RIM1, RPN11, RPN4, RVS161, SKN7, SOH1, SWS2, UBP8, VAC7, VMA3, VPS34, YAP1, YBP1, YFH1, YJR124C*	*MMS21, RAD51, RAD52, RAD54, TOP3*	**59**
RAD55 + RAD57 + RTT107	*AIM10, ASA1, ASF1, CCS1, CDH1, CLA4, CSM1, CTF18, CTF4, CTF8, DCC1, DIA2, DOC1, DST1, DUN1, EAF1, ELG1, GET2, LTE1, MMS1, MON2, NPL3, NUP60, NUP84, POL12, POL30, RAD18, RAD27, RAD6, RFA3, RFC1, RPN10, RPN8, RPT1, RRM3, SLX5, SLX8, SOD1, STP22, TSA1*		**40**
RTT107 + RAD55 (only)	*CCR4, CDC9, HUR1, PMR1, POP2, RML2, SMC5, SWI6*		**8**
RTT107 + RAD57 (only)	*CLB2, DOA1, ISU1, RPN12, RPT4, SAC3, VID22*	*PAT1*	**8**

The listed genes shared the same type of genetic interaction between at least two of *RTT107*,
RAD55, and *RAD57*. Gene deletions or loss-of-function alleles of essential genes were used in
these studies. "Negative" interactions caused either slower growth or lethality. Note that genes in
the "*RAD55 + RAD57*" row did not also share an interaction with *RTT107*.

4   Discussion

Although Rtt107 is known as a multi-BRCT scaffold with homologs in fission yeast and

vertebrates [11], its specific yet important function in limiting spontaneous genome instability

phenotypes has remained unclear. Here, we showed that the role of Rtt107 in preventing

spontaneous LOH was distinct from its well-characterized functions as a scaffold that localizes

to γH2A after induction of DNA damage. Instead, Rtt107 cooperated with the recombination

mediator Rad55, at least in part, to limit spontaneous LOH and mitotic RCOs in diploid cells.

The Rad55-S404 phosphorylation site was partially required for the Rtt107–Rad55 protein–

protein interaction and to limit spontaneous LOH and RCOs, suggesting that the physical

interaction between Rtt107 and Rad55 contributed to the suppression of these events. In contrast,

the cooperation between Rtt107 and Rad55 did not extend to prevention of all types of

spontaneous genome instability. For example, Rtt107 acted in the same pathway as Slx4 to limit

rDNA recombination. Taken together, our results indicated that Rtt107 acted with either the

recombination mediator Rad55 or its well-characterized binding partner Slx4 to limit types of

genome instability that occur spontaneously under normal growth conditions.

4.1 Rtt107 limited spontaneous genome instability in distinct pathways

The role of Rtt107 in limiting genome instability differed between various types of

instability. Rtt107 cooperated with Rad55 to prevent LOH at the mating locus and mitotic RCOs

on chromosome V in diploid cells, likely in part by binding to Rad55 in a manner dependent on

Rad55-S404 (Fig. 7A). Our data were also consistent with a model in which Rtt107 interacts

with the complete Rad55–Rad57 heterodimer by binding primarily to Rad55, and genetic

interaction data suggested that *RTT107*, *RAD55*, and *RAD57* can act in similar pathways under

normal growth conditions (Table 1). However, the Rad55-S404 site, while important for

mediating the physical interaction between Rad55 and Rtt107, was not needed to limit other

types of instability that are common in diploid cells, specifically gene conversion and possibly

BIR (Fig. 6C). These observations suggested that Rtt107 primarily acted independently of the

Rtt107–Rad55 interaction to limit these events, whereas the Rtt107–Rad55 interaction may

specifically limit LOH and mitotic RCOs. We also found that the function of Rtt107 in the

context of LOH was dependent on BRCT5/6 but independent of the Rtt107-K887 residue that

mediates interaction with γH2A (Fig. 2B) [10,19]. This was in contrast to earlier observations

indicating that *rtt107-K887M* and *rtt107ΔBRCT5/6* mutants exhibit similar phenotypes with

respect to growth during chronic MMS treatment and the recruitment of Rtt107 to induced DSBs

[10]. Interestingly, the structure of BRCT5/6 is remarkably flexible, and protein binding via

BRCT5/6 using surfaces other than the γH2A-binding pocket has been proposed previously [19].

Taken together, these findings suggested that Rtt107–BRCT5/6 may bind to a non-γH2A partner

and thus prevent LOH (Fig. 7A), and the identification of such a partner represents an

opportunity for future research. It is of interest to consider that Rad55 binds to Rtt107–BRCT1–4

[14,20], and that Rtt107 may act to bridge Rad55, or the complete Rad55–Rad57 heterodimer, to

a partner that binds BRCT5/6.

Rtt107 cooperated with Slx4 to prevent hyperrecombination at the rDNA locus in haploid

cells, while not depending primarily on Rad55 in this context. Recent work suggested that Rtt107

limits rDNA recombination at least in part as a γH2A-binding scaffold for its physical interaction

partners, including Slx4 [14]. Therefore, the cooperation of Rtt107 with Slx4 in rDNA

maintenance may involve binding to γH2A via Rtt107–BRCT5/6 (Fig. 7B). We note that Rtt107

did not rely on Slx4 to prevent spontaneous LOH (Fig. 3A), again suggesting that Rtt107 acts in

different pathways to limit different types of spontaneous genome instability.

In addition to the spontaneous genome instability phenotypes examined in this study, the

rtt107Δ mutant exhibits elevated levels of spontaneous DNA damage foci in $S/G_2/M$ phase

[15,25,38]. Interestingly, the *rtt107Δ* mutation does not exhibit broad epistasis with mutations of

canonical DNA damage repair or tolerance pathways [8,20,44,72,73]. Taken together, these

findings were consistent with a model in which Rtt107 maintains genome stability at least in part

by limiting the levels of DNA damage that occur during replication, rather than by directly

affecting a specific repair process or pathway choice. Furthermore, it is interesting that DSBs

appear to be the primary initiator of spontaneous crossovers in yeast [74–76]. The dispensability

of Rtt107 for survival after exposure to DSB-inducing ionizing radiation [8], along with the

requirement for Rtt107 in preventing spontaneous crossovers, could be reconciled by a model in

which Rtt107 prevents the occurrence of DSBs but is not essential for DSB repair.

4.2 Cooperation between Rtt107 and Rad55 in genome maintenance

Our genetic evidence suggested that Rtt107 interacted with Rad55 to limit rates of

spontaneous RCOs, and functioned at least in part with Rad55 to limit LOH. We note that the

LOH phenotype examined in this study could have been largely driven by mitotic RCO events,

and that these phenotypes would be consistent with a relationship between the Rtt107–Rad55

interaction and recombinational repair pathways, which are mediated by Rad55–Rad57.

However, it remains unclear how Rtt107 and Rad55 would cooperate to prevent spontaneous

genome instability. In complexes such as Rtt107–Slx4, Rtt107–SMC5/6, or Rtt107–Rtt101–

Mms1–Mms22, Rtt107 has commonly been proposed to act as a scaffold that recruits its binding

partners to γH2A near DNA damage lesions (Fig. 7B) [9,10,14,16,55]. In contrast, our results

presented here suggested that Rtt107–γH2A binding was actually dispensable for the function of

Rtt107 in preventing LOH, and a model in which Rtt107 simply recruits Rad55 to lesions thus

appears less applicable. Rtt107 and Rad55 also localize to different DNA damage signals, as

Rad55 associates with the Rad51–ssDNA filament during recombination [26,77], while Rad51

does not appear to assemble nonspecifically onto double-stranded DNA [78]. Rtt107 localizes to

the histone marks γH2A and H4T80ph, which flank DSB sites and are excluded from ssDNA

[7,10,78]. Therefore, while Rtt107 and Rad55 both localize near DSBs in broad terms, their

known recruitment pathways appear to target different regions around DSBs. Rad51 can be

detected in similar regions to γH2A due to homology search performed by the Rad51 filament

[78], but it is unclear whether the Rtt107–Rad55 interaction is important during homology search

or some other process. Alternatively, Rtt107 and Rad55 may cooperate at sites other than DSBs,

such as stalled replication forks. The localization of Rtt107 behind stressed replication forks has

been shown to be γH2A-dependent [13], whereas the recruitment of Rad55 to replication forks is

less well understood [79,80] and it is unclear whether Rtt107 and Rad55 can be recruited to

stressed forks at similar times.

While our findings were consistent with a relationship between the Rtt107–Rad55

interaction and recombinational repair, it is also important to note that Rtt107 does not suppress

recombination at all loci. For example, a genome-wide study that identified rDNA

hyperrecombination in the *rtt107Δ* mutant found normal rates of recombination at the *LEU2*

locus in two assays [38].

We found that the Rad55-S404 phosphorylation site was important for the physical

interaction between Rtt107–Rad55 and for limiting mitotic RCOs and LOH. However, in the

context of the LOH phenotype, we did not identify clear epistasis between the *rtt107Δ* and

rad55-S404A mutations (Fig. 6A), which suggested that the Rad55-S404 site may have roles in

Rad55's function beyond the complexes or pathways involving Rtt107. Further, it is possible that

residual binding between Rtt107 and Rad55 occurs in the *rad55-S404A* mutant, limiting the

impact of the targeted *rad55-S404A* mutation on certain genome instability phenotypes, although

this was not detectable in the assays employed here. In contrast, we observed epistasis between

the *rtt107Δ* and *rad55Δ* mutations in the context of the LOH phenotype (Fig. 3F and 4A),

suggesting that Rtt107 and Rad55 act broadly in the same pathway to prevent LOH.

It should also be noted that we identified cooperation between Rtt107 and Rad55 in

assays using diploid yeast, whereas Rtt107 did not cooperate primarily with Rad55 to prevent

rDNA recombination in haploids (Fig. 4D). Diploid and haploid yeast differ in DDR regulation,

with diploids utilizing HR over NHEJ for DSB repair [30]. Further, the sensitivity of *rad55Δ*

mutants to ionizing radiation is suppressed by heterozygosity at the mating locus, which occurs

naturally in diploids [30,31]. Therefore, the degree of cooperation between Rtt107 and Rad55 in

genome maintenance may be partly dependent on ploidy.

We identified a spontaneous LOH phenotype for an *rtt107-K426M* mutant (Fig. 2B),

suggesting that the K426-dependent interaction of Rtt107 with phosphorylated H4T80 may also

be important for preventing spontaneous LOH, possibly by affecting the timing of Rtt107

recruitment [7]. However, we note that K426 is within Rtt107–BRCT1–4, and that BRCT1–4

interacts with a number of interaction partners of Rtt107, including Rad55 [9,14,15,20].

Therefore, it remains possible that the *rtt107-K426M* mutation could also affect the Rtt107–

Rad55 complex, or other Rtt107 complexes.

4.3 Cooperation between Rtt107 and Slx4 in rDNA maintenance

We found that Rtt107 acted in the same pathway as Slx4 to maintain rDNA stability (Fig.

S2). The role of Rtt107 in rDNA maintenance may be similar to its role in GCR suppression in

haploid cells, in which Rtt107 acts at least in part as a γH2A-binding scaffold for Slx4, Mms22,

and Nse6 (Fig. 7B) [9,14,15,20]. It is not yet clear whether Rtt107 also cooperates with the Slx1

endonuclease in this context. Although an Rtt107–Slx1–Slx4 complex has been identified [15],

Rtt107 has not been previously implicated in the endonuclease function of Slx1–Slx4. In

addition, we note that the rDNA hyperrecombination phenotype of the *slx4Δ* mutant identified in

this study differs from a previous report in which the *slx4Δ* mutant exhibited lower rates of

rDNA recombination than wild-type [81]. This discrepancy may be related to measurement of

the loss of a single *ADE2* marker in the rDNA in the earlier study, rather than the combined

ADE2-CAN1 marker used here and in a recent study identifying rDNA hyperrecombination in an

slx4 mutant [14,50]. Relatedly, the *Schizosaccharomyces pombe* homolog of Rtt107, Brc1, has

been proposed to play a role in genome maintenance in regions that are susceptible to replication

stress, including the rDNA [82], likely at least in part due to recruitment of Brc1 to γH2A, with

the γH2A–Brc1 foci colocalizing with rDNA [83].

Recent work identified GCR and rDNA recombination phenotypes for mutations of the

Rtt107 interaction motifs (RIMs) that are important for the interaction between Rtt107–BRCT1–

4 and three of the binding partners of Rtt107, i.e., Slx4, Mms22, and the SMC5/6 subunit, Nse6

[14]. These *RIM* mutants each exhibited mild rDNA hyperrecombination phenotypes in

comparison to the more pronounced phenotype of the *rtt107Δ* mutant, suggesting that individual

interactions may have had limited or partially redundant contributions to the rDNA maintenance

function of Rtt107. Our results were consistent with substantial cooperation between Rtt107 and

Slx4 in rDNA maintenance, as the *slx4Δ* and *rtt107Δ* mutations had similar and epistatic effects

on rDNA stability (Fig. S2).

4.4 Distinct roles for Rtt107 in genome maintenance and DNA damage resistance

In this study, we found that the *rad9-2A* and *H2A-S129A* mutations increased the LOH

rate of the *rtt107Δ* mutant (Fig. S1 and 2C), despite suppressing the hypersensitivity of the

rtt107Δ mutant to chronic MMS treatment [16,17,22]. Previously, we reported a similar pattern

in which the *dot1Δ* mutation suppressed the MMS sensitivity of the *rtt107Δ* mutant, but did not

suppress its spontaneous LOH phenotype [25]. Each of these mutations that suppress the MMS

sensitivity of the *rtt107Δ* mutant can also dampen DNA damage checkpoint activity [53,84–87],

and the *H2A-S129A* mutation additionally suppresses the MMS-induced checkpoint

hyperactivation phenotype of the *rtt107Δ* mutant [16,17]. The lack of suppression by these

mutations of the LOH phenotype in the *rtt107Δ* mutant reported here suggested that the role of

Rtt107 in preventing spontaneous LOH was largely distinct from its role in preventing treatment-

induced checkpoint hyperactivation. This distinction was further emphasized by the observations

that the *rtt107Δ* and *rad55Δ* mutations exhibited epistasis in the context of the LOH phenotype

(Fig. 3F and 4A) and a negative genetic interaction in the context of growth during MMS

exposure (Fig. 4E) [20]. We also found that *rtt107Δ* mutation suppressed the LOH phenotype of

the *mms22Δ* mutant (Fig. 3C), despite the additive effects of *rtt107Δ* and *rtt101Δ* mutations on

MMS sensitivity observed in a growth assay [88]. Taken together, these findings indicated that

while abundant information is available on the genetic interactions of the *rtt107Δ* mutant in the

context of exposure to DNA-damaging agents, these data may not directly reflect the function of

Rtt107 in preventing spontaneous LOH.

The lack of a DNA damage sensitivity phenotype for the *rad55-S404A* mutant suggested

that the Rtt107–Rad55 interaction was not important for DNA damage resistance (Fig. 5B). This

was unsurprising, as the *rtt107Δ* mutant was less sensitive to DNA-damaging agents than the

rad55Δ mutant (Fig. 4E). In addition, a previous genome-wide screen in a different genetic

background performed in our laboratory revealed a negative genetic interaction between the

rtt107Δ and *rad55Δ* mutations in the context of CPT sensitivity [73]. However, there was no

clear negative interaction between the *rtt107Δ* and *rad55Δ* mutations at this CPT concentration

in the W303 *RAD5* genetic background, although the sensitivity of the *rtt107Δ rad55Δ* double

mutant was slightly greater than that of the *rad55Δ* single mutant (Fig. 4E).

Taken together, the findings presented here showed that Rtt107 plays multiple roles in the

prevention of spontaneous genome instability, which are often distinct from its previously

studied functions in the response to DNA-damaging treatments. Further studies are required to

determine the specific mechanisms by which Rtt107 prevents various types of genome instability

with its diverse interaction partners, including Rad55 and Slx4.

Acknowledgments

We thank G. Leung for preliminary work in strain construction; W. D. Heyer, J. Diffley, C. D.

Allis, B. Pfander, L. Aragon, and J. Krebs for generously providing yeast strains and plasmids;

and M. Aristizabal, P. Stirling, W. D. Heyer, and A. Kerr for critical reading of the manuscript

and suggestions.

Figure Captions

Fig. 1. Targeted spontaneous genome instability assays used in this study.

The locations of mutations and genomic rearrangements in each assay are shown. All assays were performed under normal growth conditions. The graph colors shown on the right correspond to the colors of graphs for each assay in subsequent figures.

Fig. 2. Rtt107–BRCT1–4 and Rtt107–BRCT5/6 were important for limiting spontaneous LOH.

A) Rtt107 structural features mutated in panel B. BRCT domains are numbered and shown in gray. Target lysines are labeled in the center of the BRCT domain in which they are located. The orange circle indicates the SQ/TQ cluster of Rtt107, targeted by the *rtt107-4A* mutation. B) Rtt107-K426 and Rtt107–BRCT5/6 contributed to genome maintenance in the bimater assay, performed using homozygous diploid strains. Mating rates and 95% confidence intervals were calculated using the maximum likelihood method, and statistical significance was assessed by the non-overlap of confidence intervals (see *Materials and Methods*). +, non-overlap of 95% confidence intervals between two strains (statistically significant); ns, overlap of confidence intervals (not significant). Strains indicated in this panel were in the W303 *RAD5* background, and were grown in SC-Leu. At least eight independent colonies were tested. C) Rtt107 prevented spontaneous LOH independently of γH2A in the bimater assay. The *H2A-S129A* mutation caused genome instability, and the *rtt107Δ H2A-S129A* double mutant phenotype was greater than those of either single mutant. Mating rates and 95% confidence intervals were calculated and compared as described for panel B. Eleven independent colonies were tested.

Fig. 3. Rtt107 limited spontaneous LOH in the same pathway as Rad55.

All panels depict bimater assays. The mating rates and 95% confidence intervals were calculated

and compared as described for the bimater assay in Figure 2B. ns, overlap of confidence

intervals (not significant). A simplified model of the corresponding physical interaction of

Rtt107 is indicated beneath each assay, and mutated complex members are shown in color.

Experiments were performed in the W303 *rad5-535* background except where noted. Note that

the scale on the *y*-axis differs between panels.

A) The *slx4Δ* mutation did not increase the rate of spontaneous LOH or affect the *rtt107Δ* mutant

phenotype. At least six independent colonies were tested. B) The *smc6-9* mutation caused far

greater instability in the bimater assay than the *rtt107Δ* mutant. The phenotype of the *rtt107Δ*

smc6-9 double mutant was not distinguishable from that of the *smc6-9* single mutant. At least

seven independent colonies were tested. C) The *mms22Δ* mutation caused instability in the

bimater assay. The *mms22Δ* mutant exhibited a higher rate of LOH than the *rtt107Δ* mutant, and

this phenotype was suppressed by the *rtt107Δ* mutation. Ten to thirteen independent colonies

were tested. D) The *rtt107Δ mms4-8A* double mutant exhibited a greater rate of LOH than either

single mutant in the bimater assay. At least seven independent colonies were tested. Note that

this experiment was performed in the W303 *RAD5* background, and that the complex model

depicted in this panel is a hypothetical arrangement. E) The *sgs1Δ* mutant exhibited a higher

LOH rate than the *rtt107Δ* mutant in the bimater assay. The *rtt107Δ sgs1Δ* double mutant

exhibited a higher LOH rate than either single mutant. Eleven independent colonies were tested.

F) The *rtt107Δ rad55Δ* double mutant exhibited a similar LOH rate to the *rtt107Δ* and *rad55Δ*

single mutants. At least 12 independent colonies were tested.

Fig. 4. Rtt107 and Rad55 limited spontaneous mitotic RCO frequency in the same pathway.

Experiments were performed in the W303 *RAD5* background. Rates and 95% confidence

intervals for the bimater assay (panel A), ALF assay (panel C), and rDNA instability assay

(panel D) were calculated and compared as described for the bimater assay in Figure 2B. +, non-

overlap of 95% confidence intervals between two strains (statistically significant); ns, overlap of

confidence intervals (not significant). A) The *rtt107Δ rad55Δ* double mutant exhibited a similar

LOH rate to the *rad55Δ* single mutant in the bimater assay. Findings were similar to those in the

W303 *rad5-535* background (Fig. 3F). Eight to 10 independent colonies were tested. B) The

rtt107Δ, rad55Δ, and *rtt107Δ rad55Δ* mutants exhibited a similar increase in spontaneous mitotic

RCO frequency. Canavanine-resistant sectored colonies were scored as representative of RCOs.

Frequencies were calculated from 13–16 independent colonies. **, $p < 0.01$, *** $p < 0.005$

compared to the wild-type (Student's *t* test). C) The *rtt107Δ* and *rad55Δ* mutants exhibited a

higher rate of instability in the haploid ALF assay. The phenotype of the *rtt107Δ rad55Δ* double

mutant showed a trend toward a higher rate than the *rtt107Δ* mutant, but this difference was not

statistically significant. Haploid strains were used to determine mating rates in the ALF assay,

and 7–9 independent colonies were tested. D) The *rtt107Δ* and *rad55Δ* mutants exhibited a

higher rate of spontaneous rDNA recombination in comparison to wild-type. The *rtt107Δ* mutant

and *rtt107Δ rad55Δ* double mutant showed a higher rate of rDNA recombination than the

rad55Δ mutant. rDNA recombination was measured as the loss of an *ADE2-CAN1* marker in the

rDNA, and eight independent colonies were tested. E) The *rtt107Δ rad55Δ* double mutant

showed greater sensitivity to MMS treatment than either single mutant. Tenfold serial dilutions

from overnight cultures were plated on media containing the indicated concentrations of DNA-

damaging agents, and then incubated for 2 days at 30°C.

Fig. 5. A *rad55-S404A* mutation largely disrupted the physical Rtt107–Rad55 interaction,

but did not clearly affect growth during chronic exposure to DNA-damaging agents.

Experiments were performed in the W303 *RAD5* background. A) The *rad55-S404A* mutation

greatly reduced the amount of Rtt107 that co-purified with Rad55, but left Rad55–Rad57

interaction largely intact. Co-purification of Rtt107 with Rad55 did not require *RAD57.* FLAG

purification was performed on log-phase cultures. Immunoblotting was performed using anti-

FLAG, anti-VSV, and anti-HA antibodies. Anti-PGK1 antibody was applied to the input fraction

as a loading control, due to the poor visibility of FLAG-Rad55 and Rad57-HA in whole cell

extract. B) The *rad55-S404A* mutation or N-terminal 3×FLAG tag on Rad55 did not affect DNA

damage sensitivity. Higher concentrations of DNA-damaging agents were used to examine the

sensitivity of the *rad55-S404A* mutant, and the *rad55Δ* mutant was included for comparison.

Tenfold serial dilutions of overnight cultures were plated on media containing the indicated

concentrations of DNA-damaging agents, and then incubated for 2 days at 30°C.

Fig. 6. The *rad55-S404A* mutation increased spontaneous rates of LOH and mitotic RCOs.

Experiments were performed in the W303 *RAD5* background. A) The *rad55-S404A* mutation

caused a higher rate of LOH in the bimater assay, but did not exhibit epistasis with the *rtt107Δ*

mutation. Mating rates were calculated from 10–12 independent colonies as described for the

bimater assay in Figure 2B. +, non-overlap of 95% confidence intervals between two strains

(statistically significant); ns, overlap of confidence intervals (not significant). B) The *rad55-*

S404A mutation increased the frequency of mitotic RCOs, and the *rad55-S404A rtt107Δ* double

mutant phenotype was similar to that of the *rtt107Δ* mutant. Frequencies were calculated from

20–21 independent colonies. *, $p < 0.05$; **, $p < 0.01$, ***, $p < 0.005$ compared to the wild-type

(Student's *t* test). C) The *rtt107Δ* mutant exhibited multiple types of spontaneous genome

instability, and these phenotypes were not significantly affected by the *rad55-S404A* mutation.

Six phenotypic classes of colony were scored using an assay for several types of mutation and

mitotic recombination events. Rates were calculated and compared as described for the bimater

assay in Figure 2B. +, non-overlap of 95% confidence intervals between two strains (statistically

significant); ns, overlap of confidence intervals (not significant). Eight to nine independent

colonies were tested for Class 1–4 events, and five to six colonies were tested for Class 5/6

events.

Fig. 7. Rtt107 acts with different partners to prevent various types of genome instability.

The dotted lines indicate physical interactions between Rtt107 and proteins that maintain

stability in the same pathway. Numbered gray boxes indicate Rtt107 BRCT domains. A) Rtt107

prevents spontaneous LOH and mitotic RCOs by cooperating with the Rad55–Rad57

heterodimer, which binds to Rtt107–BRCT1–4. The Rad55-S404 phosphorylation site

contributes to the Rtt107–Rad55 interaction. Rtt107–BRCT5/6 contributes to the function of

Rtt107 in preventing LOH events, possibly by binding to a partner other than γH2A. B) Rtt107

prevents rDNA recombination by cooperating with Slx4, which binds to Rtt107–BRCT1–4.

Rtt107 may act as a γH2A-binding scaffold that recruits Slx4 near lesions in this context.

References

1. Ciccia A, Elledge SJ. The DNA damage response: making it safe to play with knives. Mol Cell. 2011;40: 179–204. doi:10.1016/j.molcel.2010.09.019

2. Putnam CD, Kolodner RD. Pathways and mechanisms that prevent genome instability in Saccharomyces cerevisiae. Genetics. 2017;206: 1187–1225. doi:10.1534/genetics.112.145805

3. Mehta A, Haber JE. Sources of DNA double-strand breaks and models of recombinational DNA repair. Cold Spring Harb Perspect Biol. 2014;6: 016428–016428. doi:10.1101/cshperspect.a016428

4. Emerson CH, Bertuch AA. Consider the workhorse: Nonhomologous end-joining in budding yeast. Biochemistry and Cell Biology. 2016;94: 396–406. doi:10.1139/bcb-2016-0001

5. Pizzul P, Casari E, Gnugnoli M, Rinaldi C, Corallo F, Longhese MP. The DNA damage checkpoint: A tale from budding yeast. Front Genet. 2022;13: 995163. doi:10.3389/fgene.2022.995163

6. van Attikum H, Gasser SM. The histone code at DNA breaks: a guide to repair? Nature reviews Molecular cell biology. 2005;6: 757–65. doi:10.1038/nrm1737

7. Millan-Zambrano G, Santos-Rosa H, Puddu F, Robson SC, Jackson SP, Kouzarides T. Phosphorylation of histone H4T80 triggers DNA damage checkpoint recovery. Mol Cell. 2018;72: 625-635 4. doi:10.1016/j.molcel.2018.09.023

8.   Rouse J. Esc4p, a new target of Mec1p (ATR), promotes resumption of DNA synthesis

after DNA damage. The EMBO journal. 2004;23: 1188–97. doi:10.1038/sj.emboj.7600129

9.   Leung GP, Lee L, Schmidt TI, Shirahige K, Kobor MS. Rtt107 is required for recruitment

of the SMC5/6 complex to DNA double strand breaks. Journal of Biological Chemistry.

2011;286: 26250–26257. doi:10.1074/jbc.M111.235200

10.   Leung GP, Brown JAR, Glover JNM, Kobor MS. Rtt107 BRCT domains act as a targeting

module in the DNA damage response. DNA Repair. 2016;37: 22–32.

doi:10.1016/j.dnarep.2015.10.007

11.   Wan B, Hang LE, Zhao X. Multi-BRCT scaffolds use distinct strategies to support genome

maintenance. Cell Cycle. 2016;15: 2561–2570. doi:10.1080/15384101.2016.1218102

12.   Roberts TM, Zaidi IW, Vaisica JA, Peter M, Brown GW. Regulation of Rtt107 recruitment

to stalled DNA replication forks by the Cullin Rtt101 and the Rtt109 acetyltransferase. Mol

Biol Cell. 2008;19: 171–180. doi:10.1091/mbc.e07-09-0961

13.   Balint A, Kim T, Gallo D, Cussiol JR, Bastos de Oliveira FM, Yimit A, et al. Assembly of

Slx4 signaling complexes behind DNA replication forks. The EMBO Journal. 2015;34:

2182–2197. doi:10.15252/embj.201591190

14.   Wan B, Wu J, Meng X, Lei M, Zhao X. Molecular basis for control of diverse genome

stability factors by the Multi-BRCT scaffold Rtt107. Mol Cell. 2019;75: 238-251 5.

doi:10.1016/j.molcel.2019.05.035

15. Roberts TM, Kobor MS, Bastin-Shanower SA, Ii M, Horte SA, Gin JW, et al. Slx4

Regulates DNA Damage Checkpoint-dependent Phosphorylation of the BRCT Domain

Protein Rtt107/Esc4. Molecular Biology of the Cell. 2006;17: 539–548.

doi:10.1091/mbc.e05-08-0785

16. Ohouo PY, Oliveira FM, Liu Y, Ma CJ, Smolka MB. DNA-repair scaffolds dampen

checkpoint signalling by counteracting the adaptor Rad9. Nature. 2013;493: 120–124.

doi:10.1038/nature11658

17. Brown JAR, Kobor MS. Budding yeast Rtt107 prevents checkpoint hyperactivation after

replicative stress by limiting DNA damage. DNA Repair. 2019;74: 1–16.

doi:10.1016/j.dnarep.2019.01.001

18. Leung CCY, Glover JNM. BRCT domains: easy as one, two, three. Cell Cycle. 2011;10:

2461–2470. doi:10.4161/cc.10.15.16312

19. Li X, Liu K, Li F, Wang J, Huang H, Wu J, et al. Structure of C-terminal tandem BRCT

repeats of Rtt107 protein reveals critical role in interaction with phosphorylated histone

H2A during DNA damage repair. The Journal of biological chemistry. 2012;287: 9137–46.

doi:10.1074/jbc.M111.311860

20. Chin JK, Bashkirov VI, Heyer W-D, Romesberg FE. Esc4/Rtt107 and the control of

recombination during replication. DNA Repair. 2006;5: 618–628.

doi:10.1016/j.dnarep.2006.02.005

21.   Ohouo PY, Oliveira FM, Almeida BS, Smolka MB. DNA damage signaling recruits the

874   Rtt107-Slx4 scaffolds via Dpb11 to mediate replication stress response. Mol Cell. 2010;39:

875   300–306. doi:10.1016/j.molcel.2010.06.019.

876   22.   Dibitetto D, Ferrari M, Rawal CC, Balint A, Kim T, Zhang Z. Slx4 and Rtt107 control

877   checkpoint signalling and DNA resection at double-strand breaks. Nucleic Acids Res.

878   2016;44: 669–682. doi:10.1093/nar/gkv1080

879   23.   Jablonowski CM, Cussiol J, Oberly S, Yimit A, Balint A, Kim T, et al. Termination of

880   replication stress signaling via concerted action of the Slx4 scaffold and the PP4

881   phosphatase. Genetics. 2015;201: 937–949. doi:10.1534/genetics.115.181479

882   24.   Cussiol JR, Jablonowski CM, Yimit A, Brown GW, Smolka MB. Dampening DNA damage

883   checkpoint signalling via coordinated BRCT domain interactions. The EMBO journal.

884   2015;34: e201490834. doi:10.15252/embj.201490834

885   25.   Lévesque N, Leung GP, Fok AK, Schmidt TI, Kobor MS. Loss of H3 K79 trimethylation

886   leads to suppression of Rtt107-dependent DNA damage sensitivity through the translesion

887   synthesis pathway. The Journal of biological chemistry. 2010;285: 35113–22.

888   doi:10.1074/jbc.M110.116855

889   26.   Sung P. Yeast Rad55 and Rad57 proteins form a heterodimer that functions with replication

890   protein A to promote DNA strand exchange by Rad51 recombinase. Genes and

891   Development. 1997;11: 1111–1121. doi:10.1101/gad.11.9.1111

892   27.   Liu J, Renault L, Veaute X, Fabre F, Stahlberg H, Heyer W-D. Rad51 paralogues Rad55-

893   Rad57 balance the antirecombinase Srs2 in Rad51 filament formation. Nature. 2011;479:

894   245–248. doi:10.1038/nature10522

895   28.   Sugawara N, Wang X, Haber JE. In vivo roles of Rad52, Rad54, and Rad55 proteins in

896   Rad51-mediated recombination. Mol Cell. 2003;12: 209–219. doi:10.1016/s1097-

897   2765(03)00269-7

898   29.   Roy U, Kwon Y, Marie L, Symington L, Sung P, Lisby M. Rad51 paralog complex Rad55-

899   Rad57 acts as a molecular chaperone during homologous recombination. Mol Cell.

900   2020;81: 1–15. doi:10.1016/j.molcel.2020.12.019

901   30.   Symington LS, Rothstein R, Lisby M. Mechanisms and regulation of mitotic recombination

902   in *Saccharomyces cerevisiae*. Genetics. 2014;198: 795–835.

903   doi:10.1534/genetics.114.166140

904   31.   Lovett ST, Mortimer RK. Characterization of null mutants of the RAD55 gene of

905   Saccharomyces cerevisiae: effects of temperature, osmotic strength and mating type.

906   Genetics. 1987;116: 547–553. doi:10.1093/genetics/116.4.547

907   32.   Berti M, Vindigni A. Replication stress: Getting back on track. Nature Structural and

908   Molecular Biology. 2016. pp. 103–109. doi:10.1038/nsmb.3163

909   33.   Gaines WA, Godin SK, Kabbinavar FF, Rao T, VanDemark AP, Sung P. Promotion of

910   presynaptic filament assembly by the ensemble of S. cerevisiae Rad51 paralogues with

911   Rad52. Nat Commun. 2015;6: 7834. doi:10.1038/ncomms8834

912   34.   Ball LG, Zhang K, Cobb JA, Boone C, Xiao W. The yeast Shu complex couples error-free

913   post-replication repair to homologous recombination. Molecular microbiology. 2009;73:

914   89–102. doi:10.1111/j.1365-2958.2009.06748.x

915   35.   Vanoli F, Fumasoni M, Szakal B, Maloisel L, Branzei D. Replication and recombination

916   factors contributing to recombination-dependent bypass of DNA lesions by template

917   switch. PLoS Genet. 2010;6: 1001205. doi:10.1371/journal.pgen.1001205

918   36.   Bonilla B, Brown AJ, Hengel SR, Rapchak KS, Mitchell D, Pressimone CA, et al. The Shu

919   complex prevents mutagenesis and cytotoxicity of single-strand specific alkylation lesions.

920   Aguilera A, Tyler JK, editors. eLife. 2021;10: e68080. doi:10.7554/eLife.68080

921   37.   Maloisel L, Ma E, Phipps J, Deshayes A, Mattarocci S, Marcand S, et al. Rad51 filaments

922   assembled in the absence of the complex formed by the Rad51 paralogs Rad55 and Rad57

923   are outcompeted by translesion DNA polymerases on UV-induced ssDNA gaps. PLoS

924   Genet. 2023;19: e1010639. doi:10.1371/journal.pgen.1010639

925   38.   Alvaro D, Lisby M, Rothstein R. Genome-wide analysis of Rad52 foci reveals diverse

926   mechanisms impacting recombination. PLoS Genet. 2007;3: 228.

927   doi:10.1371/journal.pgen.0030228

928   39.   Yuen KWY, Warren CD, Chen O, Kwok T, Hieter P, Spencer FA. Systematic genome

929   instability screens in yeast and their potential relevance to cancer. Proceedings of the

930   National Academy of Sciences of the United States of America. 2007;104: 3925–30.

931   doi:10.1073/pnas.0610642104

40. Hendry JA, Tan G, Ou J, Boone C, Brown GW. Leveraging DNA damage response

signaling to identify yeast genes controlling genome stability. G3 (Bethesda). 2015;5: 997–

1006. doi:10.1534/g3.115.016576

41. Putnam CD, Hayes TK, Kolodner RD. Specific pathways prevent duplication-mediated

genome rearrangements. Nature. 2009;460: 984–989. doi:10.1038/nature08217

42. Putnam CD, Allen-Soltero SR, Martinez SL, Chan JE, Hayes TK, Kolodner RD.

Bioinformatic identification of genes suppressing genome instability. Proc Natl Acad Sci U

S A. 2012;109: E3251-3259. doi:10.1073/pnas.1216733109

43. Kobayashi T. Regulation of ribosomal RNA gene copy number and its role in modulating

genome integrity and evolutionary adaptability in yeast. Cellular and Molecular Life

Sciences. 2011;68: 1395–1403. doi:10.1007/s00018-010-0613-2

44. Hang LEE, Peng J, Tan W, Szakal B, Menolfi D, Sheng Z. Rtt107 is a multi-functional

scaffold supporting replication progression with partner SUMO and ubiquitin ligases. Mol

Cell. 2015;60: 268–279. doi:10.1016/j.molcel.2015.08.023

45. Princz LN, Wild P, Bittmann J, Aguado FJ, Blanco MG, Matos J, et al. Dbf4-dependent

kinase and the Rtt107 scaffold promote Mus81-Mms4 resolvase activation during mitosis.

The EMBO Journal. 2017;36: e201694831. doi:10.15252/embj.201694831

46. Ho Y, Gruhler A, Heilbut A, Bader GD, Moore L, Adams SL. Systematic identification of

protein complexes in *Saccharomyces cerevisiae* by mass spectrometry. Nature. 2002;415:

180–183. doi:10.1038/415180a

47.   Zheng Q. rSalvador: An R package for the fluctuation experiment. G3: Genes, Genomes,

953   Genetics. 2017;7: 3849–3856. doi:10.1534/g3.117.300120

954   48.   Zheng Q. Methods for comparing mutation rates using fluctuation assay data. Mutation

955   Research - Fundamental and Molecular Mechanisms of Mutagenesis. 2015;777: 20–22.

956   doi:10.1016/j.mrfmmm.2015.04.002

957   49.   Barbera MA, Petes TD. Selection and analysis of spontaneous reciprocal mitotic cross-

958   overs in Saccharomyces cerevisiae. Proceedings of the National Academy of Sciences of

959   the United States of America. 2006;103: 12819–12824. doi:10.1073/pnas.0605778103

960   50.   Fritze CE, Verschueren K, Strich R, Easton Esposito R. Direct evidence for SIR2

961   modulation of chromatin structure in yeast rDNA. EMBO J. 1997;16: 6495–509.

962   doi:10.1093/emboj/16.21.6495

963   51.   Kobor MS, Venkatasubrahmanyam S, Meneghini MD, Gin JW, Jennings JL, Link AJ. A

964   protein complex containing the conserved Swi2/Snf2-related ATPase Swr1p deposits

965   histone variant H2A.Z into euchromatin. PLoS Biol. 2004;2: 131.

966   doi:10.1371/journal.pbio.0020131

967   52.   Downs JA, Lowndes NF, Jackson SP. A role for *Saccharomyces cerevisiae* histone H2A in

968   DNA repair. Nature. 2000;408: 1001–1004. doi:10.1038/35050000

969   53.   Pfander B, Diffley JFX. Dpb11 coordinates Mec1 kinase activation with cell cycle-

970   regulated Rad9 recruitment. The EMBO journal. 2011;30: 4897–907.

971   doi:10.1038/emboj.2011.345

54.   Torres-Rosell J, Mahín F, Farmer S, Jarmuz A, Eydmann T, Dalgaard JZ, et al. SMC5 and

SMC6 genes are required for the segregation of repetitive chromosome regions. Nature Cell

Biology. 2005;7: 412–419. doi:10.1038/ncb1239

55.   Horigome C, Bustard DE, Marcomini I, Delgoshaie N, Tsai-Pflugfelder M, Cobb JA, et al.

PolySUMOylation by Siz2 and Mms21 triggers relocation of DNA breaks to nuclear pores

through the Slx5/Slx8 STUbL. Genes & Development. 2016;30: 931–945.

doi:10.1101/gad.277665.116

56.   Elserafy M, El-Khamisy SF. Choose your yeast strain carefully: the RAD5 gene matters.

Nature Reviews Molecular Cell Biology. 2018;19: 343–344. doi:10.1038/s41580-018-

0005-2

57.   Kaliraman V, Brill S. Role of SGS1 and SLX4 in maintaining rDNA structure in

Saccharomyces cerevisiae. Curr Genet. 2002;41: 389–400. doi:10.1007/s00294-002-0319-6

58.   Zappulla DC, Maharaj ASRR, Connelly JJ, Jockusch RA, Sternglanz R. Rtt107/Esc4 binds

silent chromatin and DNA repair proteins using different BRCT motifs. BMC Mol Biol.

2006;7: 40. doi:10.1186/1471-2199-7-40

59.   Endo K, Tago YI, Daigaku Y, Yamamoto K. Error-free RAD52 pathway and error-prone

REV3 pathway determines spontaneous mutagenesis in *Saccharomyces cerevisiae*. Genes

Genet Syst. 2007;82: 35–42. doi:10.1266/ggs.82.35

60.   Hoege C, Pfander B, Moldovan G-L, Pyrowolakis G, Jentsch S. RAD6-dependent DNA

repair is linked to modification of PCNA by ubiquitin and SUMO. Nature. 2002;419: 135–

141. doi:10.1038/nature00991

61.   Boiteux S, Jinks-Robertson S. DNA repair mechanisms and the bypass of DNA damage in

994   Saccharomyces cerevisiae. Genetics. 2013;193: 1025–1064.

995   doi:10.1534/genetics.112.145219

996   62.   Janke R, Kong J, Braberg H, Cantin G, Yates K JR, N.J. Nonsense-mediated decay

997   regulates key components of homologous recombination. Nucleic Acids Res. 2016;44:

998   5218–5230. doi:10.1093/nar/gkw182

63.   Srivas R, Shen JP, Yang CC, Sun SM, Li J, Gross AM. A network of conserved synthetic

lethal interactions for exploration of precision cancer therapy. Mol Cell. 2016;63: 514–525.

doi:10.1016/j.molcel.2016.06.022

64.   Ye P, Peyser BD, Pan X, Boeke JD, Spencer FA, Bader JS. Gene function prediction from

congruent synthetic lethal interactions in yeast. Molecular systems biology. 2005;1:

2005.0026. doi:10.1038/msb4100034

65.   Tong AHY, Lesage G, Bader GD, Ding H, Xu H, Xin X. Global mapping of the yeast

genetic interaction network. Science. 2004;303: 808–813. doi:10.1126/science.1091317

66.   Kuzmin E, VanderSluis B, Wang W, Tan G, Deshpande R, Chen Y. Systematic analysis of

complex genetic interactions. Science. 360: 1729. doi:10.1126/science.aao1729.

67.   Costanzo M, Baryshnikova A, Bellay J, Kim Y, Spear ED, Sevier CS. The genetic

landscape of a cell. Science. 2010;327: 425–31. doi:10.1126/science.1180823

68.   Costanzo M, VanderSluis B, Koch EN, Baryshnikova A, Pons C, Tan G. A global genetic

1012   interaction network maps a wiring diagram of cellular function. Science. 2016;353: 549–

1013   562. doi:10.1126/science.aaf1420

1014   69.   Tong AHY, Evangelista M, Parsons AB, Xu H, Bader GD, Pagé N. Systematic genetic

1015   analysis with ordered arrays of yeast deletion mutants. Science. 2001;294: 2364–2368.

1016   doi:10.1126/science.1065810

1017   70.   Collins SR, Miller KM, Maas NL, Roguev A, Fillingham J, Chu CS, et al. Functional

1018   dissection of protein complexes involved in yeast chromosome biology using a genetic

1019   interaction map. Nature. 2007;446: 806–810. doi:10.1038/nature05649

1020   71.   Pan X, Ye P, Yuan DS, Wang X, Bader JS, Boeke JD. A DNA integrity network in the

1021   yeast Saccharomyces cerevisiae. Cell. 2006;124: 1069–1081.

1022   doi:10.1016/j.cell.2005.12.036

1023   72.   Styles EB, Founk KJ, Zamparo LA, Sing TL, Altintas D, Ribeyre C. Exploring quantitative

1024   yeast phenomics with single-cell analysis of DNA damage foci. Cell Syst. 2016;3: 264-277

1025   10. doi:10.1016/j.cels.2016.08.008

1026   73.   Leung GP, Aristizabal MJ, Krogan NJ, Kobor MS. Conditional genetic interactions of

1027   RTT107, SLX4, and HRQ1 reveal dynamic networks upon DNA damage in S. cerevisiae.

1028   G3 (Bethesda). 2014;4: 1–33. doi:10.1534/g3.114.011205

1029   74.   Hum YF, Jinks-Robertson S. DNA strand-exchange patterns associated with double-strand

1030   break-induced and spontaneous mitotic crossovers in Saccharomyces cerevisiae. PLoS

1031   Genet. 2018;14: 1–26. doi:10.1371/journal.pgen.1007302

75.   Lee PS, Greenwell PW, Dominska M, Gawel M, Hamilton M, Petes TD. A fine-structure

map of spontaneous mitotic crossovers in the yeast Saccharomyces cerevisiae. PLOS

Genetics. 2009;5: e1000410.

76.   Jinks-Robertson S, Petes TD. Mitotic recombination in yeast: what we know and what we

don't know. Current Opinion in Genetics & Development. 2021;71: 78–85.

doi:10.1016/j.gde.2021.07.002

77.   Wolner B, Van Komen S, Sung P, Peterson CL. Recruitment of the recombinational repair

machinery to a DNA double-strand break in yeast. Mol Cell. 2003;12: 221–232.

doi:10.1016/S1097-2765(03)00242-9

78.   Renkawitz J, Lademann CA, Kalocsay M, Jentsch S. Monitoring homology search during

DNA double-strand break repair in vivo. Mol Cell. 2013;50: 261–272.

doi:10.1016/j.molcel.2013.02.020

79.   Herzberg K, Bashkirov VI, Rolfsmeier M, Haghnazari E, McDonald WH, Anderson S, et

al. Phosphorylation of Rad55 on Serines 2, 8, and 14 Is Required for Efficient Homologous

Recombination in the Recovery of Stalled Replication Forks. Molecular and Cellular

Biology. 2006;26: 8396–8409. doi:10.1128/MCB.01317-06

80.   Roy U, Greene EC. The Role of the Rad55–Rad57 Complex in DNA Repair. Genes.

2021;12: 1390. doi:10.3390/genes12091390

81.   Li F, Dong J, Pan X, Oum JH, Boeke JD, Lee SE. Microarray-based genetic screen defines

SAW1, a gene required for Rad1/Rad10-dependent processing of recombination

intermediates. Mol Cell. 2008;30: 325–335. doi:10.1016/j.molcel.2008.02.028

82.   Rozenzhak S, Mejía-Ramírez E, Williams JS, Schaffer L, Hammond JA, Head S. Rad3ATR

1054   decorates critical chromosomal domains with γH2A to protect genome integrity during S-

1055   phase in fission yeast. PLoS Genet. 2010;6: 1–17. doi:10.1371/journal.pgen.1001032

1056   83.   Williams JS, Williams RS, Dovey CL, Guenther G, Tainer J a, Russell P. gammaH2A binds

1057   Brc1 to maintain genome integrity during S-phase. The EMBO journal. 2010;29: 1136–

1058   1148. doi:10.1038/emboj.2009.413

1059   84.   Javaheri A, Wysocki R, Jobin-Robitaille O, Altaf M, Côté J, Kron SJ. Yeast G1 DNA

1060   damage checkpoint regulation by H2A phosphorylation is independent of chromatin

1061   remodeling. Proceedings of the National Academy of Sciences of the United States of

1062   America. 2006;103: 13771–6. doi:10.1073/pnas.0511192103

1063   85.   Giannattasio M, Lazzaro F, Plevani P, Muzi-Falconi M. The DNA damage checkpoint

1064   response requires histone H2B ubiquitination by Rad6-Bre1 and H3 methylation by Dot1.

1065   The Journal of biological chemistry. 2005;280: 9879–86. doi:10.1074/jbc.M414453200

1066   86.   Grenon M, Costelloe T, Jimeno S, Shaughnessy AO, Fitzgerald J, Zgheib O, et al. Docking

1067   onto chromatin via the Saccharomyces cerevisiae Rad9 Tudor domain. 2007; 105–119.

1068   doi:10.1002/yea

1069   87.   Wysocki R, Javaheri A, Allard S, Sha F, Côté J, Kron SJ, et al. Role of Dot1-Dependent

1070   Histone H3 Methylation in G 1 and S Phase DNA Damage Checkpoint Functions of Rad9.

1071   2005. doi:10.1128/MCB.25.19.8430

88.   Zaidi IW, Rabut G, Poveda A, Scheel H, Malmström J, Ulrich H, et al. Rtt101 and Mms1 in

budding yeast form a CUL4 DDB1 -like ubiquitin ligase that promotes replication through

damaged DNA. EMBO Rep. 2008;9: 1034–1040. doi:10.1038/embor.2008.155

89.   Sikorski RS, Hieter P. A system of shuttle vectors and yeast host strains designed for

efficient manipulation of DNA in Saccharomyces cerevisiae. Genetics. 1989;122: 19–27.

doi:10.1093/genetics/122.1.19

90.   Moore JD, Yazgan O, Ataian Y, Krebs JE. Diverse roles for histone H2A modifications in

DNA damage response pathways in yeast. Genetics. 2007;176: 15–25.

doi:10.1534/genetics.106.063792

 Supplementary Figure Captions

 Fig. S1. Rtt107 did not limit spontaneous LOH by regulating the Rad9–Dpb11 interaction.

The *rad9-2A* allele enhanced the spontaneous LOH rate of the *rtt107Δ* mutant in the bimater

assay. Mating rates were calculated and compared as described for the bimater assay in Figure

2B. +, non-overlap of 95% confidence intervals between two strains (statistically significant); ns,

overlap of confidence intervals (not significant). This experiment was performed in the W303

rad5-535 background, and 12 independent colonies were tested.

Fig. S2. Rtt107 limited rDNA recombination in the same pathway as Slx4.

The *rtt107Δ*, *slx4Δ*, and *rtt107Δ slx4Δ* mutants exhibited similar rDNA hyperrecombination

phenotypes. rDNA recombination rates and 95% confidence intervals were determined as

described in Figure 4D. +, non-overlap of 95% confidence intervals between two strains

(statistically significant); ns, overlap of confidence intervals (not significant). This experiment

was performed in the W303 *RAD5* background, and eight independent colonies were tested.

Fig. S3. Rtt107 did not primarily limit spontaneous LOH by regulating DNA damage

tolerance pathways.

Deletion of *UBC13* did not affect the rate of LOH in the wild-type or *rtt107Δ* mutant. Mating

rates were calculated and compared as described for the bimater assay in Figure 2B. +, non-

overlap of 95% confidence intervals between two strains (statistically significant); ns, overlap of

confidence intervals (not significant). This experiment was performed in the W303 *rad5-535*

background, and 11-12 independent colonies were tested.

Fig. S4. The *rad55-S404A* mutation did not affect Rad55 protein level.

Immunoblotting was performed using anti-FLAG antibody on whole-cell extract. Anti-PGK1

antibody was used as a loading control.

Fig. 1.

Fig. 2.

Fig. 3.

Fig. 4.

A.

αFLAG IP

	WT			rad55-S404A	rad57Δ
FLAG-RAD55	+	-	+	+	+
RTT107-VSV	-	+	+	+	+
RAD57-HA	-	+	+	+	-

150 —
100 —
Rtt107-VSV (co-purified)

50 —
Rad57-HA (co-purified)

50 —
FLAG-Rad55 (captured)

Input

150 —
100 —
Rtt107-VSV

50 —
Pgk1

37 —

B.

	YPD	0.015% MMS	50 mM HU	2.5 ug/mL CPT
WT				
FLAG-RAD55				
FLAG-rad55-S404A				
rad55Δ				

Fig. 5.

Fig. 6.

A.

Rad55 Rad57

Rtt107 1 2 3 4 5 6

Suppresses

Loss of heterozygosity

Mitotic crossovers

● Rad55-S404 phosphorylation

B.

Slx4

Rtt107 1 2 3 4 5 6

Suppresses

rDNA recombination

γH2A

ssDNA

● Mec1/Tel1 phosphorylation

Fig. 7.