1 **Convergent within-host evolution alters key virulence factors in a *Klebsiella pneumoniae***
2 **clone during a large hospital outbreak**
3

4 Greta Zaborskytė[1*], Karin Hjort[1], Birgitta Lytsy[2], Linus Sandegren[1✉]
5

[1]Department of Medical Biochemistry and Microbiology, Uppsala University, Uppsala, Sweden.
[2]Department of Medical Sciences, Uppsala University, Uppsala, Sweden. Current address:
Department of Laboratory Medicine, Division of Clinical Microbiology, Karolinska Institute,
Solna, Sweden.
[*]Current address: Ineos Oxford Institute for Antimicrobial Research, Department of Biology,
University of Oxford, Oxford, UK.
12

13 [✉]Corresponding author: linus.sandegren@imbim.uu.se

15 Abstract

Bacterial pathogens adapt to host niches because of within-host selective pressures, and this
evolutionary process provides valuable insights into host-pathogen interactions. However, genetic
changes underlying adaptive phenotypes are difficult to identify from data generated by genome-
wide association studies of unrelated bacterial clones. Here, we followed the evolution of a single
Klebsiella pneumoniae clone in 110 patients during a 5-year nosocomial outbreak by combining
comparative genomics with phenotypic characterization. Strong positive within-patient selection
targeted key virulence factors in isolates from infection sites. The clone repeatedly lost acute
virulence primarily via alterations in capsule and lipopolysaccharide, changed regulation of iron
uptake, and increased biofilm formation. These phenotypes represent likely niche adaptations,
mainly to the urinary tract, and some were associated with trade-offs during gastrointestinal
colonization. The substantial convergent evolution reflects the trajectories undertaken by high-risk
clones of *K. pneumoniae* and other pathogens adapting during acute and chronic infections.

29 Introduction

Opportunistic pathogens like *Klebsiella pneumoniae* often colonize the gastrointestinal tract (GI)
or nasal mucosa without causing disease[1,2]. However, different tissue compartments (e.g., the
urinary tract, lungs, blood) can be invaded due to immune suppression, indwelling medical
devices, and antibiotic treatment[3,4]. During colonization and infection, bacterial populations
encounter numerous challenges, such as obtaining essential nutrients like iron, need for binding to
cells or abiotic surfaces, and surviving attacks from the immune system. This creates a potential
for within-host selection of mutant subpopulations that are better fit for particular host niches[5–11].
The fate of such mutants depends on their survivability in the new niche and potential trade-offs
to their normal colonization and transmission routes. While "adapt-and-live" mutations maintain
pathogen transmission and are associated with globally disseminated clones, "adapt-and-die"
changes promote survival in a specific host niche but limit bacterial spread to a new host due to

trade-offs or growth in a physical location from which spread is limited[12,13]. Evolutionary studies
on *K. pneumoniae* mainly focus on horizontally transmissible genetic traits that lead to the
emergence of high-risk clones, whereas data on transient adaptations selected on the individual
host level are scarce. However, such transient adaptive changes may affect the outcome of an
infection in the individual patient, for example, by prolonging it or making it more severe[9–11].
Often, multiple diverse clones of *K. pneumoniae* circulate in the same hospital and infect
patients simultaneously[14,15]. While large-scale genomic comparisons of such unrelated bacterial
isolates can be useful, differentiation between background allelic variation and true adaptive
changes becomes complicated, thus reducing the resolution of the within-host evolutionary
trajectories. In contrast, clonal hospital outbreaks allow for a more precise analysis due to the
common genetic origin of bacteria isolated from different patients and the defined time frame[16,17].
During 2005-2010, an outbreak of a multiresistant *K. pneumoniae* clone occurred at
Uppsala University Hospital, with 320 identified patients either colonized or infected[18–20].
Transmission of this outbreak clone occurred via direct person-to-person contact and not from
contamination of hospital surfaces[19]; therefore, genetic changes under positive selection observed
among isolates should illustrate adaptive evolution to the host rather than to the environment. Here,
we combined large-scale genomic and phenotypic analysis to understand the within-host
evolutionary trajectories of the *K. pneumoniae* clone in 110 patients during asymptomatic
colonization and infection. We find strong positive selection acting on key virulence factors
especially in isolates from infections and show how the genetic changes translate into diverse
adaptive phenotypes connected to survival in the host.
62
Results
Strong positive within-patient selection targets key virulence factors at infection sites
We determined the complete genome sequence of the index isolate and 109 representative
infecting and colonizing isolates distributed throughout the outbreak (Fig. 1a and Supplementary
Fig. 1). Single nucleotide polymorphisms, insertions, and deletions were present in 407
chromosomal genes and 91 intergenic regions with a total of 653 unique genetic changes equalling
an average rate of 3.0 mutations per genome per year (Fig. 1b and Supplementary Tables 1 and 2).
When analyzing infecting and colonizing isolates separately, the infection isolates showed a 3-fold
higher mutation rate. Phylogenetic analysis generated a tree topology dominated by individual
branches and a few clusters (A-H), each with long individual branches, indicative of selection of
genetic changes within individual patients rather than spread of dominating bacterial lineages (Fig.
1c). The ratio of nonsynonymous vs. synonymous changes in protein coding genes (dN/dS)
showed a clear positive selection signal with an overall dN/dS of 5.7 and a very strong signal
(dN/dS = 93) for the 19 genes with three or more independent mutations. There was a strong
overrepresentation of nonsynonymous mutations in the branch "tips" rather than in clusters,
supporting independent acquisition of adaptative mutations among isolates (Supplementary Fig.
2).

The majority of infecting isolates were from urinary tract infections (UTIs). While rich
sugar-based carbon sources and competition with other microbiota prevail in the intestines, the
urinary tract is typically a nutrient-poor environment dominated by amino acids, limited iron, and
fierce host defences[21]. Therefore, the adaptations selected in the gut versus urinary tract would
likely display a functional bias. Independently selected genetic changes in cell envelope, iron
uptake and metabolism, and Fe-S cluster synthesis were significantly overrepresented among
infection isolates (Fig. 1d). This bias was even more pronounced on an individual gene level when
considering genes and intergenic regions (n=39) with at least two mutations acquired in parallel in
different isolates (two-tailed binomial test, p<0.0001). Adaptation to specific niches often requires
multiple mutations that produce a combination of altered traits[22]. Here, we find that genetic
changes in the most frequently mutated genes often co-occurred with up to eight top genetic targets
mutated in the same isolate (Fig. 1e, Supplementary Fig. 3 and Supplementary Table 3). This co-
occurrence was also overrepresented in infection isolates, mainly from UTIs, further supporting
strong convergent niche-specific evolution. Some convergent mutations were also acquired at
different branching points within the same phylogenetic cluster, e.g., *wcoZ* (capsule
acetyltransferase), *uvrY* (transcriptional regulator in UvrY-BarA), *envR* (efflux pump regulator),
suf (Fe-S cluster synthesis), and intergenic *fepA/fes* (siderophore uptake) (Fig. 1f). Collectively,
comparative genomics shows that changes in key virulence-associated processes were repeatedly
selected in individual patients in parallel and preferentially during infection.
99
Frequent attenuation in acute virulence and increased susceptibility to complement due to
repeatedly selected surface polysaccharide mutations
Considering the strong positive selection acting on vital virulence factors in infecting isolates, we
screened all 110 isolates for their potential to kill *Galleria mellonella* wax moth larvae. *G.*
mellonella has an innate immune system with both cellular and humoral responses similar to those
of vertebrates[23,24] and is the preferred infection model for opportunistic *K. pneumoniae,* which do
not perform well in mice models[25–27]. Based on the rate of *G. mellonella* killing and the health
status of the surviving larvae, the isolates were divided into five groups (Fig. 2a). Half of the
isolates showed complete or partial attenuation, while an increased virulence was seen for five
isolates not associated phylogenetically. One group of isolates (phylogenetic cluster B) displayed
a different infection progression, where injected larvae showed no signs of sickness up to 12 h
post-injection but died rapidly within 24 h. The attenuated group was overrepresented among
infection site isolates (Fig. 2b), strongly indicating a switch to a more chronic form of infection as
a common adaptation.
In some cases, it was easy to pinpoint the single mutations responsible for acute virulence loss
by comparison of neighbors in phylogenetic clusters (Fig. 2c). For most isolates, though, there was
more than one genetic difference between isolates with different virulence profiles, complicating
the distinction between virulence-altering changes and co-selection. There was minimal overlap
between genes and intergenic regions mutated in different virulence groups, with most unique hits
in the isolates with impaired acute virulence (Fig. 2d). These unique mutations were associated

with diverse functions, but parallel cell envelope mutations were the dominant ones, particularly
in surface polysaccharide synthesis regions (*cps* and O-antigen) (Fig. 2e, f and Supplementary
Table 4). Often, combinations of the same mutational targets were acquired in single isolates, even
with identical combinations (e.g., *wzc+manB+ompK36*). No unifying mutations were present in
the increased virulence group, reflecting diverging rather than converging genotypes associated
with a gain of function (Supplementary Table 4).
Capsule typically shields *K. pneumoniae* from complement attacks, which is an essential innate
immune response to protect against infection[28]. Therefore, a non-mucoid morphotype (capsule
loss) has been associated with impaired complement evasion, while increased mucoviscosity has
been linked to complement resistance and more invasive infections[29]. Surprisingly, here the most
hypermucoid isolates were attenuated and as sensitive to human serum as the isolates that had lost
the capsule, with up to >5 log reduction in CFU/ml within 3h (Fig. 2f). All such hypermucoid
isolates carried missense mutations in *wzc,* encoding an essential capsule co-polymerase and
regulator[30,31]. The exception was two clustering isolates with the *wzc*[Y714C] mutation. Increased
mucoviscosity due to other capsule changes, e.g., a 21-nt deletion in the capsule regulator *yrfF*,
did not reduce resistance to complement. All *wzc* mutants were isolated from infection sites,
suggesting a unique adaptation.
In most cases, reduced killing of *G. mellonella* was coupled with decreased complement
evasion (Fig. 2f). However, increased complement resistance via the O-antigen mutation
manC[Q411*] and an intergenic capsule mutation associated with *galF/cpsACP* also resulted in slower
killing of *G. mellonella*. This finding suggests that certain modifications of surface
polysaccharides could help avoid detection, thereby delaying the host response. Within some
phylogenetic clusters, sensitivity to complement via capsule loss was acquired subsequent to
attenuation in *G. mellonella* in colonizing isolates (Supplementary Fig. 4). Moreover, some
attenuated isolates had non-altered survival in serum, which suggests other factors than different
interaction with complement as the basis for virulence loss. One such contributing factor could be
a general growth defect[25], although we did not see any correlation between killing of *G. mellonella*
and the exponential growth rates *in vitro* (Supplementary Fig. 5).

Intergenic *fepA/fes* changes alter the regulation of enterobactin uptake and utilization
Iron sequestration is essential for the growth and survival of *K. pneumoniae* inside the host[32–34].
Multiple siderophore-dependent and independent systems are responsible for this process (Fig.
3a), and all of them were mutational targets among the isolates, e.g., inactivation of *fhuA*,
nonsynonymous mutations in *fepA*, gene amplifications of *feoABC*, and intergenic changes in the
fepA/fes regulon (Supplementary Table 2). Moreover, they were often co-selected with mutations
in *suf* genes, especially *sufB* and *sufC*, responsible for Fe-S cluster synthesis (Supplementary Fig.
3). We measured how well all isolates grew under iron-depleted and repleted conditions with either
FeSO$_4$ or inactivated human serum as a source of iron. While the overall growth rates and yields
were decreased in the absence of iron, we did not observe any apparent connection between
mutations and changes in iron-dependent growth (Supplementary Table 5). Therefore, seemingly

loss-of-function changes did not result in defects during iron-dependent growth, possibly because
partially redundant systems complement measurable deficits.
The *fepA/fes* region was the most common independently targeted intergenic region (Fig.
1d, f), but also intergenic regions upstream of *fepB* and *feoA* were mutated, mostly in infecting
isolates. The *fepA/fes* intergenic region contains overlapping bidirectional promoters and a Fur
(ferric uptake regulator) box that regulates the expression of the FepA enterobactin transporter and
the Fes enterobactin esterase (Fig. 3b). As expected, the index isolate regulated the expression of
both genes depending on iron levels with high expression in chelated M9 medium, low expression
in chelated M9 supplemented with $FeSO_4$, and an in-between expression in medium supplemented
with serum where most iron is bound to proteins (Fig. 3c). Different mutations in the region right
upstream of *fepA* or around and inside the Fur box either increased *fepA* expression or eliminated
the responsiveness to iron availability such that the expression of both *fepA* and *fes* remained
elevated even in high-iron conditions, most likely through disruption of the Fur-mediated
regulation (Fig. 3b). Taken together, these results illustrate that highly specific phenotypic
responses mediated by intergenic mutations associated with iron acquisition were selected during
infection.
176
Increase in biofilm formation on catheter-like surfaces were selected via mutations affecting
c-di-GMP turnover, fimbriae, and polysaccharide synthesis
Most of the infection isolates were from UTIs, where urinary catheterization promotes biofilm
growth of uropathogens, including *K. pneumoniae*[35–37]. Therefore, we tested how well the isolates
formed biofilms on silicone-coated surfaces with and without fibrinogen using the FlexiPeg
device[38]. Fibrinogen is commonly deposited on urinary catheters during catheter-induced
inflammation[39] and is required for attachment by several uropathogens[40–42]. Thirteen isolates (nine
urine/wound) had elevated levels of biofilm formation and were distributed throughout the
phylogeny; seven of them needed fibrinogen coating to display the phenotype fully (Fig. 4a).
The mutations responsible for an increased biofilm formation were in genes affecting
fimbrial adhesins, c-di-GMP signaling proteins, and polysaccharides, including capsule regulation
(Fig. 4a). Adaptive changes in the FimH tip adhesin were previously reported in *Escherichia coli*
and *K. pneumoniae* isolates from UTIs[43–46], while YfiN and YfiR were targeted in *Pseudomonas*
aeruginosa adapting during chronic lung infections[47]. The wound isolate DA25066 with extremely
high biofilm biomass had a unique 21 nt in-frame deletion in *yrfF* that encodes a negative regulator
for the Rcs two-component system, which induces capsule production[48]. Previously, a 12-nt
deletion in *yrfF* has been reported in *E. coli* isolated from keratitis[49] and selected during serial
passaging in macrophages[50]. The isolate DA69557 had multiple mutations with possible
connection to an extreme biofilm capacity previously observed by SEM[38]. One such change was
the intergenic *pgaA/csgD* mutation (poly-N-acetylglucosamine PNAG exporter/master biofilm
regulator) that was acquired in parallel both in DA69557 and part of cluster B. Moreover, at least
four different genes associated with c-di-GMP turnover, including *yfiN*, were mutated
(Supplementary Table 1).

There was no direct relationship between acute virulence features (*G. mellonella*, serum, mucoviscosity) and biofilm capacity in these isolates (Fig. 4b). In most cases, the increased biofilm capacity had evolved as the end-point adaptation; however, two clusters showed reversion of this phenotype (Fig. 4c). The blood isolate DA25061, found 1.5 months later than the phylogenetically closely related wound isolate DA25060, lost the increased biofilm phenotype due to an inactivating mutation in the transcriptional activator MrkH. Similarly, a mutation in the essential PNAG synthesis protein PgaC reverted the biofilm phenotype back to the index level in part of cluster B that otherwise showed an increase due to the intergenic *pgaA/csgD* mutation. Such observations reflect likely differences in biofilm-related fitness at different body sites (e.g., wound vs. bloodstream), but also how individual host environments shape bacterial evolution even in the same niche (e.g., urinary tract). Differences in the immune response have been suggested to affect the adaptive trajectories during asymptomatic bladder colonization with *E. coli*[6].

Increased biofilm capacity gives an advantage during antibiotic-facilitated GI colonization in mice

Since intestinal colonization is usually the dominant reservoir of spread to new hosts and also precedes infection by classical *K. pneumoniae* strains[2,51,52], we tested how a set of isolates with different evolved phenotypes compete with the index isolate when colonizing the GI tract of mice. Colonization with the natural gut microflora present was not stable in any of the competition groups, indicating that, at least under these conditions in mice, the outbreak clone is not a super colonizer (Fig. 5a). When the drinking water was supplemented with clindamycin, which causes significant disruptions to the gut microbiota and enables long-term colonization with *K. pneumoniae*[53], the bacterial load dramatically increased above 10^9 CFU/g faeces and stayed so until the end of the experiment. These results signify the role of antibiotic treatment-induced alterations to the gut microbiota in colonization with the outbreak clone, supporting its efficient spread in the hospital.

All three biofilm-forming isolates had a selective advantage over the index isolate during GI colonization. This was especially prominent for DA25066, with the most substantial biofilm formation capacity conferred by a deletion in the capsule regulator *yrfF* (Fig. 5b). Two of the biofilm-forming isolates also carried a mutation in *sbmA*, encoding the uptake protein for microcins, suggesting possible alterations in the defense against other gut bacteria. The colonizing isolate DA25116 stayed neutral, and the hypermucoid *wzc* mutant DA25071 was not as good at colonizing GI as the index isolate. Therefore, specific phenotypes (e.g., biofilm formation) selected during within-host evolution favor GI colonization, while others (e.g., *wzc*-mediated hypermucoidy) do not. This observation reflects how, given the opportunity, mutants with an advantage at infection sites could be pre-selected already during gut colonization and have an advantage in both body sites. In contrast, a hypermucoid *wzc* mutant isolated from a UTI represents a case where the mutant had adapted to the infection site so well that it no longer can colonize the gut to the same extent ("adapt and die").

Discussion

Discussion
Understanding within-host evolutionary dynamics of bacterial populations is central to infection
biology. Our study reconstructs key genomic and phenotypic events in 110 isolates of a single *K.*
pneumoniae clone during transmission between patients and adaptation to host niches. The
adaptive changes we identify as selected mainly in infecting isolates overlap with those linked to
high-risk *K. pneumoniae* clones[54–56], uropathogenic *E. coli*[57], and chronic infections by CF lung
pathogens, like *Burkholderia dolosa*[58], *B. cenocepacia*[59], and *P. aeruginosa*[47]. While genomic
analysis alone can be a powerful predictor of adaptive evolution, we show unexpected coupling to
phenotypes (e.g., *wzc*, *manB/manC*, *fhuA*) in quite a few cases. These findings highlight the
importance of connecting genotype-phenotype when assessing complex selection in the host[60,61]
even in a single clone.
We show that reduced evasion of innate immunity mechanisms (complement) and
attenuation in acute virulence was a much more common evolutionary direction than increased
virulence. Most attenuated isolates were from infection sites, suggesting that they are still
maintained rather than cleared, possibly by avoiding detection by the host. Moreover, acquisition
of parallel mutations and similar mutation-combinations strongly implies convergent adaptation
to a more chronic state, like for other opportunistic pathogens[62]. How exactly the selected
mutations mediate such a switch mechanistically in *K. pneumoniae* is a key question for future
research. The assessment of increased killing of *G. mellonella* is more difficult as the infection
progresses fast in the index isolate even at a relatively low infectious dose; therefore, there is a
possibility that we underestimate the potential for increased virulence.
Virulence changes were often connected to capsular changes. One of the most striking
cases was that *wzc*-mediated hypermucoviscosity sensitized the clone to complement and
attenuated rather than increased virulence. In contrast, a recent study associated *wzc* mutants in the
K. pneumoniae ST258 global high-risk lineage with increased macrophage resistance and higher
virulence in zebrafish larvae and murine UTIs[54]. The same authors also suggested that such
mutations are associated with blood infections, while capsule-loss mutants are associated with
UTIs. However, here we detect both morphotypes in UTIs, which could reflect differences in strain
background. Supporting the connection to UTIs, another recent study showed how urine modulates
the capsule and maintains the hypermucoid phenotype of *K. pneumoniae wzc* mutants[63].
Many seemingly loss-of-function LPS and capsule mutations altered virulence without
changing the morphotype. The most prominent case was frequent frameshifts and deletions in
manB/C, encoding the ManB phosphomannomutase and the ManC mannose-1-phosphate
guanylyltransferase. Inactivation of *manB/C* abolishes capsule production in other *K. pneumoniae*
strains[64]. Since KL51 capsule lacks mannose, in this clone, the mutations in *manB/C* probably
affect the synthesis of O3b antigen that contains three mannose residues[65]. Loss of O-antigen due
to its immunogenicity has been reported during late stages of chronic CF lung infections[66].
Inactivating *manC* mutations have specifically been connected to adaptations in *B. dolosa* and
experimentally biofilm-evolved *B. cenocepacia*[59]. Similarly, acetylation of the capsule in *K.*
pneumoniae has been associated not only with a change in immunogenicity, but also altered

adhesion to surfaces[67], and here, inactivating *wcoZ* (capsule acetyltransferase) mutations were a
common cause of virulence loss. Collectively, *manB/C* and *wcoZ* mutations could reflect selection
connected to the catheterized bladder environment.

Iron scavenging has been identified as a selective trait in bacterial pathogens, such as *B.*
dolosa[59] and *E. coli*[68]. In accordance with our observations, adaptive changes in siderophore
uptake rather than synthesis seem to be more critical in the context of infection[69]. Changes in iron
uptake in infecting site isolates were often co-selected with mutations in *suf* genes, encoding
components for the Fe-S cluster assembly complex, induced under oxidative stress and iron
starvation[70]. Genetic variation in *suf* genes was also found to be positively selected in the global
K. pneumoniae outbreak clones ST258[55] and ST147[56], and *sufB* mutations were connected with
increased iron binding[55]. Notably, mutations that could be predicted to have adverse effects on
iron acquisition did not reduce growth in low-iron conditions. This may reflect redundancy in how
K. pneumoniae obtain and use iron, as shown previously for ST258, where mutation in the
conserved *entS* (enterobactin export) led to higher expression of an alternative iron transport
system (hemin transport)[33,34]. Alternatively, seemingly detrimental loss in enterobactin synthesis
could also be attributed to the evasion of lipocalin 2, a major innate immunity protein that
specifically binds enterobactin[34]. FhuA and FepA are also receptors for phages and
bacteriocins[71,72]; therefore, loss of these proteins could be selected as protection from these factors.
In agreement with this hypothesis, one of the isolates with a *fhuA* mutation also had an inactivating
mutation in *sbmA*, encoding the transporter for microcin-type bacteriocins.

Although the dN/dS ratio is typically considered a strong indicator of positive selection[73],
our results illustrate that intergenic changes should not be neglected. We find that the intergenic
region *fepA/fes* is a prevalent mutational target, especially among UTI isolates, and is associated
with significant alterations in the regulation of enterobactin uptake and responsiveness to iron
levels. Intergenic regions related to iron acquisition were also found to be under positive selection
in ST410 carbapenem-resistant *E. coli*[68]. A study on the adaptation of *P. aeruginosa* showed that
parallel intergenic evolution predominantly targeted promoters and suggested that intergenic
mutations facilitate the evolution of essential genes by minimizing systemic detrimental effects[74].

In conclusion, we show at an unprecedented level how a clone, representing a critically
important nosocomial pathogen, underwent extensive parallel evolution during a large hospital
outbreak. We highlight the power of combining large-scale genomic analysis with phenotyping
and hope that this dataset can continue to serve as a valuable resource to expand our understanding
of the evolutionary within-host trajectories of *K. pneumoniae*.

Materials and Methods

Bacterial isolates and sampling
The first patient confirmed with the outbreak clone was identified in December 2004 and had been
treated abroad (Tehran, Iran) before being admitted to the oncology ward at Uppsala University
Hospital[18]. The isolates from the outbreak were collected from routine samples at Uppsala

University Hospital. Isolates (n=110, each from a different patient and from the first occasion it
was isolated) were randomly included in this study to cover the entire outbreak duration and to
represent the clinical and screening isolates equally. All isolates were included at times with low
incidence, and a proportion of isolates were randomly chosen from time points with high
incidence. In total, 320 patients were identified with the clone, but unfortunately, all isolates except
the ones gathered for this study were lost due to a freezer accident at the hospital, making it impos-
sible to include additional isolates in the study in retrospect. 45% (n=50) of the patients were male
and 65% (n=60) were female. The median age was 80 years (range 23–100). Clinical isolates
(n=68) were collected from urine (n=55), wounds (n=8), blood (n=4), and lungs (n=1). Colonizing
isolates (n=42) were collected from fecal samples as part of the screening for carriage of the
outbreak clone. The summary of all isolates and associated genomic and experimental data is
available as an interactive Microreact project
(https://microreact.org/project/6bg86f2tgddpz4q5xavmgh).
333
Whole-genome sequencing of outbreak isolates
The index isolate DA14734 was sequenced with long-read whole genome sequencing using the
Pacific Biosciences platform (PacBio, Menlo Park, CA). DNA was prepared from overnight
cultures using the Qiagen Genomic-Tip 100, according to the manufacturer's instructions. PacBio
sequencing was performed at Uppsala Science for Life Laboratory on an RSII system with one
SMRT cell per genome. In addition, the DA14734 genome was also sequenced with short-reads
using MiSeq (Illumina, San Diego, CA) with two times 300-bp paired-end read lengths using the
Nextera XT DNA library preparation kit (Illumina, San Diego, CA), according to the manufac-
turer´s instructions. From the index isolate, a reference genome sequence was assembled with a
combination of the PacBio reads and MiSeq reads. Assembly was performed with the CLC
Genomics workbench v.20 (CLC Bio, Qiagen), including the Microbial Genomics Pro Suite
module. The additional 109 isolates from the outbreak were also whole-genome sequenced with
the Illumina technology, and the sequences were compared to the reference genome by reference
assembly using the CLC Genomics Workbench. Variant detection was performed using CLC
Genomics in-built detection modules for single nucleotide polymorphisms (SNPs), inser-
tions/deletions, and structural variations. Variations in the chromosomal sequences were used to
construct a Maximum likelihood phylogenetic tree using the CLC Microbial Genomics Module
Epidemiology function. An initial Maximum Likelihood Phylogeny was created with a Neighbor
Joining General time reversible nucleotide substitution model. This initial tree was then manually
updated based on the exact genetic changes for each isolate with branch lengths adjusted to the
exact number of genetic changes compared to the index isolate. All sequences have been deposited
at NCBI under bio project number PRJNA857654.
356
Bioinformatic analysis of genetic changes
The mutational rate throughout the outbreak was calculated by linear regression of the number of
mutations per isolate compared to the index isolate and the respective isolate's isolation date. The

dN/dS ratio was calculated for groups of mutated genes with the same number of independent
mutations by dividing the total number of non-synonymous changes by the total number of
synonymous changes. Identification of biological functions and grouping of genes was done by a
combination of manual BLASTp[75], STRING[76], and EggNog 5.0[77] analysis combined with
literature searches. The genetic basis for phenotypic changes was determined by first classifying
all mutations as cluster or tip mutations, and then comparing tip mutations in clustering isolates
with phenotypic differences. In cases with more than one possible mutation, we focused on
repeatedly mutated genes in isolates with the same phenotype.
368
Exponential growth rate measurements in MH II and urine
Overnight (O/N) cultures were diluted 1000-fold in MH II broth, and 300 µl were transferred to
100-well honeycomb plates. Five biological replicates were used for each isolate. The plates were
incubated in a BioScreen C (Oy Growth Curves Ad Ltd) for 16 h at 37°C with shaking with OD_{600}
measurements taken every 4 min. For experiments with urine, morning urine was collected from
a healthy female volunteer on six different days. The urine was centrifuged (3800xg, 10 min),
sterile filtered (0.2 µm pore size), and aliquots were frozen at -20°C. For experiments, frozen
aliquots from different days were thawed, centrifuged again (3800xg, 10 min), and pooled
together. Exponential growth rates were analyzed using the R script-based tool BAT 2.0[78].
378
Biofilm formation
The capacity to form surface-attached biofilms was determined on the FlexiPeg biofilm device[38]
with silicone-coated pegs. To further coat the pegs with fibrinogen, the autoclaved peg lid was
incubated in a flat-bottom 96-well plate with 200 µl of 100 µg/ml fibrinogen from human plasma
(product number F3879, Sigma-Aldrich, in 0.9% NaCl or PBS) at 4°C overnight. Approximately
10^5 CFU in a total volume of 150 µl were added to a 96-well plate, where the autoclaved peg lid
was inserted, and biofilms were allowed to form statically in BHI for 48 h at 37°C with a medium
change after 24 h. To quantify the total biomass, the pegs with biofilms were stained using 0.1 %
crystal violet (CV) (Sigma-Aldrich), and the absorbance (540 nm) of the dissolved stain was
measured as described before[38].
389
Bacterial survival in human serum
O/N cultures were grown in LB and diluted 100-fold in PBS. A final dilution of ~10^5 CFU was
mixed with human serum (Sigma-Aldrich, H6914 from male AB clotted whole blood, USA origin,
sterile-filtered) in a total volume of 200 µl in a 96-well plate. Due to the natural batch variation of
human serum, serum was diluted up to 40 % when needed to achieve a similar killing effect relative
to the index and sensitive control isolates. For negative controls, serum was heat-inactivated for
30 min at 56°C. To confirm that the killing effect was due to the complement system, a control
with the complement inhibitor polyanetholesulfonic acid (Sigma-Aldrich) was done at a final
concentration of 100 µg/ml (Supplementary Fig. 5). CFU counts were performed at timepoint 0h
and after 3h static incubation at 37°C.

400
Virulence testing in *Galleria mellonella* larvae model
Research-grade wax moth *G. mellonella* larvae TruLarv™ were purchased from BioSystems
Technology Ltd (UK). O/N cultures (BHI) were diluted in PBS, and 10 µl containing 10^4 CFU
were injected via the first right proleg using a 25 µl Hamilton 7000 syringe (Model 702 RN, 22S
gauge needle). A dose of 10^4 CFU was chosen because it led to the most consistent results between
biological replicates and the best resolution between different virulence levels. For each strain,
two biological replicates (cultures started from independent colonies) were injected into 10 larvae
each. For each experiment, 10 larvae were injected with 10 µl sterile PBS to account for possible
injection trauma, and 10 larvae were used as non-injected controls. Aliquots from dilutions were
plated for CFU counts to confirm the correct infection load. Larvae were incubated at 37°C for 48
h with an hourly assessment of death during 7-12 h post-injection, followed by monitoring after
24 h, 32 h, and 48 h. Larvae were considered dead when not moving after stimulation with a sterile
pipette tip, which usually coincided with melanization. At the end of the experiment, live larvae
were given health index scores[79], and a cumulative score (0-90) was determined by adding up
scores of individual larvae for each replicate. Survival analysis was performed using the Kaplan-
Meier method and analyzed by log-rank (Mantel-Cox) test in GraphPad Prism version 9.2.0 for
macOS (GraphPad Software, San Diego, CA, USA). All isolates were categorized relative to the
index isolate DA14734.

Construction of a genetically marked index isolate for mice experiments
The index isolate DA14734 was transformed with the pSIM10-hygro λ Red recombineering
system. The aminoglycoside resistance gene *armA* was PCR amplified using Phusion High-
Fidelity DNA Polymerase (Thermo Fisher Scientific Inc.) with primers introducing a 40 bp
homology region directly outside *galK* for in-frame replacement (primers: 14734_armA_galK_F1
GCGCCGTCAGCGACGTCCATTTTCGTGAATCCGGAGTGTAAGTAAAGTGTGGTGATG
TTGA;14734_armA_galK_R1_ATCGTCCGGGGCCAGCGCGGTGGTTTGCGTTAGCATTG
TTGTCGTTTGTGAGCT. Purified *armA(galK)* DNA was electroporated into electrocompetent
DA14734 pSIM5-hygro λ Red cells washed with 10% glycerol that had the λ Red system induced
by 15 min incubation at 42°C. Successful transformants carrying *galK::armA* were selected on LA
+ gentamicin (50 mg/L) at 30°C.

Competitions during gastrointestinal colonization in mice
To assess the competitive advantage during gastrointestinal colonization in mice, the marked index
isolate DA73939 (*galK::armA*) was competed with the unmarked selected isolates. Testing all
isolates in mice was not feasible, therefore, we chose five isolates DA14742, DA25061, DA25066,
DA25071, and DA25116, representing different isolation sites, capsule status, biofilm formation
capacity, virulence in *G. mellonella* larvae, and survival in human serum. BALB/c female mice
(n=36) were acclimatized for eight days before starting the experiment. One mouse injured itself
during this period and had to be sacrificed by cervical dislocation. As a result, one competition

group (control DA14734 vs DA73939) had five instead of six mice. Overnight cultures of pair-
wise isolates were mixed 1:1 in PBS/1% bicarbonate, and 100 µl of the mixture (approximately
3×10^7 CFU) were administered via oral gavage. The procedure was repeated on two consecutive
days (at least 24 h apart). On Day 5, a third gavage with fresh bacterial cultures was given. After
that, drinking water with clindamycin (Villerton) at a final concentration of 0.5 g/L and 5 g/L
glucose was provided *ad libitum* to disrupt the endogenous microflora and facilitate colonization.
Feces from each mouse were collected every day (Day 1-Day 10), weighed, homogenized in 1 ml
sterile PBS, vortexed at full speed for 2 min, and briefly (max 5 s) spun down to remove larger
debris. Different dilutions were plated on LB agar with cefotaxime (10 mg/L) to allow growth of
all isolates and LB agar with gentamicin (50 mg/L) to select for the marked index isolate. The
number of bacteria was expressed as CFU per gram of feces. On Day 10, all mice were sacrificed
by cervical dislocation. To test for systemic spread of bacteria, spleens were extracted,
homogenized in 1 ml PBS, vortexed at full speed for 2 min, briefly spun (5 s), and plated on LB
agar with cefotaxime (10 mg/L) or gentamicin (50 mg/L). Colonized mice showed no signs of
disease or systemic infection at the end of the competitions. The competitive index was calculated
each day by dividing the output ratio in feces by the input ratio (inoculum cultures) for each mouse.
The final data was normalized to the values from the control competition to account for any
possible selective differences conferred by *galK*::*armA* and expressed as a $\log_{10}$ competitive index.
All experiments with the animals were done in accordance with the national regulations and
approved by the Regional Ethics Review Board in Uppsala (Dnr 5.8.18-15552/2019).
460
Growth under iron-depleted and repleted conditions
Iron-dependent growth in all isolates was assessed similarly as described before[34]. The iron-
depleted medium was made by treating M9 with Chelex 100 (50-100 dry mesh) resin (BioRad) for
1 h, then filter-sterilized and supplemented with 0.2% glucose, 1 mM $MgSO_4$ and 0.3 mM $CaCl_2$.
Overnight cultures were grown in BHI and diluted to 10^3 CFU/ml in M9/Chelex, M9/Chelex
supplemented with 100 µM $FeSO_4$, and M9/Chelex with 5% heat-inactivated human serum. The
growth was monitored in a BioScreen C (Oy Growth Curves Ad Ltd) for 18 h at 37°C with shaking
and OD_{600} measurements taken every 4 min. The index isolate was monitored along with the
experimental isolates for each run. The area under the curve (AUC) was calculated for each isolate
and each replicate, and then normalized to the index for each condition. AUC of each isolate was
compared to the index isolate in the respective conditions by Brown-Forsythe and Welch ANOVA
followed by Dunnett's T3 multiple comparison test.
473
RNA extraction and qPCR of *fepA* and *fes*
Overnight cultures of selected isolates were grown in BHI and diluted to 10^4 CFU/ml in
M9/Chelex, M9/Chelex supplemented with 100 µM $FeSO_4$, and M9/Chelex with 5% heat-
inactivated human serum. The cultures were grown until the mid-exponential phase, spun down,
and mixed with RNA Protect (Qiagen). RNA was extracted using the RNeasy Mini Kit (Qiagen),
and genomic DNA was removed using the TURBO DNA-free kit (Invitrogen). Complementary

DNA (cDNA) was synthesized using the High-Capacity Reverse Transcriptase kit (Applied biosystems) with approximately 500 ng of extracted RNA. The mRNA levels of *fepA* and *fes* were quantified by qPCR (Illumina Eco Real-Time PCR System), using the SYBR Green Master mix (Thermo Fisher) with the following primers: fepA_F1 CAGAACACCAACTCCAACGA; fepA_R1 CCGTTCCAGGTAACCGAGTA; fes_F1 AATCGCTGCAACGTCTCC, fes_R2 TCACGGTCGGAGGGAATA. The mRNA levels of the housekeeping gene *glnA* were assessed using primers glnA_F and glnA_R[80]. The mRNA levels were normalized to the index mRNA levels for the respective conditions using the $2^{-\Delta\Delta C_{\mathrm{T}}}$ method[81].

Sedimentation assay

Hypermucoviscosity was assessed by low-speed centrifugation as described before[64]. Overnight cultures grown in BHI (5 ml, in 50 ml Falcon tubes) were centrifuged for 5 min at 1000x g, and OD_{600} values were measured, before and after centrifugation, for the top 300 µl of supernatants. Hypermucoviscosity was expressed as the ratio between the OD_{600} of the supernatant and the OD_{600} of the corresponding culture before centrifugation.

Statistical analysis

All statistical analysis was done using GraphPad Prism version 9.2.0. for macOS (GraphPad Software, San Diego, California, USA). The respective analysis is specified for each experiment and figure in the figure legends.

Acknowledgments

We thank Evelina Kess for handling the work with mice during the colonization experiments, and the staff at the clinical microbiology lab at Uppsala University Hospital for help with collection of clinical isolates. We also thank Dan I. Andersson and Vaughn S. Cooper for comments on the manuscript draft, and Lionel Guy for help with the phylogenetic analysis.

Author contributions

G.Z. and L.S. conceived the study and designed experiments. G.Z. performed all phenotypic characterization experiments. K.H. performed whole-genome sequencing. G.Z., K.H., and L.S. performed bioinformatic analysis. B.L. provided clinical isolates and patient data. G.Z. and L.S. analyzed data and wrote the manuscript. All authors read, edited, and approved the final version of the manuscript.

References

1. Podschun, R. & Ullmann, U. Klebsiella spp. as nosocomial pathogens: Epidemiology, taxonomy, typing methods, and pathogenicity factors. *Clinical Microbiology Reviews* **11**, 589–603 (1998).

2. Gorrie, C. L. *et al.* Gastrointestinal Carriage Is a Major Reservoir of Klebsiella pneumoniae Infection in Intensive Care Patients. *Clinical Infectious Diseases* **65**, 208–215 (2017).

3. Bengoechea, J. A. & Sa Pessoa, J. Klebsiella pneumoniae infection biology: Living to
counteract host defences. *FEMS Microbiology Reviews* **43**, 123–144 (2019).

4. Gorrie, C. L. *et al.* Genomic dissection of Klebsiella pneumoniae infections in hospital
patients reveals insights into an opportunistic pathogen. *Nature Communications 2022 13:1*
13, 1–17 (2022).

5. Lieberman, T. D. *et al.* Parallel bacterial evolution within multiple patients identifies
candidate pathogenicity genes. *Nature Genetics* **43**, 1275–1280 (2011).

6. Zdziarski, J. *et al.* Host imprints on bacterial genomes-rapid, divergent evolution in
individual patients. *PLoS Pathogens* **6**, 95–96 (2010).

7. Yang, L. *et al.* Evolutionary dynamics of bacteria in a human host environment. *Proceedings*
of the National Academy of Sciences **108**, 7481–7486 (2011).

8. Jousset, A. B. *et al.* A 4.5-Year Within-Patient Evolution of a Colistin-Resistant Klebsiella
pneumoniae Carbapenemase–Producing K. pneumoniae Sequence Type 258. *Clinical*
Infectious Diseases **67**, 1388–1394 (2018).

9. Yang, Y. *et al.* Within-host evolution of a gut pathobiont facilitates liver translocation.
Nature 2022 607:7919 **607**, 563–570 (2022).

10. Lysenko, E. S., Lijek, R. S., Brown, S. P. & Weiser, J. N. Within-Host Competition Drives
Selection for the Capsule Virulence Determinant of Streptococcus pneumoniae. *Current*
Biology **20**, 1222 (2010).

11. Young, B. C. *et al.* Severe infections emerge from commensal bacteria by adaptive
evolution. *eLife* **6**, e30637 (2017).

12. Sokurenko, E. V., Gomulkiewicz, R. & Dykhuizen, D. E. Source–sink dynamics of virulence
evolution. *Nature Reviews Microbiology 2006 4:7* **4**, 548–555 (2006).

13. Culyba, M. J. & Tyne, D. V. Bacterial evolution during human infection: Adapt and live or
adapt and die. *PLOS Pathogens* **17**, e1009872 (2021).

14. Holt, K. E. *et al.* Genomic analysis of diversity, population structure, virulence, and
antimicrobial resistance in Klebsiella pneumoniae, an urgent threat to public health.
Proceedings of the National Academy of Sciences of the United States of America **112**,
E3574-81 (2015).

15. Long, S. W. *et al.* Population genomic analysis of 1,777 extended-spectrum beta-lactamase-
producing Klebsiella pneumoniae isolates, Houston, Texas: Unexpected abundance of clonal
group 307. *mBio* **8**, (2017).

16. Van Tyne, D. *et al.* Impact of antibiotic treatment and host innate immune pressure on
enterococcal adaptation in the human bloodstream. *Science translational medicine* **11**,
(2019).

17. Copin, R. *et al.* Sequential evolution of virulence and resistance during clonal spread of
community-acquired methicillin-resistant Staphylococcus aureus. *Proceedings of the*
National Academy of Sciences of the United States of America **116**, 1745–1754 (2019).

18. Lytsy, B. *et al.* The first major extended-spectrum β-lactamase outbreak in Scandinavia was
caused by clonal spread of a multiresistant Klebsiella pneumoniae producing CTX-M-15.
APMIS **116**, 302–8 (2008).
19. Ransjö, U. *et al.* Hospital outbreak control requires joint efforts from hospital management,
microbiology and infection control. *Journal of Hospital Infection* **76**, 26–31 (2010).
20. Sandegren, L., Linkevicius, M., Lytsy, B., Melhus, Å. & Andersson, D. I. Transfer of an
Escherichia coli ST131 multiresistance cassette has created a Klebsiella pneumoniae-specific
plasmid associated with a major nosocomial outbreak. *Journal of Antimicrobial*
Chemotherapy **67**, 74–83 (2012).
21. Alteri, C. J. & Mobley, H. L. T. Metabolism and Fitness of Urinary Tract Pathogens.
Microbiology Spectrum **3**, MBP-0016-2015 (2021).
22. Pál, C. & Papp, B. Evolution of complex adaptations in molecular systems. *Nat Ecol Evol* **1**,
1084–1092 (2017).
23. Browne, N., Heelan, M. & Kavanagh, K. An analysis of the structural and functional
similarities of insect hemocytes and mammalian phagocytes. *Virulence* **4**, 597–603 (2013).
24. Tsai, C. J.-Y., Loh, J. M. S. & Proft, T. Galleria mellonella infection models for the study of
bacterial diseases and for antimicrobial drug testing. *Virulence* **7**, 214–229 (2016).
25. Wand, M. E., Mccowen, J. W. I., Nugent, P. G. & Sutton, J. M. Complex interactions of
Klebsiella pneumoniae with the host immune system in a Galleria mellonella infection
model. *Journal of Medical Microbiology* **62**, 1790–1798 (2013).
26. Insua, J. L. *et al.* Modeling Klebsiella pneumoniae Pathogenesis by Infection of the Wax
Moth Galleria mellonella. *Infection and Immunity* **81**, 3552–3565 (2013).
27. Bruchmann, S., Feltwell, T., Parkhill, J. & Short, F. L. Identifying virulence determinants of
multidrug-resistant Klebsiella pneumoniae in Galleria mellonella. *Pathogens and disease* **79**,
ftab009 (2021).
28. Short, F. L. *et al.* Genomic Profiling Reveals Distinct Routes To Complement Resistance in
Klebsiella pneumoniae. *Infection and immunity* **88**, e00043-20 (2020).
29. Shon, A. S., Bajwa, R. P. S. & Russo, T. A. Hypervirulent (hypermucoviscous) Klebsiella
pneumoniae. *Virulence* **4**, 107–118 (2013).
30. Wugeditsch, T. *et al.* Phosphorylation of Wzc , a Tyrosine Autokinase , Is Essential for
Assembly of Group 1 Capsular Polysaccharides in Escherichia coli *. *The Journal of*
Biological Chemistry **276**, 2361–2371 (2001).
31. Whitfield, C., Wear, S. S. & Sande, C. Assembly of Bacterial Capsular Polysaccharides and
Exopolysaccharides. *Annual Review of Microbiology* **74**, 521–543 (2020).
32. Payne, S. M. & Neilands, I. B. Iron and Virulence in the Family Enterobacteriaceae. *CRC*
Critical Reviews in Microbiology **16**, 81–111 (1988).
33. Bachman, M. A. *et al.* Klebsiella pneumoniae yersiniabactin promotes respiratory tract
infection through evasion of lipocalin 2. *Infection and immunity* **79**, 3309–3316 (2011).
34. Holden, V. I. *et al.* Iron Acquisition and Siderophore Release by Carbapenem-Resistant
Sequence Type 258 Klebsiella pneumoniae. *mSphere* **3**, e00125-18 (2018).

35. Macleod, S. M. & Stickler, D. J. Species interactions in mixed-community crystalline
biofilms on urinary catheters. *Journal of Medical Microbiology* **56**, 1549–1557 (2007).
36. Frank, D. N., Wilson, S. S., St. Amand, A. L. & Pace, N. R. Culture-independent
microbiological analysis of Foley urinary catheter biofilms. *PLoS ONE* **4**, 1–7 (2009).
37. Flores-Mireles, A. L., Walker, J. N., Caparon, M. & Hultgren, S. J. Urinary tract infections:
epidemiology, mechanisms of infection and treatment options. *Nature Reviews Microbiology*
13, 269–284 (2015).
38. Zaborskytė, G., Wistrand-Yuen, E., Hjort, K., Andersson, D. I. & Sandegren, L. Modular
3D-Printed Peg Biofilm Device for Flexible Setup of Surface-Related Biofilm Studies.
Frontiers in Cellular and Infection Microbiology **11**, 802303 (2022).
39. Flores-Mireles, A. L. *et al.* Fibrinogen Release and Deposition on Urinary Catheters Placed
during Urological Procedures. *Journal of Urology* **196**, 416–421 (2016).
40. Flores-Mireles, A. L., Pinkner, J. S., Caparon, M. G. & Hultgren, S. J. EbpA vaccine
antibodies block binding of Enterococcus faecalis to fibrinogen to prevent catheter-
associated bladder infection in mice. *Science Translational Medicine* **6**, 254ra127-254ra127
(2014).
41. Walker, J. N. *et al.* Catheterization alters bladder ecology to potentiate Staphylococcus
aureus infection of the urinary tract. *Proceedings of the National Academy of Sciences of the*
United States of America **114**, E8721–E8730 (2017).
42. Andersen, M. J. *et al.* Inhibiting Host Protein Deposition on Urinary Catheters Reduces
Associated Urinary Tract Infections. *eLife* **11**, e75798 (2022).
43. Weissman, S. J. *et al.* Differential Stability and Trade-Off Effects of Pathoadaptive
Mutations in the Escherichia coli FimH Adhesin. *Infection and Immunity* **75**, 3548 (2007).
44. Chen, S. L. *et al.* Positive selection identifies an in vivo role for FimH during urinary tract
infection in addition to mannose binding. *Proceedings of the National Academy of Sciences*
of the United States of America **106**, 22439–22444 (2009).
45. Stahlhut, S. G. *et al.* Population variability of the FimH type 1 fimbrial adhesin in Klebsiella
pneumoniae. *Journal of Bacteriology* **191**, 1941–1950 (2009).
46. Sokurenko, E. V. *et al.* Pathogenic adaptation of Escherichia coli by natural variation of the
FimH adhesin. *Proceedings of the National Academy of Sciences of the United States of*
America **95**, 8922–8926 (1998).
47. Malone, J. G. *et al.* The YfiBNR Signal Transduction Mechanism Reveals Novel Targets for
the Evolution of Persistent Pseudomonas aeruginosa in Cystic Fibrosis Airways. *PLOS*
Pathogens **8**, e1002760 (2012).
48. Wall, E., Majdalani, N. & Gottesman, S. The Complex Rcs Regulatory Cascade. *Annual*
Review of Microbiology **72**, 111–139 (2018).
49. Van Tyne, D., Ciolino, J. B., Wang, J., Durand, M. L. & Gilmore, M. S. Novel Phagocytosis-
Resistant Extended-Spectrum β-Lactamase-Producing Escherichia coli From Keratitis.
JAMA ophthalmology **134**, 1306–1309 (2016).

50. Miskinyte, M. *et al.* The Genetic Basis of Escherichia coli Pathoadaptation to Macrophages.
PLoS Pathogens **9**, e1003802 (2013).
51. Martin, R. M. *et al.* Molecular Epidemiology of Colonizing and Infecting Isolates of
Klebsiella pneumoniae. *mSphere* **1**, e00261-16 (2016).
52. Martin, R. M. & Bachman, M. A. Colonization, Infection, and the Accessory Genome of
Klebsiella pneumoniae. *Frontiers in Cellular and Infection Microbiology* **8**, 1–15 (2018).
53. Joseph, L., Merciecca, T., Forestier, C., Balestrino, D. & Miquel, S. From Klebsiella
pneumoniae Colonization to Dissemination: An Overview of Studies Implementing Murine
Models. *Microorganisms 2021, Vol. 9, Page 1282* **9**, 1282 (2021).
54. Ernst, C. M. *et al.* Adaptive evolution of virulence and persistence in carbapenem-resistant
Klebsiella pneumoniae. *Nature Medicine* **26**, 705–711 (2020).
55. Marsh, J. W. *et al.* Evolution of Outbreak-Causing Carbapenem-Resistant Klebsiella
pneumoniae ST258 at a Tertiary Care Hospital over 8 Years. *mBio* **10**, e01945-19 (2019).
56. Martin, M. J. *et al.* Anatomy of an extensively drug-resistant Klebsiella pneumoniae
outbreak in Tuscany, Italy. *Proceedings of the National Academy of Sciences* **118**,
e2110227118 (2021).
57. Chen, S. L. *et al.* Identification of genes subject to positive selection in uropathogenic strains
of Escherichia coli: A comparative genomics approach. *Proceedings of the National
Academy of Sciences* **103**, 5977–5982 (2006).
58. Lieberman, T. D. *et al.* Genetic variation of a bacterial pathogen within individuals with
cystic fibrosis provides a record of selective pressures. *Nature Genetics 2013 46:1* **46**, 82–87
(2013).
59. Traverse, C. C., Mayo-Smith, L. M., Poltak, S. R. & Cooper, V. S. Tangled bank of
experimentally evolved Burkholderia biofilms reflects selection during chronic infections.
Proceedings of the National Academy of Sciences **110**, E250–E259 (2013).
60. Dyzenhaus, S. *et al.* MRSA lineage USA300 isolated from bloodstream infections exhibit
altered virulence regulation. *Cell Host & Microbe* (2023) doi:10.1016/j.chom.2022.12.003.
61. Damkiær, S., Yang, L., Molin, S. & Jelsbak, L. Evolutionary remodeling of global regulatory
networks during long-term bacterial adaptation to human hosts. *Proceedings of the National
Academy of Sciences* **110**, 7766–7771 (2013).
62. Gatt, Y. E. & Margalit, H. Common Adaptive Strategies Underlie Within-Host Evolution of
Bacterial Pathogens. *Molecular biology and evolution* **38**, 1101–1121 (2021).
63. Khadka, S. *et al.* Urine-mediated suppression of Klebsiella pneumoniae mucoidy is
counteracted by spontaneous Wzc variants altering capsule chain length. *mSphere* **0**, e00288-
23 (2023).
64. Dorman, M. J., Feltwell, T., Goulding, D. A., Parkhill, J. & Short, F. L. The Capsule
Regulatory Network of Klebsiella pneumoniae Defined by density-TraDISort. *mBio* **9**, 1–19
(2018).
65. Guachalla, L. M. *et al.* Discovery of monoclonal antibodies cross-reactive to novel
subserotypes of K. pneumoniae O3. *Scientific Reports* **7**, (2017).

66. Maldonado, R. F., Sá-Correia, I. & Valvano, M. A. Lipopolysaccharide modification in
Gram-negative bacteria during chronic infection. *FEMS Microbiol Rev* **40**, 480–493 (2016).
67. Hsu, C.-R. *et al.* Identification of a capsular variant and characterization of capsular
acetylation in Klebsiella pneumoniae PLA-associated type K57. *Scientific Reports* **6**, (2016).
68. Feng, Y. *et al.* Key evolutionary events in the emergence of a globally disseminated,
carbapenem resistant clone in the Escherichia coli ST410 lineage. *Communications Biology*
2019 2:1 **2**, 1–13 (2019).
69. Palacios, M., Broberg, C. A., Walker, K. A. & Miller, V. L. A Serendipitous Mutation
Reveals the Severe Virulence Defect of a Klebsiella pneumoniae fepB Mutant. *mSphere* **2**,
e00341-17 (2017).
70. Ayala-Castro, C., Saini, A. & Outten, F. W. Fe-S Cluster Assembly Pathways in Bacteria.
Microbiology and Molecular Biology Reviews **72**, 110–125 (2008).
71. Braun, V. FhuA (TonA), the career of a protein. *Journal of Bacteriology* **191**, 3431–3436
(2009).
72. Ghequire, M. G. K., Buchanan, S. K. & de Mot, R. The ColM family, polymorphic toxins
breaching the bacterial cell wall. *mBio* **9**, e02267-17 (2018).
73. Aguileta, G., Refrégier, G., Yockteng, R., Fournier, E. & Giraud, T. Rapidly evolving genes
in pathogens: Methods for detecting positive selection and examples among fungi, bacteria,
viruses and protists. *Infection, Genetics and Evolution* **9**, 656–670 (2009).
74. Hossein Khademi, S. M., Sazinas, P. & Jelsbak, L. Within-Host Adaptation Mediated by
Intergenic Evolution in Pseudomonas aeruginosa. *Genome Biology and Evolution* **11**, 1385–
1397 (2019).
75. Camacho, C. *et al.* BLAST+: architecture and applications. *BMC Bioinformatics* **10**, 421
(2009).
76. Szklarczyk, D. *et al.* The STRING database in 2021: customizable protein–protein networks,
and functional characterization of user-uploaded gene/measurement sets. *Nucleic Acids*
Research **49**, D605–D612 (2021).
77. Huerta-Cepas, J. *et al.* eggNOG 5.0: a hierarchical, functionally and phylogenetically
annotated orthology resource based on 5090 organisms and 2502 viruses. *Nucleic Acids*
Research **47**, D309–D314 (2019).
78. Thulin, M. BAT: an online tool for analysing growth curves. https://mansthulin.se/bat/
(2018).
79. Loh, J. M. S., Adenwalla, N., Wiles, S. & Proft, T. Galleria mellonella larvae as an infection
model for group A streptococcus. *Virulence* **4**, 419–428 (2013).
80. Gomes, A. É. I. *et al.* Selection and validation of reference genes for gene expression studies
in Klebsiella pneumoniae using Reverse Transcription Quantitative real-time PCR. *Sci Rep*
8, 9001 (2018).
81. Livak, K. J. & Schmittgen, T. D. Analysis of Relative Gene Expression Data Using Real-
Time Quantitative PCR and the 2−ΔΔCT Method. *Methods* **25**, 402–408 (2001).
716

Fig. 1 | Strong positive selection of genetic changes during infection. a, Overview of isolates included in the study. **b,** Genetic changes per isolate relative to the index isolate plotted over time of isolation. Red isolates (DA25065 and DA69557) were excluded from the calculation of mutations/genome/year because the number of genetic changes exceeds the mean plus two standard deviations. The insert shows infection and colonization isolates separately during the period when both types of isolates were collected (starting in September 2006). **c,** Maximum likelihood phylogeny of outbreak isolates. Clusters of isolates are shaded and assigned to letters A-H. **d,** Functional grouping of genes with at least two nonsynonymous mutations selected independently as tip mutations. Asterisks denote groups with a significant bias between infecting and colonizing isolates (binomial test, *p<0.05). The intergenic regions not assigned to any of the functional groups can be found in Supplementary Table 3. **e,** Number of colonizing and infection site isolates with at least one (1+) and 2 or more (2+) top genes and intergenic regions mutated. Bias between infection and colonization was tested using the binomial test, p values are depicted above the columns. **f,** Examples of convergent mutation selection within phylogenetic clusters.

Fig. 2 | Virulence profiles show frequent attenuation and occasional increase. a, Kaplan-Meier survival curves of *G. mellonella* larvae infected with bacterial isolates (n=20 larvae for each isolate). Assignment of virulence groups is based on larval survival and the cumulative health index scores of the surviving larvae after 48 h, compared with the index isolate (log-rank Mantel-Cox test). **b**, Distribution of isolates from infection and colonization among the virulence groups (binomial test). **c**, Single mutations identified as responsible for acute virulence loss ("*" – early stop codon, "fs" – frameshift). **d**, Venn diagram showing the overlap among the genetic targets (genes and intergenic regions) between isolates with different virulence profiles. Each genetic change was categorized as possibly contributing to the virulence based on the isolate's virulence profile and its phylogenetic location. **e**, Functional groups of unique mutations in the "no change" and "attenuation"/"slower killing" groups. Named genes acquired at least two independent mutations in attenuated/slower killing isolates. **f**, Upper panel: bacterial survival in human serum (log10 change in CFU/ml after 3 h) versus sedimentation (mucoviscosity) in different virulence groups. Lower panel: independent mutations in *cps* and O-antigen regions. Part of phylogenetic cluster B has an intergenic mutation *galF/cpsACP* conferring resistance to serum.

Fig. 3 | Regulatory changes in iron uptake and utilization. **a**, Schematic overview of *K. pneumoniae* iron uptake systems for iron bound to sidero-phores, such as catecholate-type enterobactin or hydroxamate-type ferrichrome, and free ferrous iron. **b**, Regulatory elements in the *fepA/fes* region and changes in *fepA* and *fes* mRNA levels in isolates with mutations (noted in dark red) relative to the index isolate. Comparison to the index isolate in each condition was done by unpaired t-test; significance at p<0.01 after Bonferroni correction for five comparisons. **c**, Changes in *fepA* and *fes* mRNA levels in the index isolate under low and high iron conditions. Results are shown for three biological replicates. mRNA levels in **b** and **c** are relative to the house-keeping gene *glnA* and further normalized to M9/Chelex + FeSO$_4$. Comparison between low and high iron conditions was done by one-way ANOVA followed by Dunnett's multiple comparison test; p values are depicted above the columns.

Fig. 4 | Convergent evolution of biofilm formation capacity. **a**, Phylogenetic distribution of isolates with increased surface-attached biofilm capacity relative to the index isolate. Bars indicate medians of total CV-stained biomass of 48 h biofilms grown on silicone-coated pegs without (light grey) or with fibrinogen coating (dark grey). At least four biological replicates (dots) are shown. Highlighted genetic changes are most likely responsible for the increased biofilm phenotype. **b**, Interconnection between acute virulence and survival in serum among isolates with increased biofilm formation capacity. **c**, Biofilm evolution scenarios with consecutive genetic changes illustrated on branches where mutations leading to increased (DA25060, DA25130/- DA25128, DA69557) or decreased (DA25061, DA25127) biofilm formation are shown.

Fig. 5 | Colonization of the gastrointestinal tract in mice is increased with biofilm formation. a, Overall CFU/g faeces in each mouse (n=35) during the 10-day experiment. On Day 5, clindamycin was added to the drinking water after a third oral gavage with bacteria. **b**, Top panel illustrates phenotypes for each isolate. Lower panel displays competitive indices during days 6-10 of gastrointestinal colonization. Competitive index >0 means an advantage of the tested isolate versus the index isolate DA14734, <0 – disadvantage. Dots show the competitive index in individual mice (n=6 per competition) and the trendline connects the means. Comparisons to control competition on each day were made by unpaired two-tailed t-test; significance after Bonferroni correction for multiple comparisons at p<0.01, p values depicted above the datapoints.