Optogenetic control of pheromone gradients reveals functional limits of mating behavior in budding yeast

Alvaro Banderas[1,*], Maud Hofmann[1], Celine Cordier[1], Matthias Le Bec[1], Carolina Elizondo[1], Lionel Chiron[1], Sylvain Pouzet[1], Vittore Scolari[1,2] and Pascal Hersen[1,*]

[1] Institut Curie, Université PSL, Sorbonne Université, CNRS UMR168, Physics of Cells and Cancer, 75005 Paris, France.

[2] Institut Curie, Université PSL, Sorbonne Université, CNRS UMR3664, Laboratoire Dynamique du Noyau, 75005 Paris, France.

Corresponding authors: alvaro.banderas@curie.fr and *pascal.hersen@curie.fr*

Keywords: Optogenetics, Mating, Budding Yeast, Cell-cell communication, Pheromone, Reaction-Diffusion, Gradients

Abstract

Cell-cell communication through diffusible signals allows distant cells to coordinate biological functions. Such coordination depends on the signal landscapes generated by emitter cells and the sensory capacities of receiver cells. In contrast to morphogen gradients in embryonic development, microbial signal landscapes occur in open space with variable cell densities, spatial distributions, and physical environments. How do microbes shape signal landscapes to communicate robustly under such circumstances remains an unanswered question. Here we combined quantitative spatial optogenetics with biophysical theory to show that in the mating system of budding yeast—

where two mates communicate to fuse—signal landscapes convey demographic or positional

information depending on the spatial organization of mating populations. This happens because

α-factor pheromone and its mate-produced protease Bar1 have characteristic wide and narrow

diffusion profiles, respectively. Functionally, MATα populations signal their presence as

collectives, but not their position as individuals, and Bar1 is a sink of alpha-factor, capable of both

density-dependent global attenuation and local gradient amplification. We anticipate that

optogenetic control of signal landscapes will be instrumental to quantitatively understand the

spatial behavior of natural and engineered cell-cell communication systems.

1. Introduction

Cell-cell communication through extracellular signals enables the spatiotemporal coordination of cell behavior across various scales, from single cells to tissues. The interplay between signal production, diffusion, and reaction (e.g., internalization, adsorption, and degradation) shapes the concentration landscapes that cells experience and ultimately determines a cell's responses. A classic example is the establishment of positional information in the *Drosophila melanogaster* syncytial embryo, in which an exponentially decaying extracellular *bicoid* morphogen gradient forms as a result of the specific location of emitter nuclei on the anterior region, free diffusion of the signal, and homogenous degradation of the signal throughout the embryo[1,2]. This smooth gradient is "read" by nuclei to switch on specific differentiation programs in sharply defined spatial locations.

In contrast to multicellular developmental systems where cells follow a predefined program from a well-defined initial condition, communication does not occur within closed spaces for free-living microbes. Populations of single cells often have a random spatial distribution, expand freely, explore an unknown environment, and create an ecosystem—the size and dynamics of which are the result of cellular metabolic interactions and cell-cell communication. As in development, signal landscapes depend on reaction-diffusion dynamics; however, population-level parameters such as cell density, population composition, and the spatial distribution of emitter and receiver cells are inherently variable in the context of single-cell populations. Thus, the way cells extract information and take appropriate decisions in such an uncertain environment is a very different biological problem than what is observed in embryology.

The importance of this question can be epitomized by the commonly accepted view that, in bacteria,

communication through diffusible signals provides demographic rather than positional

information. The use of demographic information by cells for decision-making is usually termed

quorum sensing[3], a mode of communication in which the signal concentration is proportional to

the population density, which allows cells—that act both as emitters and receivers—to activate

specific phenotypes once a signal threshold is sensed. Crucially, cell spatial organization and signal

diffusion (and degradation) in quorum-sensing systems have been experimentally shown to play

an important role in how much a cell can perceive the local—rather than the global—population

composition through "short-range" communication[4,5]. Short-range communication in bacteria is

not argued to provide positional information (position and orientation respect to a source), as

bacterial cells are often too small to perceive gradients. Rather, signals reflect the local population

composition which informs bacteria about the likelihood of interacting with a relevant partner.

In contrast to bacteria, in the well-known example of mating through pheromone communication

in the budding yeast *Saccharomyces cerevisiae*, information gathered by cells is usually considered

positional, i.e., gradients formed will attract cells to grow towards the position of the source and

cells possess mechanisms to perceive the direction of the gradient. Nonetheless, as in bacteria,

demographic information has also been shown to be a relevant cue in mating decision making[6]

and, furthermore, positional information seems to be available to cells only over very short

(contact) distances from partners[7].

However, unlike developmental and bacterial communication, both positional and demographic

information are entangled in the case of yeast. How yeast cells perceive demographic and

positional information to their benefit remains an unresolved question.

Mating in yeast is governed by the mating-pheromone pathway,[8,9] which relies on the secretion of

pheromones to signal the presence of different mating types (*MAT*a and *MAT*α). The *MAT*a and

*MAT*α mating-types exchange pheromones to trigger activation of their mating pathway,

orchestrating a dedicated transcriptional program, cell cycle arrest, and morphological changes

that culminate with cell and diploid formation. Distinct branches of the pheromone pathway

control transcriptional (Ste2 branch) and arrest/morphology behaviors (Far1 branch). Dose-

response experiments measuring the responses of *MAT*a cells to the purified pheromone α-factor

in bulk showed that, while the transcriptional branch of the pathway displays a Michaelian (linear

at low doses) input-output relation[10], cells transition sharply—following a Hill-type sigmoidal

response—from proliferating to growth-arrested elongated phenotypes, called mating protrusions

(or "shmoos")[11,12]. This "mating switch" **(Fig. 1A)** was notably observed in artificial gradients of α-

factor[13,14] generated using microfluidic devices. In this setting, cell cycle-arrested morphologies

were observed over a narrow range of concentrations (1–5 nM) of α-factor. Cells only displayed

chemotropic behavior, elongating their cell body up the α-factor gradient, within this concentration

range. However, importantly, it remains unknown how the amplitude and steepness of artificially

generated gradients in the context of microfluidic devices compare to the gradients that are

generated within heterogeneous, large populations of mating cells.

*MAT***a** cells secrete the Bar1 protein which degrades α-factor extracellularly. Thus, the pheromone

landscape[6] that *MAT***a** cells experience is not only determined by spatiotemporal diffusion of the

α-factor pheromone produced by *MAT*α cells, but also by the concentration of Bar1 they produce

themselves.

For a non-motile organism such as *S. cerevisiae*, evolution would be expected to produce systems

where the concentrations of α-factor that trigger mating are found only in the close vicinity of

emitter cells, so that cells do not have mating responses unless they are very close to a partner. Bar1

has been deemed responsible to exert such a function[15], by increasing the degradation rate of α-

factor and confining its activity close to the producer cells, consequently shortening the length-

scale at which mating sensing occurs to only a few micrometers (a fraction of cell size). However,

the concentration of Bar1 during mating reactions is dependent on the density of *MAT***a** cells;

therefore, the levels of Bar1 may not always be sufficient to degrade the α-factor produced by cells

of the opposing mating type. This implies that the mating switch distance can be, in theory,

extended further away from the source position (**Fig. 1B**). This scenario is plausible in outcrossing

reactions, where a *MAT*α cell within a single mating-type population can switch into a *MAT***a** cell

(and vice versa) by means of the *HO* endonuclease activity[16]. Consequently, the superimposition of

gradients produced by multiple pheromone sources represents a potential confounding factor (**Fig.**

1C): partners located at a distance from source cells cannot decide with which partner to mate[17]. To

put it differently, it remains unclear how cells extract relevant positional information from

physiological pheromone concentration landscapes[6]. Indeed, in mating discrimination assays[17],

*bar1Δ MAT***a** can discriminate real emitters from dummy (that do not produce pheromone)

emitters, but only if real emitters are in the minority. This suggests that in the absence of

degradation, pheromone halos from other cells can be a confounding factor for receiver cells.

Several theories have attributed various possible functions to Bar1 in mating populations. Three of

these theories depend on high local concentrations of Bar1 in receiver cells, acting as a localized

sinks of α-factor: (i) reducing the size of the pheromone halo around emitters (at the expense of

gradient strength) to correctly align to gradients in the presence of multiple sources[18], (ii) gradient

disentanglement from two sources in close-proximity (intra-ascus) interactions,[19] and (iii) mutual-

avoidance in *MAT***a** cells[20]. However, evidence for *MAT***a** receivers acting as pheromone sinks

during mating is currently lacking and its quantitative contribution to α-factor gradient reshaping

during mating has not been determined.

Understanding the spatial properties of α-factor-mediated communication between mating types

requires characterizing the properties of physiological α-factor gradients. Although the shape of

gradients can be inferred using numerical simulations of cells randomly arranged in space[21], the

inability to experimentally control α-factor production locally and the high response variability

among receivers preclude the construction of predictive models. To build and test predictive

models, one must first obtain quantitative measurements over a large number of cells within large

areas in which both mating types are spatially organized. Although possible, the physical

separation of mating types in microfluidic devices[22] or by micromanipulation[23] is very tedious to

achieve and hard to monitor and reproduce. Thus, while a method to obtain reliable spatial control

of α-factor production within physiological range at both large and small scales is critical, no

proven method currently exists.

Here, we address this problem using the power of optogenetics (**Fig. 1D**). We employed spatial

control of α-factor production and degradation in *Saccharomyces cerevisiae* and obtained a

quantitative description of extracellular α-factor gradients under physiological conditions.

Combined with a diffusion-reaction model, we inferred quantitative parameters that describe the

mating communication system and provide a proof of principle for the rational, quantitative

control of extracellular signal gradients within microbial cell populations.

140   2. Results

2.1 Engineered optogenetic induction of physiological α-factor levels allows spatial control of

mating.

We first aimed at engineering light-inducible strains that can produce physiological levels of

pheromone, by improving on a previous strain design[24]. For this we first engineered wild-type

MATα yeast cells using a CRISPR-based promoter replacement strategy (**see Methods**) to replace

the promoter of the two α-factor pheromone-coding genes (MF-α1 and MF-α2) with the P_{C120}

promoter, the activity of which is induced by the light-sensitive transcriptional activator EL222[25].

Additionally, to increase α-factor production and recover wild-type levels, a second copy of the

P_{C120}-MFα1 construct was inserted at an unrelated locus (**Fig. 2A, Figure S1, see Methods**) to yield

the "opto-α" strain used in this study. To quantify the levels of α-factor produced by the opto-α

strain, we used a *MAT**a*** reporter strain that harbors a super-folder green fluorescent protein (GFP)

downstream of the P_{FUS1} pheromone-responsive promoter[26] (**Fig. 2B**). This reporter strain harbors

the native untranslated terminal region (UTR) of FUS1 and carries a ubiquitination sequence that

targets super-folder GFP for faster degradation, thus providing a good estimate of the abundance

of the native transcript[26]. We stimulated mixes of opto-α and reporter strains with light using a

custom-made light-plate apparatus[27] and quantified the levels of the α-factor pheromone (relative

to the wild-type MATα strains) using two reporter strains that either carry or lack the Bar1 protease

(**Fig. 2B**).

In both cases, the levels of α-factor were light-intensity dependent. In the absence of Bar1,

pheromone levels comfortably reached the induction levels achieved by the wild type. Under the

more stringent conditions where receiver cells express Bar1, the opto-α strain reached up to ~60%

of the wild-type response. As other strains and conditions resulted in lower α-factor production

levels (**Fig. S1,S2**), we focused on the opto-α strain. Next, we checked if the levels of pheromone

produced can rescue wild-type physiological behavior. For this, we tested the capacity of *MAT***a**-

*bar1*Δ reporter cells to develop mating protrusions in response to light-induced α-factor production

by opto-α cells. With sufficient intensity of light and, like with wild-type *MAT*α emitters, we

observed formation of protrusions by *MAT***a** *bar1*Δ cells in gel-embedded mixes (**Fig. 2C, see**

Methods). Finally, we assayed diploid formation in opto-α/*MAT***a** mixes as a measure of the final

physiological output of the mating pathway (**Fig. 2 D,E).** For this, we developed an "optogenetic

mating assay" (**Fig. S3, see Methods**) based on three features. First, we used the top-agar technique

to embed mating mixtures (as used in classical growth arrest assays), which provides a

homogeneous layer where cells are located sparsely and grow as microcolonies. Second, we

employed a digital micromirror device (DMD)[28] to expose the cells to spatial light patterns

projected into the agar (**See Methods**); and finally, auxotrophic markers were used to ensure that

only diploid colonies can grow, allowing us to have a measure of light-induced mating efficiency.

Our results show that pheromone production is indeed sufficient to induce cell-cell fusion with

wild-type *MAT***a** in a light dose-dependent manner and could reach comparable maximum levels

as the wild-type *MATα* emitter (**Fig. 2 C,D**).

Taken together, these results show that the opto-α strain is a suitable candidate, comparable to the

wild-type, that produces physiological levels of pheromone production and mating behavior in

response to light. Therefore, we employed the opto-α strain as a remotely controlled signal

generator to create realistic α-factor spatial gradients within large populations of opto-α/MATa

cells.

2.2 Optogenetic induction of pheromone degradation allows specific control of growth.

It is expected that both pheromone production and pheromone degradation by the α-factor

protease Bar1 shape extracellular pheromone gradients[21]. To gain spatial control over pheromone

degradation, we constructed the opto-Bar1 *MAT***a** strain, in which the α-factor protease Bar1

expression is placed under control of the P_{C120} light-inducible promoter. We followed a similar

promoter replacement strategy as for the opto-α strain, targeting the Bar1 locus in *MAT***a** cells (**Fig.**

3A). The expression levels and activity of Bar1 protein were measured using the *MAT***a** *bar1Δ* α-

factor reporter, in the presence of purified α-factor added to the media. The levels of GFP produced

by the reporter strain were used to quantify the production of Bar1, since Bar1 degrades purified

α-factor in the culture medium (**Fig. 3A**). We observed light-dependent Bar1 activity, with the level

of the reporter response decreasing as the light intensity increased (**Fig. 3B**). We reached higher

Bar1 activity than the wild-type strain and undetectable basal (in the dark) Bar1 enzymatic activity.

Although Bar1 degrades the pheromone, its function has been long associated with increases in

mating efficiency through several candidate mechanisms[17–20,26] (see **Introduction**). However, the

presence of Bar1 is not essential for mating itself. Indeed, when performing the optogenetic mating

assay with the opto-Bar1 strain, cells located in dark regions—where *MAT***a** cells do not express

Bar1—remain proficient at forming diploids (**Fig. 3C**), with a ~2-fold difference in mating efficiency

between illuminated and dark regions (**Fig. S4**). Thus, contrary to optogenetic pheromone

induction—and as expected—optogenetic pheromone degradation does not restrict but mildly

enhances the final output of the pathway, namely the formation of diploids.

Cell cycle arrest, another physiological output of mating events, is controlled by the mating-

pheromone pathway through the FAR1 branch[29]. When Bar1 is produced and actively degrades α-

factor, one would expect to observe fewer cell cycle arrest events. Following similar methods as for

diploid formation, we used the optogenetic plate assay to measure the capacity of *MAT***a** cells to

escape cell cycle arrest as a function of Bar1 induction. We imaged an intermediate stage of the

experiment (**Fig. 3D**). A differential growth pattern was observed in the non-selective plate used

as template for replica plating in diploid-selective media (See also **Fig. S3**), as illuminated cells

(expressing Bar1) grew similarly to wild-type levels, whereas non-illuminated cells did not grow

significantly (**Fig. 3D-F).**

These results show that illuminated opto-bar1 cells recover the expected wild-type behavior in

terms of α-factor degradation activity, mating, and proliferation.

2.3 Spatial optogenetic induction of α-factor production generates distance-dependent

responses in mating populations.

After quantitative characterization of our optogenetic strains, we next moved to assess the spatial

properties of pheromone-mediated communication between $MAT\alpha$ and $MAT\mathbf{a}$ cells in an idealized

(but physiological) context. To that end, we implemented the "half-domain assay", which consists

of monitoring $MAT\alpha/MAT\mathbf{a}$ mating reactions within a spatially homogeneous random distribution

of cells of both mating types placed at the bottom of agarose-embedded "miniwells" (**Fig. 4A, see**

Methods). Half of the total surface was exposed to blue light, to optogenetically induce pheromone

production exclusively in the illuminated domain. The response of reporter cells in the dark

domain allowed us to read and quantify the diffusion-reaction properties of α-factor.

We first conducted experiments with the opto-α strain as a light-dependent pheromone generator

and the $bar1\Delta$ $MAT\mathbf{a}$ reporter strain as a pheromone receiver and fluorescent reporter of mating

behavior. Absence of Bar1 simulates natural situations where wild-type $MAT\mathbf{a}$ receivers are in the

minority (e.g., upon a mating type switching event), where Bar1 levels are not enough to modify

the global pheromone concentration. Thus, in this experiment, enzymatic degradation of α-factor

by Bar1 is absent and disappearance of the pheromone is solely determined by its intake by cells[30].

After illumination (**See Methods**), cells along a stripe located in the center of the plate, with its long

axis perpendicular to the light-dark interface, were imaged (**Fig. 4B**). A schematic of the experiment

and strains used is also shown (**Fig. 4C**). The opto-α strain was equipped with a red nuclear marker

(the HTB2 histone fused to mApple) for easy identification, which is especially important in regions

of clumping (**Fig. S5**, this new strain is named opto-α*). We used a 20x objective and an automated

stitching method (**see Methods**) to obtain images of a large stripe (5671 μm x 330 μm) of both cell

types using fluorescence and bright field microscopy at single-cell resolution (**Fig. S5**). Our

experimental results revealed a sharp phenotypic transition between illuminated and non-

illuminated areas. The sharpness of this transition (compared to the sharpness of the light border)

was assessed by running a parallel experiment in neighboring mini-wells by stimulating a control

strain that expresses YFP under the control of an equivalent EL222/P$_{C120}$ light-inducible system. This

control serves as a biological spatial imprint of the light-stimulated region, to both locate the

position of the border in the sample and determine the sharpness of the light border (**Fig. 4D,E**).

The region in the vicinity of the light-to-dark transition border illustrates the overall trend of the

mating response in the *MAT**a** bar1Δ* responder population: the GFP signal gradually vanished in

cells further and further away from the frontier in the dark domain (**Fig. 4D**). Importantly, since

we imaged cells at single-cell resolution, we clearly observed the cell morphology transitioned from

elongated shapes to round as the distance from the light to dark frontier increased (**Fig. 4E**).

Our assay demonstrates that optogenetics can be used to produce large-scale population-generated

gradients of a pheromone to infer the shape of the pheromone gradient through quantification of

gene-expression and cell-morphology outputs (**Figs. 4C,D, S5**).

2.4 Quantitative analysis reveals that population-level pheromone gradients are long-range,

shallow, and limit morphological transitions to the vicinity of the source.

Using the half-domain assay, we next sought to infer the biophysical properties of the generated

gradient. Thus, we constructed a biophysical model based on reaction-diffusion dynamics—as well

as dose-response functions—and obtained single-cell quantifications of pheromone-responses

using a combination of manual and automated single-cell segmentation and analysis (**see**

Methods). To determine the relevant diffusion-reaction parameters governing pheromone

communication between cells, we modeled the steady-state pheromone gradient profiles and the

corresponding induced cell responses. Taking advantage of the symmetry within our experimental

setting, we simplified the reaction-diffusion equations to describe a one-dimensional infinite field

of pheromone producers (**Fig. 5A**).

264
$$\frac{\partial c(x,t)}{\partial t} = D\nabla^2 c(x,t) - \Omega\, c(x,t) + \rho A(x), \tag{1}$$

where $c(x,t)$ is the concentration of α-factor, D is its diffusion coefficient, A is the pheromone

production rate, ρ is the cell density, and Ω is the rate of degradation of the pheromone by cell

uptake[30]. This equation produces the well-known solution for the diffusion equation for a point

source under uniform degradation at the steady state, $c^*(x) = \frac{A}{2\sqrt{D\Omega}} e^{-\sqrt{\Omega/D}\,x}$, which describes

gradients formed by a single cell in one dimension. For the half-domain where cells are illuminated

and therefore producing pheromone ($x < 0$), the solution depends on the cell density (**Fig. 5B**) and

the steady-state pheromone concentration profile is described as follows (see full derivation in

Supplementary Information, Section II):

274
$$c(x, \infty) = \begin{cases} \dfrac{\rho A}{\Omega} \dfrac{1}{2} e^{-\sqrt{\Omega/D}\,x} & \text{for } x \geq 0, \\ \dfrac{\rho A}{\Omega}\left(1 - \dfrac{1}{2} e^{\sqrt{\Omega/D}\,x}\right) & \text{for } x < 0. \end{cases} \tag{2}$$

From this set of equations, we can see that the concentration c at the border ($x = 0$) is greater for

the half-domain production compared to a single point-source only for the condition where the

density of producers is greater than one point-source within one decay length, $\rho > \sqrt{\Omega/D}$. In other

words, only if the pheromone diffuses far enough, so that the contributions to the gradient from

each single cell overlap.

Further, the response $\theta(x, \infty)$ of the mating pheromone pathway to the steady-state concentration

of pheromone can be modeled with a Hill function. By composing equation (2) with a Hill function

with coefficient n results in:

283
$$\theta(x, \infty) = \begin{cases} \dfrac{1}{1 + \left(C \dfrac{1}{2} e^{-\sqrt{\frac{\Omega}{D}} x} \right)^{-n}} & \text{for } x > 0, \\[4mm] \dfrac{1}{1 + \left[C \left(1 - \dfrac{1}{2} e^{\sqrt{\frac{\Omega}{D}} x} \right) \right]^{-n}} & \text{for } x < 0, \end{cases} \tag{3}$$

where $C \equiv \dfrac{\rho A}{\Omega\, EC_{50}}$ is the concentration of the ligand far away from the border $(x \to -\infty)$ in

"EC_{50}" units, namely, the concentration producing the half-maximal P$_{fus1}$ mediated gene-

expression response.

Experimental responses are measured in unknown concentration units, for this reason they have

been converted to normalized values by baseline-subtracting and normalizing them to their

maximum level before fitting the model (**see Methods**). Equivalently then, equation (3) was

normalized by its maximal value $\hat{\theta}(x, \infty) = \theta(x, \infty) / \dfrac{C^n}{1+C^n}$. This step is fundamental as the non-

normalized Hill-equation only saturates at very large concentrations $(C \gg EC_{50})$, while in all our

experiments we determined C (the pheromone concentration at the curve's plateau) to be of the

same order of magnitude as its EC_{50} (see below), *i.e.* the response curve becomes flat far from the

border in the left domain because of a constant concentration of pheromone, and not because of

saturation. Using this approach, we obtained the theoretical shape for a standard concentration

profile at various values of C and two different values of n (**Fig. 5C,D**). It is well known that no

cooperativity is involved in the gene expression output from the P$_{FUS1}$ promoter[26], we thus fixed

$n = 1$ for gene expression output.

Armed with this model, we next quantified the single-cell outputs of the activity of the mating

pheromone pathway—specifically, cell elongation (**Fig. 5E**; determined manually; **see Methods**)

and P$_{FUS1}$ dependent gene-expression (**Fig. 5F**; determined independently using standard

illumination correction and segmentation tools; **see Methods**)—in *MAT***a** cells lacking Bar1. Our

results revealed a gradient of fluorescence intensity that extends far beyond the region where the

pheromone is generated (**Fig. 5E**). Thus, the transport properties of α-factor can generate responses

at distances spanning hundreds of cell lengths, consistent with a widely diffusible molecule. The

morphological transition (**Figs. 5F**) is steeper than the gene expression response, with cells

increasing their maximal length within a narrower distance range and at a higher pheromone

concentration. This is expected, as the morphological transition is indeed switch-like[10,12].

We ran control experiments using (i) opto-α emitters mixed with the wild-type *MAT***a** reporter

(expressing Bar1) and (ii) a mixture of the wild-type *MATα* as light-independent emitters and the

*MAT***a** *bar1Δ* reporter. To ease cell segmentation, we performed a machine-learning based protocol

(See methods) to obtain a global picture of the behavior of both samples and controls. In the reaction

with wild-type expression of Bar1, basal GFP output was obtained with no detectable differences

between the light and the dark domains. In contrast, in the mix with wild-type *MATα* and the

*MAT**a** bar1Δ* reporter, strong activation was observed in both illuminated and dark domains (**Fig.**

S6). These controls further allowed us to compare the maximum GFP output activation of the opto-

α and wild-type *MATα* strains to confirm that the emission levels are close to physiological in the

half-domain assay, as well as allowing us to correct illumination inhomogeneities and background-

bleaching effects of light induction with an independent method (**Fig. S6, see Methods**).

We also observed that cells elongate, but do not preferentially grow towards (nor away from) the

large-scale gradient, as revealed by the lack of enrichment of cells with orientations close to 0° or

180° respect to the direction of the gradient (**Fig. S7**), demonstrating that—even at its steepest point

(the border)—the large-scale gradient is insufficient to induce chemotropism.

Next, we assessed the stability of the responses over time to check the adequacy of using a steady-

state approximation. We observed that outputs measured in overnight stimulations exhibited a

distance-dependent relationship that was equivalent to the same output measured after 5 hours,

indicating the shape of the gradient is not dramatically altered at later timepoints (**Fig. S8**).

Contrary to that, if no steady state was reached, the gradient would smooth out over time due to

accumulation of the pheromone in the dark domain. In addition, because we did not observe

formation of single or multiple "thin protrusions", characteristic of saturation of the pheromone

pathway[11,31], we concluded that cell responses lie within the sensitive range of the pathway, which

is indicative of the Ste2 receptor not being saturated. This further supports the idea that a steady

state is reached, because otherwise the pheromone would continue accumulating to reach

saturation, at least in its production domain.

Next, we used the biophysical model (**see Supplementary information**) to infer key quantitative

parameters of the gradient. We fitted the baseline-subtracted, normalized-to-maximum

experimental datapoints for the two observables to $\hat{\theta}(x, \infty)$ (**Fig. S9**), with n fixed to one for the

gene expression output (**see Methods**). For completeness of information, the fits are displayed on

top of the raw data (**Fig. 5E,F**). In this way, we inferred the basic properties of the gradients

generated (**Fig. 5G**): the steady-state decay length $\lambda \equiv \sqrt{\frac{D}{\Omega}} \sim 600$ μm, the steady state concentration

of α-factor for $x \ll -\lambda$, $C = 1.6\ EC_{50}$, and the Hill coefficient of the morphological response ($n = 3$).

The gradient value at light-dark transition border, the expected steepest point of the large-scale

gradient, has a value of ~1.2 EC_{50} / mm. Using an EC_{50} value for the response of 1–3 nM[10,11,26], we

obtain a value of 1.5 to 4.5 pM/μm or, roughly, ~20 pM per cell diameter (one cell diameter is ~5μm).

Such shallow gradients will produce differences in receptor occupancy between each side of the

cell an order of magnitude below the minimal ~0.5% observed in chemotropic cells in artificial

pheromone gradients[32].

Collectively, these results imply—in this spatial arrangement and in the absence of Bar1—that: *(i)*

pheromone levels characteristically decay roughly ~100 cell bodies away from producer regions or

single cells; *(ii)* that maximal steady-state pheromone levels found in typical (dense but not

crowded) mating populations are not saturating and rather very similar (less than double) to the

EC_{50} of the response (its most sensitive point); *(iii)* that cells "ignore" (filter out) large-scale

gradients (such as those produced within the spatial limit of finite emitter populations) that are

non-informative about the location of nearby emitters; and *(iv)* although the decay length of the

gradient is the same for the half-domain and a single cell (in one dimension), because the linear

density is ~100 cells/mm (~60 cells in one decay length), $C/2$ (equivalent to the pheromone levels at

the point source) would be reduced by around two orders of magnitude for a single cell. This means

that for a single cell, pheromone levels close to the EC_{50} cannot be reached, even in its immediate

vicinity. This is consistent with the need for close-proximity mechanisms and focal pheromone

emission[7] for such close-contact situations. In other words, single emitters cannot induce elongation

or chemotropism in responding cells, even in the absence of Bar1. Sufficient pheromone levels to

induce mating behavior at a distance are always produced by a collective of cells and under low

Bar1 conditions (biased ratios). Increased rates of degradation of the pheromone (e.g., in the

presence of Bar1) would scale down the gradient (See equation (3) in **Supplementary Information**

section II). These constraints over the maximum levels of pheromone that a single cell can produce

might explain why—when homogeneous Bar1 sources are present—we did not observe elongation

in either of the two domains (**Fig. S6**) and the response values stayed at basal levels, despite the

high density of emitters.

2.5 Quantitative analysis of large-scale gradients produced by opto-Bar1 receivers reveals

localization of Bar1 activity in its production domain.

Our experiments in the absence of Bar1 are adequate to explain natural situations where wild-type

*MAT***a** receivers are in the minority (e.g., upon a mating type switching event), such that Bar1 is not

abundant enough to modify the large-scale pheromone landscape. As the mating-type ratio

becomes unbiased, *MAT***a**-produced Bar1 protease becomes important. When mated under

conditions of pure diffusion-reaction—but not advection—Bar1 is believed to allow reshaping of

the pheromone landscape. The exact nature and biological function of this reshaping activity

during mating remains controversial[6]. Therefore, we used the half-domain assay with the opto-

Bar1 strain in the presence of wild-type MATα cells as emitters to shed light on the problem.

In this case, WT pheromone production is homogeneously distributed and emission of Bar1 is

localized in one half-domain (**Fig. 6A**), on the right of the light-dark transition border (**Fig. 6A, Fig.**

S10). For this assay, wild-type *MATα* emitter cells were tagged with a green nuclear tag, while

opto-Bar1 cells were tagged with a red nuclear tag. This allowed us to confirm that, as in the Petri

dish experiments (**Fig. 3D-F**, **Fig. S4**), mating events (measured as nuclei fusion events) happen in

both illuminated and non-illuminated areas and that Bar1-mediated increases in mating events

correlate with the increased initial growth of *MAT***a** (**Fig. S11**). Crucially, observations of the cell-

elongation phenotype revealed a striking difference compared to this response's profile in the opto-

α experiment (**Fig. 5**): in the opto-Bar1 case, the elongation pattern emerges far away from the

border, well into the dark (left) region of the experiment (**Fig. 6B-C**).

To understand which properties of Bar1 activity and diffusion would yield such a pheromone

profile, we explicitly modeled the contribution of Bar1 to the formation of the gradient profile

(**Supplementary Information, section III**). In this model, both α-factor and Bar1 diffuse, are

produced (in a spatially dependent manner for Bar1) and degrade. These are captured by the two

sets of parameters $(D_\alpha, A_\alpha, \Omega_\alpha)$ and $(D_\beta, A_\beta, \Omega_\beta)$. The enzymatic reaction between Bar1 and α-factor

is modelled through a Michaelis-Menten coupling term that contributes to α-factor degradation:

$$\frac{\partial c_\alpha}{\partial t} = D_\alpha \nabla^2 c_\alpha - \Omega_\alpha c_\alpha - \frac{V_m}{K_M + c_\alpha} c_\beta \cdot c_\alpha + \rho A_\alpha, \quad \text{and} \quad \frac{\partial c_\beta}{\partial t} = D_\beta \nabla^2 c_\beta - \Omega_\beta c_\beta + \rho A_\beta(\text{x}). \quad (4)$$

397 Focusing on the second of the two differential equations and considering production in the right

398 half-domain, the Bar1 spatial profile follows a similar pattern as the α-factor spatial profile in the

399 opto-α experiment. Furthermore, knowing the asymptotic limits for $c_\alpha(x)$ (See **Supplementary**

400 **Information, Section III**), it is then straightforward to numerically solve equation Eq. (4).

401 Importantly, many parameters in Eq. (4) are redundant. Further, Eq. (4) can be simplified by

402 assuming low α-factor concentrations ($c_\alpha < K_M$) reducing the number of parameters to a more

403 meaningful subset. This is justified as the K_M of Bar1 is 30 μM[29], whereas the concentrations of

404 pheromone in our experiment are estimated to be in the nM range, as shown by the opto-α

405 experiment.

406 Thus, we derive four independent dimensional parameters that fully characterize the solution: the

407 decay lengths of α-factor and Bar1 $\lambda_\alpha = \sqrt{D_\alpha/\Omega_\alpha}$ and $\lambda_\beta = \sqrt{D_\beta/\Omega_\beta}$, and the asymptotic constant

408 concentrations of α-factor in the Bar1-free and Bar1-producing domains, $c_\alpha^{max} = \frac{\rho A_\alpha}{\Omega_\alpha}$ and $c_\alpha^{min} =$

409 $\frac{1}{1+\frac{1}{K_M}\frac{V_m}{\Omega_\alpha}\frac{\rho A_\beta}{\Omega_\beta}} \cdot c_\alpha^{max}$. The steady state solution is then composed with the Hill function with

410 coefficient $n = 3$ that yields the elongation response (as in the opto-α experiment), predicting the

411 experimental response curves. Importantly, we measure the c_α^{max} parameter by comparing the

412 morphological response in the opto-Bar1 experiment (**Fig. 6C**) to the one in the opto-α experiment

413 (**Fig 5F**). This is possible since we reach a constant c_α^{max} in our experiments and this value does not

414 depend on any Bar1 parameters according to the theory. By this simple comparison we obtain a

415 lower maximum pheromone level in the opto-Bar1 experiment than in the opto-α experiment,

416 namely $c_\alpha^{max} = 1.20$ EC$_{50}$. This is obtained by taking the fitted value in the opto-α experiment as a

417 reference and considering the Hill coefficient $n = 3$ in the calculation (**see Methods**).

From theory, for any fixed c_α^{max}, it is then possible to determine the location of the half-maximal

normalized response λ_0, in λ_α units, as a function of two adimensional parameters: a measure of

Bar1 activity $c_\alpha^{min} / c_\alpha^{max}$ and a measure of Bar1 diffusion $\lambda_\beta / \lambda_\alpha$. Larger values of Bar1 diffusion

bring the half maximal response to lower negative values (towards the left side) and, similarly,

large values of Bar1 activity can do the same. Then, for each precise location of the half maximal

response we identified a curve of possible parameters in the $(c_\alpha^{min} / c_\alpha^{max}, \lambda_\beta / \lambda_\alpha)$ plane (**Fig 6D, see**

Methods). Consistent with intuition, the plot shows a clear trade-off between the diffusion and

activity of Bar1. Namely, to obtain an equivalent shift, Bar1 with high activity needs to diffuse

narrowly, whereas Bar1 with weak activity needs to diffuse widely. To mirror the opto-α

experiment and find the distance at which the half-maximal response occurs, we fitted a shifted

logistic function with the same steepness n/λ_α obtained in the opto-α experiment fit, leaving the

shift as the only free parameter (**Fig 6C**, **see Methods**). The fit gives us a value of $\lambda_0 = -1.16\,\lambda_\alpha$ for

the half maximal response location (red line in **Fig. 6C,** corresponding to the red line in **Fig. 6D**).

Calculating the diffusion coefficients of Bar1 and α-factor using the Stokes-Einstein equation results

in diffusion rates of 104 μm²/s and 361 μm²/s, respectively[21]. Thus, in theory, Bar1 diffuses about

3.5-fold slower than the pheromone (**Fig 6D**) (assuming similar degradation rates for both). This

value corresponds to a value of $c_\alpha^{min} / c_\alpha^{max}$ of 1/10 to 1/20. We then compared this expected result

to known measures of Bar1 activity to then estimate the protease's effective diffusion. For this we

made an estimation based on pure α-factor dose-response experiments where we identified the

growth arrest switch, comparing *MAT***a** wild type and its corresponding Bar1 knockout at an

equivalent cell density as in the half-domain assay (**Fig. S12**). Our estimate tells us that, under these

conditions, Bar1 reduces the levels of pheromone by at least ~100-fold, displacing the arrest switch

from its position which lies very close to the Ste2 receptor Kd (**Fig. S12**). According to our theory

(**Fig. 6C**), the ratio of the expected decay lengths for α-factor and Bar1 is $\frac{\lambda_\alpha}{\lambda_\beta} \sim 0.1$. This means that,

at the very least, Bar1 diffusion is 10-fold slower than pheromone diffusion (assuming equal

degradation rates for both). Thus, we conclude from this experiment that Bar1 tends to stay in its

production domain through mechanisms altering diffusion during mating, in contrast to α-factor,

which diffuses widely.

3. Discussion

Here, we constructed yeast strains in which pheromone production and degradation can be

controlled via optogenetics to shape extracellular signal gradients and thereby provide a better

understanding of the functions of cell-cell communication in open space. Using this method, we

show that the pheromone landscapes in sparse yeast populations are shaped by a "global excitation

- local degradation" process**,** where α-factor functions as global excitatory signal: biologically

relevant concentrations of this pheromone are built up by the contributions of many emitter cells

and receiver cells act as sinks for this signal.

We demonstrate that there is a relevant length-scale for gradient formation other than that in the

immediate vicinity of cells (1 cell body or ~5 μm); namely, the large-scale gradient formed at the

edge of finite producer populations. However, these large-scale gradients are not steep enough to

induce chemotropism (despite pheromone concentrations being high enough to trigger

elongation). Indeed, gradient values capable of inducing chemotropic behavior in the literature are

generally steep[7,13,14,33,34]. Chemotropism directed along large-scale gradients was not observed in

our half-domain assays. Thus, *MAT*a cells filter out large-scale gradients while, presumably,

tracking local small-scale gradients, given the local gradients possess sufficient steepness. Our

model allowed us to infer that the concentrations of pheromone emanating from single emitters

must be well below the mating threshold (**Fig 5H**), this means that pheromone secretion by single

cells cannot not provide positional information, because responders are not sensitive enough to

activate chemotropism (**Fig. 6E**)—even sensitized if partners are sensitized (without expressing

Bar1).

Our results also show that Bar1 diffuses at least 10-fold slower than the pheromone it degrades (**Fig**

6D,F). Such limited diffusion of Bar1 does not prevent *MAT***a** from modifying global pheromone

profiles—provided cells are at a sufficient cell density. Indeed, in our opto-Bar1 experiment (**Fig.**

6A-C), Bar1 greatly reduces the global α-factor concentration, and *MAT***a** cells do not elongate nor

arrest (**Fig S6**). Despite this, cells in close contact still mate normally (**Fig. S11**). Thus, when

separated in space, cells cannot reach the threshold for executing the morphological transition. This

is in agreement with the absence of elongation and the persistence of cell division by *MAT***a** cells in

unbiased mating reactions in the presence of advection[35]. In this sub-threshold case, it is likely that

increased cell-number, randomly oriented bipolar budding[35,36] and close-contact mechanisms[7,37] aid

mating (**Fig. 6G**). These known mechanisms are functionally very different as search strategies to

chemotropism, a function first inferred from the classical experiments by Segall in the early 1990s[34],

in which cell-cycle arrested cells in agarose extend their cell bodies towards a glass pipet that

releases pheromone. This type of chemotropism, involving cell elongation and cell-cycle arrest, is

only accurate at concentrations matching those at which cells switch[38–41]. Therefore, gradients need

to fulfill two conditions to induce this behavior: they must have a high mean pheromone

concentration to induce elongation and need to be steep enough to direct the polarisome towards

the source. Our results suggest that—in mating reactions with cell densities that would render

chemotropic cell elongation useful—there is a trade-off between the mean pheromone

concentration and the steepness of the gradient: cells are weak producers but, in principle, good

position signalers, whereas dense populations are strong signalers but weak position signalers.

Thus, our results explain why elongated *MAT***a** chemotropism is rarely observed as directed mate-

searching in mating reactions with sparse populations. Indeed, depending on the levels of Bar1,

within environments crowded with α-factor emitters, cells will either restrict pheromone levels to

the emitter vicinity and rely on local mechanisms[7,37] or exhibit random elongation due to gradient

overlapping (**Fig. 6H**). Random cell elongation could be a useful strategy under such low Bar1

conditions, as gradient overlapping will make the switch concentration proportional to the cell

density of partners, rather than their distances. Our results reveal that gradient overlapping is

significant in the absence of Bar1, implying that distance-to-source measurements by cells in the

production region cannot be disentangled from demographic measurements. Therefore, our results

highlight a limitation of chemotropism as an efficient mate-directed search strategy; namely, the

necessity of the *MAT*α-biased ratios for *MAT***a** to reach its switching threshold entails an increased

likelihood of producing local, uninformative gradients caused by increased overlapping. This

trade-off, in principle, could be balanced at some specific ratio, where Bar1 levels are enough to

spatially restrict levels of α-factor and sources are clustered to allow a high local concentration of

α-factor. This could be the case, for example, for a large microcolony of MATα encountering a close

smaller microcolony of *MAT***a** (e.g., microcolony growth after mating type switching events or

spore dispersal[42,43]).

Rappaport and Barkai[19] defined a theory for Bar1 as a perfect (or nearly perfect) α-factor sink where

most α-factor molecules that impinge upon *MAT***a** cells are absorbed, rather than released back into

the medium. This process—Bar1 dependent degradation—contrary to other mechanisms, such as

receptor endocytosis-mediated internalization, allows both absorption at the cell surface and

degradation in the media. In our experiments, when gradients were formed by the pheromone-

degrading enzyme Bar1 secreted by *MAT***a** under uniform α-factor production throughout space,

the response patterns revealed that the extent of Bar1 diffusion during the timescales relevant for

mating are very low. The evidence for Bar1 being a widely diffusible protein came from an early

study by Hicks and Herkowitz[23], who developed the "N-assay" in which α-factor gradients are

generated in agar with a heavy streak of *MAT*α cells and "tester" *MAT***a** cells are micromanipulated

and placed at different distances from the source to estimate the concentration of α-factor. When

the assay is performed on agar where a heavy streak of wild-type *MAT***a** (producing Bar1) was

allowed to stay on the opposite side of the agar for six hours and then cut out, the levels of

stimulation reduced compared to the fresh, untreated agar. Several possibilities could explain this

inconsistency. First, in the assay by Hicks and Herkowitz, Bar1 may have diffused poorly, staying

in the region close to the cut fraction, but reshaped α-factor gradients by providing a sink region—

similarly to our experiment. Second, because our assay retains physiological conditions both in and

out of the Bar1 production half-domain, the cell density of both emitters and receivers is high

(unlike in the "N-assay"). Therefore, cells could effectively lower Bar1 mobility by providing an

adsorption surface based on cell-wall mannan interactions. Indeed, Bar1 is an abundant disulfide-

bridged mannoprotein present in the cell wall of *MAT***a** cells[44] and this sink character is essential to

build models that explain the establishment of gradient disentanglement from two sources in close-

proximity intra-ascus interactions[19] and mutual-avoidance in *MAT***a** cells[20]. The latter of these

mechanisms is relevant to germinating spores during outcrossing in ecologically realistic

scenarios[43,45] and requires similar conditions as in our experiment. The cell wall-associated fraction

of Bar1 allows it to work as a sink for α-factor while retaining the capacity of the enzyme to diffuse.

Overall, our results suggest that the pheromone landscapes in "open" mating reactions are directly

shaped by population parameters and differential diffusion of Bar1 and α-factor. Functionally,

steep gradients localized at either the source or the sink can only guide chemotropism when the

population ratios are biased enough to limit global Bar1 availability. Due to its sink function, the

gradients might be steeper at a receiver cell; thus, if several receiver cells cluster, the gradients can

direct these cells away from each other[20]. Thus, gradients can effectively provide information about

the location of competitors (other *MAT***a** cells) rather than potential mates (*MAT*α cells). In addition,

local degradation of α-factor allows the asexual proliferation of unmated individuals, maximizing

the total population's fitness.

Our results open the door for creating more complex spatial patterns and the dynamic control of

pheromone production and degradation simultaneously, by diversifying the types of optogenetic

control possible. Such complexification will allow to simulate natural situations—e.g., optogenetic

induction of mating type switching—and to create synthetic communication landscapes with

useful applications in engineered living materials. Our study encourages the use of the yeast

mating system as a quantitative framework for the design and control of spatial behavior of natural

and synthetic cell-cell communication systems.

4. Methods

547   All materials described here and in the Supplementary Information are available upon reasonable

548   request. Datasets used to generate the figures and the code for the mathematical modelling can be

549   found on the public Zenodo archive 10.5281/zenodo.10572403

4.1 Strains and growth conditions

S. cerevisiae strains used in this study (**Supplementary Table 1**) are derivatives of BY4741 (*MAT***a**

his3Δ1 leu2Δ0 met15Δ0), BY4742 (MATα, *lys2Δ0,* otherwise identical to BY4741), SEY6210a[46] (*MAT***a**

leu2-3,112 ura3-52 his3Δ200 trp1Δ901 lys2-801 suc2Δ9), and SEY6210 (MATα, otherwise identical to

SEY6210a). Yeast cells were routinely grown on complete medium (Yeast peptone dextrose; YPD,

Sigma) or synthetic complete (SC) medium—composed of 6.7 g Yeast Nitrogen Base without amino

acids (Difco 291940) and 0.8 g complete supplement mixture drop-out (Formedium DCS0019) in 1

L—supplemented with 2% glucose (Merck). Cells from glycerol stocks or agar plates were

inoculated into YPD and incubated at 30°C on an orbital shaker at 250 rpm (in an Innova 4230

incubator) for 12 to 16 h. Day cultures were performed by inoculating 5 mL of SC or YPD media

with overnight cultures at a concentration of 1% *v/v*. Care was taken to reduce unwanted light

exposure as much as possible before the start of the experiments by covering the agar plates and

precultures in aluminum foil and by performing all steps without direct exposure to light (low

ambient light).

All yeast transformations were conducted following a standard lithium acetate protocol. To replace

the native promoters located upstream of the genes of interest with the EL222-dependent promoter

P_{C120}, we performed CRISPR/Cas9-based genome editing following a method described

previously[47]. Oligonucleotides used in this study are listed in **Supplementary Table 2**. The

replacement fragments used to exchange the upstream region of the relevant genes (pheromones

or Bar1) were amplified from plasmid pYTK097, which contains the original sequence of the wild-

type P_{C120} promoter from *Erythrobacter litoralis*[48], using primers containing appropriate homology

regions as overhangs. DNA sequences coding for guide-RNA (gRNA) sequences (obtained from

IDT) used to target Cas9 to the desired loci are also listed in **Supplementary Table 2**. Short and

long gRNA were hybridized with each other and ligated into plasmid pML104, which possesses a

Cas9 expression cassette and an URA3 marker for auxotrophic selection. The EL222 transcription

factor was inserted at the HIS3 locus by digesting plasmid pEZ-L105[49] with the *SfiI* restriction

enzyme. In this construct, EL222 is expressed from the constitutive P_{pgk1} promoter. Nuclear tagging

was performed by inserting mApple at the HTB2 locus by amplifying mApple from plasmid

pPH330 using primers oPH_0386 and oPH_0387 followed by transformation and selection using

G418 antibiotic selection. Nuclear tagging with mVenus was performed by replacing mApple in

the strain above using CRISPR/Cas9. The mVenus fragment was amplified from plasmid pPH_086

using the same oPH_0944 and oPH_0945 primers and guideRNAs oPH_941 and oPH_942.

4.3 Light stimulation experiments in 24-well-plates

Overnight cultures were diluted 1:50 in fresh LoFlo-SC (synthetic complete media prepared with

"low fluorescence" yeast nitrogen base lacking folic acid and riboflavin; Formedium) to reduce

media fluorescence and grown to an optical density (OD_{600}) of 0.4–0.5. Individual day cultures of

optogenetic and reporter strains were washed once with 1 volume of LoFlo-SC media, resuspended

to an OD of 0.5, mixed at a 1:1 ratio, and placed in black glass-bottom 24-well plates (CellVis). Light

stimulation was performed in a custom-made light plate apparatus based on the design by

Gerhardt et al[27] equipped with a blue LED (λ=461 nm). The final volume in all experiments was 1

mL and the incubation time was 2 h. When opto-Bar1 was used as a test strain, α-factor (Sigma)

was added at a final concentration of 1 μM. A well containing only the reporter strain was routinely

included to calculate the basal GFP expression from the P_{C120} promoter. Light intensity calibration

of the apparatus was performed as reported before[27] using a power-meter and a custom 3D-printed

adaptor. Reporter gene expression was analyzed by flow cytometry immediately after sampling.

4.4 Patterned light stimulation in Petri plates

Day cultures in SD media containing all amino acids were inoculated at 4% *v/v* with an overnight

culture and grown until the OD_{600} was ~1.0The cultures were then centrifuged at 3500 rpm for 7

min, the supernatant removed, the pellets washed once in SD medium containing all amino acids,

and the optical density adjusted to OD_{600} =1.0. These suspensions were mixed at a 1:1 ratio and 150

μL was added to 6 mL of 0.5% agar ("top" agar, kept in a water bath at 55°C).

For the "velvet" version of the assay (**Figs. 3, S4, S5, S7**), the top agar was poured over Petri plates

containing solid SC-agar (1.5% agar; 20 mL total volume) supplemented with all amino acids.,

using a leveling surface. Plates were then sealed with parafilm, keeping the lids of the plates

slightly tilted. Blue light patterning was performed with a DMD projector (DLP® LightCrafter™

4500 TexasInstrument) placed inside an incubator (Memmert UF160); the projector projects light of

the desired shape and intensity over Petri plates placed over a black surface. After light stimulation,

the plates were replicated on SD plates lacking methionine and lysine using the sterile velvet

technique, incubated for at least 24 h at 30°C, and imaged using a multi-imager (Biorad) for colony

counting.

For the "direct" version of the assay (**Figs. 2, S4, S6**), agar cell suspension was poured over 20 mL

of 1.5% agar SD media with low lysin levels (0.005 g/L) and was directly recovered for imaging and

colony counting after at least 24 h of incubation (**Fig. S4**). The basis for this simpler version of the

assay relies on the following: because the *MAT*a and *MAT*α wild-type strains are auxotrophic for

methionine and lysine, respectively (**See Supplementary Table 1**), we expected haploid growth—

and presumably mating—to depend on supplementation of these amino acids. Consequently, a

low concentration of these amino acids would allow mating, but later only select for diploid growth

as these amino acids become exhausted. We note that the methionine levels in the 150 μL cell mixes

are enough to allow mating, as adding methionine did not increase diploid growth. On the

contrary, lysine levels were optimized to determine the final concentration used (levels higher than

0.005 g/L did not affect diploid growth). Calibration of the light intensity of the device was

performed by placing a power meter at the location of the agar surface.

4.5 Half-domain gradient generation assay

Rows of wells in glass-bottom 24-well plates (CellVis) were taped in the "row" direction following

available marks in the frame that coincide with the hemisphere of the wells. Longitudinal sections

of black tape were cut and the uncut side of the tape was gently positioned facing the hemisphere

of the wells. Overnight cultures were reinoculated at 2% *v/v* in YPD. These day cultures were grown

for 4 h, washed once in 4 mL of synthetic complete media with 2% glc (SC-GLC), and then

resuspended in 1 mL of the same media. The OD600 was determined in a cuvette benchtop

spectrophotometer using 100 µL of this culture and 900 µL of SD-glc. Cultures were diluted back

to OD_{600}=0.1 in a final volume of 5 mL **SC-glc**. Concanavalin A coating was performed as follows:

wells were exposed to 300 µL of 0.2 mg/mL Concanavalin A (ConA, Sigma; 1/10 dilution of a 2

mg/mL stock that was spun down to remove protein aggregates) for 6 minutes, then the solution

was removed from each well (using a the P1000 pipet and then a P200 for the remanent), washed

once with 300 µL of sterile water (water was removed in one strong inversion movement and not

pipetted out), then 300 µL of the appropriate cell suspension mixes were gently deposited

(consistently on the west side of each well) and allowed to settle gravitationally for 35 min. During

cell settlement, fresh 1.5% *w/v* low-melting point agarose (Sigma) was melted in SD-glc in a glass

beaker with constant and gentle magnetic stirring. Then, 200 µL of the hot (70°C) agarose solution

was taken out (reverse pipetted) with a 1000 µL (blue) tip, held for 10 seconds, and the solution

was gently (5 seconds to dispense 200 µL) was deposited in the bottom "west" position,

immediately followed by gently pipetting up and down (close to the middle high, also pointing in

the west direction). We set a light exposure time of 5 h, which allowed us to observe complete

development of the output curves. The total analyzed region was roughly 5.4 mm around the

border. Wells have a diameter of 1.554 cm and cells are embedded in a total volume of 0.5 cm^3 of

agarose.

4.6 Flow cytometry

Flow cytometry measurements were performed on an LSR II cytometer (Becton-Dickinson). GFP

fluorescence was measured with a 488 nm laser using BD FACS DIVA software (BD Biosciences).

First, cells were gated in an FSC-A/SSC- to exclude debris and doublets. GFP-positive cells (either

*MAT***a** or *MATα*, depending on the experiment) were gated manually using FlowJo (Becton-

Dickinson), and the mean fluorescence of this population was used for further analysis.

4.7 Fluorescence microscopy

Imaging of the half-domain assay was performed using a Leica DMi8 inverted microscope (Leica

Microsystems) equipped with a motorized stage and a 20x objective (N.A. 0.4) using *LasX* software

(Leica Microsystems) in multi-position mode. Fluorescence images were acquired using the 470/40

(Ex), 525/50 (EM) filter for GFP and the TXR filter 560/40 (EX), 630/75 (EM) filter for mApple

proteins, respectively. Imaging was also performed using an inverted Olympus IX81

epifluorescence microscope equipped with an "Xcite exacte" light source (I = 30%) and a filter cube

(EX) 514 nm/10 (EM) 545 nm/40 (49905 – ET) with a 1000 ms exposure time.

4.8 Manual determination of cell-length

As a proxy for the morphological response to pheromone we manually measured the cell-length

for cells in the half-domain experiment. In the opto-α experiment, a total of 6995 cells were

segmented, while in the oplto-Bar1 experiment the total was 8331 cells (a larger region was sampled

in this experiment). In the range considered for fits the total numbers analyzed correspond to 6686

and 6088 cells, respectively. Lengths were measured manually using the segmented line tool in Fiji

in single cells. On each identified cell, we measured its length by drawing a line along its longest

diameter. In elongated cells we draw their central longitudinal axis using as many points as needed

to maximize precision (elongated cells are curved). Although we counted mostly non-overlapping

cells, if a mild overlap existed, or the overlap produced clearly distinguishable boundaries, cells

were considered. Prezygotes (cells having a characteristic dumbbell shape, product of cell-cell

fusion events), were not considered in the analysis. In samples with large cell-clumps, we measured

regions close to empty spaces where little overlap existed. All cell segmentation data and original

images are available in Zenodo.

4.9 Machine-learning based cell segmentation.

Manual segmentation is very precise but laborious. We therefore complemented manual

segmentation analyses using machine-learning algorithms to obtain additional output data and

analyze control half-domain assay experiments. All datasets were pre-processed using rotation,

noise insertion, pixel intensity perturbation and translation. Cell segmentation was done on

brightfield images on the half-domain assay with two machine-learning algorithms using the U-

NET architecture. The first model was used to segment elongated cells and the second model to

segment round-shaped cells. The first model was trained on 21 images during 50 epochs. Its

accuracy was estimated at 0.97 and its recall at 0.89 (Fig. S11). The second model was trained on 10

images during 100 epochs and had an estimated accuracy of 0.91 and a recall of 0.73. All the training

was performed using a GPU NVIDIA Quadro RTX 4000 (7.5GB RAM memory).

After identifying cells as such, round-shaped ones were assigned the category of "receivers" or

"emitters" using fluorescence image information. Fluorescently labeled nuclei were also segmented

using machine learning. A unique model (same for mApple tagged nuclei and for GFP-tagged

nuclei) was trained (with GFP images but as well valid for RFP segmentation) over 30 epochs and

its accuracy was estimated to be 0.98.

In the opto-α generated gradient (Fig. 5), emitters carry a mApple (red) nuclear marker, whereas

receivers do not carry a nuclear marker. Cells identified as round with overlapping nuclear

segmentation are identified as emitters. In the opto-Bar1-generated gradient (Fig. 6), emitters carry

a GFP nuclear marker and were identified equivalently. Once all cells were segmented and the

receivers (elongated and round shapes) are identified from the emitters, a CSV file that contains

information about the identity of the segmented object (receiver or emitter), its position along the

experimental axis (in µm), area (in pixels), and the mean GFP intensity in the case of fluorescent

reporter receivers is produced (Fig. 5). For the latter, background subtraction was performed as

follows: a mask was produced that is the sum of the two segmentations from the models for round

and elongated shapes. The background then corresponds to the negative part of this mask in the

brightfield image. To calculate the orientation of elongated objects, an ellipse was fitted using

Python openCV ellipse fitting function and its angle (from 0° excluded to 180°) with respect to the

x-axis was extracted (Fig. S)

The data from the nuclei segmentation was used to count haploid and diploids (independently of

the shape analysis, which filters cells) as a measure of cell proliferation along the experimental

length in the half-domain assay (**Fig. S13**). Counting nuclei allowed us to count individual cells in

regions with clumps. It also allowed the easy identification of diploid cells events in the opto-bar1

experiment (**Fig. S13**).

5.10 Quantification of the P*FUS1* transcriptional response in single cells.

Quantification of the P*FUS1*-GFP response in the half domain assay was more precisely determined

segmenting images in the GFP channel, where false positives are minimal. First the images were

corrected for illumination inhomogeneities using the BaSiC package for imageJ[50]. The background

profile obtained using this tool was equivalent to the background extraction done by ML but

performs better in the smaller lengthscale (within-image inhomogeneities). The Cell Profiler

software (Broad Institute, Cambridge, MA) was used for the quantification of the transcriptional

response. The OTSU adaptive thresholding method was used for object identification in

fluorescence images. Cell clumps were discarded with an object-size threshold and a form-factor

filter to select rounder objects. Segmentation quality was inspected visually and empirically

optimized by changing filter and threshold values. All cell segmentation data and original images

are available in Zenodo.

5.11 Data analysis

Normalization and baseline subtraction of data. Cell elongation data as well as P*FUS1* dependent

gene-expression at single cell level were provided as inputs to the downstream analysis. The

baseline of each dataset was identified as their mean value for distances to the border greater then

2700 μm on the right domain, at such distances the gradients are zero and the dataset can be

considered to having reached their asymptotic levels. After subtracting the baseline, the scaling

factor was obtained by measuring their mean value for distances to the border between 2000 and

1300 μm on the left domain, at such distances the gradients are zero and the datasets doesn't show

border effects due to reflective boundary conditions present for larger distances. Normalization

was obtained by dividing the baseline subtracted data by the scaling factor.

Fitting pheromone gradient generated by the spatial optogenetic induction of α-factor. Due to

difference in filtering, cell elongation data as well as P*FUS1* dependent gene-expression at single cell

level had to be matched. This was done by binning the data in 100 μm wide bins and taking their

mean for the downstream analysis. Fitting was done using the least squares methods on both

datasets at the same time, to extract the parameters λ, C and n by fitting gene expression to $\hat{\theta}(x, \infty)$

with Hill coefficient equal to 1 and elongation to $\hat{\theta}(x, \infty)$ with free Hill coefficient n. Only integer

values between 1 and 7 were considered in the least squares minimization procedure for the n

parameter.

Estimation of α-factor maximum concentration on the left domain c_α^{max} in the spatial optogenetic

induction of Bar1. Cell elongation data for the Bar1 induction experiment were binned and

normalized according to the same procedure used for the α-factor induction results. The

experimental data indicated a scaling factor for the Bar1 induction experiment equal to 5.75

compared to 7.20 measured in the α-factor induction experiment. Considering the measurement of

$C = 1.57 \, EC_{50}$ obtained in the α-factor induction experiment by fitting, we extracted the value of

c_α^{max} in the Bar1 induction experiment by numerically solving the following equation:

$$\frac{c_\alpha^{max^3}}{1 + c_\alpha^{max^3}} = \frac{5.75}{7.20} \cdot \frac{C^3}{1 + C^3} \tag{7}$$

746   obtaining $c_\alpha^{max} = 1.20 \, EC_{50}$.

747   *Determination of the half-maximal response in the spatial optogenetic induction of Bar1.* In order

748   to measure the distance at which the half-maximal response in the Bar1 dataset occurs, we fitted

749 the following logistic function with one parameter, the half-maximal response λ_0, to the normalized

750 baseline-subtracted data:

$$\frac{1}{1 + e^{\frac{n}{\lambda_\alpha}(x-\lambda_0)}}, \tag{8}$$

752 finding a value of $\lambda_0 = -1.16\,\lambda_\alpha$.

5.12 Numerical derivation of the half-maximal response in function of model parameters

754 In order to derive the steady state α-factor concentration profile $c_\alpha(x)$ from **Eq. 4**, we numerically

755 solved the following differential equation:

$$\frac{\lambda_\alpha^2}{c_\alpha^{max}} c_\alpha''(x) - \frac{1}{c_\alpha^{max}} c_\alpha(x) - \frac{c_\alpha^{max} - c_\alpha^{min}}{c_\alpha^{max} c_\alpha^{min}} c_\beta(x) \cdot c_\alpha(x) + 1 = 0, \tag{8}$$

757 using the steady state analytical solution to **Eq. 4** for the shape of the Bar1 concentration profile:

$$c_\beta(x) = \begin{cases} \dfrac{1}{2} e^{x/\lambda_\beta} & \text{for } x < 0, \\ \left(1 - \dfrac{1}{2} e^{-x/\lambda_\beta}\right) & \text{for } x \geq 0, \end{cases} \tag{9}$$

and conveniently setting $\lambda_\alpha = 1$, such as to have all spatial variables (x and λ_β) expressed in λ_α

units, and $c_\alpha^{max} = 1.20\,EC_{50}$ to its experimentally determined value.

We then derived numerical solutions for $c_\alpha(x)$ using the Mathematica software package and the

NDSolveValue function with Dirichlet boundary conditions c_α^{min} and c_α^{max}, with a large set of

parameters c_α^{min} and λ_β distributed equally spaced on the logarithmic scale.

For each of these solutions, we obtained the corresponding $\theta(x, \infty)$ by combining them with the

Hill function with coefficient $n = 3$. We further calculated the theoretical location of the half

maximal response λ_0 by taking the point where $\theta(x, \infty)$ crosses the value corresponding to the

mean between its minimum and its maximum values over the full domain. The resulting table of

λ_0 in function of $\lambda_\beta/\lambda_\alpha$ and $c_\alpha^{min}/c_\alpha^{max}$ was fed to the ListContourPlot function to plot **Fig 6C**. We

used the same function to highlight the contour corresponding to the experimentally determined

$\lambda_0 = -1.16\,\lambda_\alpha$ in dashed red, highlighting the set of values $\lambda_\beta/\lambda_\alpha$ and $c_\alpha^{min}/c_\alpha^{max}$ that are

compatible with the experimental data.

Acknowledgments

773   The authors would like to thank their team members for their critical reading of this manuscript.

774   This work was mainly supported by the European Research Council grant SmartCells (724813) and

775   received additional support from grants ANR-11-LABX-0038, ANR-10-IDEX-0001-02, ANR-16-

776   CE12-0025-01, and the QLife Institute ANR-17-CONV-0005

Author Contributions

778   AB conceived the research and contributed to all aspects of this work. MH, CE, SP, CC, and MLB

779   contributed to optogenetic strain construction (MH, CE), experiments (MH, CE, CC), and setup of

780   the optogenetic devices (MH, SP, MLB). LC contributed image analysis. VS contributed the

781   modeling of the pheromone reaction-diffusion dynamics and their fit to experimental data. AB, VS,

782   and PH interpreted the results. AB and PH designed the study. AB, VS and PH wrote the article.

Supplementary Information

Supplementary information contains 12 figures, the mathematical model, and two tables.

Bibliography

1.   Simsek, M. F. & Özbudak, E. M. Patterning principles of morphogen gradients. *Open Biol.* **12**, 220224 (2022).

2.   Stapornwongkul, K. S. & Vincent, J.-P. Generation of extracellular morphogen gradients: the case for diffusion. *Nat. Rev. Genet.* **22**, 393–411 (2021).

3.   Mukherjee, S. & Bassler, B. L. Bacterial quorum sensing in complex and dynamically changing environments. *Nat. Rev. Microbiol.* **17**, 371–382 (2019).

4.   van Gestel, J. *et al.* Short-range quorum sensing controls horizontal gene transfer at micron scale in bacterial communities. *Nat. Commun.* **12**, 2324 (2021).

5.   Narla, A. V., Borenstein, D. B. & Wingreen, N. S. A biophysical limit for quorum sensing in biofilms. *Proc. Natl. Acad. Sci.* **118**, e2022818118 (2021).

6.   Clark-Cotton, M. R., Jacobs, K. C. & Lew, D. J. Chemotropism and Cell-Cell Fusion in Fungi. *Microbiol. Mol. Biol. Rev.* **86**, e00165-21 (2022).

7.   Ghose, D., Jacobs, K., Ramirez, S., Elston, T. & Lew, D. Chemotactic movement of a polarity site enables yeast cells to find their mates. *Proc Natl Acad Sci U A* **118**, (2021).

8.   Bardwell, L. A walk-through of the yeast mating pheromone response pathway. *Peptides* **26**, 339–350 (2005).

9.   Shellhammer, J. P. *et al.* Quantitative analysis of the yeast pheromone pathway. *Yeast* **36**,

495–518 (2019).

10.   Yu, R. C. *et al.* Negative feedback that improves information transmission in yeast

signalling. *Nature* **456**, 755–61 (2008).

11.   Muller, N. *et al.* A Predictive Model for Yeast Cell Polarization in Pheromone Gradients.

PLOS Comput. Biol. **12**, e1004795 (2016).

12.   Malleshaiah, M. K., Shahrezaei, V., Swain, P. S. & Michnick, S. W. The scaffold protein Ste5

directly controls a switch-like mating decision in yeast. *Nature* **465**, 101–105 (2010).

13.   Brett, M.-E., DeFlorio, R., Stone, D. E. & Eddington, D. T. A microfluidic device that forms

and redirects pheromone gradients to study chemotropism in yeast. *Lab on a Chip* vol. 12 3127

(2012).

14.   Moore, T. I., Chou, C.-S., Nie, Q., Jeon, N. L. & Yi, T.-M. Robust spatial sensing of mating

pheromone gradients by yeast cells. *PLoS One* **3**, e3865 (2008).

15.   Barkai, N., Rose, M. D. & Wingreen, N. S. Protease helps yeast find mating partners. *Nature*

396, 422–423 (1998).

16.   Hanson, S. J. & Wolfe, K. H. An Evolutionary Perspective on Yeast Mating-Type Switching.

Genetics **206**, 9–32 (2017).

17.   Jackson, C. L. & Hartwell, L. H. Courtship in S. cerevisiae: both cell types choose mating

partners by responding to the strongest pheromone signal. *Cell* **63**, 1039–1051 (1990).

18.   Barkai, N., Rose, M. D. & Wingreen, N. S. Protease helps yeast find mating partners. *Nature*

396, 422–423 (1998).

19.   Rappaport, N. & Barkai, N. Disentangling signaling gradients generated by equivalent

sources. *J. Biol. Phys.* **38**, 267–278 (2012).

20.   Jin, M. *et al.* Yeast Dynamically Modify Their Environment to Achieve Better Mating

Efficiency. *Sci Signal* **4**, ra54–ra54 (2011).

21.   Diener, C. *et al.* Yeast mating and image-based quantification of spatial pattern formation.

PLoS Comput Biol **10**, e1003690 (2014).

22.   Vo, T., Shah, S. B., Choy, J. S. & Luo, X. Chemotropism among populations of yeast cells

with spatiotemporal resolution in a biofabricated microfluidic platform. *Biomicrofluidics* vol. 14

014108 (2020).

23.   Hicks, J. B. & Herskowitz, I. Evidence for a new diffusible element of mating pheromones

in yeast. *Nature* **260**, 246–248 (1976).

24.   Rojas, V. & Larrondo, L. F. Coupling Cell Communication and Optogenetics:

Implementation of a Light-Inducible Intercellular System in Yeast. *ACS Synth. Biol.* **12**, 71–82 (2022).

25.   Motta-Mena, L. B. *et al.* An optogenetic gene expression system with rapid activation and

deactivation kinetics. *Nat. Chem. Biol.* **10**, 196–202 (2014).

26.   Banderas, A., Koltai, M., Anders, A. & Sourjik, V. Sensory input attenuation allows

predictive sexual response in yeast. *Nat Commun* **7**, 12590 (2016).

27.   Gerhardt, K. P. *et al.* An open-hardware platform for optogenetics and photobiology. *Sci*

Rep **6**, 35363 (2016).

28.   Le Bec, M. *et al.* Optogenetic spatial patterning of cooperation in yeast populations. *Nat.*

Commun. **15**, 75 (2024).

29.   Nath, R. Properties of Barrier, a novel Saccharomyces cerevisiae acid protease. *Biochimie* **75**,

467–472 (1993).

30.   Kawada, D. *et al.* The yeast Arf-GAP Glo3p is required for the endocytic recycling of cell

surface proteins. *Biochim. Biophys. Acta* **1853**, 144–156 (2015).

31.   Hilioti, Z. *et al.* Oscillatory phosphorylation of yeast Fus3 MAP kinase controls periodic

gene expression and morphogenesis. *Curr Biol* **18**, 1700–1706 (2008).

32.   Arkowitz, R. a. Chemical gradients and chemotropism in yeast. *CSH Perspect Biol* **1**, a001958

(2009).

33.   Paliwal, S. *et al.* MAPK-mediated bimodal gene expression and adaptive gradient sensing

in yeast. *Nature* **446**, 46–51 (2007).

34.   Segall, J. E. Polarization of yeast cells in spatial gradients of alpha mating factor. *Proc. Natl.*

Acad. Sci. U. S. A. **90**, 8332–6 (1993).

35.   Anders, A., Colin, R., Banderas, A. & Sourjik, V. Asymmetric mating behavior of isogamous

budding yeast. *Sci. Adv.* **7**, eabf8404 (2021).

36.   Erdman, S. & Snyder, M. A filamentous growth response mediated by the yeast mating

pathway. *Genetics* **159**, 919–28 (2001).

37.   Clark-Cotton, M. R. *et al.* Exploratory polarization facilitates mating partner selection in.

Mol Biol Cell **32**, 1048–1063 (2021).

38.   Follette, P. J. & Arkowitz, R. A. Chemotropism During Yeast Mating. *Methods Mol. Biol.* **571**,

99–110 (2009).

39.   Ismael, A. & Stone, D. E. Yeast chemotropism: A paradigm shift in chemical gradient

sensing. *Cellular Logistics* vol. 7 e1314237 (2017).

40.   Stone, D. E. & Arkowitz, R. A. In Situ Assays of Chemotropism During Yeast Mating.

Methods in Molecular Biology 1–12 (2016).

41.   Arkowitz, R. A. Responding to attraction: chemotaxis and chemotropism in Dictyostelium

and yeast. *Trends Cell Biol.* **9**, 20–27 (1999).

42.   Knop, M. Evolution of the hemiascomycete yeasts: on life styles and the importance of

inbreeding. *BioEssays* **28**, 696–708 (2006).

43.   Reuter, M., Bell, G. & Greig, D. Increased outbreeding in yeast in response to dispersal by

an insect vector. *Curr Biol* **17**, R81–R83 (2007).

44.   Moukadiri, I., Jaafar, L. & Zueco, J. Identification of two mannoproteins released from cell

walls of a Saccharomyces cerevisiae mnn1 mnn9 double mutant by reducing agents. *J Bacteriol* **181**,

4741–4745 (1999).

45.   Murphy, H. a & Zeyl, C. W. Yeast sex: surprisingly high rates of outcrossing between asci.

PloS One **5**, e10461 (2010).

46.   Robinson, J. S., Klionsky, D. J., Banta, L. M. & Emr, S. D. Protein sorting in Saccharomyces

cerevisiae: isolation of mutants defective in the delivery and processing of multiple vacuolar

hydrolases. *Mol Cell Biol* **8**, 4936–4948 (1988).

47.   Laughery, M. F. *et al.* New vectors for simple and streamlined CRISPR-Cas9 genome editing

in Saccharomyces cerevisiae. *Yeast Chichester Engl.* **32**, 711–720 (2015).

48.   Rivera-Cancel, G., Motta-Mena, L. B. & Gardner, K. H. Identification of Natural and

Artificial DNA Substrates for Light-Activated LOV–HTH Transcription Factor EL222. *Biochemistry*

51, 10024–10034 (2012).

49.   Zhao, E. M. *et al.* Optogenetic regulation of engineered cellular metabolism for microbial

chemical production. *Nature* **555**, 683–687 (2018).

50.   Peng, T. *et al.* A BaSiC tool for background and shading correction of optical microscopy

images. *Nat. Commun.* **8**, 1–7 (2017).

Figure 1

Figure 1. Using optogenetics to interrogate the properties of cell-generated signaling gradients.
A. The mating switch. Yeast cells switch from budding to mating morphologies when α-factor is
present (top). This dose-response is sharp and occurs in the nM range, close to the receptor Kd
across strains. **B.** When cells with opposite mating types are located at a distance, the generated α-
factor gradient (red arrow, indicating the direction of decay) has a strength (amplitude and
steepness) that defines where the concentration is sufficient for switching-on mating behavior. The
profile of the α-factor pheromone not only depends on the local density of emitters contributing to
the gradient, but also on the presence of the extracellular α-factor peptidase Bar1, and therefore on
the density of *MAT***a** cells. When Bar1 is absent or low, the position of the mating switch is expected

to happen at a distance. In contrast, if Bar1 is high, the mating switch is displaced towards the source vicinity. **C**. Due to low Bar1 mediated degradation of α-factor when *MAT***a** populations have low density, the α-factor gradients are expected to overlap and provide sufficient intensities to trigger mating behavior at a distance, but at the cost of reduced precision due to "confusing" local maxima. **D.** Optogenetic control of α-factor and Bar1 emission allows spatial control of these two opposing activities. In the example shown, a Petri dish containing mating mixes is illuminated in specific regions allowing us to stimulate production or degradation of the pheromone (red color), depending on which genetic determinant is controlled.

Figure 2

Figure 2. Optogenetic control of extracellular α-factor production allows tight spatial control of mating.

A. The light-responsive α-factor-producing "opto-α" strain was engineered by replacing the native promoters of the two pheromone-coding genes in $MAT\alpha$ (MFα1 and MFα2) with the P_{C120} promoter, which is responsive to the light-activated LOV domain EL222 transcription factor, and by later duplicating the MFα1 (which produces the highest amount of pheromone of the two genes) light-responsive locus. **B.** The activity of the P_{FUS1} pheromone-responsive promoter in $MAT\mathbf{a}$ receivers in stirred mating reactions as a function of light intensity. In the assay, the opto-α strain is co-incubated either with $bar1\Delta$ (top left) or a wild-type $MAT\mathbf{a}$ (bottom left) P_{FUS1} GFP reporter strain. After 2 h, the mixes are assessed by flow cytometry. As a control, the reporters are incubated with wild-type $MAT\alpha$, and the baseline-subtracted relative response is calculated (right). Error bars are the SD of n=3 biological replicates. **C.** Agarose-embedded $MAT\mathbf{a}$ bar1Δ reporter strains (cells carrying cytoplasmic GFP) co-incubated with opto-α cells (carrying a red nuclear signal expressed by a mApple fluorescent protein) for 5 h under non-illuminated (left) and maximally illuminated (5.8 mW/cm², right) conditions. **D, E.** Quantification of mating efficiency through light-controlled pheromone production. Light induces cell-cell fusion in the optogenetic mating assay measured as the appearance of diploid colonies only in illuminated regions (D). Quantification of the mating efficiency at different light-doses. Efficiency is defined as relative to the activity of the wild-type $MAT\alpha$ strain (C). Error bars are the SD of n=3 biological replicates.

Figure 3

Figure 3. Optogenetic control of extracellular signal degradation allows tight spatial control of
growth arrest.

A. Promoter replacement in the Opto-Bar1 strain. EL222-mediated light-dependent expression of
the native *loci* encoding Bar1 in the *MAT***a** strain (top) and the pheromone activity assay, in which
exogenously added pheromone stimulates the sensitized GFP reporter strain (bottom). **B.** Resulting
light-dose response from the assay, measured by flow cytometry. The biological activity of the
pheromone is defined relative to the activity of the wild-type *MAT***a** strain. Error bars are the SD of
3 biological replicas**. C, D.** Diploid selective (C) and non-selective plates (D) showing mating
efficiency and growth, respectively in illuminated and dark regions. **E**. Quantification of avoidance
of cell cycle arrest in the Opto-Bar1 strain as a function of light intensity, based on the integrated
pixel intensity of a radial profile (traces shown correspond to the colored bars in D). **F.** Light-dose
dependency of cell cycle arrest in the Petri-dish assay. Growth is integrated over the entire plate
and normalized to the value of non-activated opto-Bar1 cells. Growth of the wild-type is shown as
a green dot. Error bars are the SD of two biological replicates.

Figure 4

Figure 4. Optogenetic generation of pheromone gradients through the half-domain assay.

A. Half-domain assay. Agarose-embedded cells are located at the bottom of the mini well and half of the glass surface is illuminated to induce EL222-dependent expression. **B.** To determine the position of the light border in samples along the acquisition path (red arrows), a control strain that activates EL222-dependent fluorescent protein expression is run in parallel in adjacent wells. The mean value of the border positions above and below the sample correspond to the position of the light border in the sample. **C.** Schematic representation of the experiment, close to the light-dark transition border. **D.** Microscopy image showing an overview of the mating reaction with the controls for the border location (top and bottom row) displaying strong YFP fluorescence on each side of the mating reaction (center row). Arrowheads at the bottom show where the image was stitched together. In the sample mating reaction (center), opto-α has a red nuclear marker while reporter *MAT***a** bar1delta expresses GFP when stimulated by pheromone. Samples were imaged after 5 h of light exposure. **E.** Example fields along the experimental length showing the increased abundance of elongated phenotypes and the increase in GFP levels in the illuminated region.

Figure 5

Figure 5. Quantitative properties of optogenetically generated gradients recapitulate switch-like and linear responses in *MAT*a cells. A. Theoretical, one-dimensional arrangement of pheromone sender cells on one half-domain of an infinite line and pheromone receivers in the other half-domain. **B.** Concentration profile of α-factor at steady state (Eq. 2, See Supplementary Information, section II). **C, D.** Normalized response of the receptor (Eq. 3 normalized by its maximal value, See Supplementary Information, section II) for half-domain expression of α-factor, showing predictions for responses using a Hill coefficient equal to n=1 (E panel) and n=4 (F panel). Hill coefficients greater than one correspond to a shift in the response towards a higher concentration of α-factor and a steeper response. **E.** Quantification of single-cell gene expression output and overlayed model fit, with the Hill coefficient set to 1. **F.** Quantification of single-cell gene morphological response and overlayed model fit, with a fitted Hill coefficient of 3. **G.** Population-level and single-cell estimations of pheromone-profile parameters.

Figure 6

Figure 6. Bar1-generated α-factor gradients reveal slow Bar1 diffusion. A. Schematic representation of the half-domain assay for opto-Bar1, close to the light-dark transition border. Emitter cells produce alpha factor homogeneously. Light-induced Bar1 secretion occurs with

unknown spatial distributions (bottom box). **B.** Zoomed-out depiction of the half-domain assay for Bar1 production and microscopy imaging of Bar1 emitters (cells with red nuclei), receivers (cells with green nuclei), and newly formed diploids (cells with yellow nuclei). The yellow line represents the position of the left limit of the light-dark transition border. **C.** Quantification of the single-cell morphological response (cell length). The light border is represented by the blue vertical line corresponding to the border location determined with the controls run in parallel. The blue dots correspond to the mean value in each one of 30 bins and the error bar represents its standard error. The green dashed line represents the fitted "perfect sink" model; the orange line represents the 1-parameter fit of a shifted logistic function, used to determine the distance at which the half-maximum value of the response occurs (See Supplementary Information, Section III). The inset illustrates the prediction for the "perfect sink" model, showing the sharp concentration decay of Bar1 (top) used to obtain the generated α-factor profile (bottom). **D.** Contour plot showing the results of numerical simulations using the full mathematical model, showing the relation between diffusion of Bar1 relative to diffusion of α-factor and Bar1 activity for different α-factor profiles, characterized by the distance (in units of one α-factor decay length) that the half-maximal value of the induced response shows from the border (zero). The red dashed line highlights the isoline for the experimentally determined value of such a distance (See panel B). **E-H.** Basic properties of signal landscapes in yeast mating. Pheromone secretion by single cells does not provide positional information (E), as even sensitized partners (without expressing Bar1) sense a subthreshold concentration of pheromone., insufficient to trigger chemotropism. During mating, Bar1 has a lower steady-state decay length than α-factor (F). If the mating-type ratio is unbiased (G), Bar1 levels are sufficient to modify the overall pheromone profile. Consequently, cells do not elongate at a distance. Rather, short-range mechanisms[6] (focal pheromone emission, polarisome chemotaxis) allow mating in "touching" mates. If $MAT\alpha$ is abundant and $MAT\mathbf{a}$ is not (H), Bar1 is insufficient to modify the global pheromone landscape but $MAT\mathbf{a}$ will be also able to elongate (arrow) at a distance, because $MAT\mathbf{a}$ is activated by pheromone concentrations from the contributions of several $MAT\alpha$ cells. Consequently, however, cells do not necessarily follow the right direction and elongate in random orientations.