Annotation and assessment of functional variants in regulatory regions using epigenomic data in farm animals

Ruixian Ma[1,#], Renzhuo Kuang[1,#], Jingcheng Zhang[2,#], Jiahao Sun[1], Yueyuan Xu[1], Xinbo Zhou[1], Zheyu Han[1], Mingyang Hu[4], Daoyuan Wang[1], Yuhua Fu[1], Yong Zhang[2], Xinyun Li[1], Mengjin Zhu[1], Shuhong Zhao[1,3,4], Tao Xiang[1,*], Mengwei Shi[4,*] and Yunxia Zhao[1,4,*]

[1] Key Lab of Agricultural Animal Genetics, Breeding and Reproduction of Ministry of Education and Key Laboratory of Swine Genetics and Breeding of Ministry of Agriculture, College of Animal Science and Technology, Huazhong Agricultural University, Wuhan, China.

[2] Key Laboratory of Animal Biotechnology of the Ministry of Agriculture, Northwest A&F University, Yangling, China.

[3] Hubei Hongshan Laboratory, Huazhong Agricultural University, Wuhan 430070, China.

[4] Yazhouwan National Laboratory, 8 huanjin Road, Yazhou District, Sanya City, Hainan Province, 572024, China.

[#] These authors contributed equally

Corresponding author: Tao.Xiang@mail.hzau.edu.cn; shimengwei@yzwlab.cn; yxzhao@mail.hzau.edu.cn

Abstract

Background

Understanding the functional impact of genetic variants is essential for advancing animal genomics and improving livestock breeding. Variants that disrupt transcription factor (TF) motifs provide a means to assess functional potential, but the lack of TF ChIP-seq data for farm animals presents a challenge.

Results

To address this, we curated nearly 900 epigenomic datasets from 10 farm animal species and annotated eight regulatory regions to assess how variants affect TF motifs.

Over 127 million candidate functional variants were classified into five functional

confidence categories across the species. Variants with high confidence were enriched

in eQTLs and trait-associated SNPs, showing greater potential to affect gene

expression and phenotypes. Incorporating these functional variants into genomic

prediction models improved the accuracy of Estimated Breeding Values (EBVs).

Active variants also revealed trait-related tissues, and single-cell RNA sequencing

(scRNA-seq) identified the cell types most associated with production traits. To

facilitate research, we developed the Integrated Functional Mutation (IFmut) platform,

enabling users to explore variant functions easily. Our study provides a flexible

platform and resource for studying genomic variation in farm animals, setting a new

standard for research and breeding strategies.

Conclusion

The results indicated that evaluating functional potential by annotating and

categorizing variants that interfere with transcription factor motifs can help elucidate

changes in gene expression and phenotype. By focusing on high-confidence variants

enriched in eQTL and trait-associated SNPs, it improves the accuracy of genomic

predictions in research and breeding strategies.

Keywords: Farm animals; Epigenetic annotation; Functional variants; Regulatory

regions

Introduction

Genomic variants in *cis*-regulatory elements are critical drivers of gene expression

changes, influencing phenotypes [1-4]. While many variants are neutral, some can

cause diseases or other harmful effects [5]. Identifying and characterizing variants is a

critical focus in human genetics research, especially in understanding genetic

disorders [6-8]. Large-scale initiatives like the 1000 Genomes Project have propelled

human genome exploration [9, 10]. Similarly, the elucidation of genetic variants in

farm animals, including pigs (*Sus scrofa*), cattle (*Bos taurus*), sheep (*Ovis aries*),

chickens (*Gallus gallus*), mallards (*Anas platyrhynchos*), swan geese (*Anser*

cygnoides), and goats (*Capra hircus*), as well as companion animals like rabbits

(*Oryctolagus cuniculus*) and dogs (*Canis lupus familiaris*), and working animals like

horses (*Equus asinus*), is vital for optimizing traits like production efficiency, health,

behavior, and performance [11]. Genomic selection (GS) [12] has revolutionized

animal breeding, enabling more precise selection based on traits like growth rates or

disease resistance [13-17]. However, not all variants are functionally relevant,

creating a need for systematic methods to identify and interpret functional variants.

We reviewed studies focused on identifying candidate functional variants.

CADD, primarily designed for human variants, uses supervised machine learning to

pinpoint deleterious mutations but may miss neutral or beneficial ones due to its bias

[18]. Similarly, pCADD (for pigs) and chCADD (for chickens) follow similar

algorithms but lack regulatory annotations, limiting their accuracy in non-coding

regions [19, 20]. FitCons, which estimates selective pressure, may be less effective in

detecting weakly selected mutations tied to complex traits [21]. Eigen offers

robustness but relies on high-quality datasets, and the shortage of experimentally

validated non-coding variant data hinders its reliability [22]. These methods struggle

to predict non-coding variant effects in farm animals due to their reliance on

conserved regions and limited data.

Functional genomic annotation is key to identifying variants associated with

complex traits. Genome-wide association studies (GWAS) show that many genetic

variants linked to diseases and traits lie in non-coding regions, as shown by the

Encyclopedia of DNA Elements (ENCODE) [23]. In humans, the ENCODE project

has driven considerable progress in functional variant screening, uncovering

previously unrecognized functions of several regions in the human genome. In

addition, RegulomeDB database and IGVF annotate and prioritize genetic variants in

non-coding regions based on ENCODE project, enhancing the discovery of functional

human variants and becoming a highly utilized resource for the research community

[24, 25]. Despite these advances, regulatory variants in animal genomes remain

underexplored.

To address the abovementioned concerns, we curated 877 datasets from 10

species, including pig, cattle, sheep, goat, horse, dog, rabbit, swan goose, mallard, and

chicken. These include 357 ATAC-seq, 26 DNase-seq, 363 ChIP-seq, 56 Hi-C, 69

scRNA-seq, and quantitative trait loci (QTL) datasets related to trait of six species.

Based on these, we annotated QTLs, open chromatin regions (OCRs),

nucleosome-free regions (NFRs), motifs, footprints, TF-binding sites, and H3K27ac

peaks, covering eight functional regulatory regions. We then identified candidate

functional variants and developed a classification system to evaluate their impact on

regulatory functions. We further assessed variant enrichment in expression QTLs

(eQTLs) and trait-associated SNPs, the performance of high- and

moderate-confidence variants in EBVs, variant activity across tissues, and integration

of scRNA-seq data. Finally, we created an IFmut database and a Functional

Confidence scoring system, providing a versatile public resource for researchers. This

study delivers a comprehensive database and a powerful online toolkit, advancing the

exploration of functional variants in farm animals for both research and breeding

applications.

Methods

Data collection

Genome variants data in VCF format for pig (*Sus scrofa,* susScr11.1), cattle (*Bos*

taurus, bosTau9), and chicken (*Gallus gallus,* galGal5) were downloaded from the

Ensembl database (http://ftp.ensembl.org/pub/). Genome variants for mallard (*Anas*

platyrhynchos, CAU_Wild1.0), goose (*Anser cygnoides*, GooseV1.0), goat (*Capra*

hircus, ARS1.2), rabbit (*Oryctolagus cuniculus*, OryCun2.0), dog (*Canis lupus*

familiaris, CanFam3.1), and horse (*Equus asinus*, EquCab3.0) were downloaded from

Ianimal database (https://ianimal.pro/reception/download) [26], while sheep (*Ovis*

aries, oviAri4) data were obtained from the NCBI Single Nucleotide Polymorphism

Database

(https://ftp.ncbi.nih.gov/snp/organisms/archive/sheep_9940/VCF/00-All.VCF.gz).

QTL data for ten species were downloaded from the Animal QTL database

(https://www.animalgenome.org/cgi-bin/QTLdb/). For pig, we also downloaded

significant *cis*-eQTL and *trans*-eQTL data from the PigGTEx-Portal database

(https://piggtex.ipiginc.com/#/downloads) [27].

TF ChIP-seq, H3K27ac ChIP-seq, ATAC-seq, and Hi-C data for pig, cattle,

sheep, mallard, goat, goose, rabbit, dog, horse, and chicken, and scRNA-seq data in

those eight species except goose and horse, as well as DNase-seq data for chicken,

were   downloaded   from   the   NCBI   Sequence   Read   Archive

(http://www.ncbi.nlm.nih.gov/sra/). After quality control, a total of 802 epigenomic

datasets and 69 scRNA-seq datasets were collected from multiple projects, including

75 datasets from pigs obtained from our previous study [28]. To investigate the

evolutionary sequence conservation of functional variants, we downloaded GERP

scores for pigs from Ensembl

(https://ftp.ensembl.org/pub/current_compara/conservation_scores/).

Sequencing data analysis

To adhere to ENCODE standards, we primarily followed the analysis methods

outlined in our previous study for processing ChIP-seq and ATAC-seq data [28].

ChIP-seq

Mapping and Quality control

The ENCODE ChIP-seq pipeline (https://github.com/kundajelab/chipseq_pipeline)

was used to process ChIP-seq datasets. Raw reads from each dataset were aligned to

the respective reference genomes using BWA v0.7.17 [29]. Low-quality reads

(MAPQ < 25), unmapped reads, mate unmapped reads, non-primary alignment reads,

and   duplicate   reads   were   removed   using   Picard   v1.126

(https://broadinstitute.github.io/picard) and SAMTools v1.9 [30].

Read coverage of genomic regions between filtered replicate BAM files was

computed using the multiBamSummary bins function of deepTools v2.0 [31], with a

bin size of 2 kb to assess genome-wide similarities. Pearson correlation coefficients

were calculated from the resulting read coverage matrix. Replicates with a Pearson

correlation coefficient ≥0.8 were merged, while those with coefficients <0.8 were

excluded from further analysis.

Identification of nucleosome-free regions

HOMER software was used to detect nucleosome-free regions (NFRs) [32]. Tag

directories for H3K27ac IP and input data were generated using the

makeTagDirectory command using the merged non-duplicated BAM file obtained

from the "Mapping and Quality control" procedures. Subsequently, the findPeaks

command with the -nfr option was applied to identify NFR peaks. Only datasets with

at least 10,000 peaks were considered, excluding scaffold regions.

Identification of TF binding sites and H3K27ac narrow peaks

TF binding sites and H3K27ac narrow peaks were identified using MACS2 v2.1.0 [33]

and deepTools v2.0 [31], as previously described in greater detail in our prior study

[28].

154   **ATAC-seq**

155   **Mapping, quality control, and peak calling**

156   The ENCODE ATAC-seq pipeline

157   (https://github.com/kundajelab/atac_dnase_pipelines) was used to process ATAC-seq

158   datasets. Adapter trimming was performed using Cutadapt v1.14

159   (https://cutadapt.readthedocs.io/en/stable/), followed by alignment to the reference

160   genome assemblies with Bowtie2 v2.3.4.1. Low MAPQ reads (< 25), unmapped reads,

161   mate unmapped reads, not primary alignments, reads failing platform, and duplicates

162   were removed using SAMTools v1.9 [30] and Picard v1.126

163   (https://broadinstitute.github.io/picard). Mitochondrial reads were removed using

164   BEDTools v2.26.0 [34] to generate effective reads for peak calling. Peaks were called

165   using MACS2 v2.1.0 [33] using: genome size (-g), *p*-value threshold (0.01), peak

166   model (--nomodel), shift size (--shift), extension size (--extsize), and other options

167   (--B, --SPMR, --keep-dup all, --call-summits), to generate at least 10,000 peaks for

168   further analysis.

169   **DNase-seq**

170   **Mapping, quality control, and peak calling**

171   For the DNase-seq datasets for chicken, the ENCODE DNase-seq pipeline

172   (https://github.com/kundajelab/atac_dnase_pipelines) was used, with 'dnase_seq'

173   specified to indicate DNase-seq data. The remaining steps were consistent with the

174   ATAC-seq analysis.

Identification of open chromatin regions (OCRs)

Peaks with $p < 10^{-5}$ were considered significant for further analysis. Significant narrow peaks were filtered based on replicates with high Pearson correlation coefficients ($R \geq 0.8$). Merged peaks, representing OCRs, were generated using BEDTools v2.26.0 [34], requiring at least 50% overlap between replicates. Signal tracks were generated from highly correlated replicates ($R \geq 0.8$) using MACS2 v2.1.0 [33] to visualize chromatin accessibility across the genome.

Identification of footprints in ATAC-seq and DNase-seq

Footprint analysis followed these steps: (i) broad peaks were called from merged ATAC-seq or DNase-seq data using the MACS2 v2.1.0 broad module [33, 35]; (ii) significant broad peaks ($p < 10^{-10}$ and $10^{-10} < p < 10^{-5}$ overlapping OCRs) were merged using BEDTools v2.26.0 [34]; (iii) Transcription factor footprinting was performed using the HMM-based Identification of Transcription factor footprints (HINT) framework from Regulatory Genomics Toolbox (RGT) v0.13.2 [36] with parameters adjusted for ATAC-seq or DNase-seq data (--atac-seq or --dnase-seq) and paired-end sequencing data (--paired-end). The organism information (--organism=) was also specified; and (iv) the cutoff for footprint score was set at the 20^{th} percentile of all footprints identified by the HINT framework of GRT v0.13.2 [36].

Transcription factor motif mapping in genome function region

The transcription factor motif mapping was primarily performed as follows: (i) OCR, NFR, TF binding sites, and footprint in OCR regions were merged into a BED file; (ii)

The   fasta-get-markov   command   from   the   MEME   Suite

(https://github.com/cinquin/MEME) was used to generate a .fa.bg file and the

"bedtools getfasta" command generated the corresponding .fa file from the BED file

of step (i); (iii) The fimo command in MEME Suite (--max-stored-scores 5,000,000)

was used to map motifs in the genome; and (iv) the results of fimo mapping for pig,

cattle, and sheep were filtered by $p < 5\times10^{-6}$, while for chicken, $p < 5\times10^{-7}$ was

applied.

Prediction of transcription factor motif effects

Potential functional variants (Categories 1-5) located within footprint regions were

analyzed using the motifbreakR [37] package in R v4.0. The motifDB database,

specifically JASPAR 2018 [38], was selected as the data source for predicting the

transcription factors to which the SNPs may bind.

Hi-C data processing

The Hi-C data of two-week-old LW pigs were taken from our previous study [28],

and other Hi-C data were downloaded from the GEO at http://ncbi.nlm.nih.gov/geo.

The downloaded data were processed using the HiC-Pro (v2.11.1) pipeline to produce

the ICE normalization contact matrices [39]. The insulation score of the ICE matrix

was calculated using the following options: -is 480000, -ids 320000, -im iqrMean, -ss

160000. The insulation method was utilized to define the topologically associating

domain (TAD) structure (insulation/boundaries). Hi-C loops were analyzed using

HiCCUPS v0.3.7 [40], with minor modifications, and the resolution parameter was set

to "-r 25000, 40000".

Analysis of scRNA-seq

A species-specific reference index was generated using "cellranger mkref" from 10x

Genomics Cell Ranger v7.0.1. The following reference genomes and their

corresponding filtered GTF files from Ensembl were used: pig (susScr11), cattle

(bosTau9), chicken (galGal5), sheep (oviAri4), goat (ARS1.2), rabbit (OryCun2.0),

and dog (CanFam3.1). The GTF files were filtered using "cellranger mkgtf" with

default parameters. Expression data were processed using Seurat (v5.1.0) [41], with

low-quality cells and empty droplets removed. Cells were initially filtered according

to the following criteria: a minimum of 200 expressed genes per cell and at least three

gene counts. Further quality control steps eliminated cells with fewer than 200 unique

feature counts or mitochondrial content exceeding 5%. The remaining data were used

for downstream analysis. Data normalization was performed using the NormalizeData

function with a scaling factor of 10,000 to account for library sequencing depth.

Principal component analysis (PCA) was conducted on the expression matrix of the

top 2,000 most variable genes. Dimensionality reduction and clustering were

performed using t-SNE. Cell types were identified using the CellMarker 2.0 database

[42]. An R script was written to identify the intersection between the top 25 marker

genes in each cluster and known marker genes from the database, enabling candidate

cell type identification. The average $\log_2$FC of each marker gene was normalized, and

the sum of all marker gene scores was utilized to assign a score to each cell type.

Higher scores indicated a higher likelihood of the cluster representing a specific cell

type. Final cell type identification was corroborated using prior research findings.

Variants distribution statistics

SNPs and small InDels were annotated using snpEff v4.5 software, with the reference

genome file (FASTA) and annotation files (GTF) [43].

Performance of genomic predictions

Dataset

The phenotypic dataset used for genomic prediction was obtained from a national pig

nucleus herd in North China. The study included phenotypic recordings for two

productive traits: average daily gain (ADG) from 30-100 kg and backfat thickness

(BF) at 100 kg. All phenotypic records for the traits were obtained at the same time

point, allowing a 10-kg deviation from the final body weight (100 ± 10 kg). All

phenotypes were recorded between early 2018 and October 2022. Based on the traced

pedigree, 11 pig lines were identified within the population. DNA samples were

collected from approximately 80 distantly related pigs per line, and sequencing was

performed using the DNBSEQ-T7 platform, with an average coverage of 5×. In total,

874 pigs were sequenced. After quality controls–such as a genotype missing rate

below 10%, a call rate of SNPs above 90%, and a minimum allele frequency (MAF)

above 1%–18,460,807 (18000K) SNPs were retained for analysis. Missing genotypes

were imputed using Beagle software (v5.3). Of the 874 sequenced pigs, 872 had ADG

phenotypic data, while 867 had BF recordings. Environmental factors, including

gender, herds, and physical units, were fully recorded.

Genomic Best Linear Unbiased Prediction (GLUP) models

Breeding values (EBV) for different traits were estimated using GBLUP models:

$$y = Xb + Zu + e,$$

where y indicates a column vector of phenotypic values for each trait; b represents a

vector of fixed effects, including sex effect, herd effect, and physical unit effects; u

denotes a vector of random additive genetic effects; e is a vector of residual effects.

Matrices X and Z are design matrices associated with these effects. The GBLUP

model assumes a normal distribution for the random additive and residual effects, as

$u \sim N(0, G\sigma_u^2)$, where G denotes a genomic relationship matrix constructed as

Vanraden method 1; $e \sim N(0, I\sigma_e^2)$, where I is an identity matrix. The additive

genetic variance and residual variance are indicated by σ_u^2 and σ_e^2, respectively.

Scenarios of constructing genomic relationship matrices

GBLUP models with four different genomic relationship matrices (G) were employed

to estimate GEBVs for ADG and BF traits. Four different sets of SNP markers were

utilized to develop corresponding G matrix. In scenario 1, sequenced SNP markers

with the top 1 and top 2 muscle scores (1+2 muscle, 10,544 SNPs) were calculated the

G matrix. Similarly, sequenced SNP markers with the top 1 and top 2 liver scores

(1+2 liver, 6,049 SNPs) and top 1 and top 2 adipose scores (1+2 adipose, 3,801 SNPs)

were employed to construct G matrices in scenarios 2 and 3, respectively. In scenario

4,11,000 randomly selected SNP markers were utilized to develop a G matrix.

Scenario 4 was repeated three times in the study.

Predictive Reliabilities

The mean predictive reliabilities of GEBVs were computed using the subsequent

formula [44]:

284
$$r^2 = \sum \left(1 - \frac{SEP_i^2}{\sigma_g^2}\right) / N,$$

where r^2 is the reliability of GEBVs and i indicates an individual animal i; SEP

denotes the standard error associated with the predicted GEBVs; σ_g^2 is the additive

genetic variance and N is the number of animals used.

Results

Large-scale epigenomic screening of functional variants across farm animals

To annotate genomic variation, we gathered SNPs and small InDels (2-50 bp) from

public databases for ten farm animals: pig, cattle, sheep, goat, chicken, horse, mallard,

swan goose, rabbit, and dog. This dataset, comprising nearly 570 million variants, was

classified using various sequencing methods (Fig. 1A). SnpEff software analysis

showed that over 95% of the variants were located in non-coding regions across all

species (Fig. 1B), consistent with previous findings in human studies where variants

are commonly found in intergenic and intronic regions [7, 45, 46]. Thus, identifying

functional variants from this vast pool remains a significant challenge. Given that

regulatory regions, particularly transcription factor (TF) binding sites identified by

epigenomic analyses, can inform the potential functions of variants, we aimed to

screen for potentially functional variants using epigenomic data [25, 47]. To do this,

we collected datasets from previous studies, including the FANNG project [48, 49]

and our own research [28], encompassing ATAC-seq, DNase-seq, H3K27ac ChIP-seq,

TF ChIP-seq, Hi-C, and scRNA-seq data from 10 farm animals. After rigorous

mapping and quality control, 871 datasets from 32 tissues and eight cell lines passed

quality standards, including 357 ATAC-seq, 26 DNase-seq, 363 ChIP-seq, 56 Hi-C,

and 69 scRNA-seq datasets (Fig. 1C). Specifically, we compiled 160, 199, 40, 22, 106,

213, 11, 8, 31, and 81 datasets for pig, cattle, sheep, goat, horse, chicken, mallard,

swan goose, rabbit, and dog, respectively.

Using ENCODE guidelines (https://www.encodeproject.org/), we identified

regulatory regions, resulting in over 4.6 million non-redundant genomic regions

across all species. These included open chromatin regions (OCR), H3K27ac peaks,

footprints, and nucleosome-free regions (NFR) (Supplementary Table S1).

Additionally, TF binding peak calling combined with positional weight matrices

(PWMs) identified 8.8 million non-redundant regions associated with TF binding sites

across the ten species (Supplementary Table S1). The total length of these regions

accounted for 33.55% of the pig genome (Sscrofa11.1), 43.67% for cattle

(ARS-UCD1.2), 15.00% for sheep (Oar_v4.0), 5.12% for goat (ARS1.2), 34.76% for

horse (EquCab3.0), 44.65% for chicken (Gallus_gallus-5.0), 7.94% for mallard

(CAU_Wild1.0), 4.37% for swan goose (Goose1.0), 8.93% for rabbit (OryCun2.0),

and 38.97% for dog (CanFam3.1) (Supplementary Table S2).

Genomic variants in TF binding sites, such as those cataloged in RegulomeDB,

often result in functional consequences [25]. Since the above genomic regions

containing basic regulatory and TF binding-related regions were identified through

DNA-TF interaction data, variants detected in these regions were likely to have

transcription regulation function in the farm animals. Moreover, we further identified

1,128,759~39,157,953 (4.98~47.05%) variants located in these regulatory regions

across the ten species (Supplementary Fig. S1), which we designated as potential

functional variants.

Classification approach for functional impact of variants

Gene expression is largely regulated by TFs, proteins that bind to specific [50] DNA

sequences to control gene transcription [50, 51]. Variants located within TF binding

motifs can disrupt this binding, potentially altering gene expression and impacting

phenotype (Fig. 2A). To better evaluate the likelihood that a regulatory variant

impacts transcription, we developed a functional confidence classification system

similar to that of RegulomeDB [25] (Fig. 2B; Supplementary Table S3).

Currently, only 103 TF ChIP-seq datasets exist for farm animals, most of which

target CTCF (86 datasets). Chickens have the largest number of ChIP-seq datasets,

with 17 datasets for 9 TFs. Thus, due to the lack of TF ChIP-seq data in farm animals,

the identification of potential TF motif regions relied on a combination of

ATAC-seq/Dnase-seq (i.e., OCR and footprints) and H3K27ac ChIP-seq (i.e., NFR

and H3K27ac peaks). We also incorporated QTL data, as variants in these regions

may affect agronomic traits.

Our functional confidence classification system evaluates variants based on the

number of regulatory regions they overlap with. Given the prominent association of

NFRs, OCRs and TF footprints (especially those containing fully or partially

matching recognition motifs) with transcriptional activation, variants in these regions

had the highest likelihood of affecting TF binding and gene expression, and were

therefore scored as high functional confidence variants (Category 1). Variants

meeting these criteria but never found in NFRs were subsequently scored as moderate

functional confidence (Categories 2a-2d), suggesting a moderate likelihood of

affecting TF activity. Additionally, variants located in QTL regions were assigned

higher scores (e.g., Categories 1a, 1c, 2a, and 2c), while those outside QTLs received

slightly lower scores (Categories 1b, 1d, 2b, and 2d). By contrast, variants detected in

DNase/ATAC-seq or H3K27ac ChIP-seq data but not in TF footprints or recognition

motifs were classified as low (Category 3) or minimal (Category 4) functional

confidence. Category 5 included variants detected only by H3K27ac ChIP-seq, which

may still influence transcription regulation (Fig. 2B; Supplementary Table S3).

Based on our classification approach, we identified 126,937,925 candidate

functional variants. Among these, 26,341 to 3,096,314 SNPs/small InDels were

classified as high or moderate confidence (Categories 1 and 2), representing 1.15% to

7.91% of all functional variants across the ten animals (Fig. 3C; Supplementary Fig.

S2). As expected, we observed some variants with functional potential were related to

genes known to significantly influence production traits. For example, the CCK gene

encodes cholecystokinin, which regulates digestive processes, appetite, and fat

metabolism, playing a crucial physiological role in animals [52-55]. In pig genome,

there is a SNP (rs332577759, chr13:25945714:C:T), which located in the exon of

CCK, showed significant signal in the ATAC-seq and H3K27ac ChIP-seq peaks,

leading to its annotation and classification as Category 2d (Fig. 2D). Similarly, the

SNP rs519400784 (chr22:14573586:C:CCCGAGAG) from cattle genome, located in

the homologous region of rs332577759, is also surrounded by strong signals of

ATAC-seqH3K27ac ChIP-seq, and CTCF ChIP-seq, resulting to its classification as

Category 1d (Fig. 2D).

Variants in higher functional categories are more likely associated with traits

To further explore the characteristics of variants across different functional categories,

we utilized the Ensembl Genomic Evolutionary Rate Profiling (GERP) score to assess

the evolutionary conservation of all variants [56]. Surprisingly, variants with higher

functional confidence showed significantly lower GERP scores compared to

non-functional ones, especially in Category 1 (Fig. 3A). Negative GERP scores

indicate reduced sequence conservation at these loci, suggesting higher mutation rates,

potentially due to neutral or positive selection. To ensure that these conservation

differences weren't influenced by the surrounding regulatory elements, we extended

the analysis to include 100 bp flanking regions around the variants (Fig. 3B). The

results showed that GERP scores in non-functional variants remained stable, while the

conservation scores of functional confidence variants increased significantly,

approaching the levels observed in non-functional regions. This finding supports the

notion that the lack of conservation is an inherent feature of these variants.

Previous studies suggest that artificial selection has shaped farm animal genomes,

particularly in loci linked to production traits, which tend to undergo strong positive

selection [57]. This evolutionary pattern mirrors what we observed in variants with

higher functional categories, leading us to hypothesize that artificial selection may

drive the lower conservation of these variants. To test this, we analyzed eQTL data

from over 30 pig tissues using the PigGTEx database and observed that eQTLs were

significantly more likely to be functional variants than non-functional ones (Fig. 3C)

[27]. This indicates that functional variants are more likely to influence gene

expression, a key determinant of phenotypic traits. Additionally, intersecting

functional confidence variants with known trait-associated loci revealed significant

enrichment in these regions (Fig. 3D). These findings demonstrate that our Functional

Confidence Classification system effectively identifies variants with a high likelihood

of impacting gene expression and driving phenotypic variation.

To further validate our classification system, we screened high and moderate

confidence variants (Category 1 and Category 2) in pigs and assessed their predictive

reliability for estimated breeding values (EBVs) in two traits—average daily gain

(ADG) and backfat thickness (BF)—using a large white population ($n = 874$).

Genomic BLUP modeling was performed using DMU software [58] with four

scenarios: three using functional variants from muscle, liver, or adipose tissue, and

one using 11K random variants from whole-genome pig sequencing (Fig. 3E;

Supplementary Table S4). The predictive reliability of EBVs for ADG and BF was

similar across the three tissue-based scenarios (~0.31–0.38), with the adipose tissue

scenario showing the highest reliability (~0.38), despite using the fewest markers

(3,861 SNPs). In comparison, the random SNP scenario, despite containing the

highest number of markers (11,000 SNPs), yielded the lowest reliability (~0.27). The

predictive reliability of EBVs improved by 23–46% for all three tissue types when

high and moderate functional variants were used, compared to the random SNP

scenario (Fig. 3F). This further demonstrates the utility of our Functional Confidence

Classification system in screening functional variants for genomic data in farm

animals.

Functional variants define the principal tissues of *cis*-regulatory elements and

traits

Besides evaluating the functional potential of variants, considering that the function of

variants is heavily influenced by their regulatory elements, our data also enables the

identification of the specific tissues in which these variants are active. By merging

regulatory elements, we generated a comprehensive *cis*-regulatory element (CRE)

map for seven species (the tissue numbers in other species were insufficient). The

proportion of active variants was used to estimate the tissue-specific activity of CREs,

and as expected, most CREs exhibited strong tissue specificity, being active in only

one or two tissues (Fig. 4A; Supplementary Fig. S4). This observation aligns with

findings from the human EpiMap project [59], which reported that 88% of enhancer

elements are tissue-specific. Our results further extend this conclusion, showing that

such tissue specificity is prevalent across all types of CREs.

Next, we analyzed the activity of trait-associated variants to identify the tissues

linked to production traits. In animal breeding, accurate identification of tissues

related to production traits is critical for understanding the underlying regulatory

mechanisms and improving breeding strategies [60, 61]. To this end, we quantified

trait-associated variants in each tissue and calculated their proportion relative to all

variants related to the trait. Hierarchical clustering grouped tissues into three distinct

clusters (Fig. 4B). Tissues such as adipose, spleen, muscle, and cerebellum showed

significant enrichment of trait-associated variants, indicating a strong association with

production traits (Fig. 4C). In contrast, tissues like the lung, cerebrum, and

hypothalamus showed considerably lower proportions of trait-associated variants (Fig.

4C).

To further delineate the cellular origins of these production traits, we employed

single-cell RNA-Seq (scRNA-seq) data to identify the cell types where variant-related

genes were classified as marker genes (Fig. 4D and 4E). Variants were considered

active in a cell type when their associated gene was a marker gene, reflecting the

functional relevance of the variant in that cellular context. Through calculating the

proportion of active variants across all traits, we observed some traits related to

skeletal growth, such as feed efficiency and bone lengths (hip, humerus, femur, tibia,

and ulna), were enriched in B cells and macrophages, highlighting the importance of

immune cells in skeletal development (Fig. 4D). This finding aligns with recent

research highlighting the interaction between the immune and skeletal systems, where

immune cells regulate bone remodeling via cytokine, chemokine, and growth factor

secretion [62, 63].

Moreover, several traits related to flavor and texture, such as fat percentage in

the carcass, juiciness, tenderness, shear force, chew score, and off-flavor score, were

significantly enriched in beige adipocytes. These traits, involving 33 SNPs, were

nearly all associated with the *CAST* gene (Fig. 4F). We found that *CAST* expression

was significantly higher in beige adipocytes than in other cell types. Previous studies

have shown that the *CAST* gene is linked to meat tenderness, muscle growth rate, and

post-mortem proteolysis [64], which are key factors influencing pork quality. Our

findings align with these studies and further demonstrate that the *CAST* gene's

influence on flavor-related traits is specifically localized to beige adipocytes.

Development of the Integrated Functional Mutation (IFmut) database for

screening candidate functional variants in farm animals

In order to facilitate screening for candidate functional variants in farm animals, we

integrated genomic variants with epigenomic datasets in a database, the Integrated

Functional Mutation (IFmut) database. IFmut (http://www.ifmutants.com:8212) is a

comprehensive farm animal variants database which provides functional context to

variants or regions of interest and serves as a tool to prioritize functionally important

variants (Fig. 5A). There are six modules (Category, Epigenetics, 3D Genome, Gene,

Trait, and Single cell) within IFmut that enable users to query variants of interest (Fig.

5B).

The 'Category' module integrates functional confidence categories for variants

from ten species, facilitating searches by variant ID or position (Fig. 5C). Additionally,

we also provide a tool (IfmutAnnotator) that allows users to upload a BED file

containing variants not included in our database to classify novel variants

(Supplementary Fig. S5). The "Motif" column aids in hypothesizing the variant's

potential impact on TF binding (Fig. 5D). Users can access detailed annotations by

clicking on the variant ID (Fig. 5E). The 'Epigenetics' module offers six types of

epigenetic regulatory regions (OCR, NFR, Footprint, Motif, TF-binding site, and

H3K27ac peak) across tissues or cells (Fig. 5F), and the '3D Genome' module

contains interaction data such as loops and TADs (Fig. 5G). In the 'Gene' module,

variants are linked to genes within the same TAD or chromatin loop, or to the nearest

functional gene when 3D genome data is unavailable (Fig. 5H). The 'Trait' module

enables users to search for trait-associated variants, providing insights into the genetic

mechanisms of complex traits (Fig. 5I), while the 'Single Cell' module uses curated

scRNA-seq data to analyze variant-associated gene expression across cell types (Fig.

5J). Visualization tools allow users to view epigenetic and Hi-C interaction signals

around variants and nearby genes.

In summary, IFmut offers seven key types of functional information and practical

tools, supporting the exploration of variant functionality.

Discussion

Previous research has revealed that most SNPs and small InDels are located in non-protein-coding regions of the genome [65-68], making it challenging to interpret how these variants affect function [69, 70]. Evaluating the impact of variants on TF binding sites is an effective approach for identifying functional variants in the human genome [25], but the lack of TF ChIP-seq data for farm animals has hindered this method's application in livestock research. To overcome this limitation, we utilized alternative epigenomic techniques like ATAC-seq and DNase-seq, which capture the accessible chromatin landscape and provide insights into TF binding sites. These techniques have been widely used in human and livestock research [71-73].

By leveraging epigenomic datasets, we annotated regulatory regions and developed a confidence classification system for variants based on their functional relevance. Overall, our approach was more suitable the current study of functional variants in livestock for abundant epigenetic datasets in these species, as well as the design idea of using the epigenetic data to identify TF binding sites to rank functional variants can also be transplanted to related research works on other species.

Moreover, by integrating eQTL data and trait-associated variants with our classification system, we demonstrated that higher functional confidence variants are more likely to influence gene regulation and trait expression. This enhances the practical utility of our system. For example, genomic predictions based on high- and moderate-confidence variants improved the reliability of Estimated Breeding Values

(EBVs) in pigs compared to predictions using randomly selected SNPs. Furthermore,

our analysis of active variants allowed us to identify the tissues associated with key

production traits, such as adipose, spleen, and muscle tissues, aligning with breeding

insights. Additionally, by incorporating scRNA-seq data, we pinpointed specific cell

types contributing to these traits. These findings suggest that our functional

classification system can effectively guide the identification of critical variants and

thus contribute to advancing genetic improvement in farm animals.

The lack of research into regulatory variants in farm animal genomes has

constrained progress in animal genomics, genetics, and breeding research. To address

this gap, we designed the IFmut webserver, which enables users to explore genomic

and epigenetic data related to variant functionality. Unlike other resources such as

PigBiobank, AGIDB, and IAnimal [26, 74, 75], IFmut considered the regulation of

genomic interactions that may result from SNP interference with TF. We also

incorporated Hi-C data to characterize 3D genomic interactions and utilized

single-cell data to assess the specificity of variants across cell types. Additionally,

IFmut includes a Functional Confidence classification tool for evaluating novel

variants, which is essential for advancing our understanding of how genetic variants

influence traits, ultimately supporting more informed decision-making in livestock

breeding and genetic improvement.

Conclusion

Assessing functional potentiality by annotating and classifying variants that disrupt

transcription factor motifs can explain alterations in gene expression and phenotype.

Variants with high confidence were enriched in eQTL and trait-associated SNPs, show

greater potential to influence gene expression and phenotype and help improve the

accuracy of EBVs. The establishment of the IFmut platform provides a flexible

platform and resource that sets a new standard for research and breeding strategies.

Abbreviations

TF	Transcription factor
EBVs	Estimated Breeding Values
scRNA-seq	Single-cell RNA sequencing
IFmut	Integrated Functional Mutation
GS	Genomic selection
GWAS	Genome-wide association studies
ENCODE	Encyclopedia of DNA Elements
QTL	Quantitative trait loci
OCRs	Open chromatin regions
NFRs	Nucleosome-free regions
PWMs	Positional weight matrices
GERP	Ensembl Genomic Evolutionary Rate Profiling
ADG	Average daily gain
BF	Backfat thickness
CRE	*Cis*-regulatory element
PCA	Principal component analysis
GLUP	Genomic Best Linear Unbiased Prediction

540 **Acknowledgement**

541 We thank Dr. Christopher K Tuggle (cktuggle@iastate.edu) at Department of Animal

542 Science, Iowa State University for valuable suggestions to our work.

Author contributions

Y.Z, T.X, and M.S conceived and designed the project. R.M, R.K, J.Z, J.S, Y.X, X.Z, Z.H, M.H, D.W, Y.F, Y.Z, X.L, M.Z and S.Z performed the analysis. M.S and Y.F constructed the web-based resource with input from R.M and M.H. R.M, R.K, J.Z, and M.S wrote the manuscript and prepared the figures. All authors read and approved the final manuscript.

Funding

This work was supported by the National Natural Science Foundation of China (32341051), the grant from Department of Agriculture and Rural Affairs of Hubei Province (HBZY2023B006-02), National Key R&D Young Scientists Project (2022YFD1302000), Fund of Modern Industrial Technology System of Pig (CARS-35).

Availability of data and materials

All ATAC-seq, ChIP-seq (H3K27ac and TFs), and Hi-C, as well as the candidate functional variants and their Functional Confidence scores are available at http://www.ifmutants.com:8212. The source code for calculating the Functional Confidence score for candidate functional variants is publicly available on GitHub at https://github.com/RXMa-Bio-software/the-Functional-Confidence-scoring-system.

Declarations

Ethics approval and consent to participate

Not applicable.

Consent for publication

Not applicable.

Competing interests

The authors have no competing interests to declare.

References

1. Zhang F, Gu W, Hurles ME, Lupski JR. Copy number variation in human health, disease, and evolution. *Annu Rev Genomics Hum Genet*. 2009;10:451-481.
2. Wang J, Dai X, Berry LD, Cogan JD, Liu Q, Shyr Y. HACER: an atlas of human active enhancers to interpret regulatory variants. *Nucleic Acids Res*. 2019;47(D1):D106-D112.
3. Pan Z, Yao Y, Yin H, Cai Z, Wang Y, Bai L, Kern C, Halstead M, Chanthavixay G, Trakooljul N et al. Pig genome functional annotation enhances the biological interpretation of complex traits and human disease. *Nat Commun*. 2021;12(1):5848.
4. Redon R, Ishikawa S, Fitch KR, Feuk L, Perry GH, Andrews TD, Fiegler H, Shapero MH, Carson AR, Chen W et al. Global variation in copy number in the human genome. *Nature*. 2006;444(7118):444-454.
5. Heyne HO, Karjalainen J, Karczewski KJ, Lemmela SM, Zhou W, Havulinna AS, Kurki M, Rehm HL, Palotie A, Daly MJ. Mono- and biallelic variant effects on disease at biobank scale. *Nature*. 2023; 613(7944):519-+.
6. Cherry TJ, Yang MG, Harmin DA, Tao P, Timms AE, Bauwens M, Allikmets R, Jones EM, Chen R, De Baere E et al. Mapping the cis-regulatory architecture of the human retina reveals noncoding genetic variation in disease. *Proc Natl Acad Sci U S A*. 2020;117(16):9001-9012.
7. Zhang F, Lupski JR. Non-coding genetic variants in human disease. *Hum Mol Genet.* 2015; 24(R1):R102-110.
8. Nik-Zainal S, Davies H, Staaf J, Ramakrishna M, Glodzik D, Zou X, Martincorena I, Alexandrov LB, Martin S, Wedge DC et al. Landscape of somatic mutations in 560 breast cancer whole-genome sequences. *Nature*. 2016;534(7605):47-54.
9. Devuyst O. The 1000 Genomes Project: Welcome to a New World. Perit Dial Int. 2015;35(7):676-677.
10. Collins FS, Fink L. The Human Genome Project. *Alcohol Health Res World*. 1995;19(3):190-195.

11.   O'Haire M. Companion animals and human health: Benefits, challenges, and the road ahead.
Journal of Veterinary Behavior. 2010;5(5):226-234.

12.   Meuwissen THE, Hayes BJ, Goddard ME. Prediction of total genetic value using
genome-wide dense marker maps. *Genetics*. 2001;157(4):1819-1829.

13.   Calus MP: Genomic breeding value prediction: methods and procedures. *Animal*.
2010;4(2):157-164.

14.   Brito LF, Clarke SM, McEwan JC, Miller SP, Pickering NK, Bain WE, Dodds KG, Sargolzaei
M, Schenkel FS. Prediction of genomic breeding values for growth, carcass and meat quality
traits in a multi-breed sheep population using a HD SNP chip. *BMC Genet*. 2017;18(1):7.

15.   Ma X, Christensen OF, Gao H, Huang R, Nielsen B, Madsen P, Jensen J, Ostersen T, Li P,
Shirali M et al. Prediction of breeding values for group-recorded traits including genomic
information and an individually recorded correlated trait. *Heredity (Edinb)*.
2021;126(1):206-217.

16.   Cappelloni M, Gallo M, Cesarani A. Use of threshold and linear models to estimate variance
components and breeding values for disease resistance in Italian heavy pigs. *Italian Journal of
Animal Science*. 2022;21(1):488-492.

17.   Lourenco DAL, Fragomeni BO, Tsuruta S, Aguilar I, Zumbach B, Hawken RJ, Legarra A,
Misztal. Accuracy of estimated breeding values with genomic information on males, females,
or both: an example on broiler chicken. *Genetics Selection Evolution*. 2015;47.

18.   Rentzsch P, Witten D, Cooper GM, Shendure J, Kircher M. CADD: predicting the
deleteriousness of variants throughout the human genome. *Nucleic acids research*.
2019;47(D1):D886-D894.

19.   Groß C, Derks M, Megens H-J, Bosse M, Groenen MA, Reinders M, De Ridder D. pCADD:
SNV prioritisation in Sus scrofa. *Genetics Selection Evolution*. 2020;52:1-15.

20.   Groß C, Bortoluzzi C, de Ridder D, Megens H-J, Groenen MA, Reinders M, Bosse M.
Prioritizing sequence variants in conserved non-coding elements in the chicken genome using
chCADD. *PLoS genetics*. 2020;16(9):e1009027.

21.   Gulko B, Hubisz MJ, Gronau I, Siepel A. A method for calculating probabilities of fitness
consequences for point mutations across the human genome. *Nature. genetics*.
2015;47(3):276-283.

22.   Ionita-Laza I, McCallum K, Xu B, Buxbaum JD. A spectral approach integrating functional
genomic annotations for coding and noncoding variants. *Nature genetics*. 2016;48(2):214-220.

23.   Consortium EP. The ENCODE (ENCyclopedia Of DNA Elements) Project. Science.
2004;306(5696):636-640.

24.   Consortium I. The Impact of Genomic Variation on Function (IGVF) Consortium. *arXiv
preprint arXiv*:230713708 2023.

25.   Boyle AP, Hong EL, Hariharan M, Cheng Y, Schaub MA, Kasowski M, Karczewski KJ, Park
J, Hitz BC, Weng S et al. Annotation of functional variation in personal genomes using
RegulomeDB. *Genome Res*. 2012;22(9):1790-1797.

26.   Fu YH, Liu H, Dou JW, Wang Y, Liao Y, Huang X, Tang ZS, Xu JY, Yin D, Zhu SL et al.
IAnimal: a cross-species omics knowledgebase for animals. *Nucleic Acids Research*. 2022.

27.   Teng J, Gao Y, Yin H, Bai Z, Liu S, Zeng H, Bai L, Cai Z, Zhao B, Li X. A compendium of
genetic regulatory effects across pig tissues. *bioRxiv*. 2022.

28.   Zhao Y, Hou Y, Xu Y, Luan Y, Zhou H, Qi X, Hu M, Wang D, Wang Z, Fu Y et al. A

compendium and comparative epigenomics analysis of cis-regulatory elements in the pig

genome. *Nat Commun*. 2021;12(1):2217.

29. Li H, Durbin R. Fast and accurate short read alignment with Burrows-Wheeler transform.

Bioinformatics. 2009;25(14):1754-1760.

30. Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, Marth G, Abecasis G, Durbin R,

Genome Project Data Processing S. The Sequence Alignment/Map format and SAMtools.

Bioinformatics. 2009;25(16):2078-2079.

31. Ramirez F, Dundar F, Diehl S, Gruning BA, Manke T. deepTools: a flexible platform for

exploring deep-sequencing data. *Nucleic Acids Res*. 2014;42(Web Server issue):W187-191.

32. Heinz S, Benner C, Spann N, Bertolino E, Lin YC, Laslo P, Cheng JX, Murre C, Singh H,

Glass CK. Simple combinations of lineage-determining transcription factors prime

cis-regulatory elements required for macrophage and B cell identities. *Mol Cell*.

2010;38(4):576-589.

33. Zhang Y, Liu T, Meyer CA, Eeckhoute J, Johnson DS, Bernstein BE, Nusbaum C, Myers RM,

Brown M, Li W et al. Model-based analysis of ChIP-Seq (MACS). *Genome Biol*.

2008;9(9):R137.

34. Quinlan AR, Hall IM. BEDTools: a flexible suite of utilities for comparing genomic features.

Bioinformatics. 2010;26(6):841-842.

35. Liu T. Use model-based Analysis of ChIP-Seq (MACS) to analyze short reads generated by

sequencing protein-DNA interactions in embryonic stem cells. *Methods Mol Biol*.

2014;1150:81-95.

36. Li Z, Kuo CC, Ticconi F, Shaigan M, Gehrmann J, Gusmao EG, Allhoff M, Manolov M,

Zenke M, Costa IG. RGT: a toolbox for the integrative analysis of high throughput regulatory

genomics data. *BMC Bioinformatics*. 2023;24(1):79.

37. Coetzee SG, Coetzee GA, Hazelett DJ. motifbreakR: an R/Bioconductor package for

predicting variant effects at transcription factor binding sites. *Bioinformatics*.

2015;31(23):3847-3849.

38. Khan A, Fornes O, Stigliani A, Gheorghe M, Castro-Mondragon JA, van der Lee R, Bessy A,

Cheneby J, Kulkarni SR, Tan G et al. JASPAR 2018: update of the open-access database of

transcription factor binding profiles and its web framework. *Nucleic Acids Res*.

2018;46(D1):D1284.

39. Servant N, Varoquaux N, Lajoie BR, Viara E, Chen CJ, Vert JP, Heard E, Dekker J, Barillot E.

HiC-Pro: an optimized and flexible pipeline for Hi-C data processing. *Genome Biol*.

2015;16:259.

40. Rao SS, Huntley MH, Durand NC, Stamenova EK, Bochkov ID, Robinson JT, Sanborn AL,

Machol I, Omer AD, Lander ES. A 3D map of the human genome at kilobase resolution

reveals principles of chromatin looping. *Cell*. 2014;159(7):1665-1680.

41. Hao Y, Stuart T, Kowalski MH, Choudhary S, Hoffman P, Hartman A, Srivastava A, Molla G,

Madad S, Fernandez-Granda C. Dictionary learning for integrative, multimodal and scalable

single-cell analysis. *Nature biotechnology*. 2024;42(2):293-304.

42. Hu C, Li T, Xu Y, Zhang X, Li F, Bai J, Chen J, Jiang W, Yang K, Ou Q. CellMarker 2.0: an

updated database of manually curated cell markers in human/mouse and web tools based on

scRNA-seq data. *Nucleic acids research*. 2023;51(D1):D870-D876.

43. Cingolani P. Variant Annotation and Functional Prediction: SnpEff. *Methods Mol Biol*.

2022;2493:289-314.

44.   Mrode RA, Thompson R. Linear models for the prediction of animal breeding values: CABI

publishing; 2005.

45.   Fu Y, Xu J, Tang Z, Wang L, Yin D, Fan Y, Zhang D, Deng F, Zhang Y, Zhang H et al. A gene

prioritization method based on a swine multi-omics knowledgebase and a deep learning model.

Commun Biol. 2020;3(1):502.

46.   Eichler EE. Genetic Variation, Comparative Genomics, and the Diagnosis of Disease. *N Engl*

J Med. 2019;381(1):64-74.

47.   Lee TI, Young RA. Transcriptional regulation and its misregulation in disease. *Cell*.

2013;152(6):1237-1251.

48.   Kern C, Wang Y, Xu X, Pan Z, Zhou H. Functional annotations of three domestic animal

genomes provide vital resources for comparative and agricultural research. *Nature*

Communications.

49.   Foissac S, Djebali S, Munyard K, Vialaneix N, Giuffra E. Multi-species annotation of

transcriptome and chromatin structure in domesticated animals. *BMC Biology*.

2019;17(1):108.

50.   Blackledge NP, Klose RJ. The molecular principles of gene regulation by Polycomb repressive

complexes. *Nature reviews Molecular cell biology*. 2021;22(12):815-833.

51.   Mitsis T, Efthimiadou A, Bacopoulou F, Vlachakis D, Chrousos GP, Eliopoulos E.

Transcription factors and evolution: an integral part of gene expression. *World Academy of*

Sciences Journal. 2020;2(1):3-8.

52.   Plaza A, Merino B, Sánchez-Pernaute A, Torres-García AJ, Rubio-Herrera MA, Ruiz-Gayo M.

Expression analysis of a cholecystokinin system in human and rat white adipose tissue. *Life*

sciences. 2018;206:98-105.

53.   Sun WenKui SW, Cheng Chi CC, Liu Rui LR, Chen YiHui CY, Zeng Kai ZK, Chen XiaoHui

CX, Gu YiRen GY, Li JiangLing LJ, Lv XueBin LX, Gao Rong GR. Expression patterns of

GHRL, GHSR, LEP, LEPR, SST and CCK genes in the gastrointestinal tissues of Tibetan and

Yorkshire pigs. 2019.

54.   Zhang K, Yuan Z, Bing Y, Chen X, Ding X, Chen D. Effects of active immunization against

cholecystokinin 8 on performance, contents of serum hormones, and expressions of CCK gene

and CCK receptor gene in pigs. *Endocrine*. 2007;32:338-344.

55.   Yonekura S, Kitade K, Furukawa G, Takahashi K, Katsumata N, Katoh K, Obara Y. Effects of

aging and weaning on mRNA expression of leptin and CCK receptors in the calf rumen and

abomasum. *Domestic Animal Endocrinology*. 2002;22(1):25-35.

56.   Cooper GM, Stone EA, Asimenos G, Green ED, Batzoglou S, Sidow A. Distribution and

intensity of constraint in mammalian genomic sequence. *Genome research*.

2005;15(7):901-913.

57.   Rubin C-J, Megens H-J, Barrio AM, Maqbool K, Sayyab S, Schwochow D, Wang C, Carlborg

Ö, Jern P, Jørgensen CB. Strong signatures of selection in the domestic pig genome.

Proceedings of the National Academy of Sciences. 2012;109(48):19529-19536.

58.   Per Madsen JJ. A user's guide to DMU. Version 6, release 5.2. 2013.

59.   Boix CA, James BT, Park YP, Meuleman W, Kellis M. Regulatory genomic circuitry of human

disease loci by integrative epigenomics. *Nature*. 2021;590(7845):300-307.

60.   Joslin AC, Sobreira DR, Hansen GT, Sakabe NJ, Aneas I, Montefiori LE, Farris KM, Gu J,

Lehman DM, Ober C. A functional genomics pipeline identifies pleiotropy and cross-tissue
effects within obesity-associated GWAS loci. *Nature communications*. 2021;12(1):5253.

61. Fang L, Cai W, Liu S, Canela-Xandri O, Gao Y, Jiang J, Rawlik K, Li B, Schroeder SG, Rosen
BD. Comprehensive analyses of 723 transcriptomes enhance genetic and biological
interpretations for complex traits in cattle. *Genome research*. 2020;30(5):790-801.

62. Takayanagi H. Osteoimmunology: shared mechanisms and crosstalk between the immune and
bone systems. *Nature Reviews Immunology*. 2007;7(4):292-304.

63. Mi B, Xiong Y, Knoedler S, Alfertshofer M, Panayi AC, Wang H, Lin S, Li G, Liu G.
Ageing-related bone and immunity changes: insights into the complex interplay between the
skeleton and the immune system. *Bone Research*. 2024;12(1):42.

64. Ropka-Molik K, Bereta A, Tyra M, Różycki M, Piórkowska K, Szyndler-Nędza M, Szmatoła
T. Association of calpastatin gene polymorphisms and meat quality traits in pig. *Meat science*.
2014;97(2):143-150.

65. Spielmann M, Mundlos S. Looking beyond the genes: the role of non-coding variants in
human disease. *Human molecular genetics*. 2016;25(R2):R157-R165.

66. Giral H, Landmesser U, Kratzer A. Into the Wild: GWAS Exploration of Non-coding RNAs.
Frontiers in Cardiovascular Medicine. 2018;5.

67. Okumura N, Kurosawa Y, Kobayashi E, Watanobe T, Ishiguro N, Yasue H, Mitsuhashi T.
Genetic relationship amongst the major non-coding regions of mitochondrial DNAs in wild
boars and several breeds of domesticated pigs. *Animal Genetics*. 2001;32(3):139-147.

68. Spielmann M, Mundlos S. Looking beyond the genes: the role of non-coding variants in
human disease. *Hum Mol Genet*. 2016;25(R2):R157-R165.

69. Cohen JC, Pertsemlidis A, Fahmi S, Esmail S, Vega GL, Grundy SM, Hobbs HH. Multiple
rare variants in NPC1L1 associated with reduced sterol absorption and plasma low-density
lipoprotein levels. *Proc Natl Acad Sci U S A*. 2006;103(6):1810-1815.

70. Hugot J-P, Chamaillard M, Zouali H, Lesage S, Cézard J-P, Belaiche J, Almer S, Tysk C,
O'Morain CA, Gassull M. Association of NOD2 leucine-rich repeat variants with
susceptibility to Crohn's disease. *Nature*. 2001;411(6837):599-603.

71. Buenrostro JD, Giresi PG, Zaba LC, Chang HY, Greenleaf WJ. Transposition of native
chromatin for fast and sensitive epigenomic profiling of open chromatin, DNA-binding
proteins and nucleosome position. *Nature methods*. 2013;10(12):1213-1218.

72. Halstead MM, Kern C, Saelao P, Wang Y, Chanthavixay G, Medrano JF, Van Eenennaam AL,
Korf I, Tuggle CK, Ernst CW et al. A comparative analysis of chromatin accessibility in cattle,
pig, and mouse tissues. *Bmc Genomics*. 2020;21(1).

73. Boyle AP, Davis S, Shulha HP, Meltzer P, Margulies EH, Weng Z, Furey TS, Crawford GE.
High-resolution mapping and characterization of open chromatin across the genome. *Cell*.
2008;132(2):311-322.

74. Zeng H, Zhang W, Lin Q, Gao Y, Teng J, Xu Z, Cai X, Zhong Z, Wu J, Liu Y. PigBiobank: a
valuable resource for understanding genetic and biological mechanisms of diverse complex
traits in pigs. *Nucleic acids research*. 2024;52(D1):D980-D989.

75. Zhang K, Liang J, Fu Y, Chu J, Fu L, Wang Y, Li W, Zhou Y, Li J, Yin X. AGIDB: a versatile
database for genotype imputation and variant decoding across species. *Nucleic Acids Research*.
2024;52(D1):D835-D849.

 Figure legends

 Figure 1. Workflow and genomic distribution of variants and epigenomic

 datasets collection across ten farm animals. (A) Overview of the workflow. (B)

 Proportion of variants distributed across coding and non-coding regions in the ten

 species. (C) Statistical summary of datasets categorized by tissue types across the ten

 species. The color of the inner circle represents the species, the middle circle

 represents tissues and cell lines, and the outer circle indicates the number of datasets

 corresponding to different data types. (D) The proportion of genomic variants within

 different functional regions of the ten species.

 Figure 2. Functional classification of genomic variants: approach and findings.

 (A) Schematic diagram illustrating how genomic variants influence transcription

 factor binding. (B) Design principles underlying the system for classifying variant

 functional confidence. (C) Proportions of variations from categories 1-5 across the ten

 species. (D) Signal distributions surrounding variants within the *CCK* gene in pigs

 and cattle. The top panel shows ATAC-seq and H3K27ac ChIP-seq signal peaks

 around the SNP rs332577759 (Category 2d) in heart tissue, alongside functional

 regions and interaction heatmaps in muscle. The lower panel displays rs519400784

 (Category 1d) signal intensity in cattle liver, with key functional regions highlighted

 in yellow.

 Figure 3. Conservation and impact of functionally significant variants. (A) GERP

 scores (from Ensembl) for variants across different functional categories in the pig

genome. (B) GERP scores for the 100 bp regions flanking the variants (bedtools slop

-b 100). (C) Proportion of variants overlapping with eQTLs. (D) Fisher's exact test

odds ratios for variants linked to production traits (data from Animal QTL), with

non-functional variants serving as the control. (E) Number of SNP markers used to

evaluate EBV. (F) Reliability of EBV predictions for traits like ADG and BF.

Figure 4. Variant activity in CRE and traits. (A) Proportions of CRE-linked SNP

activity across various pig tissues. (B) Heatmap showing the enrichment of production

traits based on the activity of trait-associated variants across pig tissues. (C)

Proportion of active variants linked to specific traits. Statistical significance was

determined using an unpaired Student's *t*-test. * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$,

**** $p < 0.0001$. (D) Enrichment of production traits across all cell types within

adipose. (E) T-distributed stochastic neighbor embedding (t-SNE) plot showing

scRNA-seq data from pig adipose tissue. (F) Expression levels of the *CAST* gene

across various cell types.

Figure 5. Overview of the IFmut database. (A) The IFmut homepage, displaying

information about the ten farm animals featured. (B) Graphic overview of the

modules available within IFmut. (C) Example of the variant category browser on the

"Category" page. (D) Clicking the "image" button in the "Affected motif" column

displays nucleotide conservation logos for TF recognition motifs potentially disrupted

by a variant. (E) The variant ID link redirects users to detailed variant information. (F)

Example of the epigenetics data browser for variants, available on the "Epigenetics"

page. (G) Browser showing TADs or loops encompassing the variant on the "3D

genome" page. (H) Browser the variant-related genes on the "Gene" page. (I) Trait

browser displaying traits associated with the variant on the "Trait" page. (J) Cell type

browser showing cell types linked to the variant on the "Single cell" page.

Supporting information

Supplementary Table S1. Number of regulatory regions detected in farm

animals.

Supplementary Table S2. Genomic coverage of regulatory regions in farm

animals.

Supplementary Table S3. Variant classification approach.

Supplementary Table S4. Predictive reliabilities of estimated breeding values for

traits ADG and BF.

Supplementary Fig. S1. Number of variants in different functional regions of 10

species. Including variants in the functional regions of motif, TF binding site, NFR,

footprint, OCR, H3K27ac peaks and the number of variants not in any of the

functional regions.

Supplementary Fig. S2. Statistics on the number of variants in categories 1-5 (12

subclasses) for 10 species.

Supplementary Fig. S3. Conservation calculations of functional variants and

non-functional variants for 6 species. There were significant differences in the

conservatism of functional variants and non-functional variants.

Supplementary Fig. S4. Conservation calculations of functional variants and

non-functional variants for 6 species. There were significant differences in the

conservatism of functional variants and non-functional variants.

Supplementary Fig. S5. Instructions for using the ifmutAnnotator tools in IFmut.

On the Category page of the IFmut database, we provide the ifmutAnnotator tools

annotation noval variant. The use of the ifmutAnnotator tool is divided into 3 steps,

step1, click on the browse button, step2, upload the bed file to be annotated, and step3,

download the annotation results.

A Proportions of CRE-Associated SNP Activity Across Multiple Tissues