Single-nucleus RNA sequencing reveals cellular and molecular dynamics of white and brown adipose tissue in a mouse model of type-2 diabetes

Taylah L. Gaynor,[1,2] **Ian Hsu,**[1] **Crisdion Krstevski,**[1,2] **Gabriella E. Farrugia,**[1,3,4] **Malathi S. I. Dona,**[1,3,4] **Charles D. Cohen,**[1] **Brian G. Drew,**[5,6,7] **Alexander R. Pinto***[1,2,3,4,6]

[1]Cardiac Cellular Systems; Baker Heart and Diabetes Institute; Melbourne, VIC, 3004; Australia
[2]Department of Microbiology, Anatomy, Physiology and Pharmacology of Agriculture, Biomedicine and Environment; La Trobe University; Bundoora, VIC, 3086; Australia
[3]Centre for Cardiovascular Biology and Disease/School of Agriculture, Biomedicine and Environment; La Trobe University; Bundoora, VIC, 3086; Australia
[4]Baker Department of Cardiometabolic Research, Translation, and Implementation; La Trobe University; Bundoora, VIC, 3086; Australia
[5]Molecular Metabolism and Ageing; Baker Heart and Diabetes Institute; Melbourne, VIC, 3004; Australia
[6]Baker Department of Cardiometabolic Health/ Melbourne Medical School; University of Melbourne; Parkville, VIC, 3010; Australia
[7]Central Clinical School, Monash University, Melbourne, VIC, 3004; Australia
*Correspondence: alex.pinto@baker.edu.au (A.R.P)

Summary

Excessive adipose tissue expansion is often linked with type-2 diabetes. Despite recent efforts mapping adipose tissue changes in obesity using single-cell omics, an understanding of cellular and gene expression changes in a model of type 2 diabetes, and the transcriptional circuitry controlling it, is still lacking. Here, we use single-nucleus RNA sequencing to analyze the remodeling of gonadal white and interscapular brown adipose tissue from female and male mice with or without diabetes. Analysis of 51,877 nuclei revealed altered phenotypes in every cell population in type 2 diabetes. This included an immunoregulatory response, and changes in extracellular matrix, energy generation, and hormone responses. Key transcription factors were inferred as cell-specific and non-specific nodes controlling diabetes-linked phenotypes. Finally, female-to-male population heterogeneity and gene expression differences were observed. Here we provide a resource detailing how adipose tissue remodeling, and the molecular mechanisms governing it, may contribute to cardiometabolic disease.

Keywords – Adipose, adipocytes, single-cell, single-nucleus, transcriptomics, diabetes, single-nucleus RNA sequencing, flow cytometry, microscopy

Introduction

Cardiometabolic diseases, such as type-2 diabetes mellitus (T2D), have been linked with excessive
expansion of adipose tissue[1]. Adipose depots can be broadly classified as white and brown adipose tissue
(WAT and BAT, respectively), with WAT being recognized as an endocrine organ having a primary function
in lipid storage while BAT drives thermogenesis. The anatomical location of each adipose depot can dictate
the tissue's impact systemically[2]. Specifically, the expansion of visceral WAT can exert pathogenic effects
through the release of pro-inflammatory adipokines[3]. Conversely, subcutaneous WAT and interscapular BAT
can be protective[4,5] with research actively focusing on inducing 'browning' in WAT to improve energy
expenditure and metabolic outcomes[6].

Many single-cell and spatial transcriptomics studies have provided detailed characterization of adipose depot
cellular heterogeneity[7–13] including how this is altered in obesity[14–21]. Moreover, disease-specific cell
populations in obese WAT from individuals with and without T2D have been identified[7,21]. Yet, despite these
advancements, a comprehensive analysis of both WAT and BAT in an animal model of T2D, without the
influence of confounding variables, is lacking. Further, with biological sex being more widely recognized as a
factor influencing disease processes including cardiovascular disease[22], the role of sex in adipose tissue
remodeling remains unclear.

To study the remodeling of WAT and BAT in T2D, we used single-nucleus transcriptomics to profile the entire
adipose cellular landscape of female and male mice with or without T2D (*db/db* and *db/h,* respectively).
Single-nucleus RNA-sequencing (snRNA-seq) analyses of these depots revealed complex cellular
landscapes, identifying important cell-specific gene expression patterns and regulatory circuits that govern
cell identity and phenotype. We found that T2D significantly alters the relative abundance of adipose cell
types and gene expression of virtually every adipose cell type, resulting in inflammation and tissue
remodeling. Further, we identified sex-specific differences in gene expression and cellular abundance,
providing a possible basis for varying adipose accretion. Together, these findings provide a valuable resource
for understanding WAT and BAT cellular landscapes, their changes in T2D and their potential contribution to
metabolic disorders.

Results

Leptin-receptor deficient *db/db* mice display hyperglycemia and increased fat mass

To compare adipose depots and characterize adipose remodeling induced by diabetes, we analyzed both
female and male *db/h* and leptin-receptor deficient *db/db* mice. To validate hyperglycemia in *db/db* animals,
blood glucose levels over an extended time were measured using an HbA1c test (Supplementary Table 1).
Higher blood glucose levels were seen in *db/db* mice when compared to *db/h* controls across both females
and males (*P*<0.0001 for both) with no sex-specific differences being observed between the genotypes. Next,
EchoMRI measurements of lean and fat mass relative to total body weight were taken to assess body
composition. EchoMRI showed a greater lean mass in males across both *db/h* and *db/db* genotypes
(*P*=0.0036 and *P*=0.0042, respectively). Both sexes had lower lean mass and greater fat mass in *db/db*
counterparts (*P*<0.0001 for all). Finally, *db/db* females had greater fat mass than male counterparts
(*P*=0.0017). These findings highlight T2D-related hyperglycemia and increased fat mass in *db/db* mice.

Novel high throughput nuclei isolation technique to characterize the adipose cellulome

Single-cell and spatial transcriptomics has been instrumental in shedding new light on tissue cellularity and
disease responses in multiple tissues, including adipose[7–11,16–19,21]. Due to the lipid-filled nature of mature
adipocytes, a key approach for single-cell transcriptomic profiling of adipose involves crushing tissue to
extract nuclei using a dounce homogenizer[12,13,19]. While this technique has proven useful for nuclei extraction,
a limitation of this approach is the number of samples that can be processed promptly. To address this issue,
we developed a nuclei isolation technique to isolate both mature adipocyte and stromal vascular fraction
(SVF) nuclei from samples in a high-throughput manner. The technique involves the machine-based
mechanical dissociation of snap-frozen adipose tissue, and subsequent removal of cellular debris using
fluorescence-activated cell sorting. BAT and WAT were homogenized at 30 Hz for 30 and 50 seconds,
respectively, in 10-second intervals to isolate the greatest number of nuclei (Supplementary Figure 1A). As
a result, this technique produced high-quality, intact nuclei that enabled the identification of most adipose
tissue cell populations through snRNA-seq (Figure 1A-C; Supplementary Figure 1A-B).

snRNA-seq profiling and flow cytometry reveal cellular heterogeneity of adipose depots

Using the aforementioned protocol to isolate nuclei for subsequent snRNA-seq, we were able to generate a
dataset containing 51,877 nuclei comprising 22,526 nuclei from gonadal WAT and 29,351 nuclei from
interscapular BAT after quality control filtering (Figure 1B). WAT and BAT demonstrated distinct cellular
compositions in both *db/h* and *db/db* conditions (Figures 1B-C). Cell populations were identified using their
top enriched genes and established marker genes uncovering the cellular heterogeneity of both tissues
(Figures 1D-E). Populations identified were macrophages (*Adgre1*, *Fcgr1*, *Csf1r, Cd68, Mrc1*), including a
proliferating subpopulation (*Top2a*, *Aurka*, *Mki67*), lymphocytes (*Thy1*, *Cd19*, *Cd5*, *Cd3e*), endothelial cells
(ECs; *Pecam1*, *Ly6c1*), lymphatic endothelial cells (LECs; *Cldn5, Prox1*), mesothelial cells (*Msln*), smooth
muscle cells (SMCs; *Acta2*, *Myh11*), Schwann cells (*Plp1*, *Kcna1*), pericytes (*Pdgfrb*, *Vtn*), multipotent
progenitors (*Cd34*, *Dpp4*, *Pi16*, *Creb5*), white adipocytes (*Slc7a10*), and brown adipocytes (*Ucp1*, *Cidea*).
Brown adipocytes identified in WAT likely correspond to beige or brite adipocyte subpopulations but are
referred to collectively as brown adipocytes throughout analysis. Finally, as fibroblasts and adipocyte
progenitors/pre-adipocytes share many of the same marker genes, cells expressing *Pdgfra* and *Col1a1* were
collectively referred to as fibroblasts. Epididymal cells (*Adam7*, *Pemt*, *Abcb5*, *Ces5a*), ovarian epithelial cells
(*Pax8*, *Epcam*, *Krt7*, *Cdh1*) and skeletal muscle cells (*Ttn*, *Myl1*) were identified and subsequently removed
from analysis alongside clusters with high mitochondrial RNA content and low RNA counts (see Methods).

1 **Adipose cell populations show tissue-dependent compositional and transcriptional individuality**

2 To next compare and validate the cellular compositions of *db/h* WAT and BAT, flow cytometric analysis was
3 conducted on SVF cell populations with mature adipocytes being excluded due to their fragile and buoyant
4 characteristics (Figure 2A; Supplementary Figure 1C). Populations were labelled according to their cell
5 surface marker expression (Supplementary Table 1). Due to challenges in distinguishing adipocyte
6 progenitors, preadipocytes, and fibroblasts from one another, in addition to the lack of clear cell surface
7 markers for these populations[23], they were collectively labelled as fibroblasts based on their negative
8 expression of mural cell markers. Distinct compositional differences were observed between WAT and BAT.
9 This included variations in most immune and mural cell populations with higher contributions to the viable cell
10 count being observed in WAT when compared to BAT. Of note, vascular EC (VEC) contribution was highest
11 in BAT when compared to WAT, while LECs showed the opposite trend ($P<0.0001$ and $P=0.0061$,
12 respectively). Histological validation and quantification of ECs using isolectin B4 (IB4) showed a greater
13 percentage of IB4 positive (or vascularized) tissue area in BAT corroborating the flow cytometry VEC findings
14 ($P<0.0001$), however, vascularization area per nucleus (stained with DAPI) was not different between depots
15 (Figure 2B).

16 To assess the gene expression differences between corresponding cell types from WAT and BAT, we
17 performed differential gene expression (DGE) analysis using our snRNA-seq data from *db/h* mice. Our
18 analysis found depot-dependent variations in gene expression across all cell populations with white and
19 brown adipocytes, fibroblasts, and ECs showing the most marked differences (Figure 2C). In BAT, the
20 upregulation of up to 2216 genes in brown adipocytes, 2164 genes in ECs, 1914 genes in fibroblasts, and
21 2880 genes in white adipocytes was observed and, in WAT, the upregulation of up to 3262 genes in brown
22 adipocytes, 3124 genes in ECs, 2507 genes in fibroblasts, and 3275 genes in white adipocytes was seen.
23 Gene ontology (GO) term analysis of upregulated and downregulated differentially expressed genes (DEGs)
24 uncovered the upregulation of biological pathways in BAT cell populations including *generation of precursor*
25 *metabolites and energy*, *oxidative phosphorylation*, and *ribose phosphate metabolic process* emphasizing
26 the heat-producing role of this tissue (Figure 2C). GO terms enriched in WAT included *actin filament*
27 *organization*, *cell substrate adhesion,* and *small GTPase mediated signal transduction*. GTP-mediated RHO
28 signaling has long been implicated in insulin-mediated glucose uptake and homeostasis in adipocytes,
29 highlighting the endocrine role of this tissue[24]. These findings highlight the role of adipose depot-specific
30 microenvironments in programming phenotypes of individual cell populations.

31 **Adipocyte subpopulation profiles differ between disparate adipose depots**

32 With white and brown adipocytes being present in both WAT and BAT, we subset these nuclei from the
33 dataset and conducted DGE analysis between tissues to determine possible depot-dependent roles (Figure
34 2D). White adipocytes in BAT showed greater expression of genes including *Fabp3,* of which deficient mice
35 have extreme cold sensitivity[25]. They also showed higher expression of *Ntrk3* (TRKC), a key receptor
36 maintaining cold tolerance and regulating innervation and sympathetic neuron growth in adipose tissue[26].
37 These genes suggest a possible role for white-like adipocytes in sympathetically activated BAT thermogenic
38 processes. In WAT, white adipocytes were enriched for *Lcp1*, a gene involved in leptin and glucose
39 homeostasis, and in the inhibition of WAT 'browning'[27]. Expectedly for brown adipocytes in BAT, greater
40 expression of thermogenesis-related genes including *Ucp1*, *Cidea*, *Pparg, Ppara,* and *Ppargc1b* were
41 observed while brown-like adipocytes in WAT were enriched for *Nnat, Aldh1a1*, *Cers6* and *Nrip1* (RIP140)*,*
42 genes, that when suppressed, appear to improve metabolic health via thermogenic pathways[28–31]. GO
43 enrichment analysis of DEGs uncovered pathways such as *generation of precursor metabolites and energy*,
44 *adaptive thermogenesis,* and *oxidative phosphorylation*, among others, as upregulated in adipocytes
45 (irrespective of subtype) from BAT reflecting its heat-producing capacity (Figure 2D). While in WAT, both
46 subpopulations showed expression of genes related to hormonal responses. In addition, terms relating to
47 transforming growth factor beta (TGFβ) and WNT signaling were seen. These data underscore the
48 importance of TGFβ and WNT in driving progenitor proliferation and suppressing adipogenesis[32–34]. WNT
49 signaling GO terms involved genes such as *Tle3* (Supplementary Table 2). TLE3 is a transcription factor co-
50 factor that is selective for white adipocytes by blocking the interaction between PRDM16 and PPARγ and

1 impairing thermogenesis[35]. Together these findings highlight the impact of the cellular landscapes of WAT
2 and BAT on adipocyte phenotypes.

Distinct gene regulatory circuitry govern brown and white adipose tissue cell phenotypes

Next, we determined the key gene regulatory elements that underlie differences between WAT and BAT and
the cells within them. To achieve this, we applied the *single-cell regulatory network inference and clustering*
(SCENIC) pipeline[36]. In brief, SCENIC links single-cell sequencing and cis-regulatory data to infer
transcriptional modules—regulons—comprising transcription factors (TFs) and their target genes. Cells were
first clustered based on their expression of regulons and corresponding gene regulatory networks (GRNs;
denoted as TF gene name and '(+)') showing clear clustering patterns that aligned with the previous RNA
annotations (Figure 3A; Figure 1B). Moreover, examination of the top 5 regulon markers for each cell
population in WAT and BAT demonstrated distinct top GRNs governing cell population phenotypes for brown
adipocytes and mesothelial cells, while ECs had shared marker GRNs between tissues with varying levels
of similarity observed for other cell populations (Figure 3B). Examples of regulons include Erg(+) and Tal1(+),
both important for EC homeostasis and angiogenesis[37,38], Tfec(+) and Irf5(+), both implicated in macrophage
functions and polarisation[39,40], Meox1(+) and Smad3(+), which govern fibroblast activation[41], and TGFβ-
induced fibrosis[42]. Finally, Esrrg(+), Esrra(+), and Ppara(+) are all implicated in thermogenesis and were
observed as markers for brown adipocytes. Altogether these data show an agreement between RNA-seq-
based GRN inferences with established transcriptional regulators in various cell populations. This analysis
highlights how GRN activity differs between adipose depots to yield tissue-specific cellular phenotypes.

Inflammation and macrophage accumulation in type-2 diabetes adipose depots

To next identify the impact of T2D on WAT and BAT we compared the cellular abundance of *db/h* and *db/db*
samples for each depot. Amongst the most striking changes was the increase in macrophages across both
WAT and BAT from *db/db* depots (Figure 4A) agreeing with statistical analysis of snRNA-seq counts and flow
cytometry data (Supplementary Figure 2). These cellular changes accompanied lipid droplet expansion in
adipocytes across both depots of *db/db* mice (Figure 4B). DGE analysis of WAT and BAT between *db/h* and
db/db tissues identified extensive gene expression differences in virtually all cells, with macrophages
amongst those exhibiting the greatest shifts in expression (Figure 4C). In macrophages, the top 3 GO terms
associated with upregulated genes for each cell population included *phagocytosis* and *leukocyte migration*
(Figure 4C-D). A possible consequence of lipotoxicity and adipocyte hypertrophy is the formation of
macrophage crown-like structures (CLSs). CLSs can accumulate in dysfunctional adipose depots through
macrophages clustering around dead or dying adipocytes to clear lipid and cell debris[43]. Recent work has
highlighted the importance of macrophage proliferation in the formation of these structures and subsequent
inflammation[44,45]. Indeed, histological validation using phosphohistone H3 validated the presence of
proliferative cells in the CLSs of WAT corroborating the snRNA-seq findings and affirming that macrophage
crowns exist in adipose tissue from T2D mice (Figure 4E). Moreover, subpopulation-specific expression of
proliferative markers *Top2a*, *Mki67*, and *Aurka* were observed (Figure 4F). Finally, as it is challenging to
attribute adipose tissue pathogenesis solely to a T2D-derived phenotype, we assessed the similarity of gene
expression patterns independent of obesity between our dataset and a previously published human obesity
dataset[21]. Approximately 47% of DEGs across all cell populations in *db/db* mice were unique to our dataset
and not differentially expressed in human obesity (Figure 4G).

Type-2 diabetes disrupts cell phenotypes altering adipose tissue homeostasis, energy generation and endocrine functions

Other cell populations that displayed substantially altered gene expression patterns in T2D included white
and brown adipocyte subpopulations and fibroblasts (Figure 4C-D). Notable among top GO terms
corresponding to upregulated genes in both BAT and WAT across multiple cell populations included
extracellular matrix organization and *extracellular structure organization* suggesting tissue remodeling.
Downregulated terms included *generation of precursor metabolites and energy*, and *oxidative*
phosphorylation suggesting decreased energy utilization and possible whitening of classically thermogenic
BAT. Genes that were associated with these terms included proteins essential for oxidative phosphorylation

such as ATP synthase or cytochrome bc 1 complex (Supplementary Table 3). In WAT, downregulated terms
included *cellular response to hormone stimulus*, and *hormone-mediated signaling pathway* indicating a
possible dampened hormone response in the tissue. Further, among the top 50 up- and downregulated genes
were lysosomal proteases (cathepsins; Cts genes) and leukocyte immunoglobulin-like receptor (LILR) genes
including *Lilr4b*, *Pirb* (*Lilrb3*), and *Lilrb4a* with the latter being the only LILR gene in the top 50 differentially
expressed of *db/db* BAT (Supplementary Figure 3). Moreover, in *db/db* BAT, the downregulation of heat
shock proteins (HSP genes) additionally suggests decreased thermogenesis.

snRNA-seq derived gene regulatory network inferences in type-2 diabetes

We next sought to determine which GRNs within each depot were driving the changes in adipose tissue cell
subsets in disease, focusing on adipocytes, fibroblasts, and macrophages (Figure 5A). Ranking transcription
factors based on activity, we found that the top 5 GRNs of these cell types in *db/h* mice were almost always
unique across depots. These were altered in *db/db* counterparts and exhibited signatures consistent with the
activation of known gene expression programs. For example, Cebpd(+), which is involved in cell apoptosis,
proliferation and adipocyte differentiation, was amongst the top-ranked in adipocytes, fibroblasts and
macrophage populations in T2D. Additionally, Irf4(+), a critical regulator of lipid handling and of which
deficiency induces macrophage-related inflammation[46], was amongst the top-ranked in BAT *db/h* samples
and less so for *db/db* counterparts. Nevertheless, immunohistochemical analyses detected IRF4 in BAT from
db/h and *db/db* animals (Figure 5B) despite our GRN analysis. These findings suggest possible reduced
transcriptional activity and impaired lipid handling in diabetic BAT.

Next, differentially enriched regulons were assessed. Regulons that showed common changes across both
depots included Tfeb(+), Klf13(+), Hmga2(+), and Atf3(+); all being upregulated in a disease context (Figure
5C). Regulons including Tfec(+) and Tfeb(+), among others, showed marked upregulation, particularly in
macrophages. TFEB is a transcription factor that plays an essential role in the biogenesis of lysosomal
machinery aligning with the upregulated lysosomal protease (Cts gene) expression observed above[47]
(Supplementary Figure 3). To further dissect the influence of transcriptional regulation on the concerted
immune response observed (Figure 4), gene targets linked to the GO term *leukocyte migration* were mapped
to transcription factors (Figure 5C). Fli1(+), which plays an important role in cell proliferation, activation,
differentiation, survival, and the regulation of chemokines[48], showed the greatest link to this *leukocyte*
migration with 92 target genes being associated with this regulon. Other notable regulons included Ikzf1(+)
and Spi1(+) which showed 74 and 68 genes linked to this GO term, respectively. Together, these findings
highlight the importance of transcriptional regulation for processes, such as inflammation, in T2D adipose
depots and highlight possible avenues for targeting TF candidates for therapeutic purposes.

Biological sex differences in classical adipose depots relating to adipocyte subpopulations

Given the considerable sex differences that exist in obesity and diabetes-associated disorders, we explored
potential sex-dependent mechanisms regulating adipose tissue homeostasis and diabetes-induced
remodeling. Female and male samples were barcoded using a previously described cholesterol-conjugated
oligonucleotide approach[49] (Supplementary Table 1). Samples were separated downstream based on
barcodes. Cells with undetectable barcodes were delineated using sex-specific gene expression (Females:
Xist; Males: *Ddx3y*, *Eif2s3y*, *Gm29650*, *Kdm5d*, *Uty*)[22].

Comparison of cells from females and males across each depot uncovered extensive sex-specific
differences. Sex-specific gene expression analysis found that within WAT, mesothelial cells, fibroblasts,
macrophages and white adipocytes exhibited sex-specific gene expression (Figure 6A). In BAT, gene
expression variation was primarily within adipocyte subpopulations. When the cellular composition of these
depots was considered, females and males had differing proportions of adipocyte subpopulations (Figure
6B). Specifically, *db/h* female BAT had high proportions of BA1 and BA2 cells, while *db/h* male BAT had
higher proportions of BA3 and BA4. Assessment of key genes involved in thermogenesis showed higher
expression levels in female *db/h* BAT while these dimorphisms were not as evident in a diabetic context
(Figure 6C). Analysis of the top 5 GO terms for brown adipocytes in *db/h* BAT provided some insight into
biological pathway differences showing female upregulation of *adaptive thermogenesis*, *cellular response to*

hormone stimulus and *fatty acid metabolic process* in females, and upregulation of *protein folding*, *oxidative*

phosphorylation, *ATP metabolic process,* and *proton motive force-driven ATP synthesis* pathways in males

(Figure 6D). Genes linked to these pathways encoded ATPase subunits, and heat shock proteins in males

while in females they encoded thermogenesis-related genes, including *Ppargc1a* and *Ucp1*, alongside *insulin*

receptor substrate 1 (*Irs1*) (Supplementary Tables 4 and 5). These findings align with the long recognized

increased thermogenic capacity of female BAT driven by mitochondrial size, abundance and UCP1 content[50–53].

Higher cell density and vascularization in female BAT

Finally, as many endothelial-related GO terms were noted in the analysis for female BAT we sought to assess

vascular differences between sexes (Figure 6D). Efficient blood flow to the tissue is essential for the supply

of free fatty acids for energy metabolism and generation. Differences between biological sexes have

previously been observed in gonadal WAT[54,55] yet, to date, no sex differences in BAT vascular networks have

been characterized. Staining of female and male *db/h* BAT showed more cells and IB4+ vasculature in

females (*P*=0.0091 and *P*=0.0206, respectively; Figure 6E-F). However, normalization of IB4 staining to the

number of nuclei revealed that the ratios were similar. These data indicate that, while female tissue may be

more densely packed, vascularization-to-cell ratios remain comparable. Strikingly, cell and vascularization

density sex differences were lost in T2D BAT alongside reduced gene expression differences across

adipocyte populations (Figure 6E-F). Together, these data highlight that BAT sex differences that exist in a

healthy state may be dampened or lost in disease.

1 Discussion

Pathogenic adipose tissue expansion is a well-established risk factor for cardiometabolic disease. Recent
research has been directed towards understanding how the deterioration of adipose tissue can contribute to
disease in both an obese and high-fat diet context at a single-cell level[14–21]. Limited T2D-focussed human
work has characterized insulin-regulating adipocyte subpopulations and molecularly distinct preadipocyte
populations associated with diabetes[7,13,21]. However, despite this, we are still lacking a precise understanding
of the cellular and molecular underpinnings of how these tissues change in the context of T2D. To address
this, we conducted snRNA-seq on classical adipose depots including gonadal WAT and interscapular BAT
from both *db/h* and *db/db* female and male mice.

Characterization of these depots using snRNA-seq showed that while similarity in cellular heterogeneity
existed between depots, the contribution of various cell subsets to the overall tissue cellularity differed.
Moreover, gene expression patterns differed across WAT and BAT highlighting the distinct functions of these
tissues. WAT is classically recognized for its lipid-storing functions, while BAT is known for its heat-producing
and energy-expending role. Analysis of adipocyte subtypes highlighted that cell populations have unique
functions across depots with BAT populations upregulating metabolic processes and WAT populations
upregulating WNT-related pathways. WNT signaling has important adipose-specific functions including
regulating lipogenesis, inhibiting adipogenesis[56], and the modulation of structural WNT-regulated adipose
tissue-resident cells, termed SWAT cells, which can move toward both adipogenic or progenitor-like states[57].
Finally, the assessment of GRN inferences validated the observed depot-specific gene expression patterns.

Our analysis of non-diabetic and T2D mice also revealed several features of pathogenic remodeling of
adipose. These include an increase in macrophage abundance across both WAT and BAT, consistent with
observations by others[14,16–21]. Our findings align with the more recent recognition of proliferative macrophage
importance in CLS formation[44,45,58,59] and highlight their presence in a model of T2D. Assessment of GRN
dynamics in T2D also showed a notable increase in TFEB transcription factor activity in macrophage subsets.
TFEB is an important transcription factor in the biogenesis of lysosomal machinery[47,60]. Lysosomal
degradation of hypertrophic adipocytes is a vital function of macrophage CLSs where these phagocytic cells
secrete lysosomal enzymes to create an environment wherein dying adipocytes are degraded through a
process known as exophagy. Lysosomal proteases are vital for this process with CTSB being specifically
identified as key for perilipin 1 (PLIN1; a lipid-droplet coating protein) degradation[61–63]. Indeed, these
lysosomal proteases are upregulated in *db/db* samples (most predominantly in macrophages) which was
further corroborated through Tfeb(+) expression –TFEB being a known regulator of Cts genes[47].

In line with the noted adipocyte lipid droplet expansion, BAT showed downregulation of energy generation in
T2D. BAT whitening has gathered much attention in the literature due to the hijacking of BAT-driven energy
expenditure to improve metabolic health. A wealth of research has been directed into exploring therapeutic
options to increase this expenditure including the use of cold exposure, exercise, and pharmacological and
dietary interventions[6]. Assessment of genes showing decreased expression in T2D included heat shock
proteins. Moreover, GO analysis of downregulated genes showed enrichment of genes related to vital
oxidative phosphorylation complexes, further supporting the notion of BAT moving into a lipid-storing state.

Finally, previous research has highlighted biological sex differences between adipose depots, including
functionality differences in BAT. Female BAT has greater mitochondrial abundance and size, in addition to
higher mRNA and protein levels of UCP1 and other thermogenic genes[50–53,64–66], which aligns with our data.
Further, although single-cell datasets have shown brown adipocyte populations with differing heat-producing
capacities[10,67,68], sex differences have not been characterized. Here we found that differences exist in
adipocyte subpopulation abundance across sexes. Moreover, histological analysis showed greater cellular
and vascular density in female BAT, with differences only being identified in gonadal WAT in the past[54,55].
VEGFA is long linked to BAT activity, with *Vegfa* deletion driving BAT whitening resulting in decreased
expression of thermogenic gene *Ucp1* and *Ppargc1a*[69]. Moreover, estrogen receptor 1 is known to regulate
VEGFA in adipose tissue possibly driving the sexual dimorphism[70]. However, T2D appeared to abrogate
these sex differences in vascular and gene expression.

While this study offers new insights and a resource for studying adipose, several limitations require noting.
First, the model of T2D used in this study (*db/db*) is a leptin receptor mutant meaning changes in depots may
be driven by the loss of this signaling rather than the diabetic phenotype. While we compared our data to that
of humans with obesity, data from this study should be further compared across other models of T2D. Second,
it is important to note that *db/db* mice are characterized by obesity, thus findings in the study may not be
strictly attributed to T2D alone. It has been shown, however, that *db/db* mice appear more 'diabetic' than
ob/ob mice and have less liver failure[71]. Third, mouse adipose depots, particularly BAT, inconsistently
recapitulate the human context. Thermoneutral BAT is more similar to human BAT and could be used in
future studies, particularly those considering biological sex differences[72,73]. Finally, as this dataset was
produced using nuclei, cytosolic mRNA is missing and therefore important transcriptomic information could
be absent from our dataset.

Overall, this study offers a unique insight towards adipose tissue biology and the multifaceted mechanisms
driving its maladaptive remodeling in T2D. It also informs our understanding of the impact of biological sex in
adipose homeostasis and disease. In addition to the single-cell transcriptomic resource presented here,
further analysis of different models, time points and genetic backgrounds will advance our knowledge of
adipose remodeling, potentially pointing to mechanisms that may be therapeutically manipulated to address
tissue dysfunction induced by metabolic disorders such as T2D.

Research design and methods

Animals, physiological assessment, and tissue isolation

All animal-related experiments were approved by the Alfred Research Alliance (ARA) Animal Ethics
Committee (Ethics number: E/1990/2020/B) and were performed in accordance with the National Health and
Medical Research Council of Australia. 20-week-old female and male B6.BKS(D)-Lepr[db/J] (*db/db;* referred to
as T2D) (strain #:000697, The Jackson Laboratory) or heterozygous littermates (*db/h;* referred to as ND) in
the C57BL/6J background[74–76] were housed under 12h light/dark cycles at 22°C with access to both food and
water *ad libitum*. Homozygous *db/db* mice are a well-established model of T2D, having a spontaneous
autosomal recessive global knockout for the leptin receptor. At 19-weeks of age, whole-body composition
analysis was performed on mice using Echo-MRI™ 4-in-1 700 Analyser (EchoMRI, Houston, TX, USA). Long-
term blood glucose levels were measured via HbA$_{1c}$ at endpoint (Cobas b 101 POC system, Roche
Diagnostics). At 20 weeks of age, mice were administered an intraperitoneal injection of heparin (100 units
per mouse) 20-30 minutes prior to euthanasia using CO_2 inhalation, which was followed by subsequent
cardiac exsanguination. As described previously[77], the thoracic cavity was exposed, and the right atrium was
cut. The pericardium was removed, and the inferior vena cava was cut. The apex of the left ventricle was
pierced using a 30-gauge hypodermic needle for perfusion with buffers specified below.
17

Stromal Vascular Fraction Flow Cytometry Processing

Mice were perfused with cold ethylenediaminetetraacetic acid (EDTA) at a rate of 2 mL/min/mouse before
gonadal white adipose and interscapular white and brown adipose depots were excised and snap frozen.
Tissues were washed briefly in clean PBS with 0.9 mM $CaCl_2$ (C4901, Sigma) before being minced using
curved scissors into ~2 mm pieces in a petri dish and transferred into 3 mL of enzymatic solution (2 mg/mL
collagenase IV (LS004188, Worthington Biochem) and 1 mg/mL dispase II (04942078001, Roche) in PBS
with 0.9 mM $CaCl_2$). Samples were triturated three times in 15-minute intervals using a P1000 pipette for a
total of 45 minutes then centrifuged at 400xg for 4 minutes at 4°C. The floating fraction was then carefully
removed using a P1000 pipette and discarded. Using the remaining supernatant, the pellet was resuspended
before being filtered through a 70 μm filter (43-50070-50, pluriSelect) into a tube containing PBS with 0.9 mM
$CaCl_2$. The sample was pulled through the filter twice, followed by the addition of count beads (C36950,
Invitrogen) before centrifugation at 200xg for 15 minutes at 4°C - no breaks - for debris clearance. The
supernatant was removed carefully, and the pellet was resuspended in Fx buffer (1XHBSS (14185-052,
Gibco), 2% FCS in ddH2O (10977015, Invitrogen)) before centrifugation at 400xg for 4 minutes at 4°C. The
supernatant was removed, and the pellet was resuspended in 50 μL of Fx buffer. A volume of 43μL for each
sample was transferred to 1.5 mL tubes containing Fc block (101320, BioLegend) on ice for 5 minutes and
the remaining samples were pooled and used for single color controls. Samples were then incubated in tubes
containing antibody mix (Supplementary Table 1) on ice for 20 minutes covered from light. Calcein Blue
(1:500; 65-0855-39, Invitrogen) was added to each sample and incubated at 37°C for 10 minutes. Samples
were washed using Fx buffer before centrifugation at 400xg for 4 minutes at 4°C. Supernatant was removed
and samples were resuspended in Sytox Green (1:1000; S34860, Invitrogen) in Fx buffer before being filtered
into FACS tubes (38030, StemCell) and analyzed on a BD Fortessa II flow cytometer.

Statistical Analysis

FlowJo™ V10.8 Software (for Windows BD Life Sciences) was utilized to compensate and analyze flow
cytometry data. A detailed gating strategy is outlined in Supplementary Figure 1C. For sex-independent
analyses within each depot, raw data were normalized to the average value for all *db/h* samples for that
batch. For sex-dependent analyses, the prior was conducted with only female samples for each batch.
GraphPad Prism v9.2.0 (GraphPad Software San Diego, California USA, www.graphpad.com) was
subsequently used to analyze data and conduct statistical analyses. Outliers were excluded using GraphPad
Prism's inbuilt ROUT outlier identification method (Q=1%) on data within each cell population. For between
depot analysis, data were treated as non-parametric and analyzed using a Kruskal-Wallis test. For within

depot analysis of T2D changes and sex differences, data were treated as non-parametric and analyzed using
a Mann-Whitney test. *P* values less than 0.05 were considered statistically significant.

Nuclei extraction

Mice were perfused with cold ethylenediaminetetraacetic acid (EDTA) at a rate of 2 mL/min/mouse before
gonadal white adipose and interscapular white and brown adipose depots were excised and snap frozen.
Frozen adipose tissue samples (ranging from ~25 to 34.7 mg for WAT and ~60.9 to 65.5 mg for BAT) were
mechanically homogenized in a previously published[78] nuclei purification buffer (NPB) (containing 5% bovine
serum albumin, 0.2% Nonidet™ P 40 Substitute (74385, Sigma-Aldrich), 1 mM dithiothreitol (D0632, Sigma-
Aldrich), 1X cOmplete™ EDTA-free Protease Inhibitor Cocktail (11873580001, Roche), 0.2 U/µL of Protector
RNase Inhibitor (3335399001, Roche) in Dulbecco's PBS (14190-250, Gibco)) with a 5 mm diameter
stainless steel ball (Qiagen, 69989) in Eppendorf Safe-Lock LoBind tubes (0030120.094, Eppendorf) using
a Qiagen TissueLyser II. BAT and WAT were homogenized at 30 Hz for 30 and 50 seconds, respectively, in
10-second intervals (Supplementary Figure 1A). Samples were placed on ice post-homogenization and
incubated for 15 minutes. Homogenized samples were then pulled through 20 µM pluriSelect Cell Strainers
(43-50020-03, pluriSelect) to remove debris and centrifuged at 1000xg for 5 minutes at 4°C. Pellets were
then resuspended in cold nuclear Fx Buffer (containing 1X HBSS, 2% fetal calf serum, 0.2 U/µL of Protector
RNase Inhibitor). Nuclei were barcoded using cholesterol modified barcodes and anchors as previously
described[49]. Briefly, nuclei were first incubated with the anchor-barcode mix at a final concentration of 200nM
for 5 minutes on ice before being further incubated with a co-anchor at a final concentration of 200 nM for 20
minutes (Barcodes in Supplementary Table 1). Samples were twice washed in cold nuclear Fx buffer before
samples within each group were pooled and stained with DAPI (1 µg/mL; D1306, Invitrogen) in cold nuclear
Fx buffer. Nuclei were filtered through a 35 µM filter (38030, StemCell) before being sorted on the BD
FacsARIA Fusion to remove debris (Alfred Research Alliance FlowCore).

Single-nucleus transcriptomic library preparation and sequencing

FACS isolated nuclei were counted using a hemocytometer and pooled as detailed in Supplementary Table
1 for single nucleus transcriptomic library preparation using the Chromium controller (10X Genomics).
Approximately 40,000 nuclei were loaded into each channel and processed using the Chromium Single Cell
3' v3.1 reagent kit (10X Genomics). After capture and lysis, cDNA was synthesized and amplified for 12
cycles following the manufacturer's protocol. Once amplified, the cDNA was used for downstream sequencing
on four lanes of an Illumina NovaSeq 6000 to an approximate depth of 50,000 reads and 3000 genes per
nucleus and a sequencing saturation of around 50-61.5% (see Supplementary Table 1).

Analysis of single-nucleus RNA sequencing data

Raw sequencing data in FASTQ format was processed using Cell Ranger v6.1.1 (10X Genomics) with the
inclusion of intronic reads before downstream analysis. Reads that were confidently mapped to the genome
were aligned to the *Mus musculus* mm10 transcriptome before the expression of each transcript was
quantified for each nucleus. Analysis was carried out using R v4.1 and v4.2 and Seurat v4.1.1
(satijalab.org/seurat). Nuclei with <200 and >6500 and >7000 unique genes expressed for GEX_A (BAT_dbh)
and GEX_B (BAT_dbdb), respectively, and nuclei having <200 and >7500 unique genes expressed were
excluded for GEX_C (WAT_dbh) and GEX_D (WAT_dbdb), respectively. Nuclei with a percentage of reads
mapping to mitochondrial genes greater than 2, 1, 1, and 0.5 standard deviations were removed from GEX_A,
GEX_B, GEX_C and GEX_D, respectively, to keep the top 80% of nuclei for analysis. SCTransform[79] was
used to normalize the data and regress out ribosomal protein genes (genes starting with Rps and Rpl) and
mitochondrial genes to reduce technical artifacts. Four objects were created for each sequencing run and
nuclei were clustered. Low quality clusters were removed based on nFeature_RNA, nCount_RNA and
percent.mt. Remaining cells were re-clustered, and doublets were removed using DoubletFinder[80] prior to
sample integration. Standard Seurat integration approach was performed with 4000 integration features used
and canonical correlation analysis (CCA) was utilized to find the integration anchors. Sex-based separation
was primarily conducted using the barcoding approach described above. Sex barcodes were demultiplexed
using MULTIseqDeemux() function in Seurat. Nuclei that were predicted as doublets were either further

separated based on our previously published approach[22] that considers the expression of sex-specific genes
(*Xist*, females; *Ddx3y*, *Eif2s3y*, *Gm29650*, *Kdm5d*, *Uty*, males) or discarded from analysis if they showed co-
expression. Sex information then was embedded into the integrated Seurat object. Dimensionality reduction
was conducted using 40 principal components and a resolution of 0.8 to explain variability not expected by
chance to cluster the nuclei into cell populations. Uniform Maximum Approximation and Projection (UMAP)
was used to visualize the data in two dimensions and 'FindAllMarkers' was used to identify cell clusters. As
observed in previous adipose single cell datasets, contaminating epididymal cells, ovarian epithelial cells and
skeletal muscle cells were removed from analysis. The final dataset included 51,877 nuclei across 28
samples (7 samples in each of the 4 groups) comprising 22,526 nuclei in WAT and 29,351 nuclei in BAT from
db/h and db/db samples.

Differential expression analysis

To assess differentially expressed (DE) genes between conditions and sexes, 'MASTcpmDetRate' was used.
MASTcpmDetRate is a modified version of 'MAST'[81] which considers cellular detection rate and has been
successfully used in the past for benchmarking experiments that deemed this approach superior to others[82].
First, ribosomal RNA was removed from the dataset and DE analysis was conducted. Genes with an
uncorrected P value less than 0.01 were considered statistically significant and, thus, differentially expressed
between groups. Data was primarily visualized using the ggplot2 package.

Gene ontology analysis

Gene ontology (GO) enrichment analysis of the above DE genes was performed using the 'enrichGO' function
from the clusterProfiler package (v4.4.4)[83]. DE genes were mapped to genomes from geneontology.org.
Enrichment analysis of GO Biological Process terms (GO terms) was calculated using all genes identified in
the dataset as the background gene set (with minimum and maximum gene set sizes as 10 and 500,
respectively). The 'simplify' function was used to calculate similarities between GO terms. GO terms with a
similarity higher than the 0.7 cut-off were discarded from analysis. Terms with a P value less than 0.05 were
considered statistically significant GO terms. Data was visualized using the ggplot2 package.

Gene expression overlap analysis

To determine the similarity of differentially expressed genes between mouse *db/db* and human obesity
datasets, a publicly available scRNA-seq dataset[21] of visceral WAT and present data on gonadal (visceral)
WAT samples from *db/db* mice were compared. For the human obesity dataset, samples with a BMI greater
than 30 were classed as 'obese' while those less than 25 were classed as 'lean'. DE analysis was conducted
between control and treatment groups (lean vs. obese for humans and *db/h* vs. d*b/db* for mice), as described
above. Human gene orthologs were converted to mouse orthologs using the AnnotationDbi package
(v1.58.0). DE genes with a P value less than 0.01 were isolated from each dataset and the 'overlap' function
within RVenn (v1.1.0) was used to determine both common and unique DE genes across datasets. The
number of common and unique genes were visualized using RVenn (v1.1.0), Euller (v6.1.1), VennDiagram
(v1.7.3) and Vennueler (v1.1.3).

Tissue preparation for histological validation

Age and sex-matched animals were euthanized using CO_2 inhalation cardiac exsanguination. Animals were
then perfused with PBS with 200mM KCl (P9541, Sigma-Aldrich) at a rate of 2 mL/min/mouse until the liver
cleared of blood. Mice were then perfused with 4% PFA (P6148-500G, Sigma) in PBS with 200mM KCl at a
rate of 1 mL/min/mouse. Tissue was kept in 4% PFA in PBS overnight before being stored in PBS with PBS
with 0.09% NaN_3 (S2002-25G, Sigma) at 4°C short-term. Tissue was embedded in paraffin by the Monash
Histology Platform (Monash University, Clayton, VIC, Australia).

Tissue section staining

45 FFPE blocks were cut on a microtome (RM 2235, Leica Biosystems) at 10 µm on SuperFrost plus slides then
46 dried overnight at 37°C. Paraffin was melted at 80°C for 1 minute before slides were washed three times with
47 fresh xylene for 10 minutes each (534056-4L, Honeywell). Slides were rehydrated in 100% (v/v) EtOH, 100%

(v/v) EtOH, 90% (v/v) EtOH, 70% (v/v) EtOH and 50% (v/v) EtOH for 5 minutes each before being submerged in ddH$_2$O for 1 minute. Slides were washed twice in Tris-EDTA, pH 9.0 (10 mM Tris (BIO3094T-1KG, Astral Scientific), 1 mM EDTA (EDS-500G, Sigma) in ddH2O) before being microwaved for 15 minutes in 1 L of Tris-EDTA, pH 9.0. Slides were washed twice with PBS-T (PBS, 0.05% Tween-20) before being permeabilized for 10 mins in 0.5% Triton X-100 (T9284, Sigma) in PBS. Slides were washed twice with PBS for 5 minutes each before blocking with goat block (2% goat serum (G9023-10ML, Sigma), 1% BSA (A3608-100g, Sigma), 0.1% cold fish skin (G7765-250ML, Sigma), 0.1% Triton X-100 (T9284, Sigma) for 1 hour at room temp. Slides were then incubated with primary antibody in goat block (PLIN1, 1:200, AB3526, Abcam; Isolectin B4/IB4, 1:50, L2140, Sigma-Aldrich; Interferon Regulatory Factor 4/IRF4, 1:200, PA5-115603, Invitrogen) overnight at 4°C in a humidified chamber. Sections were washed three times in PBS-T for 5 minutes each. Slides were incubated with secondary antibody in PBS-T for 1 hour at room temp. Sections were then washed three times with PBS-T with AF488 conjugated Wheat Germ Agglutinin (WGA) (29022, Jomar Life Research) for 5 minutes before a final wash with DAPI to stain nuclei in PBS-T for 10 minutes. Sections were mounted under a coverslip before imaging using a 20X multi-immersion objective and glycerol on a Nikon A1R confocal laser scanning microscope.

Statistical analysis of quantified data

For immunofluorescence-based validation, micrographs were analyzed in Fiji image analysis software[84]. Macros were utilized for IB4, nuclei and adipocyte lipid size quantification. Briefly, for tissue sections stained with IB4 and DAPI, the total tissue area and IB4 stained tissue area (determined using thresholding) were measured within a field of view. The percentage of IB4-positive tissue space was calculated. Nuclei (DAPI-positive) were selected with thresholding, quantified, and divided by the total area to give the number of nuclei per µm^2. Finally, thresholding of PLIN1-positive lipid droplets facilitated the subsequent measurement of Feret's diameter (µm) for each PLIN1-positive lipid droplet. Calculations using raw data from Fiji were conducted in Microsoft Excel before processed data were transposed into GraphPad Prism (v9.2.0). GraphPad Prism (v9.2.0) was used to conduct unpaired t-tests on comparisons between two groups and ordinary one-way ANOVAs on comparisons between three or more groups. Statistical significance was determined where $P<0.05$.

Acknowledgments

We would like to acknowledge the Boon Wurrung and Woi Wurrung (Wurundjeri) Peoples of the Kulin Nation on which our research was formulated, conducted, and reported on. We would like to pay our respects to their Elders past, present and emerging.

The authors acknowledge use of the facilities at the AMREP Flow Cytometry Platform, the Monash Histology Platform and the Baker Heart and Diabetes Institute Microscopy Platform.

Author contributions

A.R.P. supervised the study. A.R.P., G.E.F., C.K., T.L.G., and C.D.C. conceived and designed the experiments and collected the data. I.H., M.S.I.D., T.L.G., C.K., C.D.C., and A.R.P. performed the data analysis with I.H. and M.S.I.D. leading the computational analysis. A.R.P. and T.L.G. wrote the manuscript. A.R.P., B.G.D., T.L.G., G.E.F., C.K., C.D.C., I.H., B.G.D and M.S.I.D. reviewed, edited, and approved the manuscript.

Competing interests

The authors declare no competing interests.

Funding

This work was funded by grants from the National Health and Medical Research Council (NHMRC; GNT1188503) and Diabetes Australia (Y20G-PINA) to A.R.P., and Australian Government Defence Science Institute RhD Student Grant to T.L.G.. Further, T.L.G., C.D.C., C.K., and G.E.F., were supported by Australian Government Research Training Program stipends.

Ethics approval

All animal-related experiments were approved by the Alfred Research Alliance (ARA) Animal Ethics Committee (Ethics number: E/1990/2020/B) and were in accordance with NHMRC guidelines.

Data availability

All raw and processed sequencing data, and corresponding scripts, will be made available upon publication.

1. Oikonomou, E.K., and Antoniades, C. (2019). The role of adipose tissue in cardiovascular health and disease. 0–99.

2. Chait, A., and Hartigh, L.J. den (2020). Adipose Tissue Distribution, Inflammation and Its Metabolic
Consequences, Including Diabetes and Cardiovascular Disease. Frontiers in Cardiovascular Medicine *7*, 22–22.
https://doi.org/10.3389/fcvm.2020.00022.

3. Kawai, T., Autieri, M.V., and Scalia, R. (2021). Adipose tissue inflammation and metabolic dysfunction in obesity.
American Journal of Physiology - Cell Physiology *320*, C375–C391.
https://doi.org/10.1152/ajpcell.00379.2020/asset/images/large/aj-acel210014f002.jpeg.

4. Neeland, I.J., Turer, A.T., Ayers, C.R., Berry, J.D., Rohatgi, A., Das, S.R., Khera, A., Vega, G.L., McGuire, D.K.,
Grundy, S.M., et al. (2015). Body Fat Distribution and Incident Cardiovascular Disease in Obese Adults. Journal of
the American College of Cardiology *65*, 2150–2151. https://doi.org/10.1016/j.jacc.2015.01.061.

5. Becher, T., Palanisamy, S., Kramer, D.J., Eljalby, M., Marx, S.J., Wibmer, A.G., Butler, S.D., Jiang, C.S.,
Vaughan, R., Schöder, H., et al. (2021). Brown adipose tissue is associated with cardiometabolic health. Nature
Medicine 2021 27:1 *27*, 58–65. https://doi.org/10.1038/s41591-020-1126-7.

6. Machado, S.A., Pasquarelli-do-Nascimento, G., Silva, D.S.S. da, Farias, G.R., Santos, I. de O., Baptista, L.B., and
Magalhães, K.G. (2022). Browning of the white adipose tissue regulation: new insights into nutritional and metabolic
relevance in health and diseases. Nutrition & Metabolism 2022 19:1 *19*, 1–27. https://doi.org/10.1186/s12986-022-
00694-0.

7. Vijay, J., Gauthier, M.F., Biswell, R.L., Louiselle, D.A., Johnston, J.J., Cheung, W.A., Belden, B., Pramatarova, A.,
Biertho, L., Gibson, M., et al. (2020). Single-cell analysis of human adipose tissue identifies depot- and disease-
specific cell types. Nature Metabolism *2*, 97–109. https://doi.org/10.1038/s42255-019-0152-6.

8. Rajbhandari, P., Arneson, D., Hart, S.K., Ahn, I.S., Diamante, G., Santos, L.C., Zaghari, N., Feng, A.C., Thomas,
B.J., Vergnes, L., et al. (2019). Single cell analysis reveals immune cell-adipocyte crosstalk regulating the
transcription of thermogenic adipocytes. eLife *8*. https://doi.org/10.7554/elife.49501.

9. Bäckdahl, J., Franzén, L., Massier, L., Li, Q., Jalkanen, J., Gao, H., Andersson, A., Bhalla, N., Thorell, A., Rydén,
M., et al. (2021). Spatial mapping reveals human adipocyte subpopulations with distinct sensitivities to insulin. Cell
Metabolism *33*, 1869-1882.e6. https://doi.org/10.1016/j.cmet.2021.07.018.

10. Karlina, R., Lutter, D., Miok, V., Fischer, D., Altun, I., Schöttl, T., Schorpp, K., Israel, A., Cero, C., Johnson,
J.W., et al. (2021). Identification and characterization of distinct brown adipocyte subtypes in C57BL/6J mice. Life
Science Alliance *4*. https://doi.org/10.26508/lsa.202000924.

11. Shamsi, F., Piper, M., Ho, L.-L., Huang, T.L., Gupta, A., Streets, A., Lynes, M.D., and Tseng, Y.-H. (2021).
Vascular smooth muscle-derived Trpv1+ progenitors are a source of cold-induced thermogenic adipocytes. Nature
Metabolism 2021 3:4 *3*, 485–495. https://doi.org/10.1038/s42255-021-00373-z.

12. Sun, W., Dong, H., Balaz, M., Slyper, M., Drokhlyansky, E., Colleluori, G., Giordano, A., Kovanicova, Z.,
Stefanicka, P., Balazova, L., et al. (2020). snRNA-seq reveals a subpopulation of adipocytes that regulates
thermogenesis. Nature 2020 587:7832 *587*, 98–102. https://doi.org/10.1038/s41586-020-2856-x.

13. Whytock, K.L., Sun, Y., Divoux, A., Yu, G., Smith, S.R., Walsh, M.J., and Sparks, L.M. (2022). Single cell full-
length transcriptome of human subcutaneous adipose tissue reveals unique and heterogeneous cell populations.
iScience *25*, 104772–104772. https://doi.org/10.1016/j.isci.2022.104772.

14. Hill, D.A., Lim, H.W., Kim, Y.H., Ho, W.Y., Foong, Y.H., Nelson, V.L., Nguyen, H.C.B., Chegireddy, K., Kim,
J., Habertheuer, A., et al. (2018). Distinct macrophage populations direct inflammatory versus physiological changes

in adipose tissue. Proceedings of the National Academy of Sciences of the United States of America *115*, E5096–E5105. https://doi.org/10.1073/pnas.1802611115.

15. Cho, D.S., Lee, B., and Doles, J.D. (2019). Refining the adipose progenitor cell landscape in healthy and obese visceral adipose tissue using single-cell gene expression profiling. Life Science Alliance *2*, e201900561–e201900561. https://doi.org/10.26508/lsa.201900561.

16. Jaitin, D.A., Adlung, L., Thaiss, C.A., Weiner, A., Li, B., Descamps, H., Lundgren, P., Bleriot, C., Liu, Z., Deczkowska, A., et al. (2019). Lipid-Associated Macrophages Control Metabolic Homeostasis in a Trem2-Dependent Manner. Cell *178*, 686-698.e14. https://doi.org/10.1016/j.cell.2019.05.054.

17. Harasymowicz, N.S., Rashidi, N., Savadipour, A., Wu, C.-L., Tang, R., Bramley, J., Buchser, W., and Guilak, F. (2021). Single-cell RNA sequencing reveals the induction of novel myeloid and myeloid-associated cell populations in visceral fat with long-term obesity. The FASEB Journal *35*, e21417–e21417. https://doi.org/10.1096/fj.202001970r.

18. Hildreth, A.D., Ma, F., Wong, Y.Y., Sun, R., Pellegrini, M., and O'Sullivan, T.E. (2021). Single-cell sequencing of human white adipose tissue identifies new cell states in health and obesity. Nature Immunology, 1–15. https://doi.org/10.1038/s41590-021-00922-4.

19. Sárvári, A.K., Hauwaert, E.L.V., Markussen, L.K., Gammelmark, E., Marcher, A.B., Ebbesen, M.F., Nielsen, R., Brewer, J.R., Madsen, J.G.S., and Mandrup, S. (2021). Plasticity of Epididymal Adipose Tissue in Response to Diet-Induced Obesity at Single-Nucleus Resolution. Cell Metabolism *33*, 437-453.e5. https://doi.org/10.1016/j.cmet.2020.12.004.

20. Cottam, M.A., Caslin, H.L., Winn, N.C., and Hasty, A.H. (2022). Multiomics reveals persistence of obesity-associated immune cell phenotypes in adipose tissue during weight loss and weight regain in mice. Nature Communications 2022 13:1 *13*, 1–16. https://doi.org/10.1038/s41467-022-30646-4.

21. Emont, M.P., Jacobs, C., Essene, A.L., Pant, D., Tenen, D., Colleluori, G., Vincenzo, A.D., Jørgensen, A.M., Dashti, H., Stefek, A., et al. (2022). A single-cell atlas of human and mouse white adipose tissue. Nature *603*, 926–933. https://doi.org/10.1038/s41586-022-04518-2.

22. McLellan, M.A., Skelly, D.A., Dona, M.S.I., Squiers, G.T., Farrugia, G.E., Gaynor, T.L., Cohen, C.D., Pandey, R., Diep, H., Vinh, A., et al. (2020). High-Resolution Transcriptomic Profiling of the Heart During Chronic Stress Reveals Cellular Drivers of Cardiac Fibrosis and Hypertrophy. Circulation *142*, 1448–1463. https://doi.org/10.1161/circulationaha.119.045115.

23. Larsen, T.J., Jespersen, N.Z., and Scheele, C. (2018). Brown Adipose Tissue. Handb. Exp. Pharmacol. *251*, 73–84. https://doi.org/10.1007/164_2018_142.

24. Karnam, P., Standaert, M.L., Galloway, L., and Farese, R.V. (1997). Activation and Translocation of Rho (and ADP Ribosylation Factor) by Insulin in Rat Adipocytes. Journal of Biological Chemistry *272*, 6136–6140. https://doi.org/10.1074/jbc.272.10.6136.

25. Vergnes, L., Chin, R., Young, S.G., and Reue, K. (2011). Heart-type Fatty Acid-binding Protein Is Essential for Efficient Brown Adipose Tissue Fatty Acid Oxidation and Cold Tolerance. The Journal of Biological Chemistry *286*, 380–380. https://doi.org/10.1074/jbc.m110.184754.

26. Cui, X., Jing, J., Wu, R., Cao, Q., Li, F., Li, K., Wang, S., Yu, L., Schwartz, G., Shi, H., et al. (2021). Adipose tissue-derived neurotrophic factor 3 regulates sympathetic innervation and thermogenesis in adipose tissue. Nature Communications 2021 12:1 *12*, 1–18. https://doi.org/10.1038/s41467-021-25766-2.

27. Subramani, M., and Yun, J.W. (2021). Loss of lymphocyte cytosolic protein 1 (LCP1) induces browning in 3T3-L1 adipocytes via β3-AR and the ERK-independent signaling pathway. The International Journal of Biochemistry & Cell Biology *138*, 106053–106053. https://doi.org/10.1016/j.biocel.2021.106053.

28. Leonardsson, G., Steel, J.H., Christian, M., Pocock, V., Milligan, S., Bell, J., So, P.W., Medina-Gomez, G., Vidal-Puig, A., White, R., et al. (2004). Nuclear receptor corepressor RIP140 regulates fat accumulation. Proceedings of the National Academy of Sciences of the United States of America *101*, 8437–8442. https://doi.org/10.1073/pnas.0401013101/suppl_file/01013fig7.pdf.

29. Kiefer, F.W., Vernochet, C., O'Brien, P., Spoerl, S., Brown, J.D., Nallamshetty, S., Zeyda, M., Stulnig, T.M., Cohen, D.E., Kahn, C.R., et al. (2012). Retinaldehyde dehydrogenase 1 regulates a thermogenic program in white adipose tissue. Nature medicine *18*, 918–918. https://doi.org/10.1038/nm.2757.

30. Turpin, S.M., Nicholls, H.T., Willmes, D.M., Mourier, A., Brodesser, S., Wunderlich, C.M., Mauer, J., Xu, E., Hammerschmidt, P., Brönneke, H.S., et al. (2014). Obesity-induced CerS6-dependent C16:0 ceramide production promotes weight gain and glucose intolerance. Cell Metabolism *20*, 678–686. https://doi.org/10.1016/j.cmet.2014.08.002.

31. Choi, K.M., Ko, C.Y., An, S.M., Cho, S.H., Rowland, D.J., Kim, J.H., Fasoli, A., Chaudhari, A.J., Bers, D.M., and Yoon, J.C. (2023). Regulation of beige adipocyte thermogenesis by the cold-repressed ER protein NNAT. Molecular Metabolism *69*, 101679–101679. https://doi.org/10.1016/j.molmet.2023.101679.

32. Ross, S.E., Hemati, N., Longo, K.A., Bennett, C.N., Lucas, P.C., Erickson, R.L., and MacDougald, O.A. (2000). Inhibition of Adipogenesis by Wnt Signaling. Science *289*, 950–950.

33. Bennett, C.N., Ross, S.E., Longo, K.A., Bajnok, L., Hemati, N., Johnson, K.W., Harrison, S.D., and MacDougald, O.A. (2002). Regulation of Wnt signaling during adipogenesis. Journal of Biological Chemistry *277*, 30998–31004. https://doi.org/10.1074/jbc.m204527200.

34. Zamani, N., and Brown, C.W. (2011). Emerging Roles for the Transforming Growth Factor-β Superfamily in Regulating Adiposity and Energy Expenditure. Endocr. Rev. *32*, 387–403. https://doi.org/10.1210/er.2010-0018.

35. Villanueva, C.J., Vergnes, L., Wang, J., Drew, B.G., Hong, C., Tu, Y., Hu, Y., Peng, X., Xu, F., Saez, E., et al. (2013). Adipose Subtype-Selective Recruitment of TLE3 or Prdm16 by PPARγ Specifies Lipid Storage versus Thermogenic Gene Programs. Cell Metab. *17*, 423–435. https://doi.org/10.1016/j.cmet.2013.01.016.

36. Aibar, S., González-Blas, C.B., Moerman, T., Huynh-Thu, V.A., Imrichova, H., Hulselmans, G., Rambow, F., Marine, J.C., Geurts, P., Aerts, J., et al. (2017). SCENIC: Single-cell regulatory network inference and clustering. Nature methods *14*, 1083–1083. https://doi.org/10.1038/nmeth.4463.

37. Randi, A.M., Sperone, A., Dryden, N.H., and Birdsey, G.M. (2009). Regulation of angiogenesis by ETS transcription factors. Biochemical Society Transactions *37*, 1248–1253. https://doi.org/10.1042/bst0371248.

38. Deleuze, V., El-Hajj, R., Chalhoub, E., Dohet, C., Pinet, V., Couttet, P., and Mathieu, D. (2012). Angiopoietin-2 is a direct transcriptional target of TAL1, LYL1 and LMO2 in endothelial cells. PloS one *7*. https://doi.org/10.1371/journal.pone.0040484.

39. Rehli, M., Sulzbacher, S., Pape, S., Ravasi, T., Wells, C.A., Heinz, S., Söllner, L., Chartouni, C.E., Krause, S.W., Steingrimsson, E., et al. (2005). Transcription factor Tfec contributes to the IL-4-inducible expression of a small group of genes in mouse macrophages including the granulocyte colony-stimulating factor receptor. Journal of immunology (Baltimore, Md. : 1950) *174*, 7111–7122. https://doi.org/10.4049/jimmunol.174.11.7111.

40. Krausgruber, T., Blazek, K., Smallie, T., Alzabin, S., Lockstone, H., Sahgal, N., Hussell, T., Feldmann, M., and Udalova, I.A. (2011). IRF5 promotes inflammatory macrophage polarization and TH1-TH17 responses. Nature Immunology 2011 12:3 *12*, 231–238. https://doi.org/10.1038/ni.1990.

41. Alexanian, M., Przytycki, P.F., Micheletti, R., Padmanabhan, A., Ye, L., Travers, J.G., Gonzalez-Teran, B., Silva, A.C., Duan, Q., Ranade, S.S., et al. (2021). A transcriptional switch governs fibroblast activation in heart disease. Nature 2021 595:7867 *595*, 438–443. https://doi.org/10.1038/s41586-021-03674-1.

42. Khalil, H., Kanisicak, O., Prasad, V., Correll, R.N., Fu, X., Schips, T., Vagnozzi, R.J., Liu, R., Huynh, T., Lee, S.J., et al. (2017). Fibroblast-specific TGF-β-Smad2/3 signaling underlies cardiac fibrosis. The Journal of clinical investigation *127*, 3770–3783. https://doi.org/10.1172/jci94753.

43. Cinti, S., Mitchell, G., Barbatelli, G., Murano, I., Ceresi, E., Faloia, E., Wang, S., Fortier, M., Greenberg, A.S., Obin, M.S., et al. (2005). Adipocyte death defines macrophage localization and function in adipose tissue of obese mice and humans. Journal of Lipid Research *46*, 2347–2355. https://doi.org/10.1194/jlr.m500294-jlr200.

44. Amano, S.U., Cohen, J.L., Vangala, P., Tencerova, M., Nicoloro, S.M., Yawe, J.C., Shen, Y., Czech, M.P., and Aouadi, M. (2014). Local proliferation of macrophages contributes to obesity-associated adipose tissue inflammation. Cell metabolism *19*, 162–162. https://doi.org/10.1016/j.cmet.2013.11.017.

45. Zheng, C., Yang, Q., Cao, J., Xie, N., Liu, K., Shou, P., Qian, F., Wang, Y., and Shi, Y. (2016). Local proliferation initiates macrophage accumulation in adipose tissue during obesity. Cell Death & Disease 2016 7:3 *7*, e2167–e2167. https://doi.org/10.1038/cddis.2016.54.

46. Eguchi, J., Kong, X., Tenta, M., Wang, X., Kang, S., and Rosen, E.D. (2013). Interferon Regulatory Factor 4 Regulates Obesity-Induced Inflammation Through Regulation of Adipose Tissue Macrophage Polarization. Diabetes *62*, 3394–3403. https://doi.org/10.2337/db12-1327.

47. Sardiello, M., Palmieri, M., Ronza, A.D., Medina, D.L., Valenza, M., Gennarino, V.A., Malta, C.D., Donaudy, F., Embrione, V., Polishchuk, R.S., et al. (2009). A gene network regulating lysosomal biogenesis and function. Science *325*, 473–477. https://doi.org/10.1126/science.1174447/suppl_file/sardiello-som.pdf.

48. He, Y.S., Yang, X.K., Hu, Y.Q., Xiang, K., and Pan, H.F. (2021). Emerging role of Fli1 in autoimmune diseases. International Immunopharmacology *90*, 107127–107127. https://doi.org/10.1016/j.intimp.2020.107127.

49. McGinnis, C.S., Patterson, D.M., Winkler, J., Conrad, D.N., Hein, M.Y., Srivastava, V., Hu, J.L., Murrow, L.M., Weissman, J.S., Werb, Z., et al. (2019). MULTI-seq: sample multiplexing for single-cell RNA sequencing using lipid-tagged indices. Nature Methods 2019 16:7 *16*, 619–626. https://doi.org/10.1038/s41592-019-0433-8.

50. Yoshida, T., Nishioka, H., Yoshioka, K., and Kondo, M. (1987). Reduced norepinephrine turnover in interscapular brown adipose tissue of obese rats after ovariectomy. Metabolism *36*, 1–6. https://doi.org/10.1016/0026-0495(87)90054-0.

51. Yoshioka, K., Yoshida, T., Wakabayashi, Y., Nishioka, H., and Kondo, M. (1988). Reduced Brown Adipose Tissue Thermogenesis of Obese Rats After Ovariectomy. Endocrinologia Japonica *35*, 537–543. https://doi.org/10.1507/endocrj1954.35.537.

52. Quevedo, S., Roca, P., Picó, C., and Palou, A. (1998). Sex-associated differences in cold-induced UCP1 synthesis in rodent brown adipose tissue. Pflügers Archiv 1998 436:5 *436*, 689–695. https://doi.org/10.1007/s004240050690.

53. Justo, R., Frontera, M., Pujol, E., Rodríguez-Cuenca, S., Lladó, I., García-Palmer, F.J., Roca, P., and Gianotti, M. (2005). Gender-related differences in morphology and thermogenic capacity of brown adipose tissue mitochondrial subpopulations. Life Sciences *76*, 1147–1158. https://doi.org/10.1016/j.lfs.2004.08.019.

54. Kim, S.N., Jung, Y.S., Kwon, H.J., Seong, J.K., Granneman, J.G., and Lee, Y.H. (2016). Sex differences in sympathetic innervation and browning of white adipose tissue of mice. Biology of Sex Differences *7*, 1–13. https://doi.org/10.1186/s13293-016-0121-7.

55. Rudnicki, M., Abdifarkosh, G., Rezvan, O., Nwadozi, E., Roudier, E., and Haas, T.L. (2018). Female mice have higher angiogenesis in perigonadal adipose tissue than males in response to high-fat diet. Frontiers in Physiology *9*, 1452–1452. https://doi.org/10.3389/fphys.2018.01452.

56. Bagchi, D.P., Nishii, A., Li, Z., DelProposto, J.B., Corsa, C.A., Mori, H., Hardij, J., Learman, B.S., Lumeng, C.N., and MacDougald, O.A. (2020). Wnt/β-catenin signaling regulates adipose tissue lipogenesis and adipocyte-specific

loss is rigorously defended by neighboring stromal-vascular cells. Molecular metabolism *42*. https://doi.org/10.1016/j.molmet.2020.101078.

57. Palani, N.P., Horvath, C., Timshel, P.N., Folkertsma, P., Grønning, A.G.B., Henriksen, T.I., Peijs, L., Jensen, V.H., Sun, W., Jespersen, N.Z., et al. (2023). Adipogenic and SWAT cells separate from a common progenitor in human brown and white adipose depots. Nature Metabolism 2023 5:6 *5*, 996–1013. https://doi.org/10.1038/s42255-023-00820-z.

58. Haase, J., Weyer, U., Immig, K., Klöting, N., Blüher, M., Eilers, J., Bechmann, I., and Gericke, M. (2014). Local proliferation of macrophages in adipose tissue during obesity-induced inflammation. Diabetologia *57*, 562–571. https://doi.org/10.1007/s00125-013-3139-y/figures/7.

59. Zamarron, B.F., Mergian, T.A., Cho, K.W., Martinez-Santibanez, G., Luan, D., Singer, K., DelProposto, J.L., Geletka, L.M., Muir, L.A., and Lumeng, C.N. (2017). Macrophage Proliferation Sustains Adipose Tissue Inflammation in Formerly Obese Mice. Diabetes *66*, 392–406. https://doi.org/10.2337/db16-0500.

60. Palmieri, M., Impey, S., Kang, H., Ronza, A. di, Pelz, C., Sardiello, M., and Ballabio, A. (2011). Characterization of the CLEAR network reveals an integrated control of cellular clearance pathways. Human Molecular Genetics *20*, 3852–3866. https://doi.org/10.1093/hmg/ddr306.

61. Mizunoe, Y., Kobayashi, M., Hoshino, S., Tagawa, R., Itagawa, R., Hoshino, A., Okita, N., Sudo, Y., Nakagawa, Y., Shimano, H., et al. (2020). Cathepsin B overexpression induces degradation of perilipin 1 to cause lipid metabolism dysfunction in adipocytes. Scientific Reports 2020 10:1 *10*, 1–12. https://doi.org/10.1038/s41598-020-57428-6.

62. Mizunoe, Y., Sudo, Y., Okita, N., Hiraoka, H., Mikami, K., Narahara, T., Negishi, A., Yoshida, M., Higashibata, R., Watanabe, S., et al. (2017). Involvement of lysosomal dysfunction in autophagosome accumulation and early pathologies in adipose tissue of obese mice. Autophagy *13*, 642–653. https://doi.org/10.1080/15548627.2016.1274850.

63. Haka, A.S., Barbosa-Lorenzi, V.C., Lee, H.J., Falcone, D.J., Hudis, C.A., Dannenberg, A.J., and Maxfield, F.R. (2016). Exocytosis of macrophage lysosomes leads to digestion of apoptotic adipocytes and foam cell formation[S]. J. Lipid Res. *57*, 980–992. https://doi.org/10.1194/jlr.m064089.

64. Rodríguez, E., Monjo, M., Rodríguez-Cuenca, S., Pujol, E., Amengual, B., Roca, P., and Palou, A. (2014). Sexual dimorphism in the adrenergic control of rat brown adipose tissue response to overfeeding. Pflügers Archiv 2001 442:3 *442*, 396–403. https://doi.org/10.1007/s004240100556.

65. Rodríguez-Cuenca, S., Pujol, E., Justo, R., Frontera, M., Oliver, J., Gianotti, M., and Roca, P. (2002). Sex-dependent thermogenesis, differences in mitochondrial morphology and function, and adrenergic response in brown adipose tissue. Journal of Biological Chemistry *277*, 42958–42963. https://doi.org/10.1074/jbc.m207229200.

66. Rodríguez, A.M., Quevedo-Coli, S., Roca, P., and Palou, A. (2001). Sex-dependent dietary obesity, induction of UCPs, and leptin expression in rat adipose tissues. Obesity Research *9*, 579–588. https://doi.org/10.1038/oby.2001.75.

67. Cinti, S., Cancello, R., Zingaretti, M.C., Ceresi, E., Matteis, R.D., Giordano, A., Himms-Hagen, J., and Ricquier, D. (2002). CL316,243 and cold stress induce heterogeneous expression of UCP1 mRNA and protein in rodent brown adipocytes. Journal of Histochemistry and Cytochemistry *50*, 21–31. https://doi.org/10.1177/002215540205000103/asset/images/large/10.1177_002215540205000103-fig2.jpeg.

68. Song, A., Dai, W., Jang, M.J., Medrano, L., Li, Z., Zhao, H., Shao, M., Tan, J., Li, A., Ning, T., et al. (2020). Low- And high-thermogenic brown adipocyte subpopulations coexist in murine adipose tissue. Journal of Clinical Investigation *130*, 247–257. https://doi.org/10.1172/jci129167.

69. Shimizu, I., Aprahamian, T., Kikuchi, R., Shimizu, A., Papanicolaou, K.N., MacLauchlan, S., Maruyama, S., and Walsh, K. (2014). Vascular rarefaction mediates whitening of brown fat in obesity. The Journal of Clinical Investigation *124*, 2099–2099. https://doi.org/10.1172/jci71643.

70. Fatima, L.A., Campello, R.S., Santos, R.D.S., Freitas, H.S., Frank, A.P., Machado, U.F., and Clegg, D.J. (2017). Estrogen receptor 1 (ESR1) regulates VEGFA in adipose tissue. Scientific Reports 2017 7:1 *7*, 1–14. https://doi.org/10.1038/s41598-017-16686-7.

71. Suriano, F., Vieira-Silva, S., Falony, G., Roumain, M., Paquot, A., Pelicaen, R., Régnier, M., Delzenne, N.M., Raes, J., Muccioli, G.G., et al. (2021). Novel insights into the genetically obese (ob/ob) and diabetic (db/db) mice: two sides of the same coin. Microbiome *9*, 1–20. https://doi.org/10.1186/s40168-021-01097-8/figures/7.

72. Jong, J.M.A. de, Sun, W., Pires, N.D., Frontini, A., Balaz, M., Jespersen, N.Z., Feizi, A., Petrovic, K., Fischer, A.W., Bokhari, M.H., et al. (2019). Human brown adipose tissue is phenocopied by classical brown adipose tissue in physiologically humanized mice. Nature Metabolism *1*, 830–843. https://doi.org/10.1038/s42255-019-0101-4.

73. Huang, Z., Zhang, Z., Moazzami, Z., Zhu, W., Shen, S., and Ruan, H.-B. (2022). Brown adipose tissue involution associated with progressive restriction in progenitor competence. CellReports *39*, 110575–110575. https://doi.org/10.1016/j.celrep.2022.110575.

74. Coleman, D.L. (1982). Diabetes-Obesity Syndromes in Mice. Diabetes *31*, 1–6. https://doi.org/10.2337/diab.31.1.s1.

75. Coleman, D.L. (1978). Obese and diabetes: Two mutant genes causing diabetes-obesity syndromes in mice. Diabetologia *14*, 141–148. https://doi.org/10.1007/bf00429772.

76. Chen, H., Charlat, O., Tartaglia, L.A., Woolf, E.A., Weng, X., Ellis, S.J., Lakey, N.D., Culpepper, J., More, K.J., Breitbart, R.E., et al. (1996). Evidence That the Diabetes Gene Encodes the Leptin Receptor: Identification of a Mutation in the Leptin Receptor Gene in db/db Mice. Cell *84*, 491–495. https://doi.org/10.1016/s0092-8674(00)81294-5.

77. Pinto, A.R., Chandran, A., Rosenthal, N.A., and Godwin, J.W. (2013). Isolation and analysis of single cells from the mouse heart. Journal of Immunological Methods *393*, 74–80. https://doi.org/10.1016/j.jim.2013.03.012.

78. Zhu, C., Yu, M., Huang, H., Juric, I., Abnousi, A., Hu, R., Lucero, J., Behrens, M.M., Hu, M., and Ren, B. (2019). An ultra high-throughput method for single-cell joint analysis of open chromatin and transcriptome. Nat. Struct. Mol. Biol. *26*, 1063–1070. https://doi.org/10.1038/s41594-019-0323-x.

79. Hafemeister, C., and Satija, R. (2019). Normalization and variance stabilization of single-cell RNA-seq data using regularized negative binomial regression. Genome Biology *20*, 1–15. https://doi.org/10.1186/s13059-019-1874-1/figures/6.

80. McGinnis, C.S., Murrow, L.M., and Gartner, Z.J. (2019). DoubletFinder: Doublet Detection in Single-Cell RNA Sequencing Data Using Artificial Nearest Neighbors. Cell Systems *8*, 329-337.e4. https://doi.org/10.1016/j.cels.2019.03.003.

81. Finak, G., McDavid, A., Yajima, M., Deng, J., Gersuk, V., Shalek, A.K., Slichter, C.K., Miller, H.W., McElrath, M.J., Prlic, M., et al. (2015). MAST: A flexible statistical framework for assessing transcriptional changes and characterizing heterogeneity in single-cell RNA sequencing data. Genome Biology *16*, 1–13. https://doi.org/10.1186/s13059-015-0844-5/figures/6.

82. Soneson, C., and Robinson, M.D. (2018). Bias, robustness and scalability in single-cell differential expression analysis. Nature Methods 2018 15:4 *15*, 255–261. https://doi.org/10.1038/nmeth.4612.

83. Yu, G., Wang, L.G., Han, Y., and He, Q.Y. (2012). ClusterProfiler: An R package for comparing biological themes among gene clusters. OMICS A Journal of Integrative Biology *16*, 284–287. https://doi.org/10.1089/omi.2011.0118/asset/images/large/figure1.jpeg.

84. Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T., Preibisch, S., Rueden, C., Saalfeld, S., Schmid, B., et al. (2012). Fiji: an open-source platform for biological-image analysis. Nature Methods 2012 9:7 *9*, 676–682. https://doi.org/10.1038/nmeth.2019.

Figures legends

Figure 1. snRNA-seq characterization of adipose cellulome. (A) Experimental outline for novel isolation
of mature adipocyte and SVF nuclei from snap-frozen adipose tissue. **(B)** UMAP of nuclei representing the
adipose cellulome for both *db/h* and *db/db* female and male white and brown adipose tissues. **(C)**
Compositional bar chart showing proportion of each cell population in each tissue type. **(D)** Canonical
markers of broad adipose tissue cell populations represented in heat map with relative expression indicated
by red intensity. **(E)** Top 5 marker genes for each broad cell population with dot size indicating the proportion
of cells within that population expressing the marker and dot color indicating the average level of expression.
WAT, white adipose tissue; BAT, brown adipose tissue; Fibro, fibroblast; Multipot. prog., multipotent
progenitors; Meso. cells, mesothelial cells; ECs, endothelial cells; LECs, lymphatic endothelial cells; SMCs,
smooth muscle cells; Mac, macrophage; Prolif. macs, proliferating macrophages; BA, brown adipocyte; WA,
white adipocyte.

Figure 2. Cellularity and gene expression differences between disparate adipose depots. (A) Flow
cytometry data of differences in adipose cell population contribution to total viable cell count across BAT
(pink) and WAT (black). Statistical significance was assumed at P<0.05. * P< 0.05, ** P< 0.01, *** P< 0.001,
**** P< 0.0001. **(B)** Immunofluorescence staining of vasculature (endothelial cells) and nuclei using isolectin
B4 (IB4; grey) and DAPI (blue). Scale bar represents 50µm. Statistical significance was assumed at P<0.05.
**** P< 0.0001. **(C)** Lollipop plot of upregulated genes across BAT or WAT and Sankey plot of top 3 linked GO
terms for each cell population. **(D)** Volcano plots showing differentially expressed genes between white and
brown adipocyte subsets in BAT and WAT. Gene placement to the right and left indicate a positive and
negative fold difference in expression between tissues, respectively. The placement of the gene on the y-
axis indicates significance (P value). GO term bar charts indicating key terms linked to the adjacent genes
with the size of the bar representing the number of genes associated.

**Figure 3. Distinct gene regulatory networks drive cellular phenotypes within brown and white adipose
tissue depots. (A)** UMAP clustering based on regulon-specific expression patterns. Annotation of cell
populations carried over from RNA-based clustering seen in Figure 1B. **(B)** Top 5 marker regulons for each
cell population based on AUCBinary assay. Intensity of black indicates average regulon activity for each cell
population.

Figure 4. Adipose depot remodeling induced by type-2 diabetes. (A) UMAP of both *db/h* nuclei (yellow)
and *db/db* nuclei (blue). **(B)** Adipocyte lipid droplet size measured using Feret's diameter in both *db/h* and
db/db WAT and BAT. Lipid droplet stained with perilipin 1 (PLIN1; grey) and nuclei stained with DAPI (blue).
Corresponding quantification presented. Statistical significance was assumed at P<0.05. ****, *P*<0.0001. **(C)**
Sankey plots for WAT demonstrating upregulated (UR; right) and downregulated (DR; left) GO terms and
overlayed lollipop plot for associated differentially expressed genes in T2D. **(D)** Sankey plots and lollipop plot
for BAT as described previously. **(E)** Immunofluorescence staining of proliferation in crown-like structures in
db/db WAT through proliferative marker phosphohistone H3 in addition to wheat germ agglutinin (WGA) to
mark cell membranes and DAPI to mark nuclei. **(F)** Feature plots showing expression of proliferation markers
Top2a, *Mki67*, and *Aurka* in macrophage subsets. **(G)** Overlay of differentially expressed genes in all cell
populations and macrophages for visceral WAT between the present study and a human obesity study
(Emont et al. 2022).

**Figure 5. Gene regulatory network alterations in T2D adipose depots and contribution to leukocyte
migration. (A)** RSS plots demonstrating the top 5 activated regulons in *db/db* (red dots) as opposed to *db/h*
samples (blue dots) for white adipocytes, brown adipocytes, fibroblasts and macrophages for WAT (top
panel) and BAT (lower panel). **(B)** Immunofluorescence validation of transcription factor IRF4 (grey) in both
db/h and *db/db* BAT tissue with nuclei stained with DAPI (blue). Scale bar represents 25µm. **(C)** Differential
enrichment of regulons with upregulated regulons in orange and downregulated regulons in blue. Intensity of
color indicates the average expression in that cell population and dot size represents statistical significance.
Regulons with bolded text represent those shared across both depots. Bar charts adjacent to regulons
indicate the number of target genes within that regulon that are also linked to the GO term *leukocyte*
migration.

Figure 6. Sex differences in gene expression and tissue architecture across adipose depots. (A)
Lollipop plot demonstrating sex differences in gene expression between females and males in *db/h* WAT and
BAT. **(B)** Compositional bar chart demonstrating the contribution of adipose tissue cell populations to each
depot and how they differ between conditions and biological sexes. **(C)** Ridgeplots showing female and male
sample expression of key thermogenic genes between *db/h* and *db/db* conditions in BAT. **(D)** Top 5 gene
ontology terms for each cell population where dot size represents the number of genes linked to this GO term
and dot color represents the statistical significance. GO terms of interest are highlighted in bold. **(E)**
Immunofluorescence staining of BAT vascularization using isolectin B4 (IB4; red) and nuclei stained with
DAPI (grey) demonstrating differences in tissue architecture between females and males in both *db/h* and
db/db conditions. Scale bar represents 100μm. **(F)** Quantification of vascularization (% of IB4+ tissue area)
and cell density (number of DAPI+ nuclei) histology with female data points presented in pink and males in
blue. Statistical significance was assumed at P<0.05. * *P*<0.05, ** *P*<0.01, *** *P*<0.001.

Supplementary Figure 1. Optimization of nucleus isolation and flow cytometry gating strategy. (A)
Flow cytometry-based optimization of the time length required to mechanically digested at 30hz using the
Qiagen Tissue Lyser to get optimal nuclei/mg of tissue yield. **(B)** micrographs showing nuclei post Qiagen
digestion and fluorescence-activated cell sorting to remove cellular debris. Scale bar represents 20μm. **(C)**
Flow cytometry gating strategy for SVF cell population analysis. Viable and metabolically active single cells
were gated, and cell populations were delineated based on staining by antibodies specific for cell surface
markers. WAT, white adipose tissue; BAT, brown adipose tissue; FSC-H, forward scatter height; FSW-A,
forward scatter area; FSC, forward scatter width; CB, calcein blue; SG, sytox green; ECs, endothelial cells;
RMCs, resident mesenchymal cells; VECs, vascular endothelial cells; LECs, lymphatic endothelial cells.

Supplementary Figure 2. Cell population abundance changes in T2D. (A) Cell proportion differences
between *db/h* and *db/db* samples based on nuclei count in WAT and BAT. Y axis indicates frequency of total
cell count. **(B)** Contribution of adipose tissue cell populations to flow cytometry total viable cell count with
differences between *db/h* (light color) and *db/db* samples (dark color) being shown in WAT (blue) and BAT
(orange) depots. BA, brown adipocyte; WA, white adipocyte; ECs, endothelial cells; Meso. cells, mesothelial
cells; Multipot. prog, multipotent progenitors; Prolif. macs, proliferating macrophages; SMCs, smooth muscle
cells; WA, white adipocyte; LECs, lymphatic endothelial cells; VECs, vascular endothelial cells; Myel. cells,
myeloid cells; Macs, macrophages; Grans, granulocytes; Non-myel. cells, on-myeloid cells; Fibro, fibroblasts.
Statistical significance was assumed at P<0.05. * *P*<0.05, ** *P*<0.01, *** *P*<0.001, **** *P*<0.0001.

Supplementary Figure 3. Top 50 upregulated and downregulated genes in T2D WAT and BAT. Genes
up-(red) downregulated (blue) between *db/h* and *db/db* cell populations in WAT **(A)** and BAT **(B)**. Dot size
represents the fold change (FC) difference between conditions while color intensity indicates the expression
level (avg. exp. scale). ECs, endothelial cells; LECs, lymphatic endothelial cells; Meso. cells, mesothelial
cells; Multipot. prog, multipotent progenitors; SMCs, smooth muscle cells.

Figure 1. snRNA-seq characterization of adipose cellulome. (A) Experimental outline for novel isolation of mature adipocyte and SVF nuclei from snap-frozen adipose tissue. (B) UMAP of nuclei representing the adipose cellulome for both db/h and db/db female and male white and brown adipose tissues. (C) Compositional bar chart showing proportion of each cell population in each tissue type. (D) Canonical markers of broad adipose tissue cell populations represented in heat map with relative expression indicated by red intensity. (E) Top 5 marker genes for each broad cell population with dot size indicating the proportion of cells within that population expressing the marker and dot color indicating the average level of expression. WAT, white adipose tissue; BAT, brown adipose tissue; Fibro, fibroblast; Multipot. prog., multipotent progenitors; Meso. cells, mesothelial cells; ECs, endothelial cells; LECs, lymphatic endothelial cells; SMCs, smooth muscle cells; Mac, macrophage; Prolif. macs, proliferating macrophages; BA, brown adipocyte; WA, white adipocyte.

Figure 2. Cellularity and gene expression differences between disparate adipose depots. (A) Flow cytometry data of differences in adipose cell population contribution to total viable cell count across BAT (pink) and WAT (black). Statistical significance was assumed at P<0.05. * P< 0.05, ** P< 0.01, *** P< 0.001, **** P< 0.0001. (B) Immunofluorescence staining of vasculature (endothelial cells) and nuclei using isolectin B4 (IB4; grey) and DAPI (blue). Scale bar represents 50μm. Statistical significance was assumed at P<0.05. **** P< 0.0001. (C) Lollipop plot of upregulated genes across BAT or WAT and Sankey plot of top 3 linked GO terms for each cell population. (D) Volcano plots showing differentially expressed genes between white and brown adipocyte subsets in BAT and WAT. Gene placement to the right and left indicate a positive and negative fold difference in expression between tissues, respectively. The placement of the gene on the y-axis indicates significance (P value). GO term bar charts indicating key terms linked to the adjacent genes with the size of the bar representing the number of genes associated.

Figure 3. Distinct gene regulatory networks drive cellular phenotypes within brown and white adipose tissue depots. (A) UMAP clustering based on regulon-specific expression patterns. Annotation of cell populations carried over from RNA-based clustering seen in Figure 1B. (B) Top 5 marker regulons for each cell population based on AUCBinary assay. Intensity of black indicates average regulon activity for each cell population.

Figure 4. Adipose depot remodeling induced by type-2 diabetes. (A) UMAP of both db/h nuclei (yellow) and db/db nuclei (blue). (B) Adipocyte lipid droplet size measured using Feret's diameter in both db/h and db/db WAT and BAT. Lipid droplet stained with perilipin 1 (PLIN1; grey) and nuclei stained with DAPI (blue). Corresponding quantification presented. Statistical significance was assumed at P<0.05. ****, P<0.0001. (C) Sankey plots for WAT demonstrating upregulated (UR; right) and downregulated (DR; left) GO terms and overlayed lollipop plot for associated differentially expressed genes in T2D. (D) Sankey plots and lollipop plot for BAT as described previously. (E) Immunofluorescence staining of proliferation in crown-like structures in db/db WAT through proliferative marker phosphohistone H3 in addition to wheat germ agglutinin (WGA) to mark cell membranes and DAPI to mark nuclei. (F) Feature plots showing expression of proliferation markers Top2a, Mki67, and Aurka in macrophage subsets. (G) Overlay of differentially expressed genes in all cell populations and macrophages for visceral WAT between the present study and a human obesity study (Emont et al. 2022).

Figure 5. Gene regulatory network alterations in T2D adipose depots and contribution to leukocyte migration. (A) RSS plots demonstrating the top 5 activated regulons in db/db (red dots) as opposed to db/h samples (blue dots) for white adipocytes, brown adipocytes, fibroblasts and macrophages for WAT (top panel) and BAT (lower panel). (B) Immunofluorescence validation of transcription factor IRF4 (grey) in both db/h and db/db BAT tissue with nuclei stained with DAPI (blue). Scale bar represents 25μm. (C) Differential enrichment of regulons with upregulated regulons in orange and downregulated regulons in blue. Intensity of color indicates the average expression in that cell population and dot size represents statistical significance. Regulons with bolded text represent those shared across both depots. Bar charts adjacent to regulons indicate the number of target genes within that regulon that are also linked to the GO term leukocyte migration.

Figure 6. Sex differences in gene expression and tissue architecture across adipose depots. (A) Lollipop plot demonstrating sex differences in gene expression between females and males in db/h WAT and BAT. (B) Compositional bar chart demonstrating the contribution of adipose tissue cell populations to each depot and how they differ between conditions and biological sexes. (C) Ridgeplots showing female and male sample expression of key thermogenic genes between db/h and db/db conditions in BAT. (D) Top 5 gene ontology terms for each cell population where dot size represents the number of genes linked to this GO term and dot color represents the statistical significance. GO terms of interest are highlighted in bold. (E) Immunofluorescence staining of BAT vascularization using isolectin B4 (IB4; red) and nuclei stained with DAPI (grey) demonstrating differences in tissue architecture between females and males in both db/h and db/db conditions. Scale bar represents 100μm. (F) Quantification of vascularization (% of IB4+ tissue area) and cell density (number of DAPI+ nuclei) histology with female data points presented in pink and males in blue. Statistical significance was assumed at P<0.05. * P<0.05, ** P<0.01, *** P<0.001.

Supplementary Figure 1. Optimization of nucleus isolation and flow cytometry gating strategy. (A) Flow cytometry-based optimization of the time length required to mechanically digested at 30hz using the Qiagen Tissue Lyser to get optimal nuclei/mg of tissue yield. (B) micrographs showing nuclei post Qiagen digestion and fluorescence-activated cell sorting to remove cellular debris. Scale bar represents 20μm. (C) Flow cytometry gating strategy for SVF cell population analysis. Viable and metabolically active single cells were gated, and cell populations were delineated based on staining by antibodies specific for cell surface markers. WAT, white adipose tissue; BAT, brown adipose tissue; FSC-H, forward scatter height; FSW-A, forward scatter area; FSC, forward scatter width; CB, calcein blue; SG, sytox green; ECs, endothelial cells; RMCs, resident mesenchymal cells; VECs, vascular endothelial cells; LECs, lymphatic endothelial cells.

Supplementary Figure 2. Cell population abundance changes in T2D. (A) Cell proportion differences between db/h and db/db samples based on nuclei count in WAT and BAT. Y axis indicates frequency of total cell count. (B) Contribution of adipose tissue cell populations to flow cytometry total viable cell count with differences between db/h (light color) and db/db samples (dark color) being shown in WAT (blue) and BAT (orange) depots. BA, brown adipocyte; WA, white adipocyte; ECs, endothelial cells; Meso. cells, mesothelial cells; Multipot. prog, multipotent progenitors; Prolif. macs, proliferating macrophages; SMCs, smooth muscle cells; WA, white adipocyte; LECs, lymphatic endothelial cells; VECs, vascular endothelial cells; Myel. cells, myeloid cells; Macs, macrophages; Grans, granulocytes; Non-myel. cells, on-myeloid cells; Fibro, fibroblasts. Statistical significance was assumed at P<0.05. * P<0.05, ** P<0.01, *** P<0.001, **** P<0.0001.

Supplementary Figure 3. Top 50 upregulated and downregulated genes in T2D WAT and BAT. Genes up-(red) downregulated (blue) between db/h and db/db cell populations in WAT (A) and BAT (B). Dot size represents the fold change (FC) difference between conditions while color intensity indicates the expression level (avg. exp. scale). ECs, endothelial cells; LECs, lymphatic endothelial cells; Meso. cells, mesothelial cells; Multipot. prog, multipotent progenitors; SMCs, smooth muscle cells.