Transcription Factor Condensates Mediate Clustering of *MET* Regulon and Enhancement

in Gene Expression

James Lee [1,2], Leman Simpson [2,3], Yi Li [1,2], Samuel Becker [1], Fan Zou [4], Xin Zhang [3], and Lu

Bai[1,2,4*]

[1]*Department of Biochemistry and Molecular Biology,* [2]*Center for Eukaryotic Gene Regulation,*

[3]*Department of Chemistry,* [4]*Department of Physics, The Pennsylvania State University,*

University Park, PA, 16802, USA

*Corresponding author: lub15@psu.edu, (814) 863-4824

Abstract

Some transcription factors (TFs) can form liquid-liquid phase separated (LLPS) condensates.

However, the functions of these TF condensates in 3D genome organization and gene regulation

remain elusive. In response to methionine (met) starvation, budding yeast TF Met4 and a few co-

activators, including Met32, induce a set of genes involved in met biosynthesis. Here, we show

that the endogenous Met4 and Met32 form co-localized puncta-like structures in yeast nuclei upon

met depletion. Recombinant Met4 and Met32 form mixed droplets with LLPS properties *in vitro*.

In relation to chromatin, Met4 puncta co-localize with target genes, and at least a subset of these

target genes are clustered in 3D in a Met4-dependent manner. A *MET3pr*-GFP reporter inserted

near several native Met4 binding sites becomes co-localized with Met4 puncta and displays

enhanced transcriptional activity. A Met4 variant with a partial truncation of an intrinsically

disordered region (IDR) shows less puncta formation, and this mutant selectively reduces the

reporter activity near Met4 binding sites to the basal level. Overall, these results support a model

where Met4 and co-activators form condensates to bring multiple target genes into a vicinity with

higher local TF concentrations, which facilitates a strong response to methionine depletion.

Introduction

Chromosomes form extensive 3D contacts that can occur over long linear genomic distances (Marenduzzo et al. 2007; Fazary et al. 2017; Misteli 2020). A subset of these interactions, such as promoter-promoter or promoter-enhancer looping, are thought to play a central role in gene regulation (Mifsud et al. 2015; Schoenfelder and Fraser 2019). In some cases, multiple genes come together in 3D space to form "multi-gene clusters" (Mir et al. 2018; Lim and Levine 2021; Zhu et al. 2021; Uyehara and Apostolou 2023). These clusters often involve co-regulated genes responding to stress or other types of environmental cues, such as heat shock, starvation, virus infection, or developmental signals (Ferrier and Minguillón 2003; Sawado et al. 2003; McCord et al. 2011; Du and Bai 2017; Zhao and Faryabi 2023). Evidence from imaging and genomic studies suggests that these gene clusters may be related to a phenomenon called "transcription factories", where RNA polymerase II (Pol II), mediator, and nascent transcripts coalesce into distinct foci (Bartlett et al. 2006; Faro-Trindade and Cook 2006; Carter et al. 2008; Xu and Cook 2008; Sutherland and Bickmore 2009; Papantonis and Cook 2011; Rieder et al. 2012). Proteomic analysis of transcription factories showed that they are also enriched with other components in the transcription pathway, including transcription factors (TFs), histone modification enzymes, and chromatin remodelers (Lyons et al. 2023). Overall, these observations suggest that some co-regulated genes can physically cluster into sub-nuclear compartments with elevated local concentrations of transcription-related factors.

How multi-gene clusters are formed is not well understood. Since co-regulated genes are often activated by a common set of TFs (Kiran and Nagarajaram 2016; Mir et al. 2018; Tsai et al. 2019; Uyehara and Apostolou 2023), such clustering may be mediated by the DNA-protein and protein-protein interactions of the TFs. Recent studies showed that TFs like Gcn4, OCT4, and TAZ, as well as some co-factors such as the mediator and histone-acetylation reader Brd4, tend to form biomolecular condensates through liquid-liquid phase separation (LLPS) (Whyte et al. 2013; Boija et al. 2018; Sabari et al. 2018; Gibson et al. 2019; Lu et al. 2020; Lee et al. 2021). In LLPS, multivalent interactions among proteins, often through their intrinsically disordered regions (IDRs), allow them to condensate into liquid-like droplets that are separated from the aqueous phase (Banani et al. 2017; Dignon et al. 2020; Peng et al. 2020; Mitrea et al. 2022). The existence of TF / co-factor condensates is supportive of the "transcription factory" model mentioned above where transcription-related factors are locally concentrated through LLPS. So far, LLPS of TFs and co-factors are often studied *in vitro* using purified systems or *in vivo* with artificial induction. How

these proteins behave in a chromatin context at the endogenous concentration, and more
specifically, if they contribute to multi-gene cluster formation, is not very clear.

Another unresolved issue related to multi-gene clusters is their functional role in gene regulation.
Earlier studies of gene clusters found a positive correlation between clustering and gene
expression level (Li et al. 2012; Whyte et al. 2013; Zhang and Bai 2016; Mir et al. 2018; Zhu et al.
2021; Zhao and Faryabi 2023), but the mechanism of the phenomenon was not clear. This again
may be related to TF / co-factor condensates: if a gene cluster is situated within a condensate,
higher local concentrations of these factors can potentially enhance expression. This is proposed
to be the case at super-enhancers, where the condensates of co-factors like Med1 and Brd4 are
thought to drive the high transcriptional activities of the target genes. However, some reports
studying TF condensates reveal a complex, sometimes contradictory, relation to gene expression,
where the condensates are observed to have positive, negative, or neutral effects on transcription
(Lu et al. 2020; Lee et al. 2021) . In a recent study of the *GAL* genes, for example, it is proposed
that Gal4 condensation facilitates their recruitment to target genes but does not contribute to gene
activation (Meeussen et al. 2023). In general, the relation between TF condensates, multi-gene
cluster, and gene regulation needs to be further studied.

Upon met depletion (-met) in budding yeast, Met4 is recruited by several sequence-specific DNA-
binding co-activators, including Cbf1, Met31, and Met32, to bind and rapidly activated a subset of
genes (Thomas et al. 1992; Lee et al. 2010; Ouni et al. 2010; Carrillo et al. 2012). This activator
complex is further stabilized with Met28 (Kuras et al. 1997). In a previous study, we carried out a
screen to probe long-distance gene regulation that affected *MET3* promoter (*MET3pr*) activity by
inserting an insulated *MET3pr*-GFP reporter into many genomic loci and measuring GFP intensity
in -met (Du et al. 2017). At most locations, the reporter has comparable induction level; however,
it shows enhanced activity at a small subset of loci, which were defined as "transcriptional
hotspots" (Du et al. 2017). These hotspots tend to be located near endogenous Met4 target genes,
and 3C assay detects physical proximity among the hotspots. These results lead to a hypothesis
that hotspots occur at loci where *MET* genes cluster, and such clustering and transcriptional
elevation may be both mediated by the condensates of related TFs.

In this study, we test the hypothesis above using a combination of biochemistry, genetics,
genomics, and imaging approaches. We demonstrate that Met4 and Met32 form co-localized

puncta in the yeast nuclei upon met depletion. Purified Met32 forms condensate with LLPS
properties *in vitro*, with which Met4 can merge. In relation to chromatin, we show that the loci that
are associated with the Met TFs co-localize with the Met4 puncta, and at least a subset of Met4
target genes cluster in 3D nuclear space in a Met4-dependent manner. Functionally, the *MET3pr-*
GFP reporter inserted near the endogenous Met4 target sites shows higher enrichment of Met
TFs / Pol II and enhanced transcriptional activity (hotspots). A Met4 IDR truncation that reduces
Met4 puncta formation selectively affects the reporter activity at hotspots. Collectively, our results
support a model where TFs in the met response pathway form LLPS condensates, organize
multiple target genes in their vicinity, and enhance gene expression by creating a local
environment with elevated concentrations of transcription-related factors. This may represent a
common strategy for cells to concentrate limited resources to mount a strong response to stress.
107
Results
The trans-activator Met4 and its co-factor Met32 form puncta in the nucleus. Our previous
study presented evidence that the "hotspots" for *MET3pr* transcription may form 3D clusters, but
the mechanisms of clustering and transcriptional elevation are unknown. Given the recent reports
on TF condensation (Sabari et al. 2018; Lu et al. 2020; Zhang et al. 2022; Okada et al. 2023), we
hypothesize that these phenomena may be related to condensates formed by TFs in the *MET*
pathway. To test this possibility, we labeled the endogenous Met4 protein with GFP and visualized
Met4 in live cells under Airyscan confocal microscopy (**Materials and Methods**; see **Table S1** for
strain list). We observed nuclear-localized Met4-GFP under both the repressed (+met) and
activated (-met) conditions, with the latter showing higher intensity. Importantly, the Met4-GFP
displays puncta-like structures, especially under -met (**Fig. 1A & S1A**). As controls, we
constructed strains expressing a nuclear-localized free GFP or GFP fused to various chromatin-
associated factors, including TFs Cbf1 and Reb1, and Sth1, a subunit of RSC nucleosome
remodeler, which also bind chromatin with sequence specificity (Cai and Davis 1990; Ju et al.
1990; Donovan et al. 2023). The GFP signals appear more evenly distributed in these control
strains (**Fig. 1A**). To evaluate the distribution more quantitatively, we calculated the coefficient of
variance (CV) of the fluorescent intensities among the pixels inside each nuclei (**Materials and**
Methods, **Table S2**). Uneven distribution of the GFP molecules would result in large variations in
the pixel intensities and higher CVs. The CVs of Met4-GFP in both ±met conditions are
significantly higher than free GFP and other nuclear proteins (**Fig. 1B**). Since the strains in **Fig.**
1A express GFP at different levels, which directly affect CV, we also calculated the Fano number

(variance divided by the mean) to control for this variable. Met4-GFP in the -met condition has
the largest Fano number (**Fig. 1C & S1B**), which indicates that in the activated state it forms the
most punctate structures. These data support the notion that Met4 forms condensates in the
nuclei upon activation.

We next carried out the same experiment with Met32, the DNA binding co-factor of Met4. We
imaged cells containing endogenous Met32 labeled with mCherry and compared its distribution
to free mCherry. Unlike Met4, Met32-mCherry is barely visible in +met, and it is significantly
upregulated in -met (**Fig. 1D & S1C**). In the latter condition, Met32-mCherry also shows uneven
distribution with high CV and Fano number (**Fig. 1E, F & S1D**). We also imaged Met31-mCherry,
but its concentration is too low to be detected. When Met4-GFP and Met32-mCherry are tagged
in the same strain, their fluorescent signals largely overlap (**Fig. 1G & S1E**). To quantify the
overlap, we extracted the GFP and mCherry fluorescence intensities for all the pixels inside each
nucleus and calculated their correlation coefficient. As a control, we carried out the same analysis
with co-expressed Sth1-GFP and Met32-mCherry, which appear to be less co-localized (**Fig. 1G**).
Indeed, the Pearson correlation between Met4 and Met32 fluorescence on average was 0.75,
much higher than that between Sth1 and Met32 ($R = 0.19$) (**Fig. 1H**). Overall, the puncta-like
structures of Met4 and Met32 and their co-localization support the possibility that these two factors
co-condensate under the -met condition.

Met32 forms condensates with LLPS properties *in vitro*. LLPS is proposed to be a mechanism
that leads to TF condensation (Banani et al. 2017; Boeynaems et al. 2018). To test if this
mechanism applies to Met4 and Met32, we first evaluated their propensity to phase separate
using the FuzDrop algorithm (Hatos et al. 2022). All the Met activators, including Met4, Met31,
Met32, and Met28, are predicted to have higher LLPS probabilities than GFP and mCherry (**Fig.
2A**). Among these TFs, Met4 and Met32 have particularly high scores, which suggests they
function as droplet drivers that can spontaneously phase separate (Vendruscolo and Fuxreiter
2022). In addition, both factors are predicted to have large unstructured domains, a feature known
to promote LLPS (**Fig. S2A & B**).

We next purified Met4 and Met32 to study their condensation properties *in vitro*. Like many other
recombinant TFs, Met4 is prone to degradation. We managed to extract full length Met4 tagged
with 6xHis and MBP on the N- and C-terminal, respectively, although some degradation products

could not be avoided (**Fig. 2B**) (**Materials and Methods**). We imaged the purified Met4 labeled

with Alexa488 in various buffers. Met4 remains diffusive up to 5 µM at physiological salt

concentrations and forms aggregates at higher density or lower salt concentrations (**Fig. 2C**).

Although these results suggest that Met4 may not inherently undergo LLPS, we cannot rule out

the potential effects from impurities (degradation products) and tags on our observations. In

contrast, we managed to purify high-quality Met32 fused with mCherry (**Fig. 2B**). At the

physiological salt concentration with no crowding agents, Met32-mCherry forms condensates at

various concentrations from 1 to 20 µM (**Fig. 2C & S2C**). These condensates show properties

consistent with LLPS such as droplet fusion (**Fig. 2D**) and fluorescent recovery after

photobleaching (FRAP) (**Fig. 2E**). The Met32-mCherry condensates have a 64 ± 9.7s half

recovery time, comparable to well-established LLPS proteins such as FUS (5-120 seconds),

DDX4 (3-200 seconds), and hnRNPA1 (5-120 seconds) (Nott et al. 2015; Shin et al. 2017;

McSwiggen et al. 2019).

To further test potential Met4 / Met32 co-condensation, we added purified Met4 and Met32

together in 150 mM NaCl buffer and imaged in both Alexa488 and mCherry channels. The two

proteins are mixed into droplets in solution (**Fig. 2F**). In these co-localized droplets, Met32-

mCherry still undergoes FRAP recovery with a slightly slower rate (83 ± 11s half recovery time)

(**Fig. S2D**). These results are consistent with the idea that Met4 and Met32 can co-condense

through a LLPS mechanism *in vitro*, which may contribute to the Met4 / Met32 puncta formation

in vivo.

Met4-activated genes co-localize with Met4 puncta. Since Met32 is a sequence-specific DNA

binder, the Met4 / Met32 condensates can potentially contact multiple target genes. This leads to

our model in **Fig. 3A**, which predicts that 1) at least some Met4 activated genes should be co-

localized with Met4 condensates, and 2) multiple target genes should cluster in 3D. To test these

predictions, we first performed chromatin immunoprecipitation with sequencing (ChIP-seq) of

related TFs (Met4, Met32, Met28, Cbf1), as well as RNA-seq, to identify genes directly targeted

by Met4. As expected, the bindings of Met4, Met32, and Met28 are strongly induced by met

depletion, while Cbf1, an abundant pioneer factor (Kuras et al. 1996; Blaiseau and Thomas 1998;

Ouni et al. 2010; Carrillo et al. 2012; Yan et al. 2018; Donovan et al. 2019), shows constitutive

binding (**Fig. 3B & C, S3A & B**). Under the induced condition, we identified 34 regions co-bound

by all four TFs (**Fig. 3C, Table S3**), and these binding sites potentially regulate 46 genes (some

sites are located in divergent promoters). Met depletion for 2 hrs causes wide-spread changes in

the mRNA level with both up- and down-regulation (**Fig. 3D, S3C**), reflecting the fundamental role

of methionine in metabolism (Thomas and Surdin-Kerjan 1997). Importantly, 40 out of the 46

genes co-bound by all four TFs show significant increased expression under the -met condition

(**Fig. 3D, Table S4**), indicating that they are direct targets of Met4 and its co-activators.

To determine if the Met4 target genes are co-localized with the Met4 puncta, we inserted 196X

tetO repeats near two of these genes, *MET13* and *MET6*, which allow us to visualize these loci

through their association with tetR-mCherry. We also created control strains with the same array

inserted near *PUT1* and the mating locus *MAT,* two loci with no Met4 association. These

chromatin dots were imaged together with Met4-GFP under ±met conditions. Visually, *MET13*

and *MET6* loci have higher probabilities to be co-localized with Met4 puncta in -met compared to

MAT and *PUT1* (**Fig. 3E**). To test this more quantitatively, we aligned these images with the

chromatin dots at the center and averaged the surrounding GFP signals (**Materials and**

Methods). For all four loci, the averaged GFP intensities are higher near the center because the

distant pixels are more likely to be outside nuclei boundaries, where Met4 signals are absent (**Fig.**

3F & G). In +met condition, Met4-GFP has low intensities. Upon activation, we can see localized

GFP dots with heightened peak intensities at the *MET13* and *MET6* loci in comparison to controls

(**Fig. 3F & G**). These results indicate that at least some Met4 target genes are associated with

Met4 condensates upon activation, presumably through sequence-specific co-binders like Met32.

Met4-dependent clustering of *MET* genes upon induction. We next tested the second

prediction of the model in **Fig. 3A**, i.e., Met4 condensates mediate the 3D clustering of target

genes. We first carried out a Hi-C measurement under -met condition, which did not show any

significant interactions among Met4 targets (Li et al. 2023). Given the limitations of Hi-C in

detecting fine-scale interactions (Oluwadare et al. 2019), we used an assay recently developed

in our lab, Methyltransferase Targeting-based chromosome Architecture Capture (MTAC) (Li et

al. 2023), to identify potential chromosomal interactions with selected Met4 targets (**Materials**

and Methods). In this method, a 256X LacO array is inserted near the gene of interest (viewpoint,

or VP) to recruit LacI fused with DNA methyltransferase M.CviPI, which methylates proximal

cytosines in the "GC" context in *cis* and in *trans* (**Fig. 4A**). Physical proximity to the VP can thus

be evaluated based on the enrichment of the methylation signal in comparison to a control strain

containing the same LacI-M.CviPI but no LacO array.

In a previous study (Li et al. 2023), we performed MTAC assays with a VP near the *MET6* gene under ± met condition. This VP 3kb away from *MET6* captures highly-specific inter-chromosomal interactions with four other Met4-targeted genes under -met, but not +met, condition. To further illustrate this point, we carried out MTAC with *MET13* as the VP (**Fig. 4B**). Under +met condition, MTAC only detects interactions from the adjacent local chromatin (**Fig. 4C, S4A**). When the same strain was switched to the -met condition, MTAC captures two far-*cis* interactions (intra-chromosomal interactions that are over 30kb apart) with two promoter regions near Met4 targeted genes (*STR3*, *MUP1*) (**Fig. 4C, S4A**). Interestingly, these two regions were also found to interact with the *MET6* VP (Li et al. 2023). Taken together, these data support the clustering of a subset of Met4-targeted genes upon activation.

The model in **Fig. 3A** implies that the clustering of *MET* genes is mediated by Met4 / Met32. To test this idea, we carried out MTAC using the *MET6* VP ± acute auxin-induced degradation of Met4 ± met (**Fig. S4B & C**). For the -met experiment, auxin (IAA) was added 30 mins prior to the met depletion so that Met4 activation would be largely eliminated (**Fig. 4D**). This would also prevent the accumulation of Met32, as it relies on Met4 for induction (Menant et al. 2006; Lee et al. 2010; Carrillo et al. 2012). Consistent with our previous observations, very few long-distance interactions were found in the +met repressed condition, regardless of the presence or absence of Met4 (**Fig. 4E, (1) and (3)**). Upon met depletion without IAA addition, we observed significant *trans* (inter-chromosomal) interactions between the *MET6* locus with three other Met4-targeted genes (*MUP1*, *STR3*, *YKG9*) (**Fig. 4E, (2)**). Importantly, these interactions disappear when Met4 was degraded (**Fig. 4E, (4)**). These data show that the presence of Met4, and likely Met32, is required for *MET* gene clustering. This is consistent with the model in **Fig. 3A** that clustering is mediated by Met4 / Met32 condensates.

Regions near Met TF binding sites constitute *MET* "transcriptional hotspot" in haploid yeast. We next investigated the functional role of Met4 condensates in gene expression. In our previous screen, we identified the *MET13* locus on chr7 as a "transcriptional hotspot": when the *S.kud MET3pr*-GFP reporter was inserted into *MET13* or neighboring genes, it showed higher GFP expression than most of the other insertion sites upon induction (Du et al. 2017). Combined with the observation that *MET13* locus co-localizes with the Met4 puncta (**Fig. 3E-G**), this suggests a potential link between the higher reporter activity and Met4 / Met32 condensates.

However, the initial screen was performed in diploids, while all the experiments above were
carried out in haploids. We therefore first tested if *MET13* also functions as a transcriptional
hotspot in haploid yeast. We again used the *S.kud MET3pr* for this test, as the sequence of this
promoter deviates significantly with the endogenous *MET3pr*, allowing them to be differentiated
in PCR (Du et al. 2017).

We created two haploid strains with the *S.kud MET3pr*-GFP reporter inserted near the *MET13*
gene (*CWC23* and *RSM23*) and measured its transcriptional profile (**Fig. 5A**) (**Materials and**
Methods). For comparison, we also inserted the same reporter into two non-hotspot basal loci
ATG36 and *LDS2* (**Fig. 5A**). The GFP intensities from the reporter at *CWC23* and *RSM23* loci
are ~50% higher (*P*-value < 1e-4) compared to the control under the -met condition (**Fig. 5B**).
Reporter GFP mRNA levels, as well as Pol II associated with the GFP ORF, are also significantly
higher in the former strains (**Fig. 5C & D**). These results are consistent with our previous
observations in diploid cells that the *MET13* locus promotes the transcriptional activity of the
MET3pr.

Given the model in **Fig. 3A**, we reasoned that the enhanced activities of the GFP reporter may
be due to increased local concentrations of TFs, which likely lead to higher TF occupancies. To
test this idea, we performed ChIP of Met4 and Met32 followed by qPCR on the *S.kud MET3pr* at
the hotspot vs basal genomic locations because it contains a Met32 motif. Consistent with our
expectation, we observed higher enrichment of Met4 and Met32 over the *S.kud MET3pr* at hotspot
locations (**Fig. 5E & F**). Taken together, these findings confirm the existence of "hotspot" loci in
haploid yeast and suggest that it may be caused by enhanced binding of Met32 and Met4.

Reporter activities near Met4 binding sites are enhanced over a ~40kb range. Besides
MET13, other loci that colocalize with Met4 puncta, like *MET6* (**Fig. 3E-F**), should also function
as transcriptional hotspots. To test this idea, we first verified that *MET6* remains associated with
Met4 puncta after the reporter insertion. We used the *MET6* yeast strain in **Fig. 3E** and integrated
a *S.kud MET3pr* reporter nearby (**Fig. 6A**). This reporter contains a frame-shifted GFP (fsGFP)
in order not to interfere with the Met4-GFP signal. As controls we integrated the same reporter
near two genomic regions not bound by Met TFs (*PUT1* and *ATG36*). We imaged these strains
in -met and analyzed them with the same method used in **Fig. 3F**. Interestingly, despite the fact
that *S.kud MET3pr* is bound and activated by Met4, insertion of this promoter does not increase

the local Met4 intensity at *MET6* and *PUT1* (**Fig. 6B & C**, ± rep). In comparison with the reporter-

containing *PUT1* and *ATG36* loci, *MET6* again shows stronger co-localization with the Met4

puncta (**Fig. 6B & C**). This finding indicates that *S.kud MET3pr* alone is insufficient in nucleating

a new Met4 condensate or recruiting the integrated locus to an existing condensate.

We then integrated the original *S.kud MET3pr*-GFP reporter into *MET6*, *ATG36*, and *LDS2* (the

latter two are controls far from any endogenous Met4 binding sites) (**Fig. 6D**) and compared their

GFP expressions. GFP shows significantly higher intensity at *MET6* in comparison with the two

basal loci (*P*-value < 1e-4) (**Fig. 6E**). To test the generality of this observation, we inserted the

reporter into two additional genomic loci associated with Met TFs, *MET17* and *RMD6*, and the

reporter shows higher GFP expression in both loci (**Fig. 6E**). This data supports a model where

Met4 condensates associated with the endogenous *MET* genes enhance the activity of a nearby

reporter. This model also predicts that such enhancement effect should be specific to the met

pathway, i.e., inducible genes activated by other TFs should not benefit from higher local

concentrations of *MET* activators. We tested this idea by inserting into the same loci a GFP

reporter driven by either *MET17pr*, a promoter in the same pathway, or by *GAL1Spr*, a promoter

controlled by a different set of TFs in another pathway (Rohde et al. 2000; Slot and Rokas 2010).

Indeed, *MET17pr* shows the same trend as *MET3pr*, while the strengths of *GAL1Spr* remain

constant across all these loci (**Fig. 6F**). These results demonstrate the pathway-specificity of the

transcriptional hotspots.

We next conducted experiments to determine the genetic range of the transcriptional hotspot. We

inserted the GFP reporter driven by *MET3pr* or *GAL1Spr* at various distances from the Met4

binding site in the *MET6* promoter (**Fig. 6G**). *MET3pr*-GFP shows higher expression level when

inserted up to 30kb upstream and 10kb downstream of the Met4 site. Beyond this region, the

MET3pr activity returns to the basal level (**Fig. 6H**). In contrast, *GAL1Spr* shows similar activities

among these insertion sites (**Fig. 6H**). These data indicate that, at least near the *MET6* gene, the

elevated expression of the *MET3pr* can occur over a ~40kb region.

The deletion of a disordered region in Met4 reduces puncta formation and reporter

expression at transcriptional hotspots. To determine the relation between Met4 condensates

and its gene regulatory function more directly, we introduced mutations to Met4 to reduce its

propensity to form condensates. Since LLPS is often promoted by IDRs, we first performed

PONDR analysis of Met4 (Xue et al. 2010). This analysis reveals three distinct stretches of IDRs

interspersed with two previously annotated functional domains: the activation domain (aa95-144),

which interacts with mediator, and the auxiliary domain (aa312-375), which interacts with

Met31/32 (**Fig. 7A**) (Kuras and Thomas 1995). AlphaFold also predicts that Met4 is largely devoid

of folded structures (**Fig. S2A**) (Jumper et al. 2021). To probe the effect of these IDRs on Met4

puncta formation, we first truncated individual IDR stretches: ΔIDR1 (aa1-69), ΔIDR2 (aa117-359),

or ΔIDR3 (aa397-651). As Met4 is essential for cell viability in the -met media, we constructed

strains containing the endogenous unlabeled Met4 with an additional copy of truncated Met4-GFP

driven by the native *MET4* promoter. This allows us to visualize Met4 mutants without hindering

cell growth (**Fig. 7B**). Deletions of IDR2 and IDR3 both lead to significant reductions in Met4-GFP

Fano number, with ΔIDR2 having a slightly more pronounced effect (**Fig. 7B & C, S5A**). To further

specify the part of IDR2 that affects Met4 condensates, we generated shorter truncations within

IDR2, ΔIDR2.1 (aa117-135), ΔIDR2.2 (aa136-270), and ΔIDR2.3 (aa271-359). All three

truncations result in intermediate GFP Fano number between the full length Met4 and Met4-

ΔIDR2 (**Fig. 7C, S5B & C**). These findings suggest that multiple IDRs contribute to Met4

condensation, and deleting parts of the Met4 IDR can diminish, but not completely eliminate, Met4

puncta formation.

When the full length Met4 is replaced by Met4 ΔIDR2.3, the strain is viable in the absence of met,

allowing us to evaluate the functional consequence of this mutant on Met4 binding and

transcriptional activation. Given our model and supporting evidence that Met4 condensates

increase local TF concentration at Met4 targeted genes and enhance their expression, we expect

a Met4 mutant less prone to condensate will result in diminished Met4 binding and reduced

induction. We therefore compared the ChIP-seq and RNA-seq data of the full-length vs truncated

Met4. Consistent with the idea above, the Met4 ΔIDR2.3 shows decreased binding across all the

previously annotated Met TF binding sites, without generating new ChIP-seq peaks (**Fig. 7D &**

E). Interestingly, the three Met4 binding sites near *MUP1*, *STR3*, and *YKG9* that cluster with

MET6 show significantly more reduction in Met4 ChIP-seq peaks (reduced to 42±4%) than other

Met4 sites (61±8%, *P*-value = 0.02) in the Met4 ΔIDR2.3 strain. Out of the 40 potential target

genes, 22 show significant decreases in their mRNA level (**Fig. 7F, S5D, Table S4**). Importantly,

our model also predicts that decreased Met4 condensation should have more impact on target

genes that are located near Met4 puncta than those farther away. To test this idea, we again took

advantage of the *MET3pr*-GFP reporter inserted into hotspots and basal loci. We expect the

reduced Met4 condensation to have a larger impact on the reporter activity in the former case
because the reporter is only associated with Met4 puncta at hotspots (**Fig. 6B**). Indeed, after
substituting Met4 with Met4-ΔIDR2.3 in these strains, we observed strong decreases in the
reporter gene activity in hotspots but minimal effect on those in basal regions (**Fig. 7G**). These
findings collectively support a model wherein Met4 condensates cluster target genes, enhance
their activity, and give rise to the formation of transcriptional hotspots.
366
Discussion
In this study, we investigated TF distribution, 3D genome organization, and gene activation in the
met response pathway using a combination of genetics, genomics, and imaging approaches. Our
findings connect three emerging phenomena in gene regulation: TF condensates, co-regulated
gene cluster, and transcriptional hotspot. More specifically, we show that the activator Met4 and
its co-factor Met32 form puncta-like structures that co-localize with at least some Met4 target
genes. Multiple *MET* genes on different chromosomes cluster in the 3D nuclei space, and such
clustering requires the presence of Met4. Chromosomal loci near endogenous Met4 binding sites,
in one case within a 40kb range, are hotspots for *MET3pr* reporter expression. Mutations that
decrease Met4 puncta formation lead to selective reduction of the reporter transcriptional activity
at the hotspots.
378
Condensate formation of Met4 and Met32. Using live-cell imaging, we found that fluorescently
labeled Met4 and Met32 from puncta-like structures in yeast nuclei under the -met condition. This
is reflected by the higher Fano numbers in Met4 / Met32 pixel intensities in comparison to free
GFP, and in the Met4 case, other chromatin-associated factors. High resolution microscopy of
yeast TFs is challenging due to the small size of the nuclei. The low intensities of the Met4 / Met32
signals require high excitation for imaging, which also makes them prone to photo-bleaching.
Thus, we were unable to measure Met4 and Met32 puncta dynamics *in vivo*. When purified *in*
vitro, Met32-mCherry readily forms droplets with typical LLPS properties without molecular
crowder, while His- and MBP-tagged Met4 remains diffusive at physiological salt concentration.
We also found that Met4 partitions into the Met32 droplets when they are mixed, and inside cells,
the two factors show significant co-localization. These observations raise the intriguing possibility
that Met32 serves as a scaffold to nucleate the co-condensates of Met4 and other Met TFs. This
may explain why the Met4 distribution becomes more puncta-like when switched from +met to -
met condition, when Met32 becomes available inside the nuclei. However, we cannot exclude the

possibility that the absence of Met4 LLPS is due to our *in vitro* condition (tag, impurity, lack of
post-translational modification, etc.). Met4 is also reported to recruit the Mediator and SAGA
complex (Leroy et al. 2006), and these factors, together with Pol II and RNA, may also affect Met4
condensate formation *in vivo* (Henninger et al. 2021).

TF condensates and the 3D genome topology. TF condensation *in vivo* occurs in a chromatin
context, but the relation between TF condensates and 3D genome topology has not been
extensively studied. By recruiting multiple TFs, mediator, and transcription machineries into
proximity, TF binding loci can potentially serve as nucleation sites for transcription condensates.
Looped enhancers and promoters, as well as clustered genes, may also function as scaffold and
stabilize these condensates. This is best illustrated at super-enhancers (SEs), where clustered
enhancer sites tend to associate with transcriptional condensates and lead to very high activities
of their target genes (Hnisz et al. 2013; Sabari et al. 2018). Synthetic LacO array and endogenous
microsatellite repeats were also shown to promote local condensation of associated TFs (Chong
et al. 2018). Reversely, TF condensates can mediate long-distance chromosomal interactions by
contacting multiple genomic sites and therefore inducing their 3D proximity. This idea is supported
by the findings here: 1) multiple Met4-targeted genes are co-localized with Met4 condensates, 2)
at least a subset of Met4-targeted genes form clusters with long-distance intra- and inter-
chromosomal interactions, and 3) such clustering only occurs in -met activating condition in the
presence of Met4. Similar observations were made during yeast heat shock response, where the
activator Hsf1 coalesces and mediates transient clustering of heat shock genes (Chowdhary et
al. 2019). A few other mammalian TFs were also shown to contribute to 3D genome conformation.
For example, YAP/TAZ organizes their target sites into clusters that co-localize with the
YAP/TAZ/TEAD1 condensates (Lu et al. 2020). NUP98-HOXA9, a chimeric TF related to the
pathogenesis of AML, was also shown to mediate long-distance chromosomal loops between
NUP98-HOXA9 bound enhancers and oncogenes into SE-like clusters (Ahn et al. 2021).

Interestingly, a *MET3pr* reporter that is associated and activated by Met4 only co-localizes with
Met4 puncta when inserted near native Met4 binding sites. Also, adding *MET3pr* reporter gene
into these sites does not further increase the local Met4 concentration (**Fig. 6B & C**). These
observations indicate that Met4 binding is not *sufficient* for a gene to nucleate or be recruited into
a Met4 condensate. The underlying mechanism of this observation is unclear, but we speculate
that some properties of the endogenous *MET* genes may facilitate their clustering with Met4

puncta. For example, some *MET* genes, like *MET13* and *MET6*, are associated with nuclear pore
complex (NPC) even in the presence of met (Forey et al. 2021). The tethering by NPCs may
increase the chance for these genes to come into proximity, enhancing the local density of Met4
target sites to nucleate the Met4 condensates. In contrast, a reporter gene inserted into random
genomic locations may not be optimally positioned to contact NPC and/or other Met4 target genes.
Further experiments are required to test these ideas. Notably, interaction with NPC was found to
be necessary and sufficient to promote clustering of the *GAL1-10* alleles (Brickner et al. 2016).

TF condensates and gene expression. The Met4 condensates represent a local environment
with elevated Met4 concentrations, which can promote Met4 binding and lead to enhanced gene
expression. Consistent with this idea, *MET3pr* reporter inserted near these Met4 targets, which
also co-localizes with the Met4 puncta, shows increased transcriptional activity. Near *MET6*, such
hyper-activity of the reporter occurs over a 40kb range. According to previous measurement
(Bystricky et al. 2004), 40kb linear chromatin explores a 3D space with ~200 nm diameter, which
may reflect the size of the Met4 condensates. The elevation of transcription level is pathway-
specific as *GAL1Spr*-GFP inserted into the same loci does not show the same effect. This finding
agrees with the "specialized transcription factory" previously proposed (Bartlett et al. 2006) and
argues against a model where the transcription elevation solely relies on general transcription
factors like mediator or transcription machinery.

The relation between Met4 condensation and transcriptional activity is further supported by Met4
ΔIDR2.3. This Met4 mutant, in comparison to full length Met4, reduces the Met4 puncta formation
and the activities of most of its target genes in the genome. Importantly, ΔIDR2.3 does not have
strong effect on the transcriptional activity of *MET3pr* reporter inserted into basal loci, suggesting
that this truncation does not directly impact Met4 integrity and its activation function. Instead, it
selectively reduces reporter activity at transcriptional hotspots, consistent with the idea that such
reduction is due to the disruption of Met4 condensates.

In terms of mechanism, we speculate that 1) high Met4 local concentration inside the condensates
may increase the binding rate of Met4, and/or 2) multivalent interactions among Met4, Met32, and
possibly other co-activators may slow down the dissociation of Met4 from its target genes. Both
ideas are consistent with our observation that Met4 ΔIDR2.3 shows less binding than the full
length Met4. The same mechanism has been proposed for other TF condensates, and in some

cases, supported by direct kinetic evidence (Trojanowski et al. 2022). It should be noted that more

TF binding does not necessarily lead to an increase in gene expression level, for example,

transcription may be limited by a step downstream of TF binding. Consistent with this idea, a

recent study reports that Gal4 condensation facilitates its recruitment to target genes but does

not contribute to gene activation (Meeussen et al. 2023). In human cells, artificially enhanced

condensation of a TF Ews::Fli1 causes sequestering of this TF into the nucleolus, resulting in

decreased expression of their target genes (Chong et al. 2022). This indicates that the exact

relation between TF condensates and gene expression relies on the nature of the multivalent

interaction, location of condensate formation, transcription kinetics, etc., and it needs to be

analyzed on a case-by-case basis.

TF condensates and stress response. The sulfur-containing amino acid methionine is a key

metabolite for cell survival, and its depletion represents a major stress for budding yeast.

Interestingly, the TFs that were found to form condensates in yeast, include Met4, Gal4, and Hsf1,

all respond to certain types of environmental stress. Similar observations are made in higher

eukaryotes. For example, YAP forms condensates in response to osmotic stress (Cai et al. 2019;

Lu et al. 2020), and phyB in plants uses altered LLPS properties to sense light and temperature

cues (Chen et al. 2022). In contrast, constitutive TFs in yeast, like Reb1, Cbf1, and Sth1, are more

evenly distributed in the nuclei (**Fig. 1**). The condensation of stress-responding TFs may reflect a

functional need to rapidly concentrate related resources into a subset of the nuclei space for more

efficient usage. In the case of Met4, its condensates indeed allow yeast cells to mount a stronger

response to met depletion. In addition, TF condensates may accelerate TF target search and

response rate, facilitate synchronized expression of target genes, and/or enhance the activation

specificity. These potential condensate functions need to be further explored.

Materials and Methods

Plasmid and strain construction. Standard molecular methods were used to construct budding

yeast strains and plasmids (**Table S1**). The haploid strain was derived from w303a background,

and most genetic modifications here involve homologous recombination. More specifically, to

create strains with GFP-tagged endogenous Met4, the *GFP-HIS3* fragment was PCR amplified

from a plasmid (pJL1) with primers containing upstream and downstream homologous sequences

of the endogenous *MET4* gene. The amplified PCR product was transformed into haploid yeast

to insert into the *MET4* locus. A similar strategy was used to create strains with mCherry-tagged

endogenous Met32, Met4-TAP, Met32-TAP, Met28-TAP and Cbf1-TAP strains. For strains with
tetO-tetR-mCherry labeled loci, variations of plasmid pSR11 were created from containing 196X
tetO repeats with flanking homologous sequences to targeted sites were linearized and
transformed into Met4-GFP expressing haploid strains. For expression of tetR-mCherry, a
plasmid (pMY39) containing the tetR-mCherry insert was linearized and transformed into the
ADE2 genomic locus. To create Met4 truncated strains, individual plasmids containing truncated
versions of Met4 (pJL1) were created and were linearized with restriction digestion. The linearized
plasmid was then integrated into haploid yeast directly upstream of the *MET4* locus. For
endogenous replacement of *MET4* with *MET4* (ΔIDR2.3), a plasmid variant of pJL1 with *MET4*
(ΔIDR2.3) was linearized and transformed to replace the endogenous *MET4* gene.

Microscopy and image analysis. For imaging of Met4-GFP and Met32-mCherry, yeast cells
were grown in SCD + 10X Met liquid media at 30°C to OD660 ~ 0.2, washed, and then transferred
onto 5 ml of SCD - Met media for 2 hrs for induction. Afterwards the cells were put on a SCD-Met
agarose pad and put under the confocal microscope for imaging. Confocal microscopy was
performed with Zeiss LSM880 scanning laser confocal (Penn State University Huck Life Sciences
Institute) with 488 nm Argon laser (541 nm detection) and 594 nm HeNe laser (648 nm detection)
using 63x/1.4 plan apochromat objective. For single cell fluorescent images, the images were
taken with Airyscan detector and deconvoluted with Zen Black software.

The individual nuclei from single cells were segmented to extract the nuclear boundaries and pixel
fluorescent information of single nuclei with NucleiSplicer (Matlab) and subsequent nuclei
information was analyzed with NucleiAnalyzer (Matlab). Variance and standard deviation were
calculated from the nuclear pixels and plotted with OriginPro (**Table S2**). The mean GFP
intensities were calculated by subtracting the measure GFP intensities by the mean background
intensities of unlabeled cells. Coefficient of Variation (CV) values were calculated by dividing the
Std. Dev by the mean GFP values for individual cells. The Fano number was calculated by dividing
the Variance by the mean GFP. To normalize the Fano number, the mean Fano number of Free
GFP was set to a value of 1 and the rest of the conditions were divided by the mean GFP Fano
number (**Table S2**).

Single cell images of yeast cells expressing tetR-mCherry and Met4-GFP were imaged with 10 z-
stacks (0.4 μm). The z-stack images of mCherry and GFP channels were analyzed by DotTracker

(MATLAB). For individual cells, the program detects the z-stack with the highest mCherry dot

intensity and extracts the GFP image from the same z position. These images were aligned with

the mCherry dot at the center. The mean intensity projection of the GFP channel was analyzed

with ImageJ.

To measure *S.kud* Met3pr-GFP expression, yeast cells were grown in SCD+10X Met liquid media

at 30˚C to OD660 ~ 0.2, washed, and then transferred onto a SCD-Met agarose pad for induction.

After 6 hrs, the agarose pad was put under a Leica DMI6000 B fluorescent microscope for imaging.

The GFP fluorescent intensity within each cell boundary was quantified using Celltracker4

(MATLAB) (Zou and Bai 2019).

Met4 and Met32 purification. Constructs of Met32-mCherry, and Met4-MBP were inserted into

pET28b(+) backbones with an N-terminal 6x-His tag. Met4 and Met32 genes were cloned from

budding yeast genomic DNA. MBP-tag was cloned from Addgene CAT#98651 and fusion

constructs were assembled using 2xHiFi mastermix (NEB #E2621L) and transformed into DH5a

cells (NEB C2987l). A single cysteine sequence was added to the linker between Met4 and MBP

to facilitate maleimide conjugation of fluorophores for imaging. For protein expression, all

plasmids were transformed into BL21(DE3)-Sigma32 cells (courtesy of Xin Zhang) and induced

according to reported protocol for the Sigma32 cell line (Zhang et al. 2014). Bacterial protein

expression was carried out with cells grown in LB supplemented with 50 mg/L kanamycin and

100 mg/L ampicillin at 37ºC to OD_{600}=0.3, induced with 2 g/L arabinose (TCI A0515). Cells were

then grown at 30ºC until OD_{600}=0.6, induced with 1% isopropyl b-D-1-thiogalactopyranoside

(IPTG, Research Products International I56000) and grown overnight at 20ºC. Cells were

resuspended in wash buffer supplemented with 1 mM phenylmethylsulfonyl fluoride (PMSF), and

frozen before lysis.

For protein purification, bacterial cells were lysed with sonication and treated with DNaseI and

RNaseA according to published protocols for 15 mins (Carrillo et al. 2012). Met32-mCherry was

purified using Ni-NTA chromatography (Thermo Scientific HisPur resin #88222) with proteins

buffers A (20 mM Tris pH 8.0, 400 mM NaCl, 5 mM imidazole) and B (20 mM Tris pH 8.0, 400

mM NaCl, 250 mM imidazole) with a stepwise gradient of buffer B. Protein was buffer exchanged

into buffer C (20 mM HEPES pH 7.5, 400 mM NaCl, 10% glycerol, 1 mM DTT) and concentrated

to at least 80 mM using Amicon Pro centrifugal filters 10,000 NMWL (Milipore #ACS501024). The

Met4 construct was purified using Ni-NTA chromatography with proteins buffers A and B with a

stepwise gradient of buffer B. Ni-NTA elution was immediately loaded onto an amylose column
(NEB E8021S) and purified using a stepwise gradient of buffers A and D (20 mM Tris pH 8.0, 400
mM NaCl, 10 mM maltose). Protein was labeled by incubating with 0.8 molar equivalents of Alexa
Flour 488 maleimide (Invitrogen A10254). Proteins were buffer exchanged into buffer C and
concentrated to less than 50 mM (to avoid aggregation).

Western blot analysis. Met4-MBP samples were run on a 4-20% SDS-PAGE gel (BioRad
#4561094). Gels were wet transferred to a PVDF membrane (BioRad #1620177) at 80V for 3 hrs
in Tris-Glycine buffer. Transfer was confirmed by Ponceau staining (Sigma P7170-1L). Membrane
was blocked with 5% non-fat milk, washed with TBS with 1% Tween 20 (VWR 0777). Membrane
was probed with anti-MBP antibody (Thermo Fisher MA527544). Membrane was stripped using
PVDF stripping solution (DOT Scientific 65000-500) according to manufacturer instructions and
reprobed with anti-6x-His primary antibody (Thermo Fisher MA1-21315) and conducted identically
to the first probing. All incubation steps proceeded overnight. Blots were developed using Pierce
ECL substrate (Thermo Scientific 32209) with a 2 mins exposure time.

Droplet formation. Proteins were diluted from 400 mM NaCl storage buffer so that the final
concentration of the buffer was 150 mM NaCl, 20 mM HEPES pH 7.4, 2% glycerol. Samples were
plated on a microscope slide with a hydrophobic spacer (Invitrogen #S24735) between the slide
and untreated coverslip. Droplets were allowed to settle for 5 mins prior to imaging. All droplet
formation and imaging was conducted at room temperature.

Fluorescent Recovery after Photobleaching (FRAP). Bleaching was conducted after three
frames using the 594 nm laser at full power for 45 iterations. Fluorescence intensity was monitored
every 3 seconds for a total of 270 seconds per experiment. Fluorescence intensity levels were
extracted using Zeiss Zen Black software. Fluorescence intensity was normalized to percent of
pre-bleach intensity then averaged. Error bars show +/- SD of the normalized fluorescence
intensity. FRAP experiments were conducted at 20 µM Met32-mCherry. Experiments were
replicated with at least three preparations of each protein. Snapshots of the bleaching process
were processed using FIJI.

Chromatin Immunoprecipitation (ChIP). Yeast strains were grown to OD_{660} of 0.4 in 50 mL of
SCD-met medium and crosslinked with 1.39 mL of 37% formaldehyde for 20 mins at room

temperature. Crosslinking was quenched by adding 2.7 mL of 2.5 M glycine and incubation for

5 mins. Crosslinked cells were centrifuged at 1882 × g for 3 mins at 4°C, and then washed twice

with cold 1x Tris-buffer saline (TBS). Cell pellet was then resuspended in 250 µL of fresh FSPP

buffer (50 mM HEPES-KOH, pH 7.5, 140 mM NaCl, 1 mM EDTA, 1% Triton X-100, 0.1%

sodium deoxycholate, 1% protease inhibitor cocktail, and 1 mM phenylmethylsulfonyl fluoride

(PMSF) and vortexed with ~300 µL glass beads for 2 cycles of 20 mins with an intervening

10 mins in a 4°C cold room. Another 250 µL of fresh FSPP was added to each sample, and the

cap and bottom of tubes were punched with a hot needle so that cell lysate could be collected by

centrifuging at 836 × g for 5 mins at 4°C. FSPP (0.5 mL) was added to the cell lysate to resuspend

the pellet. The cell lysate was then sonicated using a 30 s on 30 s off cycle at 4°C for 7 cycles.

Sonicated cell lysate was transferred to 1.5 mL tube and centrifuged at 17,136 × g for 20 mins at

4°C. Two hundred microliters of supernatant was saved as input. Another 400 µL of supernatant

was mixed with 600 µL of FSPP buffer and 20 µL of pre-blocked Magnetic IgG beads and TAP

antibody (Thermo CAB1001) or Rpb3 antibody (Biolegend 920204) for 10–12 hrs at 4ºC.

Chromatin was incubated with the beads overnight at 4°C, and then washed sequentially at 4°C

with 1xFA-lysis buffer (50 mM HEPES-KOH, pH 7.5, 140 mM NaCl, 1 mM EDTA, 1% Triton X-

100, 0.1% sodium deoxycholate), 1xFA-lysis buffer containing 150 mM NaCl, 1xFA-lysis buffer

containing 500 mM NaCl, LiCl buffer (0.25 M LiCl, 1% NP-40, 1% sodium deoxycholate, 1 mM

EDTA, and 10 mM Tris-HCl pH 8.0), and lastly TE buffer, pH 8.0. Chromatin was then eluted with

400 µL ChIP elution buffer (50 mM NaCl, 50 mM Tris-HCl, pH 8.0, 10 mM EDTA, 1% SDS).

Samples were mixed by rotation at 30°C for 30 mins. Beads were pelleted by centrifugation at

17,136 × g and discarded. Five microliters of 20 mg/mL Proteinase K was added to supernatant;

180 µL of 1xFA-lysis buffer, 20 µL of 10% SDS and 5 µL of 20 mg/mL Proteinase K were added

into the input samples. ChIP and input samples were then incubated overnight at 65°C to reverse

crosslinking. DNA was extracted by phenol:chloroform:isoamyl alcohol (25:24:1) followed

by ethanol precipitation. Input samples were then treated with 200 µg RNase A and precipitated

with ethanol after adding 20 µg glycogen. Enrichments of target TFs were quantified by

quantitative PCR (qPCR). All ChIP experiments were done in biological duplicates.

RNA-seq. Cells were grown in 5 ml of SCD +10x met overnight. 5 ml of fresh SCD +10x was

innoculated with 5 µl of saturated culture until OD_{660} ~0.2. The culture is collected by

centrifugation and washed 3x with sterile H2O. The collected culture is then grown in SCD -met

media for 2 hrs to reach OD_{660} ~0.4 and centrifuged at 300g for 4 mins at RT. The pellet was

washed twice with sterile H2O and resuspended in 250 µl of RNA lysis buffer (10mM Tris-HCl pH8.5, 5mM EDTA, 2% SDS, 2% stock 2-mercaptoethanol) and placed on a heat block at 85ºC for 20 mins mixing every 2 mins. The mixture was centrifuged at 12,000g for 5 mins and the supernatant was transferred to a new tube. The supernatant was then mixed with 1 ml of trizol and heated at 65ºC for 20 mins with mixing in between. Standard trizol/chloroform RNA extraction protocol was followed. The extracted RNA was analyzed with Tapestation to check their integrity. RNA with RIN (>7.5) was used for downstream analysis. Messenger RNA was captured using RNA purification beads (NEB). The eluted mRNA was fragmented and denatured for first and second strand synthesis for conversion into cDNA. The sequencing libraries were constructed using conventional protocols (NEB).

Methyltransferase Targeting-based chromosome Architecture Capture (MTAC). Yeast cells were grown in 50 ml SCD +10x Met media at OD_{660} 0.3. The cells were centrifuged and washed 3 times with autoclaved H_2O before transfer to 50 ml of SCD-Met and induced with 2 nM β-eastradiol for 2 hrs. After induction the cells were centrifuged at 4000g for 5 mins and collected in 800 µl of TE and 200 µl lysis buffer (0.5 M EDTA pH 8.0, 1 M Tris pH 8.0, 10% SDS) and incubated at 65 ºC for 30 mins. The cell lysate (1ml) was then sonicated using a 30 s on 30 s off cycle at 4°C for 11 cycles. The sonicated lysate was centrifuged at 12,000g at 4 ºC for 20 mins and the supernatant was transferred to a new tube. The supernatant was mixed with 500 µl for Phenol:Chloroform:Isoamyl Alcohol (25:24:1) nucleic acid extraction. The methylated genomic DNA (5 µg) was incubated with 2 µg of anti-methylcytosine antibody in 1X IP buffer on a rotating platform 4°C for 8-12 hrs. Input DNA (1%, 50 ng) was set aside for normalization. The lysate/antibody complex was incubated with protein A/G agarose beads for 2 hrs and washed 3x with 1x IP buffer. The beads were resuspended with 210 µl digestion buffer and incubated on a rotating platform at 55 ºC for 2 hrs. The eluted DNA was precipitated in 1 ml of ethanol and 3 µl glycogen (20mg/ml) and eluted in 60 µl of TE.

For strains with Met4 Auxin-induced degradation, we tagged the endogenous Met4 protein with V5-AID (Met4-V5-AID) and introduced OsTIR1 to the cells. Depletion of the Smc2 was achieved by adding auxin (Sigma, I2886) to a final concentration of 500 µM. Strains were incubated with auxin for 30 mins prior to MTAC experiment. Depletion was confirmed by western blot.

Data availability

657 All sequencing data in this study can be accessed on GEO series GSE252386, which include 1)
658 GSE252384: ChIP-seq raw data (fastq), bed files, and bigWig files, 2) GSE252385: RNA-seq raw
659 data (fastq), counts data (featureCounts), and DESeq2 tables, and 3) GSE252985: MTAC raw
660 data (fastq), bigWig files, counts data (featureCounts) and DESeq2 tables.

Acknowledgements

We thank Dr. Joseph Reese for providing auxin-inducible degron plasmids and yeast strains. We
acknowledge all members in the Bai lab for insightful comments on the manuscript. We thank the
Microscopy Facility and the CSL Behring Fermentation Facility at Penn State for their help with
imaging and protein purification. We are grateful to Dr. Cheryl Keller at the Genomics Research
Incubator at the Huck Institutes of the Life Sciences for helping with the genomic assays. We also
want to thank the members of the Center of Eukaryotic Gene Regulation at PSU for discussions
and technical support. This work is supported by the National Institutes of Health (R35 GM139654
to L.B.) and National Science Foundation (MCB- 2016266 to L.B.).

Author Contributions

673 L.B., J.L., L.S. and Y.L. designed the experiments; J.L. performed most of the experiments and
674 data analysis with help from S.B.; L.S. and Y.L. carried out most of the experiments in Fig. 2 and
675 Fig. 4, respectively; F.Z. contributed to the image data analysis; X.Z. helped with protein
676 purification; J.L. and L.B. wrote the manuscript.

Conflict of Interest Statement

679 The authors declare no conflict of interest.

Figure Legends

Figure 1. The trans-activator Met4 and its co-factor Met32 form puncta in the nucleus. A)
Representative images of Met4-GFP in ± methionine (met) conditions. Control cells expressing

free GFP (driven by the *HOpr*), Cbf1-GFP, Reb1-GFP, and Sth1-GFP cells are shown with similar

contrasts. Scale bars represent 4µm (same as in D & G). **B & C)** The mean coefficient of variance

(CV) and Fano numbers of nuclear pixel intensities for different versions of GFPs in panel A. The

Fano numbers are normalized so that the free GFP has a Fano number 1. **D)** Representative

images of strains expressing Met32-mCherry in +/- met conditions. Control strain expressing free

mCherry (driven by the *HOpr*) is shown with similar contrasts. **E & F)** The mean CV and normalized

Fano number of pixel intensities of Met32-mCherry vs free mCherry. **G)** Representative images

of strains co-expressing Met4-GFP and Met32-mCherry (top row), or Sth1-GFP and Met32-

mCherry (bottom row). **H)** A histogram of Pearson correlation coefficient between co-expressed

Met4-GFP and Met32-mCherry signals (red bars *N*=93) or Sth1-GFP and Met32-mCherry signals

(black bars *N*=149) in each cell.

Figure 2. Met32 forms condensates with LLPS properties *in vitro*. A) Probabilities of individual

Met TFs (Met4, Met32, Met31, Met28), GFP, and mCherry to undergo liquid-liquid phase

separation from a published prediction program (Fuzdrop). **B)** Coomassie staining and western

blots of purified Met4 and Met32 proteins. **C)** Protein aggregate and droplet formation observed

for Met4-MBP and Met32-mCherry fusion proteins. Met4 and GFP are 5 µM, Met32-mCherry and

mCherry are 10 µM in 20 mM HEPES pH 7.5, 150 mM NaCl buffer. Met4 and GFP are in 20 mM

HEPES pH 7.5, 150 mM NaCl or 50 mM NaCl. Met4 is labeled with Alexa 488 (AF488). Scale bar

represent 10 µm (same as in D-F). **D)** Merging of Met32-mCherry droplets. Met32-mCherry is at

30 µM in 20 mM HEPES pH 7.5, 150 mM NaCl. **E)** Fluorescence recovery after photobleaching

(FRAP) data for 20 µM Met32 droplets. The intensity data was collected every 3 seconds for 270

seconds and normalized to percent bleaching. Inset: Representative images of Met32 FRAP. **F)**

Colocalization of Met32-mCherry droplets (10 µM) with Met4-MBP (5 µM) in 150 mM NaCl.

Figure 3. Met4-activated genes co-localize with Met4 puncta. A) Model of Met4 condensate in

the chromatin context. Upon met depletion, Met TFs may form condensates that interact with

multiple target genes, leading to the 3D clustering of these co-regulated genes. **B)** Examples of

ChIP-seq signals of Met TFs (Met4, Met32, Met28 and Cbf1) in ±met conditions. The Met TF co-

binding peak shown here is near the *MET13* gene. **C)** Heatmaps of Met4, Met32, Met28, and Cbf1

ChIP-seq peaks in ± met conditions. The peaks were clustered into ones enriched with all four

Met TFs (*N* = 34), enriched with Met4 and Cbf1 (*N* = 11), and enriched with Cbf1 only (*N* = 107).

D) Volcano plot of RNA-seq data comparing mRNA levels in ± met conditions. Most genes near

Met TFs co-binding peaks are strongly induced by met depletion (red dots). **E)** Single nuclei

images of yeast cells expressing Met4-GFP and TetR-mCherry with a TetO array integrated near
MET13 / MET6 (Met4 targets), or *MAT / PUT1 (*non-targeted *c*ontrol). Images were taken with 14
z-stacks with step size 0.4 μm. "MIP" and "same Z" show Met4-GFP images with either maximum
intensity projection among all z stacks, or with a single stack at the same z plane as "mCherry",
where the mCherry labeled TetO array shows the highest intensity. "merged" is the merged image
of the "same Z" and the "mCherry". Scale bars represent 1μm (same as in F). **F)** Average mCherry
and GFP intensities centered at the mCherry labeled TetO arrays located at different genomic
sites. **G)** The GFP intensity profile across the dot center in panel E. The GFP intensity near the
dot center is significantly higher for *MET13 / MET6* loci.

Figure 4. Met4-dependent clustering of *MET* genes upon induction. A) MTAC workflow. In a
"targeted" MTAC strain, a LacO array is integrated into a genomic locus (viewpoint, VP) and
recruits LacI-M.CviPI, an ectopic DNA methyltransferase that methylates the cytosine in a "GC"
dinucleotide in proximal DNA. LacI-M.CviPI is also expressed in a control strain with no LacO
array insertion (background methylation). Methylation in these two strains is detected by ChIP,
and methylation level in nucleosome-depleted regions (NDRs) are compared in targeted vs control
strains. Significantly higher methylation in the targeted strain indicates proximity to the VP. **B)** VP
design of the *MET13* locus. **C)** Volcano plot of MTAC signals with *MET13* as VP in ±met conditions.
Each dot represents an individual NDR, and colored dots are the NDRs that show proximity to the
VP (significantly higher methylation in the targeted vs control strains). Local (intra-chromosomal
interactions within 30kb), far-cis (intra-chromosomal interactions over 30kb), and trans (inter-
chromosomal interactions) are shown in blue, green, and orange. Same as below. **D)** Design of
Met4 depletion assay. Met4 is depleted by auxin-degron system in ±met conditions, resulting in
four conditions: (1) +met, -IAA, (2) -met, -IAA, (3) +met, +IAA, (4) -met, +IAA. β-estradiol is added
in all conditions to induce the expression of LacI-M.CviPI. **E)** Volcano plot of MTAC signals with
MET6 as VP for the four conditions in panel D. Note that long-distance interactions are detected
in condition (2) but are largely absent in other three conditions. **F)** MTAC and Met4 ChIP-seq data
at the *MET6* locus (VP) and *MUP1, STR3* and *YKG9* as interacting regions of the VP. MTAC
signal is shown in the four conditions in panel D. The ChIP enrichment of Met4 is shown in ±met
conditions. *P,* FDR-adjusted *P* value, Wald test by DESeq2. ns, non-significant.

Figure 5. Characterization of a *MET* "transcriptional hotspot" in haploid yeast. A) Schematic
of strains constructed with *S.kud MET3pr*-GFP reporter inserted near the *MET13* transcriptional
hotspot (*CWC23, RSM23*) and "basal" loci (*ATG36, LDS2*). **B & C)** Mean cellular GFP fluorescent

intensities and GFP mRNA measured by qRT-PCR in the four strains above in ±met conditions.

D) ChIP-qPCR of Rpb1 over the GFP ORF in the four strains above in -met. **E & F)** ChIP-qPCR

of Met32 and Met4 over the *S.kud MET3pr* in the four strains above in -met. The asterisks

represent *: <0.05, **: <0.01, ***: <0.001. Same as below.

Figure 6. Reporter activities near Met4 binding sites are enhanced over a ~40kb range. A)

Schematics of measuring the co-localization of the GFP reporter with Met4 puncta. *S.kud MET3pr-*

fsGFP (frameshifted GFP) reporter gene and a tetO array (196x) are inserted side by side into the

genome, in this case near the *MET6* gene. **B)** Average mCherry and GFP intensities centered at

the mCherry labeled tetO arrays near *MET6* (Met4 target) and *PUT1 / ATG36* (not Met4 targets)

in the presence of nearby reporter. Scale bars represent 1μm. **C)** The GFP intensity profile across

the dot center in panel B. For *MET6* and *PUT1,* the same type of data without the GFP reporter (-

rep) are also included. **D)** Schematic of GFP reporter insertion near three additional Met4-bound

loci, *MET17, RMD6, and MET6.* The distance and orientation of the insertion are labeled in the

diagram. **E)** Mean GFP fluorescent intensities when *S.kud MET3pr*-GFP are inserted near

indicated genes. The genes labeled in red have adjacent Met4-bound sites, while the ones labeled

in grey do not. **F)** Same as in panel E except with *MET17pr*-GFP and *GAL1Spr*-GFP reporter.

Strains with *MET17pr*-GFP were grown in -met, and the ones with *GAL1Spr*-GFP were pre-grown

in raffinose and induced by galactose for 6 hours. **G)** Schematic of the *MET3pr*-GFP reporter

inserted at various distances from the Met4 binding site near the *MET6* gene. The same

orientation was used for all the loci as indicated. **H)** Mean GFP fluorescent intensities with

MET3pr-GFP and *GAL1Spr*-GFP reporters inserted into locations indicated in panel G, in

comparison to the same reporter inserted into two control loci far from Met4 binding sites.

Figure 7. The deletion of a disordered region in Met4 reduces puncta formation and reporter

expression at transcriptional hotspots. A) PONDR disorder plot of Met4 with previously

annotated functional domains, including activation domain (aa95-144) that interacts with mediator,

inhibitory domain (aa188-235), auxiliary domain (aa312-375) that interacts with Met31/32, and

bZIP domain (aa595-660) that interacts with Cbf1. **B)** Single nuclei images of cells expressing full

length Met4 (±met) or Met4 with various truncations (-met) fused with GFP. **C)** The Fano numbers

of nuclear pixel intensities for different versions of GFPs in panel B. **D)** Heatmaps of full length

Met4 and Met4 ΔIDR2.3 ChIP-seq in –met over previously identified Met TF binding sites. **E)**

Alignment counts underneath the Met TF ChIP-seq peaks over the same sites for TAP tagged full

length Met4 and Met4 ΔIDR2.3 (sorted from high to low). **F)** Volcano plot from RNA-seq data from

cells containing full length Met4 or Met4 Δ IDR2.3 in -met. Genes associated with Met TFs (Fig. 3C) are indicated in red. **G)** Mean GFP intensities from cells expressing full length Met4 and Met4 ΔIDR2.3 with *MET3pr*-GFP reporter inserted into loci near (red) or far away from (black) Met4-target genes. The dashed lines represent basal reporter expression.

References

Ahn JH, Davis ES, Daugird TA, Zhao S, Quiroga IY, Uryu H, Li J, Storey AJ, Tsai YH, Keeley DP et al. 2021. Phase separation drives aberrant chromatin looping and cancer development. *Nature* **595**: 591-595.

Banani SF, Lee HO, Hyman AA, Rosen MK. 2017. Biomolecular condensates: organizers of cellular biochemistry. *Nat Rev Mol Cell Biol* **18**: 285-298.

Bartlett J, Blagojevic J, Carter D, Eskiw C, Fromaget M, Job C, Shamsher M, Trindade IF, Xu M, Cook PR. 2006. Specialized transcription factories. *Biochem Soc Symp*: 67-75.

Blaiseau PL, Thomas D. 1998. Multiple transcriptional activation complexes tether the yeast activator Met4 to DNA. *EMBO J* **17**: 6327-6336.

Boeynaems S, Alberti S, Fawzi NL, Mittag T, Polymenidou M, Rousseau F, Schymkowitz J, Shorter J, Wolozin B, Van Den Bosch L et al. 2018. Protein Phase Separation: A New Phase in Cell Biology. *Trends Cell Biol* **28**: 420-435.

Boija A, Klein IA, Sabari BR, Dall'Agnese A, Coffey EL, Zamudio AV, Li CH, Shrinivas K, Manteiga JC, Hannett NM et al. 2018. Transcription Factors Activate Genes through the Phase-Separation Capacity of Their Activation Domains. *Cell* **175**: 1842-1855.e1816.

Brickner DG, Sood V, Tutucci E, Coukos R, Viets K, Singer RH, Brickner JH. 2016. Subnuclear positioning and interchromosomal clustering of the GAL1-10 locus are controlled by separable, interdependent mechanisms. *Mol Biol Cell* **27**: 2980-2993.

Bystricky K, Heun P, Gehlen L, Langowski J, Gasser SM. 2004. Long-range compaction and flexibility of interphase chromatin in budding yeast analyzed by high-resolution imaging techniques. *Proc Natl Acad Sci U S A* **101**: 16495-16500.

Cai D, Feliciano D, Dong P, Flores E, Gruebele M, Porat-Shliom N, Sukenik S, Liu Z, Lippincott-Schwartz J. 2019. Phase separation of YAP reorganizes genome topology for long-term YAP target gene expression. *Nat Cell Biol* **21**: 1578-1589.

Cai M, Davis RW. 1990. Yeast centromere binding protein CBF1, of the helix-loop-helix protein family, is required for chromosome stability and methionine prototrophy. *Cell* **61**: 437-446.

Carrillo E, Ben-Ari G, Wildenhain J, Tyers M, Grammentz D, Lee TA. 2012. Characterizing the roles of Met31 and Met32 in coordinating Met4-activated transcription in the absence of Met30. *Mol Biol Cell* **23**: 1928-1942.

Carter DR, Eskiw C, Cook PR. 2008. Transcription factories. *Biochem Soc Trans* **36**: 585-589.

Chen D, Lyu M, Kou X, Li J, Yang Z, Gao L, Li Y, Fan LM, Shi H, Zhong S. 2022. Integration of light and temperature sensing by liquid-liquid phase separation of phytochrome B. *Mol Cell* **82**: 3015-3029.e3016.

821 Chong S, Dugast-Darzacq C, Liu Z, Dong P, Dailey GM, Cattoglio C, Heckert A, Banala S, Lavis L, Darzacq X
822 et al. 2018. Imaging dynamic and selective low-complexity domain interactions that control gene
823 transcription. *Science* **361**.
824 Chong S, Graham TGW, Dugast-Darzacq C, Dailey GM, Darzacq X, Tjian R. 2022. Tuning levels of low-
825 complexity domain interactions to modulate endogenous oncogenic transcription. *Mol Cell* **82**:
826 2084-2097.e2085.
827 Chowdhary S, Kainth AS, Pincus D, Gross DS. 2019. Heat Shock Factor 1 Drives Intergenic Association of Its
828 Target Gene Loci upon Heat Shock. *Cell Rep* **26**: 18-28.e15.
829 Dignon GL, Best RB, Mittal J. 2020. Biomolecular Phase Separation: From Molecular Driving Forces to
830 Macroscopic Properties. *Annu Rev Phys Chem* **71**: 53-75.
831 Donovan BT, Chen H, Eek P, Meng Z, Jipa C, Tan S, Bai L, Poirier MG. 2023. Basic helix-loop-helix pioneer
832 factors interact with the histone octamer to invade nucleosomes and generate nucleosome-
833 depleted regions. *Mol Cell* **83**: 1251-1263.e1256.
834 Donovan BT, Chen H, Jipa C, Bai L, Poirier MG. 2019. Dissociation rate compensation mechanism for
835 budding yeast pioneer transcription factors. *Elife* **8**.
836 Du M, Bai L. 2017. 3D clustering of co-regulated genes and its effect on gene expression. *Curr Genet* **63**:
837 1017-1021.
838 Du M, Zhang Q, Bai L. 2017. Three distinct mechanisms of long-distance modulation of gene expression in
839 yeast. *PLoS Genet* **13**: e1006736.
840 Faro-Trindade I, Cook PR. 2006. Transcription factories: structures conserved during differentiation and
841 evolution. *Biochem Soc Trans* **34**: 1133-1137.
842 Fazary AE, Ju YH, Abd-Rabboh HSM. 2017. How does chromatin package DNA within nucleus and regulate
843 gene expression? *Int J Biol Macromol* **101**: 862-881.
844 Ferrier DE, Minguillón C. 2003. Evolution of the Hox/ParaHox gene clusters. *Int J Dev Biol* **47**: 605-611.
845 Forey R, Barthe A, Tittel-Elmer M, Wery M, Barrault MB, Ducrot C, Seeber A, Krietenstein N, Szachnowski
846 U, Skrzypczak M et al. 2021. A Role for the Mre11-Rad50-Xrs2 Complex in Gene Expression and
847 Chromosome Organization. *Mol Cell* **81**: 183-197.e186.
848 Gibson BA, Doolittle LK, Schneider MWG, Jensen LE, Gamarra N, Henry L, Gerlich DW, Redding S, Rosen
849 MK. 2019. Organization of Chromatin by Intrinsic and Regulated Phase Separation. *Cell* **179**: 470-
850 484.e421.
851 Hatos A, Tosatto SCE, Vendruscolo M, Fuxreiter M. 2022. FuzDrop on AlphaFold: visualizing the sequence-
852 dependent propensity of liquid-liquid phase separation and aggregation of proteins. *Nucleic Acids*
853 *Res* **50**: W337-W344.
854 Henninger JE, Oksuz O, Shrinivas K, Sagi I, LeRoy G, Zheng MM, Andrews JO, Zamudio AV, Lazaris C,
855 Hannett NM et al. 2021. RNA-Mediated Feedback Control of Transcriptional Condensates. *Cell*
856 **184**: 207-225.e224.
857 Hnisz D, Abraham BJ, Lee TI, Lau A, Saint-André V, Sigova AA, Hoke HA, Young RA. 2013. Super-enhancers
858 in the control of cell identity and disease. *Cell* **155**: 934-947.
859 Ju QD, Morrow BE, Warner JR. 1990. REB1, a yeast DNA-binding protein with many targets, is essential for
860 growth and bears some resemblance to the oncogene myb. *Mol Cell Biol* **10**: 5226-5234.
861 Jumper J, Evans R, Pritzel A, Green T, Figurnov M, Ronneberger O, Tunyasuvunakool K, Bates R, Žídek A,
862 Potapenko A et al. 2021. Highly accurate protein structure prediction with AlphaFold. *Nature* **596**:
863 583-589.
864 Kiran M, Nagarajaram HA. 2016. Interaction and localization diversities of global and local hubs in human
865 protein-protein interaction networks. *Mol Biosyst* **12**: 2875-2882.

Kuras L, Barbey R, Thomas D. 1997. Assembly of a bZIP-bHLH transcription activation complex: formation of the yeast Cbf1-Met4-Met28 complex is regulated through Met28 stimulation of Cbf1 DNA binding. *EMBO J* **16**: 2441-2451.

Kuras L, Cherest H, Surdin-Kerjan Y, Thomas D. 1996. A heteromeric complex containing the centromere binding factor 1 and two basic leucine zipper factors, Met4 and Met28, mediates the transcription activation of yeast sulfur metabolism. *EMBO J* **15**: 2519-2529.

Kuras L, Thomas D. 1995. Functional analysis of Met4, a yeast transcriptional activator responsive to S-adenosylmethionine. *Mol Cell Biol* **15**: 208-216.

Lee JH, Wang R, Xiong F, Krakowiak J, Liao Z, Nguyen PT, Moroz-Omori EV, Shao J, Zhu X, Bolt MJ et al. 2021. Enhancer RNA m6A methylation facilitates transcriptional condensate formation and gene activation. *Mol Cell* **81**: 3368-3385.e3369.

Lee TA, Jorgensen P, Bognar AL, Peyraud C, Thomas D, Tyers M. 2010. Dissection of combinatorial control by the Met4 transcriptional complex. *Mol Biol Cell* **21**: 456-469.

Leroy C, Cormier L, Kuras L. 2006. Independent recruitment of mediator and SAGA by the activator Met4. *Mol Cell Biol* **26**: 3149-3163.

Li G, Ruan X, Auerbach RK, Sandhu KS, Zheng M, Wang P, Poh HM, Goh Y, Lim J, Zhang J et al. 2012. Extensive promoter-centered chromatin interactions provide a topological basis for transcription regulation. *Cell* **148**: 84-98.

Li Y, Lee J, Bai L. 2023. DNA Methylation-Based High-Resolution Mapping of Long-Distance Chromosomal Interactions in Nucleosome-Depleted Regions. *bioRxiv*: 2023.2012.2027.573430.

Lim B, Levine MS. 2021. Enhancer-promoter communication: hubs or loops? *Curr Opin Genet Dev* **67**: 5-9.

Lu Y, Wu T, Gutman O, Lu H, Zhou Q, Henis YI, Luo K. 2020. Phase separation of TAZ compartmentalizes the transcription machinery to promote gene expression. *Nat Cell Biol* **22**: 453-464.

Lyons H, Veettil RT, Pradhan P, Fornero C, De La Cruz N, Ito K, Eppert M, Roeder RG, Sabari BR. 2023. Functional partitioning of transcriptional regulators by patterned charge blocks. *Cell* **186**: 327-345.e328.

Marenduzzo D, Faro-Trindade I, Cook PR. 2007. What are the molecular ties that maintain genomic loops? *Trends Genet* **23**: 126-133.

McCord RP, Zhou VW, Yuh T, Bulyk ML. 2011. Distant cis-regulatory elements in human skeletal muscle differentiation. *Genomics* **98**: 401-411.

McSwiggen DT, Mir M, Darzacq X, Tjian R. 2019. Evaluating phase separation in live cells: diagnosis, caveats, and functional consequences. *Genes Dev* **33**: 1619-1634.

Meeussen JVW, Pomp W, Brouwer I, de Jonge WJ, Patel HP, Lenstra TL. 2023. Transcription factor clusters enable target search but do not contribute to target gene activation. *Nucleic Acids Res* **51**: 5449-5468.

Menant A, Baudouin-Cornu P, Peyraud C, Tyers M, Thomas D. 2006. Determinants of the ubiquitin-mediated degradation of the Met4 transcription factor. *J Biol Chem* **281**: 11744-11754.

Mifsud B, Tavares-Cadete F, Young AN, Sugar R, Schoenfelder S, Ferreira L, Wingett SW, Andrews S, Grey W, Ewels PA et al. 2015. Mapping long-range promoter contacts in human cells with high-resolution capture Hi-C. *Nat Genet* **47**: 598-606.

Mir M, Stadler MR, Ortiz SA, Hannon CE, Harrison MM, Darzacq X, Eisen MB. 2018. Dynamic multifactor hubs interact transiently with sites of active transcription in. *Elife* **7**.

Misteli T. 2020. The Self-Organizing Genome: Principles of Genome Architecture and Function. *Cell* **183**: 28-45.

Mitrea DM, Mittasch M, Gomes BF, Klein IA, Murcko MA. 2022. Modulating biomolecular condensates: a novel approach to drug discovery. *Nat Rev Drug Discov* **21**: 841-862.

Nott TJ, Petsalaki E, Farber P, Jervis D, Fussner E, Plochowietz A, Craggs TD, Bazett-Jones DP, Pawson T,
Forman-Kay JD et al. 2015. Phase transition of a disordered nuage protein generates
environmentally responsive membraneless organelles. *Mol Cell* **57**: 936-947.
Okada M, Fujimoto M, Srivastava P, Pandey A, Takii R, Nakai A. 2023. The Mediator subunit MED12
promotes formation of HSF1 condensates on heat shock response element arrays in heat-shocked
cells. *FEBS Lett* **597**: 1702-1717.
Oluwadare O, Highsmith M, Cheng J. 2019. An Overview of Methods for Reconstructing 3-D Chromosome
and Genome Structures from Hi-C Data. *Biol Proced Online* **21**: 7.
Ouni I, Flick K, Kaiser P. 2010. A transcriptional activator is part of an SCF ubiquitin ligase to control
degradation of its cofactors. *Mol Cell* **40**: 954-964.
Papantonis A, Cook PR. 2011. Fixing the model for transcription: the DNA moves, not the polymerase.
Transcription **2**: 41-44.
Peng L, Li EM, Xu LY. 2020. From start to end: Phase separation and transcriptional regulation. *Biochim
Biophys Acta Gene Regul Mech* **1863**: 194641.
Rieder D, Trajanoski Z, McNally JG. 2012. Transcription factories. *Front Genet* **3**: 221.
Rohde JR, Trinh J, Sadowski I. 2000. Multiple signals regulate GAL transcription in yeast. *Mol Cell Biol* **20**:
3880-3886.
Sabari BR, Dall'Agnese A, Boija A, Klein IA, Coffey EL, Shrinivas K, Abraham BJ, Hannett NM, Zamudio AV,
Manteiga JC et al. 2018. Coactivator condensation at super-enhancers links phase separation and
gene control. *Science* **361**.
Sawado T, Halow J, Bender MA, Groudine M. 2003. The beta -globin locus control region (LCR) functions
primarily by enhancing the transition from transcription initiation to elongation. *Genes Dev* **17**:
1009-1018.
Schoenfelder S, Fraser P. 2019. Long-range enhancer-promoter contacts in gene expression control. *Nat
Rev Genet* **20**: 437-455.
Shin Y, Berry J, Pannucci N, Haataja MP, Toettcher JE, Brangwynne CP. 2017. Spatiotemporal Control of
Intracellular Phase Transitions Using Light-Activated optoDroplets. *Cell* **168**: 159-171.e114.
Slot JC, Rokas A. 2010. Multiple GAL pathway gene clusters evolved independently and by different
mechanisms in fungi. *Proc Natl Acad Sci U S A* **107**: 10136-10141.
Sutherland H, Bickmore WA. 2009. Transcription factories: gene expression in unions? *Nat Rev Genet* **10**:
457-466.
Thomas D, Jacquemin I, Surdin-Kerjan Y. 1992. MET4, a leucine zipper protein, and centromere-binding
factor 1 are both required for transcriptional activation of sulfur metabolism in Saccharomyces
cerevisiae. *Mol Cell Biol* **12**: 1719-1727.
Thomas D, Surdin-Kerjan Y. 1997. Metabolism of sulfur amino acids in Saccharomyces cerevisiae.
Microbiol Mol Biol Rev **61**: 503-532.
Trojanowski J, Frank L, Rademacher A, Mücke N, Grigaitis P, Rippe K. 2022. Transcription activation is
enhanced by multivalent interactions independent of phase separation. *Mol Cell* **82**: 1878-
1893.e1810.
Tsai A, Alves MR, Crocker J. 2019. Multi-enhancer transcriptional hubs confer phenotypic robustness. *Elife
***8**.
Uyehara CM, Apostolou E. 2023. 3D enhancer-promoter interactions and multi-connected hubs:
Organizational principles and functional roles. *Cell Rep* **42**: 112068.
Vendruscolo M, Fuxreiter M. 2022. Sequence Determinants of the Aggregation of Proteins Within
Condensates Generated by Liquid-liquid Phase Separation. *J Mol Biol* **434**: 167201.

Whyte WA, Orlando DA, Hnisz D, Abraham BJ, Lin CY, Kagey MH, Rahl PB, Lee TI, Young RA. 2013. Master
transcription factors and mediator establish super-enhancers at key cell identity genes. *Cell* **153**:
307-319.
Xu M, Cook PR. 2008. Similar active genes cluster in specialized transcription factories. *J Cell Biol* **181**: 615-
623.
Xue B, Dunbrack RL, Williams RW, Dunker AK, Uversky VN. 2010. PONDR-FIT: a meta-predictor of
intrinsically disordered amino acids. *Biochim Biophys Acta* **1804**: 996-1010.
Yan C, Chen H, Bai L. 2018. Systematic Study of Nucleosome-Displacing Factors in Budding Yeast. *Mol Cell*
71: 294-305.e294.
Zhang D, Bai L. 2016. Interallelic interaction and gene regulation in budding yeast. *Proc Natl Acad Sci U S*
A **113**: 4428-4433.
Zhang H, Shao S, Zeng Y, Wang X, Qin Y, Ren Q, Xiang S, Wang Y, Xiao J, Sun Y. 2022. Reversible phase
separation of HSF1 is required for an acute transcriptional response during heat shock. *Nat Cell*
Biol **24**: 340-352.
Zhang X, Liu Y, Genereux JC, Nolan C, Singh M, Kelly JW. 2014. Heat-shock response transcriptional
program enables high-yield and high-quality recombinant protein production in Escherichia coli.
ACS Chem Biol **9**: 1945-1949.
Zhao J, Faryabi RB. 2023. Spatial promoter-enhancer hubs in cancer: organization, regulation, and function.
Trends Cancer.
Zhu I, Song W, Ovcharenko I, Landsman D. 2021. A model of active transcription hubs that unifies the roles
of active promoters and enhancers. *Nucleic Acids Res* **49**: 4493-4505.
Zou F, Bai L. 2019. Using time-lapse fluorescence microscopy to study gene regulation. *Methods* **159-160**:
138-145.