Bacterial ecology and evolution converge on seasonal and decadal scales

Robin R. Rohwer[1*], Mark Kirkpatrick[1], Sarahi L. Garcia[2,3], Matthew Kellom[4], Katherine D. McMahon[5,6†], Brett J. Baker[1,7‡]

[1]The University of Texas at Austin, Department of Integrative Biology, 2415 Speedway #C0930, Austin, TX 78712, USA
[2]Carl von Ossietzky Universität Oldenburg, Institute for Chemistry and Biology of the Marine Environment (ICBM), 26129 Oldenburg, Germany
[3]Stockholm University, Department of Ecology, Environment, and Plant Sciences, Science for Life Laboratory, 106 91 Stockholm, Sweden
[4]DOE Joint Genome Institute, Lawrence Berkeley National Laboratory, 1 Cyclotron Road, Berkeley, CA 94720, USA
[5]University of Wisconsin-Madison, Department of Bacteriology, 1550 Linden Drive, Madison, WI, 53706, USA
[6]University of Wisconsin-Madison, Department of Civil and Environmental Engineering, 1550 Linden Drive, Madison, WI, 53706, USA
[7]The University of Texas at Austin, Department of Marine Science, Port Aransas, Austin, TX 78373, USA
Correspondence: *Robin.Rohwer@gmail.com, †trina.mcmahon@wisc.edu, and ‡brett_baker@utexas.edu

Ecology and evolution are distinct theories, but the short lifespans and large population sizes of microbes allow evolution to unfold along contemporary ecological time scales. To document this in a natural system, we collected a two-decade, 471-metagenome time series from a single site in a freshwater lake, which we refer to as the TYMEFLIES dataset. This massive sampling and sequencing effort resulted in the reconstruction of 30,389 metagenomic-assembled genomes (MAGs) over 50% complete, which dereplicated into 2,855 distinct genomes (>96% nucleotide sequence identity). We found both ecological and evolutionary processes occurred at seasonal time scales. There were recurring annual patterns at the species level in abundances, nucleotide diversities (π), and single nucleotide variant (SNV) profiles for the majority of all taxa. During annual blooms, we observed both higher and lower nucleotide diversity, indicating that both ecological differentiation and competition drove evolutionary dynamics. Overlayed upon seasonal patterns, we observed long-term change in 20% of the species' SNV profiles including gradual changes, step changes, and disturbances followed by resilience. Most abrupt changes occurred in a single species, suggesting evolutionary drivers are highly specific. Nevertheless, seven members of the abundant *Nanopelagicaceae* family experienced abrupt change in 2012, an unusually hot and dry year. This shift coincided with increased numbers of genes under selection involved in amino acid and nucleic acid metabolism, suggesting fundamental organic nitrogen compounds drive strain differentiation in the most globally abundant freshwater family. Overall, we observed seasonal and decadal trends in both interspecific ecological and intraspecific evolutionary processes. The convergence of microbial ecology and evolution on the same time scales demonstrates that understanding microbiomes requires a new unified approach that views ecology and evolution as a single continuum.

Main Text

Ecology and evolution are typically considered distinct processes that unfold on different time

scales. Although microbial ecology has blossomed into a rich field spanning experimental and

natural systems, our understanding of microbial evolution relies primarily on phylogenetic

reconstructions and laboratory experiments. Phylogenies provide insight into ancient evolutionary

events[1], but lack insight into contemporary evolution[2] that occurs on directly observable time

scales. Laboratory approaches, such as the *E. coli* long-term evolution experiment[3], are able to

directly observe evolutionary processes, but lack the context of complex communities typical of

natural environments.

Here we describe a two-decade microbial time series from a freshwater lake, which allows us to

directly observe the interplay between ecology and contemporary evolution in a natural ecosystem.

We reconstructed tens of thousands of MAGs from 471 metagenomes spanning two decades with

a sampling frequency averaging twice a month. This dataset constitutes the largest and longest

metagenome dataset from a single location. We find that ecology and evolution both unfold at

short, seasonal time scales as well as longer-term decadal time scales. The similar dynamics of

inter- and intraspecies processes suggests that bacterial ecology and evolution unfold along a

continuum that lacks a clear delineation between the two theories. We conclude that microbial

ecology and evolution must be considered simultaneously and combined into a unified theory.

The TYMEFLIES dataset

We collected 471 samples over 20 years from Lake Mendota (WI, USA)[4] and obtained shotgun DNA libraries (Fig. 1a, Supplementary Table 1). We refer to these Twenty Years of Metagenomes Exploring Freshwater Lake Interannual Eco/evo Shifts as the TYMEFLIES dataset. By cross-mapping reads from ~50 metagenomes to each single-sample metagenome assembly, we obtained a total of 85,684 genome bins, 30,389 of which were medium or high quality (> 50% completeness and < 10% contamination)[5]. We clustered these 30,389 bins at 96% average nucleotide identity (ANI) and obtained 2,855 clusters from which we chose representative metagenome-assembled genomes (MAGs)[6] (Supplementary Table 2). Several previous studies have found an emergent species boundary at similar ANI cutoffs[7–9], and we observed a rapid increase in the number of clusters above the 96% ANI cutoff. For the purposes of this study, we treat these 96% ANI representative MAGs as bacterial species and refer to sub-species delineations as strains.

The representative MAGs have high estimated completeness (median 86%) and low contamination (median 0.9%) (Fig. 1b), and reflect the lake's abundant bacterial community, especially in well-sampled seasons (Fig. 1c). Using a 16S rRNA gene amplicon dataset from the same timeseries[4] as a reference for the expected community composition (Fig. 1d), we found that our representative MAGs comprise most of the abundant taxa (Fig. 1e). Moreover, we obtained 168 representative MAGs from the *Nanopelagicales* order, which is the most abundant order in Lake Mendota and accounts for 22% of the amplicon reads and 10% of the mapped metagenomic reads. Similar to SAR11 bacteria in the oceans, this freshwater lineage is abundant in lakes globally[10], difficult to culture[11], and typically has highly streamlined genomes[12].

Fig. 1. The TYMEFLIES dataset. a, Metagenomes sample dates are indicated by black vertical lines. Microbial seasons[13] are indicated by colored shading. **b**, Quality of the 2,855 reference genomes obtained after clustering to 96% ANI. We treat these reference genomes as species. **c**, Percent of metagenome reads from each sample that mapped to all reference genomes with an ANI $\geq$ 93%. Samples are grouped by season to highlight how well the reference genomes reflect each seasonal community. **d**, Rank abundance of phyla as measured by 16S rRNA gene amplicon sequencing[4]. The abundant *Nanopelagicales* order of Actinobacteria is highlighted. **e**, Abundance of phyla in the TYMEFLIES reference genomes, quantified as the mean relative abundance normalized by genome size and sequencing depth. The *Nanopelagicales* order is again highlighted.

Seasonal ecology and evolution

100 Lake Mendota has been the focus of limnological research since the late 1800s and has been

101 routinely sampled since 1984 by the North Temperate Lakes Long-Term Ecological Research

102 program (NTL-LTER)[14]. Microbial sampling began in 2000 as part of an NSF microbial

Fig. 2. Bacterial seasonality at the sub-species level. a, The percent of species with seasonality in nucleotide diversity and/or centered log ratio-transformed relative abundance. The 1,474 reference species that occurred at least 30 times were included in this analysis. **b**, A time decay plot of the Euclidean distances between the SNV profiles of an abundant species in the *Nanopelagicus* genus (ME2017-06-13_3300043469_group7_bin14). A more similar SNV profile indicates that the strain composition is more similar. Each blue point represents a pairwise comparison between two sample dates, with the time between those dates on the x-axis. The black line is a 6-month moving average, drawn to highlight the annual periodicity of strain similarities. **c**, An example of a "less diverse" bloom, where nucleotide diversity decreases as relative abundance increases. Displayed is an abundant species in the *Planktophila* genus (ME2011-09-04_3300044729_group3_bin142). **d**, An example of a "more diverse" bloom, where nucleotide diversity increases as abundance increases. Displayed is an abundant species in the *Nanopelagicaceae* family, MAG-120802 genus (ME2012-08-31_3300044613_group4_bin150). **e**, The distribution of bloom diversity patterns across the 365 species that had both seasonal abundance and seasonal nucleotide diversity.

observatory[15]. From this long history of research, we know the lake follows a consistent annual

phenology, and that phenological patterns are changing in response to climate change and invasive

species[16–19]. Rohwer *et. al*[13] found that these phenological dynamics extend to the bacterial

community. To confirm that phenological abundance patterns also exist in our more finely resolved

bacterial species, we identified annual peaks in species relative abundance using periodograms

(magnitude of Fourier transforms). After limiting this temporal analysis to the subset of

representative MAGs that occurred at least 30 times over at least 10 years, we found that 72% of

these 1,474 bacterial species have consistent seasonal abundance patterns (Fig. 2a).

To determine whether evolutionary dynamics (*i.e.* changes in allele frequency within the

population) also unfold seasonally, we mapped reads from each sample against each species'

reference genome to identify shifts in strain composition evidenced by shifts in nucleotide diversity

(π) and the profiles of single nucleotide variants (SNVs). We found that 33% of the same 1,474

species displayed consistent seasonal nucleotide diversity patterns (Fig. 2a). For a more granular

view of strain composition, we created SNV profiles for each species where each date's profile

consists of the percent reference base at each SNV position. After making our abundance cutoff

more stringent by requiring a median coverage > 10x, we calculated the Euclidean distance

between each sample's SNV profile for the most abundant 263 species. We found that 80% of these

species had consistent phenological patterns in their strain composition (Fig. 2b). This

demonstrates that phenological patterns evident in the bacterial community extend to the finest

possible taxonomic resolution. Several short-term freshwater studies have also observed changes

in strain composition on seasonal timescales[20,21]. Phenological patterns in sub-species strains

suggest that bacterial evolution, evidenced by intraspecific genomic change, also unfolds on

seasonal timescales.

Given the ubiquity of seasonal patterns in both species abundance and sub-species diversity, we

asked if they were correlated. We quantified whether a species' "bloom" in abundance consisted

of fewer strains or more strains than its baseline composition. Of the 365 species with seasonal

patterns in both abundance and nucleotide diversity (purple bars in Fig. 2a), we found both

scenarios were common; 21% of these species had less diverse blooms (Fig. 2c and yellow bars in

Fig. 2e), while 19% had more diverse blooms (Fig. 2d and green bars in Fig. 2e). Further, all

abundant phyla demonstrated an even mix of both bloom types (Fig. 2e). A lower-diversity bloom

suggests that a subset of strains outcompeted the others, while a higher-diversity bloom suggests

that micro-niches allowed rarer strains to gain abundance, resulting in higher strain diversity due

to a more even strain composition. This is in agreement with a previous study that found both

overlapping and distinct niches within freshwater bacterial species[22]. The prevalence of both

bloom diversity patterns highlights the lack of a clear boundary between ecological and

evolutionary processes since identical intraspecific and interspecific processes are occurring

simultaneously.

Long-term ecology and evolution

Long-term changes can be masked by seasonal oscillations, lost in what is referred to as the

"invisible present"[23]. The unprecedented length of the TYMEFLIES metagenome dataset provides

a unique lens into the invisible present, enabling the identification of overlayed long-term patterns.

To find long-term changes in SNV-based strain profiles, we developed a classifier trained on the

distance between each date's SNV profile and the SNV profile of that species' first occurrence in

the timeseries. We trained this classifier on 11 examples of manually identified temporal patterns,

and then applied it to all 263 most abundant species. Our classifier identified gradual change (Fig.

3a), which may arise from genetic drift or in response to a slow press disturbance, as well as abrupt

change (Fig 3b and c), which may arise if a disturbance reaches a tipping point threshold leading

to a new stable state, or from a sudden environmental shift[24,25]. Among instances of abrupt change,

we identified step changes (Fig. 3b), where the new strain composition persisted during the

remainder of our time frame, as well as patterns of disturbance with resilience (Fig. 3c), where the

strain composition recovered to baseline.

We found that 21% of the most abundant species experienced one kind of long-term change in

their strain profiles during our 20-year study period, and these changes overlayed both seasonal

and acyclical short-term dynamics (Fig. 3d). Abrupt change was almost twice as common as

gradual change (36 vs. 19 species), and resilience was only slightly more common than a lasting

step change (20 vs. 16 species) (Fig. 3d). The three long-term change patterns were found in many

abundant species distributed across phyla (Fig. 3e). Many abundant Actinobacteriota species

experienced long-term change. These long-term changes in SNV profiles reflect shifts in

intraspecific strain composition, which is typically attributed to evolutionary processes. The fact

that during our observation period over a fifth of the species experienced long-term changes in

their strain profiles emphasizes the importance of including contemporary evolutionary change in

our understanding of microbial ecology.

Fig. 3. Long-term changes in strain composition. a, An example of long-term, gradual change in strain composition. Points indicate sample dates, and distance refers to the Euclidean distance between a species' SNV profile on that sample date and that species' first available SNV profile. A species in the *Nanopelagicales* order, AcAMD-5 family is shown (ME2005-06-22_3300042363_group2_bin84). **b**, An example of an abrupt step change in strain composition in a species in the *Nanopelagicus* genus (ME2011-09-21_3300043464_group3_bin69). **c**, An example of a disturbance/resilience pattern, where an abrupt change in strain composition is followed by recovery to the original strain composition, in a species in the *Planktophila* genus (ME2015-07-03_3300042555_group6_bin161). **d**, Long-term change patterns often overlayed seasonal patterns. Of the 263 species abundant enough to observe their strain profiles, 39 had both long-term and seasonal patterns while 16 had only long-term patterns. **e**, The distribution of long-term patterns across phyla. Each species that underwent long-term change is indicated by a section of the phyla's bar, scaled by the mean abundance of that species. The sections corresponding to the examples highlighted in **a-c** are labelled.

Abrupt changes in *Nanopelagicaceae*

In general, related species did not change in unison with each other, suggesting that the drivers of evolutionary change are highly specific (Fig. 4a). One exception is an abrupt change event that impacted seven species in the *Nanopelagicaceae* (acI) family in 2012, specifically species in the *Nanopelagicus* and *Planktophila* genera (acI-B and acI-A). This is the most abundant family in Lake Mendota and in freshwaters globally[10], and the 127 *Nanopelagicaceae* reference species we recovered accounted for 8% relative abundance on average. Five of these *Nanopelagicaceae* species displayed resilience to the abrupt change, while two experienced lasting step changes in strain composition.

A myriad of possible environmental variables could have driven this event. A leading candidate is climate, which was unusually warm and dry in 2012. The lake experienced its highest epilimnion water temperatures since 1894[13] (Fig. 4b), the fifth shortest winter ice duration since 1856[26] (Fig. 4c), the 8th lowest annual discharge from its major tributary since 1976 and the second lowest peak discharge[27] (Fig. 4e). These climatic conditions led to top-down and bottom-up controls on the lake's primary productivity: the highest spring zooplankton abundance since measurements began in 1994[28] (Fig. 4d) was likely a result of the mild winter and spring allowing zooplankton populations, especially the prolific grazer *Daphnia pulicaria* to establish early, and low total phosphorus and soluble reactive phosphorus (Fig. 4f-g) was likely a result of low sediment transport associated with mild discharge events[29]. The resulting combination of high zooplankton grazing and low phosphorus, typically the limiting nutrient in lakes, may be responsible in turn for low phytoplankton biomass (Fig. 4h). Lake Mendota's dissolved organic

225   carbon (DOC) is primarily provided by phytoplankton[30], and consequently DOC was also low in

226   2012 (Fig. 4i).

Fig. 4. Abrupt changes in *Nanopelagicaceae* strain composition coincide with environmental extremes in 2012. a, Dates of all abrupt changes in strain composition arranged by phyla. Most changes were isolated events, but multiple species from two abundant genera of Actinobacteriota, *Planktophila* and *Nanopelagicus*, experienced abrupt change in 2012. Point size is scaled by species abundance. **b**, Unusually high epilimnion water temperatures during spring and summer 2012 (relative to 1894 – 2019). **c**, The preceding winter had an unusually short ice duration (relative to 1853 – 2023). **d**, Total zooplankton biomass (excluding predatory *Bythotrophes* and *Leptodora*) was unusually high, likely enabled by warm early spring temperatures (relative to 1995 – 2018). **e**, Discharge from the Yahara River, the main tributary to Lake Mendota, was unusually low and lacked high runoff events typical after storms and spring snowmelt (relative to 1989 - 2021). **f**, Sediment transport of total phosphorus, and **g**, of soluble reactive phosphorus were low (relative to 1995 – 2021). **h**, Low phytoplankton biomass, likely resulting from both high zooplankton grazing and low nutrient availability. **i**, Low dissolved organic carbon (relative to 1996 – 2022), likely a result of low phytoplankton abundance[30].

Another possible driver is the irruption of the invasive zooplankton spiny water flea (*Bythorephes*

cedertrömii) in 2009. This major disturbance resulted in a trophic cascade that decreased water

clarity[19,31], increased lake anoxia[28], and shifted the bacterial community composition[13]. Although

the abrupt changes in strain composition of seven *Nanopelagicaceae* species were observed three

years later, lag effects are common in complex ecosystems[32]. Ecosystem-wide drivers like the 2012

climate anomalies and the 2009 species invasion can have cascading and interacting effects on

nutrient and carbon dynamics, which in turn impact the bacterial community. The observed long-

term intraspecific changes suggest that such ecological drivers are also drivers of evolutionary

change, further emphasizing how ecology and evolution are intertwined.

Evolutionary signals in a *Nanopelagicus*

To understand the dynamics of abrupt evolutionary change, we further examined one of the

abundant species, a *Nanopelagicus* (acI-B), that experienced a step change in strain composition

in August 2012 (Fig. 3b). An NMDS ordination of its SNV profiles indicated the strain composition

changed abruptly at that time and settled into a new composition after a period of adjustment in

2012 and 2013 (Fig. 5a).

The relative abundance of this species was quite constant throughout our 20-year observation

period (Fig. 5b), typically with higher abundances during the spring clearwater phase. The step

change in strain composition (Fig. 3b) coincided with one in nucleotide diversity (Fig. 5c). These

patterns could result from the spread of a new strain or with an increase in the evenness of

abundance in existing strains. To distinguish between these hypotheses, we counted the number

of previously unobserved SNVs in the mapped reads of every sample. We did not see large spikes

in new SNVs in 2012 (Fig. 5d), suggesting that the step changes reflected a shift in the relative

abundances of existing strains.

Fig. 5. Step change in strain composition coincides with more genes under selection. a, An abundant *Nanopelagicus* species experienced a step change in strain composition in 2012 (ME2011-09-21_3300043464_group3_bin69, see also Fig. 3b). Samples with more similar SNV profiles appear closer on this NMDS plot. Years 2000-2011 cluster together and are distinct from years 2014-2019, which cluster separately. A sudden change in strain composition occurred on August 3, 2012. **b**, Despite the abrupt change in strain composition, the relative abundance this species remained constant over time. **c**, Concurrent with the shift in strain composition, nucleotide diversity increased and then remained high, indicating that the new equilibrium was comprised of a more diverse assemblage of strains. **d**, The absence of a spike in the number of new SNVs suggests that an increase in the evenness of existing strains occurred, rather than the introduction of new strains. **e**, Concurrent with the shift in strain composition, the number of genes under positive selection also increased (McDonald-Kreitman F-statistic p-value < 0.05). **f**, Occurrence of consistently selected genes in all the samples, in the pre-2012 period, and in the post-2012 period. X-axis indicates samples over time and Y-axis indicates genes. Shading indicates the significance level of positive selection. Amino acid-related genes and nucleic acid-related genes are indicated on the right axis. Full annotations are available in Supplementary Table 3. Note that the X-axis is evenly spaced by sample, so that years with more samples take up more space.

This interpretation is consistent with a dramatic increase in the number of genes under positive

selection that occurred at this time (Fig. 5e). As the relative abundances of some strains increase,

alleles specific to them appear to undergo partial (or "soft") selective sweeps. As the strain

composition reequilibrates, this signal dies out (Fig 5e). To identify candidate loci that may be

important to adaptation that occurred during our sampling, we sought genes that consistently

showed signals of selection over the entire timeseries, only during the pre-2012 period, and only

during the post-2012 period. Four genes were consistently selected both pre- and post-2012, four

genes were consistently selected pre-2012, and 33 genes were consistently selected post-2012. We

used gene functional predictions[33] to identify their potential metabolic pathways. Of the 33

consistently selected genes post-2012, ten are involved in amino acid metabolism or

aminoacylation, and seven are involved with nucleic acid synthesis or degradation (Fig. 5f).

Previously, the absence of biosynthesis or auxotrophies for amino acids and nucleotides has been

highlighted for microorganisms with streamlined genomes[34,35]. In the streamlined *Nanopelagicus*,

auxotrophies for various amino acids[12,36] coupled with an enrichment of transporters for many

small organic nitrogen compounds, including amino acids[12,37,38] and nucleic acid components[12,36–

38] are common. Moreover, the histidine pathway was found split between two different strains of

Nanopelagicus growing in a mixed culture[36]. Our observation of consistent selection on amino

acid and nucleic acid metabolism suggests that these genes differentiate the post-2012 strains, thus

biosynthesis, use and reuse of small organic nitrogen compounds are key in the ecology and

evolution of these globally abundant lake bacteria.

A continuum of ecology and evolution

The interface between ecology and evolution is delineated by species boundaries, but in bacteria species concepts and definitions are hotly debated[39]. Using a commonly chosen definition for microbial species boundaries, we found processes that unfold as interspecies ecological dynamics also occur in intraspecies evolution. Moreover, the time scales of these processes overlapped, as did likely environmental drivers. How microbes will respond to global changes in land use, invasive species, and climate[40] are pressing questions that require an understanding of long-term change. Few microbial timeseries are long enough to capture such dynamics, but decadal observations are an essential approach to understand how complex ecosystems respond to global change. The two-decade TYMEFLIES dataset will serve as an invaluable community resource to continue addressing these questions and move us toward a unified approach to microbial ecology and evolution. Microbial ecology and evolution must be considered simultaneously and combined into a unified theory.

Methods

Data Availability

Metagenome sequences are available through the DOE Joint Genome Institute's Genome Portal under Proposal 504350 (https://genome.jgi.doe.gov/portal/Exttemetagenomes/Exttemetagenomes.info.html), and through the NCBI Sequence Read Archive at accessions listed in Supplementary Table 1. Reference MAG sequences are available through the NCBI GenBank IDs listed in Supplementary Table 2. Environmental data is publicly available through the Environmental Data Initiative

333 (https://edirepository.org/)[41–49] and the U.S. Geological Survey's Water Data for the Nation

334 (https://waterdata.usgs.gov/nwis)[27].

Code Availability

337 Custom scripts used for data processing are available at https://github.com/rrohwer/TYMEFLIES

338 and on Zenodo.

Lake Mendota Samples

Lake Mendota is a eutrophic temperate lake located in Madison, Wisconsin (USA). Integrated

epilimnion samples were collected from the upper 12 m at a 25 m deep location referred to as the

central "deep hole" (43°05'58.2"N 89°24'16.2"W). Bacteria were collected on 0.2 μm

polyethersulfone filters (Pall Corporation), stored at -80°C, and DNA was extracted after

randomizing sample order by a single person in 2018-2019 using FastDNA Spin Kits (MP

Biomedicals). A detailed description of the study site, sample collection, and DNA extraction

procedures is provided by Rohwer and McMahon[4].

Metagenome sequencing and assembly

Samples were sequenced by the US Department of Energy Joint Genome Institute (JGI) using a

NovaSeq 6000 with an S4 flow cell. Sample metadata is available in Supplementary Table 1, and

raw sequencing data is available through the JGI Genome Portal under Proposal 504350

(https://genome.jgi.doe.gov/portal/Exttemetagenomes/Exttemetagenomes.info.html), or from the

NCBI Sequence Read Archive under accession numbers listed in Supplementary Table 1. Read

filtering was performed using standard JGI protocols[50], which are additionally detailed as metadata

paired with each sample through the JGI IMG/M website. Briefly, BBDuk[51] was used to remove

adapters and quality trim reads, and BBMap[51] was used to identify and remove common

contaminants. In our analyses we treated the resulting filtered fastq files as the metagenome reads.

Single-sample assemblies were also generated by JGI with their standard protocol[50] using

metaSPAdes[52]. These filtered fastq files and paired single-sample assemblies are available through

the JGI Genome Portal under Proposal 504350.

Obtaining and characterizing genomes

Genomes were binned out of metagenomes using the Texas Advanced Computing Center's

Lonestar6 supercomputer. Metagenomic reads were mapped back to sample assemblies using

BBMap (version 38.22)[51], sorted BAM files were created using SAMtools (version 1.9)[53], and

metagenome-assembled genomes were binned using MetaBAT2 (version 2.12.1)[54]. Metagenomic

reads from different samples were cross-mapped back to each assembly. Cross-mapping scales

exponentially, so it was performed on assemblies and sample reads broken into approximately 50-

sample groups of consecutive sample dates, with samples from the same year grouped together.

This resulted in 85,684 genome bins. CheckM2 (version 0.1.3)[5] was used to asses bin quality,

including completeness and contamination estimates, and GTDB-tk (version 2.1.1)[55] was used to

assign GTDB taxonomy (release 207)[56] to all bins. 30,389 genome bins were at least 50% complete

and less than 10% contaminated, and these bins were de-replicated to 96% ANI using dRep

(version 3.4.0)[6]. To choose 96% as our ANI cutoff, we ran dRep at ANIs ranging from 90 to 99%

and examined the resulting number of de-replicated bins, as well as the number of bins from the

same assembly that were combined. We chose 96% ANI because very few (one) of the 30,389 bins

were combined into an ANI group with a bin created from the same assembly, and because 96%

ANI was generally located right before a sudden increase in the total number of genome groups.

Our goal was to separate as many species as possible, while combining strains that were so closely

related they would compete for mapped reads. Applying a 96% ANI cutoff with dRep resulted in

2,855 representative genomes, which we treated as species in this study.

To quantify the relative abundance of each species in every sample, we mapped all sample reads

against the concatenated 96% ANI reference genomes using bbmap (version 38.22)[51], created

sorted BAM files using SAMtools (version 1.9)[53], and calculated relative abundance using coverM

(version 0.6.1)[57]. With the coverM software, we required a minimum read percent identity of 93,

proper pairs only, and excluded 1000 bp from each contig end from the calculation. CoverM

calculates relative abundance as the mean coverage divided by the mean coverage across all

genomes multiplied by the proportion of reads that mapped to the genome, thus normalizing by

genome size to estimate the fraction of cells that belong to a given species in each sample. A table

of representative MAGs along with taxonomy annotations, quality statistics, and abundance

statistics is available as Supplementary Table 2.

To further characterize the genomes, we ran inStrain (version 1.7.1)[58] using a minimum read ANI

of 93%, as recommended by the inStrain documentation given our previous choice of 96% ANI to

dereplicate genomes. This software called SNVs and calculated nucleotide diversity, among other

metrics. To identify genes we ran prodigal (version 2.6.3)[59] on each genome separately. We then

used Kofamscan (version 1.3.0)[60] to assign gene annotations from the Kyoto Encyclopedia of

Genes and Genomes (KEGG) database (release 107.1)[33]. Additional custom analyses were

performed using the R programming language (version 4.1.2)[61], and relied extensively on the

data.table R package (version 1.14.8)[62], the lubridate R package (version 1.9.3)[63], and GNU

parallel (version 'Chandrayaan')[64].

Classifying seasonal and long-term change

To classify each species' abundance pattern as seasonal or not, we started with relative abundances

as calculated by coverM (version 0.6.1)[57] and further corrected any abundance to zero if the

genome's coverage breadth was 70% or less than its expected breadth, as calculated by inStrain

(version 1.7.1)[58]. We then applied a centered log ratio transformation to the relative abundance

values using the compositions R package (version 2.0-6)[65]. After taking a daily linear interpolation

to obtain evenly spaced samples, we detrended the temporal profiles with a cubic fit. Finally, we

performed a periodogram analysis by computing the magnitude of the fast Fourier transform. If a

peak occurred within 30 days of 365 days we considered it an annual oscillation, and if any of the

top five peaks corresponded to an annual period, we classified the species as having a seasonal

abundance pattern. We applied this analysis only to the 1,474 species that occurred on least 30

dates over at least 10 years. To classify each species' nucleotide diversity pattern as seasonal or

not, we similarly performed a fast Fourier transform on its inStrain-calculated nucleotide diversity

over time. We used the same periodogram analysis to classify it as having seasonal nucleotide

diversity or not, and we applied this analysis to the same subset of 1,474 species.

To characterize blooms as more diverse or less diverse, we calculated the Pearson correlation

between centered log ratio-transformed relative abundance and nucleotide diversity for the 365

species that had both seasonal abundance and seasonal nucleotide diversity annual oscillations. We

considered it a positive correlation (more diverse blooms) if the Pearson correlation was at least

0.35 and a negative correlation (less diverse blooms) if the Pearson correlation was less than or

equal to -0.35. We repeated this analysis with up to two weeks of lag and used the highest

correlation within that window. We chose 0.35 as a reasonable cutoff after manual examination of

the first 150 species' correlations.

To calculate SNV profiles for each species, we created vectors corresponding to every SNV

position in its genome, where the value of each element was the percent of mapped reads that

matched the reference genome base at that position in each sample. SNV's were called using

inStrain[58], and we only applied this analysis to samples where the species' median coverage was

over 10x, as at coverages less than that we observed a drop in the total SNVs called. Therefore,

for both long-term and seasonal analysis of SNV profiles, we included only species that had

medium coverage over 10x on at least 30 dates over at least 10 years, which resulted in a subset

of 263 species. To identify changes in SNV profiles, we created a distance matrix for each species

based on Euclidean distances between each sample's SNV profile using the vegan R package

(version 2.6-4)[66]. From this we created a table of time elapsed and Euclidean distance between

each sample date.

To identify seasonal patterns in each species' SNV profiles, we created a daily linear interpolation

of pairwise distances between all samples, taking the mean when multiple sample pairs occurred

with the same time interval. After detrending with a cubic fit, we performed a periodogram analysis

to identify annual oscillations and the presence of seasonal patterns using the same criteria as with

our abundance and nucleotide diversity annual oscillation analysis.

To identify long-term change patterns, we subset our pairwise distance table to the distance of each

sample from the first sample. We developed a classifier for these temporal profiles of distances

between SNV profiles using 11 manually chosen species. Our classifier criteria was hierarchical:

first gradual change was identified, then step change was identified, and finally

disturbance/resilience patterns were identified. After training, the classifier was applied to all 263

species above the abundance cutoff. Gradual change was identified if a linear fit to the daily

linearly interpolated distances, excluding dates closer than a month to the starting date, resulted in

an adjusted R^2 of at least 0.55. Dates closer than a month to the starting date were excluded because

they tended to be highly similar, and a linear interpolation was applied to account for uneven

sampling dates, particularly the high frequency of summer sampling in the latter decade of the

timeseries. Possible step change locations were identified after excluding dates closer than a month

to the starting date and applying an F test to the linearly interpolated distances using the

strucchange R package (version 1.5-3)[67]. If a breakpoint was identified by the F test, the means of

measured (as opposed to interpolated) before and after distances were different (Mann-Whitnes p-

value < 0.01), and the step resulted in a new mean at least 33% higher than the previous mean, a

step change pattern was identified. Disturbance/resilience patterns were then identified using

outlier distances calculated by the default boxplot statistics in R. If a date's distance was > 1.5

times the difference between the 3rd and 1st quartile of observed distances a date was considered

an outlier, and if outlier values were maintained for at least a month the species was classified as

having a disturbance event with resilience.

Analyzing abrupt change in *Nanopelagicaceae*

To place environmental conditions in 2012 in context, historical environmental data was collected

from the North Temperate Lakes Long-Term Ecological Research program (NTL-LTER) through

the Environmental Data Initiative (EDI) interface (https://edirepository.org/) and the US

Geological Survey (USGS) Water Data for the Nation (https://waterdata.usgs.gov/nwis) using the

USGS dataRetrieval R package (version 2.7.14)[68]. EDI datasets analyzed included ice duration[26];

nutrients, pH, and carbon[41]; major ions[42]; water temperatures combined from multiple datasets[45–

49] as described in Rohwer *et al.*[13]; phytoplankton[43]; and zooplankton[44] converted to biomass as

described in Rohwer, Ladwig, *et al.*[28]. River discharge measurements were obtained from the

USGS for the Yahara River, the primary tributary into Lake Mendota (site ID: 05427718)[27]. After

exploring all parameters included in these datasets, the occurrence of a hot, dry year with low

primary productivity became apparent.

Relative abundance and nucleotide diversity of the *Nanopelagicus* MAG ME2011-09-

21_3300043464_group3_bin69 were calculated as for the seasonal analysis. New SNVs were

identified as SNV positions that were called by inStrain[58] for the first time in a given sample.

Genes under selection were identified using dN/dS and pN/pS ratios as calculated by inStrain[58]. A

McDonald-Kreitman test[69] was used to identify positively selected genes where the bias of unfixed

SNVs to be nonsynonymous was lower than the bias of fixed SNVs to be nonsynonymous

(pNpS/dNdS < 1), and positive selection was considered statistically significant when the Fisher

p-value was less than or equal to 0.05. A gene was considered consistently selected if it appeared

under significant positive selection with high frequency (in the 4th quartile). Consistently selected

genes were identified for the pre-2012 and post-2012 time periods separately.

Gene annotations were analyzed in the context of the KEGG pathways[33] they belonged to. For

each potential pathway, all genes present in the genome were visualized with KEGG Pathway

Maps (https://www.genome.jp/brite/br08901). When multiple genes that surrounded the selected

gene existed in the genome, that pathway was considered a likely annotation. When likely

pathways involved amino acid metabolism or aminoacylation, they were considered amino acid-

related. When likely pathways involved purine or pyrimidine metabolism, they were considered

nucleic acid-related.

References

1. Hug, L. A. *et al.* A new view of the tree of life. *Nature Microbiology* **1**, (2016).

2. Brennan, G. L. & Logares, R. Tracking contemporary microbial evolution in a changing ocean.

Trends in Microbiology **31**, 336–345 (2023).

3. Lenski, R. E. Experimental evolution and the dynamics of adaptation and genome evolution in

microbial populations. *ISME J* **11**, 2181–2194 (2017).

4. Rohwer, R. R. & McMahon, K. D. Lake iTag measurements over nineteen years, introducing

the limony dataset. 2022.08.04.502869 Preprint at https://doi.org/10.1101/2022.08.04.502869

(2022).

5. Chklovski, A., Parks, D. H., Woodcroft, B. J. & Tyson, G. W. CheckM2: a rapid, scalable and

accurate tool for assessing microbial genome quality using machine learning.

2022.07.11.499243 Preprint at https://doi.org/10.1101/2022.07.11.499243 (2022).

6. Olm, M. R., Brown, C. T., Brooks, B. & Banfield, J. F. dRep: a tool for fast and accurate

genomic comparisons that enables improved genome recovery from metagenomes through de-

replication. *ISME J* **11**, 2864–2868 (2017).

7. Varghese, N. J. *et al.* Microbial species delineation using whole genome sequences. *Nucleic*

Acids Research **43**, 6761–6771 (2015).

8. Jain, C., Rodriguez-R, L. M., Phillippy, A. M., Konstantinidis, K. T. & Aluru, S. High

throughput ANI analysis of 90K prokaryotic genomes reveals clear species boundaries. *Nat*

Commun **9**, 1–8 (2018).

9. Olm, M. R. *et al.* Consistent Metagenome-Derived Metrics Verify and Delineate Bacterial

Species Boundaries. *mSystems* **5**, 10.1128/msystems.00731-19 (2020).

10. Lipko, I. A. & Belykh, O. I. Environmental Features of Freshwater Planktonic Actinobacteria.

Contemp. Probl. Ecol. **14**, 158–170 (2021).

11. Kim, S., Kang, I., Seo, J.-H. & Cho, J.-C. Culturing the ubiquitous freshwater actinobacterial

acI lineage by supplying a biochemical 'helper' catalase. *ISME J* **13**, 2252–2263 (2019).

12. Neuenschwander, S. M., Ghai, R., Pernthaler, J. & Salcher, M. M. Microdiversification in

genome-streamlined ubiquitous freshwater Actinobacteria. *ISME J* **12**, 185–198 (2018).

13. Rohwer, R. R., Hale, R. J., Vander Zanden, M. J., Miller, T. R. & McMahon, K. D. Species

invasions shift microbial phenology in a two-decade freshwater time series. *Proceedings of the*

National Academy of Sciences **120**, e2211796120 (2023).

14. Magnuson, J. J., Kratz, T. K. & Benson, B. J. *Long-Term Dynamics of Lakes in the*

Landscape: Long-Term Ecological Research on North Temperate Lakes. (Oxford University

Press, 2006).

15. Yannarell, A. C., Kent, A. D., Lauster, G. H., Kratz, T. K. & Triplett, E. W. Temporal Patterns

in Bacterial Communities in Three Temperate Lakes of Different Trophic Status. *Microb Ecol*

46, 391–405 (2003).

16. Magee, M. R. & Wu, C. H. Effects of changing climate on ice cover in three morphometrically

different lakes. *Hydrol. Process.* **31**, 308–323 (2017).

17. Magee, M. R. & Wu, C. H. Response of water temperatures and stratification to changing

climate in three lakes with different morphometry. *Hydrology and Earth System Sciences* **21**,

6253–6274 (2017).

18. Snortheim, C. A. *et al.* Meteorological drivers of hypolimnetic anoxia in a eutrophic, north

temperate lake. *Ecological Modelling* **343**, 39–53 (2017).

19. Matsuzaki, S.-I. S. *et al.* Climate and food web effects on the spring clear-water phase in two

north-temperate eutrophic lakes. *Limnol. Oceanogr.* **n/a**, 1–17 (2020).

20. Okazaki, Y., Nakano, S., Toyoda, A. & Tamaki, H. Long-Read-Resolved, Ecosystem-Wide

Exploration of Nucleotide and Structural Microdiversity of Lake Bacterioplankton Genomes.

mSystems **7**, e00433-22 (2022).

21. Meziti, A. *et al.* Quantifying the changes in genetic diversity within sequence-discrete

bacterial populations across a spatial and temporal riverine gradient. *The ISME Journal* **13**, 767

(2019).

22. Garcia, S. L. *et al.* Contrasting patterns of genome-level diversity across distinct co-occurring

bacterial populations. *The ISME Journal* 1 (2017) doi:10.1038/s41396-017-0001-0.

23. Magnuson, J. J. Long-Term Ecological Research and the Invisible Present. *BioScience* **40**, 495–501 (1990).

24. Turner, M. G. *et al.* Climate change, ecosystems and abrupt change: science priorities. *Philosophical Transactions of the Royal Society B: Biological Sciences* **375**, 20190105 (2020).

25. Scheffer, M., Carpenter, S., Foley, J. A., Folke, C. & Walker, B. Catastrophic shifts in ecosystems. *Nature* **413**, 591–596 (2001).

26. Magnuson, J. J., Carpenter, S. R. & Stanley, E. H. North Temperate Lakes LTER: Ice Duration - Madison Lakes Area 1853 - current. Environmental Data Initiative https://doi.org/10.6073/PASTA/69B3391E13955392587413ECBFC7C298 (2023).

27. U.S. Geological Survey National Water Information System data available on the World Wide Web. USGS 05427718 Yahara River at Windsor, WI, Daily Discharge 00060. U.S. Geological Survey National Water Information System data available on the World Wide Web (2023).

28. Rohwer, R. R. *et al.* Increased anoxia following species invasion of a eutrophic lake. *Limnology and Oceanography Letters* **n/a**, (2023).

29. Carpenter, S. R., Booth, E. G. & Kucharik, C. J. Extreme precipitation and phosphorus loads from two agricultural watersheds. *Limnology and Oceanography* **63**, 1221–1233 (2018).

30. Berg, S. M., Peterson, B. D., McMahon, K. D. & Remucal, C. K. Spatial and Temporal Variability of Dissolved Organic Matter Molecular Composition in a Stratified Eutrophic Lake. *Journal of Geophysical Research: Biogeosciences* **127**, e2021JG006550 (2022).

31. Walsh, J. R., Carpenter, S. R. & Zanden, M. J. V. Invasive species triggers a massive loss of ecosystem services through a trophic cascade. *PNAS* **113**, 4081–4085 (2016).

32. Rastetter, E. B. *et al.* Time lags: insights from the U.S. Long Term Ecological Research Network. *Ecosphere* **12**, e03431 (2021).

33. Kanehisa, M., Furumichi, M., Sato, Y., Kawashima, M. & Ishiguro-Watanabe, M. KEGG for

taxonomy-based analysis of pathways and genomes. *Nucleic Acids Research* **51**, D587–D592

(2023).

34. Ramoneda, J., Jensen, T. B. N., Price, M. N., Casamayor, E. O. & Fierer, N. Taxonomic and

environmental distribution of bacterial amino acid auxotrophies. *Nat Commun* **14**, 7608 (2023).

35. Castelle, C. J. *et al.* Biosynthetic capacity, metabolic variety and unusual biology in the CPR

and DPANN radiations. *Nat Rev Microbiol* **16**, 629–645 (2018).

36. Garcia, S. L. *et al.* Auxotrophy and intrapopulation complementary in the 'interactome' of a

cultivated freshwater model community. *Mol Ecol* **24**, 4449–4459 (2015).

37. Garcia, S. L. *et al.* Metabolic potential of a single cell belonging to one of the most abundant

lineages in freshwater bacterioplankton. *The ISME Journal* **7**, 137 (2013).

38. Hamilton, J. J. *et al.* Metabolic Network Analysis and Metatranscriptomics Reveal

Auxotrophies and Nutrient Sources of the Cosmopolitan Freshwater Microbial Lineage acI.

MSystems **2**, e00091-17 (2017).

39. Rosselló-Móra, R. & Amann, R. Past and future species definitions for Bacteria and Archaea.

Systematic and Applied Microbiology **38**, 209–216 (2015).

40. Tiedje, J. M. *et al.* Microbes and Climate Change: a Research Prospectus for the Future. *mBio*

e00800-22 (2022) doi:10.1128/mbio.00800-22.

41. Magnuson, J. J., Carpenter, S. R. & Stanley, E. H. North Temperate Lakes LTER: Chemical

Limnology of Primary Study Lakes: Nutrients, pH and Carbon 1981 - current. Environmental

Data Initiative https://doi.org/10.6073/PASTA/325232E6E4CD1CE04025FA5674F7B782

(2023).

42. Magnuson, J. J., Carpenter, S. R. & Stanley, E. H. North Temperate Lakes LTER: Chemical

Limnology of Primary Study Lakes: Major Ions 1981 - current. Environmental Data Initiative

https://doi.org/10.6073/pasta/bb563f16c7338fdb3ddf82057ef43cc6 (2023).

43. Magnuson, J. J., Carpenter, S. R. & Stanley, E. H. North Temperate Lakes LTER:

Phytoplankton - Madison Lakes Area 1995 - current. Environmental Data Initiative

https://doi.org/10.6073/PASTA/43D3D401AF88CC05C6595962BDB1AB5C (2022).

44. Magnuson, J., Carpenter, S. & Stanley, E. North Temperate Lakes LTER: Zooplankton -

Madison Lakes Area 1997 - current. Environmental Data Initiative

https://doi.org/10.6073/PASTA/D5ABE9009D7F6AA87D1FCF49C8C7F8C8 (2022).

45. Rohwer, R. R. & McMahon, K. D. Lake Mendota Microbial Observatory Temperature,

Dissolved Oxygen, pH, and conductivity data, 2006-present. Environmental Data Initiative

https://doi.org/10.6073/PASTA/7E533C197ED8EBD27777A89A2C8D7DFE (2022).

46. Magnuson, J. J., Carpenter, S. R. & Stanley, E. H. North Temperate Lakes LTER: Physical

Limnology of Primary Study Lakes 1981 - current. Environmental Data Initiative

https://doi.org/10.6073/PASTA/316203040EA1B8ECE89673985AB431B7 (2021).

47. Magnuson, J., Carpenter, S. & Stanley, E. North Temperate Lakes LTER: High Frequency

Water Temperature Data - Lake Mendota Buoy 2006 - current. Environmental Data Initiative

https://doi.org/10.6073/PASTA/8CEFF296AD68FA8DA6787076E0A5D992 (2020).

48. Robertson, D. Lake Mendota water temperature secchi depth snow depth ice thickness and

meterological conditions 1894 - 2007. Environmental Data Initiative

https://doi.org/10.6073/PASTA/F20F9A644BD12E4B80CB288F1812C935 (2016).

49. Magnuson, J. J., Carpenter, S. R. & Stanley, E. H. Lake Mendota Multiparameter Sonde

Profiles: 2017 - current. Environmental Data Initiative

https://doi.org/10.6073/PASTA/5F15BF453851987FC030B2F07A110B21 (2021).

50. Clum, A. *et al.* DOE JGI Metagenome Workflow. *mSystems* **6**, 10.1128/msystems.00804-20

(2021).

51. Bushnell, B. *BBMap Short Read Aligner and Other Bioinformatic Tools*.

https://www.osti.gov/biblio/1241166 (2014).

52. Nurk, S., Meleshko, D., Korobeynikov, A. & Pevzner, P. A. metaSPAdes: a new versatile

metagenomic assembler. *Genome Res* **27**, 824–834 (2017).

53. Li, H. *et al.* The Sequence Alignment/Map format and SAMtools. *Bioinformatics* **25**, 2078–

2079 (2009).

54. Kang, D. D. *et al.* MetaBAT 2: an adaptive binning algorithm for robust and efficient genome

reconstruction from metagenome assemblies. *PeerJ* **7**, e7359 (2019).

55. Chaumeil, P.-A., Mussig, A. J., Hugenholtz, P. & Parks, D. H. GTDB-Tk: a toolkit to classify

genomes with the Genome Taxonomy Database. *Bioinformatics* btz848 (2019)

doi:10.1093/bioinformatics/btz848.

56. Parks, D. H. *et al.* GTDB: an ongoing census of bacterial and archaeal diversity through a

phylogenetically consistent, rank normalized and complete genome-based taxonomy. *Nucleic*

Acids Research **50**, D785–D794 (2022).

57. Aroney, S. T. N. *et al.* CoverM: Read coverage calculator for metagenomics. Zenodo

https://doi.org/10.5281/zenodo.10531254 (2024).

58. Olm, M. R. *et al.* inStrain profiles population microdiversity from metagenomic data and

sensitively detects shared microbial strains. *Nat Biotechnol* **39**, 727–736 (2021).

59. Hyatt, D. *et al.* Prodigal: prokaryotic gene recognition and translation initiation site

identification. *BMC Bioinformatics* **11**, 119 (2010).

60. Aramaki, T. *et al.* KofamKOALA: KEGG Ortholog assignment based on profile HMM and

adaptive score threshold. *Bioinformatics* **36**, 2251–2252 (2020).

61. R Core Team. *R: A Language and Environment for Statistical Computing*. (R Foundation for

Statistical Computing, Vienna, Austria, 2022).

62. Barrett, T. *et al. Data.Table: Extension of `data.Frame`*. (2024).

63. Grolemund, G. & Wickham, H. Dates and Times Made Easy with lubridate. *Journal of*

Statistical Software **40**, 1–25 (2011).

64. Tange, O. GNU Parallel 20230822 ('Chandrayaan'). Zenodo

https://doi.org/10.5281/zenodo.8278274 (2023).

65. Boogaart, K. G. van den, Tolosana-Delgado, R. & Bren, M. compositions: Compositional

Data Analysis. (2023).

66. Oksanen, J. *et al.* vegan: Community Ecology Package. (2022).

67. Zeileis, A. *et al.* strucchange: Testing, Monitoring, and Dating Structural Changes. (2022).

68. DeCicco, L., Hirsch, R., Lorenz, D., Watkins, D. & Johnson, M. *dataRetrieval: R Packages*

for Discovering and Retrieving Water Data Available from U.S. Federal Hydrologic Web

Services. (U.S. Geological Survey, Reston, VA, 2023). doi:10.5066/P9X4L3GE.

69. McDonald, J. H. & Kreitman, M. Adaptive protein evolution at the Adh locus in Drosophila.

Nature **351**, 652–654 (1991).

Acknowledgements

Long-term datasets such as TYMEFLIES rely on researchers who contribute a portion of their time and effort to future projects. This work would not be possible without the generosity of many, including Lake Mendota sampling leads Angela Kent, Tony Yannarell, Ashley Shade, Stuart Jones, Ryan Newton, Georgia Wolfe, Emily Kara Read, Lucas Beversdorf, James Mutschler, and the original Microbial Observatory lead Eric W. Triplett. We thank Peter Golightly for his advice on interpreting the genes under selection data.

Funding

This project was supported by an E. Michael and Winona Foster WARF Wisconsin Idea Fellowship (R.R.R.), the U.S. National Science Foundation (NSF) (DBI-2011002) (R.R.R.), the U.S. National Institutes of Health (RO1-GM116853) (M.Kirk.), the U.S. NSF (DEB-1831730) (M. Kirk.), the U.S. Department of Agriculture (WIS01516 and WIS01789) (K.D.M.), the U.S. NSF (DEB-0702395, DEB-1344254) (K.D.M.) and a Simons Foundation Investigator in Aquatic Microbial Ecology Award (LI-SIAME-00002001) (B.J.B.). Support for sampling was provided by the U.S. NSF North Temperate Lakes Long-Term Ecological Research program (DEB-9632853, DEB-0217533, DEB-0822700, DEB-1440297, DEB-2025982) and the U.S. NSF Microbial Observatory program (MCB-9977903, DEB-0702395). The work (proposal: https://doi.org/10.46936/10.25585/60001198) conducted by the U.S. Department of Energy Joint Genome Institute (https://ror.org/04xm1d337), a DOE Office of Science User Facility, is supported by the Office of Science of the U.S. Department of Energy operated under Contract No. DE-AC02-05CH11231. The authors acknowledge the Texas Advanced Computing Center (TACC) at The

687　University of Texas at Austin for providing HPC resources that have contributed to the research

688　results reported within this paper (http://www.tacc.utexas.edu).

Author Contributions

691　R.R.R. and K.D.M. conceptualized the research and obtained initial funding. K.D.M. and B.J.B.

692　provided resources. R.R.R. conducted field and laboratory work and curated data. RRR performed

693　analyses and created visualizations. M.Kirk. advised statistical approaches. S.L.G., M.Kell..,

694　K.D.M., and B.J.B. advised analysis approaches. R.R.R. wrote the first draft, and R.R.R. and

695　B.J.B. wrote the final draft incorporating edits provided by M.Kirk., S.L.G., M.Kell., and K.D.M.

Competing Interests

698　The authors declare no competing interests.

Supplemental Information

Supplementary Table 1. TYMEFLIES sample metadata. All the JGI, GOLD, NCBI, and
internal McMahon Lab identifiers that pair with each metagenome sample.

Supplementary Table 2. TYMEFLIES reference genome metadata. NCBI identifiers
corresponding to each reference genome, as well as the quality results from CheckM2[5], the
taxonomy results from GTDB-tk[55], and average relative abundances as calculated by coverM[57].

Supplementary Table 3. Consistently selected gene annotations. KEGG annotations of
consistently selected genes. Table row order matches heatmap row order in Fig. 5f.