1 **S-nitrosylation of EZH2 at C329 and C700 interplay with PRC2 complex assembly,**
2 **methyltransferase activity, and EZH2 stability to regulate endothelial functions**
3

Ashima Sakhuja[1], Yash T Katakia[1], Shailesh Mani Tripathi[2], Ritobrata Bhattacharyya[1], Niyati Pandya

Thakkar[1], S Ramakrishnan[1], Srinjoy Chakraborty[1], Sumukh Thakar[1], Sandeep Sundriyal[2], Shibasish

Chowdhury[1], Syamantak Majumder[1*]

[1]Department of Biological Sciences, Birla Institute of Technology and Science (BITS) Pilani, Pilani

Campus, Rajasthan, India

[2]Department of Pharmacy, Birla Institute of Technology and Science (BITS) Pilani, Pilani Campus,

Rajasthan, India

* Corresponding author

Dr. Syamantak Majumder,

Department of Biological Sciences,

Birla Institute of Technology and Science Pilani,

Pilani Campus, Room No: 3277-R, FD-III,

Pilani, Rajasthan 333 031.

Email: syamantak.majumder@pilani.bits-pilani.ac.in

Phone: +91 1596-25-5693

ABSTRACT

Nitric oxide (NO), a versatile bio-active molecule modulates cellular function through diverse mechanisms including S-nitrosylation of proteins. However, the role of this post-translational modification in regulating epigenetic pathways was very limitedly explored. Herein, we report that NO causes S-nitrosylation of selected cysteine residues of EZH2 in endothelial cells (EC) resulting in SUZ12 dissociation from EZH2 bound PRC2 complex, reduced methyltransferase activity, and diminished nuclear localization eventually hampering its stability. We detected a significant reduction in H3K27me3 upon exposure to NO as contributed by the early dissociation of SUZ12 from the PRC2 complex. Longer exposure to NO donors caused EZH2 cytosolic translocation, its ubiquitination, and further degradation primarily through the autophagosome-lysosome pathway. Through *in silico* S-nitrosylation prediction analysis and site-directed mutagenesis assay, we identified three cysteine residues namely at locations 260, 329, and 700 in EZH2 and further determined that S-nitrosylation of cysteine 329 induced EZH2 instability while S-nitrosylation of cysteine 700 abrogated EZH2's catalytic activity. A double mutant of EZH2 containing mutations at Cysteine 329 and 700 remained undeterred to NO exposure. Furthermore, reinforcing H3K27me3 in NO exposed EC through the use of an inhibitor of H3K27me3 demethylase, we confirmed a significant contribution of the EZH2-H3K27me3 axis in defining NO-mediated regulation of endothelial gene expression and migration. Molecular dynamics simulation study revealed SUZ12's inability in efficiently binding to the SAL domain of EZH2 upon S-nitrosylation of C329 and C700. Taken together, our study for the first-time reports that S-nitrosylation dependent regulation of EZH2 and its associated PRC2 complex influences endothelial homeostasis.

Keywords: Endothelial cells, Enhancer of Zeste Homolog-2, Polycomb repressive complex 2, post-translational modification, S-nitrosylation

INTRODUCTION

Polycomb repressive complex 2 (PRC2) is a chromatin-modifying group of proteins that work as repressors of gene expression. PRC2 is formed by three major classes of proteins; the functional enzymatic component of the PRC2, Enhancer of Zeste Homolog-2 (EZH2); suppressor of zeste 12 (SUZ12), a PRC2 subunit and a scaffolding protein, and embryonic ectoderm development (EED) which assembles and stabilizes the PRC2 complex.[1,2] The importance of EED and SUZ12 in maintaining the HMT activity of EZH2 and in the stability of the PRC2 complex is well studied[3,4]. EZH2 is responsible for methylating the 27[th] lysine residue of histone H3 (H3K27) which results in the repression of gene expression[2,5]. Many studies have highlighted the importance of EZH2 as a crucial player in mediating cellular differentiation during development and also has a considerable role in various disease conditions[6]. Moreover, EZH2 mediates gene silencing in embryonic stem cells thereby promoting pluripotency, and in contrast also in the activation of differentiation[6,7]. The link between EZH2 regulation via deposition of the methylation marks and cancer progression is well reported describing its role in cancer progression and metastasis[8,9]. Indeed, EZH2 dependent suppression of the tumor suppressor genes during cancer progression promotes tumorigenesis.[1,2]

The role of several post-translational modifications such as ubiquitination[10], phosphorylation[11], O-GlcNAcylation[12], sumoylation[13], and methylation[14] in regulating EZH2 and its methyltransferase activity were previously reported. Ubiquitination is known to cause a decrease in the methyltransferase activity of EZH2 leading to its degradation. EZH2 has been shown to be phosphorylated by Akt[3] and AMPK[4], which leads to suppression of its methyltransferase activity.[15] O-GlcNAcylation at serine in the SET domain of EZH2 is associated with regulating its methyltransferase activity[16]. Similarly, other mentioned PTMs at specific residues of EZH2, they can regulate its stability and catalytic activity. However, till date, no studies described how S-nitroyslation of EZH2 regulates its localization, degradation and catalytic activity.

Nitrosylation by S-NO formation is one such PTM that has a potential in regulating protein function, stability, and cellular localization. It occurs through NO, a versatile free radical that forms S-NO by selective modification of the protein at cysteine/methionine residues and mediate numerous biological functions[17]. Moreover, many cells harbors endogenous NO producing machinery including NO synthase class of enzymes. NO synthase (NOS) family of enzymes use L-arginine to endogenously produce NO which plays diverse roles in different cell types[18]. EC is one of such cell type that contain a very cell type specific NO producing machinery named as endothelial NOS (eNOS). eNOS dependent release of NO mediates various signaling cascade which are essential for endothelial migration, survival and growth. Indeed, eNOS driven release of NO uses S-nitrosylation dependent regulation of proteins to govern endothelial functions[19,20,21,22]. However, how eNOS dependent release of NO directs gene expression changes in EC through chromatin regulation remains elusive.

In this study, we addressed whether NO-mediated post-translational modification of
histone methyltransferase EZH2 could influence its catalytic activity, localization, and stability.
Through the present study, we established that S-nitrosylation of cysteine residue(s) in EZH2 protein
amends its function as an epigenetic modulator, thereby proving the role of NO as a direct modulator
of epigenetic processes and further regulating gene expression changes. Our study for the first time
connects the link between eNOS dependent NO release and its function as epigenetic modulator
through regulation of EZH2 to alter chromatin structure and thereby dictate NO dependent gene
expression changes.
119
RESULTS
Nitric oxide exposure interplayed with EZH2 including PCR2 assembly, its methyltransferase
activity, subcellular localization, and stability
NO is well-reported to regulate gene expression changes in EC[22]. However, whether such an effect is
dependent on epigenetic processes specifically via regulation of EZH2 and PRC2 complex was never
been reported. We, therefore, assessed the effect of NO exposure on EZH2 protein and its catalytic
product H3K27me3 level in EC. NO exposure time-dependently caused a reduction in the level of
EZH2 protein level specifically observing a significant reduction at 2 hours post exposure to sodium
nitroprusside (SNP, Figs. 1A,B). We also observed a significant reduction in the catalytic product
H3K27me3, however, the reduction in H3K27me3 was detected within 1 hour of SNP treatment,
much earlier than the degradation of EZH2 (Figs. 1C,D). To confirm if such an effect of NO on EZH2
is exerted through transcriptional regulation, we performed qPCR analysis of EZH2 transcripts which
revealed no alteration in EZH2 transcript level upon SNP exposure (Supplementary Fig. 1A).
We next confirm such changes in EZH2 and H3K27me3 levels *ex vivo* in rat aorta exposed
to NO. Similar to *in vitro* findings, diminished EZH2 and H3K27me3 levels were detected in rat aorta
exposed to SNP (Figs. 1E,F). Because EC possesses endogenous NO-producing machinery namely
eNOS, we next question whether activation of endogenous NO-production machinery shall also affect
the levels of EZH2 and H3K27me3. Induction of EC by bradykinin dose- and time-dependently
caused a reduction in EZH2 and H3K27me3 levels (Supplementary Figs. 1B,C). To understand the
cell-specific effect of NO on EZH2 and H3K27me3, we next exposed HEK-293, a non-EC type of
human cells to NO donor and measured the level of EZH2 and H3K27me3. Through such an
experiment, we established that NO exposure caused a reduction in EZH2 and associated H3K27me3
levels independent of the cell type under study suggesting such alteration on EZH2 and its
downstream product could be sensitive to NO in many different cell types (Supplementary Fig. 1D).
We next evaluated the effect of NO on EZH2 overexpressed through plasmid construct and confirmed
that NO exposure also affects the level of overexpressed EZH2 as detected through HA tag or by
detecting the total EZH2 (Supplementary Fig. 1E).

Because we observed a reduction in H3K27me3 level much earlier than EZH2
degradation, we next urged whether NO exposure interplays with the assembly of the PRC2 complex
which is essential for EZH2 catalytic activity to cause H3K27me3 deposition. We therefore
performed a co- immunoprecipitation experiment using EZH2 antibody and detected dissociation of
SUZ12 from EZH2 in EC exposed to SNP for only 30 minutes (Fig. 1G). All other components of the
PRC2 complex remain associated with EZH2 at least until 30 minutes of NO exposure to EC (Fig.
1G). Interestingly, the protein level of other components of PRC2 including SUZ12, AEBP2, EED,
and JARID2 in total cell lysate remain unaltered upon NO exposure to EC (Supplementary Figs. 2A,
B).

Being a histone-modifying enzyme and part of the PRC2 complex, EZH2 is primarily
localized in the nucleus. However, cytosolic translocation of EZH2 was reported by many earlier
studies which also reported its cytosolic substrates including Talin[23] and small GTPases[24] to cause
actin polymerization. We thus wanted to explore the effect of NO exposure on EZH2 localization.
Subcellular fractionation (Fig. 1H) and immunofluorescence (Fig. 1I) followed by confocal imaging
strongly indicated cytosolic translocation of EZH2 in EC upon NO exposure. Moreover, confocal
imaging also indicated recruitment of EZH2 protein to the filamentous actin (Fig. 1I). No significant
changes in localization were detected in other components of the PRC2 including SUZ12, AEBP2,
EED, and JARID2 which were primarily localized in the nucleus independent of NO treatment
(Supplementary Fig. 2C). Because the turnover of H3K27me3 is not only dependent on the methyl
transferase EZH2 but also H3K27me3 specific demethylases UTX and JMJD3, we thus detected the
quantity of UTX and JMJD3 protein in EC exposed to NO. UTX and JMJD3 protein levels remained
unaltered in EC exposed to NO for 2 hours (Supplementary Figs. 2D,E).

Nitric oxide exposure caused S-Nitrosylation of EZH2 leading to early SUZ12 dissociation and
further altering its binding partners
In the non-canonical NO signaling pathway, many proteins are post-translationally modified through
nitrosylation of cysteine/methionine/tyrosine residue thereby regulating the function of such proteins.
We, therefore, next questioned whether such could be the case with EZH2 as well. Thus, after
establishing the effect of NO exposure on EZH2's methyltransferase activity, localization, and
degradation, we next evaluated whether NO post-translationally modified EZH2 through S-
nitrosylation thereby regulating its function. Previous studies reported regulation of EZH2 localization
and function via phosphorylation of distinct residues. S-nitrosylation of EZH2 protein has never been
reported earlier and therefore we first assessed whether EZH2 is S-nitrosylated upon NO exposure.
Through the iodoTMT assay using truncated recombinant EZH2 protein (carrying residues from 429-
728 aa) in a cell-free system, we first confirmed that NO exposure caused S-nitrosylation of a
truncated form of the EZH2 protein in a cell-free system (Supplementary Fig. 3). Further to confirm
this in a cellular system, EC were exposed to SNP followed by Biotin Switch assay and

immunoprecipitation using EZH2 antibody. Such an experiment also demonstrated S-nitrosylation of

EZH2 protein in EC subjected to NO treatment (Figs. 2A,B). Further to ascertain the S-nitrosylation,

we next used a pan S-nitrosylated antibody and confirmed through an immunoprecipitation followed

by an immunoblot experiment that EZH2 in EC was indeed S-nitrosylated upon NO exposure (Fig.

2C).

189   We next performed a co-immunoprecipitation experiment using EC which was exposed to

bradykinin, a natural inducer of endogenous NO production. In doing so, we detected a robust S-

nitrosylation of EZH2 protein concurrent with the loss of EZH2 binding with SUZ12 of the PRC2

complex (Fig. 2D). Furthermore, to support this data, we also analyzed the S-nitrosylation of HA-

tagged EZH2 protein in a HEK-293 overexpression system. Through a co-immunoprecipitation

experiment using HA antibody, we again confirmed S-nitrosylation of overexpressed EZH2 protein

upon exposure to NO (Fig. 2E).

196   To exclude the possibility of the effect of SNP on EZH2 as non-specific, we next used

GSNO which is more well-accepted to impart S-nitrosylation of cellular proteins. In so doing, we

found similar reduction in the level of EZH2 after 120 minutes of GSNO exposure (Fig. 3A). As

similar to SNP treated EC, further analysis of H3K27me3 in the GSNO treated cells revealed time

dependent depletion of H3K27me3 by 60 minutes which remained significantly low at least up to 120

minutes post-GSNO treatment (Fig. 3B). We next wanted to explore the comprehensive interacting

partners of EZH2 upon S-nitrosylation to understand the effect of such post-translational

modifications on its methylation, translocation and degradation. Mass spectrometric analysis revealed

the comprehensive association map of EZH2 in control and GSNO treated cells (Fig. 3C). A total of

261 proteins were identified to be associated with EZH2 while 341 different proteins associated with

EZH2 upon GSNO exposure (Figs. 3C, D). Out of 261 proteins in control condition, 48 unique

proteins were associated with EZH2 which completely dissociated upon GSNO exposure. In contrast,

128 (out of 341 in total) unique proteins were bound to EZH2 upon GSNO exposure which were not

detected to be associated with EZH2 in untreated cells (Figs. 3C-E, Supplementary Tables 1-2).

210   We next performed a close analysis of the interactome data which indicated few

interesting association patterns of EZH2 in untreated and GSNO treated conditions. Firstly, as

observed through co-immunoprecipitation experiment, EZH2 was found to be associated with histone

H3 in untreated cells, however, such association was not detected upon GSNO exposure. Surprisingly,

nuclear-to-cytosolic shuttling protein 14-3-3 was found to be associated with EZH2 in untreated

conditions while a complete loss of association was observed upon GSNO challenge (Fig. 3E,

Supplementary Tables 1-2). Although, we expected such association with 14-3-3 to be prominent

upon GSNO exposure due to EZH2 cytosolic shuttling after S-nitrosylation, however, we observed an

opposite correlation. Such data indicated that EZH2 may be using 14-3-3 during natural cytosolic

localization while its cytosolic localization upon S-nitrosylation is likely to be driven by other

unknown factors. More interestingly, upon GSNO exposure, we detected EZH2 association with

many proteins of the endosome/lysosome/proteasome pathway proteins including several Rab family

of proteins, HSP70&90 chaperon proteins, cathepsin D lysosomal protease, lysosome-associated

membrane glycoprotein 1, proteasome subunit beta type-5/alpha type-4, lysosomal acid ceramidase

(Figs. 3E-F, Supplementary Tables 1-2). EZH2 was not found to be associated with these proteins in

control conditions.

Nitric oxide caused the degradation of EZH2 primarily through autophagosome-lysosome

pathway while inhibition of endogenous nitric oxide machinery reversed nitric oxide dependent

degradation, activity and localization of EZH2

Because we observed association of many endosomal/lysosomal and proteasomal proteins, we

therefore questioned the role of lysosomal and proteasomal degradation pathways in S-nitrosylation

dependent degradation of EZH2. The role of post-translational modification-dependent regulation of

EZH2 protein degradation is least studied. Because we observed EZH2 protein degradation upon NO

exposure, we therefore explored the key degradation machinery responsible for S-nitrosylated EZH2.

Because ubiquitination of proteins plays a key role in degradation and previous report indicated

ubiquitination mediated degradation of EZH2[25], we first assessed the level of EZH2 ubiquitination

upon NO exposure and found that S-nitrosylated EZH2 are heavily ubiquitinated upon NO exposure

(Fig. 4A). Cells use lysosomal and proteasomal degradation pathways for protein degradation, we

thus used pharmacological inhibitors of proteasomal and autophagosome-lysosome pathway to

evaluate their relative contribution towards degradation of S-nitrosylated EZH2. Inhibition of

proteasomal pathway using MG132 was unable to reverse the level of EZH2 upon NO exposure (Fig.

4B). In contrast, inhibition of autophagosome-lysosome pathway using bafilomycin A significantly

although partially reversed EZH2 degradation upon NO challenge (Fig. 4C). We then inhibited both

proteasomal and autophagosome-lysosome pathway to evaluate the effect of such combination

inhibition on EZH2 protein level. In so doing, we detected complete reversal of EZH2 protein level

upon combination inhibition of proteasomal and autophagosome-lysosomal pathways (Fig. 4D). All

these data indicated that degradation of EZH2 upon NO exposure primarily occurs through

autophagosome-lysosome pathway, however, an inhibition of the autophagosome-lysosome pathway

could likely switch S-nitrosylated EZH2 degradation through proteasomal pathway. Successful

inhibition of autophagosome-lysosomal pathway upon bafilomycin A treatment was confirmed

through increase in p62 level (Fig. 4E).

We next questioned whether inhibition of endogenous NO producing machinery could

alter the downstream effect of natural inducers of endogenous NO production machinery in EC such

as bradykinin. We exposed the EC with L-NAME, a nonselective inhibitor of all type of NO synthase

(NOS) prior to inducing with bradykinin. As observed earlier in our study, bradykinin induction

caused reduction in EZH2 level which is completely reversed upon inhibition of NOS with L-NAME

(Fig. 4F). We also performed the assay in *ex vivo* rat aorta model which further revealed reversal of

EZH2 protein degradation and protection of H3K27me3 in aortic tissues exposed to L-NAME prior to

bradykinin treatment (Figs. 4G,H). We then performed localization analysis of EZH2 in cells exposed

to bradykinin alone or in combination with L-NAME. As expected, bradykinin induction caused

cytosolic translocation of EZH2 which are also found to be localized with actin cytoskeleton,

however, inhibition of NOS family of protein in bradykinin treated EC using L-NAME restricted

EZH2 localization to nucleus (Fig. 4I).

265 Protecting the level of EZH2 downstream product H3K27me3 through inhibition of

266 demethylases reversed nitric oxide dependent effect on endothelial migration and gene

267 expression changes

EZH2's effect on cellular function is primarily dependent on its regulation of gene expression changes

through repressive H3K27me3 mark in the chromatin. Moreover, NO signaling pathway converges to

changes in expression of genes associated with endothelial survival, proliferation and migration. We

therefore wanted to investigate whether EZH2 dependent catalysis of H3K27me3 play any role in

dictating NO driven regulation of endothelial function and gene expression changes. To do so, we

took a retrograde approach in which we pre-incubated the cells with GSK-J4, a selective inhibitor of

H3K27me3 specific demethylase JMJD3 and UTX. We first confirm that inhibition of JMJD3 and

UTX reversed NO dependent reduction in H3K27me3 level (Fig. 5A). We next performed endothelial

migration using wound healing assay and observed that NO induced endothelial migration is

abrogated upon preserving the level of H3K27me3 through inhibition of demethylase JMJD3 and

UTX (Fig. 5B).

In similar experimental settings, we next detected the transcript level expression of NO-

responsive genes in EC using qPCR experiment. Based on previously reported data sets, we choose to

detect the transcript level of NO responsive genes; VEGFa, TBX20, MMP2, FGF2, KDR, TIE2, TEK,

and Angiopoietin-2[26]. Through this analysis, we detected significant increase in expression of VEGFa,

TBX20, MMP2, FGF2, and Angiopoietin-2 upon NO exposure (Fig. 5C). Pretreatment of EC with

GSK-J4 prior to NO exposure completely abrogated NO dependent increment in VEGFa, TBX20,

MMP2, FGF2, and Angiopoietin-2 transcript level (Fig. 5C). Surprisingly, although reported earlier to

be responsive to NO in EC, in the present experimental settings, we were unable to record any

changes in transcript level of KDR, TIE2, and TEK genes (Fig. 5C).

289 S-nitrosylation of EZH2 at cysteine 329 and cysteine 700 is important for its stability and

290 methyltransferase activity respectively

Upon confirming the effect of NO on S-nitrosylation of EZH2 and associated changes in its stability,

catalytic activity, PRC2 assembly and translocation, we next focused on detecting the possible

residues of EZH2 that could be S-nitrosylated. To confirm this, we used GPS-SNO

(http://sno.biocuckoo.org/) prediction tools as stipulated in methodology section. Such analysis

predicted three possible sites at cysteine 260, 329, and 700 of EZH2 protein with score of 3.158,

2.576, and 3.109 respectively which is beyond the set cut off value of 2.443 (Supplementary Fig. 4A).

Predicted cysteine residues 260 and 329 lies within the domain II of EZH2 which essentially allows

SUZ12 association with EZH2, in contrast, predicted cysteine residue at 700 lies within the catalytic

SET domain of EZH2 which is essential for its enzymatic activity.

To further decipher the role of each of these cysteine residues, we generated point mutated

constructs of these cysteine by using site-directed mutagenesis kit to convert the codon to code for

serine in places of the actual cysteine (TGT/TGC to AGT/AGC) residues (Supplementary Fig. 4B).

Using overexpression of these constructs in HEK-293 cells, we first performed biotin switch assay to

assess the S-nitrosylation of these mutated EZH2. Such analysis revealed partial loss of S-

nitrosylation of EZH2 for each of the mutants EZH2 C260S, EZH2 C329S, and EZH2 C700S.

Interestingly, a complete loss of S-nitrosylation signal was not detected with single mutation

indicating multiple cysteine residues were likely to be S-nitrosylated in EZH2 protein upon NO

exposure (Fig. 6A). We then analyzed the effect of NO exposure on EZH2 protein and its catalytic

product H3K27me3 level in cells overexpressed with WT and mutated form of the EZH2 gene. As

observed earlier, NO exposure caused significant loss of HA tagged EZH2 protein and H3K27me3

level in cells overexpressed with HA tagged EZH2 WT gene (Figs. 6B,C). A comparable loss of HA

tagged EZH2 protein and H3K27me3 level was also detected in cells overexpressed with HA tagged

EZH2 C260S mutant gene (Figs. 6B,C). Interestingly, NO exposure did not alter the level of HA

tagged EZH2 protein in cells overexpressed with HA tagged EZH2 C329S mutant gene, however,

H3K27me3 level in these cells were significantly reduced upon NO exposure (Figs. 6B,C). In

contrast, NO significantly diminished the level of HA tagged EZH2 C700S protein in cells

overexpressing HA tagged EZH2 C700S mutant, however, surprisingly, H3K27me3 level upon NO

challenge remained relatively unaltered in these cells (Figs. 6B,C).

We next decided to generate a double mutant form of EZH2 gene where both cysteine

residues at location 329 and 700 are mutated to serine thereby generating a double mutant named

EZH2 C329S C700S mutant. We then overexpressed this double mutant in HEK-293 cells followed

by treating the cells with NO. Double mutated (at cysteine 329 and 700 residues) form of the HA

tagged EZH2 was completely insensitive to NO exposure both in context to the level of HA tagged

EZH2 C329S C700S protein as well as its product H3K27me3 (Fig. 6D). In similar settings, cells

overexpressing WT EZH2 gene showed significant reduction in both HA tagged EZH2 protein and

H3K27me3 level (Fig. 6D). We next performed immunofluorescence experiment and confocal

imaging study to ascertain the effect of these individual and double mutations of EZH2 on NO driven

HA tagged EZH2 localization. In so doing, we observed that NO exposure caused significant

cytosolic delocalization of HA tagged EZH2 wt, C260S, and C700S mutated forms of HA tagged

EZH2 (Fig. 6E). Interestingly, NO was unable to induce significant cytosolic localization of HA

tagged EZH2 C329S and double mutated HA tagged EZH2 C329S C700S protein (Fig. 6E).

S-nitrosylation of EZH2 protein at C329 and C700 causes conformational changes in EZH2-SUZ12 complex leading to loose association of SUZ12 with the SAL domain of EZH2

Upon experimental validation of EZH2's S-nitrosylation and its effect on EZH2-SUZ12 interaction, catalytic activity, localization and EZH2 protein stability, we next employed molecular dynamics analysis to understand the effect of S-nitroyslation on EZH2 interaction with SUZ12 protein. Through alphafold modeling of EZH2-SUZ12, we showed the initial conformations of the EZH2-SUZ12 complex in both the EZH2 WT and EZH2 S-nitrosylated at C324 (originally C329) and C695 (originally C700) residues (Fig. 7A) while also clearly visualizing the S-nitrosylation at C324 (Fig. 7B) and C695 (Fig. 7C). The RMSD calculations elucidate the extent to which the conformations of the EZH2 WT and EZH2 S-nitrosylated complexes deviated from their initial states during the course of MD simulations. In comparison to the WT complex, a sharp rise of RMSD values is observed for S-nitrosylated complex after 100 ns indicating possible change in the overall conformations during the simulations. A system showed larger conformational changes for EZH2 S-nitrosylated than for EZH2 WT complex (Fig. 7D).

Gamblin et al. (2016) reported that residues 112 to 121 within the SAL pack engage with the SET-1 region, serving to stabilize its conformation within the active complex[27]. Importantly for regulation in the human PRC2 complex, the conserved acidic residues 584-588 of SUZ12 reciprocally pack against residues 112-121 of SAL. Consequently, we conducted measurements of the backbone RMSD for both residue ranges, 112-121 of SAL and 584-588 of SUZ12, revealing a significant fluctuation in the RMSD within the SAL region of EZH2 S-nitrosylated with little to negligible RMSD fluctuation for SUZ12 residues ranging from 584-588 (Figs. 7E,F).

Another crucial parameter of MD is root mean square fluctuation (RMSF) to determine the flexibility of protein residues. The RMSF value of both EZH2 WT and EZH2 S-nitrosylated remained relatively stable with a nearly similar trend (Supplementary Fig. 5A). Primarily, the residues near the SAL and long loop region (residue 112-121 and 345-421) display significant fluctuations. Apart from these loops and SAL region of EZH2, RMSF analysis shows stable interactions in both the simulated complexes. Subsequently, we visualized the interface hydrogen bond interactions between SAL residues 112-121 and SUZ12 residues 584-588 throughout 1 µs simulation. Interestingly, in the EZH2 WT, we observed the oxygen atom of LEU166 in EZH2 interacting through the hydrogen bond with LYS587 of SUZ12 and the HE22 atom of GLN117 in EZH2 interacting with the OD2 atom of ASP585 in SUZ12 (Figs. 7H,I) These interactions persisted throughout the simulation. In the case of S-nitrosylated mutant, these two hydrogen bond interactions were initially established and maintained up to 300 ns. However, beyond this time point, the interactions were no longer observed.

Hydrogen bonds (H-bonds) are important non-covalent interactions that help stabilize protein-ligand and protein-protein complexes. To better understand the dynamics of these

interactions at the interface, we analyzed the H-bonds formed between specific residues 112-121 of SAL and 584-588 of SUZ12. This examination allowed us to gain insights into the overall stability and binding strength of the complexes (Supplementary Fig. 5B). During the simulations, we closely observed the formation of H-bonds within a distance of 3.5 Å. Upon conducting the analysis, it was discovered that the specific residues 112-121 of SAL were found to interact with 584-588 of SUZ12 in the EZH2 WT complex. This interaction resulted in an average of 2 hydrogen bonds. Conversely, the S-nitrosylated counterpart exhibited a significantly lower average of 0.35 hydrogen bonds suggesting EZH2 WT complex have stronger interactions than EZH2 S-nitrosylated variant. This observation also aligns with the energy profile (Supplementary Fig. 5C). We further measured the distance between the oxygen atom of LEU166 in EZH2 and the hydrogen atom of LYS587 in SUZ12, as well as the distance between GLN117 in EZH2 and ASP585 in SUZ12. A sharp increase in the distance between GLN117 and the interacting residue ASP585 after 150 ns in the EZH2 S-nitrosylation complex simulation (Fig. 7G) was observed, that ultimately led to the disruption of the interaction. Such distance between the oxygen atom of LEU166 in EZH2 and the hydrogen atom of LYS587 in SUZ12, as well as the distance between the HE22 atom of GLN117 in EZH2 and its interacting atom OD2 of ASP585 in SUZ12 at 100 ns (after stabilization of the complex upon initial simulation) and at 800 ns, was also presented through the structure analysis (Figs. 7H-K).

To determine the more accurate free energy binding of all complexes, we utilized the MM/PBSA-based method. The binding free energy in this context refers to the total of all non-bonded interaction energies between the EZH2 and SUZ12 during the MD simulation, including van der Walls, electrostatic, polar solvation, and SASA energies. A more negative binding free energy indicates a stronger affinity between the EZH2 and the SUZ12. We calculated the binding free energies using 1µs MD trajectory, revealing that EZH2 WT and EZH2 S-nitrosylation have a binding free energy of -4649.807 ± 454.449 kJ/mol and -4582.981 ± 453.056 kJ/mol, Subsequently, estimation of their contributions to overall $\Delta\Delta G\_bind=66$ kJ/mol indicates that the contributions vary from one system to the next. On average, the wildtype state tends to have a marginally higher contribution (Supplementary Fig. 5C and Supplementary Table 3). Furthermore, the comparison of the binding energy decomposition between selected residue of SAL and SUZ12 in EZH2 WT and EZH2 S-nitrosylated complexes located at the binding interface reveals that the EZH2 WT predominantly contributes to binding in locally stabilizing manner, Conversely, EZH2 S-nitrosylation displays less contribution to binding energy and these effects might be the consequence of introduced nitrosylation at cysteine 324 and 695 (Supplementary Figs. 5D,E). Overall, these modeling results suggest that S-nitrosylation of the cysteine residues might induce a conformational change in the SAL region which is crucial for the regulation of the PRC2 complex. This conformational change weakens the SUZ12-SAL interaction, ultimately resulting in the destabilization of the SUZ12-EZH2 complex.

DISCUSSION

A key regulatory pathway of NO-dependent cellular function is through S-nitrosylation of protein
which is a reversible post-translational modification. Extensive studies have shown that S-
nitrosylation regulates diverse physiological and pathological processes including angiogenesis,
adaptive immunity, diabetes, heart failure, stroke etc. through modulating stability, activity,
subcellular localization, conformation change, or protein-protein interaction of target proteins[28].
Although NO signaling alters gene expression changes in vascular endothelium[29], however, the role of
NO-dependent S-nitrosylation of proteins in defining the epigenetic landscape of EC is not clearly
understood. PRC2 complex is one of the most abundant and well-studied complexes engaged in gene
repression through regulation of chromatin compaction. EZH2 being the catalytic unit of PRC2
complex catalyzes the deposition of H3K27me3 mark that causes chromatin compaction and thereby
resulting in the repression of gene expression. Nevertheless, how S-nitrosylation regulate EZH2
protein and thereby the activity of the crucial PRC2 complex was never reported. Herein, for the first
time we described that NO caused S-nitrosylation of EZH2 protein leading to inhibition of its catalytic
activity, disassembly of SUZ12 from PRC2 complex bound EZH2, cytosolic translocation of the
protein followed by ubiquitination and degradation through autophagosome-lysosome pathway.
Moreover, mass spectrometric analysis revealed S-nitrosylation of EZH2 significantly alters its
interactome preferentially allowing association with proteins of the endosome/lysosome/proteasome
pathways. We also demonstrated that natural induction of endogenous NO machinery in EC also
imparts the exact same effect on EZH2 protein. Through site-directed mutagenesis studies, we
specifically identified that S-nitrosylation of Cysteine 329 and Cysteine 700 residues were responsible
for EZH2 protein translocation/degradation and catalytic deactivation respectively. Molecular
dynamics analysis suggested structural changes in EZH2 protein upon S-nitrosylation causing
instability of EZH2-SUZ12 complex. Most importantly, we demonstrated that such S-nitrosylation of
EZH2 was responsible for NO-dependent epigenetic regulation of endothelial gene expression and
cellular migration (Figure 8).

430   Post-translational modification of EZH2 protein through phosphorylation at several sites
were reported earlier[29–32]. Such modifications of EZH2 differentially regulated the localization,
function and its association with other proteins of the PRC2 complex. For instance, AMPK
phosphorylated EZH2 at T311 to disrupt the interaction between EZH2 and SUZ12 thereby
attenuating PRC2-dependent methylation of histone H3 at Lys27 in ovarian and breast cancer.
Moreover, such phosphorylation of EZH2 by AMPK caused upregulation of PRC2 target genes, many
of which are known tumor suppressors thereby suppressing the growth of tumor cells[32]. Similarly, we
observed that S-nitrosylation of EZH2 protein at C329 and C700 caused its dissociation from SUZ12
thereby abrogating the catalytic potential of PRC2 complex. During breast cancer metastases, p38-
mediated EZH2 phosphorylation at T367 promotes EZH2 cytoplasmic localization and potentiates
EZH2 binding to vinculin and other cytoskeletal regulators of cell migration and invasion.
Interestingly, ectopic expression of a phospho-deficient T367A-EZH2 mutant is sufficient to inhibit

EZH2 cytoplasmic expression, disrupt binding to cytoskeletal regulators, and reduce EZH2-mediated

adhesion, migration, invasion, and development of spontaneous metastasis[29]. In the present study, we

observed that S-nitrosylation of EZH2 in specific at C329 caused its cytosolic localization, a transient

binding to actin cytoskeleton, ubiquitination of EZH2 protein followed by degradation through

autophagosome-lysosome pathway. A very recent study demonstrated DCAF1 mediated

phosphorylation of EZH2 at T367 to augment its nuclear stabilization and enzymatic activity in colon

cancer cells. Such DCAF1-mediated EZH2 phosphorylation followed by nuclear localization led to

elevated levels of H3K27me3 and altered expression of growth regulatory genes in cancer cells[30]. In

contrast, herein we report that S-nitrosylation of C329 residue of EZH2 caused its cytosolic

localization followed by lysosomal degradation. Moreover, S-nitrosylation at another site C700 led to

inactivation of EZH2's catalytic activity most likely due to early dissociation of SUZ12 protein from

PCR2 complex. Although, phosphorylation dependent regulation of EZH2 protein and the PRC2

complex are well documented, however, other post-translational modifications especially NO

dependent S-nitroyslation of EZH2 and its regulation of the overall function of this protein has never

been reported. Herein, we demonstrated endogenous NO producing machinery of EC as well as

external supplementation of NO using donors regulated EZH2 binding with SUZ12, its localization,

and protein stability. Such effect of NO on EZH2 contributed towards NO driven gene expression

changes and endothelial migration. These are primarily achieved by NO through S-nitroyslation of

EZH2 protein at C329 and C700 residues.

In EC endogenous eNOS dependent release of NO drives cell survival, migration and

growth typically acting through canonical and non-canonical pathways[33,34]. Canonically, NO induces

soluble guanyl cyclase (sGC) and in downstream Protein Kinase G to mediate actin remodeling

required for cell migration[35,36]. Moreover, NO signaling also promote endothelial gene expression

changes associated with endothelial survival and growth[37,38]. Such gene expression changes by NO in

EC was thought to be primarily controlled by gene transcription and mRNA translation via iron-

responsive elements[39]. Herein, for the first time, we reported that eNOS dependent NO release

instigates non-canonical pathway through S-nitroyslation of EZH2 protein and its product H3K27me3

thereby controlling the epigenetic regulation of gene expression. This was supported by our data

wherein preserving the level of EZH2 catalytic product H3K27me3 through inhibition of the

demethylases JMJD3-UTX reversed NO dependent gene expression changes in EC. Moreover, we

also observed that regulation of EZH2 by NO was independent of the cell type under study which is

also been supported through our data in HEK-293 cells exposed to NO. In parallel to well reported

function of NO[40,41] as well as EZH2[42–44] in actin cytoskeleton remodeling, we observed that NO

exposure caused transient localization of EZH2 on the actin cytoskeleton most likely prior to its

degradation through autophagosome-lysosomal pathway. In addition, preserving the level of

H3K27me3 via inhibition of demethylases reversed NO effect on cell migration. Through the present

study, we are unable to establish whether the effect of NO on endothelial migration is mainly driven

by its regulation of EZH2-actin cytoskeleton remodeling axis or via regulation of EZH2-H3K27me3-
gene expression regulation axis. Nonetheless, NO dependent remodeling of actin cytoskeleton may
not alone be driven by canonical sGC-cGMP-PKG-VASP pathway, in addition, regulation of EZH2
via NO may contribute towards its actin remodeling function.

In summary, our study for the first time reported that NO signaling pathway converges to
epigenetically regulate gene expression via controlling the stability and catalytic activity of PRC2
complex, in specific the EZH2 component of the complex. This is primarily driven by NO through S-
nitrosylation of C329 and C700 residues of EZH2 which promotes early dissociation of SUZ12 from
EZH2-PCR2 complex followed by cytosolic localization of EZH2 and its degradation through
autophagosome-lysosome pathways. We also showed that such an effect of NO on EZH2 is cell type
independent. Moreover, we also reported that NO driven gene expression changes in EC is primarily
dependent on NO regulation of EZH2 and its catalytic product H3K27me3. Our molecular dynamics
simulation study provided evidence that upon S-nitrosylation of EZH2 at C329 and C700, caused
changes in the structure of EZH2 SAL domain leading to dissociation of SUZ12 binding with EZH2.
Furthermore, through site directed mutagenesis of EZH2 at C329 and C700 depicted that S-
nitrosylation of C329 is responsible for EZH2 cytosolic translocation and degradation while S-
nitrosylation of C700 was primarily responsible for catalytic inactivation of EZH2. Our findings shed
light into a new mechanism of NO mediated regulation of EZH2 and PCR2 complex to regulate gene
expression associated with endothelial function.
498

MATERIALS AND METHODS
Cell culture
Experiments were done primarily in EA. hy926 (immortalized human umbilical vein endothelial cell),
purchased from ATCC, Manassas, USA (#CRL-2922). The cells were cultured in Dulbecco's
modified Eagle's medium (DMEM) (#AL006A; HiMedia Laboratories) supplemented with 10% fetal
bovine serum (#RM1112, FBS; HiMedia Laboratories) and 1% penicillin/streptomycin (#10378, PS;
Gibco). The cells were passaged every 2-3 days. Human umbilical vein EC (HUVEC) (#CL002-
2XT25, HiMedia Laboratories) were cultured and passaged using HiEndoXL™ EC Expansion
medium (#AL517, HiMedia Laboratories) along with 4%growth factor, 5% fetal bovine serum and
1% penicillin/streptomycin antibiotic. Human Embryonic Kidney HEK-293 cell line was a kind gift
from Prof. Uma Dubey (Department of Biological Sciences, Birla Institute of Technology and Science
Pilani, Pilani Campus, India) were maintained in Minimum Essential Media Eagle (MEM) (#AT154;
HiMedia Laboratories). All the cells were maintained at 37 ° C with 5% CO2 in a humidified
incubator.
513

Plasmid and Transfection

Transfection experiments were done with human WT PCMVHA hEZH2 plasmid (#24230, Addgene),

a kind gift from Dr. Kristian Helin (Biotech Research and Innovation Centre, University of

Copenhagen, Copenhagen, Denmark). HEK-293 cells with 80% of confluency were transfected using

250ng/ml of pCMVHA hEZH2 and Lipofectamine 2000 (#11668030, Invitrogen, Thermo Fisher

Scientific) for 4 hrs. After 48 hours of incubation, the cells were then used for immunoblotting and

immunofluorescence experiments.

Inducers and Inhibitors

Sodium Nitroprusside dihydrate (SNP, 500 µM) (#71778, Sigma Aldrich) was used as an exogenous

donor of NO in our experiments. Natural induction of NO was activated using 10 µM of Bradykinin

(BK) (#B3259; Sigma Aldrich). Inhibition of NO in cell system was achieved by treating the cells

with N_{ω}-Nitro-L-arginine methyl ester hydrochloride (L-NAME, (1 mM) (#N5751; Sigma Aldrich)

which is an analog of arginine. Bafilomycin A1 (100 nM) (#11038; Cayman Chemicals) was used as

an autophagy Inhibitor. MG132 (1 µM) (#M7449), a proteasomal inhibitor was from Sigma Aldrich.

GSK-J4 (5 µM) (#SML-0701), a H3K27me3 demethylase inhibitor was from Sigma Aldrich.

Prediction of S-nitrosylation site in EZH2 protein using GPS-SNO prediction tools

To identify the cysteine residues that are likely to be S-nitrosylated in EZH2 protein, we performed an

in silico analysis of the same by using GPS-SNO prediction tools as available at

http://sno.biocuckoo.org/. These tools have been developed by a group of scientists by manually

collecting 467 experimentally verified S-nitrosylation sites in 302 unique proteins from scientific

literature and can be used for prediction of S-nitrosylation sites. The developer performed extensive

leave-one-out validation and 4-, 6-, 8-, 10-fold cross-validations techniques to calculate the prediction

performance and system robustness[45]. More than 250 publications have used these tools for prediction

of possible S-nitrosylation sites. We next used the human EZH2 primary protein sequence available in

the addgene for hEZH2 plasmid (#24230, Addgene) and predicted the possible cysteine residues using

the GPS-SNO prediction tools. EZH2 in total has 34 cysteine residues, and according to the partially

resolved crystal structure of EZH2, none of these residues are involved in di-sulphide bonds and are

likely to be cysteine residues with free –SH group. The analysis through the GPS-SNO platform

revealed three unique sites at cysteine 260, 329, and 700 which are likely to be S-nitrosylated.

Subcellular fractionation

Following the treatment with SNP for the given time point conditions, EA. hy926 cells were washed

with phosphate□buffered saline (PBS) followed by scraping. The cell pellet was resuspended in

ice□cold PBS containing 0.1% Nonidet P□40 (NP□40) and 1% protease inhibitor (#P8340; Sigma

Aldrich) following centrifugation. After 10 mins of centrifugation, the supernatant was collected as

the cellular and cytoplasmic fraction, and the nuclear fraction was further pelleted down. The cellular

and nuclear lysate obtained was sonicated for 10 s (×2) followed by Immunoblotting experiments.

Immunoblotting

Cells grown to 80% confluency were provided treatment conditions followed by washing with

sterile1xPBS. They were then incubated in RIPA buffer (#9806; Cell Signaling Technology)

containing protease inhibitor for lysis after washing with sterile 1X PBS. After scraping, the cells

were subjected to repeated cycles of sonication for 10 s (×2). This was followed by a centrifugation

step at 10,000 g for 10 min to pellet down the cell debris and collect the supernatant. Protein

quantification was done by Bradford assay. After this, the protein samples were mixed with 5×

laemmli buffer and heated at 100°C for 10 mins. They were then processed for SDS polyacrylamide

gel electrophoresis with 10–250 kDa prestained protein ladder (#PG-PMT2922; Genetix) used for

molecular weight reference. Proteins were transferred onto a nitrocellulose membrane (Bio-Rad) at 15

V, 2.5 A for 35 min. The membrane was blocked with 5% skimmed milk for 1 h. The membrane was

incubated overnight at 4°C with the primary antibodies: EZH2 mAb (1:1000; #5246), JARID2 Rabbit

mAb (1:1000; #13594), SUZ12 Rabbit mAb (1:1000; #3737), EZH1 Rabbit mAb (1:1000; #42088),

EED Rabbit mAb (1:1000; #51673), AEBP2 Rabbit mAb (1:1000; #14129). H3K27me3 (1:1000),

UTX (1:1000), JMJD3 (1:1000), HA☐Tag Rabbit mAb (1:500; #3724), GAPDH Rabbit mAb

(1:2000; #5174), β-actin (1:2000; #3700), H3 Rabbit mAb (1:2000, #4499) (Cell Signaling

Technology), S-Nitrocysteine mAb (1:500; #94930) Ubiquitin mAb(1:1000;#3933). Afterward, the

blots were washed with TBS-T(x3) and incubated with a secondary peroxidase☐conjugated antirabbit

or mouse IgG antibody (1:2000) (#7074 or #7076, Cell Signaling Technology) overnight. These

conjugation reactions were detected using the Clarity™ (#1705061) or Clarity™ Max Western ECL

Substrates (#1705062) (Bio☐Rad).

Immunofluorescence and Confocal Microscopy

Following SNP treatment in EA.hy926 & transfected HEK-293 cells were fixed using 2%

paraformaldehyde (10 mins). The coverslips were then treated with 0.1% Triton X☐100 (5 min) for

permeabilizing the cells. Blocking was done with 2% bovine serum albumin (BSA) for 60 mins at

room temperature. Afterward, cells were incubated with primary antibodies EZH2 Rabbit mAb

(1:1000) or HA-tag (Alexa Fluor™ mAb 647 Conjugate) (1:2000; #37297) overnight at 4°C. The

cells were co-stained with Alexa Fluor™ Plus 480 conjugated antirabbit IgG secondary antibody

(1:4000; #A32732) (Thermo Fisher Scientific) for 1 h followed with phalloidin for staining F-actin

(1:5000; #A22287 or #A12379, Thermo Fisher Scientific) for 30 min. At last, the cells were stained

with 1 µM of DAPI (#D9542; Sigma-Aldrich) for 10 min for nuclear staining. Imaging was done

using Zeiss ApoTome 2.0 microscope (Carl Zeiss) and fully Spectral Confocal Laser Scanning

Microscope (#LSM 880Carl Zeiss).

Coimmunoprecipitation Experiments

Protein extraction was carried out using 1x RIPA buffer. After washing the Dynabeads (SureBeads™

Protein A Magnetic Beads, #1614013, BioRad) in PBS-T(x3), they were preincubated with the EZH2

mAb, Ha Tagged mAb, ubiquitin mAb or S-Nitrocysteine mAb (1:50) for 2hrs. The beads were then

washed and incubated overnight with a protein cell lysate containing 1000 µg of total protein for

pulldown of the desired protein from the cell lysate. The cell lysate was then heated at 70° C for 10

mins and magnetized to be separated from the beads. This was then processed for immunoblotting

experiments.

Ex vivo **experiments using rat aortic tissues**

The Institutional Animal Ethics Committee of BITS Pilani, Pilani Campus approved all experimental

procedures for the rodent studies. Ex vivo experiments were done using Male Wistar rats aged 12–16

weeks. After giving anesthesia, the rats were dissected from the ventral end. PBS was used to perfuse

the heart and aorta, eliminating blood cells from the vessels. The primary aortas were then collected,

and the fatty tissue layers were removed carefully. For acquiring the aortic explants, the aorta was cut

in a size of 4-5 mm cylindrical pieces. Following a PBS wash, the explants were cultured in

HiEndoXL™ EC expansion medium and incubated in the complete growth medium for 12 hrs before

initiating the treatment. Afterward, the tissue fractions were homogenized and subjected to sonication

after suspending them in RIPA lysis buffer. The protein was estimated using Bradford assay followed

by immunoblotting studies.

RNA Isolation, cDNA Synthesis and qPCR

Reverse transcriptase-quantitative polymerase chain reaction (RT-qPCR) was performed to measure

different gene expressions at the transcription level. Upon reaching 80% confluency EA.hy926 cells

underwent GSK-J4 treatment for 4 hr followed by SNP exposure for 2hrs. After that, RNA was

isolated from the cells using Trizol Reagent (#15596, TRIzol™ Reagent; Life Technologies, Thermo

Fisher Scientific). RNA isolation was succeeded by cDNA preparation from 1µg of total RNA using

iScript™ cDNA Synthesis Kit (#1708891; Bio-Rad Laboratories, Hercules, CA, United States). The

quality and quantity of the RNA was measured using a Nano-Drop spectrophotometer (SimpliNano;

GE Lifesciences). Before cDNA synthesis, DNA contamination was removed by pre-treating isolated

RNAs with the DNase. This was followed by Real-time PCR where iTaq™ Universal SYBR® Green

Supermix (#1725124; Bio-Rad Laboratories) was used with a total master mix volume of 10 µl and

GAPDH was taken as the housekeeping gene. Data analysis was done by calculating delta-delta Ct.

Details of the primers are as given in Supplementary Table 4.

Biotin Switch Assay

EA.hy926 cells or EZH2 WT/mutant overexpressing HEK-293 treated with NO donor SNP (500 µM)

for 2 hours were processed with the help of a Biotin Switch Assay kit (Abcam) using the standard

manufacturer protocol. With this, all the "S-NO" (S-nitrosylated) groups were replaced with Biotin,

628 forming an S-Biotin complex, which was detected by incubation with the streptavidin bound HRP

629 reagent. After this, the samples were immunoprecipated with EZH2 specific antibody using the

630 above-mentioned protocol. This was followed by dot blot and immunoblotting experiments for the

631 detection of the S-nitrosylation of the EZH2 protein.

Scratch wound healing Assay

633 EA.hy926 cells were grown in a 24-well plate. They were pre-treated with GSK-J4 for 4 hrs following

634 SNP treatment at regular intervals(0-24hrs) to measure the wound healing under different treatment

635 conditions or in combination. Imaging was done at definite intervals to calculate the wound healed by

636 cell migration. The wound was created in a straight line. Another perpendicular wound to the first one

637 was drawn to create a cross-shaped wound using a 1mm microtip.

Site-directed mutagenesis reaction

Phusion Site-Directed Mutagenesis Kit (#F541Thermo Fisher Scientific™) was used to insert point

mutations at the specific positions in the pCMV-HA hEZH2 plasmid as mentioned in the

manufacturer's protocol. Prior to this, the primers were phosphorylated using T4 Polynucleotide

Kinase (#EK0031, Thermo Fisher Scientific™). After the ligation step, the mutated product was

transformed in the competent Dh5-Alpha E. coli cells. The sequence of primers for inserting point

mutations at the predicted sites to convert cysteine to serine are provided in Supplementary Table 5.

Transformation experiments and Plasmid Isolation

The ligation product from the SDM reaction was mixed with 100µl of competent DH5-Alpha cells

and kept on ice for 25 mins. This was followed by heat shock at 42° C for 30 seconds. After adding

900 µl of SOC medium, the product was kept at 37° C for 1 hr. This was followed by centrifugation at

3000 rpm for 5 mins. 800 µl of supernatant was removed, and the pellet was resuspended in the

remaining 100 µl media. The product was then spread on the LB Agar plate with the help of a

spreader and kept overnight at 37° C. A single colony picked from the plate was used to inoculate 5

ml of Luria Broth (LB) containing ampicillin, which was kept overnight at 200 rpm and 37˚C. This

was used for plasmid isolation, which was performed using the manufacturer's protocol (#12123,

Qiagen Plasmid Minikit). Isolated plasmid was then sent for Sanger Sequencing to confirm the

insertion of point mutations at the desired location and then used for further transfection experiments.

Detection of S-Nitrosylation using iodoTMT labelling ™ Reagent

EZH2 recombinant protein (#MBS2097714, MyBioSource) were treated with GSNO (100µM) for 30

mins. The samples were then processed for iodoTMT labelling with Pierce ™ S-Nitrosylation

Western Blot kit (#90105, Thermo Fisher Scientific™). A total of 5µg/ml of EZH2 recombinant

protein sample was prepared for both control and GSNO treated conditions in 100µl of HENS Buffer.

Next, MMTS was added to each, following vortexing for 1 min and incubating it at room temperature

for 30 mins to block all free thiols. The protein was then precipitated by adding 600µl of pre-chilled (-

20°C) acetone and freezing the samples at -20°C for 1 hr for MMTS removal. The samples were

centrifuged at 4°C for 10 minutes (at 12000g). The tubes were inverted to decant the acetone and the

white pellet was dried for 10-15 mins. The pellet was resuspended in 100µl of HENS Buffer and 2µl

of the iodoTMTzero labeling reagent was added followed by the addition of 4µl of 1M sodium

ascorbate and vortexing briefly to mix and incubating it for 2 hrs at room temperature. The samples

were then processed for immunoblotting experiments with anti-TMT antibody.

Sample Preparation for Mass Spectroscopy analysis

EA.hy926 cells were treated with GSNO (100µM) for 30 mins followed by lysis with RIPA buffer.

Protein extraction was carried out using 1x RIPA buffer. After washing the Dynabeads (SureBeads™

Protein A Magnetic Beads, in PBS-T(x3), they were pre-incubated with the EZH2 mAb, (1:50) for

2hrs. The beads were then washed and incubated overnight with a protein cell lysate containing

1000ug of total protein for pulldown of EZH2 from the cell lysate. The cell lysate was then heated at

70° C for 10 mins and magnetized to be separated from the beads. Dynabead bound

immunoprecipitated proteins were eluted by heating the samples in 1X PBS. Eluted protein per

sample was used for digestion and reduced with 5 mM TCEP and further alkylated with 50 mM

iodoacetamide and then digested with Trypsin (1:50, Trypsin/lysate ratio) for 16 h at 37 °C. Digests

were cleaned using a C18 silica cartridge to remove the salt**.** Protein per sample was used for

digestion and reduced with 5 mM TCEP and further alkylated with 50 mM iodoacetamide and then

digested with Trypsin (1:50, Trypsin/lysate ratio) for 16 h at 37 °C. Digests were cleaned using a C18

silica cartridge to remove the salt and dried using a speed vac. The dried pellet was resuspended in

buffer A (2% acetonitrile, 0.1% formic acid).

Mass Spectrometric Analysis of Peptide Mixtures

Experiments were performed on an Easy-nlc-1000 system coupled with an Orbitrap Exploris mass

spectrometer. 1µg of peptide sample were loaded on C18 column 15 cm, 3.0µm Acclaim PepMap

(Thermo Fisher Scientific) and separated with a 0–40% gradient of buffer B (80% acetonitrile, 0.1%

formic acid) at a flow rate of 500 nl/min) and injected for MS analysis. LC gradients were run for 110

minutes. MS1 spectra were acquired in the Orbitrap (Max IT = 60ms, AGQ target = 300%; RF Lens =

70%; R=60K, mass range = 375−1500; Profile data). Dynamic exclusion was employed for 30s

excluding all charge states for a given precursor. MS2 spectra were collected for top 20 peptides. MS2

(Max IT= 60ms, R= 15K, AGC target 100%).

Data Processing

All samples were processed and RAW files generated were analyzed with Proteome Discoverer (v2.5)

against the Uniprot Human database. For dual Sequest and Amanda search, the precursor and

fragment mass tolerances were set at 10 ppm and 0.02 Da, respectively. The protease used to generate

peptides, i.e. enzyme specificity was set for trypsin/P (cleavage at the C terminus of "K/R: unless

followed by "P"). Carbamidomethyl on cysteine as fixed modification and oxidation of methionine

and N-terminal acetylation were considered as variable modifications for database search. Both

peptide spectrum match and protein false discovery rate were set to 0.01 FDR.

Model and Protein Preparation for *in silico* analysis

The crystal structure of the Human Polycomb Repressive Complex 2 (PDB ID: 5HYN) was acquired

from the Protein Data Bank[27]. This complex exhibits multiple missing loops, particularly within the

EZH2 residues at positions 1-9, 182-210, 249-256, and 345-421. To address these structural gaps, the

EZH2 structure was downloaded from AlphaFold, assigned an AFDB accession code AF-Q15910-F1,

and possessed an average pLDDT score of 76.25 (Uniport ID: Q15910)[46]. Subsequently, the EZH2

alphafold model was superposed to the original PDB file (5HYN) to obtain the PRC2 complex.

Finally, EED, H3K79M, and JARID2 K116m3 proteins were deleted to keep only the EZH2-SUZ12

complex for further modelling studies. The EZH2-SUZ12 complex was pre-processed using Protein

Preparation Workflow tool in Schrödinger suite 2022[47]. Hydrogens were added, charges were

assigned, bond orders were refined, and all water molecules and non-standard residues were deleted

before proceeding. The protein backbone was minimized by employing OPLS 2005 force field.

Further residues Cys324 and Cys695 were S-nitrosylated by Vienna-PTM[48].

Molecular dynamics (MD) simulations

The charges of both wild-type and mutant complexes were neutralized by placing a total of 10 Na^+

ions at positions with high electronegative potential. Both complex and counter ions were then placed

in a pre-equilibrated cubic box of SPC/E water molecules. The periodic box of water was extended to

a distance of 10 Å from the protein complex and counter ions. Another 442 NaCl molecules were

added to the system to maintain a 150 mM salt concentration. The prepared systems were subjected to

50,000 steps of steepest descent energy minimization. The structural fluctuations and stability of the

relaxed protein complex were analyzed by time-dependent molecular dynamics (MD) simulation

studies using GROMACS 5.1.5. The particle mesh Ewald method (PME) was used for the calculation

of electrostatic interactions[49]. Periodic boundary conditions were imposed in all directions. The long-

range electrostatic interactions have been calculated without any truncation, while a 10 Å cutoff was

applied to Lennard–Jones interactions. The LINCS-like algorithm were employed to restrain
hydrogen-containing bonds only and handle long-range electrostatic interactions. SHAKE algorithm
was applied to constrain the bond involving hydrogens[50]. The temperature was controlled at 300 K
using Langevin dynamics with the collision frequency 1. A time step of 2 fs was used and the
structures were saved at every 10 ps interval for the entire duration of the MD run. Equilibrium phase
was comprised of two short 1 ns simulations in NVT and NPT ensembles, utilizing a Berendsen
thermostat and Parrinello-Rahman barostat respectively at 300K and 1 bar pressure. The unrestrained
1 μs simulation with NPT ensembles at 300 K was considered as production simulation.
741

742 **Structure Analysis**
743 Built-in features of GROMACS, such as gmx rms, rmsf, and hbond, were employed to assess the
744 root-mean-square deviation (RMSD), root-mean-square fluctuation (RMSF), and hydrogen bonds
745 (Hbond), respectively, throughout the trajectory. Visualization of the structures was performed using
746 PYMOL and VMD software[51,52].
747

Binding energy calculations using MM/PBSA:
The MM-PBSA protocol was employed to determine the effect of PTMs addition on the binding of
subunits within a chosen EZH2-SUZ12 complex. The free binding energies of each complex were
computed utilizing the equations provided below. Briefly, the provided set of equations were applied
to represent an imaginary AB dimer, where A corresponds to EZH2 and B to SUZ12.
$$\Delta Gbind = GAB - (GA + GB)$$
Where GAB is the binding free energy of the EZH2-SUZ12 complex, GA is the binding free energy
of the EZH2, and GB is the binding free energy of the SUZ12.
$\Delta\Delta G\_bind = \Delta G\_bindABnitrosylation - \Delta G\_bindABwildtype$
To evaluate the impact of incorporating post-translational modifications (PTMs) into the complex on
their binding, we compute the disparity in binding free energy (ΔΔGbind) between the S-nitrosylated
and wildtype complexes using Equation 2. The binding free energy was determined by utilizing the
1μs trajectory obtained from molecular dynamics simulations, and calculations were performed on
every 500 frames extracted from the last 500 ns of the trajectory using the g_mmpbsa tool integrated
with GROMACS[53–55].
763

Statistics Analysis
All the values are expressed as the mean ± SD. An unpaired t-test was used to determine the statistical
differences in comparisons between the two groups and one-way ANOVA with a false discovery rate
(FDR) method was used for comparison between multiple groups. GraphPad Prism software
(GraphPad Prism 9) was used to perform statistical analyses. A p-value (<0.05) was considered
statistically significant.

Data availability

771 All data can be obtained from the corresponding author on reasonable request.

Acknowledgment

This work was supported by the Core Research Grant from Science and Engineering Research Board-Department of Science and Technology, Govt. of India (CRG/2022/002209) to SM. This work was also partly supported by a Competitive Research Grant from the Department of Biotechnology, Govt. of India (BT/PR33144/MED/30/2170/2019) to SM. AS is supported by a Senior Research Fellowship from CSIR, Govt. of India (File No- 09/719(0107)/2019-EMR-I). Y.T.K., N.P.T. and S.M.T. are supported by a graduate fellowship from BITS Pilani. S.T. was supported by a graduate fellowship from BITS Pilani. S.R. is supported by a graduate fellowship from Department of Science and Technology-Innovation in Science Pursuit for Inspired Research fellowship (DST/INSPIRE/03/2019/000582). We gratefully acknowledge the technical assistance of Mr. Suman Kumar (Confocal Facility, BITS Pilani, Pilani Campus) and BITS Pilani, Pilani Campus for providing the high-performance computing (HPC) facility for simulation study.

Author contributions

AS designed and performed the experiments, analyzed the data, and wrote the first draft of the manuscript. YTK, RB, NP, SR, SC, and ST performed experiments and analyzed data. SMT, SS, and Shiabsish performed all the molecular simulation studies and wrote the associated methodology and results section of the manuscript for the same. SM secured the funding, designed the experiments, supervised the study and wrote the final draft of the manuscript.

Competing interests

The authors declare no competing interests.

References

1. Tan, J. Z., Yan, Y., Wang, X. X., Jiang, Y. & Xu, H. E. EZH2: Biology, disease, and structure-based drug discovery. *Acta Pharmacologica Sinica* vol. 35 161–174 Preprint at https://doi.org/10.1038/aps.2013.161 (2014).
2. Kuzmichev, A., Nishioka, K., Erdjument-Bromage, H., Tempst, P. & Reinberg, D. Histone methyltransferase activity associated with a human multiprotein complex containing the Enhancer of Zeste protein. *Genes Dev* **16**, 2893–2905 (2002).
3. Schlesinger, Y. *et al.* Polycomb-mediated methylation on Lys27 of histone H3 pre-marks genes for de novo methylation in cancer. *Nat Genet* **39**, 232–236 (2007).
4. Cao, R. *et al.* Role of histone H3 lysine 27 methylation in Polycomb-group silencing. *Science* **298**, 1039–1043 (2002).
5. Siamwala, J. H. *et al.* Nitric Oxide Reverses the Position of the Heart during Embryonic Development. *Int J Mol Sci* **20**, (2019).

6.   Lee, S. H. *et al.* The role of EZH1 and EZH2 in development and cancer. *BMB Rep* **55**, 595–
601 (2022).
7.   Chen, Y. H., Hung, M. C. & Li, L. Y. EZH2: a pivotal regulator in controlling cell
differentiation. *Am J Transl Res* **4**, 364 (2012).
8.   Gan, L. *et al.* Epigenetic regulation of cancer progression by EZH2: From biological insights
to therapeutic potential. *Biomark Res* **6**, 1–10 (2018).
9.   Yan, K. S. *et al.* EZH2 in Cancer Progression and Potential Application in Cancer Therapy: A
Friend or Foe? *Int J Mol Sci* **18**, (2017).
10.   Wang, X. *et al.* A covalently bound inhibitor triggers EZH 2 degradation through CHIP
⬜mediated ubiquitination . *EMBO J* **36**, 1243–1260 (2017).
11.   Ghobashi, A. H. *et al.* Activation of AKT induces EZH2-mediated β-catenin trimethylation in
colorectal cancer. *iScience* **26**, 107630 (2023).
12.   Chu, C. S. *et al.* O-GlcNAcylation regulates EZH2 protein stability and function. *Proc Natl
Acad Sci U S A* **111**, 1355–1360 (2014).
13.   Riising, E. M., Boggio, R., Chiocca, S., Helin, K. & Pasini, D. The Polycomb Repressive
Complex 2 Is a Potential Target of SUMO Modifications. *PLoS One* **3**, e2704 (2008).
14.   Li, Z. *et al.* Methylation of EZH2 by PRMT1 regulates its stability and promotes breast cancer
metastasis. *Cell Death & Differentiation 2020 27:12* **27**, 3226–3242 (2020).
15.   Alghamdi, T. A. *et al.* Histone H3 Serine 10 Phosphorylation Facilitates Endothelial
Activation in Diabetic Kidney Disease. *Diabetes* **67**, 2668–2681 (2018).
16.   Lo, P. W. *et al.* O-GlcNAcylation regulates the stability and enzymatic activity of the histone
methyltransferase EZH2. *Proc Natl Acad Sci U S A* **115**, 7302–7307 (2018).
17.   Gould, N., Doulias, P. T., Tenopoulou, M., Raju, K. & Ischiropoulos, H. Regulation of protein
function and signaling by reversible cysteine s-nitrosylation. *Journal of Biological Chemistry*
288, 26473–26479 (2013).
18.   Wu, G., Meininger, C. J., McNeal, C. J., Bazer, F. W. & Rhoads, J. M. Role of L-Arginine in
Nitric Oxide Synthesis and Health in Humans. *Adv Exp Med Biol* **1332**, 167–187 (2021).
19.   Iwakiri, Y. *et al.* Nitric oxide synthase generates nitric oxide locally to regulate
compartmentalized protein S-nitrosylation and protein trafficking. *Proc Natl Acad Sci U S A*
103, 19777–19782 (2006).
20.   Heiss, C., Rodriguez-Mateos, A. & Kelm, M. Central role of eNOS in the maintenance of
endothelial homeostasis. *Antioxid Redox Signal* **22**, 1230–1242 (2015).
21.   Tran, N. *et al.* Endothelial Nitric Oxide Synthase (eNOS) and the Cardiovascular System: in
Physiology and in Disease States. *Am J Biomed Sci Res* **15**, 153 (2022).
22.   Hemish, J., Nakaya, N., Mittal, V. & Enikolopov, G. Nitric oxide activates diverse signaling
pathways to regulate gene expression. *J Biol Chem* **278**, 42321–42329 (2003).
23.   Gunawan, M. *et al.* The methyltransferase Ezh2 controls cell adhesion and migration through
direct methylation of the extranuclear regulatory protein talin. *Nat Immunol* **16**, 505–516
(2015).
24.   Su, I. H. *et al.* Polycomb group protein ezh2 controls actin polymerization and cell signaling.
Cell **121**, 425–436 (2005).
25.   Yu, Y. L. *et al.* Smurf2-mediated degradation of EZH2 enhances neuron differentiation and
improves functional recovery after ischaemic stroke. *EMBO Mol Med* **5**, 531–547 (2013).
26.   Matrone, G. *et al.* Nuclear S-nitrosylation impacts tissue regeneration in zebrafish. *Nat
Commun* **12**, (2021).
27.   Justin, N. *et al.* Structural basis of oncogenic histone H3K27M inhibition of human polycomb
repressive complex 2. *Nature Communications 2016 7:1* **7**, 1–11 (2016).
28.   Foster, M. W., Hess, D. T. & Stamler, J. S. Protein S-nitrosylation in health and disease: a
current perspective. *Trends Mol Med* **15**, 391–404 (2009).
29.   Anwar, T. *et al.* p38-mediated phosphorylation at T367 induces EZH2 cytoplasmic
localization to promote breast cancer metastasis. *Nat Commun* **9**, (2018).
30.   Ghate, N. B. *et al.* Phosphorylation and stabilization of EZH2 by DCAF1/VprBP trigger
aberrant gene silencing in colon cancer. *Nat Commun* **14**, (2023).
31.   Yan, J. *et al.* EZH2 phosphorylation by JAK3 mediates a switch to noncanonical function in
natural killer/T-cell lymphoma. *Blood* **128**, 948–958 (2016).

32. Wan, L. *et al.* Phosphorylation of EZH2 by AMPK Suppresses PRC2 Methyltransferase
Activity and Oncogenic Function. *Mol Cell* **69**, 279-291.e5 (2018).
33. Kapil, V. *et al.* The Noncanonical Pathway for In Vivo Nitric Oxide Generation: The Nitrate-
Nitrite-Nitric Oxide Pathway. *Pharmacol Rev* **72**, 692–766 (2020).
34. Tejero, J., Shiva, S. & Gladwin, M. T. Sources of Vascular Nitric Oxide and Reactive Oxygen
Species and Their Regulation. *Physiol Rev* **99**, 311–379 (2019).
35. Francis, S. H., Busch, J. L. & Corbin, J. D. cGMP-dependent protein kinases and cGMP
phosphodiesterases in nitric oxide and cGMP action. *Pharmacol Rev* **62**, 525–563 (2010).
36. Gobeil, F. *et al.* Nitric oxide signaling via nuclearized endothelial nitric-oxide synthase
modulates expression of the immediate early genes iNOS and mPGES-1. *J Biol Chem* **281**,
16058–16067 (2006).
37. Parenti, A. *et al.* Nitric oxide is an upstream signal of vascular endothelial growth factor-
induced extracellular signal-regulated kinase1/2 activation in postcapillary endothelium. *J Biol*
Chem **273**, 4220–4226 (1998).
38. Cooke, J. P. & Losordo, D. W. Nitric oxide and angiogenesis. *Circulation* **105**, 2133–2135
(2002).
39. Förstermann, U. *et al.* Nitric oxide synthase isozymes characterization, purification, molecular
cloning, and functions. *Hypertension* (1994) doi:10.1161/01.HYP.23.6.1121.
40. Yu, J. *et al.* Endothelial nitric oxide synthase is critical for ischemic remodeling, mural cell
recruitment, and blood flow reserve. *Proc Natl Acad Sci U S A* **102**, 10999–11004 (2005).
41. Morales-Ruiz, M. *et al.* Vascular endothelial growth factor-stimulated actin reorganization and
migration of endothelial cells is regulated via the serine/threonine kinase Akt. *Circ Res* **86**,
892–896 (2000).
42. Su, I. H. *et al.* Polycomb group protein ezh2 controls actin polymerization and cell signaling.
Cell **121**, 425–436 (2005).
43. Gunawan, M. *et al.* The methyltransferase Ezh2 controls cell adhesion and migration through
direct methylation of the extranuclear regulatory protein talin. *Nat Immunol* **16**, 505–516
(2015).
44. Titelbaum, M. *et al.* Ezh2 harnesses the intranuclear actin cytoskeleton to remodel chromatin
in differentiating Th cells. *iScience* **24**, (2021).
45. Xue, Y. *et al.* GPS-SNO: computational prediction of protein S-nitrosylation sites with a
modified GPS algorithm. *PLoS One* **5**, (2010).
46. AlphaFold Protein Structure Database. https://alphafold.com/entry/Q15910..
47. Schrödinger Release 2022-2: Protein Preparation Wizard; Epik, Schrödinger, LLC, New York,
NY, 2016; Impact, Schrödinger, LLC, New York, NY, 2016; Prime, Schrödinger, LLC, New
York, NY, 2019.
48. Margreitter, C., Petrov, D. & Zagrovic, B. Vienna-PTM web server: a toolkit for MD
simulations of protein post-translational modifications. *Nucleic Acids Res* **41**, (2013).
49. Harvey, M. J. & De Fabritiis, G. An Implementation of the Smooth Particle Mesh Ewald
Method on GPU Hardware. *J Chem Theory Comput* **5**, 2371–2377 (2009).
50. Ryckaert, J. P., Ciccotti, G. & Berendsen, H. J. C. Numerical integration of the cartesian
equations of motion of a system with constraints: molecular dynamics of n-alkanes. *J Comput*
Phys **23**, 327–341 (1977).
51. The PyMOL Molecular Graphics System, Version 1.8 Schrödinger, LLC.
https://www.sciepub.com/reference/159710.
52. Humphrey, W., Dalke, A. & Schulten, K. VMD: Visual molecular dynamics. *J Mol Graph* **14**,
33–38 (1996).
53. Kumari, R., Kumar, R. & Lynn, A. g_mmpbsa--a GROMACS tool for high-throughput MM-
PBSA calculations. *J Chem Inf Model* **54**, 1951–1962 (2014).
54. Van Der Spoel, D. *et al.* GROMACS: Fast, flexible, and free. *J Comput Chem* **26**, 1701–1718
(2005).
55. Tripathi, S. M., Akash, S., Rahman, M. A. & Sundriyal, S. Identification of synthetically
tractable MERS-CoV main protease inhibitors using structure-based virtual screening and
molecular dynamics potential of mean force (PMF) calculations. *J Biomol Struct Dyn* (2023)
doi:10.1080/07391102.2023.2283780.

Figure 1. Nitric oxide caused cytosolic localization and degradation of EZH2 protein accompanied with an early reduction in H3K27me3 levels due to dissociation of SUZ12. (A-D) Immunoblots analysis of EZH2 (n=4) and H3K27me3 (n=5) proteins levels on exposure to SNP (500 μmol/L) in EA.hy926 cells for variable times. (E-F) Immunoblot analysis using lysate of rat aortic explants exposed to SNP (500 μmol/L) for 2hrs. (n=3) (G) EA.hy926 cells treated with SNP (500 μM) for 30 minutes were subjected to co-immunoprecipitation using EZH2 antibody followed by immunoblotting to show the association of EZH2 with other subunits of the PRC2 complex and histone H3. (n=3) (H) Immunoblotting for EZH2 in nuclear and cytosolic fractions from EA.hy926 cells treated with SNP (500 μmol/L for 30 & 60 mins). GAPDH and histone H3 showed the purity of cytosolic and nuclear fractions respectively. (n=3) (I) Immunofluorescence followed by confocal imaging of EA.hy926 cells exposed to 1 hr of SNP to show cytosolic translocation of EZH2 (green). F-actin (red) is stained with phalloidin-Alexa Fluor 555. DAPI staining is shown in blue (Scale bar: 10 μm). (n=3) Values represent the mean ± SD. *p < 0.05, and **p < 0.01 by unpaired t-test for two groups.

Figure 2. External nitric oxide supplementation or endogenous induction of nitric oxide producing machinery caused S-nitrosylation of EZH2 and dissociation of SUZ12 and histone H3. (A) Dot blot of protein lysates collected from EA.hy926 cells exposed to SNP (500 μmol/L) for 30 min and processed through biotin switch assay followed by immunoprecipitation with EZH2 antibody. Blots were incubated with Streptavidin-HRP followed by developing with chemiluminescence substrate for visualization. (n=3) (B) Same samples were run through SDS-PAGE followed by transfer to nitrocellulose membrane followed by incubation with Streptavidin-HRP and developing the blot with chemiluminescence substrate for visualization. (n=3) (C-D) EA.hy926 cells were exposed to either SNP (500 μmol/L, C) or bradykinin (10 μmol/L, D) for 30 minutes followed by immunoprecipitation with EZH2 antibody and further immunoblotting to show the presence of S-nitrosylation of EZH2 using Nitro-Cysteine antibody. (n=3) (E) Plasmid containing HA tagged EZH2 were transfected in HEK-293 cells followed by exposing to SNP (500 μmol/L) for 30 minutes. Cell

lysates were then immunoprecipitated using HA antibody followed by immunoblotting with
respective antibodies. (n=3)

Figure 3. GSNO exposure caused reduction in EZH2 and H3K27me3 level along with altering
the interacting partners of EZH2. (A-B) Immunoblotting for EZH2 (A) and H3K27me3 (B) in
cultured EA.hy926 cells upon exposure to GSNO (100 µmol/L) for different time points (0, 30, 60,
and 120 minutes). (n=4) (C) Venn-diagram show the percentage of common/overlapping proteins in
control and 30 minutes GSNO (100 µmol/L) exposed EA.hy926 cells as analyzed through orbitrap
mass spectrophotometer. (D) Data showing the number of total and unique interacting proteins
present in EA.hy926 cells upon exposure to GSNO (100 µmol/L) for 30 minutes. (E) Heat map based
visualization of all the uniquely interacting proteins in control and GSNO treated EA.hy926 cells. (F)
Heat map based visualization of uniquely interacting proteins in Control and GSNO treated groups
with a PEP score $\geq$ 5. Values represent the mean $\pm$ SD. *$p < 0.05$, and **$p < 0.01$ by unpaired t-test
for two groups.

Figure 4. Elevated ubiquitination of EZH2 on exposure to Nitric oxide in endothelial cells to
cause its degradation primarily through the autophagosome-lysosome pathway. (A) EA.hy926
cells exposed to SNP (500 µmol/L) for 30 minutes followed by co-immunoprecipitation to pulldown
EZH2. This pulldown samples were then subjected to immunoblotting with ubiquitin antibody to
show the presence of ubiquitinated EZH2. (B) Immunoblotting analysis of EZH2 in EA.hy926 cells
pretreated with MG-132(MG) (1 µmol/L) for 2 hrs followed by exposing to SNP (500 µmol/L) for
another 2 hrs. (n = 4) (C) Immunoblotting for EZH2 in cultured EA.hy926 cells pretreated with
Bafilomycin A1(Baf) (100 nmol/L) for 2 hours followed by exposing to SNP (500 µmol/L) for
additional 2 hrs. (n=4) (D) Immunoblotting for EZH2 protein in cultured EA.hy926 cells pretreated
with both MG-132 (1 µmol/L) and bafilomycinA1 (100 nmol/L) for 2 hrs followed by exposing to
SNP (500 µmol/L) for additional 2 hrs. (n=4) (E) Immunoblotting to show the presence of elevated
p62 in EA.hy926 cells upon treatment with bafilomycin A1 (100 nmol/L) to confirm disruption of
autophagosome-lysosome pathway. NOS inhibitor treatment reversed NO dependent reduction in
EZH2 and H3K27me3 level. (F) Immunoblotting for EZH2 with protein lysate collected from
cultured EA.hy926 cells pretreated with L-NAME (1 mmol/L) for 1 hr followed by exposing to
bradykinin(BK) (10 µmol/L) for additional 2 hrs. (n=3) (G-H) Rat aortic rings were pretreated with L-
NAME (1 mmol/L) for 1 hr followed by exposing to bradykinin (10 µmol/L) for additional 2 hrs.
Immunoblot analysis were carried out for EZH2 (G) and H2K27me3 (H) in these tissue lysates. (n=3)
(I) EZH2 (green) immunostaining along with F☐actin (Red) staining of EA.hy926 cells after
pretreating with L-NAME (1 mmol/L) for 1 hr followed by exposing to bradykinin (10 µmol/L) for

additional 2hrs. DAPI staining is shown in blue. (Scale bar: 10 µm) (n=3) Values represent the mean

± SD. *p < 0.05, and **p < 0.01 by unpaired t-test for two groups.

Figure 5. GSK-J4, a demethylase inhibitor protects H3K27me3 level in NO exposed endothelial

cells and reversed NO-dependent cell migration and gene expression changes. (A)

Immunoblotting for H3K27me3 in EA.hy926 cells pretreated with GSK-J4 (5 µmol/L) for 4 hrs

followed by challenging them with SNP (500 µmol/L) for additional 2 hours (n=4). (B) Scratch

wound healing assay to show the healing rate of EA.hy926 cells which are pretreated with GSK-J4 (5

µmol/L) for 4 hrs followed by treatment with SNP (500 µmol/L). Healing rate was followed until 24

hrs post-SNP treatment. Images were acquired using bright field microscope adapted with a camera

for phase contrast imaging. (n=4) (C) RT-qPCR analysis to measure the transcript level expression of

VEGFa, TBX20, MMP2, FGF2, KDR, TIE2, TEK, and Angiopoietin in EA.hy926 cells pretreated

with GSK-J4 (5 µmol/L) for 4 hrs followed by challenging them with SNP (500 µmol/L) for

additional 4 hrs (n=4). Values represent the mean ± SD. *p < 0.05, **p < 0.01, and ***p < 0.001 by

unpaired t-test for two groups.

1006

Figure 6. Point mutated EZH2 protein responded differentially to nitric oxide in the context of

its effects on the stability and catalytic activity of EZH2. (A) Biotin Switch assay followed by dot-

blot and Ponceau staining of HA immunoprecipitated cell lysates of HEK-293 cells overexpressing

WT and mutant EZH2 (having point mutations at cysteines 260, 329, and 700) followed by

challenging with SNP (500 µmol/L) for 2 hrs to show the levels of S-nitrosylated HA tagged EZH2.

(n=3) (B-C) Immunoblotting for HA (EZH2) (B) and H3K27me3 (C) in cell lysates collected from

HEK-293 cells containing wild type or mutated versions of EZH2 (C260S, C329S, C700S) and were

exposed to SNP (500 µmol/L) for 2 hrs. (n=4) (D) Immunoblotting experiments to show the levels of

HA (EZH2) and H3K27me3 in HEK-293 cells transfected with wild-type and the double mutant

version of EZH2 (created by inserting point mutations to convert cysteine to serine at positions C329

and C700) and exposed to SNP (500 µmol/L) for 2 hrs. (n=4). (E) Immunostaining for HA (EZH2)

(Red) along with F☐actin (green) staining in HEK-293 cells to localize HA tagged EZH2 in wild-type

and mutated versions, following their exposure to SNP (500 µmol/L) for 2 hrs. DAPI staining is

shown in blue. (Scale bar: 10 µm) (n=3) Values represent the mean ± SD. ***p < 0.001 by unpaired t-

test for two groups.

1022

1023   **Figure 7. Performing Molecular Dynamic (MD) simulation studies through GROMACS 5.1.5**

1024   **version using WT and S-nitrosylated EZH2 at C324 and C695.** (A-C) Structural illustration of the

1025   initial structural alignment of the wildtype and S-nitrosylated EZH2 protein of the EZH2-SUZ12

1026   complex (A). The magenta and green color represents the EZH2 chain of both WT and S-nitrosylated

version, while the cyan and red color indicates the SUZ12 chain. It also provides a close-up view of S-nitrosylation at position C324 of EZH2 in the S-nitrosylated protein (B) and also zooms in on the S-nitrosylation at position C695 of EZH2 in the S-nitrosylated protein (C). (D-E) Backbone RMSD of whole EZH2 WT and EZH2 S-nitrosylated (D). Backbone RMSD of EZH2 WT and EZH2 S-nitrosylated specifically within the residue ranges between 112-121 of SAL (E). Backbone RMSD of residues in SUZ12 ranging between 584-588 (F). (H-K) Structural illustration of EZH2-SUZ12 complex near the SAL (112-121) domain of EZH2 in both EZH2 WT and EZH2 S-nitrosylated forms at 100 ns and 800 ns of simulation. Structural data reflects the distance between unique residues of EZH2 (LEU116 and GLN117) and SUZ12 (ASP585 and LYS587) to reflect potential for hydrogen bonds which were lost at 800 ns only in EZH2-SUZ12 complex having S-nitrosylated form of EZH2 at C324 and C695.

Figure 8. Schematic depicting the effect of EZH2 S-nitrosylation on its stability, translocation and catalytic activity of Polycomb Repressive Complex-2. PRC-2 consists mainly of EZH2, SUZ12, and EED, residing primarily in the nucleus. The major subunit of PRC2 i.e. histone methyltransferase EZH2 localizes primarily in the nucleus and is responsible for deposition and maintenance of the levels of H3K27me3. These cells also have a limited cell migration capacity. On exposure of these cells to NO, S-Nitrosylation of specific cysteine residues on EZH2 occur leading to a decline in their stability which is marked by reduction in EZH2 protein levels along with dissociation of SUZ12 from EZH2 bound PRC2 complex. Upon disassembly, S- Nitrosylated EZH2 translocate from the nucleus to the cytosol where it undergoes ubiquitination followed by its degradation primarily through the autophagosome-lysosome pathway. Additionally, there is a significant effect on the catalytic activity of EZH2, which is shown by reduction in the deposition of the H3K27me3 marks on histone H3. NO dependent S-nitrosylation of EZH2 also causes a spike in the migratory activities of EC due to EZH2 recruitment to the F-actin and likely contributing towards remodeling prior to degradation. The schematic diagram was prepared using BioRender (Agreement number: WB26FEOYBU).

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7

Figure 8