A *Mycobacterium tuberculosis* secreted virulence factor disrupts host snRNP biogenesis

Komal Chauhan[1], Nishat Passricha[1], Yogita Kapoor[2], Dipanwita Datta[1], Naresh Arora[1], Mrinmoy Das[1], Kavya Rao[1], Lakshyaveer Singh[1], Ravi Dutt[1], Raman Deep Sharma[1], Binayak Sarkar[3], Basanti Malakkar[3], Haroon Kalam[1], Prince Saini[4], Lalita Mehra[5], Prasenjit Das[5], Vineet Ahuja[6], Amit Singhal[7,8], Vinay Nandicoori[2, 3] and Dhiraj Kumar[1]*

[1]Cellular Immunology Group, International Centre for Genetic Engineering and Biotechnology, Aruna Asaf Ali Marg, New Delhi-110067, India
[2]Centre for Cellular and Molecular Biology, Hyderabad, India
[3]National Institute of Immunology, Aruna Asaf Ali Marg, New Delhi-110067, India
[4]Department of Biological Sciences, Indian Institute of Science Education and Research Mohali, Punjab 140306, India
[5]Department of Pathology, All India Institute of Medical Sciences, New Delhi-110012, India
[6]Department of Gastroenterology, All India Institute of Medical Sciences, New Delhi-110012, India
[7]A*STAR Infectious Diseases Labs (A*STAR ID labs), Agency for Science, Technology and Research (A*STAR), Singapore, Singapore.
[8]Singapore Immunology Network (SIgN), A*STAR, Singapore, Singapore.

Running title: Regulation of host RNA splicing by Mtb

*Correspondence- dhiraj@icgeb.res.in

Abstract

Mycobacterium tuberculosis (Mtb) infection of human macrophages triggers extensive

dysregulation of host alternative splicing (AS). Here, we present an unexpected mechanism

exploited by the Mtb to achieve the same. We identify several bacterial secretory proteins

that interact with the host splicing factors (SFs) and alter select RNA splicing events *in vitro*,

in cell and during *ex vivo* infections. We show bacterial proteins disrupt AS by impairing

Small nuclear ribonucleoprotein (snRNP) biogenesis, primarily by interacting with and

sequestering U5snRNA and SNRPF, key spliceosome constituents. These interactions are

driven, in part, by one of the identified Mtb proteins, Hsr1, whose access to the host cytosol

is ESX1-dependent. The Mtb*Δhsr1* is unable to interact and sequester U5snRNA or SNRPF

and fails to alter host AS. Consequently, Mtb*Δhsr1* shows compromised survival and

pathogenesis in macrophages and mice. Lungs of infected mice show distinct SNRPF

staining, which is hsr1 dependent. Similar distinctive SNRPF staining is also noted in human

tuberculous granulomas. Moreover, monocytes from tuberculosis patients show substantial

splicing dysregulation, indicating splicing defects as a key determinant behind systemic

tuberculosis pathology. We propose mycobacterial exploitation of the evolutionarily

conserved mechanism of snRNP biogenesis, aided its adaptation for human pathogenesis and

provides a unique therapeutic opportunity.

Introduction

Tuberculosis (TB) is one of the leading causes of human mortality due to infectious diseases.[1] *Mycobacterium tuberculosis* (Mtb), the causative organism, is known to alter a variety of host cellular processes that eventually help establish the infection and cause disease. Thus, cellular mechanisms like signaling, autophagy, apoptosis, inflammation etc., have been extensively explored for potential host-directed therapy (HDT) against tuberculosis [2-5]. Mtb infection also alters host gene expression, which determines the long-term host responses against infection. In fact, the pathogen regulates host gene expression at multiple tiers, including epigenetic, transcriptional and post-transcriptional stages [6-9]. Previously, reports from our group and other showed that infection of human macrophages with Mtb triggers a massive alteration in the pattern of host RNA splicing[10,11]. Altered RNA splicing can impact the target proteins function, localization, stability and stoichimetry[12]. We reported increased instances of transcripts which either do not get translated or are substrate of a process called nonsense-mediated decay[13], were upregulated as a result of alternative splicing in the infected macrophages[8,11,13]. Curiously, genes typically belonging to functional classes that are activated upon infection are impacted by AS[8,9,12]. This targeted blunting of host response by bacterial infection favours the bacteria[11]. Incidentally, infection of macrophages with *Salmonella typhimurium* or *Lysteria monocytogenes* were also reported to alter isoform usage, in addition to a widespread shortening of 3'UTR[14]. The macrophage transcriptome also appears to get spliced differently during other inflammatory conditions [15,16]. However, the mechanism determining differential isoform usage in most of these conditions is unclear. Given the fundamental function of RNA splicing in shaping the host proteome and its perturbation by Mtb during macrophage infections, insights into the mechanism of infection-dependent alternative RNA is critical to understand molecular details of tuberculosis pathogenesis[9]

Host RNA splicing is a complex and fundamental physiological process in all eukaryotes.

RNA splicing is performed by a multimolecular and dynamic spliceosome complex, which

employs a large number of proteins to ensure the accuracy of splicing to form mature

mRNAs [17-19]. One of the key components is U-rich U snRNAs, which form protein-RNA

complexes known as small nuclear ribonucleoproteins (snRNPs)[20-22]. The snRNPs are

recruited stepwise onto the pre-mRNA exon-intron boundaries to form a functional

spliceosome, join the adjoining exons, and splice out the intervening intron [18,23].

Like other bacterial pathogens, Mtb also secretes virulence factors to alter host physiology [24-

29]. Since mycobacterial secretory proteins in the infected macrophages have access to both

cytosol and the nucleus [24-26,30], we asked whether secreted Mtb proteins could impact

spliceosome complex assembly and, consequently, host RNA splicing. This study extensively

maps the interactions between mycobacterial secretory proteins and host proteins involved in

RNA splicing. We identify the Mtb proteins capable of altering the assembly of spliceosome

complex by exploiting an evolutionarily conserved quality control mechanism of RNA

splicing in the eukaryotes.

Results

A targeted high-throughput yeast-2-hybrid (Y2H) screen to identify *Mtb*-human protein interactions

Eukaryotic RNA splicing is a highly conserved, multi-step process involving a large number of proteins and protein-RNA complexes (herein referred to as splicing factors or SF). Our hypothesis was to test whether Mtb secretory proteins could directly interact with the SFs to interfere in the host RNA splicing. For this, we performed a high-throughput one-to-one yeast-2-hybrid assay. The bait library (binding domain or BD) comprised of mycobacterial secreted proteins while the prey library (activation domain or AD) comprised the protein-coding isoforms of genes belonging to functional classes "RNA splicing" and "Spliceosome function", making it a unique custom library of human transcripts (Table S1). Briefly, our BD library contained 384 *Mtb* genes and also included 37 putative negative controls (Fig. S1A-B Table S1 and S2)[24,27-29,31].

Four independent mating reactions were performed between 384 BD and 162 AD clones to obtain high-quality interaction data, leading to 2,48,832 mating reactions. The schematic of the Y2H strategy followed for the study is shown in Fig. 1A. In total, 123 Mtb proteins showed interactions with one or more SFs, together consisting of 439 incidences of interactions, including 121 autoactivation incidences (Table S3). The BD clones showing autoactivation were excluded from the list of interactors. A few negative control list proteins also showed interactions, including 56 autoactivation incidences (Table S3).

The interactome between *Mtb* secretory proteins and host SFs, as observed by the Y2H screen, is shown in Fig. 1B (Table S4). The interactions spanned across all classes of host genes involved in RNA splicing, including RNA polymerases, Sm proteins, SMN complex proteins, U2 snRNPs, LSm proteins, HnRNPs, U4/U6.U5 snRNP tricomplex proteins and non-snRNP proteins (Fig. 1C). The Venn-Diagram shows the number of Mtb proteins

interacting with the host SFs (Fig. 1C). Overall, host proteins from RNA Pol II subunits, Sm

Core, catalytic step II factors, HnRNPs and SMN complex (Fig. S1C) and Mtb proteins from

information pathway, virulence factors and conserved hypothetical proteins functional class

showed most interactions (Fig. S1D). To note, there were many instances where different

proteins of similar functional sub-class shared their interacting partner with *Mtb* (Fig. 1D).

Interaction of Mtb proteins with the splice variants of human proteins

Since the human library consisted of all protein-coding isoforms of the corresponding genes,

among the 162 AD clones that were screened, a total of eighteen genes were represented by

more than one transcript (Fig. S2A). Among them POLR2H, SYF2, SNRPD2, LSm7,

RBMX, LUC7L2, DCAF4, PPIH and U2AF1 showed interaction with Mtb protein in Y2H

(Fig. S2A). The analysis of interaction patterns observed for the different transcripts of the

same gene revealed unique insights (Fig. S2B). It shows the gain or loss of interacting

partners as a consequence of the gain or loss of exons (Fig. S2B). Most strikingly, truncated

isoforms for three genes showed an unusually high number of interactions with the Mtb

proteins (Fig. S2B). In our earlier report, we noted a significant increase in the expression of

truncated protein-coding isoforms of many host genes, which appeared to help Mtb. Possibly,

the truncated spliced variants take confirmations that allow it to interact with more partners.

However, this remains a preliminary observation and needs a more focused study to

specifically understand the underlying mechanisms and consequences.

***In vitro* splicing assays point to a direct role of *Mtb* proteins in regulating host-RNA**

splicing.

We asked whether the mycobacterial proteins interacting with the host SFs, either through

Y2H or MS approach, could also alter host RNA splicing. For functional assays, we selected

transcripts from a handful of genes like *ACSL1*, *ACADVL*, *DDX11*, *DDX42* and *NFYA*, each

representing a specific exon exclusion event upon H37Rv infection (Fig 1E)[11]. These

transcripts were specifically upregulated in H37Rv infected macrophages but not in H37Ra

infected ones[11]. We designed primers to screen for the specific splicing events (Fig. S3A).

The selected transcripts from the above genes showed an exon exclusion event specifically

upregulated upon H37Rv infection (Fig. S3B). The increase was not due to an increased

transcriptional event since the exon inclusion primers did not show any change in expression

(Fig. S3C). Moreover, using primers that spanned the junction between flanking exons after

excluding the middle exon, we reconfirmed that the increased expression of the transcripts

noted above was indeed due to an increased exon exclusion event (Fig. S3D). The first set of

functional assays involved *in vitro* splicing assays. For *in vitro* splicing assays, we generated

minigene constructs for four specific isoforms for the genes *ACSL1*, *ACADVL*, *DDX11* and

DDX42. The schematic details for *in vitro* splicing assay are shown in Fig. 1E and Fig.S3E.

We verified the substrate RNAs produced via *in vitro* transcribing the minigene constructs,

undergo splicing and form the desired product in the presence of splicing-competent nuclear

extract from THP-1 macrophages (Fig. S3F).

Next, to test whether Mtb proteins could directly interfere with RNA splicing, we purified

several Mtb proteins identified in the Y2H screen and spliceosome complex pulldown assays

(Fig. 1, Fig. S1, Fig. S3G) and used them in independent in *vitro* splicing assays. Most Mtb

proteins did not have any impact on the inclusion or exclusion of the test exon in the *in vitro*

assays, a few mycobacterial proteins appeared to cause the exclusion of test exon in the case

of ACADVL minigene substrate (Fig. 1F). Thus, Mtb secretory proteins could indeed

selectively alter the choice of inclusion/exclusion of specific exons. While the *in vitro* assay

provided evidence for the possible role of Mtb proteins in altering the host RNA splicing, it

did not capture the mechanism for global impact on RNA splicing as reported earlier and

which was our major interest [11].

***Mycobacterium smegmatis*-based knock-in system to verify the role of *Mtb* proteins in regulating host RNA splicing**

As an alternate approach, we recapitulated the infection-like scenario, by generating 19 non-pathogenic *M. smegmatis* (*Msmeg*) lines inducibly expressing Mtb proteins identified through Y2H (*Rv1390, Rv0170, Rv0639, Rv0780, Rv0288, Rv1094, Rv3864, Rv1566c, Rv1643, Rv3849* and *Rv1435c*) (Fig S4A). These were sub-cloned under the IVN-inducible promotor to develop corresponding knock-in *Msmeg::Rv* lines (Fig. 2A). We next used each of the *Msmeg::Rv* lines to infect THP-1 macrophages and analyzed specific RNA splicing events of a set of target genes through RT-PCR using the exclusion primers (Fig. S3A), against transcripts from *ACSL1*, *ACADVL*, *DDX11*, *DDX42* and *NFYA* (Fig. S4A). After the first biological experiment, out of 11 genes screened 6 genes (Rv1566c, Rv1643, RV1435c, Rv3849, Rv1390) showed IVN-induction dependent AS event in one or more of the five test exons analysed (Fig. S4A). Out of these Rv1390 did not show consistent patterns during the subsequent biological replicates (Fig. S4B). The *Msmeg::Rv* lines expressing *Rv1566c*, *Rv1435c* or *Rv1643* showed significant exon skipping events for one or more test exons analyzed across three biological replicates (Fig. 2B). Another *Msmeg::Rv* line, expressing Rv3849, also showed consistent AS events across three biological replicates (Fig. S4B); unlike Rv1566c, Rv1435c and Rv1643, we could not confirm its secretion from the *Msmeg::Rv* line and therefore Rv3849 was not taken forward for further studies (Fig. S4C). Most importantly, we observed insignificant changes in the AS for the target transcripts when *Msmeg* alone or upon IVN-induction were used for infection (Fig. S4D). Thus, with the limitation of using only five test exons, each Mtb protein expressed in *Msmeg* impacted the splicing of a unique and overlapping subset of test exons analyzed (Fig. 2B). Despite the limitations, the results from this surrogate approach suggests that select Mtb proteins were able to alter specific host RNA splicing events during infection in THP-1 macrophages.

In-cell splicing reporter assays illustrate the direct role of *Mtb* proteins in regulating host-RNA splicing.

We next developed an *in-cell* splicing reporter assay to test the three Mtb proteins identified in the results in Fig. 2B. The reporters included a constitutive pRint vector and a set of test pGint vector[32]; the latter was modified to incorporate the custom exon-intron boundaries from our list of splicing events (Fig. 2C). In HEK293T cells, stably expressing pRint, pGint constructs (pGint_ACADVL, pGint_DDX11, pGint_DDX42, pGint_ACSL1 and pGint_NFYA) were co-transfected with plasmids encoding *Rv1566c, Rv1435c and Rv1643* respectively, in separate experiments (Fig. 2D). We used a plasmid encoding *Rv1252c* as a non-interacting control (NIC) for this experiment. We measured the %pGint⁺ cells (GFP⁺) in the pRint⁺ gated population (Fig. 2D). As shown in Figure 2E, transfecting NIC marginally but consistently increased the splicing event for *ACADVL* and *ACSL1*. However, transfecting *Rv1566c*, *Rv1435c,* and *Rv1643* further increased the exon exclusion event in *ACADVL* but not in *ACSL1* (Fig. 2E). NIC transfection had no impact on the exon exclusion event for *DDX11*, *DDX42* and *NFYA* (Fig. 2E). In contrast, expressing *Rv1566c*, *Rv1435c* or *Rv1643* led to a significant increase in the exon exclusion event for *DDX11*, *DDX42* and *NFYA* (Fig. 2E). We also analyzed pRint expression in pGint⁺-population across all the conditions, which did not show any significant differences, suggesting no impact on the basal splicing event as a result of ectopically expressing Mtb proteins in 293T cells (Fig. S5A). Taken together, through the validation experiments in figures 1-2, *in vitro* assays, *ex vivo* infection assays, and *in-cell* reporter-based assays, we demonstrate that the Mtb proteins identified in this study were capable of systemically altering host RNA splicing in a targeted fashion.

Verification of physical interaction between Mtb proteins and host splicing factors

The three Mtb proteins identified above (Rv1435c, Rv1566c and Rv1643) were next verified for their physical interaction with the human SFs in a mammalian system. Mtb proteins

(Flag-tagged) and human SFs (3X-HA-tagged) were co-transfected in HEK293T cells, and

co-immunoprecipitation was performed using anti-HA and anti-FLAG antibodies,

respectively. The Y2H interaction combinations thus verified included PRPF18-Rv1435c,

SNRPD2-Rv1435c, GEMIN3-Rv1435c, POLR2H-Rv1566c, and SNRPD2-Rv1643 (Fig. 2F).

The combination of SNRPD2-Rv1566c, while showed interactions in the Y2H screen, did not

show interaction in the co-IP experiment (Fig. S5B). We also included a few combinations

that did not show interaction in the Y2H screen as the additional negative control. Thus, co-

transfecting *SNRPF* with *Rv1566c* in HEK293T cells did not result in their co-

immunoprecipitation (Fig. S5B). On the other hand, co-transfecting *SNRPF* with *Rv1435c*

showed only limited interaction (Fig. S5B). While Rv1435c could pulldown SNRPF, the

reverse was not true (Fig. S5B). For a couple of Rv1435c interactions, we also verified the

interaction by exchanging the BD and AD clones in Y2H (Fig. S5C). Overall, from the above

set of results, we conclude that the Mtb proteins, which were able to alter targeted host-RNA

splicing in the functional assays, also showed physical interactions with the host splicing

factors in the mammalian system. Finally, each of these three proteins was able to alter exon

inclusion for ACADVL test transcript in the *in vitro* splicing assay (Fig. S5D).

Mtb infection impacts levels and distribution of splicing-associated proteins in

macrophages.

The results above together show that select Mtb proteins could interact with host SFs and

alter the host RNA splicing. However, these results did not provide any insights into the

possible mechanisms. We next analyzed *in vivo* assembled spliceosome complexes from

Mtb-infected macrophages by label-free quantitative proteomics. The spliceosome complexes

were pulled down using anti-SNRPB (Y12) antibody from nuclear and cytosolic lysates of

the infected and uninfected macrophages to analyse the abundance and distribution of host

spliceosome proteins (Table S5). In the nuclear fractions of the Mtb-infected cells, there was

a significant depletion of SR proteins like SRSF3, SRSF7, Sm Core proteins like SNRPC,

SNRPB, SNRPN and SNRPD2 and some HNRNP proteins, as compared to the uninfected

ones (Fig. 3A). Other proteins like RPL4, SF3B2 and SYNCRIP followed a reversed pattern

in the infected cells, getting more enriched in the cytosol unlike the uninfected cells (Fig.

3A). Similarly, proteins like EIF5A and EIF5AL1 showed increased nuclear enrichment in

the infected macrophages, in contrast to the uninfected cells (Fig. 3A). With the limited

example of proteins identified in the immunoprecipitated spliceosome complexes, it was

evident that a diverse set of SFs were either getting depleted or mislocalized in the Mtb-

infected macrophages.

Mtb infection dramatically alters the levels and distribution of proteins involved in

RNA splicing.

To specifically investigate the levels and distribution of host SFs in Mtb-infected cells, we

systematically analyzed a representative set of SFs involved at different stages of RNA

splicing (Fig. 3B). We monitored three different parameters- MFI/cell, foci/cell and the

cytosolic to the nuclear ratio (Cyt/Nuc) for these proteins. (Fig. 3C-H). We also included

some transcription-related proteins for this analysis, since splicing begins while the RNA is

still being transcribed [33,34]. Both POLR2G and RBP8 (transcription related proteins) showed,

unexpectedly, increased nuclear retention (S6A). The U1 snRNP protein SF1 showed

increased MFI and increased cytosolic retention upon infection (Fig. 3C). The U5 snRNP

component DDX23 showed a decrease in MFI upon infection (Fig. 3D). The SMN complex

proteins GEMIN8 and GEMIN3 showed differential MFI and foci/cell (Fig. 3E-F). A couple

of SFs like PRPF4 (U4/U5.U6 associated) and SNRP200 (U5 associated) did not show

alteration in any of the three parameters analyzed (Fig. S6B). Among the Sm proteins, LSM7

showed decreased MFI/cell in the infected macrophages and SNRPF showed an increased

cytosolic retention (Fig. 3G-H). However, a more perplexing observation was the direct

interaction of SNRPF with Mtb phagosomes (Fig. 3H). It also appeared that SNRPF was

getting sequestered in the cytoplasm due to its strong interaction with Mtb phagosomes (Fig.

3I). The results in Figure 3 together indicated that the components of the RNA splicing

machinery, specifically snRNP complex biogenesis, could be dysregulated in the Mtb-

infected macrophages.

U5 snRNA directly interacts with *Mtb* phagosomes

The nuclear-cytosolic transport of U-rich snRNAs is fundamental to their assembly into U

snRNPs, which, together with different snRNPs, form the spliceosome complex[35]. The

snRNP complex assembly requires cytosolic transport of the snRNAs, which then interact

with SMN complex and Sm Core proteins to form snRNP before entering back into the

nucleus to form spliceosome complexes (Fig. 4A). Since SNRPF, a part of the pentameric

partial Sm ring, which first recognizes the Sm-sites present on snRNAs before the assembly

of the full heptameric ring takes place with the help of the SMN complex [20], was increasingly

retained in the cytoplasm of Mtb-infected macrophages, we anticipated that some snRNAs

could also show dysregulation in the infected macrophages. We analyzed the U2 and U5

snRNAs by fluorescence in situ hybridization (FISH) in the Mtb-infected macrophages. The

U2 FISH did not show any aberrant pattern for the parameters- MFI/cell, foci/cell and

Cyt/Nuc ratio, except for an increased MFI/cell at 24 hours post-infection (hpi, Fig. 4B).

Unlike U2 FISH, U5 FISH showed remarkable patterns in the Mtb-infected macrophages

(Fig. 4C). There was a decrease in MFI/cell and foci/cell at 24 hpi; however, a substantial

increase was noted in all three parameters at 48 hpi (Fig. 4C). At 48 hpi, there was significant

cytosolic retention of U5snRNA (Fig. 4C). More surprisingly, as in the case of SNRPF, we

noted direct interaction of U5snRNAs with Mtb in the cytosol (Fig. 4D), which quantitatively

increased from 24 to 48 hpi (Fig. 4D). The direct interaction of Mtb with SNRPF and

U5snRNA suggests binding and cytosolic sequestration of these specific splicing components

by Mtb. This would also mean stoichiometric disbalance among the components of snRNPs

in the infected macrophages, a condition which is known to cause widespread splicing defects

[9,36]. Interestingly, we verified that the expression of both U5 snRNA (by qPCR) and SNRPF

(by immunoblot) consistently increased post-infection, which could be a compensatory

mechanism to sustain the snRNP biogenesis in the Mtb-infected cells (Fig. 4E-F). Therefore,

disruption of snRNP biogenesis appears to serve as the major mechanism that triggers large-

scale splicing alterations in the Mtb-infected macrophages.

_Rv1435c_ deletion strain of Mtb fails to alter the splicing of target transcripts.

We next asked whether the Mtb proteins identified in this study were directly responsible for

dysregulating snRNP biogenesis. To test the above, we deleted one of the three identified

Mtb proteins, _Rv1435c,_ in H37Rv using recombineering (Fig. S6C)[37]. This protein is

annotated as a secretory protein and showed positive results in most of the interaction and

splicing assays. We confirmed the deletion of _Rv1435c_ by PCR (Fig. S6D) and also generated

a complemented strain _ΔRv1435c::WT_ using an integrative vector (see Methods). We first

verified the impact of _Rv1435c_ deletion on the splicing of five target transcripts used in this

study. All five target transcripts, _ACADVL, DDX11, DDX42, ACSL1_ and _NFYA_, showed

significantly reduced test exon exclusion events for these transcripts upon infection with the

H37Rv_ΔRv1435c_ strain compared with the WT Mtb (Fig. 5A). Moreover, in the cells infected

with H37Rv_ΔRv1435c_::_WT_, the exon exclusion event was restored to the same extent or

higher than the WT infected cells (Fig. 5A). These results conclusively show that Rv1435c

can directly alter host RNA splicing during Mtb infection. We designated the Mtb protein

Rv1435c as the host splicing regulator 1 or Hsr1.

H37Rv_Δhsr1_ alters infection-induced global alternative splicing.

We next wanted to see whether the loss of _hsr1_ causes global recovery in the infection-

induced RNA splicing events. We performed RNAseq analysis from the THP-1

macrophages, uninfected or infected with WT or H37Rv*Δhsr1* strains and analyzed the

relative abundance of different splice variants of the protein-coding genes. The clustergram in

figure 5B shows that H37Rv infection causes increased expression of a large number of

transcripts, most of which were effectively restored in H37Rv*Δhsr1* infected cells (Fig. 5B).

Interestingly, H37Rv*Δhsr1*-infected cells show increased expression of another unique set of

transcripts, likely represented those splicing events, which were possibly inhibited during WT

H37Rv infection (Fig. 5B). Curiously, a large number of transcripts showing upregulation in

the Mtb-infected macrophages actually represented NMD substrates or processed transcripts

(Table S6), which usually don't form a functional protein. Expression of splice variants that

are either truncated in length or do not code for a functional protein is one of the ways

through which infection-induced alternative splicing helps Mtb evade host responses [11]. The

results here show a direct role of Mtb protein Hsr1 in manipulating host RNA splicing events

favoring the pathogen.

Mtb *hsr1* directly perturbs snRNP biogenesis by sequestering U5snRNAs to alter host

RNA splicing.

We next investigated how the loss of Mtb *hsr1* contributed to the reversal in the splicing

events in the macrophages. We analyzed U5snRNA interaction and distribution in Mtb*Δhsr1*

infected cells by FISH. In concurrence with the altered RNA splicing events reported above,

Mtb*Δhsr1* infected cells did not show cytosolic retention of U5snRNA (Fig. 5C). On the

similar line, the interaction between Mtb and U5snRNA was almost entirely lost in Mtb*Δhsr1*

infected cells (Fig. 5C). Like U5 snRNA, even SNRPF interaction with Mtb*Δhsr1* was

significantly lower than that observed with the WT Mtb (Fig. 5D). The effect of *hsr1* deletion

on the regulation of U5snRNA and SNRPF was specific since we did not observe any

difference in any of the parameters analysed for U2snRNA (Fig. S6E). These results imply

that Mtb Hsr1 perturbs host snRNP biogenesis by sequestering the U5snRNA and SNRPF in

the cytosol, leading to widespread RNA splicing defects.

Hsr1-mediated impact on host RNA splicing is ESX1-dependent

While we did observe secretion of Hsr1 from Mtb as well as when expressed in *Msmeg*, it

was not evident whether it also gets secreted inside the macrophages during infection. We

generated anti-sera against Hsr1 in mice to check its secretion and localization during

infection in macrophages. Interestingly, we could observe significant Hsr1 secretion in the

cytosol of THP-1 macrophage infected with the WT or *Δhsr1::WT* Mtb (Fig. 5E). The Hsr1

signal was almost completely lost in the cells infected with *Δhsr1* (Fig. 5E). Curiously, when

we used the *ΔCE* Mtb strain (lacking both *Esx1*/ESAT6 and *Esx2*/CFP10), we could not see

any cytosolic staining for Hsr1. The Hsr1 staining in THP-1 cytosol was restored when the

complemented strain *ΔCE::WT* was used for infection (Fig. 5E). Expectedly, THP-1

macrophages infected with Mtb strains lacking the *esx1* system failed to alter RNA splicing

for the target genes like ACSL1 and ACADVL (Fig. 5F). Consistent with the above

observation, the Mtb *ΔCE* strain could not alter SNRPF cytosolic/nuclear distribution (Fig.

S6F). These results suggest that the Hsr1-mediated impact on host RNA splicing reported in

this study is ESX1-dependent.

Evolutionary perspective of hsr1 in mycobacterial virulence

On UniProtKB, Rv1435c is characterized as a conserved proline, glycine, valine-rich secreted

protein. There are several orthologs present in diverse mycobacterial species, like *M.*

marinum, where it is classified as a conserved hypothetical protein; in opportunistic

pathogenic species like *M. paratuberculosis, M. avium,* and *M. helveticum,* the identified

orthologue codes for secreted beta-xylosidase enzyme (Fig. S7A-B). *In M. smegmatis*, the

orthologue, with very poor similarity, encodes a translation initiation factor (IF-2). In a few

pathogenic species like *M. lacus and M. heidelbergense,* it codes for a beta-xylosidase (Fig.

S7A-B). Thus, most *hsr1* orthologues in the genus *Mycobacterium* encode a secreted protein,

and their functions are better characterized in non-pathogenic or opportunistic pathogens. It is

likely that the ability of hsr1 to alter host splicing events was likely a gain-of-function event

during the course of Mtb co-evolution as a human pathogen.

Mtb Hsr1 is a virulence factor.

Since Mtb*Δhsr1* infection could not alter RNA splicing events as in WT Mtb-infected cells,

we wondered whether loss of *hsr1* also impacts bacterial pathogenicity. The Mtb*Δhsr1*

showed reduced intra-macrophage survival compared to the WT Mtb (Fig. 6A, S8A). The

Mtb*Δhsr1* strain, as such, did not have a growth defect *in vitro*. Rather, it grew faster than the

WT cells in liquid culture which was consistent with a decrease in the bacterial cell length

(Fig. S8B-C). Curiously, *Δhsr1::WT* grew inside the THP-1 macrophages even better than the

WT Mtb, pointing to Hsr1 being an important virulence factor (Fig. 6A, Fig. S8A).

Interestingly, *Δhsr1* showed increased targeting to the lysosomes and also the cell had higher

cathepsin D staining (Fig. S8D). Since Mtb*Δhsr1* was unable to bind and sequester SNRPF,

we wondered whether overexpressing SNRPF during WT Mtb infection could have an impact

on Mtb CFU. Overexpressing SNRPF in human macrophages caused a relative decline in

cytosolic retention of SNRPF and a reduced intracellular CFU of WT Mtb (Fig. S8E),

indicating that SNRPF sequestration by Hsr1 serves as the virulence mechanism of Hsr1. On

the contrary, overexpression of SNRPD2 did not have any impact on Mtb CFU (Fig. S8F).

Next, to assess the role of Hsr1 in bacterial virulence *in vivo*, we infected BALB/c mice with

WT, Mtb*Δhsr1* and Mtb*Δhsr1::WT* strains and analyzed bacterial CFU at two and four weeks

post-infection. At both two- and four-weeks post-infection, Mtb*Δhsr1* had fewer CFUs in the

lungs of the animals (Fig. 6B). Unlike in THP-1 macrophages, the complemented strain

Mtb*Δhsr1::WT* was not able to rescue the bacterial CFU (Fig. 6B). At both 2- and 4- weeks

post-infection, analysis of a panel of eight cytokines in the lung homogenates showed an

increase in cytokine levels in WT infected animals, which, to some extent, were diminished

in the Mtb*Δhsr1* infected mice (Fig. S8G). The lung homogenates from mice infected with

Mtb*Δhsr1::WT* did not revert the expression to the WT levels (Fig. S8G). The lung tissue

sections from mice that were infected for four weeks showed strong SNRPF staining in case

of WT and *Δhsr1::WT* Mtb infection (Fig. 6C). The SNRPF staining was strikingly lower in

uninfected mice or those infected with Mtb*Δhsr1* (Fig. 6C). Remarkably, SNRPF staining in

the WT or *Δhsr1::WT* Mtb infected mice appeared mostly cytosolic in nature. Together, the

ex-vivo assays in macrophages and *in vivo* assays corroborate Mtb Hsr1 as a virulence factor

which helps its survival in the animals, largely by sequestering the host SNRPF.

Macrophages in human TB granulomas show altered snRNP complex component

localization.

Specific SNRPF cytosolic retention in THP1 macrophage and in the lungs of infected mice

prompted us to explore the clinically relevant human samples. Tissue sections from the

confirmed human TB granulomas belonging to abdominal TB cases were stained with

SNRPF. To our surprise, all granuloma lesions showed strong SNRPF staining (Fig. 6D,

S9A). Moreover, it was evident that there were more cytosolic SNRPF than in the nucleus

(Fig. 6D). Lesions that were highly necrotic showed even higher SNRPF levels (Fig. 6D).

Multinucleated giant cells, a hallmark of TB granulomas, showed intense staining for

cytosolic SNRPF (Fig. S9A). Unlike the granuloma lesions, in healthy tissue sections,

SNRPF staining was not remarkable (Fig. S9B). Finally, we also stained for another sm Core

protein, i.e., SNRPD2, in the granulomatous lesions or healthy sections. In both cases,

SNRPD2 staining was conspicuously nuclear and indistinctive (Fig. S9C). Thus, these results

highlight an unusual SNRPF-dependent dysregulation in human TB lesions. No previous

report of such rampant dysregulation of snRNP biogenesis in human TB granulomas exists.

Monocytes from human pulmonary TB patients show pervasive regulation of alternative splicing.

Next, from an RNAseq dataset from monocytes of 20 PTB cases and 19 healthy contacts, we analyzed and compared the relative abundance of different transcripts [38]. We focused our analysis on four major functional classes- RNA splicing, immune response, inflammation and metabolism (Fig. 6E, Table S7). As can be seen from the clustergrams, transcript level expression of genes belonging to each of these functional classes was able to classify the PTB and HC subjects (Fig. 6E). In each of the cases, there were two major clusters of transcripts- one showing increased expression in the TB cases (Cluster 1) and the other in the control cases (Cluster 2). We analyzed the biotypes of the transcripts present in these clusters across all the groups. In the RNA splicing class, cluster 2 was much bigger than cluster 1, showing more downregulation events in the TB cases (Fig. 6E). Moreover, the small number of transcripts that were upregulated in TB subjects were either not the protein-coding isoforms or encoded a truncated protein isoform (Fig. 6F). A similar analysis performed for other functional classes revealed systemic splicing defects (Fig. 6F). In inflammation, immune response and metabolism classes, while a large number of transcripts were upregulated in the PTB subjects, a significant number of those genes did not code for a protein or encoded a truncated isoform of the corresponding protein (Fig. 6F). For example, the upregulated genes in the immune response class show increased expression of the transcripts of genes like TLR4, THBS1, OAS, OASL3, GRN, CXCL16, etc. (Fig. 6G), however for each of them one or more splice variants with highly truncated protein length were also upregulated, which could significantly blunt the host responses. Comparison of monocyte expression data with those from H37Rv-infected THP-1 (Fig. 5B) showed around 45 commonly upregulated transcripts; however, almost a third of those genes did not code for a protein-coding variant (Fig. S9D, Table S8). Taken together, these observations make it evident that by perturbing

453 host RNA splicing, the mechanism for which we report in this study, Mtb exerts a pervasive

454 influence on host physiology and contributes towards systemic pathologies witnessed in TB

455 patients.

Discussions

RNA splicing is a conserved eukaryotic process which impacts cellular proteome diversity [10,39]. Also, its dysregulation is associated with diverse pathologies [40]. This study was designed to understand the mechanism of Mtb infection-mediated alteration of host RNA splicing[9]. Hijacking of host RNA splicing machinery by viral pathogens, including influenza virus and SARS-CoV2 is well-known [41-44]. However, very little is understood in the context of bacterial pathogens. Identification of several Mtb proteins that could directly interact with and alter host RNA splicing is a significant leap in our understanding of host-pathogen interactions. Similarly, the observation that different spliced variants could interact differently with Mtb proteins adds a new dimension to host-pathogen interactions. There are only limited efforts to systematically study and map out how alternative splicing events could contribute to the functional divergence in the proteome [45]. However, our results underscore that more efforts are needed to understand the consequences of this fundamental aspect of eukaryotic regulation.

The dysregulation in the abundance and distribution of SFs in the infected macrophages provided crucial insights into the underlying mechanism. The U snRNPs biogenesis is fundamental to eukaryotic gene expression, given their critical role in RNA splicing [46-48]. For snRNPs to form, pre-snRNAs from the nucleus are transported to the cytoplasm, where they interact with the multi-molecular SMN complex and Sm Core ring [46-48]. The snRNA-protein complex comprising SMN complex and Sm Core ring, is transported back to the nucleus where they mature into snRNPs and participate in the splicing reactions [49,50]. SMN complex, including SMN1/2 dimer and Gemin proteins, charge snRNAs with the Sm core ring [51]. A loss or altered stoichiometry in either the Gemins or SMN1 or 2, can lead to widespread splicing defects in different tissues [20,52-54].

Thus, while a few Mtb proteins could alter RNA splicing of test transcripts in the *in vitro*

assays, the dysregulation of SMN complex proteins and Sm core proteins reported in this

study appears to be the major driver for the global alteration of AS in Mtb-infected

macrophages. To our surprise, we also noted direct interactions between SNRPF and U5

snRNA with Mtb, which suggests potential sequestration of these key regulatory components

by Mtb. While several viral proteins are reported to directly interact with the SFs [9,36,44,55],

there is no precedence of such interaction of SFs or snRNAs directly with a viral or bacterial

pathogen.

There could be concerns about the specificity of this regulation since targeted dysregulation

of splicing factors (SFs) can lead to a pervasive impact on alternative splicing. However,

studies suggest it does not alter the splicing of all introns in the cell [19,56,57]. Rather, the impact

of one splicing factor dysregulation could be very distinct and selective from another SF

dysregulation [19]. At the same time, there is some functional congruence, too, in that a specific

SF could regulate AS of functionally related genes [19,56,57].

Experiments using the *hsr1* deleted Mtb strain provide definite proof that Mtb Hsr1 regulates

host RNA splicing in an ESX1-dependent manner. More strikingly, the loss of *hsr1* impacted

the ability of Mtb to interact and sequester U5snRNA or SNRPF. It is pertinent to ask why

Mtb targets U5snRNA and no other snRNAs or at least the U2snRNAs, which we analyzed.

During splicing, U5snRNPs recognize the acceptor splice site [58]. Several viral proteins

interact with the U5 components to disrupt the normal RNA splicing [36,55]. Thus, targeting of

U5snRNA through interaction reveals somewhat conserved pathogenic mechanisms across

diverse human pathogens. The decline in Mtb*Δhsr1* survival in macrophages and in mice

clearly indicates that the ability to disrupt snRNP biogenesis is a desirable phenotype for

tuberculosis pathogenesis. To note, the orthologues of *hsr1* are secreted xylose hydrolysing

enzymes, which is probably required by non-pathogenic, or opportunistic pathogenic bacteria

in search of diverse carbon sources. Therefore, the gain of function to interact with host SFs

appears to be an important adaptation during the evolution of Mtb as a human pathogen.

Interestingly, all snRNAs, with a few exceptions, while synthesized in the nucleus, must

travel to the cytosol for assembly before they are transported back to the nucleus for

participation in the splicing reaction [49]. The cytosolic export of pre-snRNAs ensures that

immature snRNAs do not get incorporated into the spliceosomal complexes in the nucleus,

which can alter host RNA splicing catastrophically [35,59]. This evolutionarily conserved quality

control step appears to be co-opted by the pathogenic mycobacteria during its evolution as the

human pathogen.

Tuberculosis granulomas evolve through a complex interplay between the bacterial factors

and host machinery, which largely benefits the pathogen [60]. Therefore, the intense SNRPF

staining of human tuberculosis granulomas is of great interest. Similarly, the monocyte

RNAseq data from the patients emphasize the pervasiveness of infection-induced alteration in

host RNA splicing. Incidentally, differentially expressed transcript isoforms were recently

reported in an LTBI cohort and identified those associated with resistance to TST/IGRA

conversion [61]. There are systemic inflammation, metabolic dysregulation and muscle loss

reported in TB patients [62], some of which could potentially be a consequence of the large-

scale splicing defects observed in these individuals.

To conclude, with the limitation of only select splicing events analyzed, we present a new

paradigm of pathogen-mediated perturbation of host physiology and show the hijacking of

this evolutionarily conserved mechanism by Mtb to blunt host responses. We propose that

inhibition of molecular interactions between the host and the pathogenic partners identified in

this study could be exploited for developing a novel class of therapeutics.

Acknowledgements:

The work was supported by the DBT-Wellcome Trust India Alliance Senior Fellowship to DK (IA/S/17/1/503071). The work was also partly supported by grants from SERB (EMR/2016/005296 to DK) and the Department of Biotechnology, Govt of India (BT/IC-06/003/91-Flagship program to DK). All work on virulent Mtb was performed in the Tuberculosis Aerosol Challenge Facility (TACF), a national facility at ICGEB, New Delhi, supported by the Department of Biotechnology, Govt. of India.

Author Contributions:

Original conceptualization: DK; Experiments: KC, NP, YK, DD, NA, PS, BM, HK, LS, RD, RDS, LM, BS, PY; Investigation: KC, DD; Analysis and Visualization: KC, MD, PY, DD, NSB, DK; Resources: VN, AS, VA, PD; Supervision: DK; Writing-original draft: DK, KC, MD, NP; Writing—review & editing: VN, AS, NSB, PD, DK

543 **Figures and Figure Legends**

Figure 1: Mycobacterial secretory proteins and host spliceosome proteins interactome.

A) The schematic flow chart for high-throughput one-to-one Y2H assay in 1536-well format. Protein pairs showing interaction in 3 or 4 biological replicates were considered as positive interactions.

B) Mtb proteins -host splicing factor interactome- the positive interactions from the Y2H screen were used to generate the network shown here. The diamond and circle represent the host and Mtb proteins, respectively, and the size corresponds to the number of interactions observed. The proteins coloured yellow in the network are discussed in detail in the text.

C) The Venn diagrams show proteins belonging to the major spliceosome components and their corresponding overlap with the interacting Mtb proteins.

D) A more descriptive representation of the Venn diagrams, shown above in Fig. 1C is presented in the table as a list of interactions.

E) Transcripts that were used for making the minigene constructs to test the role of Mtb proteins in host RNA splicing. Only exons are shown, and two different transcripts for each of the five genes are shown. The red bars indicate the overlapping sequences, and the skipped exons are shown in yellow.

F) On the left, a schematic of the minigene cassettes for DDX42 and ACADVL is shown. The constitutive exons are shown in green, while the alternative exons are shown in blue. *In vitro* transcribed products of minigenes were used for in-vitro splicing assays in the presence of purified Mtb proteins Rv3867, Rv0360, Rv1566c and Rv1471.

Figure 2: *Ex vivo* infection assays and reporter assays to test the role of Mtb proteins in host RNA splicing

A) The schematic representation of infecting THP-1 macrophages with *M. smegmatis* strains that express mycobacterial proteins under IVN inducible promotor. Isoform-specific qPCR for the truncated isoforms of ACADVL, DDX11, DDX42, ACSL1 and NFYA was used to assess the impact on alternative splicing.

B) Expression of spliced isoforms of the five target genes (as indicated) in THP-1 macrophages upon infection with WT or *Rv1566c, Rv1435c* and *Rv1643* knock-in *M smegmatis* lines. *M smegmatis* lines were induced with IVN (indicated dose) for 12 hours before infection. Data from three independent biological experiments. The p-values shown are from One-way ANOVA.

C) Designing of the custom *in vivo* fluorescent splicing reporter. Schematic for the constitutive splicing reporter pRint with a short adenovirus intron intervening the DsRed gene with a strong 5' and 3' splice sites. In the pGint splicing constructs, the alternative exon of ACADVL, DDX11, DDX42, ACSL1 and NFYA, along with the flanking 100bp intronic sequences, were inserted within eGFP reading frame. The exon disrupts the reading frame, however, skipping of the test exon reconstitutes EGFP fluorescence.

D) The pRint expressing stable HEK293T cell lines were co-transfected with a pGint construct and pDest-FLAG construct encoding proteins Rv1252c (NIC), Rv1566c, Rv1435c, Rv1643, or empty vector control, followed by Flow cytometry. In the DsRed$^+$ gate, %FITC$^+$ population was calculated to measure the extent of the test exon-skipping event.

E) %FITC$^+$ population for each of the indicated reporter constructs in the presence of indicated mycobacterial protein, non-interacting control or vector controls. Each dot represents an independent biological replicate (N=3), indicating p values from One-way ANOVA.

F) Validation of physical interactions between indicated pairs of *Mtb* proteins (FLAG-tagged) and host proteins (HA-tagged) in HEK293T cells. Data is representative of two or more biologically independent transfection and pulldown experiments.

Figure 3: Dysregulation of host RNA splicing machinery in Mtb-infected macrophages

A) Heatmap shows massive dysregulation in the level and cellular distribution of human proteins involved in RNA splicing and identified in MS of the spliceosome complex pulldown from the Mtb infected THP-1 macrophages 48 hours post-infection.

B) A schematic model depicting the ordered pathway of mRNA transcription by RNAP II and splicing catalyzed by spliceosome complex. The U1snRNP binds the 5' splice site and U2 snRNP binds the branch point adenine to form complex A. U4/U5.U6 snRNP binds to form complex B and causes conformational changes to form activated complex B. This is followed by the first transesterification reaction releasing U1 and U4 snRNP and the second transesterification reaction forming complex C releasing the lariat intron and U2, U5 and U6 snRNP. The U snRNP are the major players in splicing catalysis and their biogenesis is mediated by the SMN complex and the Sm proteins in the cytoplasm.

C-H) The proteins were visualized by indirect immunofluorescence (green), the nucleus was visualized by DAPI staining (blue) and mCherry-expressing bacteria (red) was used for infection. Each dot in the bar graphs represents a field analyzed, and an average of a hundred cells were quantified. The confocal images were analyzed for three different parameters- overall MFI (MFI/cell), number of Foci (Foci/cell) and cytosolic to the nuclear ratio (Cyt/Nuc) for all major splicing-related proteins upon Mtb infection such as C) U1 associated proteins (SF1), D) U5 associated proteins (DDX23), E) SMN complex proteins (GEMIN8) F) SMN complex proteins (GEMIN3) and G) Sm Core proteins (LSM7) and H) Sm Core proteins (SNRPF)

I) Image showing SNRPF colocalizing with bacterial phagosomes. The 3D reconstruction of the nucleus, SNRPF foci and bacteria (see methods) was done to show the SNRPF Foci at <0.2 microns distance from the bacterial phagosomes. Scale bar: 2 μm and p values indicate One-way ANOVA.

Figure 4: Dysregulation in SMN complex and Sm proteins impact U snRNA localization and overall assembly of U snRNPs.

A) Schematic representation of U snRNA nucleo-cytoplasmic trajectory and assembly of Sm core ring on snRNA by SMN complex to form snRNPs

B) Representative images for RNA FISH against U2 snRNA in THP-1 macrophages upon Mtb infection 24- and 48-hrs post-infection. The U2 snRNA probe was labelled with Alexa 488 (green), the nucleus was stained with DAPI (blue) and mCherry-expressing H37Rv (red) was used for infection. The plots corresponding to overall MFI (MFI/cell), number of Foci (Foci/cell) and cytosolic to nuclear ratio (Cyt/Nuc) of U2 snRNA are shown here. Each dot represents a field analyzed, and an average of a hundred cells are quantified. The p-values is from one-way ANOVA.

C) Representative images for RNA FISH against U5 snRNA in THP-1 macrophages upon Mtb infection 24- and 48-hrs post-infection. The U5 snRNA probe was labelled with Alexa 488 (green), the nucleus was stained with DAPI (blue) and mCherry-expressing H37Rv (red) was used for infection. The plots corresponding to overall MFI (MFI/cell), number of Foci (Foci/cell) and cytosolic to nuclear ratio (Cyt/Nuc) of U5 snRNA are shown here. Each dot represents a field analyzed, an average of a hundred cells are quantified, and p values are from one-way ANOVA.

D) Direct interaction between H37Rv (red) and U5snRNA at 24- and 48-hrs post-infection. For the plot at the right, the number of U5snRNA puncta within 0.2 microns from H37Rv surface was counted, p values from one-way ANOVA. Representative images for colocalization of Sm core protein SNRPF (green) to the H37Rv phagosome (red). The 3D reconstruction and spot analysis are done as previously described.

E) Expression of U5 snRNA in THP-1 macrophages, 12 and 48 hpi with H37Rv. Data from three independent biological experiments.

F) Immunoblot showing SNRPF protein expression in the whole cell lysates of uninfected (UI) and H37Rv-infected THP-1 macrophages, 48 hpi. Data is representative of two independent biological experiments.

Figure 5: Role of H37Rv hsr1 protein in snRNP biogenesis and altering host RNA splicing.

A) Q-PCR for the target transcripts as mentioned after infection with H37Rv, H37Rv*Δhsr1* and H37Rv*Δhsr1::WT* strain. Data from three biological replicates (N=3), *p<0.05, **p<0.01 (one-way ANOVA).

B) Clustergram of RNAseq data obtained from uninfected, or THP1 macrophages infected with H37Rv or H37Rv*Δhsr1* strains. Transcripts showing an infection-dependent increase (red box), hsr1-triggered increase (purple) or hsr1-blocked transcripts (turquoise) are shown. The distribution of different transcript biotypes across the three groups is shown in the aligned plot towards the right.

C) Representative image for the U5snRNA FISH in THP-1 macrophages infected with WT H37Rv or H37Rv*Δhsr1* strains. The plots at the right show the cytosolic to nuclear ratio and the ratio of interacting to non-interacting U5snRNA with WT H37Rv or H37Rv*Δhsr1*. **p<0.01 (one-way ANOVA).

D) Representative images of SNRPF staining in THP-1 macrophages infected with WT H37Rv or H37Rv*Δhsr1* strains. For the plot at the right, the number of SNRPF puncta within 0.2 microns from the H37Rv surface was counted as interacting and shows the ratio of interacting to non-interacting SNRPF with WT H37Rv or H37Rv*Δhsr1*. *p<0.05 (one-way ANOVA).

E) Representative images of THP1-macrophages stained with DAPI (nucleus) and Rv1435c antisera (green) upon infection with WT H37Rv, H37Rv*Δhsr1,* H37Rv*ΔCE* and H37Rv*ΔCE::WT* and H37Rv*Δhsr1::WT.* Images are representative of two independent biological experiments. Scale bar-2 μm.

F) Expression of truncated isoform as indicated upon infection with WT H37Rv*,* H37Rv*ΔCE* and H37Rv*ΔCE::WT* at 48hpi from three biologically independent experiments. The p values are from One-way ANOVA.

Figure 6: Regulation of host alternative splicing as a virulence mechanism

A) THP-1 macrophages were infected with WT H37Rv, H37RvΔ*hsr1*, H37RvΔ*hsr1::WT* strains. At 48 hours post-infection, bacterial load was examined using CFU plating. Data represent fold change in CFU at 48 hrs compared to 0 hr post-infection. (N=3), p values are from one-way ANOVA.

B) BALB/C mice (6-8 weeks old) were infected with 200 CFU of WT H37Rv, H37RvΔ*hsr1*, H37RvΔ*hsr1::WT* in separate experiments. At 2- and 4 weeks post-infection, bacterial load in the lungs of the animals were determined by CFU plating. P values are from one-way ANOVA.

C) The 4 week post-infection lung tissue sections of WT H37Rv, H37RvΔ*hsr1*, H37RvΔ*hsr1::WT* were stained with DAPI (nucleus) and SNRPF. Scale bar: 20 µm

D) Human intestinal tuberculosis granulomatous lesions were stained with SNRPF. The arrows indicate multinucleated giant cells with intense staining for cytosolic SNRPF. Black arrows show areas of intense SNRPF staining in the granulomas and in the giant cells.

E) RNAseq data from the monocytes of pulmonary TB cases (N=20) and healthy contacts (N=19) was analyzed for expression of differential alternatively spliced transcripts. Genes belonging to the indicated functional classes were selected for the cluster analysis. Cluster 1 in each case shows transcripts upregulated in PTB cases.

F) For each of the functional classes shown in Fig. 6D, the relative distribution of transcript biotypes (as indicated) is plotted and binned separately for those that were up- (cluster 1, red) or down- (cluster 2, blue) regulated in PTB cases.

G) The transcripts from the list of genes in the immune response class (Fig. 6D) were further analyzed for their length with respect to the longest encoded isoform of the corresponding gene. Only genes with more than one transcript in the list were chosen for this analysis. The red dotted line separates cluster 1 and cluster 2 transcripts. Transcripts of DYSF and RNASET2 are present in both clusters.

References

in *WHO consolidated guidelines on tuberculosis: Module 3: diagnosis - rapid diagnostics for tuberculosis detection WHO Guidelines Approved by the Guidelines Review Committee*   (2021).

Pisu, D., Huang, L., Grenier, J. K. & Russell, D. G. Dual RNA-Seq of Mtb-Infected Macrophages In Vivo Reveals Ontologically Distinct Host-Pathogen Interactions. *Cell Rep* **30**, 335-350 e334 (2020). https://doi.org/10.1016/j.celrep.2019.12.033

Matteucci, K. C., Correa, A. A. S. & Costa, D. L. Recent Advances in Host-Directed Therapies for Tuberculosis and Malaria. *Front Cell Infect Microbiol* **12**, 905278 (2022). https://doi.org/10.3389/fcimb.2022.905278

Young, C., Walzl, G. & Du Plessis, N. Therapeutic host-directed strategies to improve outcome in tuberculosis. *Mucosal Immunol* **13**, 190-204 (2020). https://doi.org/10.1038/s41385-019-0226-5

Kumar, D. & Rao, K. V. Regulation between survival, persistence, and elimination of intracellular mycobacteria: a nested equilibrium of delicate balances. *Microbes Infect* **13**, 121-133 (2011). https://doi.org/10.1016/j.micinf.2010.10.009

Karim, A. F. *et al.* Express path analysis identifies a tyrosine kinase Src-centric network regulating divergent host responses to Mycobacterium tuberculosis infection. *J Biol Chem* **286**, 40307-40319 (2011). https://doi.org/10.1074/jbc.M111.266239

Khan, N. *et al.* M. tuberculosis Reprograms Hematopoietic Stem Cells to Limit Myelopoiesis and Impair Trained Immunity. *Cell* **183**, 752-770 e722 (2020). https://doi.org/10.1016/j.cell.2020.09.062

Kalam, H., Singh, K., Chauhan, K., Fontana, M. F. & Kumar, D. Alternate splicing of transcripts upon Mycobacterium tuberculosis infection impacts the expression of functional protein domains. *IUBMB Life* **70**, 845-854 (2018). https://doi.org/10.1002/iub.1887

Chauhan, K., Kalam, H., Dutt, R. & Kumar, D. RNA Splicing: A New Paradigm in Host-Pathogen Interactions. *J Mol Biol* **431**, 1565-1575 (2019). https://doi.org/10.1016/j.jmb.2019.03.001

Liu, Y. *et al.* Impact of Alternative Splicing on the Human Proteome. *Cell Rep* **20**, 1229-1241 (2017). https://doi.org/10.1016/j.celrep.2017.07.025

Kalam, H., Fontana, M. F. & Kumar, D. Alternate splicing of transcripts shape macrophage response to Mycobacterium tuberculosis infection. *PLoS Pathog* **13**, e1006236 (2017). https://doi.org/10.1371/journal.ppat.1006236

Zhang, X. O., Fu, Y., Mou, H., Xue, W. & Weng, Z. The temporal landscape of recursive splicing during Pol II transcription elongation in human cells. *PLoS Genet* **14**, e1007579 (2018). https://doi.org/10.1371/journal.pgen.1007579

Hug, N., Longman, D. & Caceres, J. F. Mechanism and regulation of the nonsense-mediated decay pathway. *Nucleic Acids Res* **44**, 1483-1495 (2016). https://doi.org/10.1093/nar/gkw010

Pai, A. A. *et al.* Widespread Shortening of 3' Untranslated Regions and Increased Exon Inclusion Are Evolutionarily Conserved Features of Innate Immune Responses to Infection. *PLoS Genet* **12**, e1006338 (2016). https://doi.org/10.1371/journal.pgen.1006338

Bhatt, D. M. *et al.* Transcript dynamics of proinflammatory genes revealed by sequence analysis of subcellular RNA fractions. *Cell* **150**, 279-290 (2012). https://doi.org/10.1016/j.cell.2012.05.043

16 Pandya-Jones, A. *et al.* Splicing kinetics and transcript release from the chromatin
compartment limit the rate of Lipid A-induced gene expression. *RNA* **19**, 811-827
(2013). https://doi.org/10.1261/rna.039081.113
17 Wahl, M. C., Will, C. L. & Luhrmann, R. The spliceosome: design principles of a
dynamic RNP machine. *Cell* **136**, 701-718 (2009).
https://doi.org/10.1016/j.cell.2009.02.009
18 Wilkinson, M. E., Charenton, C. & Nagai, K. RNA Splicing by the Spliceosome.
Annu Rev Biochem **89**, 359-388 (2020). https://doi.org/10.1146/annurev-biochem-
091719-064225
19 Papasaikas, P., Tejedor, J. R., Vigevani, L. & Valcarcel, J. Functional splicing
network reveals extensive regulatory potential of the core spliceosomal machinery.
Mol Cell **57**, 7-22 (2015). https://doi.org/10.1016/j.molcel.2014.10.030
20 Zhang, R. *et al.* Structure of a key intermediate of the SMN complex reveals
Gemin2's crucial function in snRNP assembly. *Cell* **146**, 384-395 (2011).
https://doi.org/10.1016/j.cell.2011.06.043
21 Karijolich, J. & Yu, Y. T. Spliceosomal snRNA modifications and their function.
RNA Biol **7**, 192-204 (2010). https://doi.org/10.4161/rna.7.2.11207
22 Strasser, A., Dickmanns, A., Luhrmann, R. & Ficner, R. Structural basis for m3G-
cap-mediated nuclear import of spliceosomal UsnRNPs by snurportin1. *EMBO J* **24**,
2235-2243 (2005). https://doi.org/10.1038/sj.emboj.7600701
23 Lee, Y. & Rio, D. C. Mechanisms and Regulation of Alternative Pre-mRNA Splicing.
Annu Rev Biochem **84**, 291-323 (2015). https://doi.org/10.1146/annurev-biochem-
060614-034316
24 Penn, B. H. *et al.* An Mtb-Human Protein-Protein Interaction Map Identifies a Switch
between Host Antiviral and Antibacterial Responses. *Mol Cell* **71**, 637-648 e635
(2018). https://doi.org/10.1016/j.molcel.2018.07.010
25 Portal-Celhay, C. *et al.* Mycobacterium tuberculosis EsxH inhibits ESCRT-dependent
CD4(+) T-cell activation. *Nat Microbiol* **2**, 16232 (2016).
https://doi.org/10.1038/nmicrobiol.2016.232
26 Sharma, G., Upadhyay, S., Srilalitha, M., Nandicoori, V. K. & Khosla, S. The
interaction of mycobacterial protein Rv2966c with host chromatin is mediated
through non-CpG methylation and histone H3/H4 binding. *Nucleic Acids Res* **43**,
3922-3937 (2015). https://doi.org/10.1093/nar/gkv261
27 Mehra, A. *et al.* Mycobacterium tuberculosis type VII secreted effector EsxH targets
host ESCRT to impair trafficking. *PLoS Pathog* **9**, e1003734 (2013).
https://doi.org/10.1371/journal.ppat.1003734
28 de Souza, G. A., Leversen, N. A., Malen, H. & Wiker, H. G. Bacterial proteins with
cleaved or uncleaved signal peptides of the general secretory pathway. *J Proteomics*
75, 502-510 (2011). https://doi.org/10.1016/j.jprot.2011.08.016
29 Malen, H., Berven, F. S., Fladmark, K. E. & Wiker, H. G. Comprehensive analysis of
exported proteins from Mycobacterium tuberculosis H37Rv. *Proteomics* **7**, 1702-1718
(2007). https://doi.org/10.1002/pmic.200600853
30 Yaseen, I., Kaur, P., Nandicoori, V. K. & Khosla, S. Mycobacteria modulate host
epigenetic machinery by Rv1988 methylation of a non-tail arginine of histone H3. *Nat*
Commun **6**, 8922 (2015). https://doi.org/10.1038/ncomms9922
31 Malakar, B. *et al.* Phosphorylation of CFP10 modulates Mycobacterium tuberculosis
virulence. *mBio* **14**, e0123223 (2023). https://doi.org/10.1128/mbio.01232-23
32 Bonano, V. I., Oltean, S. & Garcia-Blanco, M. A. A protocol for imaging alternative
splicing regulation in vivo using fluorescence reporters in transgenic mice. *Nat Protoc*
2, 2166-2181 (2007). https://doi.org/10.1038/nprot.2007.292

33 Herzel, L., Ottoz, D. S. M., Alpert, T. & Neugebauer, K. M. Splicing and
transcription touch base: co-transcriptional spliceosome assembly and function. *Nat*
Rev Mol Cell Biol **18**, 637-650 (2017). https://doi.org/10.1038/nrm.2017.63
34 Oesterreich, F. C. *et al.* Splicing of Nascent RNA Coincides with Intron Exit from
RNA Polymerase II. *Cell* **165**, 372-381 (2016).
https://doi.org/10.1016/j.cell.2016.02.045
35 Becker, D. *et al.* Nuclear Pre-snRNA Export Is an Essential Quality Assurance
Mechanism for Functional Spliceosomes. *Cell Rep* **27**, 3199-3214 e3193 (2019).
https://doi.org/10.1016/j.celrep.2019.05.031
36 Boudreault, S. *et al.* Reovirus mu2 protein modulates host cell alternative splicing by
reducing protein levels of U5 snRNP core components. *Nucleic Acids Res* **50**, 5263-
5281 (2022). https://doi.org/10.1093/nar/gkac272
37 van Kessel, J. C. & Hatfull, G. F. Recombineering in Mycobacterium tuberculosis.
Nat Methods **4**, 147-152 (2007). https://doi.org/10.1038/nmeth996
38 Del Rosario, R. C. H. *et al.* Histone acetylome-wide associations in immune cells
from individuals with active Mycobacterium tuberculosis infection. *Nat Microbiol* **7**,
312-326 (2022). https://doi.org/10.1038/s41564-021-01049-w
39 Kornblihtt, A. R. *et al.* Alternative splicing: a pivotal step between eukaryotic
transcription and translation. *Nat Rev Mol Cell Biol* **14**, 153-165 (2013).
https://doi.org/10.1038/nrm3525
40 Bradley, R. K. & Anczukow, O. RNA splicing dysregulation and the hallmarks of
cancer. *Nat Rev Cancer* (2023). https://doi.org/10.1038/s41568-022-00541-7
41 Liao, K. C. & Garcia-Blanco, M. A. Role of Alternative Splicing in Regulating Host
Response to Viral Infection. *Cells* **10** (2021). https://doi.org/10.3390/cells10071720
42 Wang, C. *et al.* Abnormal global alternative RNA splicing in COVID-19 patients.
PLoS Genet **18**, e1010137 (2022). https://doi.org/10.1371/journal.pgen.1010137
43 Manet, E. *et al.* Modulation of alternative splicing during early infection of human
primary B lymphocytes with Epstein-Barr virus (EBV): a novel function for the viral
EBNA-LP protein. *Nucleic Acids Res* **49**, 10657-10676 (2021).
https://doi.org/10.1093/nar/gkab787
44 Thompson, M. G. *et al.* Viral-induced alternative splicing of host genes promotes
influenza replication. *Elife* **9** (2020). https://doi.org/10.7554/eLife.55500
45 Louadi, Z. *et al.* DIGGER: exploring the functional role of alternative splicing in
protein interactions. *Nucleic Acids Res* **49**, D309-D318 (2021).
https://doi.org/10.1093/nar/gkaa768
46 Neuenkirchen, N. *et al.* Reconstitution of the human U snRNP assembly machinery
reveals stepwise Sm protein organization. *EMBO J* **34**, 1925-1941 (2015).
https://doi.org/10.15252/embj.201490350
47 Shaw, D. J., Eggleton, P. & Young, P. J. Joining the dots: production, processing and
targeting of U snRNP to nuclear bodies. *Biochim Biophys Acta* **1783**, 2137-2144
(2008). https://doi.org/10.1016/j.bbamcr.2008.07.025
48 Massenet, S., Pellizzoni, L., Paushkin, S., Mattaj, I. W. & Dreyfuss, G. The SMN
complex is associated with snRNPs throughout their cytoplasmic assembly pathway.
Mol Cell Biol **22**, 6533-6541 (2002). https://doi.org/10.1128/MCB.22.18.6533-
6541.2002
49 Gubitz, A. K., Feng, W. & Dreyfuss, G. The SMN complex. *Exp Cell Res* **296**, 51-56
(2004). https://doi.org/10.1016/j.yexcr.2004.03.022
50 Pellizzoni, L. Chaperoning ribonucleoprotein biogenesis in health and disease. *EMBO*
Rep **8**, 340-345 (2007). https://doi.org/10.1038/sj.embor.7400941

51 Yong, J., Kasim, M., Bachorik, J. L., Wan, L. & Dreyfuss, G. Gemin5 delivers
snRNA precursors to the SMN complex for snRNP biogenesis. *Mol Cell* **38**, 551-562
(2010). https://doi.org/10.1016/j.molcel.2010.03.014
52 Schlaen, R. G. *et al.* The spliceosome assembly factor GEMIN2 attenuates the effects
of temperature on alternative splicing and circadian rhythms. *Proc Natl Acad Sci U S*
A **112**, 9382-9387 (2015). https://doi.org/10.1073/pnas.1504541112
53 Zhang, Z. *et al.* SMN deficiency causes tissue-specific perturbations in the repertoire
of snRNAs and widespread defects in splicing. *Cell* **133**, 585-600 (2008).
https://doi.org/10.1016/j.cell.2008.03.031
54 Cartegni, L., Hastings, M. L., Calarco, J. A., de Stanchina, E. & Krainer, A. R.
Determinants of exon 7 splicing in the spinal muscular atrophy genes, SMN1 and
SMN2. *Am J Hum Genet* **78**, 63-77 (2006). https://doi.org/10.1086/498853
55 De Maio, F. A. *et al.* The Dengue Virus NS5 Protein Intrudes in the Cellular
Spliceosome and Modulates Splicing. *PLoS Pathog* **12**, e1005841 (2016).
https://doi.org/10.1371/journal.ppat.1005841
56 Wang, Z. & Burge, C. B. Splicing regulation: from a parts list of regulatory elements
to an integrated splicing code. *RNA* **14**, 802-813 (2008).
https://doi.org/10.1261/rna.876308
57 Grosso, A. R. *et al.* Tissue-specific splicing factor gene expression signatures. *Nucleic*
Acids Res **36**, 4823-4832 (2008). https://doi.org/10.1093/nar/gkn463
58 Artemyeva-Isman, O. V. & Porter, A. C. G. U5 snRNA Interactions With Exons
Ensure Splicing Precision. *Front Genet* **12**, 676971 (2021).
https://doi.org/10.3389/fgene.2021.676971
59 Jia, Y., Mu, J. C. & Ackerman, S. L. Mutation of a U2 snRNA gene causes global
disruption of alternative splicing and neurodegeneration. *Cell* **148**, 296-308 (2012).
https://doi.org/10.1016/j.cell.2011.11.057
60 Ramakrishnan, L. Revisiting the role of the granuloma in tuberculosis. *Nat Rev*
Immunol **12**, 352-366 (2012). https://doi.org/10.1038/nri3211
61 Simmons, J. D. *et al.* Differentially expressed transcript isoforms associate with
resistance to tuberculin skin test and interferon gamma release assay conversion.
PLoS One **18**, e0284498 (2023). https://doi.org/10.1371/journal.pone.0284498
62 Mehdiratta, K. *et al.* Kupyaphores are zinc homeostatic metallophores required for
colonization of Mycobacterium tuberculosis. *Proc Natl Acad Sci U S A* **119** (2022).
https://doi.org/10.1073/pnas.2110293119
897

Figure 1

Figure 1: Mycobacterial secretory proteins and host spliceosome proteins interactome.

A) The schematic flow chart for high-throughput one-to-one Y2H assay in 1536-well format. Protein pairs showing interaction in 3 or 4 biological replicates were considered as positive interactions.

B) Mtb proteins -host splicing factor interactome- the positive interactions from the Y2H screen were used to generate the network shown here. The diamond and circle represent the host and Mtb proteins, respectively, and the size corresponds to the number of interactions observed. The proteins coloured yellow in the network are discussed in detail in the text.

C) The Venn diagrams show proteins belonging to the major spliceosome components and their corresponding overlap with the interacting Mtb proteins.

D) A more descriptive representation of the Venn diagrams, shown above in Fig. 1C is presented in the table as a list of interactions.

E) Transcripts that were used for making the minigene constructs to test the role of Mtb proteins in host RNA splicing. Only exons are shown, and two different transcripts for each of the five genes are shown. The red bars indicate the overlapping sequences, and the skipped exons are shown in yellow.

F) On the left, a schematic of the minigene cassettes for DDX42 and ACADVL is shown. The constitutive exons are shown in green, while the alternative exons are shown in blue. *In vitro* transcribed products of minigenes were used for in-vitro splicing assays in the presence of purified Mtb proteins Rv3867, Rv0360, Rv1566c and Rv1471.

Figure 2

Figure 2: *Ex vivo* infection assays and reporter assays to test the role of Mtb proteins in host RNA splicing

A) The schematic representation of infecting THP-1 macrophages with *M. smegmatis* strains that express mycobacterial proteins under IVN inducible promotor. Isoform-specific qPCR for the truncated isoforms of ACADVL, DDX11, DDX42, ACSL1 and NFYA was used to assess the impact on alternative splicing.

B) Expression of spliced isoforms of the five target genes (as indicated) in THP-1 macrophages upon infection with WT or *Rv1566c, Rv1435c* and *Rv1643* knock-in *M smegmatis* lines. *M smegmatis* lines were induced with IVN (indicated dose) for 12 hours before infection. Data from three independent biological experiments. The p-values shown are from One-way ANOVA.

C) Designing of the custom *in vivo* fluorescent splicing reporter. Schematic for the constitutive splicing reporter pRint with a short adenovirus intron intervening the DsRed gene with a strong 5' and 3' splice sites. In the pGint splicing constructs, the alternative exon of ACADVL, DDX11, DDX42, ACSL1 and NFYA, along with the flanking 100bp intronic sequences, were inserted within eGFP reading frame. The exon disrupts the reading frame, however, skipping of the test exon reconstitutes EGFP fluorescence.

D) The pRint expressing stable HEK293T cell lines were co-transfected with a pGint construct and pDest-FLAG construct encoding proteins Rv1252c (NIC), Rv1566c, Rv1435c, Rv1643, or empty vector control, followed by Flow cytometry. In the DsRed$^+$ gate, %FITC$^+$ population was calculated to measure the extent of the test exon-skipping event.

E) %FITC$^+$ population for each of the indicated reporter constructs in the presence of indicated mycobacterial protein, non-interacting control or vector controls. Each dot represents an independent biological replicate (N=3), indicating p values from One-way ANOVA.

F) Validation of physical interactions between indicated pairs of *Mtb* proteins (FLAG-tagged) and host proteins (HA-tagged) in HEK293T cells. Data is representative of two or more biologically independent transfection and pulldown experiments.

Figure 3

Figure 3: Dysregulation of host RNA splicing machinery in Mtb-infected macrophages

A) Heatmap shows massive dysregulation in the level and cellular distribution of human proteins involved in RNA splicing and identified in MS of the spliceosome complex pulldown from the Mtb infected THP-1 macrophages 48 hours post-infection.

B) A schematic model depicting the ordered pathway of mRNA transcription by RNAP II and splicing catalyzed by spliceosome complex. The U1snRNP binds the 5' splice site and U2 snRNP binds the branch point adenine to form complex A. U4/U5.U6 snRNP binds to form complex B and causes conformational changes to form activated complex B. This is followed by the first transesterification reaction releasing U1 and U4 snRNP and the second transesterification reaction forming complex C releasing the lariat intron and U2, U5 and U6 snRNP. The U snRNP are the major players in splicing catalysis and their biogenesis is mediated by the SMN complex and the Sm proteins in the cytoplasm.

C-H) The proteins were visualized by indirect immunofluorescence (green), the nucleus was visualized by DAPI staining (blue) and mCherry-expressing bacteria (red) was used for infection. Each dot in the bar graphs represents a field analyzed, and an average of a hundred cells were quantified. The confocal images were analyzed for three different parameters- overall MFI (MFI/cell), number of Foci (Foci/cell) and cytosolic to the nuclear ratio (Cyt/Nuc) for all major splicing-related proteins upon Mtb infection such as **C)** U1 associated proteins (SF1), **D)** U5 associated proteins (DDX23), **E)** SMN complex proteins (GEMIN8) **F)** SMN complex proteins (GEMIN3) and **G)** Sm Core proteins (LSM7) and **H)** Sm Core proteins (SNRPF)

I) Image showing SNRPF colocalizing with bacterial phagosomes. The 3D reconstruction of the nucleus, SNRPF foci and bacteria (see methods) was done to show the SNRPF Foci at <0.2 microns distance from the bacterial phagosomes. Scale bar: 2 μm and p values indicate One-way ANOVA.

Figure 4: Dysregulation in SMN complex and Sm proteins impact U snRNA localization and overall assembly of U snRNPs.

A) Schematic representation of U snRNA nucleo-cytoplasmic trajectory and assembly of Sm core ring on snRNA by SMN complex to form snRNPs

B) Representative images for RNA FISH against U2 snRNA in THP-1 macrophages upon Mtb infection 24- and 48-hrs post-infection. The U2 snRNA probe was labelled with Alexa 488 (green), the nucleus was stained with DAPI (blue) and mCherry-expressing H37Rv (red) was used for infection. The plots corresponding to overall MFI (MFI/cell), number of Foci (Foci/cell) and cytosolic to nuclear ratio (Cyt/Nuc) of U2 snRNA are shown here. Each dot represents a field analyzed, and an average of a hundred cells are quantified. The p-values is from one-way ANOVA.

C) Representative images for RNA FISH against U5 snRNA in THP-1 macrophages upon Mtb infection 24- and 48-hrs post-infection. The U5 snRNA probe was labelled with Alexa 488 (green), the nucleus was stained with DAPI (blue) and mCherry-expressing H37Rv (red) was used for infection. The plots corresponding to overall MFI (MFI/cell), number of Foci (Foci/cell) and cytosolic to nuclear ratio (Cyt/Nuc) of U5 snRNA are shown here. Each dot represents a field analyzed, an average of a hundred cells are quantified, and p values are from one-way ANOVA.

D) Direct interaction between H37Rv (red) and U5snRNA at 24- and 48-hrs post-infection. For the plot at the right, the number of U5snRNA puncta within 0.2 microns from H37Rv surface was counted, p values from one-way ANOVA. Representative images for colocalization of Sm core protein SNRPF (green) to the H37Rv phagosome (red). The 3D reconstruction and spot analysis are done as previously described.

E) Expression of U5 snRNA in THP-1 macrophages, 12 and 48 hpi with H37Rv. Data from three independent biological experiments.

F) Immunoblot showing SNRPF protein expression in the whole cell lysates of uninfected (UI) and H37Rv-infected THP-1 macrophages, 48 hpi. Data is representative of two independent biological experiments.

Figure 5: Role of H37Rv hsr1 protein in snRNP biogenesis and altering host RNA splicing.

A) Q-PCR for the target transcripts as mentioned after infection with H37Rv, H37Rv*Δhsr1* and H37Rv*Δhsr1::WT* strain. Data from three biological replicates (N=3), *p<0.05, **p<0.01 (one-way ANOVA).

B) Clustergram of RNAseq data obtained from uninfected, or THP1 macrophages infected with H37Rv or H37Rv*Δhsr1* strains. Transcripts showing an infection-dependent increase (red box), hsr1-triggered increase (purple) or hsr1-blocked transcripts (turquoise) are shown. The distribution of different transcript biotypes across the three groups is shown in the aligned plot towards the right.

C) Representative image for the U5snRNA FISH in THP-1 macrophages infected with WT H37Rv or H37Rv*Δhsr1* strains. The plots at the right show the cytosolic to nuclear ratio and the ratio of interacting to non-interacting U5snRNA with WT H37Rv or H37Rv*Δhsr1*. **p<0.01 (one-way ANOVA).

D) Representative images of SNRPF staining in THP-1 macrophages infected with WT H37Rv or H37Rv*Δhsr1* strains. For the plot at the right, the number of SNRPF puncta within 0.2 microns from the H37Rv surface was counted as interacting and shows the ratio of interacting to non-interacting SNRPF with WT H37Rv or H37Rv*Δhsr1*. *p<0.05 (one-way ANOVA).

E) Representative images of THP1-macrophages stained with DAPI (nucleus) and Rv1435c antisera (green) upon infection with WT H37Rv, H37Rv*Δhsr1,* H37Rv*ΔCE* and H37Rv*ΔCE::WT* and H37Rv*Δhsr1::WT*. Images are representative of two independent biological experiments. Scale bar-2 μm.

F) Expression of truncated isoform as indicated upon infection with WT H37Rv, H37Rv*ΔCE* and H37Rv*ΔCE::WT* at 48hpi from three biologically independent experiments. The p values are from One-way ANOVA.

Figure 6

Figure 6: Regulation of host alternative splicing as a virulence mechanism

A) THP-1 macrophages were infected with WT H37Rv, H37RvΔ*hsr1*, H37RvΔ*hsr1::WT* strains. At 48 hours post-infection, bacterial load was examined using CFU plating. Data represent fold change in CFU at 48 hrs compared to 0 hr post-infection. (N=3), p values are from one-way ANOVA.

B) BALB/C mice (6-8 weeks old) were infected with 200 CFU of WT H37Rv, H37RvΔ*hsr1*, H37RvΔ*hsr1::WT* in separate experiments. At 2- and 4 weeks post-infection, bacterial load in the lungs of the animals were determined by CFU plating. P values are from one-way ANOVA.

C) The 4 week post-infection lung tissue sections of WT H37Rv, H37RvΔ*hsr1*, H37RvΔ*hsr1::WT* were stained with DAPI (nucleus) and SNRPF. Scale bar: 20 μm

D) Human intestinal tuberculosis granulomatous lesions were stained with SNRPF. The arrows indicate multinucleated giant cells with intense staining for cytosolic SNRPF. Black arrows show areas of intense SNRPF staining in the granulomas and in the giant cells.

E) RNAseq data from the monocytes of pulmonary TB cases (N=20) and healthy contacts (N=19) was analyzed for expression of differential alternatively spliced transcripts. Genes belonging to the indicated functional classes were selected for the cluster analysis. Cluster 1 in each case shows transcripts upregulated in PTB cases.

F) For each of the functional classes shown in Fig. 6D, the relative distribution of transcript biotypes (as indicated) is plotted and binned separately for those that were up- (cluster 1, red) or down- (cluster 2, blue) regulated in PTB cases.

G) The transcripts from the list of genes in the immune response class (Fig. 6D) were further analyzed for their length with respect to the longest encoded isoform of the corresponding gene. Only genes with more than one transcript in the list were chosen for this analysis. The red dotted line separates cluster 1 and cluster 2 transcripts. Transcripts of DYSF and RNASET2 are present in both clusters.

References

in *WHO consolidated guidelines on tuberculosis: Module 3: diagnosis - rapid diagnostics for tuberculosis detection WHO Guidelines Approved by the Guidelines Review Committee* (2021).

Pisu, D., Huang, L., Grenier, J. K. & Russell, D. G. Dual RNA-Seq of Mtb-Infected Macrophages In Vivo Reveals Ontologically Distinct Host-Pathogen Interactions. *Cell Rep* **30**, 335-350 e334 (2020). https://doi.org/10.1016/j.celrep.2019.12.033

Matteucci, K. C., Correa, A. A. S. & Costa, D. L. Recent Advances in Host-Directed Therapies for Tuberculosis and Malaria. *Front Cell Infect Microbiol* **12**, 905278 (2022). https://doi.org/10.3389/fcimb.2022.905278

Young, C., Walzl, G. & Du Plessis, N. Therapeutic host-directed strategies to improve outcome in tuberculosis. *Mucosal Immunol* **13**, 190-204 (2020). https://doi.org/10.1038/s41385-019-0226-5

Kumar, D. & Rao, K. V. Regulation between survival, persistence, and elimination of intracellular mycobacteria: a nested equilibrium of delicate balances. *Microbes Infect* **13**, 121-133 (2011). https://doi.org/10.1016/j.micinf.2010.10.009

Karim, A. F. *et al.* Express path analysis identifies a tyrosine kinase Src-centric network regulating divergent host responses to Mycobacterium tuberculosis infection. *J Biol Chem* **286**, 40307-40319 (2011). https://doi.org/10.1074/jbc.M111.266239

Khan, N. *et al.* M. tuberculosis Reprograms Hematopoietic Stem Cells to Limit Myelopoiesis and Impair Trained Immunity. *Cell* **183**, 752-770 e722 (2020). https://doi.org/10.1016/j.cell.2020.09.062

Kalam, H., Singh, K., Chauhan, K., Fontana, M. F. & Kumar, D. Alternate splicing of transcripts upon Mycobacterium tuberculosis infection impacts the expression of functional protein domains. *IUBMB Life* **70**, 845-854 (2018). https://doi.org/10.1002/iub.1887

Chauhan, K., Kalam, H., Dutt, R. & Kumar, D. RNA Splicing: A New Paradigm in Host-Pathogen Interactions. *J Mol Biol* **431**, 1565-1575 (2019). https://doi.org/10.1016/j.jmb.2019.03.001

Liu, Y. *et al.* Impact of Alternative Splicing on the Human Proteome. *Cell Rep* **20**, 1229-1241 (2017). https://doi.org/10.1016/j.celrep.2017.07.025

Kalam, H., Fontana, M. F. & Kumar, D. Alternate splicing of transcripts shape macrophage response to Mycobacterium tuberculosis infection. *PLoS Pathog* **13**, e1006236 (2017). https://doi.org/10.1371/journal.ppat.1006236

Zhang, X. O., Fu, Y., Mou, H., Xue, W. & Weng, Z. The temporal landscape of recursive splicing during Pol II transcription elongation in human cells. *PLoS Genet* **14**, e1007579 (2018). https://doi.org/10.1371/journal.pgen.1007579

Hug, N., Longman, D. & Caceres, J. F. Mechanism and regulation of the nonsense-mediated decay pathway. *Nucleic Acids Res* **44**, 1483-1495 (2016). https://doi.org/10.1093/nar/gkw010

Pai, A. A. *et al.* Widespread Shortening of 3' Untranslated Regions and Increased Exon Inclusion Are Evolutionarily Conserved Features of Innate Immune Responses to Infection. *PLoS Genet* **12**, e1006338 (2016). https://doi.org/10.1371/journal.pgen.1006338

Bhatt, D. M. *et al.* Transcript dynamics of proinflammatory genes revealed by sequence analysis of subcellular RNA fractions. *Cell* **150**, 279-290 (2012). https://doi.org/10.1016/j.cell.2012.05.043

16 Pandya-Jones, A. *et al.* Splicing kinetics and transcript release from the chromatin
compartment limit the rate of Lipid A-induced gene expression. *RNA* **19**, 811-827
(2013). https://doi.org/10.1261/rna.039081.113
17 Wahl, M. C., Will, C. L. & Luhrmann, R. The spliceosome: design principles of a
dynamic RNP machine. *Cell* **136**, 701-718 (2009).
https://doi.org/10.1016/j.cell.2009.02.009
18 Wilkinson, M. E., Charenton, C. & Nagai, K. RNA Splicing by the Spliceosome.
Annu Rev Biochem **89**, 359-388 (2020). https://doi.org/10.1146/annurev-biochem-
091719-064225
19 Papasaikas, P., Tejedor, J. R., Vigevani, L. & Valcarcel, J. Functional splicing
network reveals extensive regulatory potential of the core spliceosomal machinery.
Mol Cell **57**, 7-22 (2015). https://doi.org/10.1016/j.molcel.2014.10.030
20 Zhang, R. *et al.* Structure of a key intermediate of the SMN complex reveals
Gemin2's crucial function in snRNP assembly. *Cell* **146**, 384-395 (2011).
https://doi.org/10.1016/j.cell.2011.06.043
21 Karijolich, J. & Yu, Y. T. Spliceosomal snRNA modifications and their function.
RNA Biol **7**, 192-204 (2010). https://doi.org/10.4161/rna.7.2.11207
22 Strasser, A., Dickmanns, A., Luhrmann, R. & Ficner, R. Structural basis for m3G-
cap-mediated nuclear import of spliceosomal UsnRNPs by snurportin1. *EMBO J* **24**,
2235-2243 (2005). https://doi.org/10.1038/sj.emboj.7600701
23 Lee, Y. & Rio, D. C. Mechanisms and Regulation of Alternative Pre-mRNA Splicing.
Annu Rev Biochem **84**, 291-323 (2015). https://doi.org/10.1146/annurev-biochem-
060614-034316
24 Penn, B. H. *et al.* An Mtb-Human Protein-Protein Interaction Map Identifies a Switch
between Host Antiviral and Antibacterial Responses. *Mol Cell* **71**, 637-648 e635
(2018). https://doi.org/10.1016/j.molcel.2018.07.010
25 Portal-Celhay, C. *et al.* Mycobacterium tuberculosis EsxH inhibits ESCRT-dependent
CD4(+) T-cell activation. *Nat Microbiol* **2**, 16232 (2016).
https://doi.org/10.1038/nmicrobiol.2016.232
26 Sharma, G., Upadhyay, S., Srilalitha, M., Nandicoori, V. K. & Khosla, S. The
interaction of mycobacterial protein Rv2966c with host chromatin is mediated
through non-CpG methylation and histone H3/H4 binding. *Nucleic Acids Res* **43**,
3922-3937 (2015). https://doi.org/10.1093/nar/gkv261
27 Mehra, A. *et al.* Mycobacterium tuberculosis type VII secreted effector EsxH targets
host ESCRT to impair trafficking. *PLoS Pathog* **9**, e1003734 (2013).
https://doi.org/10.1371/journal.ppat.1003734
28 de Souza, G. A., Leversen, N. A., Malen, H. & Wiker, H. G. Bacterial proteins with
cleaved or uncleaved signal peptides of the general secretory pathway. *J Proteomics*
75, 502-510 (2011). https://doi.org/10.1016/j.jprot.2011.08.016
29 Malen, H., Berven, F. S., Fladmark, K. E. & Wiker, H. G. Comprehensive analysis of
exported proteins from Mycobacterium tuberculosis H37Rv. *Proteomics* **7**, 1702-1718
(2007). https://doi.org/10.1002/pmic.200600853
30 Yaseen, I., Kaur, P., Nandicoori, V. K. & Khosla, S. Mycobacteria modulate host
epigenetic machinery by Rv1988 methylation of a non-tail arginine of histone H3. *Nat*
Commun **6**, 8922 (2015). https://doi.org/10.1038/ncomms9922
31 Malakar, B. *et al.* Phosphorylation of CFP10 modulates Mycobacterium tuberculosis
virulence. *mBio* **14**, e0123223 (2023). https://doi.org/10.1128/mbio.01232-23
32 Bonano, V. I., Oltean, S. & Garcia-Blanco, M. A. A protocol for imaging alternative
splicing regulation in vivo using fluorescence reporters in transgenic mice. *Nat Protoc*
2, 2166-2181 (2007). https://doi.org/10.1038/nprot.2007.292

33 Herzel, L., Ottoz, D. S. M., Alpert, T. & Neugebauer, K. M. Splicing and
transcription touch base: co-transcriptional spliceosome assembly and function. *Nat*
Rev Mol Cell Biol **18**, 637-650 (2017). https://doi.org/10.1038/nrm.2017.63
34 Oesterreich, F. C. *et al.* Splicing of Nascent RNA Coincides with Intron Exit from
RNA Polymerase II. *Cell* **165**, 372-381 (2016).
https://doi.org/10.1016/j.cell.2016.02.045
35 Becker, D. *et al.* Nuclear Pre-snRNA Export Is an Essential Quality Assurance
Mechanism for Functional Spliceosomes. *Cell Rep* **27**, 3199-3214 e3193 (2019).
https://doi.org/10.1016/j.celrep.2019.05.031
36 Boudreault, S. *et al.* Reovirus mu2 protein modulates host cell alternative splicing by
reducing protein levels of U5 snRNP core components. *Nucleic Acids Res* **50**, 5263-
5281 (2022). https://doi.org/10.1093/nar/gkac272
37 van Kessel, J. C. & Hatfull, G. F. Recombineering in Mycobacterium tuberculosis.
Nat Methods **4**, 147-152 (2007). https://doi.org/10.1038/nmeth996
38 Del Rosario, R. C. H. *et al.* Histone acetylome-wide associations in immune cells
from individuals with active Mycobacterium tuberculosis infection. *Nat Microbiol* **7**,
312-326 (2022). https://doi.org/10.1038/s41564-021-01049-w
39 Kornblihtt, A. R. *et al.* Alternative splicing: a pivotal step between eukaryotic
transcription and translation. *Nat Rev Mol Cell Biol* **14**, 153-165 (2013).
https://doi.org/10.1038/nrm3525
40 Bradley, R. K. & Anczukow, O. RNA splicing dysregulation and the hallmarks of
cancer. *Nat Rev Cancer* (2023). https://doi.org/10.1038/s41568-022-00541-7
41 Liao, K. C. & Garcia-Blanco, M. A. Role of Alternative Splicing in Regulating Host
Response to Viral Infection. *Cells* **10** (2021). https://doi.org/10.3390/cells10071720
42 Wang, C. *et al.* Abnormal global alternative RNA splicing in COVID-19 patients.
PLoS Genet **18**, e1010137 (2022). https://doi.org/10.1371/journal.pgen.1010137
43 Manet, E. *et al.* Modulation of alternative splicing during early infection of human
primary B lymphocytes with Epstein-Barr virus (EBV): a novel function for the viral
EBNA-LP protein. *Nucleic Acids Res* **49**, 10657-10676 (2021).
https://doi.org/10.1093/nar/gkab787
44 Thompson, M. G. *et al.* Viral-induced alternative splicing of host genes promotes
influenza replication. *Elife* **9** (2020). https://doi.org/10.7554/eLife.55500
45 Louadi, Z. *et al.* DIGGER: exploring the functional role of alternative splicing in
protein interactions. *Nucleic Acids Res* **49**, D309-D318 (2021).
https://doi.org/10.1093/nar/gkaa768
46 Neuenkirchen, N. *et al.* Reconstitution of the human U snRNP assembly machinery
reveals stepwise Sm protein organization. *EMBO J* **34**, 1925-1941 (2015).
https://doi.org/10.15252/embj.201490350
47 Shaw, D. J., Eggleton, P. & Young, P. J. Joining the dots: production, processing and
targeting of U snRNP to nuclear bodies. *Biochim Biophys Acta* **1783**, 2137-2144
(2008). https://doi.org/10.1016/j.bbamcr.2008.07.025
48 Massenet, S., Pellizzoni, L., Paushkin, S., Mattaj, I. W. & Dreyfuss, G. The SMN
complex is associated with snRNPs throughout their cytoplasmic assembly pathway.
Mol Cell Biol **22**, 6533-6541 (2002). https://doi.org/10.1128/MCB.22.18.6533-
6541.2002
49 Gubitz, A. K., Feng, W. & Dreyfuss, G. The SMN complex. *Exp Cell Res* **296**, 51-56
(2004). https://doi.org/10.1016/j.yexcr.2004.03.022
50 Pellizzoni, L. Chaperoning ribonucleoprotein biogenesis in health and disease. *EMBO*
Rep **8**, 340-345 (2007). https://doi.org/10.1038/sj.embor.7400941

51 Yong, J., Kasim, M., Bachorik, J. L., Wan, L. & Dreyfuss, G. Gemin5 delivers
snRNA precursors to the SMN complex for snRNP biogenesis. *Mol Cell* **38**, 551-562
(2010). https://doi.org/10.1016/j.molcel.2010.03.014
52 Schlaen, R. G. *et al.* The spliceosome assembly factor GEMIN2 attenuates the effects
of temperature on alternative splicing and circadian rhythms. *Proc Natl Acad Sci U S*
A **112**, 9382-9387 (2015). https://doi.org/10.1073/pnas.1504541112
53 Zhang, Z. *et al.* SMN deficiency causes tissue-specific perturbations in the repertoire
of snRNAs and widespread defects in splicing. *Cell* **133**, 585-600 (2008).
https://doi.org/10.1016/j.cell.2008.03.031
54 Cartegni, L., Hastings, M. L., Calarco, J. A., de Stanchina, E. & Krainer, A. R.
Determinants of exon 7 splicing in the spinal muscular atrophy genes, SMN1 and
SMN2. *Am J Hum Genet* **78**, 63-77 (2006). https://doi.org/10.1086/498853
55 De Maio, F. A. *et al.* The Dengue Virus NS5 Protein Intrudes in the Cellular
Spliceosome and Modulates Splicing. *PLoS Pathog* **12**, e1005841 (2016).
https://doi.org/10.1371/journal.ppat.1005841
56 Wang, Z. & Burge, C. B. Splicing regulation: from a parts list of regulatory elements
to an integrated splicing code. *RNA* **14**, 802-813 (2008).
https://doi.org/10.1261/rna.876308
57 Grosso, A. R. *et al.* Tissue-specific splicing factor gene expression signatures. *Nucleic*
Acids Res **36**, 4823-4832 (2008). https://doi.org/10.1093/nar/gkn463
58 Artemyeva-Isman, O. V. & Porter, A. C. G. U5 snRNA Interactions With Exons
Ensure Splicing Precision. *Front Genet* **12**, 676971 (2021).
https://doi.org/10.3389/fgene.2021.676971
59 Jia, Y., Mu, J. C. & Ackerman, S. L. Mutation of a U2 snRNA gene causes global
disruption of alternative splicing and neurodegeneration. *Cell* **148**, 296-308 (2012).
https://doi.org/10.1016/j.cell.2011.11.057
60 Ramakrishnan, L. Revisiting the role of the granuloma in tuberculosis. *Nat Rev*
Immunol **12**, 352-366 (2012). https://doi.org/10.1038/nri3211
61 Simmons, J. D. *et al.* Differentially expressed transcript isoforms associate with
resistance to tuberculin skin test and interferon gamma release assay conversion.
PLoS One **18**, e0284498 (2023). https://doi.org/10.1371/journal.pone.0284498
62 Mehdiratta, K. *et al.* Kupyaphores are zinc homeostatic metallophores required for
colonization of Mycobacterium tuberculosis. *Proc Natl Acad Sci U S A* **119** (2022).
https://doi.org/10.1073/pnas.2110293119
894