A *Mycobacterium tuberculosis* secreted virulence factor disrupts host snRNP biogenesis

Komal Chauhan[1], Nishat Passricha[1], Yogita Kapoor[2], Dipanwita Datta[1], Naresh Arora[1], Mrinmoy Das[1], Kavya Rao[1], Lakshyaveer Singh[1], Ravi Dutt[1], Raman Deep Sharma[1], Binayak Sarkar[3], Basanti Malakkar[3], Haroon Kalam[1], Prince Saini[4], Lalita Mehra[5], Prasenjit Das[5], Vineet Ahuja[6], Amit Singhal[7,8], Vinay Nandicoori[2, 3] and Dhiraj Kumar[1]*

[1]Cellular Immunology Group, International Centre for Genetic Engineering and Biotechnology, Aruna Asaf Ali Marg, New Delhi-110067, India
[2]Centre for Cellular and Molecular Biology, Hyderabad, India
[3]National Institute of Immunology, Aruna Asaf Ali Marg, New Delhi-110067, India
[4]Department of Biological Sciences, Indian Institute of Science Education and Research Mohali, Punjab 140306, India
[5]Department of Pathology, All India Institute of Medical Sciences, New Delhi-110012, India
[6]Department of Gastroenterology, All India Institute of Medical Sciences, New Delhi-110012, India
[7]A*STAR Infectious Diseases Labs (A*STAR ID labs), Agency for Science, Technology and Research (A*STAR), Singapore, Singapore.
[8]Singapore Immunology Network (SIgN), A*STAR, Singapore, Singapore.

Running title: Regulation of host RNA splicing by Mtb

*Correspondence- dhiraj@icgeb.res.in

Abstract

We earlier reported that *Mycobacterium tuberculosis* (*Mtb*) alters host RNA splicing to help its intracellular growth. Here, we report an unprecedented mechanism whereby a secreted virulence factor from *Mtb* interferes with the biogenesis of key spliceosomal components, causing an altered RNA splicing pattern. A high-throughput yeast-2-hybrid screen identified several *Mtb*-secreted proteins that can interact with the host RNA splicing factors (SFs). Through custom-designed in-cell assays, we show that one of those proteins, Rv1435c, targets specific exon-skipping events to alter RNA splicing. We show that Rv14345c or **h**ost **s**plicing **r**egulator 1 (hsr1) facilitates direct interaction between *Mtb* phagosomes and U5 snRNA and SNRPF, key components of the snRNPs. Genetic deletion of *hsr1* reverses the specific exon-skipping events caused by WT *Mtb* infection. The *Δhsr1* strain shows compromised growth during *ex vivo* infection in the macrophages and *in vivo* infection in the mice. Tissue sections from the WT *Mtb* or *Δhsr1*-infected mice showed significant hsr1-dependent SNRPF staining, a phenomenon also noted in the human intestinal tuberculosis (ITB) biopsies. We infer that hsr1 is a virulence factor which alters RNA splicing by interacting with U5snRNA and SNRPF. The splicing regulators from the host and pathogen are novel targets for anti-tuberculosis therapy.

Introduction

Tuberculosis (TB) caused by the bacterial pathogen *Mycobacterium tuberculosis* (*Mtb*) is a considerable challenge to the global sustainable development goals on health. With more than 10 million new TB cases and nearly 1 million deaths globally, the conventional approaches for prevention and control are increasingly looking insufficient to eradicate TB. The emerging paradigm on novel TB therapy envisages exploiting host-directed therapy (HDT). The advantage of HDT is its perceived universality in targeting drug-resistant or drug-sensitive infections. Also, HDTs, combined with antibiotics, are expected to hinder the development of new drug-resistant strains. Over the past two decades, several studies identified and reported host-dependency factors that help the intracellular growth and survival of *Mtb*, mostly by modulating the anti-bacterial functions of the hosts [1-7]. In a genome-wide analysis of identifying such host factors, proteins from diverse functional classes, including gene expression, were reported to have pro-bacterial functions [6].

The transcriptional response of the host cells to *Mtb* infections has been studied extensively[8-12]. While the immediate response of host cells following infection with *Mtb* involves processes like trafficking, signalling and metabolic alterations, the long-term shaping of host response is decided by reshaping the transcriptional landscape of the cells[13,14]. Eukaryotic gene expression involves several post-transcriptional regulatory events, including capping, poly-adenylation, RNA splicing, etc. We previously showed that in the *Mtb*-infected macrophages, several genes classically known for their anti-bacterial functions were spliced in a manner that did not allow their translation or the polypeptides thus formed were truncated[13]. Moreover, the alternatively spliced transcripts often lacked critical functional domains due to exon exclusion events[14,15]. Several other studies together converge to the notion that RNA splicing could be a central vulnerability for the pathogens to exploit.

However, the molecular mechanism that allows pathogens, including *Mtb*, to alter host RNA splicing in a targeted manner remains poorly understood. RNA splicing is an evolutionarily conserved mechanism among eukaryotes and involves the assembly of a large multi-molecular complex called spliceosome at the exon-intro junctions[16-22]. The key components of the spliceosomes are small nuclear ribonuclear proteins or snRNPs, comprising snRNA and associated proteins[23-31]. Interestingly, snRNPs are formed through complex protein-RNA interactions spanning across the nucleus and the cytosol[29,32]. In addition, accessory proteins like hnRNPs, SR proteins and SRSF proteins play an important role in recognizing the exon-

intron junctions and splicing reactions[17,19,33,34]. Cumulatively, there are plenty of potential

steps where secreted virulence proteins from *Mtb* could intercept the RNA splicing

machinery, thereby altering the course of downstream reactions. Secreted virulence factors of

Mtb are classically known to interfere with cytosolic events like trafficking, antigen

presentation, bioenergetics and nuclear events like epigenetic regulations [35-39].

To understand how *Mtb* alters host RNA splicing, we hypothesized that certain secreted

virulence factors of *Mtb* could interact with and alter the assembly and/or recognition of

spliceosome complexes or their components. Here, through a high throughput yeast-2-hybrid

screening between secreted *Mtb* proteins and host splicing-related proteins or splicing factors

(SFs), we identify several interacting partners between the two. We rationally narrowed it

down to one *Mtb* protein, Rv1435c, to understand the molecular mechanism in detail. Using a

genetic knockout of Rv1435c, we show that cytosolic interactions with snRNP biogenesis

components allowed *Mtb* to alter the stoichiometry of snRNPs in the cells, eventually altering

the RNA splicing process.

Results

Dysregulated spliceosome complex composition in *Mtb*-infected macrophages

Previously, we reported a targeted alteration of host RNA splicing pattern upon *Mtb* infection

in the macrophages[13]. We also showed how altered RNA splicing impacted cellular function

since a lot of target genes were spliced in a manner that led to the mRNAs, which were either

non-functional (lacking specific functional domains), unstable or could not get translated

(Fig. 1A)[14] . These alterations were reported as advantageous for the pathogen inside the host

macrophages. To understand how RNA splicing was specifically altered in the infected

macrophages, we systematically analyzed the RNA splicing process in these cells. RNA

splicing requires the assembly of a large multimolecular complex, called the spliceosome, in

a step-wise manner, which helps in the recognition, excision of the introns and stitching of

the exons together (Fig. 1B). To analyze the assembly of spliceosome complexes in the *Mtb*

infected macrophages, we decided to purify and analyze the *in vivo* assembled spliceosome

complexes from these cells. THP-1 macrophages were infected with H37Rv (the laboratory

strain of *Mtb*) at 1:10 MOI, and at 48 hours post-infection (h.p.i.), cytosolic and nuclear

extracts were prepared without disrupting the assembled spliceosome complexes (see

methods). The spliceosome complexes were then immunoprecipitated using the anti-SNRPB

antibody (Y-12) followed by mass spectrometry analysis (Fig. 1C). The detailed list of

identified peptides and proteins is shown in Table S1. The quantitative analysis of

spliceosome-related proteins is shown as the heatmap in Fig. 1D. Several host splicing

proteins showed distinct levels and nucleo-cytoplasmic distribution in the *Mtb*-infected

macrophages compared to the uninfected cells (Fig. 1D). Thus, proteins like RPL4, SF3B2

and SYNCRIP showed contrasting nucleo-cytoplasmic distribution in the infected cells (Fig.

1D). In contrast, proteins like SNRNPs and HNRNPs showed severe depletion of the nuclear

pool in the infected macrophages (Fig. 1D). The SNRNP proteins were particularly intriguing

since they are critical for the assembly of small nuclear ribonuclear proteins

(snRNPs)[23,26,40,41]. Dysregulation of these SFs suggests a systemic perturbation of the splicing

machinery in *Mtb*-infected macrophages.

Identification of *Mtb* proteins that could interact with host spliceosome complex

proteins

Since a large number of spliceosome-associated proteins were dysregulated, we hypothesized

that the virulent factors secreted by *Mtb* in the infected macrophages could likely be

interacting with some of these proteins and responsible for their dysregulation. To test this

hypothesis, we decided to screen whether any secreted protein from *Mtb* could interact with

the components of host RNA splicing machinery using the yeast-2-hybrid (Y2H) method.

The *Mtb* genes encoding the secreted proteins were part of the bait library (binding domain or

BD vector), while the human spliceosome-associated proteins encoding genes constituted the

prey library (activation domain or AD vector). In total, we had 384 *Mtb* genes in the BD

library, including 37 negative controls (Table S2). For the secreted protein list, we relied

mainly on our recent secretome study[42] in addition to other *Mtb* secretome studies[36,43-45]

(Table S2). The AD library was comprised of the protein-coding isoforms of genes belonging

to functional classes "RNA splicing" and "Spliceosome function" (~162 clones), making it a

unique custom library of human transcripts (Table S2). In the BD library, we also included a

few negative controls (proteins which are never reported in the secretome analysis, Table S3).

Four independent mating reactions between 384 BD clones and 162 AD clones, amounting to

about 2,48,832, were conducted to obtain the interaction. The schematic for the Y2H assay is

shown in Fig. 2A. In total, 123 Mtb proteins showed interactions with one or more SFs,

together consisting of 439 incidences of interactions (Table S4). Out of these 439, 121

interactions were a result of autoactivation (Table S4). The BD clones showing autoactivation

were excluded from the list of interacting proteins. A few negative control list clones also

showed interactions, including 56 autoactivation incidences (Table S3). The interacting BD

clones that did not show autoactivation were considered true interactors and mapped as a

network to represent the *Mtb*-host spliceosome interactome, comprising 54 human SFs and

117 *Mtb* proteins (Fig. 2B, Table S5).

The interactions spanned across all classes of host proteins that regulate RNA splicing,

including RNA polymerases, Sm proteins, SMN complex proteins, U2 snRNP proteins, LSm

proteins, U4/U5.U6 tricomplex proteins and non-snRNP proteins (Fig. 2C). The Venn

diagram in Fig. 2C shows number of *Mtb* proteins interacting with proteins from each of

these classes discussed above. Overall, host proteins from RNA Pol II subunits, Sm core,

catalytic step II factors, HnRNPs and SMN complex and *Mtb* proteins from information

pathway, virulence factors and conserved hypothetical proteins functional classes showed

maximum interactions (Fig. S1A). Interestingly, host proteins from similar functional steps in

RNA splicing shared their interacting partners from the *Mtb* (Fig. 2D). The Mtb proteins

showing interaction also represented diverse functional classes (Fig. S1B).

Finally, due to the inclusion of protein-coding splice variants of host SFs in the AD library,

eighteen genes were represented by more than one transcript in the interactions list (Fig. S2).

Out of those, POLR2H, SYF2, SNRPD2, LSm7, RBMX, LUC7L2, DCAF4, PPIH and

U2AF1 showed interactions with the *Mtb* proteins (Fig. S2). Curiously, the spliced variants

with a truncated coding region interacted with many *Mtb* proteins compared to their full-

length counterparts (Fig. S2). While this observation suggests looking at protein homeostasis

in the *Mtb*-infected macrophages more carefully, we decided to remain focused on our

objective of understanding the mechanism of infection-induced alternative splicing.

Mtb secretory protein Rv1435c could alter host RNA splicing

Several *Mtb* secretory proteins showed promising interactions in the Y2H assays above.

Before taking any of these proteins for further mechanistic explorations, it was important to

verify them on two accounts- one, if the identified *Mtb* secreted protein could alter host RNA

splicing and two, whether the identified interacting partners in the Y2H assay also interacted

with each other in a mammalian system. We picked up a few *Mtb* proteins, one showing a

very high number of interactors (Rv1643, 22 interactions), another with an intermediate

number of interactors (Rv1566c, 13 interactors) and a third with a low number of interactors

(Rv1435c, 5 interactors) (Table S3). To test whether these *Mtb* proteins could alter host RNA

splicing in a targeted manner, we first developed a few assays. For the functional assays, it

was critical to choose target genes for which splicing events were to be measured. We chose

around five target genes from the host, including *ACSL1, ACADVL, DDX11, DDX42* and

NFYA, which consistently showed exon exclusion events upon *Mtb* infection[13]. We designed

primers to score for the test exon exclusion event, exon inclusion event, and a set of primers

from the exon junctions (Fig. S3A). The exon exclusion primers clearly showed, upon *Mtb*

infection, an increase in the expression of selected transcripts of the genes where the test exon

was excluded (Fig. S3B). On the contrary, with the exon inclusion primers, except for the

ACSL1 transcript, others either did not show any regulation (ACADVL and DDX42) or

showed a decrease in expression upon *Mtb* infection (DDX11 and NFYA) (Fig. S3C).

Finally, primers designed from the junction of exons 1 and 3 conclusively showed that the

selected transcripts showed clear cases of *Mtb* infection-induced exon exclusion (Fig. S3D).

Next, we used a splicing reporter assay, which included a constitutive reporter construct

pRint and a test construct *pGint* [46] (Fig. 3A). The *pGint* reporter construct was modified to

include the test exon (middle exon) along with the flanking 100bp intronic sequences (Fig.

3A). The HEK293T cells stably expressing *pRint* were transfected with *pGint_ACADVL,*

pGint_ACSL1, pGint_DDX11, pGint_DDX42 and *pGint_NFYA* along with either a control

non-interacting *Mtb* protein encoding plasmid (NIC) or plasmid encoding for *Rv1435c,*

Rv1566c or Rv1643 (Fig. 3B). Expressing NIC alone led to a marginal but consistent increase

in the %pGint⁺ cells in the case of *DDX11* and *ACSL1* (Fig. 3C). Rv1435c and Rv1566c

showed a significant increase in %pGint⁺ cells in the case of all except *ACSL1* over and

above corresponding NIC expression (Fig. 3C). Unlike Rv1566c and Rv1435c, for Rv1643,

significant increase compared to the NIC were noted in the case *ACADVL, DDX11* and

DDX42 but not in the case of *NFYA* (Fig. 3C). Despite the limitation that we used only a

handful of target RNA splicing, we could show that the three selected *Mtb* proteins were able

to alter host RNA splicing.

In the above assay, among the three *Mtb* proteins analyzed, *Rv1566c* and *Rv1435c* showed a

stronger influence on exon exclusion events. Among the two, Rv1435c also contains a

signalP, which is essential for secretion through the classical bacterial secretory pathway [43,44].

Secondly, while some function for Rv1566c was assigned earlier [45,47], Rv1435c remains an

uncharacterised protein. Therefore, we decided to pursue further characterization of Rv1435c

in regulating host RNA splicing events.

First, we verified the physical interactions between Rv1435c and its Y2H counterparts from

the human SFs. Out of five interacting SFs, we picked up two Rv1435c interacting SFs,

GEMIN3 or DDX20 and PRPF18, to verify interactions in the mammalian cells. Flag-tagged

Rv1435c was co-expressed along with HA-tagged human counterparts in HEK293T cells,

followed by co-immunoprecipitation. Consistent with the results from the Y2H screen, both

anti-Flag and anti-HA pulldowns were able to co-immunoprecipitate each other, suggesting

strong interaction of Rv1435c protein with the two human proteins GEMIN3 and PRPF18

(Fig. 3D). Therefore, Rv1435c was able to induce exon exclusion event in the *in-cell* reporter

assay and also showed physical interaction with human SFs in the mammalian cells.

A surrogate infection model using *M smegmatis* knock-in line shows a direct role of

Rv1435c in altering host RNA splicing during infection

Next, to test whether Rv1435c could also impact these exon exclusion events during

infection, we used the non-pathogenic *Mycobacterium smegmatis* (*Msmeg*) strain. We cloned

Rv1435c under a Tet-inducible promoter and expressed it in the *Msmeg*, making it

Msmeg::Rv1435c (Fig. 3D). We confirmed that Rv1435c was expressed and secreted by the

modified *Msmeg* upon induction (Fig. S4A). Next, we infected THP-1 macrophages with

Msmeg, Msmeg::vector or Msmeg::Rv1435c, which were either untreated or treated with

ATC before infecting the cells (Fig. 3D). Specific exon exclusion events were then tested

using RT-PCR assays against all the five target events described above. For *ACADVL,*

DDX42, ACSL1 and *NFYA*, while we did observe a trend, the increase in exon exclusion

event when *Msmeg::Rv1435c* line was induced by anhydrotetracycline (ATC), it was not

very consistent and significant (Fig. S4B). In the case of DDX11, infection with *Msmeg*

alone, *Msmeg::vector* or *Msmeg::Rv1435c* did not induce the exon exclusion event (Fig. 3E).

However, when *Msmeg::Rv1435c* was induced overnight with ATC, there was a significant

increase in the case of *DDX11* exon exclusion event (Fig. 3E). Overall, results from the above

sections show that Rv1435c from *Mtb* interacts with select host SFs and also alters specific

host RNA splicing events.

***Mtb* Rv1435c is a secretory protein with no assigned function**

Once the role of Rv1435c in altering host RNA splicing was verified, we were curious to

know what was known about this protein or its homologues from other species. The *Mtb*

Rv1435c is a 202 amino acid-long polypeptide; a Pfam search revealed the presence of

signalsP (1-33) at the N-terminal, indicating its secretion through the classic sec-mediated

pathway (Fig. S4C). The remaining polypeptide segment (34-171) is classified as an

extracytoplasmic region with a disordered region towards the C-terminal (172-202, Fig.

S4C). Similar structural features were also revealed by AlphaFold Protein Structure

Database[48] (Fig. S4D). Sequence search revealed the presence of its orthologues in several

pathogenic and non-pathogenic species of the genus *Mycobacterium* (Fig. S4E). Classifying

these orthologues through a dendrogram analysis revealed an interesting pattern. In truly

pathogenic species like *Mtb* and *M. marinum*, this gene is classified as a secreted protein with

no function assigned (Fig. S4F). In opportunistic pathogenic species, like *M. helveticum, M.*

paratuberculosis, M. shinjukuense and a few other unclassified species, this gene has been

assigned the function of secreted beta-xylosidase (Fig. S4F). Even in a few truly pathogenic

species like *M. lacus* and *M. heildelbergense*, this gene is characterized as secreted beta-

xylosidase. It is possible that altering RNA splicing through interaction with the host

spliceosome complex emerged as a function during the evolution of the pathogenic species.

***MtbΔRv1435c* fails to alter target gene alternative splicing during infection**

Next, to establish that Rv1435c could be directly involved in altering the host RNA splicing

during *Mtb* infection, we deleted the *Rv1435c* gene in *Mtb* using recombineering[49] to generate

the knockout strain *MtbΔRv1435c* (Fig. 4A). The knockout (KO) of *Rv1435c* was verified by

PCR using specific primers as described in the methods and also by RT-PCR to test the

expression of Rv1435c RNA in the KO cells (Fig. 4B and 4C). We also generated an

Rv1435c complemented strain by electroporating the WT *Rv1435c* encoding plasmid in the

MtbΔRv1435c cells to generate *MtbΔRv1435c::WT*. Next, we infected THP-1 macrophages

with the WT *Mtb*, *MtbΔRv1435c or MtbΔRv1435c::WT* and assayed for the exon exclusion

events in the five target genes, as described earlier. While the WT *Mtb* infection, expectedly,

increased the exon exclusion event for each of the five target genes, in *MtbΔRv1435c*-

infected THP-1 macrophages, the exon exclusion events were significantly downregulated

(Fig. 4C). In the THP-1 macrophages infected with *MtbΔRv1435c::WT*, the exon exclusion

event was restored to the levels observed during WT *Mtb* infection (Fig. 4C). These results

indicate that Rv1435c protein of *Mtb* can alter the targeted host RNA splicing events in the

infected macrophages and therefore, we named it as **h**ost **s**plicing **r**egulator 1 or hsr1.

***Mtb* hsr1 binds to and sequesters U5snRNA in the cytosol**

Next, we wanted to understand the mechanism of hsr1-mediated alteration in host RNA

splicing. In the spliceosome complex pulldown results (Fig. 1) and the Y2H screening results

(Fig. 2), several proteins from the snRNP complexes were present. This included proteins

from the SMN complex like GEMINs, DDX proteins and those from the heptameric core that

are recruited to the 3' sm Site of snRNAs for the biogenesis of snRNPs. The biogenesis of

snRNPs follows an evolutionarily conserved nucleo-cytoplasmic shuttling, where snRNAs,

upon being synthesized in the nucleus, are exported to the cytosol (Fig. 5A). In the cytosol,

snRNAs bind with the SMN complex, followed by the recruitment of sm Core proteins[50]

(Fig. 5A). We anticipated that snRNP biogenesis, due to its cytosolic trajectory, could be a

potential target of *Mtb* proteins-mediated alteration. We analyzed the levels of two snRNAs-

U2 and U5 snRNAs, in the *Mtb*-infected macrophages using Fluorescence In Situ

Hybridization (FISH). While the U2 snRNA showed no noticeable pattern (Fig. S5A), we

noted a significant regulation in U5 snRNA levels in *Mtb*-infected macrophages at 24 and 48

hpi (Fig. 5B). While there was a significant decline in U5 snRNA levels at 24 hpi, we noted a

significant increase in U5 snRNA levels at 48 hpi (Fig. 5B-C). To our surprise, we also

observed a direct interaction between *Mtb* phagosomes and U5 snRNA (Fig. 5D). The U5

snRNA interaction with the *Mtb* phagosomes increased over time from almost no noticeable

interaction at 24 hpi to significantly high interaction at 48 hpi (Fig. 5D). It appeared that *Mtb*

phagosomes were binding and sequestering the U5 snRNA in the cytosol (Fig. 5D). Next, to

test whether hsr1 was by anyway involved in this binding, we infected THP-1 macrophages

with WT *Mtb* or *Δhsr1* and quantified the FISH signals to compare the interaction between

U5 snRNA and phagosomes (Fig. 5E). We noted a significant decline in U5 snRNA

interaction with the phagosomes in the case of *Δhsr1* infection (Fig. 5E). The above results

suggest that hsr1 was responsible for binding with U5 snRNA in the cytosol. Any defect in

snRNP biogenesis is known to alter the global RNA splicing pattern[16,30,31,51-54] . Therefore, it

was likely that the altered RNA splicing events regulated by hsr1 could be due to its

interaction with the U5 snRNA.

Mtb *hsr1* interacts with SNRPF, a key sm Core protein in the cytosol

The hsr1-dependent interaction of *Mtb* phagosomes with U5 snRNA raised the question of

whether the interaction involved only the U5 snRNAs or reflected the binding of the entire

U5snRNP complex. The snRNP complexes, as discussed above, comprise proteins from

SMN complexes like GEMINs, SMNs, etc and the heptameric ring of sm Core proteins,

including SNRPB, SNRPD1, SNRPD2, SNRPC, SNRPF, etc. We analyzed some of the

marker proteins of snRNPs using confocal microscopy, including DDX23, GEMIN8,

GEMIN3, Lsm7, SNRPD2 and SNRPF (Fig. 6A and S6). While some of these, like DDX23,

GEMIN8, SNRPD2 and Lsm7, showed some random interactions with the *Mtb* phagosomes,

none of them showed a pattern similar to that of U5 snRNA (Fig. S6). The only protein which

showed U5 snRNA-like interaction with *Mtb* phagosome was SNRPF (Fig. 6A). Moreover,

we further verified that SNRPF interaction with *Mtb* phagosomes was also hsr1 dependent

since there was a sharp decline of their interaction when *Δhsr1* strain was used for infection

(Fig. 6B). Curiously, SNRPF was also part of our Y2H prey library; however, we did not see

any interaction between SNRPF and hsr1 in the Y2H assay. Since our criteria for selecting

positive interactions were very stringent, we sought to test whether these two proteins interact

in the coimmunoprecipitation assay. In the HEK293T cells overexpressing Rv1435c-FLAG

and SNRPF-HA, we observed that FLAG immunoprecipitation was able to pulldown

SNRPF-HA (Fig. 6C). However, since other smCore proteins line SNRPD2 did not show

noticeable interaction with *Mtb* phagosomes, we believe that Mtb phagosomes, though hsr1

specifically binds with U5 snRNA and SNRPF. These results suggest that *Mtb* hsr1 facilitates

the phagosome interaction with U5 snRNA and the smCore protein SNRPF, likely

contributing to an altered host RNA splicing.

***Mtb* hsr1 secretion and, therefore, its impact on host RNA splicing is ESX1-dependent**

The hsr1 protein was a part of the list of *Mtb* proteins present in the culture filtrate, indicating

their secretion [42]. However, whether hsr1 also gets secreted during macrophage infection is

unknown. Secretion during infection could be a critical factor for the ability of hsr1 to

interact with the human spliceosome components. We generated antisera against hsr1 and

confirmed its specificity by immunoblotting and microscopy before using it to examine the

secretion of hsr1 during macrophage infection (Fig. S6A). In the uninfected macrophages,

expectedly, we could not see any signal for hsr1; however, in the WT *Mtb*-infected

macrophages, we could see several hsr1 foci on *Mtb* phagosomes and in the cytosol (Fig.

6D). When we used the Δ*hsr1*, there was no noticeable signal of hsr1 protein in the

phagosomes or in the cytosol (Fig. 6D). When we used the complemented strain Δ*hsr1::WT,*

the signal for hsr1 on the bacterial phagosome and in the cytosol was restored (Fig. 6D). We

next question, how *Mtb* hsr1 was secreted during macrophage infections. For access to

cytosol, the type VII secretion system ESX is critical, which is known to form pores in the

phagosome membrane, allowing access to the cytosol[37,45]. To test that, we used a strain

lacking CFP10 and ESAT6 (Δ*CE*), the two critical components of the ESX1 secretion system

[42,55,56]. In Δ*CE*-infected macrophages, we could not observe any signal for hsr1 in the cytosol

of infected cells (Fig. 6D). Moreover, even secretion of hsr1 appeared to be ESX1-dependent

since there was no hsr1 signal from the phagosomes (Fig. 6D). This was surprising since

sequence analysis predicted a putative signalP sequence at the N-terminal of hsr1, making it a

substrate of sec system[42]. Finally, when we used a complemented strain Δ*CE::WT*, where

both ESAT6 and CFP10 were restored, we could see hsr1 signal, both at the phagosomes and

in the cytosol (Fig. 6D). These results suggest that ESX1 pathway plays a central role in hsr1

secretion. The logical question next was to test whether that also impacts exon exclusion

events in the host target genes through alternative splicing. We compared specific exon

exclusion events in the case of ACADVL and ACSL1, and in both cases, Δ*CE* strain

infection behaved similar to Δ*hsr1* infection, whereas Δ*CE::WT* complemented strain showed

exon exclusion similar to the WT *Mtb* or Δ*hsr1::hsr1* infection (Fig. 6E). With the above

results, we conclude that hsr1-mediated impact on host RNA splicing was dependent on its

secretion in ESX1-dependent manner.

***Mtb* hsr1 is a virulence factor and acts by sequestering SNRPF**

Results so far show that without hsr1, *Mtb* could not alter the splicing events in the manner

caused by the WT strain. We wondered whether this loss of function also impacted the intra-

macrophage trajectory of *Mtb*. In *ex vivo* infected THP-1 macrophages, the growth of the

Δhsr1 strain was markedly reduced compared to the WT *Mtb* (Fig. 7A). In agreement with

that, *Δhsr1* strain infected macrophages showed increased Cathepsin D staining and also an

increase in the co-localization of *Mtb* with Cathepsin D, compared to the WT *Mtb* (Fig. 7B).

This suggests that *Δhsr1* strains fail to execute the phagosome maturation arrest as is known

for the WT *Mtb* [57]. Since hsr1 was also responsible for binding with SNRPF in the cytoplasm

of THP-1 macrophages, we next overexpressed SNRPF in THP-1 macrophages and infected

them with WT *Mtb*, expecting that stoichiometric abundance of SNRPF should overcome the

impact of hsr1 during WT infection and would help better control of bacterial growth. In

THP-1 macrophages expressing SNRPF and infected with WT *Mtb*, we noted increased MFI

for SNRPF and a reduction in the bacterial CFU (Fig. 7C). No such effect was seen when

another sm Core protein SNRPD2 was overexpressed in these cells (Fig. S6B). We next

compared the in vivo infection of Wt and *Δhsr1* strains in the mice model (Fig. 7D). Upon

infection in mice, the *Δhsr1* strain was significantly compromised in the establishment of

infection compared to the WT *Mtb*, as reflected by a reduced bacterial burden in the lungs of

infected mice at 2- and 4- weeks post-infection (Fig. 7E). We also analyzed SNRPF

immunostaining in the lung sections of uninfected WT *Mtb*-infected and *Δhsr1*-infected mice

with anti-SNRPF antibody. Consistent with the results observed during *ex vivo* infection in

THP-1 macrophages, the lung sections from WT *Mtb*-infected mice showed significant

accumulation of SNRPF in the extranuclear space, suggesting binding and possible

sequestration in the cytoplasm (Fig. 7F). This cytosolic staining of SNRPF was not observed

in the lung sections from mice that were infected with *Δhsr1* strain (Fig. 7F). Thus, cytosolic

sequestration of SNRPF by *Mtb* protein hsr1 is an important virulence mechanism of *Mtb*.

Altered RNA splicing in monocytes from human TB subjects and distinct SNRPF
dysregulation in the human TB granulomas

To understand the dysregulation of RNA splicing in human TB subjects, we re-analyzed

RNAseq data from the monocytes of human pulmonary TB subjects and corresponding

healthy contacts[58]. The dataset comprised monocyte transcriptomes from 20 PTB subjects

and 19 healthy controls. We analyzed this data set for transcript-level quantification (see

methods, table S6). Using all the significantly regulated transcripts, we could observe distinct

clustering of PTB subjects and healthy controls in the PCA analysis (Fig. S6C). Among the

large number of transcripts showing significant regulation, we focused specifically on

transcripts belonging to functional classes like RNA splicing, metabolism, immune response

and inflammation. As can be seen from the heat maps in Fig. 8A, based on the level of

transcript expression within each of the functional classes, the PTB group could be

distinctively identified from the healthy controls. This remarkable observation highlights

systemic RNA splicing alterations in tuberculosis patients. Finally, to assess whether an

SNRPF-dependent mechanism plays a role in altering RNA splicing in human subjects, we

screened human intestinal TB (ITB) granulomas for SNRPF by immunohistochemistry.

Consistent with the earlier results from *ex vivo* infection in THP-1 macrophages and *in vivo*

infection in mice, the human ITB granulomas showed extensive SNRPF staining (Fig. 8B).

Moreover, these staining were more pronounced in the multinucleate giant cells and

neighbouring histiocytes (Fig. 8B and S6D). Additionally, we could observe strong cytosolic

staining in the granulomas (Fig. 8B). No such distinct staining of SNRPF was noted in the

control tissues (Fig. S6E). Similarly, there was no specific staining difference for another sm

Core protein SNRPD2 between the control and ITB sections, again underscoring the

specificity of the SNRPF-mediated mechanism (Fig. S8). Taken together, using results from

ex vivo infection, *in vivo* infections and from the human TB subjects, we conclude that

SNRPF sequestration and consequent alteration in the host RNA splicing aids bacterial

growth and survival within the host, thereby contributing to its pathogenicity.

Discussions

The ability of *Mtb* to adapt to the hostile immunological environment in the hosts is remarkable. Almost every physiological process relevant to host immune responses appears to be targeted by the *Mtb,* enabling the pathogen's survival and consequent pathology. This study specifically looked at RNA splicing, a post-transcriptional gene regulation mechanism in the eukaryotes, to understand how *Mtb* achieves the targeted impairment of the seemingly complex process of RNA splicing in the host. As shown earlier by us and others, infection-induced altered RNA splicing helps the bacteria control host responses by preventing the synthesis of target genes in a fully functional form [13,15,59-61]. Considering that RNA splicing is fundamental to eukaryotic gene expression, which in turn is critical for shaping the cellular responses to infection, the mechanistic understanding of this process could have a transformational impact on our ability to develop novel strategies towards controlling TB.

Since RNA splicing is a multi-step process involving the assembly of a large multi-molecular complex, our working hypothesis was that certain secreted proteins from *Mtb* could interact with and, therefore, interfere in the assembly of spliceosome complexes, thereby altering the pattern of RNA splicing. Proteomic analysis of the spliceosome complexes from the infected macrophages and the Y2H screen between bacterial secretome and host SFs, through independent approaches, did converge to some dysregulation at the level of snRNPs, the key components of spliceosome complexes. The assembly of snRNPs require snRNAs to get transported to the cytoplasm, where they interact with the components of the SMN complex and the heptameric ring called sm Core proteins, together forming the snRNPs, that enter the nucleus and participate in the RNA splicing reactions[23,24,31,50,53]. While the interaction network obtained by Y2H was dense and practically impossible to verify all those interactions through independent techniques, we rationally chose a few *Mtb* genes identified through the Y2H screen for further validation. With the customized assays developed for the purpose, we narrowed down on Rv1435c or hsr1, as the major regulator of host RNA splicing during infection. Apart from our observation of its ability to alter host RNA splicing, there is no defined function of Rv1435c in the literature, making it an attractive target for mechanistic explorations.

Interestingly, the orthologues of Rv1435c in other mycobacterial species are classified as secreted xylosidase, an activity lacking in the *Mtb* hsr1. Mechanistically, hsr1 allowed *Mtb* to bind in the cytosol with two key components of snRNPs, U5 snRNA and SNRPF.

Experiments with SMN complex proteins like GEMIN3 or GEMIN8, smCore proteins like SNRPD2 and snRNAs like U2 snRNA reveal that this interaction was most likely selective with U5 snRNA and SNRPF, and it did not represent the interaction of *Mtb* with whole snRNP complex. Through studies in unrelated systems like neurodegenerative diseases and cancer, it is shown that stoichiometric disbalance of snRNP components like snRNA, SMN complex proteins or smCore proteins can dramatically alter the snRNP complex biogenesis and impact RNA splicing[30,31,34,50,54,62-64]. U5 snRNA is classically known for its role in recognition of the 3' splice site and, therefore, under depleting U5 snRNA condition, there could be a fierce competition between various 3' acceptor splice sites, causing weaker splice junctions to lose out and not get incorporated into the final processed mRNA. Similarly, SNRPF is part of several snRNP complexes, and its dysregulation due to cytoplasmic binding with *Mtb* may have a widespread impact on RNA splicing [31,54,63]. Crucially, the nucleo-cytoplasmic shuttling of snRNA for snRNP biogenesis is an evolutionarily conserved quality control mechanism[53]. However, it appears that *Mtb*, through its course of evolution, learnt to disrupt this process without actually going to the nucleus. The snRNP biogenesis defect appears to be operational in the lungs of *Mtb*-infected mice and the human TB granuloma sections. Since human PTB subjects also showed massive alterations in the RNA splicing pattern, to the extent that they could classify healthy from the PTB subjects, it is plausible that this mechanism could cause systemic RNA splicing defect, underlying many pathological states.

Interestingly, very limited reports highlight bacterial manipulation of host RNA splicing [13,15,65]. A large number of studies focus on chronic pathology like neurodegenerative disorders or cancer[51,64]. Among infectious diseases, the majority of studies relate to viral infections. However, the mechanism of viral infection-mediated RNA splicing alterations differs from the one reported in this study.

Loss of hsr1 in *Mtb* resulted in a compromised growth of bacteria inside human macrophages or in mouse lungs. This observation makes *Mtb* hsr1 a novel virulence factor. While we cannot rule out additional mechanisms, however, since unlike the WT *Mtb*, the *Δhsr1* strains cannot alter host RNA splicing, we propose that losing the ability to alter host RNA splicing compromises the virulence of *Mtb*. This observation makes host RNA splicing regulators and the mycobacterial factors regulating that as a new set of host- and pathogen-directed therapy candidates against TB, respectively.

Acknowledgements:

The work was supported by DBT-Wellcome Trust India Alliance Senior Fellowship to DK

(IA/S/17/1/503071). The work was also partly supported by grants from SERB

(EMR/2016/005296 to DK) and the Department of Biotechnology, Govt of India (BT/IC-

06/003/91-Flagship program to DK).

Author Contributions:

Original conceptualization: DK; Experiments: KC, NP, YK, DD, NA, PS, BM, HK, LS, LM;

Investigation: KC; Analysis and Visualization: KC, MD, DK; Resources: VN, AS, VA, PD;

Supervision: DK; Writing-original draft: DK, KC, MD, NP; Writing—review & editing: VN,

AS, PD, DK

Conflict of Interest:

The authors declare no conflict of interest.

References:

1. Augenstreich, J. & Briken, V. Host Cell Targets of Released Lipid and Secreted Protein Effectors of Mycobacterium tuberculosis. *Front Cell Infect Microbiol* **10**, 595029 (2020). https://doi.org/10.3389/fcimb.2020.595029

2. Chandra, P. *et al.* Mycobacterium tuberculosis Inhibits RAB7 Recruitment to Selectively Modulate Autophagy Flux in Macrophages. *Sci Rep* **5**, 16320 (2015). https://doi.org/10.1038/srep16320

3. de Martino, M., Lodi, L., Galli, L. & Chiappini, E. Immune Response to Mycobacterium tuberculosis: A Narrative Review. *Front Pediatr* **7**, 350 (2019). https://doi.org/10.3389/fped.2019.00350

4. Ernst, J. D. The immunological life cycle of tuberculosis. *Nat Rev Immunol* **12**, 581-591 (2012). https://doi.org/10.1038/nri3259

5. Flynn, J. L. & Chan, J. Immunology of tuberculosis. *Annu Rev Immunol* **19**, 93-129 (2001). https://doi.org/10.1146/annurev.immunol.19.1.93

6. Kumar, D. & Rao, K. V. Regulation between survival, persistence, and elimination of intracellular mycobacteria: a nested equilibrium of delicate balances. *Microbes Infect* **13**, 121-133 (2011). https://doi.org/10.1016/j.micinf.2010.10.009

7. Zhai, W., Wu, F., Zhang, Y., Fu, Y. & Liu, Z. The Immune Escape Mechanisms of Mycobacterium Tuberculosis. *Int J Mol Sci* **20** (2019). https://doi.org/10.3390/ijms20020340

8. Karim, A. F. *et al.* Express path analysis identifies a tyrosine kinase Src-centric network regulating divergent host responses to Mycobacterium tuberculosis infection. *J Biol Chem* **286**, 40307-40319 (2011). https://doi.org/10.1074/jbc.M111.266239

9. Koul, A., Herget, T., Klebl, B. & Ullrich, A. Interplay between mycobacteria and host signalling pathways. *Nat Rev Microbiol* **2**, 189-202 (2004). https://doi.org/10.1038/nrmicro840

10. Lavalett, L. *et al.* Alveolar macrophages from tuberculosis patients display an altered inflammatory gene expression profile. *Tuberculosis (Edinb)* **107**, 156-167 (2017). https://doi.org/10.1016/j.tube.2017.08.012

11. Lesho, E. *et al.* Transcriptional responses of host peripheral blood cells to tuberculosis infection. *Tuberculosis (Edinb)* **91**, 390-399 (2011). https://doi.org/10.1016/j.tube.2011.07.002

12. McGarvey, J. A., Wagner, D. & Bermudez, L. E. Differential gene expression in mononuclear phagocytes infected with pathogenic and non-pathogenic mycobacteria. *Clin Exp Immunol* **136**, 490-500 (2004). https://doi.org/10.1111/j.1365-2249.2004.02490.x

13. Kalam, H., Fontana, M. F. & Kumar, D. Alternate splicing of transcripts shape macrophage response to Mycobacterium tuberculosis infection. *PLoS Pathog* **13**, e1006236 (2017). https://doi.org/10.1371/journal.ppat.1006236

14. Kalam, H., Singh, K., Chauhan, K., Fontana, M. F. & Kumar, D. Alternate splicing of transcripts upon Mycobacterium tuberculosis infection impacts the expression of functional protein domains. *IUBMB Life* **70**, 845-854 (2018). https://doi.org/10.1002/iub.1887

15   Chauhan, K., Kalam, H., Dutt, R. & Kumar, D. RNA Splicing: A New Paradigm in Host-
Pathogen Interactions. *J Mol Biol* **431**, 1565-1575 (2019).
https://doi.org/10.1016/j.jmb.2019.03.001
16   Artemyeva-Isman, O. V. & Porter, A. C. G. U5 snRNA Interactions With Exons Ensure
Splicing Precision. *Front Genet* **12**, 676971 (2021).
https://doi.org/10.3389/fgene.2021.676971
17   Lee, Y. & Rio, D. C. Mechanisms and Regulation of Alternative Pre-mRNA Splicing.
Annu Rev Biochem **84**, 291-323 (2015). https://doi.org/10.1146/annurev-biochem-
060614-034316
18   Matera, A. G. & Wang, Z. A day in the life of the spliceosome. *Nat Rev Mol Cell Biol*
15, 108-121 (2014). https://doi.org/10.1038/nrm3742
19   Papasaikas, P., Tejedor, J. R., Vigevani, L. & Valcarcel, J. Functional splicing network
reveals extensive regulatory potential of the core spliceosomal machinery. *Mol Cell*
57, 7-22 (2015). https://doi.org/10.1016/j.molcel.2014.10.030
20   Schlautmann, L. P. & Gehring, N. H. A Day in the Life of the Exon Junction Complex.
Biomolecules **10** (2020). https://doi.org/10.3390/biom10060866
21   Wahl, M. C., Will, C. L. & Luhrmann, R. The spliceosome: design principles of a
dynamic RNP machine. *Cell* **136**, 701-718 (2009).
https://doi.org/10.1016/j.cell.2009.02.009
22   Wilkinson, M. E., Charenton, C. & Nagai, K. RNA Splicing by the Spliceosome. *Annu
Rev Biochem* **89**, 359-388 (2020). https://doi.org/10.1146/annurev-biochem-091719-
064225
23   Gubitz, A. K., Feng, W. & Dreyfuss, G. The SMN complex. *Exp Cell Res* **296**, 51-56
(2004). https://doi.org/10.1016/j.yexcr.2004.03.022
24   Karijolich, J. & Yu, Y. T. Spliceosomal snRNA modifications and their function. *RNA
Biol* **7**, 192-204 (2010). https://doi.org/10.4161/rna.7.2.11207
25   Liu, J. L. & Gall, J. G. U bodies are cytoplasmic structures that contain uridine-rich
small nuclear ribonucleoproteins and associate with P bodies. *Proc Natl Acad Sci U S
A* **104**, 11655-11659 (2007). https://doi.org/10.1073/pnas.0704977104
26   Patel, S. B. & Bellini, M. The assembly of a spliceosomal small nuclear
ribonucleoprotein particle. *Nucleic Acids Res* **36**, 6482-6493 (2008).
https://doi.org/10.1093/nar/gkn658
27   Pellizzoni, L. Chaperoning ribonucleoprotein biogenesis in health and disease. *EMBO
Rep* **8**, 340-345 (2007). https://doi.org/10.1038/sj.embor.7400941
28   Shaw, D. J., Eggleton, P. & Young, P. J. Joining the dots: production, processing and
targeting of U snRNP to nuclear bodies. *Biochim Biophys Acta* **1783**, 2137-2144
(2008). https://doi.org/10.1016/j.bbamcr.2008.07.025
29   Yong, J., Kasim, M., Bachorik, J. L., Wan, L. & Dreyfuss, G. Gemin5 delivers snRNA
precursors to the SMN complex for snRNP biogenesis. *Mol Cell* **38**, 551-562 (2010).
https://doi.org/10.1016/j.molcel.2010.03.014
30   Zhang, R. *et al.* Structure of a key intermediate of the SMN complex reveals Gemin2's
crucial function in snRNP assembly. *Cell* **146**, 384-395 (2011).
https://doi.org/10.1016/j.cell.2011.06.043
31   Zhang, Z. *et al.* SMN deficiency causes tissue-specific perturbations in the repertoire
of snRNAs and widespread defects in splicing. *Cell* **133**, 585-600 (2008).
https://doi.org/10.1016/j.cell.2008.03.031

32   Strasser, A., Dickmanns, A., Luhrmann, R. & Ficner, R. Structural basis for m3G-cap-
mediated nuclear import of spliceosomal UsnRNPs by snurportin1. *EMBO J* **24**, 2235-
2243 (2005). https://doi.org/10.1038/sj.emboj.7600701

33   Wang, Z. & Burge, C. B. Splicing regulation: from a parts list of regulatory elements to
an integrated splicing code. *RNA* **14**, 802-813 (2008).
https://doi.org/10.1261/rna.876308

34   Cartegni, L., Hastings, M. L., Calarco, J. A., de Stanchina, E. & Krainer, A. R.
Determinants of exon 7 splicing in the spinal muscular atrophy genes, SMN1 and
SMN2. *Am J Hum Genet* **78**, 63-77 (2006). https://doi.org/10.1086/498853

35   Pai, A. A. *et al.* Widespread Shortening of 3' Untranslated Regions and Increased
Exon Inclusion Are Evolutionarily Conserved Features of Innate Immune Responses
to Infection. *PLoS Genet* **12**, e1006338 (2016).
https://doi.org/10.1371/journal.pgen.1006338

36   Penn, B. H. *et al.* An Mtb-Human Protein-Protein Interaction Map Identifies a Switch
between Host Antiviral and Antibacterial Responses. *Mol Cell* **71**, 637-648 e635
(2018). https://doi.org/10.1016/j.molcel.2018.07.010

37   Portal-Celhay, C. *et al.* Mycobacterium tuberculosis EsxH inhibits ESCRT-dependent
CD4(+) T-cell activation. *Nat Microbiol* **2**, 16232 (2016).
https://doi.org/10.1038/nmicrobiol.2016.232

38   Sharma, G., Upadhyay, S., Srilalitha, M., Nandicoori, V. K. & Khosla, S. The interaction
of mycobacterial protein Rv2966c with host chromatin is mediated through non-CpG
methylation and histone H3/H4 binding. *Nucleic Acids Res* **43**, 3922-3937 (2015).
https://doi.org/10.1093/nar/gkv261

39   Yaseen, I., Kaur, P., Nandicoori, V. K. & Khosla, S. Mycobacteria modulate host
epigenetic machinery by Rv1988 methylation of a non-tail arginine of histone H3.
Nat Commun **6**, 8922 (2015). https://doi.org/10.1038/ncomms9922

40   Klimesova, K. *et al.* TSSC4 is a component of U5 snRNP that promotes tri-snRNP
formation. *Nat Commun* **12**, 3646 (2021). https://doi.org/10.1038/s41467-021-
23934-y

41   Neuenkirchen, N. *et al.* Reconstitution of the human U snRNP assembly machinery
reveals stepwise Sm protein organization. *EMBO J* **34**, 1925-1941 (2015).
https://doi.org/10.15252/embj.201490350

42   Malakar, B. *et al.* Phosphorylation of CFP10 modulates Mycobacterium tuberculosis
virulence. *mBio* **14**, e0123223 (2023). https://doi.org/10.1128/mbio.01232-23

43   de Souza, G. A., Leversen, N. A., Malen, H. & Wiker, H. G. Bacterial proteins with
cleaved or uncleaved signal peptides of the general secretory pathway. *J Proteomics*
75, 502-510 (2011). https://doi.org/10.1016/j.jprot.2011.08.016

44   Malen, H., Berven, F. S., Fladmark, K. E. & Wiker, H. G. Comprehensive analysis of
exported proteins from Mycobacterium tuberculosis H37Rv. *Proteomics* **7**, 1702-
1718 (2007). https://doi.org/10.1002/pmic.200600853

45   Mehra, A. *et al.* Mycobacterium tuberculosis type VII secreted effector EsxH targets
host ESCRT to impair trafficking. *PLoS Pathog* **9**, e1003734 (2013).
https://doi.org/10.1371/journal.ppat.1003734

46   Bonano, V. I., Oltean, S. & Garcia-Blanco, M. A. A protocol for imaging alternative
splicing regulation in vivo using fluorescence reporters in transgenic mice. *Nat Protoc*
2, 2166-2181 (2007). https://doi.org/10.1038/nprot.2007.292

47 Luan, X. *et al.* High Immunogenicity of a T-Cell Epitope-Rich Recombinant Protein
Rv1566c-444 From Mycobacterium tuberculosis in Immunized BALB/c Mice, Despite
Its Low Diagnostic Sensitivity. *Front Immunol* **13**, 824415 (2022).
https://doi.org/10.3389/fimmu.2022.824415

48 Varadi, M. *et al.* AlphaFold Protein Structure Database in 2024: providing structure
coverage for over 214 million protein sequences. *Nucleic Acids Res* **52**, D368-D375
(2024). https://doi.org/10.1093/nar/gkad1011

49 van Kessel, J. C. & Hatfull, G. F. Recombineering in Mycobacterium tuberculosis. *Nat*
Methods **4**, 147-152 (2007). https://doi.org/10.1038/nmeth996

50 Massenet, S., Pellizzoni, L., Paushkin, S., Mattaj, I. W. & Dreyfuss, G. The SMN
complex is associated with snRNPs throughout their cytoplasmic assembly pathway.
Mol Cell Biol **22**, 6533-6541 (2002). https://doi.org/10.1128/MCB.22.18.6533-
6541.2002

51 Wood, K. A., Eadsforth, M. A., Newman, W. G. & O'Keefe, R. T. The Role of the U5
snRNP in Genetic Disorders and Cancer. *Front Genet* **12**, 636620 (2021).
https://doi.org/10.3389/fgene.2021.636620

52 Will, C. L. *et al.* A novel U2 and U11/U12 snRNP protein that associates with the pre-
mRNA branch site. *EMBO J* **20**, 4536-4546 (2001).
https://doi.org/10.1093/emboj/20.16.4536

53 Becker, D. *et al.* Nuclear Pre-snRNA Export Is an Essential Quality Assurance
Mechanism for Functional Spliceosomes. *Cell Rep* **27**, 3199-3214 e3193 (2019).
https://doi.org/10.1016/j.celrep.2019.05.031

54 Jia, Y., Mu, J. C. & Ackerman, S. L. Mutation of a U2 snRNA gene causes global
disruption of alternative splicing and neurodegeneration. *Cell* **148**, 296-308 (2012).
https://doi.org/10.1016/j.cell.2011.11.057

55 Groschel, M. I., Sayes, F., Simeone, R., Majlessi, L. & Brosch, R. ESX secretion
systems: mycobacterial evolution to counter host immunity. *Nat Rev Microbiol* **14**,
677-691 (2016). https://doi.org/10.1038/nrmicro.2016.131

56 Stoop, E. J., Bitter, W. & van der Sar, A. M. Tubercle bacilli rely on a type VII army for
pathogenicity. *Trends Microbiol* **20**, 477-484 (2012).
https://doi.org/10.1016/j.tim.2012.07.001

57 Vergne, I. *et al.* Mycobacterium tuberculosis phagosome maturation arrest:
mycobacterial phosphatidylinositol analog phosphatidylinositol mannoside
stimulates early endosomal fusion. *Mol Biol Cell* **15**, 751-760 (2004).
https://doi.org/10.1091/mbc.e03-05-0307

58 Del Rosario, R. C. H. *et al.* Histone acetylome-wide associations in immune cells from
individuals with active Mycobacterium tuberculosis infection. *Nat Microbiol* **7**, 312-
326 (2022). https://doi.org/10.1038/s41564-021-01049-w

59 Denisenko, E. *et al.* Transcriptionally induced enhancers in the macrophage immune
response to Mycobacterium tuberculosis infection. *BMC Genomics* **20**, 71 (2019).
https://doi.org/10.1186/s12864-019-5450-6

60 Louadi, Z. *et al.* DIGGER: exploring the functional role of alternative splicing in
protein interactions. *Nucleic Acids Res* **49**, D309-D318 (2021).
https://doi.org/10.1093/nar/gkaa768

61 Bradley, R. K. & Anczukow, O. RNA splicing dysregulation and the hallmarks of
cancer. *Nat Rev Cancer* (2023). https://doi.org/10.1038/s41568-022-00541-7

62   Boudreault, S. *et al.* Reovirus mu2 protein modulates host cell alternative splicing by
reducing protein levels of U5 snRNP core components. *Nucleic Acids Res* **50**, 5263-
5281 (2022). https://doi.org/10.1093/nar/gkac272
63   De Maio, F. A. *et al.* The Dengue Virus NS5 Protein Intrudes in the Cellular
Spliceosome and Modulates Splicing. *PLoS Pathog* **12**, e1005841 (2016).
https://doi.org/10.1371/journal.ppat.1005841
64   Almstead, L. L. & Sarnow, P. Inhibition of U snRNP assembly by a virus-encoded
proteinase. *Genes Dev* **21**, 1086-1097 (2007). https://doi.org/10.1101/gad.1535607
65   Corre, M. *et al.* Alternative splicing induced by bacterial pore-forming toxins
sharpens CIRBP-mediated cell response to Listeria infection. *Nucleic Acids Res* **51**,
12459-12475 (2023). https://doi.org/10.1093/nar/gkad1033
699
700

Figure Legends:

Figure 1: Dysregulation of host RNA splicing machinery in Mtb-infected macrophages

A) The schematic representation of perturbed alternative splicing resulting in exon skipping events upon *Mtb* infection.

B) A schematic model depicting the ordered pathway of mRNA transcription by RNAP II and splicing catalysed by spliceosome complex. The U1snRNP binds the 5' splice site and U2 snRNP binds the branch point adenine to form complex A. U4/U5.U6 snRNP binds to form complex B and causes conformational changes to form activated complex B. This is followed by the first transesterification reaction, releasing U1 and U4 snRNP and the second transesterification reaction forming complex C, releasing the lariat intron and U2, U5 and U6 snRNP. The U snRNP are the major players in splicing catalysis.

C) The schematic of the spliceosome complex pulldown from the *Mtb* infected THP-1 macrophages 48 hours post-infection for Mass Spectrometry.

D) Heatmap shows massive dysregulation in the level and cellular distribution of human proteins involved in RNA splicing and identified in MS of the spliceosome complex pulldown from the *Mtb* infected THP-1 macrophages 48 hours post-infection. The three boxes depict proteins that show depletion (green), depletion and mislocalisation (blue) and mislocalisation (red) patterns in Mtb-infected cells compared to the uninfected cells.

Figure 2: Mycobacterial secretory proteins and host spliceosome proteins interactome.

A) The schematic flow chart for high throughput one-to-one Y2H assay in 1536 well format. Protein pairs showing interaction in 3 or 4 biological replicates were considered as positive interactions.

B) *Mtb* proteins -host splicing factor interactome- the positive interactions from the Y2H screen were used to generate the network shown here. The diamond and circle represent the host and *Mtb* proteins, respectively, and the size corresponds to the number of interactions observed. The proteins coloured yellow in the network are described further in the subsequent sections.

C) The Venn diagrams show proteins belonging to the major spliceosome components and their corresponding overlap with the interacting *Mtb* proteins.

Figure 3: *Ex vivo* infection assays and reporter assays to test the role of *Mtb* proteins in host RNA splicing

A) Designing of the custom *in vivo* fluorescent splicing reporter. Schematic for the constitutive splicing reporter pRint with a short adenovirus intron intervening the DsRed gene with a strong 5' and 3' splice sites. In the pGint splicing constructs, the alternative exon of ACADVL, DDX11, DDX42, ACSL1 and NFYA, along with the flanking 100bp intronic sequences, were inserted within eGFP reading frame. The exon disrupts the reading frame, however, skipping of the test exon reconstitutes EGFP fluorescence.

B) In cell reporter assay: the pRint expressing stable HEK293T cell lines were co-transfected with a pGint construct and pDest-FLAG construct encoding proteins Rv1252c (NIC), Rv1566c, Rv1435c, Rv1643, or empty vector control, followed by Flow cytometry. In the DsRed$^+$ gate, the %FITC$^+$ population was calculated to measure the extent of the test exon-skipping event.

C) %FITC$^+$ population for each of the indicated reporter constructs in the presence of indicated mycobacterial protein, non-interacting control, or vector controls. Each dot represents an independent biological replicate (N=3), indicating p values from One-way ANOVA.

D) Validation of physical interactions between indicated pairs of Rv1435c proteins (FLAG-tagged) and host proteins – (HA-tagged) in HEK293T cells. Data is representative of two or more biologically independent transfection and pulldown experiments.

E) The schematic representation of infecting THP-1 macrophages with *Msmeg* strains that express Rv1435c under IVN inducible promotor followed by RTPCT for assessing target gene splicing event.

F) Isoform-specific qPCR to assess the impact of Rv1435c expressing *Msmeg* infection in THP-1 macrophages on exon skipping event in DDX11, N=3, indicating p values from One-way ANOVA.

Figure 4: Role of Mtb Rv1435c protein in altering host RNA splicing.

A) The schematic of knocking out the *Rv1435c* or *hsr1* gene in H37Rv using the recombineering approach. Positive colonies were screened with the primers F1-F3 (forward) & R1-R3 (reverse), as shown by the arrows.

B) Agarose gel electrophoresis to confirm the replacement of 609bp product (*Rv1435c*) with 1.4Kb hygromycin cassette in the knockout line (F1-R1), absence of hyg in the WT (F2-R2 and F3-R3) and control segment (*rbpA*).

C) Q-PCR to show the loss of expression of *Rv1435c* in the knockout strain

D) Q-PCR for the target transcripts as mentioned after infection with H37Rv, H37Rv*Δhsr1* and H37Rv*Δhsr1::WT* strain. Data from three biological replicates (N=3), indicating p-values from a one-way ANOVA analysis.

Figure 5: Direct interaction of U5 snRNA with Mtb phagosomes.

A) Schematic representation of U snRNA nucleo-cytoplasmic trajectory and assembly of Sm core ring on snRNA by SMN complex to form snRNPs

B) Representative images for RNA FISH against U5 snRNA in THP-1 macrophages upon *Mtb* (mCherry-H37Rv) infection at 24- and 48-hrs post-infection. The U5 snRNA probe was labelled with Alexa 488 (green), and the nucleus was stained with DAPI (blue).

C) Plots show quantification of U5 snRNA FISH MFI in uninfected and *Mtb*-infected macrophages at 24 ad 48 hpi.

D) Direct interaction between H37Rv (red) and U5snRNA at 24- and 48-hrs post-infection. U5snRNA puncta within 0.2 microns from the H37Rv surface are shown as green dots on the 3-D constructed Mtb contours. The 3D reconstruction of the nucleus, U5 snRNA

789   foci and bacteria (see methods) was done to show the U5 snRNA Foci at <0.2 microns
790   distance from the bacterial phagosomes. The quantification of interacting U5 snRNA
791   punctas is shown in the plot at the right. Scale bar: 2 μm.

E) Representative image for the U5snRNA FISH in THP-1 macrophages infected with
     mCherry expressing WT H37Rv or H37Rv*Δhsr1* strains. The plots at the right show
     the ratio of interacting to non-interacting U5snRNA with WT H37Rv or H37Rv*Δhsr1*.
     Data is from three biological experiments, two fields each, representing p-value from a
     Student's t-test analysis.

Figure 6: *Mtb* hsr1 secretion is ESX1-dependent, and impacts host RNA splicing by sequestering SNRPF

A) Image showing SNRPF colocalising with bacterial phagosomes. The 3D reconstruction
     of the nucleus, SNRPF foci and bacteria (see methods) was done to show the SNRPF
     Foci at <0.2 microns distance from the bacterial phagosomes. Scale bar: 2 μm and p
     values indicate Student's t-test.

B) Representative images of SNRPF staining in THP-1 macrophages infected with WT
     H37Rv or H37Rv*Δhsr1* strains. For the plot below, the number of SNRPF puncta within
     0.2 microns from the H37Rv surface was counted as interacting and shown as the ratio
     of interacting to non-interacting SNRPF with WT H37Rv or H37Rv*Δhsr1*. Data is from
     three biological replicates with two or more fields per slide, representing p-values from
     Student's t-test.

C) Validation of physical interactions between Rv1435c (FLAG-tagged) and SNRPF (HA-
     tagged) in HEK293T cells using co-immunoprecipitation. Data is representative of two
     or more biologically independent transfection and pulldown experiments.

D) Representative images of THP1-macrophages stained with DAPI (nucleus) and
     Rv1435c antisera (green) upon infection with mCherry expressing WT H37Rv, *Δhsr1,*
     Δhsr1::WT, ΔCE and *ΔCE::WT* strains*.* Images are representative of two independent
     biological experiments. Scale bar-2 μm.

E) The exon-skipping event in ACADVL and ACSL1 transcripts upon infection with WT
     H37Rv*,* H37Rv*ΔCE* and H37Rv*ΔCE::WT* at 48hpi from three biologically independent
     experiments. The p values are from One-way ANOVA.

Figure 7: Regulation of host alternative splicing as a virulence mechanism

A) THP-1 macrophages were infected with WT H37Rv or *Δhsr1* strains. At 48 hours post-
     infection, bacterial load was examined using CFU plating. Data represent fold change
     in CFU at 48 hrs compared to 0 hr post-infection. (N=3), p values are from Student's t-
     test.

B) The plot shows quantification for staining of cathepsin D in THP1 macrophages
     infected with WT H37Rv and *Δhsr1*at 48hpi. The different parameters plotted show
     overall MFI (MFI/cell) and the ratio of cathepsin D foci interacting with bacterial
     phagosomes (int/non-int). Each dot represents a field analyzed, and an average of a
     hundred cells are quantified, representing p-values from one-way ANOVA (MFI) or
     Student's t-test (Int/non-Int).

C) The plot shows quantification (MFI/cell) for staining of SNRPF in WT THP1 macrophages and THP1::*SNRPF* infected with WT H37Rv at 48hpi. Each dot represents a field analyzed, and an average of a hundred cells are quantified. The plot on the right shows the bacterial survival after 48 hours post-infection in WT THP1 and THP1::*SNRPF*. Each dot represents an independent biological experiment. Indicated p values represent Student's t-test.
D) Schematic for study design using mice model of infection
E) BALB/C mice (6-8 weeks old) were infected with 200 CFU of WT H37Rv and *Δhsr1* in separate experiments. At 2- and 4- weeks post-infection, bacterial load in the lungs of the animals was determined by CFU plating. N=6, representing p-values from Student's t-test.
F) The 4-week post-infection lung tissue sections of uninfected, WT H37Rv and Δhsr1 were stained with DAPI (nucleus) and anti-SNRPF antibody. Scale bar: 20 μm

Figure 8: Altered RNA splicing and distinct SNRPF dysregulation in the granulomas of human TB subjects.

A) RNAseq data from the monocytes of pulmonary TB cases (N=20) and healthy contacts (N=19) was analyzed to assess the expression of differential alternatively spliced transcripts. Genes belonging to the indicated functional classes were selected for the cluster analysis. Cluster 1 in each case shows transcripts upregulated in PTB cases.
B) Biopsy sections from human intestinal tuberculosis granulomatous lesions were stained with anti-SNRPF antibody. The arrows indicate multinucleated giant cells with intense staining for cytosolic SNRPF. Black arrows show areas of intense SNRPF staining in the granulomas and in the giant cells.

Figure 9: A model to show *Mtb*-mediated alteration of host RNA splicing by blocking the snRNP biogenesis and post-transcriptional gene regulation in the macrophages.

A

pre-mRNA

Uninfected | Mtb infection

Alternative splicing

Functional isoform | mRNA | Truncated isoform
Non-sense mediated decay

Anti-bacterial | Pro-bacterial

B

RNA Pol II

U1

U2

U1 U2
Pre-Spliceosome
(Complex A)

Spliced mRNA

U2
U5 U6
Post-catalytic Spliceosome

U1 U2
U5
U4 U6
Pre-catalytic Spliceosome
(complex B)

U2
U5 U6
Complex B*

U2
U5 U6
Lariat
Catalytic Spliceosome
(Complex C)

C

THP1 macrophages

Infection

H37Rv

Mild lysis,
Cytosolic and
Nuclear extract

IP *In vivo* assembled
complex (Anti-Y12)
LC-MS

D

Spliceosome Complex proteomics

EIF5A
EIF5AL1
TADA2B
RPL35A
SRSF3
ALYREF
SNRPC
SNRPB
SNRPN
SNRPA
SNRPD2
HNRNPL
HNRNPA3
HNRNPH3
HNRNPDL
MAGOHB
MAGOH
TRA2B
SRSF7
HNRNPA2B1
RPL4
SF3B2
SYNCRIP
ILF3
HMGA1
EIF4A3
LMNA

Depletion

Depletion and mislocalization

Mislocalization

Cyt Nuc   Cyt Nuc
Inf   UI

1.0
0.5
0
-0.5
-1.0

A

B

C

Figure 3

A

healthy controls (n=19)
PTB (n=20)

B

SNRPF staining (ITB biopsy)

1   **Supplementary information for the manuscript:**

2   **A *Mycobacterium tuberculosis* secreted virulence factor disrupts host**
3   **snRNP biogenesis**
4
5   Chauhan, K et al.
6
Supplementary methods
Supplementary Figures and Legends to Figures S1-S6
Supplementary tables S1-S7

Ethics statement: Studies on human samples were approved by the IEC of AIIMS, New Delhi

(Ref no. IEC-304/02-06-2017). For animal experiments, approval was obtained from the

Institutional Animal Ethics Committee, ICGEB, New Delhi (ICGEB/IAEC/22092022/CI-28).

Biological materials: Tissue sections from Intestinal TB biopsy samples used in the study

were obtained from the Department of Pathology, AIIMS, New Delhi. Access to these

materials is subject to institutional guidelines. **Clinical diagnosis of ITB subjects:** The

diagnoses depend on clinical features of nocturnal fever, weight loss, diarrhoea, and

radiological suspicion of short-segment bowel thickenings, narrowed ileocaecal region,

significant mesenteric lymphadenopathy, and/or necrosis. In such cases, an intestinal biopsy is

taken, or a segment of the intestine is resected if there is a feature of intestinal obstruction due

to fibrosis. All our cases underwent intestinal resection. Histological examination is critical in

such cases, and demonstration of large, confluent, multiple, necrotic granulomas with

multinucleated giant cells in deeper layers of the intestine are suggestive of ITB, as we have

demonstrated in our cases. ZN stain has a very low sensitivity, varying from 5-10%. Such cases

are confirmed by either the Gene-Xpert test, Mycobacterial culture, or where such tests are not

possible, by ATT trial followed by re-evaluation at two months and six months. All our cases

underwent either a *Mycobacterial* culture or a gene-Xpert test.

Antibodies and reagents

The antibodies used in the immunofluorescence experiments were as follows: SNRPD2

(#NBP2-95236), SNRPF(#NBP1-57463), LSm7 (#NBP1-92084), Gemin3 (#NBP2-52454)

and Gemin8(#NBP1-83925) from Novus biologicals.DDX23(#A300-693A) from Bethyl

Laboratories were used. PKH26 red and PKH67 green fluorescent labelling dyes were used

from Sigma-Aldrich. The spliceosome complex pull-down was done using an anti-SNRPB

antibody (Y-12, MA513449). The antibodies used in immune-blotting experiments were as

tubulin (Abcam plc; ab6046), histone (Abcam plc; ab18521), IRDye 800CW Goat anti-mouse

IgG secondary antibody (LI-COR Biosciences)and IRDye 680RD goat anti-rabbit IgG

secondary antibody(LI-COR Biosciences). The co-immunoprecipitation was done using

EZView Red Anti-HA affinity gel (E6779), Anti-FLAG M2 Affinity gel (A2220) and

immunoblots were detected using Anti-HA antibody, Mouse monoclonal (H3663) and

Monoclonal Anti-FLAG M2 antibody mouse (F1804) from Sigma-Aldrich.

Cell culture

Mammalian cells: THP1 monocytes (ATCC) were maintained in RPMI 1640 media (Cell

Clone), and HEK293T cells (ATCC) were maintained in DMEM media (Cell Clone). Cells

were grown at 37°C with a 5% CO2 atmosphere. All the media was supplemented with 10%

fetal bovine serum. HEK293T cells were transfected with plasmids at a 30-50% confluency

using jetPRIME buffer (Polyplus #712-60) as per manufacturer protocol. The siRNA was

transfected using the Dharmafect reagent. HEK293T constitutively expressing PRINT plasmid

was selected in the presence of blasticidin (Gibco) 20 μg/ml concentration.

Bacterial Cells: *M. tuberculosis H37Rv*, *M. smegmatis* mc^{2}155 and mutant strains were grown

at 37°C to log phase in Middlebrook 7H9 media with 0.05% Tween80, 10% ADC Enrichment,

and 0.2% glycerol. Plasmids were selected with 50 mg/ml kanamycin or hygromycin

depending upon the resistance marker.

Infection

THP-1 cells were activated into macrophages by treatment with 32 nM PMA (Sigma) for 24

hours, followed by another 24 hours for recovery. The THP-1-derived macrophages were

infected with *Mtb* strains at an MOI of 10 for 4 hours, followed by 2 hours of amikacin (200

μg/μl) treatment. *M. smegmatis* infection was performed at an MOI of 10 for 1 hour, followed

by 2 hours of amikacin treatment. Amikacin treatment clears all the extracellular bacteria,

which are washed following the treatment, and this time point is considered as 0 hours of

infection throughout the manuscript.

RNA isolation and qPCR

RNA was isolated at indicated time points using MDI mammalian Total RNA isolation kit

(MTRK250) as per the manufacturer's protocol. The RNA was quantified using a nanodrop

spectrophotometer. 1000ng RNA was used to make cDNA using the Iscript cDNA synthesis

kit (BioRad #170-8891), and 2 μl of 1:5 diluted cDNA was used to set qPCR on the BioRad

CFX384 well plate qPCR machine. Primers were custom synthesized from Sigma Aldrich
Chemicals Ltd. A list of primers used in this study is provided in the table S7.

Transformation of yeast cell

The yeast competent cells were prepared by harvesting yeast cells by centrifugation at 400 rpm
for 5 min at 25 °C. The supernatant was discarded, and the pellet was washed and resuspended
in filter-sterilized SORB buffer (100 mM Lithium acetate, 100 mM Tris HCl, pH 8.0, 1 mM
EDTA, pH 8.0, 1 M sorbitol pH 8.0) to which denatured salmon sperm DNA was added (10
mg/ml). The bait (pGBDKT7 vector) plasmids were mixed gently with the Y187 competent
cells. Then PEG mixture was added and incubated at 30 °C for 30 min, 42 °C for 30 min and
4 °C for 30 min in a thermocycler machine. Tubes were taken out and plated on 15 cm diameter
SC-TRP plates (Cat No. Clonetech 630413). Plates were incubated at 30°C for 2 to 7 days. A
similar protocol was used for the prey (pGADT7 vector) plasmid in the AH109 yeast strain on
SC-Leu medium (Cat No. Clonetech 630414).

Y2H assay between bait and prey libraries

The bait library was prepared by plating the Y187 yeast strain (bait) transformed with Mtb
proteins in 384 well patterns on the Singer Robot Plate. The AH109 (prey) yeast strain
transformed with host spliceosome protein was grown in SC-leu broth medium and equally
plated on a robotic compatible square plate. Using 384 pinned robotic pad bait libraries were
patched on another YAPD plate. Each plate was then patched with individual cells containing
prey plasmid and mixed well using the robotic system. This step was performed 4 times as four
independent mating steps, thereby making it a total of 1536 spots per plate. The plate containing
mixed colonies was kept for mating of both strains at 30 °C for 36 hours. After 36 hours these
colonies were selected on SC-Trp-Leu (Cat No. Clonetech 630417) plates for the selection of
those cells which contain both plasmids (bait as well as prey). After 72 hours these plates were
replicated on SC-Trp-Leu-His (3 drops-out, Clonejet 630419) and SC-Trp-Leu-His-Ade (4
drop-outs, Clonetech 630428) for the screening of interaction between the protein which was
fused with bait and prey respectively. The colonies that appear on both 3-DO and 4-DO plates
represented very specific and strong interactions between the bait and prey proteins.

Co-immunoprecipitation and Immunoblotting

The cells were lysed using a non-denaturing IP lysis Buffer (20 mM Tris-Cl, 137 mM NaCl, 2
mM EDTA, 10% glycerol, 1% NP-40, pH 8) and centrifuged at 14000 rpm for 10 min at 4 °C.

The supernatant was precleared using IgG beads, and 300 µg of protein was used for IP. The

IP was performed using FLAG-tagged beads (Sigma) and HA-tagged beads (Sigma) at 4 °C

for 4 hours, followed by three washes with IP buffer. The beads were then boiled in 2X SDS

loading dye. The protein samples were subjected to SDS PAGE followed by transfer on

nitrocellulose membrane (BioRad) using BioRad Turbo transfer blot. The blocking was

performed by Odessey blocking buffer (LI-COR Biosciences) in 1:1 dilution with 1X PBS at

room temperature. The blots were immunoblotted with primary antibodies, followed by IR-

tagged secondary antibodies (LI-COR Biosciences). Blot imaging was done using Odessey IR

Imaging System (LI-COR Biosciences).

Immunofluorescence

For immunofluorescence experiments, ~0.2x 10^6 cells were plated on 12 mm coverslips per

well in a 24- well plate. Cells were infected with m-cherry expressing *Mtb*, and other strains

were labelled with PKH67 fluorescent dye (Sigma). The cells were fixed using 4% v/v

paraformaldehyde for 20 min at room temperature for indicated time points. The cells were

incubated with blocking buffer (0.02% Triton-X100, 3% w/v BSA in 1XPBS) for 1 hour. The

cells were washed thrice with 1X PBST and incubated with primary antibodies for 3 hours.

The cells were washed thrice with 1XPBST and incubated with secondary Alexa Flour (488,

555, 590 or 670nm) tagged antibodies (Invitrogen). The nucleus was stained using DAPI (300

nM) for 10 min. The cells were mounted on slides with Pro-Long Gold antifade reagents (Life

Technologies). All the incubation and staining were done at room temperature. Images were

acquired by NIS-Elements software using the Nikon A1R laser scanning confocal microscope

equipped with a Nikon Plan Apo VC ×20, NA 0.75, and Plan Apo VC ×100 oil, NA 1.40

objectives were used. Serial confocal sections, 0.5 µm thick, were acquired with a z-stack

spanning 10–15 µm to form a composite image. Images were analysed using Imaris 9.6.6

imaging software using several tools like deconvolution, 3D-reconstruction and shortest

distance calculation amongst others, NIS-Elements and image J software.

Fluorescent reporter splicing assay using flow cytometry

For Flow cytometry experiments, ~0.5 x 10^6 HEK293T cells constitutively expressing pRINT

vector (Addgene plasmid # 24218) were plated on 12 well plates. These were co-transfected

with 800ng of modified pGINT vector (Addgene plasmid # 24217) with exon skipping cassette

and *Mtb* protein-expressing vectors at a ratio of 1:1 using jetPRIME buffer. The cells were

fixed using 4% v/v paraformaldehyde in 1X PBS for 20 minutes at room temperature. Then,

the cells were scraped and resuspended in 1X PBS. The data were acquired using BD

FACSDiva acquisition software in BD FACS Fortussa cytometer (BD).

Spliceosome complex Co-IP

For the pulldown experiment, ~200 million THP-1 derived macrophages were infected with

H37Rv at an MOI of 10 and samples were processed 48-hour post-infection. The nuclear and

cytoplasmic extract was prepared using the Active Motif Nuclear Complex Co-IP kit

(catalogue No.54001) per the manufacturer's guidelines. The nuclear extract and cytoplasmic

extract were passed through a 0.2-micron filter. 500 µg of nuclear extract and cytoplasmic

extracts were used for pulldown using 10 µg Y-12 antibody (MA5-13449, Thermo Fisher) or

IgG-control (Sigma) antibody for each condition. The Y-12 antibody was crosslinked using

DMP (20 mM) to Protein A-Sepharose beads for 1 hour (beads were washed prior with 0.2 M

sodium borate, pH 9.0) followed by 0.2 M ethanolamine incubation for 1 hour at room

temperature. The Nuclear and Cytosolic extracts were then incubated with crosslinked beads

overnight at 4 °C. The beads were washed thrice with wash buffer (0.025 M Tris, 0.15 M NaCl,

pH 7.2). The protein was finally eluted in 50 µl of an elution buffer containing primary amine

pH 2.8. To neutralize the low pH of the elution buffer, 1 M Tris (pH 9.5) was added to the

collection tube.

Mass Spectrometric Data analysis for immune-precipitated spliceosome complex

Sample Preparation- An equal volume of eluted protein sample (50ul) was used for digestion

and reduced with 5 mM TCEP and further alkylated with 50 mM iodoacetamide and then

digested with Trypsin (1:50, Trypsin/lysate ratio) for 16 h at 37 °C. Digests were cleaned using

a C18 silica cartridge to remove the salt and dried using a speed vac. The dried pellet was

resuspended in buffer A (2% acetonitrile, 0.1% formic acid). An equal amount of peptide (1ug)

was injected into the analytical column. *Mass Spectrometric Analysis of Peptide Mixtures*- All

the experiments were performed using EASY-nLC 1000 system (Thermo Fisher Scientific)

coupled to Thermo Fisher-QExactive plus equipped with nano electrospray ion source. 1ug

peptide was loaded on C18 column 50 cm, 3.0µm Easy-spray column (Thermo Fisher

Scientific) at column heating temp 40 C. Peptides were eluted with a 0–40% gradient of buffer

B (80% acetonitrile, 0.1% formic acid) at a flow rate of 300 nl/min) and injected for MS

analysis. LC gradients were run for 90 minutes MS1 spectra were acquired in the Orbitrap

(Max IT = 50ms, AGC target = 3e6; R=70K, mass range = 350-1750; Profile data). Dynamic

exclusion was employed for 30s excluding all charge states for a given precursor. MS2 spectra

were collected for top 15 peptides. MS2 (Max IT= 100ms, R= 17.5K, AGC target 1e5). *Data*

Processing- All samples were processed and RAW files generated were analysed with

Proteome Discoverer (v2.2) against the Uniprot Human and Mtb database. For dual Sequest

and Amanda search, the precursor and fragment mass tolerances were set at 10 ppm and 0.5

Da, respectively. The protease used to generate peptides, i.e. enzyme specificity was set for

trypsin/P (cleavage at the C terminus of "K/R: unless followed by "P"). Carbamidomethyl on

cysteine as fixed modification and oxidation of methionine, N-terminal acetylation were

considered as variable modifications for database search. Both peptide spectrum match and

protein false discovery rate were set to 0.01 FDR. Only statistically significant proteins (Using

ANNOVA p-value cut off 0.05) that were found in all three biological replicates across and in

all experimental groups were used to generate heatmap and for further analysis.

RNA FISH

RNA-fluorescence *in situ* hybridization (RNA FISH) was performed using previously

described snRNA probes [1]. The sequences of the oligonucleotide used for FISH were as

follows: U5 snRNA, 5′-CTC TCC ACG GAA ATC TTT AGT AAA AGG CGA AAG ATT

TAT ACG ATT TGA AGA G-3′; The U5 snRNA probes were labelled with Alexa Flour 647

(Invitrogen) at their 5' end. Next, the cells were fixed with 4% v/v paraformaldehyde for 20

min at room temperature. Next, cells were permeabilized and incubated with RNA-FISH probe

for three hours at 40 °C by using the ViewRNA™ ISH Cell Assay Kit system (Thermo

Fischer). Subsequently, cells were immune-stained and mounted using a Pro-Long gold

antifade (Life technologies).

Bacterial transformations

For the generation of *M. smegmatis* mutants expressing *H37Rv* proteins, 50 ml of an early

exponential phase culture of *M. smegmatis* mc^{2}155 was washed twice and concentrated ice-

cold 10% ultrapure glycerol (Invitrogen). Concentrated cells were electroporated with

approximately 10 μg of pNit plasmid. Electroporation was carried out at 2500 V, 25 μF, and

1000 Ω in the Biorad gene pulsar. Electroporated cells were allowed to recover at 37 °C in 7H9

Middlebrook broth and plated on 7H11 supplemented with 10% OADC and kanamycin (25

μg/ml; Sigma).

Mtb survival assay

For CFU experiments, ~0.01 million THP-1 macrophages were seeded in a 96-well plate and *Mtb* infection was done at an MOI of 10. The lysis was done by incubating with 0.06% SDS for 5 min at indicated time points. The lysed cells were diluted to 1:10 and 1:50 in 7H9 media and 10 µl of the dilutions were plated in a drop-wise fashion on a 7H11 plate supplemented with 10% OADC. The plates were incubated for approximately around 3 weeks for colony enumeration at 37 °C.

Expression and purification of recombinant proteins

The Mtb proteins were cloned in the pEZY19 vector and transformed into E. coli BL21(DE3) competent. Protein expression was induced with 0.1 mM isopropyl b-D-1-thiogalactopyranoside (IPTG) for 16-19 hours. Collected frozen cells were lysed in a lysis Buffer (50 mM Tris-HCl pH. 8.0 containing 300 mM NaCl,10% Glycerol, 1mm DTT, 1mm PMSF,1mm BHCL,0.1% Triton X-100, 10mM imidazole and lysozyme 1mg/ml) followed by sonication (0.5 s pulse and 0.9 s rest) at 2 min intervals to prevent overheating. After centrifugation, the proteins were purified from the supernatant using nickel-nitrilotriacetic acid (Ni-NTA) affinity chromatography (Cat. No.- 30410, Sigma). The recombinant proteins were eluted from Ni-NTA resins by elution buffer (50 mM Tris-HCl pH. 8.0 and 300 mM NaCl) containing 300mm Imidazole. The purified protein was concentrated using 10Kda Centricon (ACS501024, Millipore) subjected to gel permeation chromatography on GE Healthcare Superdex 75 Increase 10/300 GL where the monomeric fraction of the protein was eluted. The final protein was stored in 50 mM Tris-HCl pH. 8.0 containing 300 mM NaCl,10% Glycerol buffer at -80˚C.

BALB/c aerosol challenge

All mice experiments were carried out in the Tuberculosis Aerosol Challenge Facility (TACF, ICGEB, New Delhi, India). BALB/c mice have been housed in individually ventilated cages contained within the biosafety level 3 enclosure. The animal holding area of the BSL3 lab is maintained at 20–25 °C, 30–60% humidity and 12–12 h of the light–dark cycle. An aerosol challenge of ~100 CFUs was given to the animals in a Wisconsin–Madison chamber according to the standardized protocol in the TACF facility. Two animals were selected randomly and humanely euthanized 24 h post aerosol challenge to check for infection establishment. The lungs and spleen tissues were harvested and homogenized to enumerate CFU. Tissue lysates

were serially diluted and plated on Petri plates containing Middlebrook 7H11 agar (Difco)

supplemented with 10% OADC (Becton, Dickinson) and 0.5% glycerol.

RNA-Seq analysis

Reads were quality filtered using Trimmomatic (v.0.39, LEADING:10 TRAILING:10

SLIDING WINDOW:4:15 MINLEN:75) [2] and were assessed for quality using FastQC (v.10.1)

[3] and visualised with MultiQC (v.1.9)[4]. Salmon (v.1.8, –numGibbsSamples 30) [5] was used to

quantify quality filtered reads against the human genome (GRCh38, primary assembly) with

Gencode annotation (v.40) [6]. Profiles were imported using tximeta (v.1.10) [7] and [8] (v.1.22) and

filtered to retain all transcripts from only protein-coding genes. Statistical analysis was done

using swish methodology from fishpond (v.1.8) [9] and significantly differentially expressed

transcripts (DETs) were defined based on *qvalue* ≤ 0.1. Furthermore, Isoform usage was

analysed using IsoformSwitchAnalyzeR (v.1.18) [10] with reduceToSwitchingGenes = FALSE,

reduceFurtherToGenesWithConsequencePotential = FALSE, No dIFcutoff and Isoform

fraction cuffoff. Heatmap of the DETs was generated with [11] using mean z-scaled normalised

scaled TPM values and clustered based on Euclidean distance and Ward.D2 methodology.

Clusters were defined manually. Enrichment analysis was done separately on overexpressed

and repressed DETs from each pairwise comparisons using enrichR (v.3.0) [12] with KEGG

Human 2021 and Gene Ontology (GO) [13] (Biological Process (BP), Cellular Component (CC)

and Molecular Function (MF)) 2021 database figures were generated using ggplot2 (v.3.3.5)

[14]. Analyses were done in R statistical software environment (v.4.1) [15].

Generation of Rv1435c mutant

Rv1435c mutant was generated using recombineering methodology which requires an Allelic

exchange substrate [16]. To construct that, 836bp upstream (LHS) and 1Kb downstream (RHS)

regions of the *Rv1435c* locus (retaining ~170 bp of the gene at both ends) were amplified from

Mtb (*H37Rv* genome) with the help of primers having appropriate restriction endonuclease sites

(LHS-DraIII; RHS-SfiI). The flanks were then digested and ligated with other two fragments-

Hygromycin resistant cassette and *oriE*+cosλ with compatible ends to create a four-piece AES.

The latter was then digested with SnaBI sites to release the linearised fragment consisting of

LHS-*Hyg^r*-RHS and electroporated into recombineering proficient *Rv::pNitET* (Kan^r) strain

that overexpresses RecET recombinases upon over-expression using Isovaleronitrile. Mutant

colonies were analysed using different primer sets to confirm legitimate recombination at the

required locus and cured of the pNitET plasmid by negative selection on 2% sucrose-containing

7H11-hyg^r agar plates.

Primers used:

Rv1435c 5' Flank forward primer with DraIII & SnaBI site:

5'CACCTTTTCACAAAGTGTACGTACGTGGTGCACGACGCATTGGA 3'

Rv1435c 5' Flank reverse primer with DraIII site:

5' TTTTTTTTCACTTCGTGCGCCGGCGGCTACCGGCGCGT 3'

Rv1435c 3' Flank forward primer with SfiI site:

5' CACCTTTTGGCCTAGATGGCCGGGCCGGTCCTGGCTGGTGCT 3'

rv1435c 3' Flank reverse primer with SfiI site and SnaB1 site:

5' TTTTTTTTGGCCACTTAGGCCTACGTAGGGATCGCTGGGAGCGTGGTT 3'

Generation of Rv1435c complementation construct pSCW-F-Rv1435c

Rv1435c was cloned along with its native promoter (500bp upstream of the gene) between ScaI

and HindIII sites using an integrative vector, pSCW-F (C-terminal 3X FLAG tag), which

facilitates the integration of the gene at Tweety attP sites in the *Mtb* genome.

Primers used:

FP *Rv1435c* (with UPSTREAM 500bp) ScaI

5' FP : 5' CACCAGTACTATGCGCCAGCGTGCTGTTG 3'

RP *Rv1435c* HindIII, w/o stop codon

5'TTTTAAGCTTGCGCGCACCTGCGGGC 3'

Immunohistochemical staining: The paraffin-embedded specimens were cut into 4–5 μm-

thick sections, deparaffinized in xylene for 5–10 min, and dehydrated through a series of graded

alcohols (100%, 100%, 95% and 80%). Tissue sections were boiled at 100^{O}C for 30 min in

10mM citrate buffer (pH 6.0), followed by cooling and treatment with blocking solution (100%

methanol containing 0.3% H_2O_2) for 30 min to block any endogenous peroxidase activity. They

were incubated with SNRPF or SNRPD2 primary antibody (1:600, pH 9) overnight at 4°C,

followed by incubation with secondary antibody (Ultra trek anti polyvalent kit with HRP

conjugated secondary kit, Scy Tek Laboratories, Logan, USA) for 30 mins at room

temperature. The slides were thereafter stained with 3,3′-Diaminobenzidine (DAB). Finally,

the sections were counterstained with Hematoxylin and Eosin and observed under a

microscope. **Interpretation of IHC results:** SNRPF staining in >10% of cells was considered

positive. Both cytoplasmic and nuclear staining were looked for. The spatial distribution of the

stained sections, in terms of granulomas areas and areas away from granulomas, were noted

down to identify polarization of staining, if any.

Legends to Figures S1-S9:

Figure S1: Mycobacterial secretory proteins and host spliceosome proteins interactome.

A) The percentage of the interaction shown by spliceosome-related proteins belonging to different spliceosome protein categories

B) The percentage of the interacting Mtb proteins belonging to the different functional classifications.

Gene	No. of isoform in Y2H	Interacting transcript	Interaction
SNRP40	2	0	No
PQBP1	2	0	No
NCBP2	2	0	No
POLR2H	4	4	19/10/4/1
SF3B1	2	0	No
SYF2	2	2	2 (Rv1643, Rv2927c)
DCAF4	2	1	2 (Rv1643, Rv2927c)
SNRPD2	2	1	16 (Rv1643, Rv2927c, Rv1566c)
LSm7	2	1	26 (Rv3864, Rv1643, Rv2927c, Rv1566c)
LSm4	2	0	No
SNRPG	3	0	No
RBPMS	2	0	No
SF1	3	0	No
CSTF	2	0	No
LUC7L2	2	1	4 (Rv1643, Rv2927c)
RBMX	2	1	8 (Rv3849)
U2AF1	3	1	2 (Rv1643, Rv2927c)
PPIH	3	1	1

305
306 **Figure S2: Interaction of Mtb proteins with the splice variants of human proteins.**
307

A) A descriptive list of the human spliceosome proteins, represented by more than one isoform in the AD library, along with their interactions with the *Mtb* protein, is presented in the table.

B) The table shows the distinct spliced mRNA isoforms that are classified as loss of exon, mutually exclusive exon and loss of the functional domain. The isoforms of SYF2, differing by only one exon, show interaction with the same *Mtb* proteins, suggesting no contribution of alternatively spliced exons towards interaction. Only one of the isoforms of genes which show exon skipping, like DCAF4 and PPIH, or mutually exclusive exons like U2AF1 and LUC7L2, showed interaction with any *Mtb* proteins, suggesting the involvement of alternative exons in the interactions. The truncated isoforms of SNRPD2, LSm7 and RBMX show a loss of functional domains and multiple interactions with Mtb proteins, while the longer isoforms for these genes did not show any interactions. Similarly, the truncated isoforms of POLR2H show more interaction.

Figure S3: Altered RNA splicing in *Mtb*-infected THP-1 macrophages

A) The schematic shows the primer design strategy for exon-inclusion, exon-exclusion and exon-junction primers.

B) Expression of five target genes (as indicated) in THP-1 macrophages upon infection with H37Rv using exon exclusion primers at 48 hpi. Data from three independent biological experiments.

C) Expression of five target genes (as indicated) in THP-1 macrophages upon infection with H37Rv using exon inclusion primers at 48 hpi. Data from three independent biological experiments.

D) Expression of five target genes (as indicated) in THP-1 macrophages upon infection with H37Rv using exon junction primers at 48 hpi. Data from three independent biological experiments.

Figure S4: Rv1435 secretion from the modified *M.smegmatis* strains

A) The protein expression in modified *M. smegmatis* strains expressing Rv1435c upon induction with 2.5µM and 5 µM IVN was checked by immunoblotting in

whole cell lysate and culture supernatant (SN). As control, the modified *M. smegmatis* strains expressing pknB were also induced with IVN and its expression in the supernatant or whole cell lysate was measured by immunoblotting.

B) Isoform-specific qPCR to assess the impact of Rv1435c expressing *Msmeg* infection in THP-1 macrophages on exon skipping event in *ACADVL*, *DDX42*, *ACSL1* and *NFYA*; N=3, indicating p values from One-way ANOVA.

C) Domain organization of Rv1435c depicting signalP, extracytoplasmic and disordered regions.

D) Predicted structure of Rv1435c from AlphaFold Protein Structure Database

E) Multiple sequence alignment of Rv1435c protein in different mycobacterial species using CLUSTAL Omega. The red bar indicates the conserved sequence among different species.

F) A phylogenetic tree corresponding to the multiple sequence alignment is shown in S6C.

Figure S5: Dysregulation of splicing proteins upon Mtb infection.

A) Representative images for RNA FISH against U2 snRNA in THP-1 macrophages upon *Mtb* (mCherry-H37Rv) infection at 24- and 48-hrs post-infection. The U5 snRNA probe was labelled with Alexa 488 (green). Plots on the right show quantification of U2 snRNA FISH MFI in uninfected and *Mtb*-infected macrophages at 24 ad 48 hpi.

370   B) Representative image for staining of DDX23, GEMIN3, GEMIN8, LSm7 and
371   SNRPD2 in uninfected and H37Rv infected THP-1 macrophages 48 h post-
372   infection.
373
374

375
376 **Figure S6: Regulation of host alternative splicing as a virulence mechanism**
377
A) Western blot showing the specificity of anti-sera raised against Rv1435c.
B) The plot represents the bacterial CFU in WT THP-1 and THP-1::SNRPD2 at 48hr
post-infection.
C) PCA analysis of significant transcripts (splice isoforms) identified by RNA-Seq in
monocytes of Human TB patients.
D) Human intestinal tuberculosis granulomatous lesions were stained with SNRPF.
Intense cytosolic staining in multinucleated giant cells (black arrow) is shown.
E) Representative image for IHC staining of SNRPF in gut sections from healthy
controls.

F) Representative image for IHC staining of SNRPD2 in gut sections from healthy
controls and in intestinal tuberculosis granulomatous lesions from diseased
individuals.
390
391

393
394
395
396   **Supplemental Tables:**
397
Table S1: Spliceosome proteins identified in Y-12 pulldown MS analysis
Table S2: List of Mtb genes in the bait library (BD library) and human splicing factor coding
transcripts in the prey library (AD library).
Table S3: List of genes included in the BD library as negative controls
Table S4: Yeast-2-hybrid interaction matrix for AD and BD proteins
Table S5: Two-column file for the interaction network shown in Fig 2B.
Table S6: RNAseq data shown in Fig. 8A.
Table S7: List of primers
406

407
408

Supplementary References

1. Liu, J. L. & Gall, J. G. U bodies are cytoplasmic structures that contain uridine-rich small nuclear ribonucleoproteins and associate with P bodies. *Proc Natl Acad Sci U S A* **104**, 11655-11659 (2007). https://doi.org/10.1073/pnas.0704977104

2. Bolger, A. M., Lohse, M. & Usadel, B. Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics* **30**, 2114-2120 (2014). https://doi.org/10.1093/bioinformatics/btu170

3. Andrews, S. (2017).

4. Ewels, P., Magnusson, M., Lundin, S. & Kaller, M. MultiQC: summarize analysis results for multiple tools and samples in a single report. *Bioinformatics* **32**, 3047-3048 (2016). https://doi.org/10.1093/bioinformatics/btw354

5. Patro, R., Duggal, G., Love, M. I., Irizarry, R. A. & Kingsford, C. Salmon provides fast and bias-aware quantification of transcript expression. *Nat Methods* **14**, 417-419 (2017). https://doi.org/10.1038/nmeth.4197

6. Frankish, A. *et al.* GENCODE reference annotation for the human and mouse genomes. *Nucleic Acids Res* **47**, D766-D773 (2019). https://doi.org/10.1093/nar/gky955

7. Love, M. I. *et al.* Tximeta: Reference sequence checksums for provenance identification in RNA-seq. *PLoS Comput Biol* **16**, e1007664 (2020). https://doi.org/10.1371/journal.pcbi.1007664

8. Martin Morgan, V. O., Jim Hester, Hervé Pagès. SummarizedExperiment: SummarizedExperiment container. *R package version 1.28.0* (2022). https://doi.org/10.18129/B9.bioc.SummarizedExperiment

9. Zhu, A., Srivastava, A., Ibrahim, J. G., Patro, R. & Love, M. I. Nonparametric expression analysis using inferential replicate counts. *Nucleic Acids Res* **47**, e105 (2019). https://doi.org/10.1093/nar/gkz622

10. Vitting-Seerup, K. & Sandelin, A. IsoformSwitchAnalyzeR: analysis of changes in genome-wide patterns of alternative splicing and its functional consequences. *Bioinformatics* **35**, 4469-4471 (2019). https://doi.org/10.1093/bioinformatics/btz247

11. Kolde, R. Pheatmap: pretty heatmaps. *R package version v.1.0.12* (2019).

12. Chen, E. Y. *et al.* Enrichr: interactive and collaborative HTML5 gene list enrichment analysis tool. *BMC Bioinformatics* **14**, 128 (2013). https://doi.org/10.1186/1471-2105-14-128

13. Ashburner, M. *et al.* Gene ontology: tool for the unification of biology. The Gene Ontology Consortium. *Nat Genet* **25**, 25-29 (2000). https://doi.org/10.1038/75556

14. Wickham, H. ggplot2: Elegant Graphics for Data Analysis. *Springer-Verlag New York* (2016).

15. Team, R. C. R: A Language and Environment for Statistical Computing. (2022).

16. van Kessel, J. C. & Hatfull, G. F. Recombineering in Mycobacterium tuberculosis. *Nat Methods* **4**, 147-152 (2007). https://doi.org/10.1038/nmeth996

17. Xu, C. *et al.* Structural insights into Gemin5-guided selection of pre-snRNAs for snRNP assembly. *Genes Dev* **30**, 2376-2390 (2016). https://doi.org/10.1101/gad.288340.116