One-step N-terminomics based on isolation of protein N-terminal peptides from LysargiNase digests by tip-based strong cation exchange chromatography

Kazuya Morikawa[1], Hiroshi Nishida[1], Koshi Imami[1,2], Yasushi Ishihama[1,3]

1) Department of Molecular Systems BioAnalysis, Graduate School of Pharmaceutical Sciences, Kyoto University, Kyoto 606–8501, Japan

2) Proteome Homeostasis Research Unit, RIKEN Center for Integrative Medical Sciences, Yokohama 230-0045, Japan

3) Laboratory of Clinical and Analytical Chemistry, National Institute of Biomedical Innovation, Health and Nutrition, Ibaraki, Osaka 567-0085, Japan.

*Correspondence and requests for materials should be addressed to Y.I. (email: yishiham@pharm.kyoto-u.ac.jp).

Keywords

Protein terminal peptides, LysargiNase, Strong cation exchange chromatography, StageTip

In Brief (80 words)

Large-scale isolation of protein N-terminal peptides from LysargiNase digests was achieved by one-step strong cation exchange chromatography with disposable and parallel-processable pipette-tip columns. Novel cleavage sites for signal or transit peptides and non-canonical translation initiation sites were identified by this tip-based method. Together with two other methods similarly consisting of protease digestion and one-step tip LC, we have established a simple and comprehensive N- and C-terminomics platform based on CHAMP (chromatographic amplification of protein terminal peptides) approaches.

Highlights

- One-step isolation of protein N-terminal peptides from LysargiNase digests by tip-based strong cation exchange chromatography.
- Highly efficient isolation of protein N-terminal peptides by tip LC comparable to HPLC.
- Identifying novel post-translational cleavage sites and non-canonical translation initiation sites.
- Complementary terminome coverage by the three CHAMP methods consisting of protease digestion and one-step tip LC**.**

Abstract

We have developed a one-step isolation method for protein N-terminal peptides from LysargiNase digests by pipette tip-based strong cation exchange (SCX) chromatography. This CHAMP-N (CHromatographic AMplification of Protein N-terminal peptides) method using disposable and parallel-processable SCX tips instead of conventional HPLC SCX columns facilitates simple, sensitivie and high-throughput N-terminomic profiling without sacrificing identification numbers and selectivity achieved by the HPLC-based method. By applying the CHAMP-N method to HEK293T cells, we identified novel cleavage sites for signal and transit peptides, and non-canonical translation initiation sites. Finally, for proteome-wide terminomics, we have established a simple and comprehensive N- and C-terminomics platform based on three different tip-based approaches including CHAMP-N, in which protease digestion and one-step isolation by tip LC are commonly used to achieve complementary terminome coverage.

Introduction

Proteins from a single gene have various sequences and modifications generated by several mechanisms, including alternative splicing, mutation, translational control, proteolysis and co- or post-translational modifications. These diverse protein isoforms are called proteoforms (1–3). Each proteoform has different function, localization, and stability (4). For example, it has been reported that proteoforms derived from non-canonical translation initiation sites affect tumorigenesis (5–7), and that proteoforms cleaved at different sites of the amyloid precursor protein are involved in the development of Alzheimer's disease (8). Each of these proteoforms resulting from different translation initiation sites or proteolytic cleavages has a different protein terminal sequence. Moreover, localization and stability are affected by the signal or transit peptide sequences and N- or C-degrons (9), present at the protein termini. Thus, the comprehensive analysis of protein termini is the starting point for understanding the functions of diverse proteoforms.

Comprehensive analysis of protein termini, or terminomics, is based on bottom-up proteomics, in which protein terminal peptides are isolated from other digested non-terminal peptides and subjected to liquid chromatography/tandem mass spectrometry (LC/MS/MS). Many methods to isolate protein N- and C-terminal peptides have been developed. COFRADIC (combined fractional diagonal chromatography) (10) and TAILS (terminal amine isotopic labeling of substrates) (11) are widely known as N-terminal peptide isolation methods. Even recently, new methods for N-terminal peptide isolation have been reported (12). However, all of these methods require complex and time-consuming procedures involving chemical modifications that block amines, and the efficiency and selectivity of the

chemical modifications are also issues. We have recently developed simple isolation methods for protein N- and C-terminal peptides that consist only of enzymatic digestion of proteins followed by chromatographic separation; CHromatographic AMlification of Protein N- and C-terminal peptides (CHAMP-N (13), CHAMP-C (14)) methods. With these methods, terminal peptides can be isolated in a single step at high selectivity without complex steps as in the conventional methods. In the CHAMP-C method, C-terminal peptides can be easily isolated from V8 protease digests using pipette-tip columns packed with CeO_2 particles. On the other hand, the CHAMP-N method, which uses HPLC-based strong cation exchange (SCX) chromatography for LysargiNase digests, can be fully automated, but it requires dedicated equipment and has low-throughput because it analyzes one sample at a time in series.

In this study, we developed a simple, high-throughput protein N-terminal peptide isolation method using pipette-tip based SCX tips instead of HPLC columns from LysargiNase digests, and applied it to the identification of novel post-translational cleavage sites (signal peptide and transit peptide cleavage sites) and non-canonical translation initiation sites. Furthermore, we established a comprehensive terminomics platform by combining the LysargiNase/SCX tip-based CHAMP-N method with the V8 protease/CeO_2 tip-based CHAMP-C method (14) and the trypsin/SCX tip-based CHAMP-NC method (15–19).

Experimental procedures
Materials.

LysargiNase was purchased from Merck Millipore (Burlington, VT). Empore cation-SR disks and SDB-XC Empore disks were purchased from GL Sciences (Tokyo, Japan). UltraPure Tris Buffer and fetal bovine serum were purchased from Thermo Fisher Scientific (Waltham, MA). Protease inhibitors were purchased from Sigma-Aldrich (St. Louis, MO). TPCK-treated sequencing-grade modified trypsin was obtained from Promega (Madison, WI). Polyethylene frit was purchased from Agilent Technologies (Santa Clara, CA). Water was purified using a Millipore Milli-Q system (Bedford, MA). All other chemicals were purchased from FUJIFILM Wako (Osaka, Japan), unless otherwise specified.

Cell Culture, Protein Extraction, and Lysarginase Digestion.

HEK293T cells from the RIKEN BRC cell bank (Ibaraki, Japan) were cultured in Dulbecco's modified Eagle's medium with 10% fetal bovine serum at 37 °C, 5% CO_2. Proteins were extracted with a phase-transfer surfactant (PTS protocol), as described previously (20), except that LysargiNase was used instead of trypsin/LysC. Briefly, HEK293T cells were suspended in PTS lysis buffer consisting of protease inhibitors, 12 mM sodium

deoxycholate (SDC), 12 mM sodium *N*-lauroylsarcosinate (SLS) in 100 mM Tris–HCl buffer (pH 9). The cell lysate was diluted tenfold with 10 mM $CaCl_2$ solution and incubated overnight with LysargiNase [enzyme/substrate ratio = 1:50 or 1:100 (w/w)] at 37 °C. After digestion, an equal volume of ethyl acetate was added, and the solution was acidified with trifluoroacetic acid (TFA). Then, the peptides were desalted with SDB-StageTips (21, 22).

Strong Cation Exchange Chromatography.

SCX chromatography was performed using disposable pipette tip-based SCX-StageTips (22), prepared as follows: double SCX membrane from Empore cation-SR disks was stamped out using a blunt-end 16-gauge syringe needle and placed into a P200 pipette tip. Mobile phases were prepared with aqueous 30% acetonitrile in buffers with various acid or salt solutions (Table S1). Prior to use, SCX-StageTips were activated with aqueous 30% acetonitrile in 1 M NaCl, and were equilibrated with 300 μL of the mobile phase. Ten micrograms of the desalted peptide samples were dried in a SpeedVac concentrator (Thermo Fisher Scientific), reconstituted with the mobile phase, and loaded onto each SCX-StageTip. The eluted fractions were collected. After SpeedVac concentration, each sample was reconstituted in 4% acetonitrile with 0.5% TFA and injected onto a nanoLC/MS/MS system equipped with an Orbitrap Fusion Lumos mass spectrometer (Thermo Fisher Scientific) or Orbitrap Exploris 480 mass spectrometer (Thermo Fisher Scientific), as described below. To compare HPLC-based and tip-based CHAMP-N, peptide isolation experiments were performed in triplicate using 80 μg x 3 of LysargiNase-digested HEK293T peptides. A quarter of the isolated peptides were injected onto nanoLC/MS/MS using Orbitrap Fusion Lumos and measurements were performed in triplicate.

Isolation of C-terminal and N, C-terminal peptides by CHAMP-C and CHAMP-NC.

C-terminal peptide isolation by CHAMP-C (14) and N, C-terminal peptide isolation by CHAMP-NC (15, 19) were performed as described previously. In brief, C-terminal peptide isolation was performed using CeO_2 tips, which was loaded with methanol-suspended CeO_2 particles and centrifuged on a P200 pipette tip with a polyethylene frit. Ten micrograms of the V8 protease-digested peptide samples were reconstituted with 40% acetonitrile containing 50 mM 5-Aminovaleric acid-HCl and 1M NaCl, and loaded onto each CeO_2 tip. N, C-terminal peptide isolation was performed using SCX-StageTips. Ten micrograms of the trypsin/Lys-C-digested peptide samples were reconstituted with 30% acetonitrile containing 0.125 % TFA, and loaded onto each SCX-StageTip. In both methods, the eluted fractions were collected.

NanoLC/MS/MS Analysis.

The LC/MS/MS analyses were performed on an HTC-PAL autosampler (CTC Analytics, Zwingen, Switzerland) and an Ultimate 3000 pump (Thermo Fisher Scientific), combined with an Orbitrap Fusion Lumos mass spectrometer or an Ultimate 3000 liquid chromatograph combined with an Orbitrap Exploris 480 mass spectrometer (Thermo Fisher Scientific). The LC mobile phases consisted of solvent A (0.5 % acetic acid) and solvent B (0.5 % acetic acid and 80 % acetonitrile).

For the Orbitrap Fusion Lumos system, the peptides were separated on self-pulled needle columns (150 mm, 100 μm ID) packed with Reprosil-Pur 120 C18-AQ 3 μm (Dr. Maisch, Ammerbuch, Germany) (23), and separated by a linear gradient set as follows: 5-10 % B in 5 min, 10-40 % B in 60 min, and 40-99 % B in 5 min, followed by 99 % B for 10 min. The flow rate was 500 nL/min. The electrospray voltage was set to 2.4 kV in the positive mode. The mass range of the survey scan was from 300 to 1,500 m/z with resolution of 120,000, standard automatic gain control (AGC) target, and maximum injection time of 50 ms. The first mass of MS/MS scan was set to 110 m/z with a resolution of 15,000, 100% normalized AGC target, and maximum injection time of 50 ms. The fragmentation was performed by higher energy collisional dissociation with a normalized collision energy of 30 %. The dynamic exclusion time was set to 20 s.

For the Orbitrap Exploris 480 system, the peptides were separated on self-pulled needle columns (250 mm, 100 μm ID) packed with Reprosil-Pur 120 C18-AQ 1.9 μm (Dr. Maisch, Ammerbuch, Germany) at 50 °C in a column oven (Sonation), and separated by a linear gradient set as follows: 5 % B in 8.3 min, 5-19 % B in 92.2 min, 19-29 % B in 34.5 min, 29-40 % B in 15 min, and 40-99 % B in 0.1 min, followed by 99 % B for 4.9 min. The flow rate was 400 nL/min. The electrospray voltage was set to 2.4 kV in the positive mode. The mass spectrometric analysis was carried out with the FAIMS Pro interface. The FAIMS mode was set to a standard resolution, and the total carrier gas flow was 4.0 L/min. The compensation voltage (CV) was set to –40, –60, and – 80, and the cycle time of each CV experiment was set to 1 s. The mass range of the survey scan was from 300 to 1,500 m/z with resolution of 60,000, 300% normalized AGC target, and auto maximum injection time. The first mass of MS/MS scan was set to 120 m/z with a resolution of 30,000, standard AGC target, and auto maximum injection time. The fragmentation was performed by higher energy collisional dissociation with a normalized collision energy of 30 %. The dynamic exclusion time was set to 20 s.

Data Analysis.
The raw MS data files acquired with FAIMS were converted into an MzXML format using FAIMS MzXML Generator (https://github.com/coongroup/FAIMS-MzXML-Generator) (24). The obtained mzxml files as well as the raw MS data files acquired using the Orbitrap Fusion

Lumos mass spectrometer were searched by MaxQuant (ver. 1.6.17.0) (25), and the database search was performed with Andromeda (26) against the SwissProt database (ver. 2020-10, 42,372 human protein entries). For most of the CHAMP-N experiments, search parameters included two missed cleavage sites for LysargiNase-digested peptides, while LysargiNase-digestion with N-terminal free semi-specificity was set for the searching of Neo-N-termini by CHAMP-N. Two missed cleavage sites for trypsin-digested peptides were set for the CHAMP-NC experiments, and two missed cleavage sites for V8 protease-digested peptides were set for the CHAMP-C experiments. Protein N-terminal acetylation and methionine oxidation were set as variable modifications, and cysteine carbamidomethylation was set as a fixed modification. For the searching of Neo-N-termini by CHAMP-N, peptide N-terminal acetylation was set instead of protein N-terminal acetylation. The peptide mass tolerance was 4.5 ppm, and the MS/MS tolerance was 20 ppm. The false discovery rate was set to 0.01 at the peptide-spectrum match level and protein level.

Experimental Design and Statistical Rationale

All experiments were performed using aliquots of the same HEK293T digest to minimize confounders by steps prior to isolation of the terminal peptides. Isolation of terminal peptides was prepared with n = 3, and in most cases a single LC/MS/MS measurement was performed, whereas for comparison with the HPLC-based method LC/MS/MS measurements were performed in triplicate for each sample. No statistical analysis was performed in this study.

RESULTS AND DISCUSSION

Optimization of Isolation Conditions for N-terminal Peptides Using Pipette-tip SCX Columns.

This study aimed to establish the tip-based CHAMP-N method, in which N-terminal peptides are isolated with high selectivity using a pipette-tip microcolumn (Figure S1A) instead of the conventional HPLC-based method (13). In this method, proteins are initially digested with LysargiNase, a digestive enzyme that cleaves the N-terminal side of Lys and Arg. Peptides digested with LysargiNase have 0 charge on the acetylated N-terminal peptides and +1 charge on the unmodified N-terminal peptides, whereas the non-N-terminal peptides have at least +2 charge under acidic conditions. Based on the charge/orientation retention model (17), non-N-terminal peptides with two positive charges in close proximity are strongly retained in the SCX. Thus, due to the relatively weak retention, acetylated N-terminal peptides with 0 charge, as well as unmodified N-terminal peptides with +1 charge, and N-terminal peptides with +2 charge (e.g., His-containing peptides) with charges at

distant positions can be simultaneously isolated (Figure S1B). We have previously reported the CHAMP-N method on a pipette-tip microcolumn (15, 19). However, we found that N-terminal peptides with +2 charge could not be isolated under the isolation conditions with formic acid (FA) used in our previous report (Figure S2A, B). Even when the isocratic elution volume was increased, N-terminal peptides with +2 charge were hardly isolated (Figure S2C, D). In this study, we first investigated the eluent conditions to elute N-terminal peptides with high selectivity and without loss of +2 charge N-terminal peptides using HEK293T cell-derived proteins.

Salt- and pH-based elution are two major modes for peptide separation by SCX chromatography (27, 28). For isolating protein N-terminal peptides by SCX-HPLC (13), we employed salt-based elution under acidic conditions. Previously, Adachi *et al.* reported the superiority of strong acid-based elution compared to salt-based elution in an SCX-StageTip-based fractionation (29). Therefore, we compared salt-based elution with acid-based elution. For the salt-based elution, we used the same eluents as those used in the HPLC-based method (13). For the acid-based elution, an eluent containing 0.2 % TFA, which is a stronger acid than FA used in the above experiment, was used. Isocratic elution was conducted for collecting 100 µL of the eluent for each fraction, and six fractions were collected and measured by LC/MS/MS. As a result, the peptides eluted in the order of the charge/orientation retention model under both conditions, and protein N-terminal peptides were successfully isolated (Figure 1A). Although the elution profiles were similar under both conditions, more N-terminal peptides were identified by acid-based elution than by salt-based elution (Figure 1B, C). This result is consistent with the previous report in which acid-based elution showed higher separation efficiency than salt-based elution in peptide fractionation (29).

When the TFA concentration was increased from 0.2 % to 0.5 %, N-terminal peptides were isolated with smaller elution volume. After the fraction 3, the selectivity decreased sharply (Figure 2A, B), and 96.4 % of all N-terminal peptides identified, or 1794 N-terminal peptides, were identified from the fractions 1 or 2 (Figure 2C). These results indicate that N-terminal peptides can be isolated selectively and without loss of N-terminal peptides by collecting fractions 1 and 2 (100 µL in total) under the 0.5 % TFA condition.

Based on the N-terminal peptide isolation conditions by salt- and acid-based elution described above, we compared the identification number and selectivity of N-terminal peptides in a single LC/MS/MS run without pre-fractionation. As a result, more N-terminal peptides were identified by acid-based elution than by salt-based elution, as was the case with fractionation (Figure S3A). N-terminal peptides could be isolated with a high selectivity of more than 95 % under all conditions (Figure S3A, B). The optimal conditions were defined as those of acid-based elution with 0.5 % TFA, which allowed identification of a larger

number of N-terminal peptides with smaller elution volume. Next, we compared the conditions with 0.5 % TFA with those with FA previously reported (15, 19). Compared to the conventional method with 2.5 % FA, using this method with 0.5 % TFA, more N-terminal peptides could be identified (Figure S3C). In addition, N-terminal peptides with +2 charge could be isolated by this method (Figure S3D), whereas they could hardly be isolated by our previous method. In subsequent experiments, acid-based elution with 0.5% TFA conditions were used.

Comparison of Conventional HPLC-based Method and Novel Tip-based Method.

Next, we compared the performance of the new tip-based method with that of the previously reported HPLC-based method (13). The results of the HPLC-based method for comparison were obtained by reanalyzing the previously published data (PXD010551) using the same database search parameters as the tip-based method. The HPLC-based method identified 1875 N-terminal peptides on average, whereas the tip-based method successfully identified 1985 N-terminal peptides on average in a single LC/MS/MS run (Figure 3A). The tip-based method showed relatively lower variability among replicates, which may be attributed to the fact that, unlike the HPLC-based method, multiple tips can be used for parallel processing. On the other hand, N-terminal peptides with two positive charges were detected at a higher rate by the HPLC-based method (Figure 3B). These results suggest that it is difficult to separate non-N-terminal peptides with two positive charges close to each other from N-terminal peptides with two positive charges far from each other, and that the HPLC column is superior to the tip-based method in separation. Thus, the tip-based method is inferior to the HPLC-based method in a few aspects. However, the tip-based method was able to isolate N-terminal peptides with a high selectivity of more than 92%, and considering the advantages of easy operation and high throughput, we have successfully established a practical and simple protocol.

Identification of Proteolytic Cleavage Sites and Translation Initiation Sites.

We applied this tip-based CHAMP-N method to the detection of neo-N-termini derived from proteolysis or non-canonical translation initiation. In "semi-specific search free N-terminus" for the MaxQuant search, peptides whose N-termini do not match the specificity of the enzyme, neo-N-terminal peptides, can be additionally identified. This means that it is possible to detect neo-N-termini caused by downstream shifts of the translation initiation site or endogenous proteolysis, which differ from the native N-terminal sequence in the database.

As a result, over 900 neo-N-terminal peptides were identified in at least one replicate (Figure S4A). Among them, 108 of the neo-N-termini were acetylated. Since most acetylation of protein N-termini occurs co-translationally (30), these neo-N-termini are suggested to be

non-canonical translation initiation sites. Actually, 85 of 108 were acetylated at Met or the next residue of Met (Figure 4A). The removal of the initiator methionine at the protein N-terminus and acetylation is dependent on the next residue of the Met (30). Interestingly, the frequency of the next residue of Met in 85 acetylated neo-N-termini was similar to that in the native N-termini (Figure 4B). Non-canonical translation initiation sites are known not only from unannotated methionine (AUG) but also from AUG-like codons called near-cognate codons (31–33). Among the acetylated neo-N-termini identified in this study, near-cognate initiation codons, especially CUG-initiated translation initiation site candidates, were also identified in two proteins (Figure S4B). These two candidate translation initiation sites were not identified by a recently reported approach for accurately identifying translation initiation sites, including initiation at near-cognate codons (34). This approach uses the results of three N-terminomics methods on HEK293T samples, and our method identified those that were not found there.

Another feature of the neo-N-termini was found to be a large number of neo-N-terminal peptides derived from the N-terminal region of the total length of the protein (Figure S4C). We also found that proteins with neo-N-termini were the most enriched with signal and transit peptides (Figure 4C). These results suggest that the neo-N-termini generated by the cleavage of signal or transit peptides can be detected. In fact, when we checked whether the lengths of the signal peptide and transit peptide sequences annotated in UniProt matched the position of the neo-N terminus identified in this study, 24 neo-N termini matched the signal peptide and 19 neo-N termini matched the transit peptide (Figure 4D). For example, COX6A1 protein localized to mitochondria has a transit peptide at residues 1 through 24, and a neo-N-terminal peptide from residue 25 was identified (Figure S5A).

Among the neo-N-termini that did not match the cleavage position annotated in UniProt, there were several neo-N-termini that deviated a few residues from the cleavage position. Although annotated in UniProt, some of the neo-N-termini have no experimental evidence, and it is possible that the neo-N-termini identified in this study are the true cleavage positions. Therefore, we used TargetP 2.0 (35) as a transit peptide prediction model and SignalP 6.0 (36) as a signal peptide prediction model to search for candidate true cleavage positions. Although UniProt considers results from multiple prediction tools in the absence of experimental evidence, it is unclear whether the latest versions of SignalP and TargetP are used. The latest version, SignalP 6.0, is a signal peptide prediction model using protein language model and has better prediction performance than SignalP 5.0 based on deep learning. Based on these prediction models and experimental data from this study, we attempted to identify the true cleavage site candidates.

As a result, 7 neo-N-termini matched the predicted cleavage sites by SignalP 6.0 (Table 1), and 11 neo-N-termini matched the predicted cleavage sites by TargetP 2.0 (Table 2).

Most of these 18 proteins have annotated signal or transit peptide sequences on UniProt, but there is no experimental evidence for all but two. Therefore, for these proteins, the neo-N-terminal positions identified in this study are true candidates for cleavage positions. For example, FKBP9 protein has a signal peptide from residue 1 to 24 annotated on UniProt, but according to the SignalP 6.0 prediction, residues 1 to 29 are the signal peptide, and a neo-N-terminal peptide from residue 30 was actually identified (Figure S5B). Of the 18 neo-N termini matched in this study, 12 are also registered in TopFIND (37), a database of protein termini, strengthening the possibility that these termini are the true cleavage positions. It is expected that more experimental information on protein termini will be obtained by this method or existing terminomics methods in the future, leading to further improvement of the prediction model accuracy. Synergistic effects are also expected, such as experimental validation using the improved prediction model.

As mentioned above, most of the neo-N-termini identified in this study are derived from the N-terminal region of the total protein length, but the majority of the neo-N-termini are located less than 10 residues from the native N-terminus (Figure S4D). Typical signal and transit peptides are 10 to 50 residues in length, but shorter peptides are cleaved. In particular, the neo-N-terminus, starting at residue 3, was the most abundant, identified in 70 proteins. In 12 of them, the N-terminus was acetylated, suggesting that it may be excessively cleaved by methionine aminopeptidase, which removes the initiator methionine during translation. We also found that some proteins had more than 10 neo-N-terminal peptides in one protein (Figure S4E). Many of these neo-N-terminal peptides were peptide ladders with one amino acid residue shaved off each (Figure S4F), suggesting cleavage by exopeptidase. These findings suggest that excessive cleavage by methionine aminopeptidase and cleavage by exopeptidase may occur frequently in the cell.

Comprehensive Protein N- and C-terminome Analysis Using Three CHAMP Methods.

We recently reported the CHAMP-C method (14), which is a tip-based method for isolation of C-terminal peptides and the CHAMP-NC method (15, 19), which is a tip-based version of a previously known method for simultaneous isolation of N- and C-terminal peptides (Figure 5A). The CHAMP-C method consists only of protein digestion by V8 protease followed by metal oxide-based ligand-exchange (MOLEX) chromatography. When digested by V8 protease, which cleaves on the C-terminal side of Asp and Glu, the non-C-terminal peptides have two carboxy groups at their C-termini, whereas the C-terminal peptides have only one carboxy group. Therefore, using MOLEX, the dicarboxylates of the non-C-terminal peptide forms a stable chelate with metal atoms, allowing the isolation of the weaker retained C-terminal peptides. On the other hand, the CHAMP-NC method consists only of tryptic digestion followed by SCX separation. When digested by trypsin, which

cleaves at the C-terminal side of Lys and Arg, the non-terminal peptides have more than +2 charge while the acetylated N-terminal and C-terminal peptides have +1 charge under acidic conditions. Therefore, using SCX, the acetylated N- and C-terminal peptides, which are weakly retained, can be isolated because the non-terminal peptides with two positive charges are strongly retained by SCX. Note that, unlike the CHAMP-N method, unmodified N-terminal peptides with +2 charge are not isolated by this method.

We performed a comprehensive protein terminome analysis of HEK293T cell-derived proteins using these three CHAMP methods (Figure 5B). In each CHAMP method, HEK293T cell-derived proteins were analyzed in triplicate, and 10 µg of protein sample was used for terminal peptide isolation in each replicate. In total, 2254 protein C-termini and 1901 protein N-termini were identified (Figure 5C). Since each CHAMP method is based on a different digestion enzyme and isolation mechanism, the overlapping rate in identified protein termini between methods is low, and the combination of the three CHAMP methods provides a more comprehensive analysis than using each alone (Figure 5C). As for protein C-termini, C-terminal peptides identified by CHAMP-NC method tend to have acidic amino acid residues near the peptide C-termini, and such C-terminal peptides are not easily identified by CHAMP-C method because they are relatively strongly retained by MOLEX (Figure S6A). On the other hand, C-terminal peptides with Lys and Arg near the peptide C-termini are too short to be identified by the CHAMP-NC method using trypsin digestion (Figure S6B). As for protein N-termini, many N-terminal peptides identified only by the CHAMP-N method are not acetylated or have extra basic amino acids, i.e., extra positive charge (Fig. S6C-E). Trypsin-digested N-terminal peptides always have Lys or Arg, while LysargiNase-digested N-terminal peptides have less positive charge derived from Lys and Arg. Therefore, with the CHAMP-N method, it is possible to isolate peptides with extra positive charge, which makes the retention in SCX stronger. For LysargiNase-digested N-terminal peptides identified by the CHAMP-N method, the net charge (Z) under acidic conditions correlated with the charge state of the precursor ion, the number of b- and y-ions after fragmentation, and the score in database searches by MaxQuant. This suggests that N-terminal peptides with Z = 0 are difficult to identify. On the other hand, trypsin-digested N-terminal peptides identified by the CHAMP-NC method are always Z ≥ 1. Therefore, combining terminomics methods based on different digestion enzymes and isolation mechanisms can improve coverage and allow for more comprehensive identification of protein termini by complementing each other.

Conclusions

We have succeeded in developing a tip-based CHAMP-N method that can be parallelized while maintaining the performance (identification number and selectivity) of the HPLC method. We have applied this method to identify novel post-translational cleavages (signal

peptides and transit peptides) and non-canonical translation initiation sites, including initiation from near-cognate codons. In combination with the other two tip-based CHAMP methods, one-step terminomics can now be performed very easily. The CHAMP method is expected to be widely used as a standardized method for terminomics in the future.

Data Availability.

The raw MS data and analysis files have been deposited with the ProteomeXchange Consortium (http://proteomecentral.proteomexchange.org) via the jPOST partner repository (https://jpostdb.org/) with the data set identifier PXD048917.

Supplemental data – This article contains supplemental data.

ACKNOWLEDGEMENTS

We would like to thank the members of the Department of Molecular Systems BioAnalysis for fruitful discussion.

Funding and additional information

This work has been funded by the JST Strategic Basic Research Program CREST (No. JPMJCR1862), AMED-CREST program (No.JP18gm1010010), and JSPS Grants-in-Aid for Scientific Research 21H02459, 23H04924 to Y.I. and 20H03241 to K. I., and JST FOREST (No. JPMJFR214L) to K.I.

Author contributions—K.M., H.N., K.I., and Y. I. designed research; K.M., and H.N. performed research; K.M. analyzed data; K.M., H.N., K.I., and Y. I. wrote the paper.

Conflict of interest—The authors declare no competing interests.

Abbreviations—The abbreviations used are: SCX, strong cation exchange; CHAMP-N, chromatographic amplification of protein N-terminal peptides; ACN, acetonitrile; COFRADIC, combined fractional diagonal chromatography; TAILS, terminal amine isotopic labeling of substrates; CHAMP-C, chromatographic amplification of protein C-terminal peptides; PTS, phase-transfer surfactants; SDC, sodium deoxycholate; SLS, sodium *N*-lauroylsarcosinate; Tris–HCl; tris(hydroxymethyl)amino- methane hydrochloride, TFA, trifluoroacetic acid; AGC, automatic gain control; CV, compensation voltage; FA, formic acid; iMet, initiator methionine; CHAMP-NC, chromatographic amplification of protein N- and C-terminal peptides; MOLEX, metal oxide-based ligand-exchange; Nt-peptide, N-terminal peptide.

REFERENCES

1. Smith, L. M., Thomas, P. M., Shortreed, M. R., Schaffer, L. V., Fellers, R. T., LeDuc, R. D., Tucholski, T., Ge, Y., Agar, J. N., Anderson, L. C., Chamot-Rooke, J., Gault, J., Loo, J. A., Paša-Tolić, L., Robinson, C. V., Schlüter, H., Tsybin, Y. O., Vilaseca, M., Vizcaíno, J. A., Danis, P. O., and Kelleher, N. L. (2019) A five-level classification system for proteoform identifications. *Nat. Methods* 16, 939–940

2. Aebersold, R., Agar, J. N., Amster, I. J., Baker, M. S., Bertozzi, C. R., Boja, E. S., Costello, C. E., Cravatt, B. F., Fenselau, C., Garcia, B. A., Ge, Y., Gunawardena, J., Hendrickson, R. C., Hergenrother, P. J., Huber, C. G., Ivanov, A. R., Jensen, O. N., Jewett, M. C., Kelleher, N. L., Kiessling, L. L., Krogan, N. J., Larsen, M. R., Loo, J. A., Ogorzalek Loo, R. R., Lundberg, E., MacCoss, M. J., Mallick, P., Mootha, V. K., Mrksich, M., Muir, T. W., Patrie, S. M., Pesavento, J. J., Pitteri, S. J., Rodriguez, H., Saghatelian, A., Sandoval, W., Schlüter, H., Sechi, S., Slavoff, S. A., Smith, L. M., Snyder, M. P., Thomas, P. M., Uhlén, M., Van Eyk, J. E., Vidal, M., Walt, D. R., White, F. M., Williams, E. R., Wohlschlager, T., Wysocki, V. H., Yates, N. A., Young, N. L., and Zhang, B. (2018) How many human proteoforms are there? *Nat. Chem. Biol.* 14, 206–214

3. Smith, L. M., Agar, J. N., Chamot-Rooke, J., Danis, P. O., Ge, Y., Loo, J. A., Paša-Tolić, L., Tsybin, Y. O., Kelleher, N. L., and Consortium for Top-Down Proteomics (2021) The Human Proteoform Project: Defining the human proteome. *Sci Adv* 7, eabk0734

4. Schlüter, H., Apweiler, R., Holzhütter, H.-G., and Jungblut, P. R. (2009) Finding one's way in proteomics: a protein species nomenclature. *Chem. Cent. J.* 3, 11

5. Sato, K., Masuda, T., Hu, Q., Tobo, T., Gillaspie, S., Niida, A., Thornton, M., Kuroda, Y., Eguchi, H., Nakagawa, T., Asano, K., and Mimori, K. (2019) Novel oncogene 5MP1 reprograms c-Myc translation initiation to drive malignant phenotypes in colorectal cancer. *EBioMedicine* 44, 387–402

6. Hann, S. R., Dixit, M., Sears, R. C., and Sealy, L. (1994) The alternatively initiated c-Myc proteins differentially regulate transcription through a noncanonical DNA-binding site. *Genes Dev.* 8, 2441–2452

7. Hann, S. R., King, M. W., Bentley, D. L., Anderson, C. W., and Eisenman, R. N. (1988) A non-AUG translational initiation in c-myc exon 1 generates an N-terminally distinct protein whose synthesis is disrupted in Burkitt's lymphomas. *Cell* 52, 185–195

8. Chow, V. W., Mattson, M. P., Wong, P. C., and Gleichmann, M. (2010) An overview of APP processing enzymes and products. *Neuromolecular Med.* 12, 1–12

9. Varshavsky, A. (2019) N-degron and C-degron pathways of protein degradation. *Proc. Natl. Acad. Sci. U. S. A.* 116, 358–366

10. Gevaert, K., Goethals, M., Martens, L., Van Damme, J., Staes, A., Thomas, G. R., and Vandekerckhove, J. (2003) Exploring proteomes and analyzing protein processing by mass spectrometric identification of sorted N-terminal peptides. *Nat. Biotechnol.* 21, 566–569

11. Kleifeld, O., Doucet, A., auf dem Keller, U., Prudova, A., Schilling, O., Kainthan, R. K., Starr, A. E., Foster, L. J., Kizhakkedathu, J. N., and Overall, C. M. (2010) Isotopic labeling of terminal amines in complex samples identifies protein N-termini and protease cleavage products. *Nat. Biotechnol.* 28, 281–288

12. Hanna, R., Rozenberg, A., Saied, L., Ben-Yosef, D., Lavy, T., and Kleifeld, O. (2023) In-

Depth Characterization of Apoptosis N-Terminome Reveals a Link Between Caspase-3 Cleavage and Posttranslational N-Terminal Acetylation. *Mol. Cell. Proteomics* 22,

13. Chang, C.-H., Chang, H.-Y., Rappsilber, J., and Ishihama, Y. (2021) Isolation of Acetylated and Unmodified Protein N-Terminal Peptides by Strong Cation Exchange Chromatographic Separation of TrypN-Digested Peptides. *Mol. Cell. Proteomics* 20, 100003

14. Nishida, H., and Ishihama, Y. (2022) One-Step Isolation of Protein C-Terminal Peptides from V8 Protease-Digested Proteins by Metal Oxide-Based Ligand-Exchange Chromatography. *Anal. Chem.* 94, 944–951

15. Tsumagari, K., Chang, C.-H., and Ishihama, Y. (2021) Exploring the landscape of ectodomain shedding by quantitative protein terminomics. *iScience* 24, 102259

16. Gauci, S., Helbig, A. O., Slijper, M., Krijgsveld, J., Heck, A. J. R., and Mohammed, S. (2009) Lys-N and trypsin cover complementary parts of the phosphoproteome in a refined SCX-based approach. *Anal. Chem.* 81, 4493–4501

17. Alpert, A. J., Petritis, K., Kangas, L., Smith, R. D., Mechtler, K., Mitulović, G., Mohammed, S., and Heck, A. J. R. (2010) Peptide orientation affects selectivity in ion-exchange chromatography. *Anal. Chem.* 82, 5253–5259

18. Helbig, A. O., Gauci, S., Raijmakers, R., van Breukelen, B., Slijper, M., Mohammed, S., and Heck, A. J. R. (2010) Profiling of N-acetylated protein termini provides in-depth insights into the N-terminal nature of the proteome. *Mol. Cell. Proteomics* 9, 928–939

19. Tsumagari, K., Chang, C.-H., and Ishihama, Y. (2021) A protocol for analyzing the protein terminome of human cancer cell line culture supernatants. *STAR Protoc* 2, 100682

20. Masuda, T., Tomita, M., and Ishihama, Y. (2008) Phase transfer surfactant-aided trypsin digestion for membrane proteome analysis. *J. Proteome Res.* 7, 731–740

21. Rappsilber, J., Ishihama, Y., and Mann, M. (2003) Stop and go extraction tips for matrix-assisted laser desorption/ionization, nanoelectrospray, and LC/MS sample pretreatment in proteomics. *Anal. Chem.* 75, 663–670

22. Rappsilber, J., Mann, M., and Ishihama, Y. (2007) Protocol for micro-purification, enrichment, pre-fractionation and storage of peptides for proteomics using StageTips. *Nat. Protoc.* 2, 1896–1906

23. Ishihama, Y., Rappsilber, J., Andersen, J. S., and Mann, M. (2002) Microcolumns with self-assembled particle frits for proteomics. *J. Chromatogr. A* 979, 233–239

24. Hebert, A. S., Prasad, S., Belford, M. W., Bailey, D. J., McAlister, G. C., Abbatiello, S. E., Huguet, R., Wouters, E. R., Dunyach, J.-J., Brademan, D. R., Westphall, M. S., and Coon, J. J. (2018) Comprehensive Single-Shot Proteomics with FAIMS on a Hybrid Orbitrap Mass Spectrometer. *Anal. Chem.* 90, 9529–9537

25. Cox, J., and Mann, M. (2008) MaxQuant enables high peptide identification rates, individualized p.p.b.-range mass accuracies and proteome-wide protein quantification. *Nat. Biotechnol.* 26, 1367–1372

26. Cox, J., Neuhauser, N., Michalski, A., Scheltema, R. A., Olsen, J. V., and Mann, M.

(2011) Andromeda: a peptide search engine integrated into the MaxQuant environment. *J. Proteome Res.* 10, 1794–1805

27. Marino, F., Cristobal, A., Binai, N. A., Bache, N., Heck, A. J. R., and Mohammed, S. (2014) Characterization and usage of the EASY-spray technology as part of an online 2D SCX-RP ultra-high pressure system. *Analyst* 139, 6520–6528

28. Xu, J., Gao, J., Yu, C., He, H., Yang, Y., Figeys, D., and Zhou, H. (2016) Development of Online pH Gradient-Eluted Strong Cation Exchange Nanoelectrospray-Tandem Mass Spectrometry for Proteomic Analysis Facilitating Basic and Histidine-Containing Peptides Identification. *Anal. Chem.* 88, 583–591

29. Adachi, J., Hashiguchi, K., Nagano, M., Sato, M., Sato, A., Fukamizu, K., Ishihama, Y., and Tomonaga, T. (2016) Improved Proteome and Phosphoproteome Analysis on a Cation Exchanger by a Combined Acid and Salt Gradient. *Anal. Chem.* 88, 7899–7903

30. Aksnes, H., Ree, R., and Arnesen, T. (2019) Co-translational, Post-translational, and Non-catalytic Roles of N-Terminal Acetyltransferases. *Mol. Cell* 73, 1097–1114

31. Ingolia, N. T., Lareau, L. F., and Weissman, J. S. (2011) Ribosome profiling of mouse embryonic stem cells reveals the complexity and dynamics of mammalian proteomes. *Cell* 147, 789–802

32. Lee, S., Liu, B., Lee, S., Huang, S.-X., Shen, B., and Qian, S.-B. (2012) Global mapping of translation initiation sites in mammalian cells at single-nucleotide resolution. *Proc. Natl. Acad. Sci. U. S. A.* 109, E2424–32

33. Chen, J., Brunner, A.-D., Cogan, J. Z., Nuñez, J. K., Fields, A. P., Adamson, B., Itzhak, D. N., Li, J. Y., Mann, M., Leonetti, M. D., and Weissman, J. S. (2020) Pervasive functional translation of noncanonical human open reading frames. *Science* 367, 1140–1146

34. Ichihara, K., Matsumoto, A., Nishida, H., Kito, Y., Shimizu, H., Shichino, Y., Iwasaki, S., Imami, K., Ishihama, Y., and Nakayama, K. I. (2021) Combinatorial analysis of translation dynamics reveals eIF2 dependence of translation initiation at near-cognate codons. *Nucleic Acids Res.* 49, 7298–7317

35. Almagro Armenteros, J. J., Salvatore, M., Emanuelsson, O., Winther, O., von Heijne, G., Elofsson, A., and Nielsen, H. (2019) Detecting sequence signals in targeting peptides using deep learning. *Life Sci Alliance* 2,

36. Teufel, F., Almagro Armenteros, J. J., Johansen, A. R., Gíslason, M. H., Pihl, S. I., Tsirigos, K. D., Winther, O., Brunak, S., von Heijne, G., and Nielsen, H. (2022) SignalP 6.0 predicts all five types of signal peptides using protein language models. *Nat. Biotechnol.* 40, 1023–1025

37. Fortelny, N., Yang, S., Pavlidis, P., Lange, P. F., and Overall, C. M. (2015) Proteome TopFIND 3.0 with TopFINDer and PathFINDer: database and analysis tools for the association of protein termini to pre- and post-translational events. *Nucleic Acids Res.* 43, D290–7

Table 1. Proteins with signal peptide cleavage sites consistent with SignalP 6.0 predictions

UniProt accession number	Gene name	SignalP 6.0	Neo-Nt position in this study	Experimental evidence in UniProt	UniProt signal peptide	TopFIND 4.1
O00115	DNASE2	1-16	17	No	1-18	-
P13667	PDIA4	1-24	25	No	1-20	+
P04843	RPN1	1-24	25	No	1-23	+
Q14257	RCN2	1-25	26	No	1-22	+
Q9Y3Q3	TMED3	1-27	28	No	1-23	+
O95302	FKBP9	1-29	30	No	1-24	-
P11047	LAMC1	1-35	36	No	1-33	-

Neo-N-termini matched predicted cleavage sites by SignalP 6.0. UniProt information is current as of October 2023.

Table 2. Proteins with transit peptide cleavage sites consistent with TargetP 2.0 predictions

UniProt accession number	Gene name	TargetP 2.0	Neo-Nt position in this study	Experimental evidence in UniProt (2023.08)	UniProt transit peptide (2023.08)	TopFIND 4.1
P82673	MRPS35	1-22	23	No	1-?	+
Q8TD30	GPT2	1-24	25	No	-	+
Q9Y2Q9	MRPS28	1-28	29	No	1-71	+
P49419	ALDH7A1	1-29	30	No	1-26	-
Q9BX68	HINT2	1-30	31	No	1-17	+
P49590	HARS2	1-34	35	No	1-33	+
P54886	ALDH18A1	1-43	44	No	-	+
Q7Z6M4	MTERF4	1-43	44	Yes	1-42	-
P06576	ATP5F1B	1-46	47	Yes	1-47	+
P82650	MRPS22	1-53	54	No	-	+
Q9H2K0	MTIF3	1-55	56	No	1-31	+

Neo-N-termini matched predicted cleavage sites by TargetP 2.0 UniProt information is current as of October 2023.

Figure 1

Figure 1. Comparison of salt-based and acid-based isocratic elution of protein N-terminal peptides. SCX tip-based separation of LysargiNase-digested HEK293T peptides under isocratic conditions with salt (10 mM KCl, 7.5 mM potassium phosphate buffer at pH 2.2 and 30% ACN) or acid (0.2% TFA and 30% ACN). Z is the charge number at acidic pH, which is based on the number of basic residues per peptide, such as unmodified N-terminus, Lys, Arg and His. Each fraction was eluted at 100 µL. The Orbitrap Fusion Lumos system was used. (A) Distributions of Z values of identified peptides. (B) Numbers of identified N-terminal peptides. N-terminal peptide selectivity (%) was calculated as the sum of the signal intensity of all identified N-terminal peptides divided by that of all identified peptides. (C) Numbers of identified peptides in any of the six fractions.

Figure 2

Figure 2. Optimization of the elution volume for SCX tip-based separation of LysargiNase-digested HEK293T peptides under acid-based isocratic condition (0.5% TFA and 30% ACN). Z is the charge number at acidic pH, which based on the number of basic residues per peptide, such as unmodified N-terminus, Lys, Arg and His. Each fraction was eluted at 50 µL. The Orbitrap Fusion Lumos system was used. (A) Numbers of identified protein N-terminal peptides. N-terminal peptide selectivity (%) was calculated as the sum of the signal intensity of all identified N-terminal peptides divided by that of all identified peptides. (B) Distributions of Z values of identified peptides. (C) Overlaps of identified N-terminal peptides in each fraction.

Figure 3. Comparison of HPLC-based and tip-based isolation of protein N-terminal peptides. SCX HPLC or SCX tip separation of LysargiNase-digested HEK293T peptides. Z is the charge number at acidic pH, which based on the number of basic residues per peptide, such as unmodified N-terminus, Lys, Arg and His. The HPLC is the results of a reanalysis of our previous report using Orbitrap Fusion Lumos (JPST000422). For each replicate, three LC/MS/MS runs were performed. The Orbitrap Fusion Lumos system was used. (A) Numbers of identified N-terminal peptides. N-terminal peptide selectivity (%) was calculated as the sum of the signal intensity of all identified N-terminal peptides divided by that of all identified peptides. (B) Distributions of Z values of identified peptides.

Figure 4

Figure 4. Application of tip-based CHAMP-N to identification of translation initiation sites and cleavage sites of transit peptides and signal peptides. Peptides identified in at least one of the three replicates were analyzed. The Orbitrap Exploris 480 system was used. (A) Numbers of identified Nt-acetylated Nt-peptides. The Nt-peptides are categorized into three classes. Ac-MX: The Nt-peptides acetylated at the first (initiator) methionine. M/Ac-X: The Nt-peptides acetylated at the next residue of methionine. Other: The Nt-peptides without methionine around the acetylated residue. (B) The amino acid frequency at the second residue next to iMet of acetylated protein N-termini based on the absence or presence of iMet. (C) Enrichment analysis using UniProt Keywords for the proteins with neo-Nt-peptides. (D) Numbers of transit peptide or signal peptide cleavage site matched to UniProt annotations or unmatched neo-Nt-peptides.

Figure 5

Figure 5. Comprehensive analysis of protein N- and C-termini using the three CHAMP methods. The Orbitrap Exploris 480 system was used. (A) Schematic illustration of protein N- and C-terminal peptide enrichment. (B) Numbers of identified N- and C-terminal peptides. N- or C-terminal peptide selectivity (%) was calculated as the sum of the signal intensity of all identified N- and C-terminal peptides divided by that of all identified peptides. (C) Overlaps of identified N- and C-terminal peptides in each CHAMP method. Peptides identified in at least one of the three replicates were analyzed.

Figure S1

Figure S1. Schematic illustration of protein N-terminal peptide enrichment. (A) Photograph of a SCX-StageTip column. (B) Proteins are digested with LysargiNase, and tip-based SCX chromatography is performed to trap the non-N-terminal peptides. The non-N-terminal peptides are retained in the SCX column, whereas the protein N-terminal peptides flow through the SCX column.

Figure S2

Figure S2. Insufficient separation of protein N-terminal peptides under formic acid-based isocratic condition. SCX tip-based separation of LysargiNase-digested HEK293T peptides under formic acid-based isocratic condition. Z is the charge number at acidic pH, which is based on the number of basic residues per peptide, such as unmodified N-terminus, Lys, Arg and His. The Orbitrap Fusion Lumos system was used. (A) Numbers of identified protein N-terminal peptides. N-terminal peptide selectivity (%) was calculated as the sum of the signal intensity of all identified N-terminal peptides divided by that of all identified peptides. (B) Distributions of Z values of identified peptides. (C) Numbers of identified protein N-terminal peptides with fractionation.(D) Distributions of Z values of identified peptides with fractionation.

Figure S3

Figure S3. Comparison of salt-based and acid-based isocratic elution of protein N-terminal peptides without fractionation. SCX tip-based separation of LysargiNase-digested HEK293T peptides under isocratic conditions with salt or acid. Z is the charge number at acidic pH, which based on the number of basic residues per peptide, such as unmodified N-terminus, Lys, Arg and His.(A) & (C) Numbers of identified N-terminal peptides. N-terminal peptide selectivity (%) was calculated as the sum of the signal intensity of all identified N-terminal peptides divided by that of all identified peptides. (B) & (D) Distributions of Z values of identified peptides.

Figure S4

Figure S4. Characteristics of the neo-Nt-peptides. Peptides identified in at least one of the three replicates were analyzed. The Orbitrap Exploris 480 system was used. (A) Numbers of identified peptides with semi-specific search free N-terminus. (B) translation initiation site candidates with near-cognate codons. (C) Neo-Nt starting position of identified neo-Nt-peptides. (D) Zoom into the 3-50 neo-Nt starting position. Transit peptide or signal peptide cleavage site matched peptides are shown in orange. (E) Numbers of neo-Nt-peptides per protein. (F) List of proteins with ≥10 neo-Nt-peptides. Peptide ladder sequences (peptides with three or more consecutive start positions) are shown in red.

Figure S5

Figure S5. MS/MS spectra of the peptides matched to transit peptide or signal peptide cleavage site . The Orbitrap Exploris 480 system was used. (A) MS/MS spectrum of the peptide matched to transit peptide cleavage site annotated in UniProt. (B) MS/MS spectrum of the peptide matched to signal peptide cleavage site predicted by SignalP 6.0.

Figure S6

Figure S6. Characteristics of the peptides identified by each of the three CHAMP methods. Peptides identified in at least one of the three replicates were analyzed. The Orbitrap Exploris 480 system was used. (A) The residue distance distribution of the nearest Asp or Glu residue to the identified peptide sequence (including residues next to the V8 protease digestion site) in CHAMP-NC and CHAMP-C experiment. (B) The residue distance distribution of the nearest Lys or Arg residue to the identified peptide sequence (including residues next to the trypsin digestion site) in CHAMP-NC and CHAMP-C experiment. (C), (D) & (E) Numbers of identified N-terminal peptides. (F) Distributions of precursor ion charge state of identified Nt-peptides by CHAMP-N. (F) Distributions of precursor ion charge state of identified peptides by CHAMP-N. (G) Relationship between charge number at acidic pH and b- and y-ion count of Nt-peptides by CHAMP-N. (H) Relationship between charge number at acidic pH and Andromeda score (MaxQuant) of Nt-peptides by CHAMP-N.