Changes in responses of auditory cortex neurons upon falling asleep and awakening

Amit Marmelshtein[1,2*], Barak Lavy[3*], Barak Hadad[2], Yuval Nir[1,2,3,4]

1. Sagol School of Neuroscience, Tel Aviv University; Tel Aviv 6997801, Israel
2. Department of Physiology & Pharmacology, School of Medicine, Tel Aviv University, Tel Aviv 6997801, Israel
3. Department of Biomedical Engineering, Faculty of Engineering, Tel Aviv University, Tel Aviv 6997801, Israel
4. The Sieratzki-Sagol Center for Sleep Medicine, Tel Aviv Sourasky Medical Center, Tel Aviv, Israel.

*These authors contributed equally

Corresponding author: Yuval Nir, ynir@tauex.tau.ac.il

Abstract

The neural processes accompanying the process of falling asleep are only partially understood. At the cortical level, previous studies have shed light on aspects of both spontaneous neural activity and sensory responses that change between wakefulness and sleep across species. For example, sleep-related changes in early auditory cortex include reduction in spontaneous firing rates, increases in the occurrence of post-onset silent (OFF) periods, and elevated population synchrony. However, it remains unknown whether changes occur abruptly around behavioral state transitions of sleep onset and awakening. Here, we recorded spontaneous and sound-evoked neuronal spiking activity in early auditory cortex along with polysomnography during thousands of episodes when rats fell asleep or woke up. We found that when falling asleep, stimulus-induced neuronal silent periods (OFF periods), characteristic of non-rapid eye movement (NREM) sleep, increased within few seconds around sleep onset. Moderate reductions in spontaneous firing rate also occurred within seconds. By contrast, an increase in neuronal population synchrony built up gradually over dozens of seconds until reaching maximal levels. Similar effects, but with opposite direction, were observed around awakenings when stimulus-induced silent periods were suppressed within few seconds. The results highlight stimulus-induced neuronal silence as a key indicator of transitions between vigilance states. More generally, these findings emphasize the added value of going beyond monitoring ongoing activity and perturbing the nervous system to reveal its state - an insight that could also help guide development of more sensitive non-invasive monitors of falling asleep in humans.

Keywords

A1, rats, NREM, OFF periods, click-trains

Introduction

Every night we drift into sleep, becoming disconnected as our conscious experience of the sensory environment largely subsides. This profound shift in our mental state raises fascinating questions regarding the ways by which brain states shape our behavior and subjective experiences. Previous work in rodents has investigated how different brain states affect sensory processing (Harris and Thiele, 2011; Lee and Dan, 2012; Mcginley et al., 2015). In the auditory domain, such research revealed how factors such as arousal, locomotion, sleep pressure, ongoing behavioral tasks, and attention influence auditory responses at the neuronal level (Atiani et al., 2009; Edeline et al., 2001; Harris and Thiele, 2011; Issa and Wang, 2008; Marmelshtein et al., 2023; Nir et al., 2013; Sela et al., 2020).

In a recent investigation, we examined the responses to sounds within the early auditory cortex of rats across vigilant wakefulness, sleep deprivation, NREM sleep, and REM sleep. Our findings revealed robust modulation in specific aspects of neuronal activity, particularly in population synchrony and post-onset responses (Marmelshtein et al., 2023). In contrast, other features such as spontaneous firing rates only showed modest modulation, while onset responses were largely unaffected by vigilance states (Marmelshtein et al., 2023). While these changes were examined over the scale of dozens of minutes or even hours, the dramatic and relatively abrupt brain state changes accompanying sleep onset and offset offer a potentially valuable model for understanding how changes in brain states can rapidly influence behavior and perception, irrespective of confounding factors that may operate at slower time scales.

Falling asleep is a continuous and dynamic process, reflecting a progressing sequence of changes in electrical, physiological, behavioral, and phenomenological phenomena (Lacaux et al., 2023; Ogilvie, 2001) governed by diminished activity in subcortical wake-promoting circuits co-occurring with increased sleep-promoting neuromodulation (Nir and de Lecea, 2023). Changes in heart rate, blood flow, eye movements, reaction times, as well as reports of subjective experience take minutes to develop and coincide with sequential changes in electroencephalogram (EEG) activity. Such changes range from diminished alpha oscillations and increased cortical theta activity, to the occurrence of sharp-waves, and ultimately to bona-fide sleep events such as K-Complexes and sleep spindles (Adamantidis et al., 2019; Hori et al., 1994). These changes lead one to consider falling sleep as an inherently gradual and continuous process, which does not necessarily occur at any single moment

(Ogilvie, 2001). On the other hand, measures of behavioral responsiveness, and possibly some EEG phenomena, do change rapidly around sleep onset, possibly at the order of a few seconds (Ogilvie, 2001; Strauss et al., 2022). This raises the question of whether there is a functionally-relevant abrupt binary switch within the extended process of falling asleep. We sought to address this open question by testing whether changes in cortical neuronal activities accompanying sleep onset exhibit abrupt or gradual dynamics.

To date, studies examining sleep onset with high temporal resolution (seconds) were typically performed non-invasively in humans. These studies have primarily focused on high-level neural aspects such as the timing and topographical changes in different EEG components and functional magnetic resonance imaging (fMRI) measurements (Fukunaga et al., 2006; Hori et al., 1994). Auditory EEG responses around sleep onset reveal that signatures of low-level auditory processing, as reflected in early EEG components or responses related to stimulus-specific adaptation, are not robustly modulated by sleep onset (Strauss et al., 2015). By contrast, measures of high-level processing such as P300, responses linked to predictive coding, and measures of baseline EEG complexity and connectivity, are strongly suppressed upon falling asleep (Strauss et al., 2015, 2022). However, non-invasive studies lack the ability to investigate processes at the neuronal level. On the other hand, studies focusing on sleep-related changes in cortical sensory responses at the neuronal level (Hayat et al., 2022; Issa and Wang, 2008, 2011, 2013; Marmelshtein et al., 2023; Sela et al., 2016, 2020) have not investigated modulations within seconds around state transitions. In this study, we aimed to bridge this gap by investigating, with high-temporal resolution, changes in neuronal activity in the auditory cortex as rats fell asleep or woke up. Our findings reveal that post-onset silence (or firing) undergoes substantial and abrupt changes within few seconds upon falling asleep (or awakening). At the same time, other aspects of auditory processing such as elevated population synchrony, vary at a slower time scale (tens of seconds or minutes), reflecting gradual aspects of the extended sleep onset process.

Results

 We investigated how sensory processing changes during the transitions between wakefulness and sleep by conducting sixteen experiments on adult male Wister rats (n = 6) implanted with microwire arrays targeting the auditory cortex, as well as monitoring with polysomnography including electroencephalography (EEG), electromyography (EMG), and synchronized video (Methods, Figure 1A-C). Rats were situated within a computer-controlled motorized running wheel placed inside an acoustic chamber for a total duration of 10 hours, commencing at light onset. In the first part of the experiment (5 hours), the wheel was periodically rotated for 4-7 seconds each time, alternating with idle intervals of 36-39 seconds (Figure 1D). In the second part of the experiment (recovery), rats were allowed to sleep undisturbed (while still in the wheel apparatus) for an additional 5 hours, during which they naturally transitioned between sleep and wakefulness. Throughout the entire 10h experimental period, we continuously presented auditory stimuli, 500ms-long click-trains at 2, 10, 20, 30 and 40Hz (Figure 1C) with an average inter-stimulus interval of 2s, played from an overhead free-field speaker (Methods) so that many of these sounds occurred around episodes of falling asleep or awakenings.

Figure 1. Experimental Setup. (A) (left): Acoustic chamber with an ultrasonic speaker and video monitoring, in which rats were confined to a motorized running wheel, while EEG, EMG, and

extracellular neural activity were measured continuously. (right): Motorized running wheel (Lafayette
instrument), to which the rats were confined. (B) Histology of microwire traces targeting the auditory
cortex. (C) Click-trains at varying rates presented throughout the experiment (D) Schematic depiction
of the experimental paradigm. For the first 5h (zeitgeber time [ZT] 0–5), rats were intermittently
forced to perform a short running bout with breaks allowing them to fall asleep. It was followed by 5h
of undisturbed recovery sleep opportunity (zeitgeber time [ZT] 5–10). Auditory stimuli were presented
throughout the entire experiment.
110

In the first 5h part of the experiment (periodic wheel rotation paradigm), rats often (44.3%) fell asleep
during idle intervals and awakened with each wheel rotation, thereby presenting numerous isolated
episodes of falling asleep (n=2,971 total, 186 ± 12.1 per session in n=16 sessions, mean ± SEM, Figure
2A, 2B, 2D). The second 5h part of the experiment (recovery sleep opportunity) also resulted in a
substantial number of falling asleep episodes (n=2,996 total, 187± 16.6 per session, mean ± SEM,
Figure 2A, 2C, 2D), as well as many spontaneous awakenings (n=2,718 total, 170± 17.2 per session,
mean ± SEM, Figure 2A, 2B, 2D). In total, data analysis was performed for 5967 episodes of falling
asleep and 2718 episodes of awakening.

119
Figure 2. Capturing moments of falling asleep with polysomnography. (A) Representative hypnogram
of a typical 10h session. Rows mark epochs of wakefulness (cyan), NREM sleep (black), and REM sleep
(purple). Red dashed lines mark two example moments of a rat falling asleep – either between wheel
rotations or during subsequent recovery sleep. (B) Representative Polysomnography of a trial of a rat
falling asleep during two consecutive 40s intervals between brief (5s) wheel rotations (vertical gray
bars). Rows (top to bottom) mark EMG, EEG, and automatically scored vigilance state (top – wake,
cyan; bottom – NREM sleep, black) as in panel A. (C) Same as B, but for an example of a rat falling
asleep during the recovery sleep opportunity period. (D) Count distribution across all sessions (n = 16)
of falling-asleep instances during the periodic wheel rotation paradigm, as well as falling-asleep
instances and awakenings during the recovery sleep opportunity period (left to right, respectively).
130

Identifying episodes of falling asleep and awakening

We first set out to automatically identify each falling asleep episode. To this end, we implemented the neural network-based automated AccuSleep algorithm (Barger et al., 2019) (Methods, Figure 3). The algorithm was trained on 1h-long manually-scored recovery period data containing intervals of wakefulness, non-rapid eye movement (NREM) sleep, and rapid eye movement (REM) sleep. We focused on the first hour of the recovery sleep opportunity that likely represents the deepest NREM sleep, in an effort to take a conservative approach in recognizing NREM and falling asleep episodes that refrains from subsequent analysis of borderline cases. Assessing the performance of the automatic sleep scoring method on the three sessions where sleep scoring was conducted fully (~10h each) we found that 93 ± 1.09 % of automatically scored 1s epochs correctly matched their corresponding manually scored labels (Figure 3D).

Figure 3. Automated sleep scoring by neural network for extracting falling asleep epochs. (A) Comparison of manual and automated sleep scoring of a representative session ~9h (top and bottom hypnograms, respectively). Red rectangle marks an example 6-minute segment of vigilance and sleep state changes expanded in panel B. (B) Polysomnography of a representative 6-minute segment. Rows (top to bottom) mark EMG, EEG, and manually and automatically scored vigilance state, as in A. Red rectangle marks further focus on 75 seconds expanded in panel C. (C) Same as in B, but for a shorter 75-second segment. (D) Confusion matrix between the vigilance states scored by automatic and manual sleep scoring over 3 sessions (~28h). Along the diagonal is the percentage of the automatically scored labels (column) that match the manually scored labels (row). Outside of the diagonal is the percentage of the automatically scored labels that do not match the manually scored labels but match the label of the relevant row.

Falling asleep suppresses sound-evoked post-onset neuronal firing within seconds

To analyze how neural processing and sensory responses in the auditory cortex changed around falling asleep with fine temporal resolution, we defined three intervals around wake-sleep transitions: "wakefulness" (-20 to -5sec before falling asleep), "First seconds of NREM" (+5 to +10sec after falling asleep), and "NREM" (>+60 after falling asleep). Based on our previous study on state-dependent auditory processing (Marmelshtein et al., 2023), we hypothesized that features of auditory cortical processing such as onset responses will maintain their magnitude when falling asleep, whereas features such as post-onset neuronal firing and population synchrony may be modulated during the process.

Figure 4A provides an example neuronal response of a representative unit to click stimuli, with trials vertically arranged according to their timing relative to the moment of falling asleep (top to bottom on y-axis). As can be seen for this unit (Figure 4A), as well as the average response (Figure 4B, n=152 units, 7 sessions), the onset response magnitude did not change around falling asleep (comparing wakefulness and first seconds of NREM). In contrast, the post-onset period (30-80 ms following sound onset) showed differences already during the first seconds of NREM sleep, in line with our hypothesis.

Expanding the analysis to all units (n=317) and multiple features of auditory processing (Figure 4C-E) revealed that onset response did not significantly change upon falling asleep, i.e. when comparing wakefulness and NREM sleep (Figure 4E, gray shading, -6.08% ± 4.65%, p = 0.19, t_{316} = -1.31, LME [Linear Mixed Effects Model]). Spontaneous firing rate and population synchrony exhibited moderate modulation (Figure 4C-E, gray shadings; -14.6% ± 5.81%, p = 0.013, t_{316} = -2.51 for reduction in spontaneous firing; 15.6% ± 4.26%, p = 2.9 × 10^{-4}, t_{316} = 3.67 for elevation in population synchrony, LME). The most notable change was observed for post-onset neuronal firing (n = 152 units), which was strongly modulated between wakefulness and NREM, as can be seen in Figure 4A-D (gray shadings) and as evident quantitatively (-56% ± 4.3%, p = 1.2 × 10^{-26}, t_{151} = -13.1, Figure 4E).

We further examined whether changes occur already within the first few seconds after falling asleep or, alternatively, reflect changes due to more consolidated NREM sleep within tens of seconds. As can be seen in the example unit in Figure 4A (dark blue) and the population response (Figure 4B, dark blue), post-onset firing was already significantly modulated within seconds after sleep onset. Quantifying this for the entire dataset (Figure 4C-E, n=317 units) revealed a sharp variation in post-onset firing within seconds of falling asleep (-51.2% ± 4.94%, p = 2.5 × 10^{-19}, t_{151} = -10.4, comparing wakefulness to first seconds of NREM). Changes in spontaneous firing rate, although more moderate, also occurred within seconds (-17.8% ± 5.47%, p = 1.2 × 10^{-3}, t_{316} = -3.26). In contrast, changes in

population synchrony were slower to occur and did not show significant modulation in the first seconds of NREM (2.26% ± 5.82%, p = 0.69, t_{316} = 0.38). Analysis of variance among the distinct features of

Figure 4. Post-onset firing changes dramatically within first seconds of falling asleep. (A) Representative raster and peristimulus time histogram (PSTH) for a unit in response to a click. Raster trials (rows) are ordered according to stimulus time relative to the proximate falling asleep instance. Azure, dark blue, and black shadings mark if the stimulus occurred while the animal was awake, in the first seconds after falling asleep, or during NREM sleep >60s after falling asleep, respectively. (B) Peristimulus time histogram of the mean normalized response to a click of all responsive units (n = 152, 7 sessions) during wakefulness, first seconds of NREM sleep, and NREM sleep following 60 seconds after the moment of falling asleep. (C) Modulation Index of activity and response features depending on stimulus time relative to the moment of falling asleep for all units (n = 152, 7 sessions for Post-Onset FR and n = 317, 15 sessions, for the other features). Azure, dark blue, and black shadings

cortical auditory processing features confirmed that they are differentially modulated within the first second of NREM ($p = 2.1 \times 10^{-4}$, n = 6 animals, Friedman test). In addition, upon entering a deeper state of NREM sleep, processing features were also differentially modulated ($p = 1.1 \times 10^{-4}$, n = 6 animals, Friedman test). Pairwise comparisons revealed that post-onset firing was modulated significantly more strongly than the other three features ($p < 2.1 \times 10^{-10}$, df = 151, for all three features within first seconds of NREM, and for all three features within NREM state, $p < 1.6 \times 10^{-14}$, df = 151, LME). The slower buildup of population synchrony can be appreciated most clearly when contrasting Figure 4C (robust changes in temporal dynamics at a scale of tens of seconds) with Figure 4D (no changes in temporal dynamics in first seconds). Indeed, a direct comparison established that the difference between the first seconds of NREM and deeper NREM was significantly larger for population synchrony modulation than for post-onset modulation ($p = 0.023$, df = 151, $t_{151} = 2.3$, LME).

Awakening restores sound-evoked post-onset neuronal firing within seconds

Does cortical auditory processing show an opposite profile upon awakening to that observed when falling asleep? To test this, we compared neuronal activity in the seconds just before awakening ("Sleep", -20 to -5sec before awakening) and just after awakening ("First seconds of wakefulness, +5 to +10sec after awakening). Figure 5A provides a representative example of a unit firing around awakening. As can be seen, post-onset activity rapidly changes from silence during sleep to firing in wakefulness, whereas population synchrony is minimally affected at these time-scales.

Quantifying these effects across the entire dataset (n=315 units, Figure 5A-D), we found a modest albeit significant increase in onset firing rate that occurred both in the first seconds of wakefulness, (11.7% ± 3.82%, $p = 2.6 \times 10^{-3}$, $t_{314} = 3.04$, LME). Similarly, we found moderately higher modulation indices representing elevation in spontaneous firing rate (19.8% ± 6.72%, $p = 3.5 \times 10^{-3}$, $t_{314} = 2.94$). As observed upon falling asleep, changes in post-onset firing were most pronounced and occurred within the first seconds of state transitions (n = 145 units, 42.3% ± 5.46%, $p = 1.6 \times 10^{-12}$, $t_{144} = 7.74$). Population synchrony, as was the case when falling asleep, did not significantly change within the first

few seconds around state transitions (-5.2% ± 4.2%, p = 0.22, t_{314} = -1.23). Together, awakening exhibits a "mirror-image" of the changes in cortical auditory activity to those found around falling asleep, with post-onset firing remaining the aspect that changes most strongly and rapidly. Analysis of variance among the distinct features of cortical auditory processing confirmed that they are differentially modulated within the first second of wakefulness (p = 3.3×10^{-4}, n = 6 animals, Friedman test). Pairwise comparisons revealed that post-onset firing was modulated significantly more strongly than the other three features (p < 3.8×10^{-6}, df = 144, for all three features, LME).

Figure 5 post-onset firing changes dramatically within first seconds of awakening. (A) Representative raster and peristimulus time histogram (PSTH) for a unit in response to a click. Raster trials (rows) are ordered according to the trial's stimulus time relative to the proximate awakening instance. Black and blue shadings mark if the stimulus occurred while the animal was asleep or in the first seconds after waking up, respectively. (B) Peristimulus time histogram of the mean normalized response to a click of all responsive units (n = 145, 7 sessions) during sleep and first seconds of wakefulness. (C) Modulation Index of activity and response features depending on stimulus time relative to the moment of awakening. Black and blue shadings mark "sleep" and "first seconds of wakefulness" periods as in A. Yellow shading represents the standard error of the mean (SEM). (D) Modulation Index of activity and response features between "first seconds wakefulness" and "sleep" states across units and sessions (n = 145, 7 sessions for Post-Onset FR and n = 315, 15 sessions, for the other features). Bars and black error bars represent the mean±SEM estimated by the Linear Mixed Effects model. Small grey markers represent individual units (red for the representative unit in panel A). Large grey markers represent individual session mean (with each shape representing an individual animal).

Discussion

We studied how neuronal spiking activity and responses to clicks in early rat auditory cortex change in the seconds around falling asleep and waking up. We found that a robust increase in the occurrence of post-onset neuronal silent periods (OFF states) and moderate reductions in spontaneous firing rates - both known to occur in NREM sleep (Marmelshtein et al., 2023) - changed abruptly around the first seconds of falling asleep. These changes in activity and responsivity during the first (~5sec) moments of sleep reached their full magnitude as observed in later consolidated NREM sleep (>60 sec from the moment of falling asleep). By contrast, an increase in neuronal population synchrony only changed substantially during later (>60 sec) NREM sleep without exhibiting robust modulations around the seconds of falling asleep. Onset response magnitude was unaffected by falling asleep or awakening, echoing previous studies showing its invariance during sleep (Issa and Wang, 2008; Nir et al., 2013). During the first seconds after awakenings, we found opposite changes in neuronal activity and responsivity (rapid and robust decrease in the occurrence of post-onset silent periods, and rapid moderate elevation in spontaneous firing rates). These changes showed identical temporal dynamics, but with opposite directionality, to those observed when falling asleep. Together, the results indicate that some changes in auditory cortex activity occur within seconds around wake-sleep transitions whereas specific aspects such as population synchrony vary with slower dynamics. Auditory-induced silent OFF periods emerge as the aspect that changes most robustly around state-transitions, showing abrupt and large changes in its magnitude within seconds.

How do the present results relate to previous studies on changes in auditory responses across states of arousal? A recent study examined the effect of gradual propofol anesthesia on auditory processing (Bergman et al., 2022). There, observed changes were gradual and tightly locked to increasing anesthetic concentration rather than to more abrupt loss of behavioral responsiveness. Furthermore, we recently showed that the degree of post-onset neuronal firing decreased gradually with mounting sleep pressure across several hours of sleep deprivation (Marmelshtein et al., 2023). Thus, gradual changes in arousal often lead to gradual changes in auditory processing. Here, however, we were able to examine, for the first time, abrupt transitions in arousal states around falling asleep and waking up episodes. Under these conditions, we find that sharp step-like changes can occur at a timescale of few seconds, as demonstrated for the decrease in post-onset firing accompanying sleep onset. Thus, we suggest that arousal-related changes in neural responses can be either gradual or abrupt, depending on the nature of the change in arousal state. Accordingly, gradual changes in arousal state (e.g. mounting sleep pressure or deepening anesthesia), will be associated with gradual changes in neural

responses, whereas abrupt changes in arousal state (e.g. falling asleep or awakening) will be associated
with sharp changes in neural responses.

Beyond neuronal correlates in specific pathways such as the auditory cortex, what do our findings
reveal more generally about falling asleep – is it fundamentally a gradual or abrupt process? While
extensive evidence shows that many electrical, physiological and behavioral phenomena change
gradually in the span of minutes (Eban-Rothschild et al., 2016; Lacaux et al., 2023; Ogilvie, 2001), some
EEG phenomena and specific measures of behavioral responsiveness do show fast temporal dynamics
(Ogilvie, 2001; Strauss et al., 2022). We find evidence supporting both of these seemingly contradicting
themes. On the one hand, post-onset neuronal silence developed rapidly, within just few seconds upon
sleep onset. Exposing these fast dynamics relied critically on our ability to sleep score our data at high
temporal resolution (1 second). On the other hand, population synchrony displayed different temporal
dynamics, developing gradually on the time scale of one minute. Similarly, measures of EEG
information and connectivity were also found to change on such a slow time-scale (Strauss et al.,
2022). This implies that these two measures may reflect different underlying neural processes both
accompanying sleep onset, one that is continuous and gradual, and another abrupt.

The functional significance of increase in post-onset neuronal silence remains unclear. It is likely that
neuronal silence as captured in extracellular recordings reflects a state of neuronal bistability where
neurons alternative between active (up) and hyperpolarized silent (down) states , a key feature of sleep
and unresponsive states (Marmelshtein et al., 2023; Massimini et al., 2007; Vyazovskiy et al., 2013,
2011). While such bistability does not interfere with short-lived localized onset responses, it can
disrupt complex neural processing by breaking-off sustained and widespread propagation of sensory
signals (Massimini et al., 2005; Pigorini et al., 2015). Indeed, states of drowsiness and sleep associated
with increased bistability affect processing requiring integration beyond local activity patterns.
Accordingly, drowsiness preferentially disrupts perceptual discrimination, rather than perceptual
detection (Xu et al., 2023). Moreover, measures of complex neural processing, such as predictive
coding, are preferentially affected by sleep onset (Strauss et al., 2015), and baseline complexity
measures were maximally attenuated in NREM sleep when stimuli evoked a k-complex, reflecting high
neuronal bistability (Andrillon et al., 2016).

Certain limitations in our study require explicit acknowledgment. First, it is not trivial to generalize
results about the precise dynamics around state transitions from rodents to humans and other species.
Since rodents exhibit more fragmented sleep and shorter bouts of sleep and wakefulness (Lo et al.,
2004), the first few seconds of sleep or wakefulness in rats may or may not be directly comparable to
the first few seconds of longer sleep bouts. Nevertheless, the fact that some aspects of auditory

processing changed robustly and abruptly (post-onset firing) whereas other aspects changed more

modestly (ongoing firing rates) or more gradually (population synchrony) goes beyond differences

across species. Second, while our recordings focused on (early) auditory cortex, it remains unknown

what is the spatial scale of these effects, and to what extent they occur across other regions. On one

hand, the observed changes in auditory responses may be inherited from earlier auditory stations such

as the auditory thalamus, a possibility that we cannot rule out with this dataset. On the other hand,

changes in neural processing may be pervasive and reflect general effects on cortical processing.

Further studies are needed to examine to what extent such findings can be generalized to other

sensory modalities and cortical areas. Finally, some limitations relate to our extracellular recordings,

discussed also previously (Marmelshtein et al., 2023); these include a bias for larger pyramidal cells

with high baseline firing rates as well as lack of information on cortical layer and cell types, possibly

biasing our results towards specific neuronal subpopulations.

In conclusion, our investigation reveals the temporal dynamics and magnitude of modulation in

different features of auditory processing upon falling asleep and awakening. Our findings extend

previous studies in showing that specific neural signatures of auditory responses, particularly the

degree of post-onset firing, are robustly influenced by vigilance states and their transitions

(Marmelshtein et al., 2023). A rapid and robust dramatic increase in post-onset neuronal silence,

occurring within seconds of falling asleep, contrasts with the slower modulation observed in

population synchrony, and emerges as a robust signature of altered responsivity of the sleeping brain

to external sensory stimuli.

Methods

Animals

The study involved electrophysiology experiments in six adult male Wistar rats (weight: 300-350g, age: >12 weeks). Rats were kept in transparent Perspex cages with food and water available ad libitum. The ambient temperature was maintained between 20-24 Celsius, and there was a 12:12 hours light/dark cycle with light onset at 10:00 AM. All experimental procedures, including animal handling, wheel cycle paradigm, and surgery, were conducted in accordance with the National Institutes of Health's Guide for the care and use of laboratory animals and approved by the Institutional Animal Care and Use Committee of Tel Aviv University.

Surgery and electrode implantation

Before surgery, DiI fluorescent dye (DiIC18, Invitrogen) was applied to microwire arrays under microscopic control to aid in localization. Generally, surgical procedures followed our previous protocols (Sela et al., 2020). First, isoflurane (4%) was used to induce general anesthesia, and animals were then placed in a stereotactic frame (David Kopf Instruments). During the rest of the surgery, anesthesia (isoflurane, 1.5-2%) and a 37C body temperature (closed-loop heating pad system, Harvard Apparatus) were maintained. Animals were given antibiotics (Cefazolin, 20 mg/kg i.m.), analgesia (Carpofen, 5 mg/kg i.p.), and dexamethasone (0.5 mg/kg, i.p.). Their scalp was shaved, and liquid gel (Viscotears) was applied to protect their eyes. Subcutaneous infusion of lignocaine (7 mg/kg) was performed before incision, and then the skull was exposed and cleaned. Two frontal screws (one on each hemisphere, 1mm in diameter) and a single parietal screw (left hemisphere) were placed in the skull for recording EEG. Two screws were placed above the cerebellum, serving as reference and ground. Two single-stranded stainless-steel wires were inserted to the neck muscles to record EMG. EEG and EMG wires were to a head-stage connector (Omnetics). Dental cement was applied to cover all screws and wires. A small craniotomy was performed over the right hemisphere, and the dura was meticulously dissected. An auditory cortex-targeting 16-electrode microwire array (Tucker-Davis Technologies, TDT, 33 or 50mm wire diameter, 6-6.5 mm long, 15 tip angle; arrays consisting of 2 rows 3 8 wires, with 375mm medial-lateral separation between rows and 250mm anterior–posterior separation within each row) was implanted diagonally (angle of 28 degrees, see Figure 1B) with insertion point center coordinates of P: -4.30mm, L: 4.5mm relative to Bregma, and inserted until a final depth of 4.6mm was reached. Silicone gel (Kwik-Sil; World Precision Instruments) was applied to cover the craniotomy, and Fusio (Pentron) was used to fix the microwire array in place. After surgery, chloramphenicol 3% ointment was applied topically, and buprenorphine (0.025 mg/kg s.c.) was given

systemically for additional post-operative analgesia. Dexamethasone (1.3 mg/kg) was given with food

in the days following the surgery to reduce pain and inflammation around implantation.

Histology

After conducting the experiments, we confirmed the position of electrodes in 4 out of 6 animals

through histology, as shown in Figure1B. Animals were anesthetized with 5% isoflurane and perfused

with 4% paraformaldehyde. The brains were refrigerated in paraformaldehyde for a week and then cut

into 50-60mm serial coronal sections using a vibrating microtome (Leica Biosystems). The sections

were stained with fluorescent cresyl violet/Nissl (Rhenium). The histological verification ensured that

the electrodes were located within areas Au1/AAF/AuD as defined by Paxinos and Watson (Paxinos

and Watson, 2006).

Electrophysiology

Data were acquired using a RZ2 processor (TDT). Microwire extracellular activity was digitally sampled

at 24.4 kHz (PZ2 amplifier), while EEG and EMG were pre-amplified (RA16LI , TDT) and sampled at

256.9 Hz (PZ2 amplifier, TDT). Spike sorting was performed with "wave_clus" (Quiroga et al., 2004),

which utilized a detection threshold of five SDs and automatic superparamagnetic clustering of wavelet

coefficients. The clusters were then manually selected, refined, and tagged as either multi-units (MU)

or single-units (SU) based on their stability throughout the recording, quality of separation from other

clusters, consistency of spike waveforms, and inter-spike interval distributions (N_{MU} = 332, N_{SU} = 88),

as described in (Nir et al., 2015). We verified the accurate localization of the auditory cortex, which

includes A1, the anterior auditory field, or the dorsal auditory cortex, using histology (Figure 1B) and

by analyzing the units' response latency to auditory stimuli (Marmelshtein et al., 2023).

Experimental Design

Following surgery, the subjects were housed separately and introduced to the motorized running

wheel for a few hours every day for habituation (Figure 1A). They were then gradually introduced to

the periodic wheel rotations to be performed in the first experimental phase (Figure 1D, see below).

We conducted 16 experimental sessions in the following manner. At the start of the day (10 AM), the

rats were moved from their home cage to a motorized running wheel (Figure 1A, Model 80860B,

Lafayette Instrument) placed within a sound-attenuation chamber (-55dB, H.N.A) and underwent five

hours of periodic wheel rotation. During the first experimental phase, the wheel was rotated for 3

seconds every 36-39 seconds, which constituted intervals when rats could behave freely (run, groom,

or fall asleep). In the second experimental phase, rats were allowed to rest undisturbed for an

additional 5 hours in the (fixed) wheel, which constituted a recovery sleep opportunity (Figure 1D).

The rats were kept in the wheel apparatus under the ultrasonic speaker for the entire session. Auditory stimulation was delivered intermittently throughout both the sleep deprivation and recovery sleep periods, irrespective of the wheel's movement.

Auditory stimulation

We used Matlab (MathWorks) to compile sound waveforms that were then transduced into voltage signals using a high-sampling rate sound card (192 kHz, LynxTWO, Lynx). These signals were amplified (SA1, TDT) and played through a magnetic speaker (MF1, TDT), mounted 60 cm above the motorized running wheel. We conducted two different auditory experiments over separate sessions/days:

Auditory paradigm A

This paradigm was used in 7 sessions in 6 animals: Stimuli included click trains and a set of Dynamic Random Chords (DRC)(Linden et al., 2003). Click trains lasted 500ms and were played at rates of either {2, 10, 20, 30, 40} clicks/sec (Figure 1C). A typical 10-hour session contained 2000 blocks, each consisting of a single DRC stimulus and a single repetition of each click train (presented at random order), with an inter-stimulus interval of 2s and ±0.25s jitter.

Auditory paradigm B

This paradigm was used in 9 sessions in 5 animals: The stimuli used in the study included a 40 clicks/second click-train and a set of DRC stimuli. Each 10-hour session had 600 blocks, with each block consisting of a single DRC stimulus and 4 repetitions of the 40 Hz click train, presented in random order. The inter-stimulus interval was 2 seconds, with a ±0.25-second jitter. Both paradigms included an 8-second inter-stimulus interval every 2 minutes.

Acoustic sound-level measurements

We used a high-quality ultrasonic microphone (model 378C01, PCB Piezotronics) to measure sound levels at 12 positions around the circumference of the wheel apparatus (anterior-posterior axis) and at five positions along its width (left-right axis) with a spacing of 2cm. This covered the area where animals were present during the experimental sessions. We found that the root-mean-square (RMS) intensity in response to 2-second white noise sounds had minimal variance at different positions along the anterior-posterior axis, with a mean absolute deviation of 0.7±0.3dB (±Std. Dev) and a maximum difference of 2.4dB between any two positions. Similarly, along the left-right axis, we recorded a mean absolute difference of 0.4±0.3dB (±Std. Dev.) and a maximum difference of 1.5dB. When we examined sound levels at different frequency bands (1, 2, 4, 8, 16, 32, and 64 KHz, in response to the same white noise stimulus) at different positions, we found similarly low variation (all below 4dB)).

Sleep scoring and identification of state transitions

In each session (n=16), at least one hour (the last hour, representing recovery sleep opportunity period) was manually sleep scored, employing visual inspection of EEGs, EMGs, and video/behavior as in previous studies (Rodriguez et al., 2016; Sela et al., 2020; Vyazovskiy et al., 2011). In an additional three sessions, manual sleep scoring of the entire 8-9h experiment was performed and used for assessing the performance of automatic sleep scoring (Figure 3A-D).

Before any sleep scoring, we excluded any periods when the wheel was moving (forced running bouts during the first experimental phase). Next, we categorized different time periods to either active-wakefulness with behavioral activity (e.g., locomotion, grooming) as confirmed by video, quiescent-wakefulness (low-voltage high-frequency EEG activity and high tonic EMG with occasional phasic activity), NREM sleep (high-amplitude slow wave activity and low tonic EMG activity), REM sleep (low-amplitude wake-like frontal EEG co-occurring with theta activity in parietal EEG and flat EMG, Figure 2A-C), or unknown periods not analyzed further (e.g. state transitions, to conservatively avoid these epochs in subsequent analysis).

Next, the 1h+ manually-scored data in each session was used to train an automated sleep-scoring algorithm using convolutional neural networks (CNNs) (Barger et al., 2019). The algorithm was fed with the manual sleep scoring vector along with the frontal EEG signal (low-pass filtered < 60Hz), and the EMG signal (band-pass filtered 10Hz – 125Hz). Manually scored labels, originally tagged in 1ms resolution, were re-labeled into 1sec epochs, where each epoch was labeled according to the most frequent manual label in that interval. For each rat, a personalized CNN was trained using the manually labeled data of all the sessions of the same rat (1-4 sessions per rat). Weights for mixture z-scoring, for both training and classification, were calculated using the relative prevalence of each state in the data used for training. Epoch length was 1 sec. The CNNs trained with consideration of 9 epochs surrounding each epoch scored (four epochs preceding and four epochs following the current epoch). Calibration data for mixture z-scoring for classification was also provided as the training data. Using the relevant rat's CNN for each session, the entire data set was classified into the sleep scoring labels. After applying the classification method, we designed a post-processing method for the labels to meet our goal for early detection of true fall-asleep (awakening) transitions. In which case, for each session, the epochs labels were examined from the last one to the first one. Epochs with low certainty levels in all states, i.e., below 50%, were set to be the same as the label of the following epoch (the later in time). Then, epochs that were labeled differently from their two neighboring (preceding and following) epochs, and their neighbors had the same label, were changed to match their neighbors. Any epochs containing times when the wheel was moving were excluded from the analysis.

Extracting falling asleep and awakening episodes, based on the automatic sleep scoring, was

performed as follows. We considered a falling asleep instance as a continuous sequence (≥2 sec) of

NREM epochs (epoch = 1 sec), preceded by a continuous sequence (≥2 sec) of quiescent-wakefulness

epochs. The beginning of the first NREM epoch was considered, for our matter, the moment of falling

asleep. For the subsequent analysis of neural activity, a symmetrical time window around the falling

asleep moment was taken. The window extended as long as possible without including periods of rat

movement, wheel movement, or epochs that are not part of quiescent-wakefulness or NREM

sequences. Falling asleep instances were discarded from the subsequent analysis if more than one

falling asleep transition occurred within the same 36-39sec interval between consecutive wheel cycles.

Awakening instances were extracted only from the spontaneous sleep opportunity phase. We

considered an awakening instance in a similar way, a continuous sequence of quiescent-wakefulness

epochs (≥2 sec), preceded by a continuous sequence (≥2 sec) of NREM epochs. The beginning of the

first quiescent-wakefulness epoch was considered, for our matter, the moment of awakening. For the

subsequent analysis of neural activity, a symmetrical time window around the awakening moment was

taken. The window extended as long as possible without including periods of rat movement, or epochs

that are not part of quiescent-wakefulness or NREM sequences. Falling asleep and awakening

instances with a time window shorter than 3 sec were not analyzed.

QUANTIFICATION AND STATISTICAL ANALYSIS

Analysis of auditory responses across states

Auditory stimuli presented during the times of wheel movement or while the rat was active

(locomotion, grooming for example) were excluded from analysis.

Analysis of responses to click trains (Figures 4C, 4D, 4E, 5C, 5D)

To measure spontaneous firing rate, we calculated the average firing rate during the [-500,0]ms

window leading up to the click-train stimuli. For post-onset firing rate, we calculated the average firing

rate during the [30,80]ms window. We analyzed onset response to different click rates using smoothed

peri-stimulus time histograms (PSTH) with a Gaussian kernel (σ=2ms). The onset response was

determined by extracting the highest firing rate during the [0,50]ms window of the smoothed PSTH.

Population synchrony was defined as population coupling (Okun et al., 2015) and calculated as the

correlation of each unit firing to that of the entire neuronal population average in 50ms bins during

the baseline period ([-1000,0]ms). We calculated population coupling for each trial baseline period

and then averaged for all trials in a given condition.

Feature modulation between states (Figures 4E, 5D)

For each session, all auditory stimulation trials that appeared around falling asleep (or awakening) episodes were sorted according to their time relative to the moment of falling asleep (or awakening). We defined three vigilance states in the falling asleep process: "wakefulness" / "First seconds of NREM" / "NREM", with different time ranges relative to the moment of falling asleep (-20s < t < -5s, 5s < t < 10s, 60s < t; respectively). Two states were also defined for awakening episodes: "Sleep" / "First seconds of wakefulness" (-20s < t < -5s, 5s < t < 10s; respectively). Auditory trials around moments of falling asleep / awakenings were sorted into these states.

For each feature of auditory processing (spontaneous firing, onset response, population synchrony, post-onset firing), we pooled trials from all relevant click-trains presented (2,10,20,30,40 Hz): for spontaneous firing rate and population synchrony we examined baseline activity of all click trains; for onset firing rate, responses to all click trains; and for post-onset firing rate click trains of 2 Clicks/sec and 10 Clicks/sec (allowing sufficient inter click interval to not interfere with the post-onset period). Within the states and relevant stimuli pooled for each features the unit response among all trials was averaged resulting PSTH. The response analysis, as described for each feature was performed on each PSTH, gaining features value for all states. To compare responses between the vigilance states, modulation index was calculated between the feature's value in the first seconds of NREM state (first seconds of wakefulness, for awakening) to the value in the wakefulness state (Sleep, for awakening) and between the feature's value in the NREM state (wakefulness, for awakening) to the value in wakefulness state (Sleep, for awakening).

Modulation index between two conditions for all measures above was calculated as:

$$\frac{ConditionA - ConditionB}{max(ConditionA, ConditionB)} \times 100$$

Feature analysis along arousal state transitions (Figures 4A, 4C, 4D, 4E, 5A, 5C, 5D)

For each session and per each feature, auditory trials within falling asleep (awakening) episodes were extracted and sorted by time relative to the moment of falling asleep (awakening) as before, yielding a sorted unit response raster (Figure 4A). For each unit, the responses for adjacent trials (in terms of time relative to the moment of falling asleep/awakening) were smoothed (except for population synchrony) by a moving average window (uniformly weighted), with the size of 50 trials and an overlap of 49 trials. Due to unevenly spaced stimuli's relative times, new time stamps were created by smoothing the original stimuli relative times with the same method. Resulting in a modified raster where each row represents a 50-trial averaged response and its average stimuli time. Thereafter, a

modulation index was calculated between the averaged value of the feature in each (averaged) time

stamp compared to that during "wakefulness" ("Sleep" for awakening) state. For population synchrony,

the feature values were first calculated for each trial, and then feature values and time stamps were

smoothed using a 50-trial window, gaining the state value.

The continuous modulation index for each feature was averaged among all units within a session to

gain the sessions' mean continuous modulation index. Next, for each session and feature, the

continuous index was interpolated, using the 'interp1' function with 'pchip' as the interpolation

method (MATLAB, MathWorks), to gain evenly spaced and mutual values for all sessions. Thereafter,

the continuous values were averaged into a grand mean of all sessions of continuous modulation index

along the falling asleep (awakening) process (figures 4C-D, 5C).

Rejecting non-significant units

Units with an average spontaneous firing rate <0.5 spikes/sec in any of the vigilance states were

excluded from the analysis. Units without a significant response to a single-click stimulus were also

excluded from our analysis. The significance of the response was calculated using the Monte Carlo

Permutation Test, which was conducted as follows. A unit's PSTH, in the time interval of [-500ms,

30ms] relative to the beginning of each 2 Hz and 10 Hz Clicks/sec trial, underwent minimal temporal

smoothing (gaussian kernel of $\sigma = 2$). The significance was examined separately for the trials before

the moment of falling asleep (or awakening) and for the trials after the moment of falling asleep (or

awakening). Among the trials of each group, the PSTHs in the baseline interval ([-500ms,0] before click

onset) were averaged, as well as the activity in the onset interval ([0, 30ms] after click onset). The

difference between the maximum of the averaged PSTH in the time window of the onset and the

baseline was calculated. After that, we shuffled between the baselines and the onsets of the smoothed

PSTHs (that is, randomly, for 50% of the trials, an exchange was made between the time window of

the baseline and the time window of the onset), and then again, we averaged over all the trials, and

calculated the difference in the firing rate in the time windows of onset to baseline. This process was

performed 1000 times to calculate the surrogate distribution from which the p-value of the real onset

response magnitude was derived. Units whose response was not significant, for both before and after

falling asleep (or awakening), were also excluded from the analysis.

Statistics

We used a linear mixed-effects model (LME) to analyze electrophysiological neural data due to its

nested and hierarchical nature (Aarts et al., 2014; Makin and De Xivry, 2019). This model was

necessary to account for non-independencies in measures from different units obtained in the same

electrode, experimental session, or animal. We included animal identity as a random effect, as well as

experimental session and microwire electrode as nested random effects within each animal. We

obtained model parameters using the 'fitlme' function (Matlab, MathWorks) and utilized restricted

maximum likelihood estimation.

When presenting average effects per session, we only considered sessions that had a minimum of 3

units. Throughout the main text, all effect sizes are presented as mean±SEM for all units unless

specified otherwise. To evaluate variance across different conditions (>2), we conducted a Friedman

test (similar to a non-parametric repeated measures ANOVA) on data summarized at the animal level.

This involved averaging all the units for each of the 6 animals.

Acknowledgments

We thank Dr. Noa Bar-Ilan Regev for administrative assistance. Supported by the Israel Science Foundation (ISF) grants 1326/15 and 1557/22 and by the European Research Council grant ERC-2019-CoG 864353.

Author contributions

A.M. and Y.N. designed the project. A.M. built experimental setup, compiled stimuli, and performed experiments. B.L. led data analysis with support from A.M. and B.H. Y.N. supervised the project and secured funding. All authors wrote the paper.

Declaration of interests

The authors declare that no financial and non-financial competing interests exist.

References

Aarts, E., Verhage, M., Veenvliet, J. V., Dolan, C. V., and Van Der Sluis, S. (2014). A solution to dependency: Using multilevel analysis to accommodate nested data. Nat. Neurosci. *17*, 491–496.

Adamantidis, A.R., Gutierrez Herrera, C., and Gent, T.C. (2019). Oscillating circuitries in the sleeping brain. Nat. Rev. Neurosci. *20*, 746–762.

Andrillon, T., Poulsen, A.T., Hansen, L.K., L É Ger, D., and Kouider, S. (2016). Neural markers of responsiveness to the environment in human sleep. J. Neurosci. *36*, 6583–6596.

Atiani, S., Elhilali, M., David, S. V., Fritz, J.B., and Shamma, S. a. (2009). Task Difficulty and Performance Induce Diverse Adaptive Patterns in Gain and Shape of Primary Auditory Cortical Receptive Fields. Neuron *61*, 467–480.

Barger, Z., Frye, C.G., Liu, D., Dan, Y., and Bouchard, K.E. (2019). Robust, automated sleep scoring by a compact neural network with distributional shift correction. PLoS One *14*, 1–18.

Bergman, L., Krom, A.J., Sela, Y., Marmelshtein, A., Hayat, H., Regev, N., and Nir, Y. (2022). Propofol Anesthesia Concentration Rather Than Abrupt Behavioral Unresponsiveness Linearly Degrades Responses in the Rat Primary Auditory Cortex. Cereb. Cortex.

Eban-Rothschild, A., Rothschild, G., Giardino, W.J., Jones, J.R., and de Lecea, L. (2016). VTA dopaminergic neurons regulate ethologically relevant sleep–wake behaviors. Nat. Neurosci. *19*, 1–14.

Edeline, J.M., Dutrieux, G., Manunta, Y., and Hennevin, E. (2001). Diversity of receptive field changes in auditory cortex during natural sleep. Eur. J. Neurosci. *14*, 1865–1880.

Fukunaga, M., Horovitz, S.G., van Gelderen, P., de Zwart, J.A., Jansma, J.M., Ikonomidou, V.N., Chu, R., Deckers, R.H.R., Leopold, D.A., and Duyn, J.H. (2006). Large-amplitude, spatially correlated fluctuations in BOLD fMRI signals during extended rest and early sleep stages. Magn. Reson. Imaging *24*, 979–992.

Harris, K.D., and Thiele, A. (2011). Cortical state and attention. Nat. Rev. Neurosci. *12*, 509–523.

Hayat, H., Marmelshtein, A., Krom, A.J., Sela, Y., Tankus, A., Strauss, I., Fahoum, F., Fried, I., and Nir, Y. (2022). Reduced neural feedback signaling despite robust neuron and gamma auditory responses during human sleep. Nat. Neurosci. *25*, 935–943.

Hori, T., Hayashi, M., and Morikawa, T. (1994). Topographical EEG changes and the hypnagogic

experience.

Issa, E.B., and Wang, X. (2008). Sensory responses during sleep in primate primary and secondary

auditory cortex. J Neurosci *28*, 14467–14480.

Issa, E.B., and Wang, X. (2011). Altered neural responses to sounds in primate primary auditory

cortex during slow-wave sleep. J. Neurosci. *31*, 2965–2973.

Issa, E.B., and Wang, X. (2013). Increased neural correlations in primate auditory cortex during slow-

wave sleep. J. Neurophysiol. *109*, 2732–2738.

Lacaux, C., Strauss, M., Bekinschtein, T., and Oudiette, D. (2023). Embracing sleep onset complexity.

Lee, S.H., and Dan, Y. (2012). Neuromodulation of Brain States. Neuron *76*, 109–222.

Linden, J.F., Liu, R.C., Sahani, M., Schreiner, C.E., and Merzenich, M.M. (2003). Spectrotemporal

structure of receptive fields in areas AI and AAF of mouse auditory cortex. J. Neurophysiol. *90*, 2660–

2675.

Lo, C.C., Chou, T., Penzel, T., Scammell, T.E., Strecker, R.E., Stanley, H.E., and Ivanov, P.C. (2004).

Common scale-invariant patterns of sleep-wake transitions across mammalian species. Proc. Natl.

Acad. Sci. U. S. A. *101*, 17545–17548.

Makin, T.R., and De Xivry, J.J.O. (2019). Ten common statistical mistakes to watch out for when

writing or reviewing a manuscript. Elife *8*.

Marmelshtein, A., Eckerling, A., Hadad, B., Ben-Eliyahu, S., and Nir, Y. (2023). Sleep-like changes in

neural processing emerge during sleep deprivation in early auditory cortex. Curr. Biol. *33*, 2925-

2940.e6.

Massimini, M., Ferrarelli, F., Huber, R., Esser, S.K., Singh, H., and Tononi, G. (2005). Breakdown of

cortical effective connectivity during sleep. Science (80-.). *309*, 2228–2232.

Massimini, M., Ferrarelli, F., Esser, S.K., Riedner, B.A., Huber, R., Murphy, M., Peterson, M.J., and

Tononi, G. (2007). Triggering sleep slow waves by transcranial magnetic stimulation. Proc. Natl.

Acad. Sci. U. S. A. *104*, 8496–8501.

Mcginley, M.J., Vinck, M., Reimer, J., Batista-brito, R., Zagha, E., Cadwell, C.R., Tolias, A.S., Cardin,

J.A., and Mccormick, D.A. (2015). Waking State : Rapid Variations Modulate Neural and Behavioral

Responses. Neuron *87*, 1143–1161.

Nir, Y., and de Lecea, L. (2023). Sleep and vigilance states: Embracing spatiotemporal dynamics.

Neuron *111*, 1998–2011.

Nir, Y., Vyazovskiy, V. V, Cirelli, C., Banks, M.I., and Tononi, G. (2013). Auditory Responses and

Stimulus-Specific Adaptation in Rat Auditory Cortex are Preserved Across NREM and REM Sleep.

Cereb. Cortex bht328-.

Ogilvie, R.D. (2001). The process of falling asleep. Sleep Med. Rev. *5*, 247–270.

Paxinos, G., and Watson, C. (2006). The rat brain in stereotaxic coordinates: hard cover edition

(Elsevier).

Pigorini, A., Sarasso, S., Proserpio, P., Szymanski, C., Arnulfo, G., Casarotto, S., Fecchio, M., Rosanova,

M., Mariotti, M., Lo Russo, G., et al. (2015). Bistability breaks-off deterministic responses to

intracortical stimulation during non-REM sleep. Neuroimage *112*, 105–113.

Quiroga, R.Q., Nadasdy, Z., and Ben-Shaul, Y. (2004). Unsupervised spike detection and sorting with

wavelets and superparamagnetic clustering. Neural Comput. *16*, 1661–1687.

Rodriguez, X.A. V, Funk, X.C.M., Vyazovskiy, X.V. V, Nir, X.Y., Tononi, G., and Cirelli, X.C. (2016). Why

Does Sleep Slow-Wave Activity Increase After Extended Wake ? Assessing the Effects of Increased

Cortical Firing During Wake and Sleep. *36*, 12436–12447.

Sela, Y., Vyazovskiy, V. V., Cirelli, C., Tononi, G., and Nir, Y. (2016). Responses in rat core auditory

cortex are preserved during sleep spindle oscillations. Sleep *39*, 1069–1082.

Sela, Y., Krom, A.J., Bergman, L., Regev, N., and Nir, Y. (2020). Sleep differentially affects early and

late neuronal responses to sounds in auditory and perirhinal cortices. J. Neurosci. *40*, 2895–2905.

Strauss, M., Sitt, J.D., King, J.R., Elbaz, M., Azizi, L., Buiatti, M., Naccache, L., Van Wassenhove, V., and

Dehaene, S. (2015). Disruption of hierarchical predictive coding during sleep. Proc. Natl. Acad. Sci. U.

S. A. *112*, E1353–E1362.

Strauss, M., Sitt, J.D., Naccache, L., and Raimondo, F. (2022). Predicting the loss of responsiveness

when falling asleep in humans. Neuroimage *251*.

Vyazovskiy, V. V., Olcese, U., Cirelli, C., and Tononi, G. (2013). Prolonged wakefulness alters neuronal

responsiveness to local electrical stimulation of the neocortex in awake rats. J. Sleep Res. *22*, 239–

250.

Vyazovskiy, V. V, Olcese, U., Hanlon, E.C., Nir, Y., Cirelli, C., and Tononi, G. (2011). Local sleep in

awake rats. Nature *472*, 443–447.

Xu, Y., Wokke, M., Noreika, V., Bareham, C., Jagannathan, S., Georgieva, S., Trentin, C., and

Bekinschtein, T. (2023). Effects of alertness on perceptual detection and discrimination. BioRxiv

2023.03.21.533623.