Invariant sets for switched systems to tackle bacterial resistance

Alejandro Anderson[1], Matthew W. Kinahan[2], Alejandro H. Gonzalez[3],
Klas Udekwu[*2], and Esteban A. Hernandez-Vargas[†1]

[1]Department of Mathematics and Statistical Science, University of Idaho, USA
[2]Department of Biological Sciences, Bioinformatics and Computational Biology, University of Idaho, USA
[3]University of Littoral (UNL), Institute of Technological Development for the Chemical Industry (INTEC) and National Scientific and Technical Research Council (CONICET), Argentina

Abstract

The use of chemotherapeutics agents in the treatment of bacterial infections have resulted in the emergence of multidrug resistant pathogens. Clinically, with single and even multiple drug intervention strategies, pathogens have developed resistance to one or all drugs utilized. This leads to the reasonable conclusion that the primary effect of any (finite) amount of drugs is delaying resistance development as opposed to prevention. Importantly, it has been shown that in sequential exposure of pathogens to antibiotics, evolution of resistance to some drugs may increase sensitivity toward others previously used, a phenomenon known as collateral sensitivity. This suggests that multidrug resistance could be avoided by an adequate use of the available drugs. Without a framework to do this however, each bug:drugs interaction network would need to be assembled blindly, an arduous experimental process. This study develops a framework for describing qualitatively collateral sensitivity networks, accordingly the interactions between emerging drug-resistant variants are modeled and a dynamic analysis is conducted to predict failure or success of sequential drug therapies.

1   Introduction

Antimicrobial drug resistance (AMR) was uncovered in the 1950s and it is well known that the deployment of antimicrobial without prescription has increased the AMR crisis [15, 26, 38, 40, 45, 39]. AMR has mainly been addressed by the discovery of new classes of antibiotics but it was reported that the development of antimicrobials

---

[*]kudekwu@uidaho.edu
[†]esteban@uidaho.edu

might not keep up with the escalation of AMR in the next decades [19, 4, 12]. In addition, the combination of drugs necessary to treat AMR has triggered the emergence of multidrug-resistant (MDR) infections [29, 5, 51, 50] further aggravating the AMR crisis. Accordingly, control strategies that extend the life span of existing antibiotics and reduce the MDR are currently being studied [14, 47, 23, 8]. One approach has been focused on exploiting synergistic and antagonistic effects between antimicrobials [22], like orthogonal sensitivities that arise from the evolution to resistance which is known as collateral sensitivity (CS). CS was first described for Escherichia coli in 1952 [63] and it was observed in microbes infections and in cancer as well [57]. The CS phenomenon occurs when the genetic changes accrued on the develop of resistance towards one agent simultaneously decrease fitness to a second agent [34] and it is thought to be caused by a trade-off of pleiotropic resistance mutations [54]. More importantly, CS can be used to design drug combinations therapy: for instance by including mutual CS drugs in drug-cycling regimen [13, 28, 61]. To further understand the extent of CS, laboratory evolution has been deployed to map out CS profiles between the known antibiotics for specific drug-resistant strains [36, 53, 35, 27]; genome sequencing to characterize the components that contribute to CS [58, 41, 7, 20]; and the development of mathematical modeling that provides insight into the collateral dynamics network and supports control strategies to reduce the growth of MDR [37, 6].

Mathematical modeling establishes a high level of control involving factors such as pharmacodynamic (PD) and pharmacokinetic (PK) [65], optimal time of drug exposure [68], between others with relevance on the effects antibiotics have on bacterial growth, inhibition, killing, and mutation [60, 30, 9, 66]. Regarding collateral effects, mathematical models have been developed to evaluate *sequential drug regimens* in vitro and silico [69, 52], to assess the *robustness* of collateral sensitivity [52], to exploit its statistical structure and designing optimal policies [43], to evaluate possible *reversion of evolution towards resistance* [48] and to compare *cycling vs mixing* treatments [55]. None of these models however tackled the CS phenomenon with a combinatorial mutation network which has already been used for multidrug resistance (without collateral effects) on cancer [33], for which it was assumed that a mutation that confers resistance to one drug does not confer resistance to any of the other drugs in use[1]; framework that leads to a dead-end towards MDR. The epistatic interactions between genes involved in CS is a complex and not fully understood phenomenon but some magnitudes and directions affected by this phenomenon can be measured and framed within combinatorial mutation networks and - by dynamic network-based analysis [46] and the set-control theory [11] - we can determine drug combinations that contain the infection.

Eliminating an infection requires driving the system states to the healthy equilibrium at the origin. This is referred to as the regulation problem in control [17, 18, 70]. However, stabilizing the origin is not always feasible, especially in the context of evolutionary dynamics that lead to resistance. In such cases, more flexible control goals, such as invariant sets, become invaluable [1, 10]. Invariant sets, unlike fixed equilibria, enhance controllability by offering alternative stable regions and they provide safety zones to maintain infection suppression. Such invariant sets play a fundamental stabi-

---

[1]In the supplementary material of [33] it was noted that the combinatorial mutation network can be extended to include cross-resistance, even though this comment was not elaborated upon.

lization role as the only regions formally stabilizable under Lyapunov analysis [59]. In this study, we introduce a combinatorial mutation network that models qualitative collateral sensitivity data for the assessment of sequential drug regimens. We employed a switched system framework [42], extensively utilized in biomedical control problems [24, 62, 2] and recently adapted for sequential drug regimens with mutation dynamics [64], with the goal of targeting phenotype states associated with specific collateral effects to improve treatment success in chronic infections. The model suggests that collateral sensitivity profiling can forecast the emergence and proliferation of MDR strains. Furthermore, we explore invariant regions for resistance evolution population trajectories that determine the conditions under which mutual collateral sensitivity possesses the capacity to contain a chronic infection.

2 Evolving collateral effects in response to antibiotic exposure

The European Committee on Antimicrobial Susceptibility Testing (EUCAST) defines a microorganism as susceptible to a level of antimicrobial activity associated with a high likelihood of therapeutic success. Conversely, resistance is defined as a high likelihood of therapeutic failure. These categories are usually defined by breakpoints determined in standard phenotype test systems [31, 3]. Ideally, clinical breakpoints distinguish between patients who are likely or unlikely to respond to antimicrobial treatment.

For convenience, the minimum inhibitory concentration (MIC) can be used as a parameter of antibiotic action [60], since it was reported there is a linear impact between an increase in the MIC and hospital moralities [16]. It is defined as follows.

Definition 1 (Minimum Inhibitory Concentration - MIC). *The MIC breakpoint is defined as the lowest concentration of a drug that will inhibit the visible growth of a microorganism.*

Collateral sensitivity can be defined as a decrease in the MIC of the antibiotic-resistant strain relative to the wild-type. Conversely, an increase in the MIC relative to the wild-type defines a *collateral (or cross) resistance*. On the other hand microbiology laboratories uses clinical breakpoints to categorize microorganisms as clinically susceptible or resistant dependent on the quantitative antimicrobial susceptibility as indicated by the MIC value [32, 49]. Figure 1 illustrates the scenario for two hypothetical drugs A and B, for the case of combining antibiotics in sequential order (not for mixtures drugs). According to Figure 1 the MIC of the wild-type (Variant $V0$) indicates a profile of susceptibility to both drugs, after $V0$ is expose to drug A the MIC increase respect to A and respect to B, i.e. drug A shows cross resistance. On the other hand, after $V0$ is expose to drug B the MIC increase respect to B but it decrease respect to A, i.e. drug B shows collateral sensitivity. To determine the susceptible/resistant profile of emerging variants following drug exposure, collateral sensitivity effects can is formalized in next section using a combinatorial mutation network and, then linked the network with a switched system to model the population trajectories under drug sequential therapies.

Figure 1: Drug-driven vectors describe the direction the MIC evolves under drug exposure. The drug-driven vector of A (red) shows cross-resistance and the vector of B (blue) indicates collateral sensitivity. According to the position of the MIC and the breakpoints, the naive strain $V0$ is sensitive to both drugs, the variant $V1$ is resistant to both drugs and the variant $V2$ is resistant to drug B but sensitive to drug A.

3 Materials and methods

Consider N drugs $\{\sigma_1, \cdots, \sigma_N\} = \Sigma$. Σ defines a drug concentration space $\mathcal{C} \subseteq \mathbb{R}^N_{\geq 0}$, where $c = (c_1, \cdots, c_N) \in \mathcal{C}$ is such that c_i represents concentration of drug σ_i for all $i = 1, \cdots, N$ (see Figure 1 for 2 drugs $\Sigma = \{A, B\}$). We define a MIC N-dimensional vector that quantifies collateral sensitivity and cross-resistance of all drugs on Σ for a particular microorganism V, $MIC(V) \in \mathcal{C}$, as follows

$$MIC(V) = (MIC_{\sigma_1}(V), \cdots, MIC_{\sigma_N}(V)) \in \mathcal{C},$$

where $MIC_{\sigma_i}(V)$ is the MIC of drug σ_i for microorganism V. The breakpoints of Σ, $Br_\Sigma \in \mathcal{C}$, is given by

$$Br_\Sigma = (Br_{\sigma_1}, \cdots, Br_{\sigma_N}) \in \mathcal{C},$$

where Br_{σ_i} represents a maximum concentration of drug σ_i allowed to used for all $i = 1, \cdots, N$. Then, the breakpoints and the MIC of Σ state the following.

Definition 2. *[Sensitive/Resistant strains] Consider a microorganism V and N drugs $\Sigma = \{\sigma_1, \cdots, \sigma_N\}$. It is said that V is sensitive/resistant to drug σ_i, with $i = 1, \cdots, N$, if the i^{th} element of the vector $MIC(V) - Br_\Sigma$, is negative/non-negative.*

The phenomenon of a microorganism V_i exposed over time to drug σ until it converges to a resistant variant V_j (a different phenotypic state) can be represented by the following graph

$$V_i \xrightarrow{\sigma} V_j,$$

which can be measured by the following *drug-driven vector*.

Definition 3 (Drug-driven vector). *Consider the following graph $V_i \xrightarrow{\sigma} V_j$. The drug-driven vector, $\vec{v}_\sigma \in \mathbb{R}^N$, is defined by*

$$\vec{v}_\sigma := MIC(V_j) - MIC(V_i).$$

Negative elements of $\vec{v}_\sigma$ represent collateral sensitivity, unintendedly produced by drug σ; positive elements represent cross-resistance, unintendedly produced by drug σ, and null elements nonchange in relation to the MIC of the naive variant V_i. In addition, considering the breakpoints of Σ and Definition 2, the vector $\vec{v}_\sigma$ determines the susceptible/resistant profile of all emerging variants.

3.1 State-space modeling

To obtain a formal dynamical model describing the mutation/variation path of a pathogen when exposed to a sequence of N possible drugs (i.e., when applying one drug at a time), we associate each possible combination of sensible/resistant variant to a state in a state-space. This way, the potential dimension of the state space is 2^N, accounting for the fact that each variant can be either sensitive or resistant to each of the drugs. Formally, an state x_i, corresponding to variant V_i, for some $i \in \{1, \cdots, 2^N\}$, is given (for instance) by $x_i := (\sigma_{1,S}, \sigma_{2,R}, \cdots, \sigma_{N,S})$, meaning that variant V_i is susceptible to drug σ_1, resistant to drug σ_2, $\cdots$, and susceptible to drug σ_N or, the same, $MIC(V_i) := (MIC_{\sigma_1}(V_i), MIC_{\sigma_2}(V_i), \cdots, MIC_{\sigma_N}(V_i))$ fulfill the conditions

$$
\begin{aligned}
MIC_{\sigma_1}(V_i) &< Br_{\sigma_1}, \\
MIC_{\sigma_2}(V_i) &\geq Br_{\sigma_2}, \\
&\cdots, \\
MIC_{\sigma_N}(V_i) &< Br_{\sigma_N}.
\end{aligned}
$$

We denote the complete model containing the 2^N states as the classical one.

However, if the model is intended to represent the system behavior just for realistic initial conditions - i.e., initial conditions that can be reached starting from other realistic initial conditions and feasible paths - the dimension (and complexity) of the model can be significantly reduced. Suppose that the original complete system of dimension 2^N is divided into two subspaces of dimension L and E, respectively (L denotes "latent" and E denotes 'explicit' subspace), such that $L + E = 2^N$. And suppose also that the L-dimensional subspace cannot be reached, by any drug sequence, from its complementary subspace of dimension E, containing this latter all the realistic initial conditions. This way, the E-dimensional subspace is an invariant subspace containing all the initial biologically feasible conditions: L-dimensional subspace can not be the result of a mutation path promoted by the collateral sensitivity and cross-resistance profiles of the chosen drugs.

To exemplify this point, let us consider the hypothetical case of a pathogen and two drugs described in Figure 1, where drug A shows cross-resistance with respect to drug B and drug B presents collateral sensitivity with respect to drug A. According to this simple scheme, the complete 2^N states are denoted as $A_S B_S$, $A_R B_S$, $A_s B_R$ and $A_R B_R$, denoting all possible variant combinations according to their sensitivity/resistance to each drug. However, starting from the (wild type) variant V_0, which

is a $A_S B_S$ state, and a biologically feasible initial condition, only variants V_1, which is $A_S B_R$, and variant V_2, which is $A_R B_R$, can be reached by applying sequentially drugs A and B (in any order and during any time period). State variant $A_R B_S$ constitutes in this example the L-dimensional (with $L = 1$) latent subspace to which the system cannot evolve, while the complementary 3-dimensional subspace is the invariant subspace. Figure 2 shows a graph $V_0 \xrightarrow{\sigma} V_i$ for every $\sigma \in \Sigma := \{A, B\}$ and $i = 1, 2$, of this example. Our main hypothesis supporting this approach is that, given that the collateral sensitivity and cross-resistance profiles of the drugs are considered fixed for the model, they necessarily determine its structure (its dimension and potential behavior).

The general way to obtain the reduced model is by applying each drug at a time to each state that emerges from the previous application of each drug. The procedure is stopped when the network converges to a closed graph, as shown in the right plot of Figure 2.

Figure 2: Classic combinatorial mutation network describing resistant and sensitive states for drugs A and B resulting in a total of 2^N states (left) vs a combinatorial mutation network accounting for collateral sensitivity and cross-resistance of drugs A and B (right).

A network with n sates $(x_1, x_2, \cdots, x_n)$ can be linked to a switched system, as demonstrated in a prior study [64] and by this we can investigate the impact of sequential drug exposures on population dynamics. The effect of therapy is modeled by drug-induced death rate framework, i.e., the growth rate for variant x_i under exposure of drug σ is given by $\alpha_i^\sigma > 0$ with a carry capacity of $K_i^\sigma > 0$; death and mutation from x_i to x_j rates are given by δ_i^σ and μ_{ij}^σ, respectively. The combinatorial mutation network is given by the binary matrix $m_{i,j}^\sigma \in \{0, 1\}$, where $m_{i,j}^\sigma = 1$ implies an arrow from x_i to x_j, and $m_{i,j}^\sigma = 0$ there is no connection. For a discretization of time $k \geq 0$, the following switched system describes the nonlinear dynamics between states.

$$x_i(k+1) = \alpha_i^{\sigma(k)} x_i(k) \left(1 - \frac{x_i(k)}{K_i^{\sigma(k)}}\right) - \delta_i^{\sigma(k)} x_i(k) + \mu_{ij}^{\sigma(k)} \sum_{j=1}^n m_{i,j}^{\sigma(k)} x_j(k). \quad (1)$$

The states of the system are given by $x = (x_1, x_2, \cdots, x_n)$, and the control input by σ, for all time $k > 0$.

Remark 1. *For switched systems, mode selection from the finite set can be formulated as a dynamic programming problem, but the solution is difficult to implement and may not account for constraints [25, 24]. More applicable control approaches involve*

mixed-integer optimization [67], state-dependent switching rules [21], and receding horizon switching signals [2].

The switch of drugs shifts the balance of birth and death such that the colony shrinks or escapes. It is assumed that during single-drug therapy with, say, drug σ_d, mutations lead to the emergence of a variant resistant to this drug, i.e., state $\sigma_{1,?}, \cdots, \sigma_{d,S}, \cdots, \sigma_{1,?}$ varies/mutates to state $\sigma_{1,?}, \cdots, \sigma_{d,R}, \cdots, \sigma_{1,?}$ whatever ? is S or R. Consequently, the resistant phenotype proliferates while the sensitive strains will be killed at a constant rate, ultimately culminating in treatment failure. Also, to simplify the model, all sensitive cells are killed at the same rate as the naive strain.

Remark 2. *The dynamics framework we consider here is that under the pressure of any drug there is an evolution towards resistance to the drug. Thus monotherapy eventually fails to contain overall infection levels leading to treatment failure as resistant strains dominate. Within this framework, combination drug regimens are required to prevent uncontrolled spread of the infection. Our goal is to determine optimal switching protocols that constrain the infection despite ongoing evolution of drug resistance.*

3.2   Invariance analysis to contain drug-resistant infection

In the framework proposed by Eq. 1 it is required to use combinations of agents with mutual collateral sensitivity to prevent population trajectory escape, because of the constant evolution towards resistance. However, this requirement is not a sufficient condition to ensure that population trajectories will not escape. To see this, consider Figure 3, where three illustrative dynamics show the effects of drug A (red) and drug B (blue) with mutual collateral sensitivities on the two populations $A_R B_S$ and $A_R B_S$. Depending on the birth, death, and mutation rates, (a) the behavior of the population can achieve perfect balance and the total population can be maintained inside a target window; (b) the population can experience shrinkage so it will remain inside the target window; and (c) the escape of the population, because it is not possible to keep the states inside the target region.

The assessment of this problem has been successfully tackled by set-control theory [10]. Set-control involves examining regions within the state space that feasibly bound system trajectories, such regions are known as invariant sets. Examining invariant sets for our dynamical system can help to determine if a trajectory will remain bounded or escape in this framework.

Let us assume there is a therapeutic window $\mathbb{T}$ on the states space, where we want to maintain the population of microorganisms, of course we assume that $0 \in \mathbb{T}$. A control invariant set (CIS) inside $\mathbb{T}$, for the switched system on Eq. 1 can be defined as follows.

Definition 4 (Control Invariant Set - CIS)**.** *A set $\Omega \subset \mathbb{T}$ on the states space is said to be a control invariant set of the switched system (1) if for every state $x(k) \in \Omega$ there is a feasible mode $\sigma \in \Sigma$ such that the next state follows the condition $x(k+1) \in \Omega$.*

The invariant set is a generalization of the steady-state concept and its characterization implies there is a feasible sequence of drugs that keep the population bounded

Figure 3: **Illustrative dynamic evolution of mutual collateral sensitivity**. Exposure to drug A reduces concentration of $A_S B_R$ cells (sensitive to A and resistant to B) but allows the escape of $A_R B_S$. Oppositely, exposure to drug B reduces concentration of $A_R B_S$ but induces the escape of $A_S B_R$. a) A proper balance between dynamic of drugs A and B implies that the total population can be feasible bounded. b) The best scenario of mutual collateral sensitivity implies the asymptotic stability of the origin. c) Despite a mutual collateral sensitivity it may not be feasible to prevent the escape of the infection.

inside Ω obviating the need for identifying the specific drug sequence, its order and exposure times, which is related with a complex and challenging optimal control problem. Instead, by simply ensuring the existence of a CIS, we can establish that the infection can be feasibly containing. Certainly, the presence of a multidrug-resistant (MDR) strain within the dynamical network implies the nonexistence of an invariant set within a therapeutic window.

To characterize a CIS, the concept of controllable set can be used [2]. The controllable set of Ω is given by all the states that can be driven by a feasible mode $\sigma \in \Sigma$ in one step to Ω, and it can be formally defined by the following.

$$\mathcal{S}(\Omega) := \{x(k) \in \mathbb{R}^n_{\geq 0} : \exists\, \sigma \in \Sigma \text{ s.t. } x(k+1) \in \Omega\}.$$

Next result is well established on set-control theory.

Property 1. *If $\Omega \subseteq \mathcal{S}(\Omega)$ then Ω is a CIS.*

Proof. Suppose that $x(k) \in \Omega$ for some time-step $k \geq 0$. Since $\Omega \subseteq \mathcal{S}(\Omega)$, then $x(k) \in \mathcal{S}(\Omega)$ as well. By definition of the controllable set $\mathcal{S}(\Omega)$ there exists $\sigma \in \Sigma$ such that $x(k+1) \in \Omega$, which conclude the proof. $\qquad\square$

Various methods for characterizing invariant sets for a linear switched system were proposed in [56]. Based on the ideas presented there, we have developed a method for characterizing CIS in our framework, as detailed in the supplementary material.

4 Results

We developed a model framework to consider the collateral effects of antibiotics listed in Table A1 against the resistance evolution of P. aeruginosa. Bacterial population dynamics include distinct phenotype states, the number and nature of these states are defined by the collateral sensitivity and cross-resistance profile of the antibiotics selected for the treatment. The states consist of subpopulation groups that respond differently to various drugs, exhibiting mutations that lead to resistance to the drug in use, as well as collateral effects on other drugs. The framework was then used to explore how collateral sensitivity can help reduce the development of resistance in a population by cyclic treatments. Moreover, we studied the scope of orthogonal sensitivities needed to prevent the infection from spreading due to generated resistance using set-control theory; by studying specific areas in the population space - called invariant sets - where the behavior of all subpopulations can be controlled. We assessed methods to find bounded invariant sets for the switched system associated with different treatments, and we demonstrate that the existence of this set implies the infection can be suppressed (or at least contained) through a feasible sequence of drugs.

4.1 Sequential drug therapy simulation

We have simulated a combinatorial mutation network to describe the interactions between strains resistant and sensitive to 3 drugs. We compare a cyclic treatment for two types of networks. The first is a conventional network with 3 generic drugs A, B and C; in this case there are 2^3 different phenotypic states (as described in Figure 4-a), the drug-driven resistance leads to mutations that follow a dead-end path to the multidrug resistance strain $A_R B_R C_R$, which is inevitable. The second network is designed according methodology proposed on Section 3.1, for the antibiotics Aztreonam (AZE), Ciprofloxacin (CIP), Tobramycin (TOB) from Table A1, with $A = AZE$, $B = CIP$, and $C = TOB$. The collateral sensitivity and cross-resistance profiles of these drugs according Figure A1 (generated in [27]) produce a particular network, where the directions of mutations take into account all the side effects. Figure 4-b shows the network for these drugs, the red arrows indicate that there is cross-resistance and the blue arrows indicate that there is collateral sensitivity. As there are 3 drugs, one drug can exhibit cross-resistance concerning one drug and collateral sensitivity concerning the other; in these cases the black arrow is used. Both networks were linked to the dynamics model in Equation 1. This allowed simulating bacterial growth under $50hr$ cyclic dosing of drugs A, B, C over $600hr$ (as described in Figure 4-c). Importantly, Figure 4-b shows that drug interactions alter potential mutation pathways, so not all theoretically possible phenotype strains manifest in the network. The bacteria cannot mutate in a way that becomes fully resistant to all drugs, what cannot be avoided in the classic network presented in Figure 4-a. This is a crucial point as it enables us to select the right drugs to prevent the emergence of multiresistant strains.

Figure 4: Dynamic evolution of a classic mutation network vs a collateral sensitivity network simulations for three drugs with cyclic treatment. We consider two cases: a) for a combinatorial mutation network given by generic drugs without collateral effects (a). And b) for a combinatorial mutation network that considers collateral effects between drugs Aztreonam ($A = AZE$), Ciprofloxacin ($B = CIP$), Tobramycin ($C = TOB$).

4.2 Invariant set for Levofloxacin and Ampicillin

The antibiotics Levofloxacin and Ampicillin show mutual collateral sensitivity when used in PAO1 wild type [27], thus providing an ideal scenario to test whether the model generated by these two antibiotics admit invariants within a therapeutic window. The mutual collateral sensitivity of the drugs LEV and AMP generates the combinatorial mutation network shown in Figure 5-a, with the states $A_S B_S$ (sensitive to both drugs), and the states $A_S B_R$ and $A_R B_S$. The network indicates the direction of mutations under the effect of each drug. Note that the system has 3 states and one of them is sensitive to both drugs (wild type $A_S B_S$), therefore this state converges to the origin when either drug A or drug B is applied. Thus, we can disregard this dimension, and

represent the invariant sets in 2 dimensions; $A_S B_R$ and $A_R B_S$. In this way, we have better visibility of the sets. To calculate a control invariant for the system associated

Figure 5: Invariant set for combination of drugs Levofloxacin (LEV) and Ampicillin (AMP). From the profiles of collateral sensitivity and cross resistance of drugs LEV and AMP the network describing the interactions between variants $LEV_S AMP_S$, $LEV_R AMPC_S$ and $LEV_S AMPC_R$ is computed. There exists invariant sets of the mathematical model associated to the network which implies the infection can be bounded with drugs LEV and AMP. Invariant Set Ω (green) inside T. Note that the controllable set $\mathcal{S}(\Omega)$ (dashed-lines) contains the invariant set for all cases.

with the network, we use Algorithm 1 proposed in the Supplementary Material for the linearized model. Four cases were simulated, in which different growth, death, and mutation rates were considered (see Table 2). For cases one (Figure 5-b), two (Figure 5-c), and four (Figure 5-e) we chose a therapeutic window given by the set $\mathbb{T}_1 = \{0 \leq A_R B_S \leq 1, 0 \leq A_S B_R \leq 1\}$, and the algorithm finds an invariant within $\mathbb{T}_1$, as Figure 5 shows the controllable sets contains set Ω in each case, fulfilling Property 1. For case 3 (Figure 5-d) the algorithm does not find an invariant within $\mathbb{T}_1$, but it converges for $\mathbb{T}_2 = \{0 \leq A_R B_S \leq 1, 0 \leq A_S B_R \leq 0.5\}$.

4.3 Invariant set for Tobramycin, Carbenicillin and Colistin

Here we consider 3 drugs; Tobramycin, Carbenicillin, and Colistin, whose collateral effects were analyzed in [27] for the treatment of the PAO1 wild-type strain. According to Imamovic et. at. these drugs exhibit collateral sensitivity and cross-resistance both. The combinatorial mutation network for these drugs is shown in Figure 6-a, with 4 states: the wild type $A_S B_S C_S$ sensitive to all drugs ($A = TOB$, $B = CAR$, $C = COL$), and three other states that mutate among themselves according the collateral effects of the drug used, $A_S B_R C_S$, $A_R B_S C_S$ and $A_R B_S C_R$.

Through Algorithm 1 we proved that there exists an invariant for the linear dynamic system associated with this set of drugs, which is plotted in Figure 6-b in three dimensions (the dimension of $A_S B_S C_S$ is not considered because this state always decays), the parameters from Table 3 were used.

Figure 6: Normalized cell population of variants $A_R B_S C_R$, $A_R B_S C_S$ and $A_S B_R C_S$ inside the invariant set Ω for drugs Tobramycin (TOB), Carbenicillin (CAR) and Colistin (COL).

The invariant set Ω found and represented in Figure 6-b implies that the infection can be feasibly contained through some sequence given by the drugs TOB, CAR and COL. This means that even though there is constant evolution towards resistance, the dynamics and interaction between these drugs are enough to contain the infection.

Technically, if a state,

$$x(k) = (A_S B_R C_S(k), A_R B_S C_S(k), A_R B_S C_R(k)) \in \Omega,$$

representing population at time $k \geq 0$, one drug in $\Sigma = \{TOB, CAR, COL\}$ implies that

$$x(k+1) = (A_S B_R C_S(k+1), A_R B_S C_S(k+1), A_R B_S C_R(k+1)) \in \Omega,$$

and so on indefinitely.

5   Discussion

Our finding shows that a sequential drug treatment that aims to eliminate a population of distinct bacterial phenotype all evolving resistance to the current drug requires drug synergy. Without collateral sensitivity between agents, single or combined antibiotics are bound to fail by enabling persistence of at least one resistant bacterial variant. In this current study, we have used a birth/death model that captures interactions between distinct variants of bacteria using the collateral sensitivity and cross-resistance profile of the drugs targeting specific species. The interactions were modeled by combinatorial mutation networks where the mutation rates between the variants can be specified. We modeled the drug effect by discrete events affecting the birth, death, and mutation rates of a continuous bacterial population. Systems containing both continuous and discrete-time dynamics and characterized by abrupt changes in at least one state variable at certain time instants, are called hybrid systems [44] and can be used to analyze qualitative properties of the population evolution.

Our model enables predicting the effect of sequential drug treatments, taking into account the antagonistic and synergistic interactions between the drugs involved in the process. This proposed model can be used to study advanced control strategies and determine optimal combinations and sequences of drugs. It can also be used to determine which combinations lead to the proliferation of multidrug-resistant strains, as well as to analyze the trade-offs between the ongoing evolution of resistance to the current drug and the evolution toward sensitivity to potential drugs. In the latter, we focused some of our dynamic analysis. To this end, we employed set-control theory to characterize stable regions within the feasible state space. These regions could include steady states, multiple equilibria, or invariant sets. However, our system had only healthy equilibria (e.g., the origin) with no relevance in this context. Therefore, we proposed to use the concept of control invariant sets to assess when the total population could be feasibly maintained within a therapeutic window, as an invariant set captured the regions of the state space where the system can feasibly remain indeterminately. We provided a method to characterize invariant sets for our particular system and computed several invariant sets for a population of distinct phenotypes of P. aeruginosa and its sensitivity profile for 24 relevant antibiotics [27]. In this way, the existence of a nonempty invariant set within a therapeutic window for a specific system (determined by the drugs chosen for therapy) ensures that drug resistance could be feasibly contained. Furthermore, the nonexistance of such a set implies that the population can not

be feasibly contained inside the therapeutic window, resulting in resistance prevailing over synergy.

Our dynamic analyses offer valuable qualitative and quantitative insights into the collateral effects on population dynamics. We can reliably identify drug combinations leading to treatment failure and those enabling the controlled dynamics of all subpopulations, depending on specific birth, death, and mutation rates. To expand our findings, we must incorporate PD/PK factors, account for stochastic processes, address resource competition, and delve into parameter estimation. Although these additional factors introduce a layer of complexity to the dynamic analysis, also would allow us to effectively leverage our results in modeling in-vitro population growth, and to ensure the effectiveness of combining pharmacology and evolution for sustaining pathogen suppression.

Acknowledgment of Support and Disclaimer

This material is based upon work supported by the National Science Foundation under Grant No. 2315862.

References

[1] A. Anderson, P. Abuin, A. Ferramosca, E. Hernandez-Vargas, and A. Gonzalez. Cyclic control equilibria for switched systems with applications to ecological systems. *International Journal of Robust and Nonlinear Control*, 2021.

[2] Alejandro Anderson, Alejandro H Gonzalez, Antonio Ferramosca, and Esteban A Hernandez-Vargas. Discrete-time MPC for switched systems with applications to biomedical problems. *Communications in Nonlinear Science and Numerical Simulation*, 95:105586, 2021.

[3] Jennifer M Andrews. Determination of minimum inhibitory concentrations. *Journal of antimicrobial Chemotherapy*, 48(suppl_1):5–16, 2001.

[4] Giannamaria Annunziato. Strategies to overcome antimicrobial resistance (amr) making use of non-essential target inhibitors: A review. *International journal of molecular sciences*, 20(23):5844, 2019.

[5] Cesar A Arias, Diana Panesso, Danielle M McGrath, Xiang Qin, Maria F Mojica, Corwin Miller, Lorena Diaz, Truc T Tran, Sandra Rincon, E Magda Barbu, et al. Genetic basis for in vivo daptomycin resistance in enterococci. *New England Journal of Medicine*, 365(10):892–900, 2011.

[6] Linda BS Aulin, Apostolos Liakopoulos, Piet H van der Graaf, Daniel E Rozen, and JG Coen van Hasselt. Design principles of collateral sensitivity-based dosing strategies. *Nature Communications*, 12(1):5691, 2021.

[7] Camilo Barbosa, Roderich Römhild, Philip Rosenstiel, and Hinrich Schulenburg. Evolutionary stability of collateral sensitivity to antibiotics in the model pathogen pseudomonas aeruginosa. *Elife*, 8:e51481, 2019.

[8] Michael Baym, Laura K Stone, and Roy Kishony. Multidrug evolutionary strategies to reverse antibiotic resistance. *Science*, 351(6268):aad3292, 2016.

[9] Anna Camilla Birkegård, Tariq Halasa, Nils Toft, Anders Folkesson, and Kaare Græsbøll. Send more data: a systematic review of mathematical models of antimicrobial resistance. *Antimicrobial Resistance & Infection Control*, 7(1):1–12, 2018.

[10] F. Blanchini and S. Miani. *Set-Theoretic Methods in Control*. Systems & Control: Foundations & Applications. Springer International Publishing, 2015.

[11] Franco Blanchini and Stefano Miani. *Set-theoretic methods in control*, volume 78. Springer, 2008.

[12] Tobias Bollenbach. Antimicrobial interactions: mechanisms and implications for drug discovery and resistance evolution. *Current opinion in microbiology*, 27:1–9, 2015.

[13] Erwin M Brown and Dilip Nathwani. Antibiotic cycling or rotation: a systematic review of the evidence of efficacy. *Journal of antimicrobial chemotherapy*, 55(1):6–9, 2005.

[14] Remy Chait, Allison Craney, and Roy Kishony. Antibiotic interactions that select against resistance. *Nature*, 446(7136):668–671, 2007.

[15] John Crofton and DA Mitchison. Streptomycin resistance in pulmonary tuberculosis. *British medical journal*, 2(4588):1009, 1948.

[16] John S Esterly, Jamie Wagner, Milena M McLaughlin, Michael J Postelnick, Chao Qi, and Marc H Scheetz. Evaluation of clinical outcomes in patients with bloodstream infections due to gram-negative bacteria according to carbapenem mic stratification. *Antimicrobial agents and chemotherapy*, 56(9):4885–4890, 2012.

[17] Mirko Fiacchini and Marc Jungers. Necessary and sufficient condition for stabilizability of discrete-time linear switched systems: A set-theory approach. *Automatica*, 50:75–83, 01 2014.

[18] Jose C Geromel and Patrizio Colaneri. Stability and stabilization of continuous-time switched linear systems. *SIAM Journal on Control and Optimization*, 45(5):1915–1930, 2006.

[19] Howard S Gold and Robert C Moellering Jr. Antimicrobial-drug resistance. *New England journal of medicine*, 335(19):1445–1453, 1996.

[20] Patrick R Gonzales, Mitchell W Pesesky, Renee Bouley, Anna Ballard, Brent A Biddy, Mark A Suckow, William R Wolter, Valerie A Schroeder, Carey-Ann D Burnham, Shahriar Mobashery, et al. Synergistic, collaterally sensitive β-lactam combinations suppress resistance in mrsa. *Nature chemical biology*, 11(11):855–861, 2015.

[21] Daniel Görges, Michal Izák, and Steven Liu. Optimal control and scheduling of switched systems. *IEEE Transactions on Automatic Control*, 56(1):135–140, 2010.

[22] Matthew D Hall, Misty D Handley, and Michael M Gottesman. Is resistance useless? multidrug resistance and collateral sensitivity. *Trends in pharmacological sciences*, 30(10):546–556, 2009.

[23] Robert EW Hancock. Collateral damage. *Nature Biotechnology*, 32(1):66–68, 2014.

[24] Esteban Hernandez-Vargas, Patrizio Colaneri, Richard Middleton, and Franco Blanchini. Discrete-time control for switched positive systems with application to mitigating viral escape. *International journal of robust and nonlinear control*, 21(10):1093–1111, 2011.

[25] Esteban A. Hernandez-Vargas, Patrizio Colaneri, and R.H. Middleton. Switching strategies to mitigate HIV mutation. *Control Systems Technology, IEEE Transactions on*, 22:1623–1628, 07 2014.

[26] King K Holmes, David W Johnson, and Thomas M Floyd. Studies of venereal disease: I. probenecid-procaine penicillin g combination and tetracycline hydrochloride in the treatment of penicillin-resistant gonorrhea in men. *JAMA*, 202(6):461–466, 1967.

[27] Lejla Imamovic, Mostafa Mostafa Hashim Ellabaan, Ana Manuel Dantas Machado, Linda Citterio, Tune Wulff, Soren Molin, Helle Krogh Johansen, and Morten Otto Alexander Sommer. Drug-driven phenotypic convergence supports rational treatment strategies of chronic infections. *Cell*, 172(1-2):121–134, 2018.

[28] Lejla Imamovic and Morten OA Sommer. Use of collateral sensitivity networks to design drug cycling protocols that avoid resistance development. *Science translational medicine*, 5(204):204ra132–204ra132, 2013.

[29] Michael D Iseman. Treatment of multidrug-resistant tuberculosis. *New England Journal of Medicine*, 329(11):784–791, 1993.

[30] Alfonsas Juška, Genovaitė Gedminienė, and Rūta Ivanec. Growth of microbial populations. mathematical modeling, laboratory exercises, and model-based data analysis. *Biochemistry and Molecular Biology Education*, 34(6):417–422, 2006.

[31] G Kahlmeter, DFJ Brown, FW Goldstein, AP MacGowan, JW Mouton, Inga Odenholt, A Rodloff, C-J Soussy, M Steinbakk, F Soriano, et al. European committee on antimicrobial susceptibility testing (eucast) technical notes on antimicrobial susceptibility testing, 2006.

[32] Gunnar Kahlmeter, Derek FJ Brown, Fred W Goldstein, Alasdair P MacGowan, Johan W Mouton, Anders Österlund, Arne Rodloff, Martin Steinbakk, Pavla Urbaskova, and Alkiviadis Vatopoulos. European harmonization of mic breakpoints for antimicrobial susceptibility testing of bacteria. *Journal of antimicrobial chemotherapy*, 52(2):145–148, 2003.

[33] Natalia L Komarova and Dominik Wodarz. Drug resistance in cancer: principles of emergence and prevention. *Proceedings of the National Academy of Sciences*, 102(27):9714–9719, 2005.

[34] Viktória Lázár, Ana Martins, Réka Spohn, Lejla Daruka, Gábor Grézal, Gergely Fekete, Mónika Számel, Pramod K Jangir, Bálint Kintses, Bálint Csörgő, et al. Antibiotic-resistant bacteria show widespread collateral sensitivity to antimicrobial peptides. *Nature microbiology*, 3(6):718–731, 2018.

[35] Viktória Lázár, István Nagy, Réka Spohn, Bálint Csörgő, Ádám Györkei, Ákos Nyerges, Balázs Horváth, Andrea Vörös, Róbert Busa-Fekete, Mónika Hrtyan, et al. Genome-wide analysis captures the determinants of the antibiotic cross-resistance interaction network. *Nature communications*, 5(1):4352, 2014.

[36] Viktória Lázár, Gajinder Pal Singh, Réka Spohn, István Nagy, Balázs Horváth, Mónika Hrtyan, Róbert Busa-Fekete, Balázs Bogos, Orsolya Méhi, Bálint Csörgő, et al. Bacterial evolution of antibiotic hypersensitivity. *Molecular systems biology*, 9(1):700, 2013.

[37] Bruce R Levin and Klas I Udekwu. Population dynamics of antibiotic treatment: a mathematical model and hypotheses for time-kill and continuous-culture experiments. *Antimicrobial agents and chemotherapy*, 54(8):3414–3426, 2010.

[38] Stuart B Levy. The challenge of antibiotic resistance. *Scientific American*, 278(3):46–53, 1998.

[39] Stuart B Levy and Bonnie Marshall. Antibacterial resistance worldwide: causes, challenges and responses. *Nature medicine*, 10(Suppl 12):S122–S129, 2004.

[40] StuartB Levy. Microbial resistance to antibiotics: an evolving and persistent problem. *The Lancet*, 320(8289):83–88, 1982.

[41] Apostolos Liakopoulos, Linda BS Aulin, Matteo Buffoni, Efthymia Fragkiskou, JG Coen van Hasselt, and Daniel E Rozen. Allele-specific collateral and fitness effects determine the dynamics of fluoroquinolone resistance evolution. *Proceedings of the National Academy of Sciences*, 119(18):e2121768119, 2022.

[42] Daniel Liberzon. *Switching in systems and control*. Springer Science & Business Media, 2003.

[43] Jeff Maltas and Kevin B Wood. Pervasive and diverse collateral sensitivity profiles inform optimal strategies to limit antibiotic resistance. *PLoS biology*, 17(10):e3000515, 2019.

[44] Alexey S Matveev and Andrey V Savkin. *Qualitative theory of hybrid dynamical systems*. Springer Science & Business Media, 2012.

[45] Douglas L Mayers, Jack D Sobel, Marc Ouellette, Keith S Kaye, and Dror Marchaim. *Antimicrobial Drug Resistance: Clinical and Epidemiological Aspects, Volume 2*, volume 2. Springer, 2017.

[46] Chandrakala Meena, Chittaranjan Hens, Suman Acharyya, Simcha Haber, Stefano Boccaletti, and Baruch Barzel. Emergent stability in complex network dynamics. *Nature Physics*, pages 1–10, 2023.

[47] Jean-Baptiste Michel, Pamela J Yeh, Remy Chait, Robert C Moellering Jr, and Roy Kishony. Drug interactions modulate the potential for evolution of resistance. *Proceedings of the National Academy of Sciences*, 105(39):14918–14923, 2008.

[48] Portia M Mira, Kristina Crona, Devin Greene, Juan C Meza, Bernd Sturmfels, and Miriam Barlow. Rational design of antibiotic treatment plans: a treatment strategy for managing evolution and reversing resistance. *PloS one*, 10(5):e0122283, 2015.

[49] Johan Willem Mouton, DFJ Brown, P Apfalter, R Cantón, CG Giske, M Ivanova, AP MacGowan, A Rodloff, C-J Soussy, M Steinbakk, et al. The role of pharmacokinetics/pharmacodynamics in setting clinical mic breakpoints: the eucast approach. *Clinical Microbiology and Infection*, 18(3):E37–E45, 2012.

[50] Daniel M Musher, Mark E Dowell, Virginia D Shortridge, Robert K Flamm, James H Jorgensen, Pierre Le Magueres, and Kurt L Krause. Emergence of macrolide resistance during treatment of pneumococcal pneumonia. *New England Journal of Medicine*, 346(8):630–631, 2002.

[51] Michael M Mwangi, Shang Wei Wu, Yanjiao Zhou, Krzysztof Sieradzki, Herminia de Lencastre, Paul Richardson, David Bruce, Edward Rubin, Eugene Myers, Eric D Siggia, et al. Tracking the in vivo evolution of multidrug resistance in staphylococcus aureus by whole-genome sequencing. *Proceedings of the National Academy of Sciences*, 104(22):9451–9456, 2007.

[52] Daniel Nichol, Joseph Rutter, Christopher Bryant, Andrea M Hujer, Sai Lek, Mark D Adams, Peter Jeavons, Alexander RA Anderson, Robert A Bonomo, and Jacob G Scott. Antibiotic collateral sensitivity is contingent on the repeatability of evolution. *Nature communications*, 10(1):334, 2019.

[53] Tugce Oz, Aysegul Guvenek, Sadik Yildiz, Enes Karaboga, Yusuf Talha Tamer, Nirva Mumcuyan, Vedat Burak Ozan, Gizem Hazal Senturk, Murat Cokol, Pamela Yeh, et al. Strength of selection pressure is an important parameter contributing to the complexity of antibiotic resistance evolution. *Molecular biology and evolution*, 31(9):2387–2401, 2014.

[54] Csaba Pál, Balázs Papp, and Viktória Lázár. Collateral sensitivity of antibiotic-resistant microbes. *Trends in microbiology*, 23(7):401–407, 2015.

[55] Rafael Peña-Miller and Robert Beardmore. Antibiotic cycling versus mixing: the difficulty of using mathematical models to definitively quantify their relative merits. 2010.

[56] Mara Perez, Alejandro Anderson, Pablo Abuin, Esteban A Hernandez-Vargas, Alejandro H Gonzalez, and Marcelo Actis. Computing control invariant sets for waiting-time switched systems: A study case of glucose regulation. *IEEE Control Systems Letters*, 2023.

[57] Kristen M Pluchino, Matthew D Hall, Andrew S Goldsborough, Richard Callaghan, and Michael M Gottesman. Collateral sensitivity as a strategy against cancer multidrug resistance. *Drug Resistance Updates*, 15(1-2):98–105, 2012.

[58] Nicole L Podnecky, Elizabeth GA Fredheim, Julia Kloos, Vidar Sørum, Raul Primicerio, Adam P Roberts, Daniel E Rozen, Ørjan Samuelsen, and Pål J Johnsen. Conserved collateral antibiotic susceptibility networks in diverse clinical strains of escherichia coli. *Nature Communications*, 9(1):3673, 2018.

[59] James Blake Rawlings, David Q Mayne, and Moritz Diehl. *Model predictive control: theory, computation, and design*, volume 2. Nob Hill Publishing Madison, WI, 2017.

[60] Roland R Regoes, Camilla Wiuff, Renata M Zappala, Kim N Garner, Fernando Baquero, and Bruce R Levin. Pharmacodynamic functions: a multiparameter approach to the design of antibiotic treatment regimens. *Antimicrobial agents and chemotherapy*, 48(10):3670–3676, 2004.

[61] Carola EH Rosenkilde, Christian Munck, Andreas Porse, Marius Linkevicius, Dan I Andersson, and Morten OA Sommer. Collateral sensitivity constrains resistance evolution of the ctx-m-15 β-lactamase. *Nature communications*, 10(1):618, 2019.

[62] Adel Settati and Aadil Lahrouz. Stationary distribution of stochastic population systems under regime switching. *Applied Mathematics and Computation*, 244:235–243, 2014.

[63] Waclaw Szybalski and Vernon Bryson. Genetic studies on microbial cross resistance to toxic agents i. cross resistance of escherichia coli to fifteen antibiotics. *Journal of bacteriology*, 64(4):489–499, 1952.

[64] Josephine NA Tetteh, Sorin Olaru, Hans Crauel, and Esteban A Hernandez-Vargas. Scheduling collateral sensitivity-based cycling therapies toward eradication of drug-resistant infections. *International Journal of Robust and Nonlinear Control*, 33(9):4824–4842, 2023.

[65] Klas I Udekwu and Howard Weiss. Pharmacodynamic considerations of collateral sensitivity in design of antibiotic treatment regimen. *Drug Design, Development and Therapy*, pages 2249–2257, 2018.

[66] JG Coen Van Hasselt and Ravi Iyengar. Systems pharmacology: defining the interactions of drug combinations. *Annual review of pharmacology and toxicology*, 59:21–40, 2019.

[67] Xuping Xu and Panos J Antsaklis. Results and perspectives on computational methods for optimal control of switched systems. In *International Workshop on Hybrid Systems: Computation and Control*, pages 540–555. Springer, 2003.

[68] Nara Yoon, Robert Vander Velde, Andriy Marusyk, and Jacob G Scott. Optimal therapy scheduling based on a pair of collaterally sensitive drugs. *Bulletin of mathematical biology*, 80:1776–1809, 2018.

[69] Mari Yoshida, Sabrina Galinanes Reyes, Soichiro Tsuda, Takaaki Horinouchi, Chikara Furusawa, and Leroy Cronin. Time-programmable drug dosing allows the manipulation, suppression and reversal of antibiotic drug resistance in vitro. *Nature communications*, 8(1):15589, 2017.

[70] Guisheng Zhai, Bo Hu, Kazunori Yasuda, and Anthony N Michel. Stability analysis of switched systems with stable and unstable subsystems: an average dwell time approach. *International Journal of Systems Science*, 32(8):1055–1061, 2001.

A Appendix

A.1 Antibiotics for P. aeruginosa

Antibiotic	Abbreviation	Class	Target
Amikacin	AMI	aminoglycoside	protein synthesis, 30S
Gentamicin	GEN	aminoglycoside	protein synthesis, 30S
Tobramycin	TOB	aminoglycoside	protein synthesis, 30S
Ciprofloxacin	CIP	quinolone	DNA gyrase
Levofloxacin	LEV	quinolone	DNA gyrase
Ampicillin	AMP	b-lactam (penicillin)	cell wall
Piperacillin	PIP	b-lactam (penicillin)	cell wall
Carbenicillin	CAR	b-lactam (penicillin)	cell wall
Ticarcillin	TIC	b-lactam (penicillin)	cell wall
Aztreonam	AZE	b-lactam (monobactam)	cell wall
Cefepime	CFP	b-lactam (cephalosporin)	cell wall
Cefuroxime	CFX	b-lactam (cephalosporin)	cell wall
Ceftazidime	CFZ	b-lactam (cephalosporin)	cell wall
Meropenem	MER	b-lactam (carbapenem)	cell wall
Imipenem	IMI	b-lactam (carbapenem)	cell wall
Minocycline	MIN	tetracycline	protein synthesis, 30S
Doxycycline	DOX	tetracycline	protein synthesis, 30S
Azithromycin	AZY	macrolide	protein synthesis, 50S
Erythromycin	ERI	macrolide	protein synthesis, 50S
Clarithromycin	CLA	macrolide	protein synthesis, 50S
Colistin	COL	polymyxin	lipopolysaccharide
Fosfomycin	FOS	fosfomycin	cell wall biogenesis
Rifampicin	RIF	rifamycin	RNA synthesis
Trimethoprim	TMS	antifolate	combination folic acid pathway

Table A1: List of antibiotics used in [27].

Imamovic et al. exposed P. *aeruginosa* strain PAO1 to 24 clinically significant antibiotics from Table A1, increasing the concentration over 10 days such that by day 10 all strains could grow in the presence of the antibiotic when the antibiotic concentration was above the EUCAST breakpoint for P. *aeruginosa*. Dose response curves were performed with the 23 other antibiotics to elucidate collateral sensitivity or collateral resistance interactions showed in Figure A1.

Figure A1: Heat-map representation of collateral sensitivity and resistance from [27].

A.2 Invariant characterization

By neglecting the logistic term and considering a constant mutation rate Eq. (1) can be linearized as follows.

$$\frac{dx_i}{dt} = (\alpha_i^\sigma - \delta_i^\sigma)x_i(t) + \mu \sum_{j=1}^{n} m_{i,j}^\sigma x_j(t) \tag{2}$$

This can be written by the following equations.

$$\begin{pmatrix} \dot{x}_1(t) \\ \dot{x}_2(t) \\ \vdots \\ \dot{x}_2(t) \end{pmatrix} = \begin{pmatrix} \alpha_1^\sigma - \delta_1^\sigma & 0 & \cdots & 0 \\ 0 & \alpha_2^\sigma - \delta_2^\sigma & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & \alpha_n^\sigma - \delta_n^\sigma \end{pmatrix} \begin{pmatrix} x_1(t) \\ x_2(t) \\ \vdots \\ x_2(t) \end{pmatrix} + \tag{3}$$

$$+\mu \begin{pmatrix} 0 & m_{1,2}^\sigma & \cdots & m_{1,n}^\sigma \\ m_{2,1}^\sigma & 0 & \cdots & m_{2,n}^\sigma \\ \vdots & \vdots & \ddots & \vdots \\ m_{n,1}^\sigma & m_{n,2}^\sigma & \cdots & 0 \end{pmatrix} \begin{pmatrix} x_1(t) \\ x_2(t) \\ \vdots \\ x_2(t) \end{pmatrix} \tag{4}$$

Which is equivalent to

$$\begin{aligned} \dot{x} &= (A_\sigma + \mu M_\sigma)x, \\ &= \Lambda_\sigma x \end{aligned} \tag{5}$$

where x belong to the state constraint $\mathbb{X} \subset \mathbb{R}_{\geq 0}^n$ and $\sigma \in \Sigma$. Note that $\Lambda_\sigma^{-1}\Omega$ is well defined even if matrix Λ_σ is singular.

Lemma 2. *A set $\Omega \subset \mathbb{X}$ is an invariant set for linear system* (5) *if* $\Omega \subseteq \cup_{\sigma \in \Sigma} \Lambda_\sigma^{-1} \Omega$.

Algorithm 1: Invariant set within $\mathbb{T}$ for N drugs

Data: $\Lambda_\sigma, \mathbb{T}, \ \ \sigma = 1, 2, \ldots, N$
Result: Invariant Ω

$\Omega(0) = \mathbb{T}$;
while $\Omega(0) \not\subseteq \cup_{\sigma \in \Sigma} \Lambda_\sigma^{-1} \Omega(0)$ **do**
for $\sigma = 1 : N$ **do**
$\Omega(\sigma) = \Lambda_\sigma^{-1} \Omega(\sigma - 1) \cap \mathbb{T}$;
end
$\Omega(0) = \Omega(N)$;
end
return $\Omega = \Omega(0)$

Proposition 3. *Algorithm 1 converges to the largest invariant set of the switched system* (5) *contained in* $\mathbb{T}$, *or to an empty set.*

A.3   Parameters of simulation

Simulations of Figure 4. The states are given by $x_1 = AsBsCs$, $x_2 = ArBrCs$, $x_3 = AsBrCs$, $x_4 = AsBsCr$, $x_5 = AsBrCr$. The mutation rate and carrying capacity are considered the same for all states, $\mu_i = 10^{-4}$ and $K_i = 10^6$ for all $i = 1, \cdots, 5$, respectively. The growth and clearance parameters are given by Table A2.

PARAMETERS	AZTREONAM	CIPROFLOXACIN	TOBRAMYCIN
α_1	0.22	0.22	0.22
δ_1	0.25	0.25	0.25
α_2	0.27	0.27	0.22
δ_2	0.25	0.25	0.25
α_3	0.22	0.27	0.22
δ_3	0.25	0.25	0.25
α_4	0.22	0.22	0.27
δ_4	0.25	0.25	0.25
α_5	0.22	0.27	0.27
δ_5	0.25	0.25	0.25

Table A2: Model parameters for drugs on Case 5.