BootCellNet, a resampling-based procedure, promotes unsupervised identification of cell populations via robust inference of gene regulatory networks.

Yutaro Kumagai

Cellular and Molecular Biotechnology Research Institute, National Institute of Advanced Industrial Science and Technology, Higashi 1-1-1, Tsukuba, Ibaraki 305-8565, Japan

Correspondence:

Yutaro Kumagai,

Cellular and Molecular Biotechnology Research Institute, National Institute of Advanced Industrial Science and Technology, Higashi 1-1-1, Tsukuba, Ibaraki 305-8565, Japan.

E-mail: yutaro.kumagai@aist.go.jp.

Abstract

Recent advances in measurement technologies, particularly single-cell RNA sequencing (scRNA-seq), have revolutionized our ability to acquire large amounts of omics-level data on cellular states. As measurement techniques evolve, there has been an increasing need for data analysis methodologies, especially those focused on cell-type identification and inference of gene regulatory networks (GRNs). We have developed a new method named BootCellNet, which employs smoothing and resampling to infer GRNs. Using the inferred GRNs, BootCellNet further infers the minimum dominating set (MDS), a set of

genes that determines the dynamics of the entire network. We have demonstrated that

BootCellNet robustly infers GRNs and their MDSs from scRNA-seq data and facilitates

unsupervised identification of cell clusters using scRNA-seq datasets of peripheral blood

mononuclear cells and hematopoiesis. It has also identified COVID-19 patient-specific

cells and their potential regulatory transcription factors. BootCellNet not only identifies

cell types in an unsupervised and explainable way but also provides insights into the

characteristics of identified cell types through the inference of GRNs and MDS.

Author Summary

Single-cell omics technologies, such as single-cell RNA-seq (scRNA-seq), are

instrumental in identifying novel cell subsets that are involved in various biological

processes and diseases. These technologies, however, require further development in data

analysis, especially in areas focused on cell-type identification and inference of

interactions between genes. The problem of cell-type identification essentially involves

clustering, which necessitates a balance between distinguishing different cell states and

grouping similar ones together. Current clustering methods still suffer from uncertainty

in determining the appropriate number of clusters and in explaining why some cells are

clustered together but others are separated. The inference of interactions between genes,

gene regulatory network (GRN), remains challenging due to the noisy nature of scRNA-

seq. We have developed BootCellNet, a method that infers GRNs and a set of genes

dominating the network dynamics and utilizes the set to cluster cells and identify cell

types. The method addresses challenges in GRN identification and clustering methods

simultaneously and will facilitate the generation of the working hypotheses from a large

amount of scRNA-seq data.

Introduction

Recent advances in measurement technologies have revolutionized our ability to acquire

large amounts of omics-level data. Among these technologies, single-cell RNA

sequencing (scRNA-seq) has become particularly widespread. The technique allows for

an unbiased investigation of the distribution and status of various cell types within a

sample, providing insights into the dynamic and heterogeneous nature of cells.

As measurement techniques evolve, there has been an increasing need for data

analysis methodologies[1]. One significant challenge is the definition of cell types and

states from scRNA-seq data. This requires a balance between distinguishing different cell

states and grouping similar ones together. In this regard, the problem essentially boils

down to one of clustering. In the analysis of scRNA-seq, cells are clustered in an

unsupervised manner by using various algorithms. Among those, *k*-means clustering,

hierarchical clustering, spectral clustering, and Louvain clustering are examples of

methods utilized frequently. However, these methods do not provide an answer for the

appropriate number of clusters and an explanation of why some cells are clustered

together, but others are separated, leaving uncertainty in the analysis. An alternative to

unsupervised clustering is the use of supervised methods. Typically, cells are labeled by

using an annotated dataset. However, this approach faces difficulties in identifying novel

cell types not included in the given dataset.

The inference of gene regulatory networks (GRNs) from transcriptomic data like

scRNA-seq represents another major challenge. Successful unraveling of the complex

interactions and regulatory mechanisms that control gene expression within cells will

offer profound insights into cellular functions and disease mechanisms. Numerous studies

are dedicated to this field, employing a range of computational approaches[2]. Despite

the efforts, GRN inference is still challenging, due to the noisy nature of scRNA-seq,

characterized by dropouts and strong batch effects.

Once a GRN has been inferred, an essential next step is identifying genes that

are crucial for the overall control and regulation of the network. Understanding which

genes have a significant influence on the network can offer critical insights into the

underlying biological processes and potential targets for therapeutic intervention[3].

There are several methods to explore these pivotal elements within GRNs, such as the

selection of genes based on network centrality measures[4]. Among them, one of the

promising approaches relies on the concept of the minimum dominating set (MDS).

The MDS is a graph theoretical concept, defined as the smallest subset of vertices

in a graph such that every other vertex is adjacent to at least one vertex in this subset.

MDS plays a crucial role in controlling complex networks[5]. In the context of GRNs,

this translates to a minimal set of genes whose regulatory influence extends across the

entire network. In GRNs, MDS can be used to identify master regulatory genes[6]. In

protein interaction networks, MDS proteins are found to be enriched with essential,

cancer-related, and virus-targeted genes, showing a higher impact on network resilience

than hub proteins[7].

In this study, we introduce BootCellNet, a method that employs smoothing and

resampling procedures to reduce the noise in scRNA-seq data. We showed that this

approach robustly infers GRN and MDS. Since GRN controls the whole transcriptome

by definition, and MDS, in theory, governs the dynamics of the whole GRN, the

expression pattern of genes in MDS may reflect the entire transcriptome. Leveraging this

premise we utilized MDS for cell clustering. Because clusters are given based on the

MDS, the method gives a rational explanation of why some cells are clustered together

while others are not. Applying our method to multiple scRNA-seq datasets confirmed its

effectiveness. This addresses the aforementioned challenges in GRN identification and

clustering methods simultaneously.

This paper is organized as follows. In the first three sections, we introduced the

BootCellNet procedure and applied it to scRNA-seq data of peripheral blood

mononuclear cells. The first two sections showed the necessity of steps in BootCellNet

for robust GRN inference and MDS computation. The following section showed that

robustly computed MDS offers consistent clustering results, and the result was compared

to the results of supervised labeling and unsupervised clustering. In the next section, we

applied the method to scRNA-seq data of hematopoiesis to show if the method can be

applied to continuous spectra of cell states and help to infer developmental course. The

following section is devoted to the introduction of BootCellNet2, in which the unique set

of control nodes is used instead of MDS, and the clustering is automated. We showed that

the use of the new set of nodes and the automated clustering provides more robust

clustering results. Validation using CITE-seq data was also performed in the following

section to compare to unsupervised clustering. Finally, we applied BootCellNet2 to

scRNA-seq data of cells from COVID-19 patients. In each application, the relevance of

clustering results was evaluated by its robustness, and the biological relevance of the

obtained clusters was evaluated through the expert-based labeling of the obtained clusters.

Results.

BootCellNet enables robust GRN inference.

One of the problems in GRN inference from scRNA-seq data is its noisy and unstable

nature, making the inferred GRN unreliable. To overcome the noise, BootCellNet

employs two procedures, smoothing of scRNA-seq data and resampling-shuffling of data

to estimate *p*-values of genes and their regulations, as detailed in Methods (Figure 1).

To confirm the importance of each step, we first compared GRNs constructed

with or without smoothing. GRNs were inferred 10 times each with or without smoothing,

and appearances of genes/nodes and regulations/edges were counted (Figure 2A). As

shown in Figure 2B, the number of appeared genes was almost the same in GRNs with

smoothing as in those without smoothing before selection using *q*-values. After the

selection of genes using *q*-values, however, the number of genes was higher in GRNs

with smoothing. This result indicates that the smoothing procedure promotes the selection

of consistent genes and stable inference of GRN.

The above result shown in Figure 2B also implies that the smoothing alone is

sufficiently effective in extracting reliable genes/nodes because the resampling-based

selection of significant genes/nodes did not alter the number of genes/nodes drastically.

We examined the regulations/edges in the networks with or without resampling-based

selection. As shown in Figure 2C, GRNs with a resampling-based selection have a higher

burden of regulations/edges that appeared 10 times, and a lower burden of those that

appeared only once, indicating that the selection efficiently enriches regulations/edges

with high consistency.

BootCellNet enables robust MDS inference.

Once we have robustly inferred GRN, we further infer how the GRN regulates

gene expression by identifying MDS in the GRN. We implemented exact MDS

computation based on an integer programming formulation. GRNs inferred above were

subjected to the computation of MDS, and then the appearance of genes in each MDS

was counted. As shown in Figure 3A, the resampling-based selection increased the

number of genes in MDS that appeared in all 10 specimens. Moreover, the sizes of MDS

in GRNs with selection were larger than those without selection (Figure 3B), despite the

selection having reduced the number of genes/nodes and regulations/edges. Furthermore,

in the GRNs with the selection, MDS genes with a low degree rank appeared more

consistently than in the GRNs without the selection (Figure 3C). These results indicate

that the resampling-based selection of nodes and edges leads to consistent GRN structures

and robust inference of MDS.

BootCellNet determines cell type clusters in combination with the computation of MDS.

Since MDS, in theory, controls the dynamics of the whole network, the expression pattern

of genes in MDS may represent the whole transcriptome. Thus, it is assumed that the

expression pattern of genes in MDS can classify cells. To answer if GRN and MDS that

are robustly inferred by the BootCellNet can facilitate rational cell type classification, we

tested the idea on the PBMC scRNA-seq dataset. We first obtained small clusters of cells,

by using Louvain algorithm with high resolution. The resolution of 10 gave 63 clusters

of cells (Figure 4A). Hereafter, we call these original clusters "clumps" to distinguish

them from clusters obtained later.

Then we constructed GRN and MDS was computed as shown in Figure 4B. We

calculated the average expression of MDS genes in each cell clump and clustered the cell

clumps as well as the genes by using hierarchical clustering with Ward's method (Figure

4C, left). By visual inspection of the heatmap of the MDS expression, we determine the

number of clusters of clumps as 13 (Figure 4C, right). As a comparison, we also carried

out the same by using a set of genes with variable expression instead of MDS (Figure 4D).

Of 2000 variable genes, we sampled 100 genes (around three times the number of MDS,

containing 36.0 genes on average; Figure 3B) and performed the same procedure as for

MDS. We then applied the number of clusters of clumps as 13, as in the procedure for

MDS. The clustering was compared with each other and adjusted Rand indexes (ARIs)

were calculated. As shown in Figure 4E, MDS-based clustering had higher ARIs among

the 10 specimens than clustering based on 100 variable genes, indicating the robustness

of MDS-based clustering.

To evaluate the obtained clusters, we picked one specimen shown in Figure 5A

and checked the marker genes, Gene Ontology (GO) terms, and enriched TFBS in each

cluster in the specimen. Marker genes, GO terms, and enriched TFBSs are listed in

Supplementary Tables 1, 2, and 3, respectively. Cluster 1, 2, 3, and 9 exhibited

characteristics of T cells. Cluster 1 is associated with the signature of CD4 T cells,

whereas Cluster 2 is associated with that of CD8 T cells. Clusters 3 and 7 exhibit

characteristics of activated CD4 and CD8 T cells, respectively. Cluster 8 is also associated

with T cells, and based on the MDS expression pattern, it exhibits high RUNX3

expression, indicating the cluster is CD8 T cells[8]. Cluster 5 is associated with NK cell-

like signatures. Cluster 4 shows B cell characteristics, marked by CD19 and

immunoglobulin genes. Clusters 6, 10, and 11 feature markers whose promoters are

enriched with PU.1/SPI1 transcription factor binding site (TFBS), suggesting that these

clusters represent myeloid cell populations. CLEC10A and CD1C genes emerged as

markers of Cluster 10, pointing to myeloid dendritic cells (DCs)[9]. This aligns with the

cluster's association with myeloid DC signatures. Cluster 12, marked with PDGFRA and

GATA1, is linked to a platelet signature. Cluster 13, identified by SPIB, TCF4, and IRF7

as markers, all of which are well-known TFs involved in plasmacytoid DC (pDC)

development and function[10]. Consistently, the cluster is associated with pDC gene

signature, indicating that the cluster represents pDC population.

To compare this result with reference-based supervised cell labeling, the cells

were labeled by using SingleR with Human Cell Atlas dataset in celldex R package. The

main dataset, encompassing 37 cell types, labeled the PBMC scRNA-seq data with 7 cell

types: T cells, B cells, monocytes, platelets, NK cells, CMP, GMP, and pre-B cells (data

not shown). The fine dataset contains 157 subtypes, resulting in 19 cell types as shown in

Figure 5B. Most labels are consistent with the result from BootCellNet. However, the

reference-based labeling misclassified certain populations, such as clusters of myeloid

DCs (Cluster 10) and pDCs (Cluster 13) as monocytes and CD34- GMP/Pre-B cells,

respectively. Moreover, it tended to subdivide cells into smaller populations. NK cells,

represented by Cluster 5 in MDS-based clustering, were subdivided into several clusters

with similar markers, such as killer-cell receptors (KLRs) and granzymes (GZMs). The

activated CD8 T cell Cluster 7 in MDS-based clustering was mislabeled as a CD4 T cell.

B cells, represented by Cluster 4 in MDS-based clustering, were also divided into several

subpopulations, most with nearly identical markers. A population, B cell with memory

phenotype, has unique markers IGHA1/2, implying class switching which is a

characteristic of activated B cells. This implies that some populations might not be

identified by MDS-based clustering.

The result obtained by BootCellNet was also compared to that of unsupervised

Louvain clustering (Figure 5C). We chose a resolution of 1.0 because it gave 14 clusters,

a similar number to that of BootCellNet. Although the results were mostly similar, we

found two notable differences. First, a cluster of pDC was missing. Those cells were

mixed with cDC2 to form a cluster (Cluster 11 in Figure 5C). Second, Louvain clustering

could distinguish memory B cells from naïve B cells (Clusters 9 and 3 in Figure 5C,

respectively), while BootCellNet could not and mixed into one cluster (Cluster 4 in Figure

5A) as described above. These results imply that unsupervised clustering including

Louvain clustering is effective enough to identify cell types, though there's no guiding

principle to choose resolution and no rational explanation to separate or merge cells at the

step of clustering. We will discuss these points later.

BootCellNet identifies MDS in GRN. This provides insights into how the

differentiation and/or the function of cell types are regulated by checking the expression

of genes in MDS as shown in Figure 5D. Comparing the B cell cluster 4 and the pDC

cluster 13, TCF4, a TF important for pDC development[10], was expressed more in

Cluster 13. Cluster 12, representing platelets, has a high expression of GFI1B which

regulates platelet function[11]. We found CEBPB, a TF that controls

monocyte/macrophage differentiation, was higher in Clusters 6 and 11 of

monocyte/macrophage compared to Cluster 10 of DCs. Id2 is expressed more in the NK

cell cluster 5, compared to Clusters 1, 2, 3, 7, 8, and 9, all of which are clusters of T cells.

Consistently, the TF is essential for NK cell maturation[12]. Cluster 7 also shows high

Id2 expression, thought to a lesser extent. This is consistent with reports that Id2 is

important for CD8+ memory T cell development[13] because the cluster represents an

activated CD8 T cell population. Further, by assessing RUNX3 expression, we

determined Cluster 2, 7, and 8, which show higher RUNX3 expression, to be CD8 T cells,

whereas Cluster 1, 3, and 9 are identified as CD4 T cells, given that RUNX3 is important

for CD8 T cell development[8]. By examining RORA expression in Cluster 1 and 3, we

identified Cluster 3 as activated T cells, since RORA is notably expressed in Th17

population[14].

These findings collectively suggest that MDS-based clustering offers robust and

accurate cell type identification, largely aligning with reference-based supervised labeling,

while providing insights into the regulatory mechanisms of the cells through the

identification of GRN and MDS.

248 BootCellNet facilitates inference of developmental courses of

249 hematopoiesis.

The above result indicates that BootCellNet can identify cell types from scRNA-seq data

containing multiple discrete cell types. To examine if BootCellNet can help to infer

continuous spectra of cell states, we applied BootCellNet to scRNA-seq hematopoiesis

data. The scRNA-seq data of myeloid cell development[15] was subjected to BootCellNet.

The procedure resulted in a GRN of 110 TFs and MDS of 22 TFs (Figure 6A). The MDS

includes TFs that are known for their importance in myeloid/erythroid development, such

as Cebpe, Gata2, Gfi1b, Irf8, Jdp2, Klf1, Meis1, Myb, Sfpi1, and Sox4, further suggesting

the robust inference GRN and its dominating TFs by BootCellNet.

The cells were clustered into 9 as shown in Figure 6B. By examining the

expressed MDS TFs (Figure 6C), the marker genes (Supplementary Table 4), and

associated GO terms (Supplementary Table 5), we identified Cluster 1, 4, 6, and 8 as stem

cells, Cluster 2 as erythrocytes, Cluster 5 as megakaryocytes, Cluster 3 as monocytes, and

Cluster 7 as neutrophils. Cluster 9 seems to be lymphocytes and dendritic cells, which are

removed in the analyses performed in a report using the data[16].

Cluster 1 has Flt3 as a marker, as well as Sox4, Runx1, Cebpa, Lmo4, and Gata2,

which are TFs known to be expressed in hematopoietic stem cells. Cluster 3, 4, and 7 are

high in Sfpi1 expression, indicating that those clusters are myeloid lineages. However,

the expression of Sox4 distinguishes Cluster 4 from others as more progenitor-like cells.

Cluster 3 is associated with terms related to innate immune responses, and has Csf1r as a

marker gene, as well as Irf8. Cluster 7 is associated with myeloid cell-related terms and

has Fcnb and Elane genes as markers. The mentioned marker genes are characteristics of

monocytic and neutrophilic cells, respectively[17], supporting the identification of

Cluster 3 as monocytes and Cluster 7 as neutrophils. Comparing Cluster 3 and 7, the

former is high in Irf8 expression, whereas the latter has higher expression of Jdp2 and

Cebpe. This expression pattern of TFs is also a characteristic that distinguishes monocytes

and neutrophils[17, 18]. The importance of higher expression of Cebpe in the neutrophil

lineage has been suggested by using CellOracle, which is a cell lineage analysis method

based on GRN and a mathematical dynamical system model[16]. The result demonstrates

that BootCellNet efficiently infers MDS and their roles in development, without using

mathematical dynamical system models.

We inferred the lineage trajectories by using the clusters identified by

BootCellNet. Slingshot[19] was applied to the data, setting Cluster 1 as a starting point.

This resulted in 4 lineage curves (Figure 6D; one lineage is omitted, see the Figure Legend

for details). The result recapitulates known developmental trajectories and aligns with the

analysis performed by using a different method[16]. One trajectory represents the

development of erythrocytes spanning from Cluster 1 to 2 through 8 and 6. The others

represent that of monocytes and neutrophils spanning from Cluster 1, through 4, to Cluster

3 and 7, respectively. This result suggests that cell cluster identification by BootCellNet

facilitates lineage trajectory inference.

Automated cell type cluster identification by BootCellNet2.

Two ambiguous points remain in the BootCellNet procedure. First, MDS is not

necessarily uniquely determined. Second, we determined the number of clusters by visual

inspection of the hierarchical clustering heatmap.

To circumvent the first problem, we use the idea of critical nodes and intermittent

nodes. In the case where multiple MDSs exist, one can define critical nodes and

intermittent nodes, which appear in all the MDSs or at least in one MDS[20]. A set of

critical and intermittent nodes is uniquely determined. By using those sets, we could

circumvent possible inconsistencies derived from the non-uniqueness of MDS.

To address the second point, we utilize multiscale bootstrapping to determine

statistically significant clusters. The method provides approximately unbiased confidence

for each clustering[21]. By providing a cutoff p-value in advance, an R package pvclust

extracts statistically confident clusters of cell clumps. However, this leaves other clumps

unclustered. Thus we compare the clusters given by pvclust with the clusters obtained by

hierarchical clustering with varying numbers of clusters, and determine the number of

clusters.

We implemented the above methods (Figure 7A), and the procedure was applied

to the PBMC dataset. Using the same dataset as in Figure 2, we first checked whether the

new procedure consistently gives a unique set of nodes. We counted the appearance of

nodes, both critical and intermittent, in 10 specimens. As shown in Figure 7B, the number

of nodes that consistently appeared 10 times was not significantly different from that of

the former procedure. This indicates that the uniquely determined set of nodes is still

influenced by noises and resampling in GRN reconstruction.

313   We then performed clustering of cell clumps using the critical and intermittent

nodes instead of MDS. Also, we utilized multiscale bootstrap to determine the number of

clusters. The R package pvclust gives clusters with relatively low approximately unbiased

p-values. Based on the clustering, we determined the number of clusters as described in

detail in Methods and shown briefly in Figure 7A. The obtained clusters were compared

with each other and ARI was calculated. As shown in Figure 7B, the automated clustering

procedure along with the calculation of critical and intermittent nodes (hereafter we call

the nodes as control nodes) gave clustering results highly similar to each other. The

numbers of clusters were stable, 16 or 17 except for one 14. This indicates that the

procedure offered consistent clustering results. If we did not use the unique set of control

nodes for clustering but instead used MDS, the number of clusters had more variations.

ARIs were significantly higher in the combination of the automated clustering procedure

and the use of the unique set of nodes than others (Figure 7C). This suggests the

importance of the use of the unique set of nodes, although the set had variation as shown

above. Taken together, these results indicate that the use of the automated clustering

procedure together with the unique control node set offers more consistent clustering than

the former procedure.

330   To evaluate the newly obtained clusters, we picked the same specimen as shown

in Figure 5A. The resulting GRN with critical nodes and intermittent nodes is shown in

Figure 8A. The GRN contains 186 TFs with 22 critical and 33 intermittent TFs. Using the

nodes, cell clumps were clustered, obtaining 17 clusters. The clusters are mostly the same

as those in Figure 5A. We checked the marker genes and associated GO terms

(Supplementary Tables 6 and 7, respectively). Based on the marker genes and GO terms,

the cell clusters were labeled as shown in Figure 8B. Clusters of T cells were more

separated than in Figure 5A, and the novel clustering result enabled the identification of

naïve, activated, central memory, and effector memory subsets, along with Cluster 1, 4,

and 11, which represent naïve CD4 T cell subsets. As in the above analysis, we made use

of markers as well as the set of nodes to explain why those clusters are separated (Figure

8C). Cluster 1 is high in JUNB which is critical for the survival of T cells[22], while

Cluster 11 is high in NFE2L2/NRF2 which interfers activation of T cells[23] and Cluster

4 is not high for either, suggesting differential activation states of the three naïve T cell

clusters. These results together suggest that the novel procedure BootCellNet2 offers

automated and robust clustering and identification of biologically meaningful cell types.

348 BootCellNet2 identified cell type clusters complying with
349 CITE-seq only from transcriptome data.

We further validated BootCellNet2 using CITE-seq data, which allows us to evaluate

clustering results with antibody-derived tags. The BootCellNet2 procedure was applied

to transcriptome data from the CITE-seq study reported[24]. The analysis identified a set

of 29 control nodes, consisting of 19 critical nodes and 10 intermittent nodes

(Supplementary Figure 1), and generated 10 clusters as shown in Figure 9A. We labeled

the clusters based on their marker genes and associated GO terms for each cluster

(Supplementary Tables 8 and 9, respectively), as well as CITE-seq results (Figure 9C). In

comparison, we also performed clustering using the Louvain algorithm with a resolution

of 0.7, which resulted in 10 clusters, the same number of clusters as BootCellNet2. The

clusters were labeled as shown in Figure 9D, according to the marker genes, associated

GO terms, and CITE-seq results (Supplementary Tables 10 and 11, and Figure 9E,

respectively). The results suggest that the two methods offered largely comparable results.

A few key differences, however, were found. BootCellNet2 identified distinct

states of T cells similar to those observed in another PBMC dataset analysis shown above

(see Figure 8B), which was not the case with Louvain clustering. Moreover, CITE-seq

results suggest clear separation in CD4 and CD8 expressions within the T cell clusters

identified by BootCellNet2 (Cluster 1, 2, 3, and 5, as shown in Figure 9B); Clusters 3 and

5 exhibited low CD4 and high CD8, whereas Clusters 1 and 2 showed high CD4 and low

CD8 expression. The difference between the states of CD4 and CD8 T cells was also

evident from CD45RA expression known to be high in naïve T cells (high in Cluster 1

and low in Cluster 2) and CD57 expression known to be high in activated T cells (high in

Cluster 3 and low in Cluster 5), respectively. Expression of HMGB2, which is highly

expressed in effector and memory T cells compared to naïve T cells[25], further supports

these findings (Figure 9C). Additionally, platelets were identified by BootCellNet2 but

not by Louvain. These results collectively imply that BootCellNet2 clustering achieves

more accurate cell type delineation, aligning more closely with antibody-derived tag

results than unsupervised clustering.

Identification of characteristic cell types in COVID-19 patients

by BootCellNet2.

We applied BootCellNet2 to scRNA-seq data of bronchoalveolar lavage fluid cells from

COVID-19 patients[26] to test whether BootCellNet2 can identify cell types in scRNA-

seq data where discrete and continuous cell types/states are mixed. The cells were

clustered into 32 in the original article. BootCellNet2 gave the GRN of 161 TFs with 21

critical nodes and 13 intermittent nodes (Figure 10A). The automated clustering

procedure offered 17 clusters of the cell clumps (Figure 10B). Marker genes were

identified for each cluster, and the GO terms associated with the markers of each cluster

were checked (Supplementary Tables 12 and 13, respectively).

Clusters 1, 2, 3, 5, 8, 13, 15, and 16 were associated with myeloid cell-related

signatures. Compared to the clusters in the original report (reproduced in Supplementary

Figure 2, encircled as "macrophage/monocyte", "neutrophil", and "DC"), the clustering

by BootCellNet2 has a smaller number of clusters while identifying myeloid cells with

different characteristics.

Clusters 1, 3, and 8 were associated with inflammatory immune responses and/or

characteristics of macrophage/monocyte/DC thus identified as myeloid cells. Clusters 1,

3, and 8 have high expression of FOS, IRF1, and NFKB2, which are TFs involved in

inflammatory immune response (Figure 10C). The clusters are significantly enriched with

cells from severe patients (Figure 10D). The associated terms are also related to

inflammatory responses. In contrast, Clusters 2, 5, 13, 15, and 16 are enriched with cells

from mild and/or healthy individuals. Clusters 5, 6, 13, 14, and 16 have HLA genes as

markers and are associated with DC/monocyte signatures, implying the clusters contain

antigen-presenting myeloid cells. Clusters 5, 13, and 16 have either MARCO, CEBPA,

and PPARG as markers, indicating that the clusters are alveolar macrophages since

PPARG is an essential TF for alveolar macrophage development[27]. Clusters 5, 6, 13,

14, and 16 did not show high expression of inflammatory TFs such as FOS, IRF1, and

NFKB2, in contrast to Clusters 1, 4, and 8. This implies that the myeloid cells in healthy

individuals or patients with mild symptoms are less inflammatory than those in

individuals with severe disease, consistent with former reports showing that inflammation

and cytokine storms are the cause of severe COVID-19[28].

Cluster 2 is associated with myeloid and dendritic cell signatures. It has CD19

as a marker. Indeed, the cluster may contain pDC, myeloid DC, and B cells, which were

clustered separately in the original report (Supplementary Figure 2), suggesting a

probable failure of the BootCellNet2 procedure in the identification of cell types.

Clusters 4 and 15 are associated with T cell signature and have genes for components of

T cell receptors such as CD3D/E/G, LCK, LAT, and ZAP70 as markers. They also have

CD8A/B and RUNX3 as markers, indicating that those clusters are CD8 T cell

populations. Further, all of them have genes for granzymes and IFN-g as markers,

suggesting that the clusters represent activated CD8 T cells.

To examine the differences among those activated T cells clustered separately,

we compared Cluster 4 to Cluster 15 and found that the latter, which is enriched with cells

from patients with mild symptoms, but not the former, which is enriched with cells from

patients with severe symptoms, has ID2, TCF7, and CCR5 as markers, which are highly

expressed in effector memory CD8 T cells[13, 29, 30]. We also examined the expression

of IL7R (encoding IL-7Ralpha) and PDCD1 (encoding PD-1) and found that Cluster 15

has both as markers while Cluster 4 has only PDCD1, indicating that Cluster 4 is

exhausted T cells [31]. Notably, examination of the expression of the control nodes in the

GRN also showed that HMGB2, which is indispensable for the development of exhausted

T cells[25], is higher in Cluster 4 than in Cluster 15 (Figure 10C). These results

collectively suggest that induction of effector memory CD8 T cell upon SARS-CoV-2

infection is critical for controlling disease to mild symptoms, and an increase of exhausted

T cells is detrimental for disease control, consistent with former reports[26, 32, 33].

Taken together, the results indicate that BootCellNet2 successfully identifies

clusters of cells specific to patients with different symptoms, and offers explanations of

the difference in symptoms that are consistent with preceding reports.

Discussion.

In the current study, a method named BootCellNet has been developed, which combines

smoothing, resampling, GRN inference, computation of a set of genes controlling the

entire network, and clustering. We demonstrated the robustness of our method,

particularly given the probabilistic steps involved and the presence of noise within the

data. We also performed an expert-based evaluation of the clusters by manually labeling

the clusters. This ensured that despite the probabilistic nature of our method, the clusters

identified are not only statistically robust but also biologically meaningful.

Since the method consists of multiple steps, the choice of algorithms will affect

the results of the procedure. We used k-NN smoothing to reduce the noise in scRNA-seq.

We performed the smoothing on the 2-dimensional UMAP plane, but one can do the same

in a higher dimension[34]. One can also use different smoothing methods, such as k-NN

graph-based MetaCell algorithm[35], or data imputation methods such as MAGIC[36]

and DeepImpute[37].

In BootCellNet, GRN reconstruction is performed with the ARACNe method,

because of its low computational load. As in the current study, NestBoot utilizes a nested

bootstrap to control FDR in GRN inference[38], and they showed that the bootstrapping

procedure improved the accuracy of the GRN inference by various inference methods

such as GENIE3[39]. Thus, the incorporation of methods other than ARACNe in

BootCellNet may improve the GRN inference accuracy.

The use of MDS as a set of genes that controls the dynamics of the whole

network needs some caution. The controllability of a network by its MDS requires a

premise that each MDS node can control its adjacent nodes individually[5]. In the case

where nodes are TFs, this may be justified by assuming that a TF controls each target

gene through individually controlled accessibility, although the assumption should be

verified independently.

Nevertheless, we demonstrated that the utilization of expression levels of only

several tens of MDS genes or the unique set of control nodes was sufficient for robust

clustering of cells while using one hundred genes with variable expression was not. This

implies that MDS can provide a dimensionality reduction of scRNA-seq data.

We inferred the network of TF genes in the current study, because we used

transcriptome data, which is principally controlled by TFs. But this could be extended by

incorporating not only TFs, but also co-factors, post-transcriptional regulators, and

soluble mediators that control TF expression and/or activity. The incorporation of these

factors also conforms to the picture where transcriptome activity is controlled principally

on the mRNA level. Indeed, some post-transcriptional regulators, such as ZFP36L1 and

ZFP36L2 were included and found to be a part of MDS. The incorporation of more factors

will help to uncover dynamic picture of the regulatory network.

Despite the room for improvement and caution, and although the method is

relatively simpler than other sophisticated methods, we demonstrated that BootCellNet

effectively worked for a variety of datasets with different characteristics. Notably, the

proposed method excels in explainability, not only clustering cells but also explaining

why they are clustered based on the GRN, and providing characterization of what type of

population they represent. Along with automating the determination of the number of

clusters, the method resolves the common problem in unsupervised clustering which does

not provide an answer for the appropriate number of clusters and/or an explanation of

why some cells are clustered together but others are separated. For example, the clusters

can be explained by factors such as GFI1B, CEBPB, and RUNX3 in clustering PBMC

data, by Cebpe in clustering hematopoiesis data, and by HMGB2 in clustering COVID-

19 data, allowing the identification of what type of cells the clusters represent and what

characteristics they possess, even within similar cell types, without looking at GO terms,

using information from GRN/MDS. This demonstrates that BootCellNet can be a useful

complementary for other methods of clustering, supervised labeling, and lineage

inference.

Also, BootCellNet, especially BootCellNet2, is suitable for automation. The

method will be utilized for automated and routine annotation of large-scale scRNA-seq

cell data. Moreover, BootCellNet addresses an increasing need for explainable automated

data processing[40], since it has a firm basis in statistics and network dynamics. Although

there is a strict need for manual inspection and experimental verification[1], the proposed

method will accelerate the generation of the working hypotheses from a large amount of

scRNA-seq data.

Materials and Methods

Outline of the BootCellNet procedure.

The proposed procedure is outlined in Figure 1. The procedure, BootCellNet, utilizes

smoothing of scRNA-seq expression data and resampling to reduce noise in the dataset

and to facilitate a robust inference of gene regulatory network (GRN), in combination

with computation of minimum dominating set (MDS). Using the MDS, BootCellNet

performs clustering of cells to identify cell types and states.

BootCellNet2 is outlined in Figure 7A. The procedure utilizes a unique set of

control nodes consisting of critical nodes and intermittent nodes, instead of MDS, and the

clustering is automated as described below.

Estimation of statistical significance by using resampling.

The nature of the BootCellNet procedure enables statistical evaluation of inferred GRNs.

BootCellNet repeatedly samples cells from the smoothed scRNA-seq dataset and

performs GRN inference. Some nodes (genes) and edges (regulatory interactions)

frequently show up in each resampled set, while others do only occasionally. At the same

time, we shuffle gene names and perform the same inferences repeatedly to evaluate the

occurrence of false positives.

We first performed the k-NN smoothing on 2-dimensional UMAP space. The k

is set to a value small enough to find a cell population having about k cells in a dataset.

In this regard, the k determines the resolution of the analysis. For given k and 2-

dimensional UMAP space, for each cell, k-nearest neighbors in the UMAP space with

Euclidean distance are selected, and the average of the expression levels is computed to

obtain k-NN smoothed cells. The decimal part of the averaged expression levels was

truncated to obtain expression levels in integers.

Next, we resample N k-NN smoothed cells for M times to infer GRN or for $\hat{M}$

times to calculate the support values of appearances of genes/nodes and regulations/edges.

In the current study, the resampling number N was set to from one-third to half of the total

number of cells in the dataset, but no more than 20000 to reduce the computation load.

For each resampled set, we compute variable expression genes and then infer GRN. We

also perform shuffling of gene names in each resampled set to calculate the support.

Let the resampled sets be $X_m = \{x_{m1}, x_{m2}, \ldots, x_{mN}\}$, where $m = 1, 2, \ldots, \widehat{M}$, and

x be a vector of k-NN smoothed gene expressions of a cell. Shuffling of gene names in

X_m for K times is performed to generate nested resampling sets X_{mk} ($k = 1, 2, \ldots K$)[41].

Then we obtain $S(X_{mk}; \theta)$, which takes 1 if a hypothesis θ is supported by the set X_{mk}

or 0 if not. In the current case, θ is a hypothesis that we find a gene as a variable

expression gene or one that we find a regulatory relationship (i.e. an edge of GRN) in a

GRN inferred from X_{mk}. The parameter to be estimated is $\lambda(\theta) = \frac{1}{\widehat{M}K} \sum_{m,k} S(X_{mk}; \theta)$,

representing how often we occasionally have θ. Once we estimate the parameter, we can

compute a p-value by supposing the event obeys a Poisson distribution with parameter

λM. However, the rate of occasional appearances of genes is equal to (the number of

variable expression genes selected) / (the number of total genes), and thus there's no need

to compute by resampling. Hence, we computed the λ only for regulations/edges in GRNs.

The obtained p-values were adjusted by the Benjamini-Yekutieli method to obtain q-

values. Nodes and edges with $q<$1e-05 were considered significant, and non-significant

nodes and edges were removed.

In the implementation of this paper, resampling of cells was performed using a

function named BootCellNet, and shuffling of gene names was carried out in a function

named AggregateBCN. For inferring GRN, resampling was done with $M = 100$, but the

shuffling of gene names was not performed ($K = 0$). In the estimation of the support,

resampling was performed with $\hat{M} = 1000$, and gene name shuffling was carried out with

$K = 100$, obtaining 10^5 resampled specimens.

Inference of gene regulatory network.

For each resampled specimen, 2000 genes with variable expression levels were selected

by using FindVariableFeature function in Seurat[42] with vst method. From the set of

genes, we only focused on transcription factors (TFs). The list of TFs was retrieved from

a reported list of human TFs[43] in the case of human dataset analysis, and from

AnimalTFDB 4.0[44] in the case of mouse dataset analysis. The TFs with variable

expression levels were subjected to GRN inference.

GRN is inferred by using ARACNe method[45] implemented in an R package

minet[46] with the estimation of mutual information by Miller-Madow asymptotic bias

corrected empirical estimator and an equal width discretization, as recommended in a

report[47].

Computation of minimal dominating set (MDS).

The minimum dominating set (MDS) is a graph theoretical concept, which is the smallest

subset of vertices in a graph such that every other vertex is adjacent to at least one vertex

in this subset. MDS can be computed exactly by using an integer programming

formulation described elsewhere[48]. We implemented the method using the R package

lpSolve. Although the problem is known as an NP-hard problem, the number of genes in

the GRN is typically less than 200 and thus MDS is computable in a laptop computer. We

also implemented the computation of critical and intermittent nodes. To obtain the nodes,

the computation of MDS is repeated for the defined times (15 times as default, by setting

num.bin.solns = 15 in the function lp of the lpSolve package), and nodes identified as

MDS in all the computations are defined as critical nodes, and those in at least one

computation are defined as intermittent nodes.

Clustering of cell clumps.

In the BootCellNet procedure, we start from small clusters and aggregate them. The

starting small clusters (here we call them clumps to distinguish them from the final

clusters) were computed by using Louvain algorithm with a high resolution implemented

in Seurat. The resolution was determined so that the size of the smallest clump was not

much lower than k (we use 80 % of k as a default procedure), which is the parameter used

in the smoothing.

After obtaining the cell clumps, expression levels of MDS genes were averaged

over each cluster. Using the expression levels, the clumps are hierarchically clustered by

using Ward's algorithm with a value of Pearson correlation subtracted from 1 as a distance.

The number of clusters of clumps was determined by visual inspection.

In the BootCellNet2 procedure, an R package pvclust is used to compute

approximately unbiased p values for clusters of clumps with the multiscale bootstrap

algorithm. A function pvpick in the pvclust package extracts clusters having p values less

than a given cutoff to offer a high-confident clustering. ARIs are computed between the

high-confident clustering result given by pvclust/pvpick and those given by a function

rect.hclust with varying numbers of clusters. Then the number of clusters of clumps is

determined as the lowest number with a maximal ARI (see Figure 7A).

Single-cell RNA-seq data processing.

The peripheral blood mononuclear cell (PBMC) dataset was provided by 10x Genomics

(https://cf.10xgenomics.com/samples/cell-

exp/3.1.0/manual_5k_pbmc_NGSC3_ch5/manual_5k_pbmc_NGSC3_ch5_filtered_feat

ure_bc_matrix.tar.gz). Expression levels and metadata were aggregated and processed

with a standard workflow by using Seurat. The processed data containing 3707 cells was

then subjected to BootCellNet with $k = 10$ and $N = 3000$. The starting 63 cell clumps were

given with a resolution of 10, which was determined according to the criterion described

above.

The myeloid/erythroid development scRNA-seq dataset[15] was retrieved from

a repository (Paul_Cell_MARSseq_GSE72857.RData,

https://github.com/theislab/scAnalysisTutorial). The data, comprised of 2730 cells, were

processed with a standard workflow by using Seurat. The processed data were subjected

to BootCellNet procedure with $k = 10$ and $N = 2000$. The starting 62 cell clumps were

obtained with a resolution of 10.

The CITE-seq data of PBMC[24] was retrieved from NCBI GEO under

accession number GSE100866. The data is comprised of scRNA-seq data and antibody-

derived tag sequencing data. These data were aggregated and processed with a standard

workflow by using Seurat. The processed data containing 7690 cells after filtering was

then subjected to BootCellNet with $k = 10$ and $N = 5000$. The starting 66 cell clumps were

obtained by using Louvain clustering with a resolution of 9.

The single-cell RNA-seq data of COVID-19 patients[26] were obtained from

UCSC Cell Browser (http://cells.ucsc.edu/covid19-balf/nCoV.rds). The data is comprised

of 66452 cells in total of bronchoalveolar lavage fluid from 4 healthy individuals, 3 mild

and 6 severe patients. The data were analyzed by using Seurat and were subjected to

BootCellNet with $k = 20$ and $N = 20000$. The starting 113 cell clumps were given by using

Louvain clustering with a resolution of 10.

As a comparison, supervised cell type annotation was performed by using

SingleR[49] with Human Primary Cell Atlas fine subtype labels in an R package

celldex[49].

625 Transcription factor binding site (TFBS), Gene Ontology (GO)

626 enrichment and lineage analyses.

627 Marker genes were obtained by using FindAllMarkers function in Seurat. We used

628 Homer[50] for TFBS and GO analysis of the marker genes of each cluster. Promoters are

629 defined as from -500 to +200 of transcription start sites. For lineage analysis,

630 Slingshot[19] was utilized with default parameters.

632 Statistical comparison of clustering results.

633 Adjusted Rand index[51] was calculated to compare two clustering results.

636 Author Contributions

637 Conceptualization, algorithm development and coding, data analysis, funding acquisition,

638 manuscript preparation; Y. K.

Competing Interests

None declared.

Data availability

The code for BootCellNet is deposited in the author's GitHub (https://github.com/oursvallee/BootCellNet). All the data used are publicly available as described in Methods.

Acknowledgment.

The authors thank Drs. Yasuyuki Kida, Nobuhito Mori, and Kentaro Kawata for discussions, Mr. Akihiko Fujii for technical assistance, and Dr. Yasuko Ozaki for secretarial assistance. This research has partly been supported by Kakenhi 22H04925 (PAGS) and 23H04486, and by a Grant from International Joint Usage/Research Center, the Institute of Medical Science, the University of Tokyo. Computational time was in part provided by the NIG supercomputer at ROIS National Institute of Genetics and by Human Genome Center, the Institute of Medical Science, the University of Tokyo.

References

1.	Heumos L, Schaar AC, Lance C, Litinetskaya A, Drost F, Zappia L, et al. Best practices for single-cell analysis across modalities. Nat Rev Genet. 2023;24(8):550-72. doi: 10.1038/s41576-023-00586-w. PubMed PMID: 37002403; PubMed Central PMCID: PMCPMC10066026 Janssen, and received consulting fees from Chan-Zuckerberg Initiative. F.J.T. consults for Immunai Inc., Singularity Bio B.V., CytoReason Ltd and Omniscope Ltd, and has ownership interest in Dermagnostix GmbH and Cellarity. M.G.J. consults for and has ownership interests in Vevo Therapeutics. L. Heumos has received

speaker's honorarium from Vesalius Therapeutics. Single-Cell Best Practices Consortium: M.G.J. consults for and has ownership interests in Vevo Therapeutics. R.P. is co-founder of Ocean Genomics, Inc. The other authors declare no competing interests.

2. Nguyen H, Tran D, Tran B, Pehlivan B, Nguyen T. A comprehensive survey of regulatory network inference methods using single cell RNA sequencing data. Brief Bioinform. 2021;22(3). doi: 10.1093/bib/bbaa190. PubMed PMID: 34020546; PubMed Central PMCID: PMCPMC8138892.

3. Guo WF, Zhang SW, Zeng T, Akutsu T, Chen L. Network control principles for identifying personalized driver genes in cancer. Brief Bioinform. 2020;21(5):1641-62. doi: 10.1093/bib/bbz089. PubMed PMID: 31711128.

4. Badia IMP, Wessels L, Muller-Dott S, Trimbour R, Ramirez Flores RO, Argelaguet R, et al. Gene regulatory network inference in the era of single-cell multi-omics. Nat Rev Genet. 2023;24(11):739-54. doi: 10.1038/s41576-023-00618-5. PubMed PMID: 37365273.

5. Nacher JC, Akutsu T. Dominating scale-free networks with variable scaling exponent: heterogeneous networks are not difficult to control. New Journal of Physics. 2012;14(7):073005.

6. Nazarieh M, Wiese A, Will T, Hamed M, Helms V. Identification of key player genes in gene regulatory networks. BMC Syst Biol. 2016;10(1):88. doi: 10.1186/s12918-016-0329-5. PubMed PMID: 27599550; PubMed Central PMCID: PMCPMC5011974.

7. Wuchty S. Controllability in protein interaction networks. Proc Natl Acad Sci U S A. 2014;111(19):7156-60. doi: 10.1073/pnas.1311231111. PubMed PMID: 24778220; PubMed Central PMCID: PMCPMC4024882.

8. Egawa T, Tillman RE, Naoe Y, Taniuchi I, Littman DR. The role of the Runx transcription factors in thymocyte differentiation and in homeostasis of naive T cells. J Exp Med. 2007;204(8):1945-57. doi: 10.1084/jem.20070133. PubMed PMID: 17646406; PubMed Central PMCID: PMCPMC2118679.

9. Collin M, Bigley V. Human dendritic cell subsets: an update. Immunology. 2018;154(1):3-20. doi: 10.1111/imm.12888. PubMed PMID: 29313948; PubMed Central PMCID: PMCPMC5904714.

10. Reizis B. Plasmacytoid Dendritic Cells: Development, Regulation, and Function. Immunity. 2019;50(1):37-50. doi: 10.1016/j.immuni.2018.12.027. PubMed PMID: 30650380; PubMed Central PMCID: PMCPMC6342491.

11. Stevenson WS, Morel-Kopp MC, Chen Q, Liang HP, Bromhead CJ, Wright S, et al. GFI1B mutation causes a bleeding disorder with abnormal platelet function. J Thromb Haemost. 2013;11(11):2039-47. doi: 10.1111/jth.12368. PubMed PMID: 23927492.

12.   Li ZY, Morman RE, Hegermiller E, Sun M, Bartom ET, Maienschein-Cline M,
et al. The transcriptional repressor ID2 supports natural killer cell maturation by
controlling TCF1 amplitude. J Exp Med. 2021;218(6). doi: 10.1084/jem.20202032.
PubMed PMID: 33857289; PubMed Central PMCID: PMCPMC8056751 the submitted
work. No other disclosures were reported.
13.   Cannarile MA, Lind NA, Rivera R, Sheridan AD, Camfield KA, Wu BB, et al.
Transcriptional regulator Id2 mediates CD8+ T cell immunity. Nat Immunol.
2006;7(12):1317-25. doi: 10.1038/ni1403. PubMed PMID: 17086188.
14.   Yang XO, Pappu BP, Nurieva R, Akimzhanov A, Kang HS, Chung Y, et al. T
helper 17 lineage differentiation is programmed by orphan nuclear receptors ROR alpha
and ROR gamma. Immunity. 2008;28(1):29-39. doi: 10.1016/j.immuni.2007.11.016.
PubMed PMID: 18164222; PubMed Central PMCID: PMCPMC2587175.
15.   Paul F, Arkin Y, Giladi A, Jaitin DA, Kenigsberg E, Keren-Shaul H, et al.
Transcriptional Heterogeneity and Lineage Commitment in Myeloid Progenitors. Cell.
2015;163(7):1663-77. doi: 10.1016/j.cell.2015.11.013. PubMed PMID: 26627738.
16.   Kamimoto K, Stringa B, Hoffmann CM, Jindal K, Solnica-Krezel L, Morris SA.
Dissecting cell identity via network inference and in silico gene perturbation. Nature.
2023;614(7949):742-51. doi: 10.1038/s41586-022-05688-9. PubMed PMID: 36755098;
PubMed Central PMCID: PMCPMC9946838.
17.   Kwok I, Becht E, Xia Y, Ng M, Teh YC, Tan L, et al. Combinatorial Single-Cell
Analyses of Granulocyte-Monocyte Progenitor Heterogeneity Reveals an Early Uni-
potent   Neutrophil   Progenitor.   Immunity.   2020;53(2):303-18   e5.   doi:
10.1016/j.immuni.2020.06.005. PubMed PMID: 32579887.
18.   Maruyama K, Fukasaka M, Vandenbon A, Saitoh T, Kawasaki T, Kondo T, et al.
The transcription factor Jdp2 controls bone homeostasis and antibacterial immunity by
regulating osteoclast and neutrophil differentiation. Immunity. 2012;37(6):1024-36. doi:
10.1016/j.immuni.2012.08.022. PubMed PMID: 23200825.
19.   Street K, Risso D, Fletcher RB, Das D, Ngai J, Yosef N, et al. Slingshot: cell
lineage and pseudotime inference for single-cell transcriptomics. BMC Genomics.
2018;19(1):477. doi: 10.1186/s12864-018-4772-0. PubMed PMID: 29914354; PubMed
Central PMCID: PMCPMC6007078.
20.   Nacher JC, Akutsu T. Analysis of critical and redundant nodes in controlling
directed and undirected complex networks using dominating sets. Journal of Complex
Networks. 2014;2(4):394-412. doi: 10.1093/comnet/cnu029.
21.   Shimodaira H, Hasegawa M. Multiple comparisons of log-likelihoods with
applications   to   phylogenetic   inference.   Molecular   biology   and   evolution.

1999;16(8):1114.

22.   Hsieh T, Sasaki D, Taira N, Chien H, Sarkar S, Seto Y, et al. JunB Is Critical for
Survival of T Helper Cells. Front Immunol. 2022;13:901030. doi:
10.3389/fimmu.2022.901030. PubMed PMID: 35837408; PubMed Central PMCID:
PMCPMC9273772.

23.   Muri J, Kopf M. Redox regulation of immunometabolism. Nat Rev Immunol.
2021;21(6):363-81. doi: 10.1038/s41577-020-00478-8. PubMed PMID: 33340021.

24.   Stoeckius M, Hafemeister C, Stephenson W, Houck-Loomis B, Chattopadhyay
PK, Swerdlow H, et al. Simultaneous epitope and transcriptome measurement in single
cells. Nat Methods. 2017;14(9):865-8. doi: 10.1038/nmeth.4380. PubMed PMID:
28759029; PubMed Central PMCID: PMCPMC5669064.

25.   Neubert EN, DeRogatis JM, Lewis SA, Viramontes KM, Ortega P, Henriquez
ML, et al. HMGB2 regulates the differentiation and stemness of exhausted CD8(+) T cells
during chronic viral infection and cancer. Nat Commun. 2023;14(1):5631. doi:
10.1038/s41467-023-41352-0. PubMed PMID: 37704621; PubMed Central PMCID:
PMCPMC10499904.

26.   Liao M, Liu Y, Yuan J, Wen Y, Xu G, Zhao J, et al. Single-cell landscape of
bronchoalveolar immune cells in patients with COVID-19. Nat Med. 2020;26(6):842-4.
doi: 10.1038/s41591-020-0901-9. PubMed PMID: 32398875.

27.   Schneider C, Nobs SP, Kurrer M, Rehrauer H, Thiele C, Kopf M. Induction of
the nuclear receptor PPAR-gamma by the cytokine GM-CSF is critical for the
differentiation of fetal monocytes into alveolar macrophages. Nat Immunol.
2014;15(11):1026-37. doi: 10.1038/ni.3005. PubMed PMID: 25263125.

28.   Vanderbeke L, Van Mol P, Van Herck Y, De Smet F, Humblet-Baron S, Martinod
K, et al. Monocyte-driven atypical cytokine storm and aberrant neutrophil activation as
key mediators of COVID-19 disease severity. Nat Commun. 2021;12(1):4117. doi:
10.1038/s41467-021-24360-w. PubMed PMID: 34226537; PubMed Central PMCID:
PMCPMC8257697.

29.   Zhou X, Yu S, Zhao DM, Harty JT, Badovinac VP, Xue HH. Differentiation and
persistence of memory CD8(+) T cells depend on T cell factor 1. Immunity.
2010;33(2):229-40. doi: 10.1016/j.immuni.2010.08.002. PubMed PMID: 20727791;
PubMed Central PMCID: PMCPMC2928475.

30.   Kohlmeier JE, Miller SC, Smith J, Lu B, Gerard C, Cookenham T, et al. The
chemokine receptor CCR5 plays a key role in the early memory CD8+ T cell response to
respiratory virus infections. Immunity. 2008;29(1):101-13. doi:
10.1016/j.immuni.2008.05.011. PubMed PMID: 18617426; PubMed Central PMCID:

PMCPMC2519120.

31.   Seo W, Jerin C, Nishikawa H. Transcriptional regulatory network for the

establishment of CD8(+) T cell exhaustion. Exp Mol Med. 2021;53(2):202-9. doi:

10.1038/s12276-021-00568-0. PubMed PMID: 33627794; PubMed Central PMCID:

PMCPMC8080584 Pharmaceutical, MSD and Bristol-Myers Squibb and research

funding from Taiho Pharmaceutical, Daiichi-Sankyo, Kyowa Kirin, Zenyaku Kogyo,

Oncolys BioPharma, Debiopharma, Asahi-Kasei, Sysmex, Fujifilm, SRL, Astellas

Pharmaceutical, Sumitomo Dainippon Pharma and BD Japan outside of this study. The

other authors declare no competing interests.

32.   Odak I, Barros-Martins J, Bosnjak B, Stahl K, David S, Wiesner O, et al.

Reappearance of effector T cells is associated with recovery from COVID-19.

EBioMedicine. 2020;57:102885. doi: 10.1016/j.ebiom.2020.102885. PubMed PMID:

32650275; PubMed Central PMCID: PMCPMC7338277.

33.   Wauters E, Van Mol P, Garg AD, Jansen S, Van Herck Y, Vanderbeke L, et al.

Discriminating mild from critical COVID-19 by innate and adaptive immune single-cell

profiling of bronchoalveolar lavages. Cell Res. 2021;31(3):272-90. doi: 10.1038/s41422-

020-00455-9. PubMed PMID: 33473155; PubMed Central PMCID: PMCPMC8027624.

34.   Wagner F, Yan Y, Yanai I. K-nearest neighbor smoothing for high-throughput

single-cell RNA-Seq data. BioRxiv. 2017:217737.

35.   Baran Y, Bercovich A, Sebe-Pedros A, Lubling Y, Giladi A, Chomsky E, et al.

MetaCell: analysis of single-cell RNA-seq data using K-nn graph partitions. Genome Biol.

2019;20(1):206. doi: 10.1186/s13059-019-1812-2. PubMed PMID: 31604482; PubMed

Central PMCID: PMCPMC6790056.

36.   van Dijk D, Sharma R, Nainys J, Yim K, Kathail P, Carr AJ, et al. Recovering

Gene Interactions from Single-Cell Data Using Data Diffusion. Cell. 2018;174(3):716-

29 e27. doi: 10.1016/j.cell.2018.05.061. PubMed PMID: 29961576; PubMed Central

PMCID: PMCPMC6771278.

37.   Arisdakessian C, Poirion O, Yunits B, Zhu X, Garmire LX. DeepImpute: an

accurate, fast, and scalable deep neural network method to impute single-cell RNA-seq

data. Genome Biol. 2019;20(1):211. doi: 10.1186/s13059-019-1837-6. PubMed PMID:

31627739; PubMed Central PMCID: PMCPMC6798445.

38.   Morgan D, Tjarnberg A, Nordling TEM, Sonnhammer ELL. A generalized

framework for controlling FDR in gene regulatory network inference. Bioinformatics.

2019;35(6):1026-32. doi: 10.1093/bioinformatics/bty764. PubMed PMID: 30169550.

39.   Huynh-Thu VA, Irrthum A, Wehenkel L, Geurts P. Inferring regulatory networks

from expression data using tree-based methods. PLoS One. 2010;5(9). doi:

10.1371/journal.pone.0012776. PubMed PMID: 20927193; PubMed Central PMCID:
PMCPMC2946910.

40. Occhipinti A, Verma S, Doan LMT, Angione C. Mechanism-aware and
multimodal AI: beyond model-agnostic interpretation. Trends Cell Biol. 2024;34(2):85-
9. doi: 10.1016/j.tcb.2023.11.002. PubMed PMID: 38087709.

41. Hinkley D, Shi S. Importance sampling and the nested bootstrap. Biometrika.
1989;76(3):435-46.

42. Hao Y, Hao S, Andersen-Nissen E, Mauck WM, 3rd, Zheng S, Butler A, et al.
Integrated analysis of multimodal single-cell data. Cell. 2021;184(13):3573-87 e29. doi:
10.1016/j.cell.2021.04.048. PubMed PMID: 34062119; PubMed Central PMCID:
PMCPMC8238499.

43. Joung J, Ma S, Tay T, Geiger-Schuller KR, Kirchgatterer PC, Verdine VK, et al.
A transcription factor atlas of directed differentiation. Cell. 2023;186(1):209-29 e26. doi:
10.1016/j.cell.2022.11.026. PubMed PMID: 36608654; PubMed Central PMCID:
PMCPMC10344468.

44. Shen WK, Chen SY, Gan ZQ, Zhang YZ, Yue T, Chen MM, et al. AnimalTFDB
4.0: a comprehensive animal transcription factor database updated with variation and
expression annotations. Nucleic Acids Res. 2023;51(D1):D39-D45. doi:
10.1093/nar/gkac907. PubMed PMID: 36268869; PubMed Central PMCID:
PMCPMC9825474.

45. Margolin AA, Nemenman I, Basso K, Wiggins C, Stolovitzky G, Dalla Favera
R, et al. ARACNE: an algorithm for the reconstruction of gene regulatory networks in a
mammalian cellular context. BMC Bioinformatics. 2006;7 Suppl 1(Suppl 1):S7. doi:
10.1186/1471-2105-7-S1-S7. PubMed PMID: 16723010; PubMed Central PMCID:
PMCPMC1810318.

46. Meyer PE, Lafitte F, Bontempi G. minet: A R/Bioconductor package for inferring
large transcriptional networks using mutual information. BMC Bioinformatics.
2008;9:461. doi: 10.1186/1471-2105-9-461. PubMed PMID: 18959772; PubMed Central
PMCID: PMCPMC2630331.

47. de Matos Simoes R, Emmert-Streib F. Influence of statistical estimators of
mutual information and data heterogeneity on the inference of gene regulatory networks.
PLoS One. 2011;6(12):e29279. doi: 10.1371/journal.pone.0029279. PubMed PMID:
22242113; PubMed Central PMCID: PMCPMC3248437.

48. Nacher JC, Akutsu T, editors. Analysis on critical nodes in controlling complex
networks using dominating sets. 2013 International Conference on Signal-Image
Technology & Internet-Based Systems; 2013: IEEE.

49.   Aran D, Looney AP, Liu L, Wu E, Fong V, Hsu A, et al. Reference-based analysis

of lung single-cell sequencing reveals a transitional profibrotic macrophage. Nat

Immunol. 2019;20(2):163-72. doi: 10.1038/s41590-018-0276-y. PubMed PMID:

30643263; PubMed Central PMCID: PMCPMC6340744.

50.   Heinz S, Benner C, Spann N, Bertolino E, Lin YC, Laslo P, et al. Simple

combinations of lineage-determining transcription factors prime cis-regulatory elements

required for macrophage and B cell identities. Mol Cell. 2010;38(4):576-89. doi:

10.1016/j.molcel.2010.05.004. PubMed PMID: 20513432; PubMed Central PMCID:

PMCPMC2898526.

51.   Hubert L, Arabie P. Comparing partitions. Journal of classification. 1985;2:193-

218.

Figure Legends.

Figure 1. The BootCellNet procedure. Starting from scRNA-seq data (top left), k-NN

smoothing is carried out to reduce noise. From the smoothed data, N cells are sampled

and GRNs are computed M times. Parallelly, sampling and GRN construction are

performed $\hat{M}$ times with shuffling gene names K times, resulting in $\hat{M}K$ shuffled

specimens. Using the shuffled data, statistically significant genes and their regulatory

relationships are selected. The resulting GRN is subjected to the computation of MDS. At

the same time, cells in the original scRNA-seq data are clustered into small clumps by

using Louvain clustering with a high resolution. Averaged expression levels of genes in

MDS are calculated and used for the clustering of cell clumps. The resulting clusters are

regarded as cell types.

Figure 2. Smoothing and resampling in BootCellNet are important for the robust

selection of genes and regulations. (A) Schematics of the computer experiment. (B)

Numbers of genes/nodes in GRNs generated 10 times with or without smoothing and with

or without selection. (C) Distribution of number of appearances of edges in 10 specimens,

with (red) or without (gray) resampling-based selection.

Figure 3. BootCellNet robustly identifies MDS. (A) Distribution of the number of the

appearance of genes/nodes in 10 MDS specimens. (B) Numbers of genes in MDS with

(red) or without (gray) selection. (C) Relationship between rank of mean degree and

number of appearances in 10 MDS specimens of each gene.

Figure 4. BootCellNet robustly identifies cell types. (A) Clumps of cells by using

Louvain clustering with a resolution of 10. (B) GRN and its MDS (red squares) identified

by BootCellNet. (C) An example of the heatmap representation of hierarchical clustering

of cell clumps shown in panel A (left) and the resulting clusters (right) based on MDS.

(D) An example of the heatmap of hierarchical clustering of cell clumps (left) and the

resulting clusters (right) based on 100 genes with variable expression. (E) Adjusted Rand

indexes calculated among 10 MDS-based clusterings (left) and 10 variable gene-based

ones (right) were shown.

Figure 5. Cell types in peripheral blood mononuclear cells identified by BootCellNet

and its comparison to reference-based labeling and unsupervised clustering. (A) The

resulting clusters obtained by BootCellNet and their annotations described in the main

text were shown. (B) Annotation of cells obtained by SingleR with the fine

HumanCellAtlas dataset. (C) Clustering result obtained by Louvain clustering

implemented in Seurat, with a resolution of 1.0. (D) Heatmap of average expression of

MDS genes in each cluster obtained by BootCellNet. Color codes for columns are the

same as those in panel A.

Figure 6. BootCellNet facilitates the inference of the course of hematopoiesis. (A)

GRN and its MDS (red squares) identified by BootCellNet. (B) The resulting cell clusters.

(C) Heatmap of average expression of MDS genes in each cluster obtained by

BootCellNet. Color codes for columns are the same as those in panel B. (D) Lineage

curves were obtained by Slingshot using the clusters by BootCellNet. One lineage curve,

which spans from Cluster 1 to Cluster 9 is omitted, since Cluster 9 encompasses lymphoid

cells.

Figure 7. BootCellNet2 and its validation. (A) Outline of the procedure. Upto the GRN

construction, the procedures are the same. From the constructed GRN, the unique set of

control nodes consisting of critical nodes and intermittent nodes was computed. Cell

clumps are obtained in the same manner as BootCellNet. Averaged expression levels of

genes in the unique set are calculated and used for the clustering of cell clumps. Here the

clustering was done with multiscale bootstrap to obtain highly confident clusters. The

resulting high-confident clustering results are compared to clustering with varying

numbers of clusters. The optimal number of clusters is determined as the lowest number

of clusters with maximal similarities. (B) Number of appearances of genes in the unique

set of control nodes or MDS in 10 specimens. No statistically significant difference was

observed by Welch's two-sided *t*-test. (C) Adjusted Rand indexes calculated among 10

automated clusterings with (left) or without (right) using the unique set of control nodes

are shown. The numbers above the heatmaps represent the number of clusters determined

by the BootCellNet2 procedure. (D) Adjusted Rand indexes calculated among 10

automated clusterings with (left) or without (middle) using the unique set of control nodes,

along with those without using either automated clustering and the unique set (right) are

shown. *p < 5x10^-5 by Welch's two-sided *t*-test.

Figure 8. Identification of cell type clusters by BootCellNet2. (A) GRN and its unique

set of control nodes (critical nodes as red squares, intermittent nodes as red circles)

identified by BootCellNet2 are shown. (B) The resulting cell clusters. Annotations for

each cluster described in the main text were shown. (C) Heatmap of average expression

of genes in the unique set in each cluster obtained by BootCellNet2. Color codes for

columns are the same as those in panel B.

Figure 9. BootCellNet2 gives clustering that comply more with CITE-seq results. (A)

The cell clusters obtained by BootCellNet2. Annotations for each cluster according to GO

terms, marker genes, and antibody-derived tags were shown. (B) Heatmap of average

expression of antibody-derived tags in each cluster obtained by BootCellNet2. Color

codes for columns are the same as those in panel A. (C) Heatmap of average expression

of the genes in the unique control set in each cluster obtained by BootCellNet2. Color

codes for columns are the same as those in panel A. (D) The cell clusters obtained by the

Louvain algorithm with a resolution of 0.7. Annotations for each cluster according to GO

terms, marker genes, and antibody-derived tags were shown. (E) Heatmap of average

expression of antibody-derived tags in each cluster obtained by the Louvain algorithm.

Color codes for columns are the same as those in panel D.

Figure 10. Identification of cell types in bronchoalveolar lavage fluid scRNA-seq

data by BootCellNet2. (A) GRN and its MDS (red squares) identified by BootCellNet.

(B) The resulting cell clusters. (C) Heatmap of average expression of MDS genes in each

cluster obtained by BootCellNet. Color codes for columns are the same as those in panel

B. (D) Distribution of cells from each class of patients in each cluster. Statistical

significances of enrichment of cells from each origin were calculated by Fisher's exact

test with Bonferroni correction. Asterisks were shown when p<5x10^-4.

Supplementary Figure 1. GRN reconstructed from CITE-seq data and its unique set of

control nodes (critical nodes as red squares, intermittent nodes as red circles) identified

by BootCellNet2 are shown.

Supplementary Figure 2. The original clustering result of BAL scRNA-seq in Liao

et al. The result was reproduced from the deposited data and is shown (left). Cell type

labeling based on the original paper is shown. As a comparison, the MDS-based clustering

(same as Figure 9B) is also shown.

Supplementary Table 1. Marker genes of each MDS-based cluster of PBMCs. The

output of FindAllMarkers function in Seurat was printed as a table.

Supplementary Table 2. Gene Ontology terms for marker genes of each MDS-based

cluster of PBMCs. Each sheet corresponds to each cluster.

Supplementary Table 3. TFBSs enriched in promters of marker genes of each MDS-

based cluster of PBMCs. Each sheet corresponds to each cluster.

Supplementary Table 4. Marker genes of each MDS-based cluster of cells

undergoing hematopoiesis. The output of FindAllMarkers function in Seurat was

printed as a table.

Supplementary Table 5. Gene Ontology terms for marker genes of each MDS-based

cluster cluster of cells undergoing hematopoiesis. Each sheet corresponds to each

cluster.

Supplementary Table 6. Marker genes of each BootCellNet2 clustering results of

PBMCs. The output of FindAllMarkers function in Seurat was printed as a table.

Supplementary Table 7. Gene Ontology terms for marker genes of PBMCs. Each

sheet corresponds to each cluster.

Supplementary Table 8. Marker genes of each BootCellNet2 clustering results of

PBMCs based on CITE-seq. The output of FindAllMarkers function in Seurat was

printed as a table.

Supplementary Table 9. Gene Ontology terms for marker genes as in Supplementary

Table 8. Each sheet corresponds to each cluster.

Supplementary Table 10. Marker genes of each Louvain clustering results of PBMCs

based on CITE-seq. The output of FindAllMarkers function in Seurat was printed as a

table.

Supplementary Table 11. Gene Ontology terms for marker genes as in

Supplementary Table 10. Each sheet corresponds to each cluster.

Supplementary Table 12. Marker genes of each BootCellNet2 cluster of BAL cells

from COVID-19 patients. The output of FindAllMarkers function in Seurat was printed

as a table.

Supplementary Table 13. Gene Ontology terms for marker genes of BAL cells from

COVID-19 patients. Each sheet corresponds to each cluster.

1002

scRNA-seq data

k-NN smoothing

resampling *N* cells (x *M* times)

selection

resampling (x $\hat{M}$ times) + shuffling (x *K* times)

clustering with high resolution

determine the clusters

Identify MDS

check expression of MDS genes in each clumps

identification of cell types

Figure 1

Figure 2

Figure 3

Figure 4

A

	1
CD8 T	2
2	3
Activated CD8 T	4
8	5
CD8 T	6
7	7
	8
cDC1	9
cDC2	10
monocyte	11
	12
B cell	13

NK

CD4 T

Activated CD4 T

platelet

pDC

B

K_cell:CD56hiCD62L+
NK_cell
NK_cell:IL2

T_cell:CD8+_effector_memory
T_cell:CD4+_Naive
T_cell:CD4+_central_memory
T_cell:CD8+
T_cell:CD4+
T_cell:CD4+_effector_memory

CMP
Monocyte:CD16-Platelets
Pre-B_CMPCD34-
Monocyte:CD16-

B_cell:Plasma_cell
B_cell:Memory
B_cell:immNaive

C

	0
	1
	2
	3
	4
	5
	6
	7
	8
	9
	10
	11
	12
	13

D

row Z-score
-3   3

Figure 5

Figure 6

Figure 7

A

B

critical

intermittent

naive CD4 T

naive CD4 T

naive CD8 T

memory
CD8 T

naive CD4 T

central memory
CD4 T

activated
CD8 T

effector memory
CD4 T

NK

effector memory
CD8 T

cDC1

platelet

pDC

cDC2

monocyte

B cell

C

-2   row Z-score   2

Figure 8

Figure 9

Figure 10

critical

intermittent

Supplementary Figure 1

Original clustering in Liao et al.

BootCellNet2 (same as Figure 10B)

Supplementary Figure 2