Systems Analysis of Immune Changes after B-cell Depletion
in Autoimmune Multiple Sclerosis

Jessica Wei[1,2]*, Jeonghyeon Moon[1,2]*, Yoshiaki Yasumizu[1,2,3], Le Zhang[1], Khadir Radassi[1], Nicholas Buitrago-Pocasangre[1], M. Elizabeth Deerhake[1], Nicolas Strauli[4], Chun-Wei Chen[4], Ann Herman[4], Rosetta Pedotti[5], Catarina Raposo[5], Isaiah Yim[6], Jenna Pappalardo[1,2], Erin E. Longbrake[1], Tomokazu S. Sumida[1], Pierre-Paul Axisa[1,2,7#†], David A. Hafler[1,2,8#†]

Affiliations

1. Department of Neurology, Yale School of Medicine, New Haven, CT, USA
2. Department of Immunobiology, Yale School of Medicine, New Haven, CT, USA
3. Department of Experimental Immunology, Immunology Frontier Research Center, Osaka University, Suita, Osaka, Japan
4. Genentech, San Francisco, CA
5. Roche, Basel, Switzerland
6. Department of Biomedical Engineering, Yale University, New Haven, CT, USA
7. Centre de recherches en cancérologie de Toulouse (CRCT), Inserm, Toulouse, France
8. Broad Institute, Boston, MA, USA

* equal contribution
# equal contribution
† correspondence: pierre-paul.axisa@inserm.fr and david.hafler@yale.edu

Abstract

Multiple sclerosis (MS) is a complex genetically mediated autoimmune disease of the central nervous system where anti-CD20-mediated B cell depletion is remarkably effective in the treatment of early disease. While previous studies investigated the effect of B cell depletion on select immune cell subsets using flow cytometry-based methods, the therapeutic impact on patient immune landscape is unknown. In this study, we explored how a therapy-driven "*in vivo perturbation*" modulates the diverse immune landscape by measuring transcriptomic granularity with single-cell RNA sequencing (scRNAseq). We demonstrate that B cell depletion leads to cell type-specific changes in the abundance and function of CSF macrophages and peripheral blood monocytes. Specifically, a CSF-specific macrophage population with an anti-inflammatory transcriptomic signature and peripheral CD16$^+$ monocytes increased in frequency post-B cell depletion. This was accompanied by increases in TNFα messenger RNA and protein in monocytes post-B cell depletion, consistent with the finding that anti-TNFα treatment exacerbates autoimmune activity in MS. In parallel, B cell depletion induced changes in peripheral CD4$^+$ T cell populations, including increases in the frequency of TIGIT$^+$ regulatory T cells and marked decreases in the frequency of myelin peptide loaded-tetramer binding CD4$^+$ T cells. Collectively, this study provides an exhaustive transcriptomic map of immunological changes, revealing different mechanisms of action contributing to the high efficacy in B cell depletion treatment of MS.

Introduction

While B cell depletion is efficacious in the treatment of various autoimmune diseases including rheumatoid arthritis and type 1 diabetes[1,2], it has conferred remarkable therapeutic benefits in early relapsing remitting multiple sclerosis (MS)[3]. While presumably pathogenic myelin-reactive T cells[4,5] with loss of regulatory T cell (Treg) function[6-8] have been established as mechanistic drivers of MS pathogenesis, accumulating evidence directly implicates B cells as key contributors to loss of immune tolerance. The central role of B cells in MS pathophysiology is substantiated by the infiltration of B cells into MS lesions and CSF[9], the presence of ectopic meningeal B cell follicles adjacent to areas of focal cortical demyelination[10,11], and the detection of oligoclonal IgG bands in the CSF as a diagnostic marker[12,13].

Previous studies investigated the effect of B cell depletion on specific immune subsets using hypothesis driven flow cytometry-based methods[14,15]. In this study, we sought to explore the diverse immune landscape and cell states with "*in vivo* perturbation" in patients with MS by measuring immune phenotypes with scRNAseq transcriptomic granularity. B cell depletion in patients was accomplished with systematic removal of naïve and memory B cells through a humanized anti-CD20 antibody, ocrelizumab. We performed 5' scRNAseq on 18 paired peripheral blood mononuclear cells (PBMC) and paired CSF samples obtained from incident MS patients pre- and post- B cell depletion treatment followed by flow cytometry validation of protein expression. The high dimensional single-cell data set allowed for simultaneous interrogation of the diverse immune populations in patient blood and CSF across tissue types, disease states and treatment status.

Our data revealed increased frequency of a CSF macrophage population that exhibited anti-inflammatory transcriptomic signatures post-B cell depletion therapy. In the periphery, we discovered that CD16[+] monocytes showed the highest upregulation of transcriptomic TNFα/NFκB signatures after ocrelizumab treatment compared to other cell types. The transcriptional changes were confirmed with increases in TNFα protein in monocytes consistent with the finding that anti-TNFα treatment increases MS disease activity. Moreover, changes in the myeloid compartment were accompanied by increases in TIGIT expressing FoxP3[+] Tregs and decreases in the frequency of circulating, myelin reactive CD4[+] T cells. Thus, we defined the immune cell states associated with remission and demonstrate that B cell depletion achieves therapeutic efficacy in early MS through modulating in parallel the innate and adaptive immune compartments.

Results

CD14+CD68+ CSF cells increase in frequency post B cell depletion therapy

All patients were treatment naïve to any immunomodulatory therapy with new onset relapsing

remitting MS (Supplementary Table 1). To elucidate the effects of anti-CD20 treatment on the

CNS microenvironment, we interrogated five CSF samples from people with relapsing remitting

MS (RRMS) pre- and post- B cell depletion treatment and performed an additional comparison

with six CSF samples from age-matched healthy donors. All pre-treatment samples were

collected from treatment-naïve patients, while post-treatment CSF samples were obtained at

different time points: two at 6-month post-treatment, two at 12-month post-treatment, and one at

18-month post-treatment (Supplementary Table 1). After QC of low-quality cells, scRNAseq

yielded 60,704 total single cells from 16 CSF samples, including 15,122 cells from six healthy

donor samples, 28,493 cells from five MS treatment-naïve samples and 17,089 cells from five MS

post- treatment samples (Fig 1a).

After normalization and Harmony batch correction (see Methods), CSF cells were classified

into 17 clusters (Fig 1b). To assess potential donor and sample variability, the frequencies of all

immune populations in the CSF within each patient were profiled. Across all five patients, pre-

treatment MS samples had consistently lower cell frequencies in the CD14+CD68+ myeloid-1 (Mac

1) cluster compared to healthy donors, and the frequencies of this cluster subsequently increased

in all patients after B cell depletion treatment (Fig 1c and Supplementary Fig1). In parallel, from

all five patients, 1536 B cells were found pre-treatment and only 188 B cells were detected post

treatment, as expected due to the mode of action of anti-CD20 therapies. To further identify the

immune subset most affected by B cell depletion therapy, we used MELD[16] to quantify the effect

of B cell depletion treatment on all immune cell clusters in the CSF and discovered that the same

Mac 1 cluster was the most enriched cell type after treatment (Fig 1d). While the dendritic cell

clusters exhibited similar trends of increased frequency post-treatment as the Mac 1 cluster, we

prioritized downstream analyses on the Mac 1 subset as it had the highest MELD enrichment

score and the least MELD score variability across patients.

CD14+CD68+ CSF cells are CSF- specific macrophages with microglia gene signatures

Previous studies reported CD14+ CSF cells in various disease settings[17-19]. We aimed to

further evaluate the myeloid transcriptomic signatures in our patient CSF samples to define the

CD14⁺CD68⁺ myeloid clusters. Mac 1 cluster exhibited high levels of microglia (*TREM2, SPI1,*

CD68, MEF2C) and monocyte (*CD14, FCGR3A, CSF1R, HLA-DRA*) transcriptomic signatures,

while lacking hallmark microglial genes such as *SALL1*, *P2RY12, FCRLS* and *TMEM119*. In

addition to pan-macrophage lineage markers *HLA-DR* and *CD14*, Mac 1 cluster also exhibited

high expression of *APOE, CSF1R*, and genes that are expressed in extra parenchymal CSF

macrophages such as *CST, TGFB1, MS4A7, LYZ, CLEC7A*. The mixture of microglial-like and

monocytic genes and the abundant expression of *C1Q* and *HLA* class II genes classified this

cluster as CSF-specific macrophages, distinct from monocytes, microglia and macrophages in

other CNS tissues (Fig 1e)[20]. Interestingly, memory CD4 cluster 5 exhibited a similar (albeit

muted) signature, leading to the initial co-clustering of those two clusters despite coming from two

separate lineages (Supplementary Fig 2a, b and Methods).

To further delineate the CSF macrophage subset, we computed pan-macrophage (*CD44,*

CCR2, CD45, CD206, CD163, CD274, CD169, MYB), CNS macrophage (*TGFBI, MS4A7,*

MS4A6C, LYZ2, CD163, P2RX7, CST, CLEC7A), and microglia (*P2Y12R, TMEM119, TREM2,*

CD115, CD172A, CD91, SPI1, FCRLS, SALL1, HEXB, SIGLECH, SLC2A5) subset-specific gene

modules to evaluate module scores on all myeloid clusters in the CSF. The post-treatment

enriched Mac 1 cluster scored the highest on the microglia module compared to other myeloid

clusters. In contrast, CD14⁺CD68⁺ myeloid-2 (Mac 2) cluster scored higher on the pan-

macrophage module compared to others (Fig 1f). Collectively, these transcriptomic signatures

reflect the phenotypic diversity of the macrophage compartment in the CSF, and that anti-CD20

treatment increased the frequency of a specific subset of microglia-like macrophages within the

CSF.

B cell depletion induces an anti-inflammatory phenotype in CSF macrophages

With the recent identification of CSF-specific macrophages[20] and the limited availability of

patient CSF samples, the role of CSF macrophages in homeostasis and during MS pathogenesis

remains unclear. To better understand the treatment effect of B cell depletion on CSF

macrophage immunophenotype, we performed differential gene expression (DE) analysis of pre-

and post- treatment MS samples along with healthy donor CSF (Fig 2a). In the enriched Mac 1

cluster, hierarchical clustering on differentially expressed genes in MS pre-treatment vs healthy

donor cells revealed extensive changes, segregating cells from the two groups and highlighting

alterations in MS patients (Supplementary Fig 3a). In contrast, only 82 genes were differentially

expressed between MS pre- and post- treatment CSF macrophages, therefore grouping CSF

macrophages from patients together by hierarchical clustering despite treatment status

(Supplementary Fig 3b). Among the 60 genes upregulated after B cell depletion are *EGR1* that

represses macrophage activation[21], *ZFP36* and *ZFP36L1* that modulate post-transcriptional

regulation of immune responses and oxidative phosphorylation genes (*NDUFA13, UQCRB,*

NDUFB1, ATP5F1E) that are associated with anti-inflammatory macrophages[22,23]. In addition,

genes involved in migration and endocytosis (*SLC11A1, ITGB2, CSF1R, DAB2, NUMB, RAB11A,*

ANXA1, APOE) were downregulated after B cell depletion (Fig 2a).

We observed increases in HLA class I and class II mRNA expression in Mac 1 cells from

baseline MS patient CSF as compared to healthy CSF Mac 1 cells. These increases in MHC

expression from patients with MS decreased after B cell depletion (Fig 2b). In contrast, CD83, a

macrophage immune checkpoint marker that contributes to the resolution of inflammation and

can induce Tregs in experimental models of MS[24-26], was downregulated in MS pre-treatment

cells, and recovered to healthy donor levels post- B cell depletion treatment (Fig 2c). We next

applied the classical and alternative macrophage activation paradigm to delineate the

neuroinflammatory state of CSF macrophages. Post- treatment Mac 1 cells exhibited

transcriptomic downregulation in pro-inflammatory programming (*CCR7, JAK1, STAT1, IL1B,*

TNF, TLR4, CD86) and upregulation of anti-inflammatory genes (*IL-10, TGFB, CLEC7A*) In

addition, we observed decrease of macrophage scavenger receptors (*CD163, MRC1, MSR1*) (Fig

2d). Taken together, the decreases in MHC class I and class II gene module scores suggest that

B cell depletion reduces the probability of T cell activation through CSF macrophage antigen

presentation. The increased expression in *CD83, TGFB, IL10*, and the oxidative phosphorylation

pathway in CSF macrophages post-treatment suggests that B cell depletion therapy promotes the

resolution of inflammatory phenotype in CSF macrophages to restore homeostasis. Lastly, we

computed transcriptomic signature scores based on peripheral monocytic gene modules and

applied them to the CSF myeloid populations. We found that the enriched Mac 1 cluster scored

the highest on the intermediate (*HLADR, CD14, CD11C, CD68, FCGR3A, CX3CR1, CSF1R,*

TLR4) and nonclassical monocyte (*FCGR3A, CX3CR1, SLAN, CSF1R, CXCR1, CXCR4*) gene

modules, whereas Mac 2 scored higher on the classical monocyte (*CD14, CCR2, CCR5, SELL,*

CD36, CD33, CD64) gene module (Fig 2e). Thus, the transcriptomic resemblance of CSF

macrophages to intermediate monocytes prompted us to investigate whether B cell depletion

therapy modulates intermediate monocyte frequency in the periphery.

Increased abundance of CD16+ monocytes post B cell depletion in MS patient PBMC

We next investigated whether alterations observed in the CSF post-B cell depletion are recapitulated in peripheral blood. We performed immune profiling with cryopreserved PBMC using scRNAseq from 18 newly diagnosed treatment-naïve patients with relapsing remitting RRMS whose samples were collected for both pre- and 6 months post-treatment (Supplementary Table 1, Fig 3a). In our unsupervised analysis we retained 38 communities, which we assigned to coarse-grained immune cell types of interest (naïve T cells, memory CD4+ T cells, cytotoxic lymphocytes, B cells, myeloid cells), and 18 fine-grained cell-types, as described in Supplementary Fig 4a, b and Fig 1b. We classified communities into main lineages based on marker gene inspection and scoring cells against reference datasets using singleR software (Supplementary Fig 4c)[27].

Using MELD, we observed computed differences in abundance between the pre- and post-treatment samples. As expected, ocrelizumab treatment significantly decreased the number of B cells except for plasmablasts, which are known to downregulate CD20 that leads to loss of sensitivity to ocrelizumab-mediated depletion (Fig 3b). Given our CSF data, we next focused our attention on myeloid cells. We observed cellular enrichment in two myeloid clusters: CD16+ monocytes and plasmacytoid dendritic cells (pDCs) (Fig 3c). We confirmed these changes by formally testing for variations in frequency across all donors and observed that the increased frequency of CD16+ monocytes was conserved across donors (Fig 3d and Supplementary Fig 5a). The CD16+ monocytes cluster was the only cluster with detectable levels of *FCGR3A*, the gene encoding CD16 (Supplementary Fig 5b and d), and displayed markers associated with intermediate and non-classical monocytes. Scoring against the Monaco Immune reference with singleR revealed strong enrichment in intermediate and non-classical reference transcriptomes (Supplementary Fig 5c). To confirm this change in abundance we measured circulating frequencies of various monocyte subpopulations using flow cytometry (Supplementary Fig 6a). While there was no significant difference after B cell depletion in CD14+CD16- monocytes, there was a significant increase in CD14+CD16+ monocytes with treatment (Fig 3e).

Increased activation and TNFα production in CD16+ monocytes post B cell depletion

We next determined whether CD16+ monocytes harbor an altered transcriptomic state after B cell depletion. We computed differentially expressed genes while controlling for inter-individual variation using a generalized linear mixed model (glmm, as implemented in NEBULA[28].

We observed an increased expression of soluble molecules (*CCL5, CXCL8, TNFA*), surface
receptors (*CD83, ITGB2, ITGA2B*), and NFκB signaling pathway (*TRAF1, NFKB2, REL, RELB,*
NFKBIA) (Fig 4a). CD16$^+$ monocytes showed downregulation in major transcription factors
(*RXRA, IKZF1, KLF4, IRF4*) and CD81, a marker that facilitates monocytes homing to the CNS
in EAE[29]. Given the activated transcriptomic signature observed in CD16$^+$ monocytes, we
validated changes at the protein level and observed downregulation of CD81, and upregulation
of monocyte activation marker HLA-DR using flow cytometry (Fig 4b).

Given the presence of differentially expressed genes from the NFκB signaling pathway, we
next performed GSEA analysis in the enriched CD16$^+$ monocyte cluster using the Hallmark gene
sets collection. Consistent with these findings, we observed enrichment in TNF-NFκB pathway
(Fig 4c), as well as downregulation related to JAK-STAT signaling gene sets (IL-2, STAT5, IL-6,
JAK, STAT3, interferon alpha and interferon gamma signaling) post-treatment. Finally, to
determine whether these pathway modulations can also be observed in CSF macrophages
(described in Figure 2), we created custom gene sets based on PBMC DE and ran GSEA on the
enriched Mac 1 CSF macrophages to test for gene signature enrichment. We observed a positive
enrichment of blood-upregulated genes in the CSF, while blood-downregulated genes showed no
significant enrichment in the CNS (Fig 4d). These data suggest that while B cell depletion therapy
upregulates similar biological signatures in the CNS and periphery, the treatment-mediated *in vivo*
perturbation of biological pathways is tissue-dependent and informed by the local environment.

B cell depletion induces ubiquitous response to TNFα in PBMC
We next assessed whether the observed upregulation of TNF-NFκB pathway post-
treatment is cell-type specific. We computed differentially expressed genes for pre- vs post- B cell
depletion in all clusters and used DE results to run GSEA analysis. We observed TNF-NFκB
pathway activation across most immune cell types post-ocrelizumab treatment (Fig 5a). However,
the downregulation of JAK-STAT related pathways was restricted to CD16$^+$ monocytes, and the
remaining or repopulating B cells post-treatment showed increased expression of various gene
sets related to cell survival (P53 pathway, apoptosis), metabolism (cholesterol homeostasis) and
JAK-STAT signaling (IL2 STAT5 signaling, IL6 JAK STAT3 signaling). Interestingly, clustering
communities based on GSEA leading edge genes similarity showed that the NFκB signaling
enrichment was most similar between B cells and myeloid cells while T lymphocytes formed a

separate cluster, suggesting that transcriptomic responses to NFκB signaling differ between those
lineages (Fig 5b).

Despite the ubiquitous TNF-NFκB pathway activation across cell types, CD16$^+$ monocytes
showed the highest pre-treatment transcriptomic expression of TNFα and sustained the
upregulated expression after treatment (Fig 5c). We also detected an increased expression of
TNFα in B cells cluster 3, MAIT, and DC clusters. To confirm changes in TNFα expression at the
protein level, we enriched CD14$^+$ or CD16$^+$ monocytes from cryopreserved PBMCs and showed
that LPS-stimulated monocytes from MS patients expressed more TNFα post B cell depletion (Fig
5d). Increases of TNFα production by LPS-stimulated CD14$^+$ cells was similarly observed in MS
patients treated with ocrelizumab by another group[30]. However, there was no significant difference
in TNFα ELISA using culture supernatant (data not shown).

B cell depletion therapy reprograms the CD4$^+$ T helper cell compartment in both CSF and
PBMC

Though CD4$^+$ T cells are thought to play a central role in MS pathophysiology, limited
changes were observed in the CD4$^+$ T cell compartment using singleR, MonacoImmune, and
manual cellular annotation methods. Therefore, we applied a recently developed framework that
captures a more granular classification and qualitative assessment of CD4$^+$ T cells based on
scRNAseq data (Fig 6a)[31]. This novel framework assigned CD4 T cells into five major clusters
(cluster Layer 1 or L1) and 18 minor clusters (cluster L2) by Symphony reference mapping (Fig
6b, Supplementary Fig 7a and 8a), and measured the activities of 12 pre-defined transcriptomic
gene programs of CD4$^+$ T cells using non-negative matrix factorization projection (NMFproj).

At the major cluster L1 level, we did not detect any significant cell frequency changes in both
CSF and PBMC (Supplementary Fig 7b and 8b). However, at the minor cluster L2 level, there
was a significant reduction of CD4$^+$ T effector memory (Tem) expressing T peripheral helper (Tph)
markers (Tem -Tph; *PDCD1^{lo}CXCR5$^+$*) in both CSF (padj= 1.88 x 10^{-2}) and PBMC (padj= 9.41x10^{-6}
^{6}) tissues after B cell depletion treatment (Fig 6c and d, Supplementary Fig 7c and 8c). In CSF
alone, the frequency of CD4$^+$ T central memory (Tcm) expressing T follicular helper (Tfh) markers
(Tcm -Tfh; *PDCD1$^+$CXCR5$^+$*) was significantly reduced (padj=9.36x10^{-3}) while the frequency of
Tcm-Th0 was increased (padj=0.0472), suggesting a shift toward a naïve phenotype post- B cell
depletion in the CNS (Fig 6c and d, Supplementary Fig 7c and 8c). We also observed a significant
increase of CD4$^+$ naive T cells expressing *SOX4* (Tnaive *SOX4*; *SOX4$^+$PECAM1$^+$*) (padj=4.29x10^{-3}
^{3}) in the blood, which is a recent thymic emigrant population[32], indicating that the peripheral blood

CD4[+] T cell pool has been replenished by newly generated CD4[+] T cells after the treatment (Fig
6d, Supplementary Fig 8c).

We then assessed the changes in gene program activity quantified by NMFproj in each L2
subpopulation. We observed a significant reduction in cell types for NMF6 (Tfh-Feature or Tfh-F;
MAF, *CXCR5*) and NMF11 (Th1-F; *GZMK*, *EOMES*) post-treatment in both blood and CSF. (Fig
6e and f, Supplementary Data 1 and 2). Because NMF6 (Tfh-F) is predominantly high in Tcm-Tfh,
Tem-Tph, and intermediate Treg (Treg Int), the decrease of NMF6 (Tfh-F) in both tissues indicates
that B cell depletion treatment reduces their frequencies and potentially represses the
repopulation of these CD4 subtypes quantitatively and qualitatively. An increase of NMF2 (Th17-
F; *RORC*, *CCR6*) in the Tem population was observed in CSF, whereas NMF2 signature
decreased in the blood (Fig 6e and f, Supplementary Data 1 and 2). We also observed increased
NMF10 (Tissue-F; *JUNB*, *NFKBIA*) in blood after treatment (Fig. 6f, Supplementary Data 2).
These observations suggested a redistribution of the CD4[+] T subsets between the periphery and
CNS. Altogether, these findings demonstrate that B cell depletion notably alters the CD4[+] T cell
compartment by reducing specific T cell populations such as Treg Int, Tcm-Tfh, and Tem-Tph and
modifying effector gene expression profiles such as repression of NMF6 (Tfh-F) and NMF11 (Th1-
F), which may be associated with its therapeutic efficacy in MS.

B cell depletion increases suppressive Tregs
Loss of Treg function has been repeatedly observed in patients with MS[8,33], and we showed
MS susceptibility variant modulate can modulate Treg function[34]. We sub-clustered Tregs from
scRNAseq PBMC data and compared L2 subpopulation frequencies to examine Treg alterations
in more detail. We observed a significant decrease in naïve Tregs (Treg Naive) (*FOXP3, CCR7*,
padj= 3.13×10^{-2}) and Treg Int (*FOXP3*, *FCRL3*, padj=6.79×10^{-3}) and an increase in effector Treg
(Treg Eff) (*HLA-DR*s, *CD74*, padj=2.33×10^{-3}) in post-treatment samples. (Fig 7a and b). We also
examined gene expression differences in the whole Treg population and found that *HLA-DR*s and
CD74, which are the markers of Treg Eff, were increased. In contrast, *FCRL3*, a marker of Treg
Int, was decreased after treatment (Fig 7c, Supplementary Fig 9a and b). These data suggest
that B cell depletion skewed the Treg function toward the effector phenotype. Next, we examined
the potential mediators of myeloid-Treg interactions using ligand-receptor prediction analysis (Fig
7d). Since TNF is known to enhance the function of Tregs through interaction with TNFR2, which
is abundant in Tregs[35,36], we hypothesized that monocytes upregulated *TNF* expression following
B cell depletion and promoted Treg expansion through TNFR2 signaling. We measured Treg

frequency using flow cytometry in MS patient PBMC pre- and post- ocrelizumab treatment. We observed a significant increase in the frequency (p < 0.0001) of Tregs (Fig 7e) in post-treatment PBMC samples. TIGIT protein is highly expressed in Treg Eff[37] (Supplementary Fig 9b and c) and has been shown to associate with increases in functional activity in human and mice[38,39]. Therefore, we measured the frequency of TIGIT expressing Tregs after B cell depletion and observed a significant increase post- ocrelizumab treatment (Fig 7f). In summary, these data indicate that B cell depletion corrects for the loss of immune regulation in MS by increasing Treg frequency and effector function.

B cell depletion decreases myelin tetramer binding CD4$^+$ T cells

We and others have shown increases in the frequency of inflammatory myelin reactive T cells recognizing a number of myelin antigens, presumably as a consequence of epitope spreading, in the circulation of MS patients. We have demonstrated the utility of using a panel of MHC class II tetramers loaded with myelin epitopes from myelin basic protein (MBP), proteolipid protein (PLP), and myelin oligodendrocyte glycoprotein (MOG) to directly measure the frequency of autoreactive T cells in the disease[4,5,40,41]. To increase the accuracy of detecting autoreactive T cells, we used both PE- and APC-conjugated tetramers with the same myelin composition (Supplementary Fig 6b). We initially examined a cohort of patients pre- and post-B cell depletion and observed a marked decrease in the frequency of myelin PE- and APC-tetramer double positive (Tetramer DP) CD4$^+$ T cells (Fig 7g). In these data, decreased frequencies of CD45RA-CXCR5$^+$ cells and CCR6$^+$CXCR3$^-$ cells were observed in myelin tetramer binding CD4$^+$ T cells (Fig 7h). Longitudinal kinetic analysis using fresh PBMC derived from a separate MS cohort (Supplementary Table 1) revealed significant decreases in the frequency of these autoreactive CCR6$^+$CXCR3$^+$ CD4$^+$ T cells at 6 months post-treatment with no significant changes in CCR6$^+$CXCR3$^+$ cells in non-tetramer reactive CD4$^+$ T cells. This trend continued through 52 weeks (Fig 7i). These results indicate that B cell depletion directly or indirectly regulates the frequency of presumably pathogenic myelin reactive T cells.

Discussion

We performed an unbiased analysis of the immune landscape using scRNAseq after *in*
vivo perturbation of the immune system with B cell depletion therapy in treatment-naïve MS
patients. Ocrelizumab, a humanized anti-CD20 monoclonal antibody, induces a systematic
removal of naïve and memory B cells with ~98% efficacy in preventing new CNS lesions[3]. Here,
we demonstrate that B cell depletion induces cell type-specific recovery of CSF macrophages
and intermediate monocytes and reprograms them toward transcriptional phenotypes that
resemble healthy donor macrophage and monocytes. Specifically, we identified a novel anti-
inflammatory transcriptomic signature in post-treatment MS CSF macrophages, promoting the
resolution of inflammation and maintain homeostasis with upregulation of CD83, TGFβ, and IL-
10. CD16$^+$ monocytes showed the highest level of TNFα messenger RNA expression pre-
treatment, and B cell depletion further increased TNFα protein expression in these monocytes.
Furthermore, after B cell depletion, we observed shifts in T cell populations, including decreased
frequencies of Tem-Tph and Treg Int, along with Th1- and Tfh-type gene programs. Notably, we
found that B cell depletion increased TIGIT$^+$ regulatory T cell frequency and decreased myelin
tetramer-binding CD4$^+$ T cell frequency in the blood compartment. Our study provides an
extensive transcriptomic map of immunological changes through the simultaneous interrogation
of the diverse immune populations perturbed by *in vivo* B cell depletion. This approach enables
us to discover distinct cell-type specific mechanisms of action consistent with the remarkable
efficacy of B cell depletion therapy in MS.

The changes in monocyte frequency after B cell depletion were unexpected and revealed
the power of an unbiased systems approach in dissecting the multitudes of immune-regulatory
mechanisms of a highly efficacious therapy. We hypothesize that the dichotomous roles of
monocytes in the pathogenesis of MS are due to the nature in which myeloid phenotype and
function are orchestrated by the metabolic requirements in surrounding tissue environments.
Soluble factors and cytokines shape macrophage differentiation and activate molecular programs
that either exacerbate or attenuate disease[42]. Systemic depletion of pathogenic B cells likely
reduces the inflammatory states of multiple immune cell types in a feedback loop manner, and
the subsequent recalibration of the cytokine milieu allows macrophages to differentiate in a
steady-state environment and engage in disease-resolving transcription program to restore
homeostasis. Given our study is limited to one post-treatment timepoint, it is possible that the
changes we describe will normalize on the long term. Our ability to detect functional changes in

the CSF compartment were limited by the small sample size, variable post-treatment time-points,

and a small number of cells composing the CSF macrophages clusters compared to other

immune populations. However, a previous study by Mrdjen et al. dissected changes in myeloid

populations in the EAE model and observed CNS-specific alterations of macrophages and

microglia phenotypes during the disease course[43]. Interestingly, they noted that border-

associated macrophages (BAMs) share similar profiles with pDCs compared to other myeloid

subsets. We similarly observed that increased abundance of CSF macrophages was

accompanied by increased abundance of pDCs (Fig 1).

While the ontogeny of CSF macrophages has not been well elucidated, we posit that some

of the enriched macrophage population could derive from blood monocytes as we observed some

similarity in transcriptomic kinetics between enriched CSF macrophages and enriched CD16+

monocytes post- B cell depletion. With autoimmune exacerbations in patients with MS, peripheral

blood monocytes receive inflammatory cytokine signals and cross the blood- brain barrier with T

cells leading to CNS lesions[44,45]. While macrophages are the dominant cell type in these lesions,

blood monocytes also extravasate into the CSF and differentiate into CSF macrophages[46].

However, it is critical to not hyperactivate CSF macrophages in the process of phagocytosing

waste to prevent downstream immune activation of autoreactive lymphocytes[47]. With B cell

depletion, the decrease in myeloid inflammatory cytokines enables CSF macrophages to receive

pro-resolving signals to execute their homeostatic function in mediating tissue repair[48]. Our data

suggest an anti-inflammatory phenotype in CSF macrophage after B cell depletion treatment and

demonstrate that anti-CD20 depletion therapy restores macrophage gene signatures similar to

those of healthy controls. Importantly, the inclusion of healthy donor CSF samples allows us to

conclude that anti-CD20 treatment reprograms CSF macrophages toward a homeostatic/healthy

cell state.

In the peripheral immune compartment, we uncovered a ubiquitous TNFα/NFκB activation

signature across a wide range of circulating immune cells post- B cell depletion. CD14$^+$CD16$^+$

monocytes have been shown to be a potent producer of TNFα[49], and our study suggests that TNF

is a pleiotropic cytokine that potentially exerts anti-inflammatory effects after B cell depletion. It

has not escaped our attention that B cell depletion confers moderate clinical effectiveness in

treating rheumatoid arthritis whereas anti-TNFα clearly worsen MS immunopathogenesis[50]. While

the increased production of TNFα with a clinically effective treatment appears to be counter-

intuitive, the beneficial role of TNFα in MS is substantiated by clinical trials results showing anti-

TNFα treatment in patients led to significant worsening of disease activity[51]. Moreover, molecular

dissection of MS risk allele rs1800693 located in the gene encoding the TNFR1 revealed the

associated variant that codes for a soluble form of TNFR1, which mimics TNF blocking

molecules[52,53], again consistent with the observation that TNFα blockade leads to increased

disease activity. In the context of chronic inflammation, this scenario where TNFα bears an anti-

inflammatory role is reminiscent of the anti-tumor immune response. Specifically, in the tumor

microenvironment, constant exposure to TNFα leads to immunosuppressive responses involving

Tregs, B regulatory cells and myeloid-derived suppressor cells[54], and blocking TNF leads to

improved response to immune checkpoint blockade in an orthotopic melanoma mouse model[55].

Additionally, identifying the precise signaling events leading to the observed transcriptomic

changes is challenging given that the main TNFα signaling pathway is through NFκB, a highly

ubiquitous signaling pathway. Thus, we cannot exclude other receptors signaling through NFκB

are participating to the transcriptomic alterations observed post-treatment. Nevertheless, our rich

dataset provides a non-biased road map which can now be investigated in animal models to better

elucidate the more detailed mechanisms associated with B cell depletion.

Several of the established cell annotation methods that we employed were unable to

detect changes in the CD4 T cell compartment. The biologically relevant signals could potentially

be obfuscated due to the small frequency of myelin-specific T cells. Applying reference mapping

with finer granular reference and gene program quantification by NMF, we were able to detect B

cell depletion- mediated modulation of various T cell subsets. While we were able to confirm the

increase of effector Treg frequency both by transcriptomic and protein expression, it will be of

interest to more precisely identify whether there are differences in specific myelin antigens and

other antigens, as recent studies indicate EBV infection of B cells is associated with the onset of

MS[56].

Our T cell data suggests several mechanisms in which B cell depletion can lead to

modulation of T cell functionality. One potential model is that TNFα from myeloid cells engages

TNFR2 on Tregs leading to suppression of autoreactive T cells after B cell depletion in RRMS

patients. Alternatively, as first shown by Lanzavecchia et al.[57], B cells may be the key antigen

presenting cell and their depletion may result in loss of autoreactive effector or memory T cells.

Similarly, the decrease of MHC expression on myeloid cells may also support this hypothesis.

Rather than identifying one mechanism for the high clinical efficacy of B cell depletion in MS, our

unbiased "*in vivo* perturbation" investigation uncovered a number of changes in the immune

circuitry. This is perhaps not surprising as the genetic architecture of MS and other autoimmune

diseases suggests that multiple pathways are involved in disease pathogenesis[58]. In summary,

our unbiased systems analysis identified a series of immunoregulatory pathways induced by B

cell depletion. We propose that for B cell depletion to be such a globally effective therapy in MS,

essentially subverting the autoimmune attack of the CNS, several mechanisms of action must be

involved to maintain homeostasis. It is likely that different immunosuppressive pathways become

activated among patients, leading to marked decreases of autoreactive myelin reactive T cells in

the blood compartment. Our analyses comparing immune cells in CSF and blood also highlights

shared vs distinct changes across compartments, suggesting regulation of CNS homing

mechanisms is affected by anti-CD20 therapies. Considering the potential of disease-mediating

and homeostatic functions in the myeloid compartment, future analyses can be designed with a

myeloid focus using fresh tissue for higher sensitivity in protein detection. Nevertheless, these

datasets and observations provide a critical starting point that will require well-designed *in vitro*

and *ex vivo* assays and appropriate animal models to fully elucidate the perturbational effects of

B cell depletion on the functionality of the immune system.

Methods

Ethic Statement

This study was approved by the Institutional Review Board at Yale University. CSF and blood samples were obtained from healthy donors and MS patients with informed consent.

Patient Cohorts

All patients had early onset relapsing remitting MS and had not been on previous immunomodulatory treatments. A small subset of patients had received IV solumedrol within 3 months of blood draw. Eighteen patients undergoing single cell RNA seq studies had CSF analysis prior to the initiation of treatment, and five of those subjects had repeat lumbar punctures, as outlined in the results section. A total of six age-matched healthy controls had lumbar punctures, and those results were previously reported[44]. An additional four subjects had flow cytometric analysis only. Patient characteristics are summarized in Supplementary Table 1.

Sample preparation for scRNAseq

Fresh patient CSF samples were centrifuged, and cells were immediately processed using 10x Genomics 5Pv1 chemistry. Samples were collected prior to infusion of B cell depletion therapy. In the CSF sample cohort, four of the patients were administered ocrelizumab B cell depletion treatment, and one was treated with rituximab. Patient PBMCs were isolated from whole blood using Lymphoprep (STEMCELL) density gradient centrifugation. All patients were administered ocrelizumab B cell depletion in the PBMC cohort. Cryopreserved patient-matched pre-treatement and 6-month post-treatment PBMCs were thawed and processed within the same experimental batch using 10x Genomics 5Pv1 chemistry. For PBMCs, TCR libraries were generated along with the gene expression libraries.

scRNAseq QC

PBMC and CSF libraries were sequenced at 20,000 read pairs per cell on Illumina NovaSeq instrument. Fastq files were processed using cellranger version 3.1.0 mapping to GRCh38 human reference genome. Alignment and quantitation were performed with the "cellranger count" command for each emulsion (using the 2020-A 10x genomics human reference), to generate unique molecular identifier (UMI) count matrices.

For CSF, Data QC was performed in R using the Seurat package. Low quality cells were
filtered out based on mitochondria percent, UMI counts and number of features for individual
samples. Samples were then merged, log10-transformed, and batch corrected using Harmony.

For PBMCs, we first filtered extreme outliers by excluding droplets with less than 1500
UMI counts, or less than 850 unique genes detected. As distribution of those parameters varied
across emulsions, we median-centered the log10-transformed number of unique genes detected
and removed low quality droplets with less than 1,100 unique genes detected or more than 2.5%
mapping of UMI counts mapping to mitochondrial genes. We also removed potential doublets by
filtering out droplets with more than 2,600 unique genes detected.

scRNAseq analysis
Dimensionality reduction and clustering
For cells passing quality control, we normalized UMI counts by dividing each count by the
total number of counts per cell. We then multiplied normalized counts by 10,000 and added a
pseudo count of 1 before log-transformation. We computed the stabilized variance of each gene
using the variance-stabilizing transformation (VST) and retained genes with stabilized variance >
1 for principal component analysis (PCA). Genes mapping to the T cell receptor (TCR), the B cell
receptor (BCR) and the Y chromosome were excluded from PCA analysis. We computed the first
50 principal components (PCs) using a partial singular value decomposition method, based on
the implicitly restarted Lanczos bidiagonalization algorithm (IRLBA), as implemented in the *Seurat*
R package[59]. To correct for systematic differences across samples, we applied harmony
integration[60] to the first 50 PC loadings and retained 30 harmony-corrected PCs to build nearest
neighbor graphs for visualization using Uniform Manifold Approximation and Projection (UMAP)
(minimum distance = 0.5, spread = 10), and community detection using Louvain algorithm, as
implemented in *Seurat*. We also computed a relative likelihood of cells being observed in specific
experimental conditions using Manifold Enhancement of Latent Dimensions (MELD)[16].

We embedded cells into 2 UMAP dimensions and applied Louvain algorithm. We
annotated cluster cell types based on individual gene expression and the SingleR automatic
annotation package using the MonacoImmuneData PBMC reference[61]. We tested for variation in
clusters frequency in the CSF and in the blood separately by modelling the per-sample cluster
frequencies using a beta distribution in a generalized linear model framework, as implemented in
the *betareg* R package[62].

Differential gene expression (DE)

For DE testing at the single cell level in CSF, we tested each gene individually using a
generalized linear model approach as implemented in the *speedglm* R package, using a poisson
distribution and a log link function[34]. We excluded low expression genes based on a UMI count
per cell < 0.005, as well as ribosomal, mitochondrial, TCR, BCR. We evaluated differences in
counts with treatment predictor, and a [number of UMIs detected] covariate to account for
differences in library size. We then computed shrunk log fold changes using adaptative shrinkage
methods implemented in the *ashr* R package[63] (using a mixture of normal distributions) and FDR
(Benjamini & Hochberg method) across all genes tested. Genes with FDR < 0.05 were considered
differentially expressed. For differential expression testing at the single cell level in PBMCs, we
used a negative binomial mixed linear model as implemented in the NEBULA package[28]. We then
used shrunken log fold changes as a ranking metrics to run geneset enrichment analysis (GSEA).

Treg volcano and ligand-receptor analyses
Volcano plot displaying differential expression analysis performed using *nebula* comparing pre-
and post-treatment Treg population. Among genes with differential expression (BH < 0.05) and
average expression >0.1, those that encode surface proteins (based on Cell Surface Protein Atlas
surfaceome protein database) were selectively labeled[64]. NicheNet (*nichenetr*)[65] was used to
identify predicted ligand-receptor interactions between myeloid populations and Tregs, will a
particular focus on potential ligands that are differentially regulated in myeloid cells with B cell
depletion treatment. Tregs were selected as the "receiver cell type", including all expressed genes
as potential receptors. Myeloid cells were selected as the "sender cell type", limiting the set of
potential ligands to the combined list of genes differentially expressed with B cell depletion
treatment in myeloid cell clusters (see *NEBULA* analysis). Predicted ligand-receptor interactions
were displayed as a heat map in which ligands (expressed by myeloid cells) were plotted against
receptors (expressed by Tregs) and weighted by prior interaction potential.

CD4$^+$ T cell automatic labeling and quantification of gene programs
The PBMC and CSF data were processed using the pipeline developed in the previous study[31] to
assign CD4$^+$ T cell clusters. This pipeline employs Azimuth[37] for the extraction of CD4$^+$ T cells and
uses Symphony[66] for predicting CD4$^+$ T cell clusters. For interpretability, 'Treg Act' has been
renamed to 'Treg Int' from the original literature. We tested for variation in cluster frequency by

modeling the per-sample cluster frequencies using a beta distribution in a generalized linear

model framework, as implemented in the *betareg* R package. For the assessment of TIGIT protein

expression, we used CITE-seq data from PBMCs deposited in GSE164378 and performed

reference mapping using the pipeline. Additionally, a 12-dimensional qualitative evaluation was

conducted on the extracted CD4[+] T cells using NMFproj[67]. We applied a generalized linear model

to assess feature changes per cluster[31].

Flow cytometry analysis

Frozen PBMCs were used for flow cytometry validation, except longitudinal myelin tetramer

staining were performed on fresh PBMCs (n=4). Patient peripheral blood mononuclear cells were

stained with a ViaKrome 808 Fixable Viability dye following the manufacturer's instructions. Cells

were then labeled with surface antibodies for 30 min at 4°C. For intracellular staining, cells were

fixed and permeabilized with BD Cytofix/Cytoperm Buffer (BD Biosciences) for 10 min at room

temperature, then washed with phosphate-buffered saline. Intracellular proteins were stained in

permeabilization buffer (eBioscience) for 30 min at 4 °C. Antibody details are provided in

Supplementary Table 2. For TNF staining, monocytes were enriched from cryopreserved PBMCs

using EasySep™ Human Monocyte Enrichment Kit without CD16 Depletion Kit (STEMCELL

technologies). Enriched monocytes were stimulated with 100ng/ml LPS for 4h at 37 °C before

staining. To investigate myelin tetramer reactive T cells, APC- or PE- conjugated tetramers which

were composed by DRB1*15:01 (loaded with MBP, MOG and PLP) or DRB1*04:01 (loaded with

MOG and PLP) were used[4,41]. Myelin tetramers were incubated with cells for 30 min at 37°C

before staining with antibodies. Cells were acquired on a BD Symphony flow cytometer with

FACSDiva (BD Pharmingen) and data were analyzed with FlowJo software v.10 (Treestar).

Peptide loading

Biotinylated monomers were diluted to a concentration of 0.5 mg/mL with 0.1 M phosphate buffer

and incubated with 0.4 mg/ml of at 37°C for 72 h in the presence of 2.5 mg/ml n-Octyl β-D-

glucopyranoside (OG) and 1 mM Pefabloc SC (Sigma–Aldrich, St. Louis, MO). Peptide loaded

monomers were subsequently conjugated into tetramers using R-PE streptavidin (ThermoFisher

Scientific, Waltham, MA) or fluorochromes of interest at a molar ratio of 8:1. Myelin peptide

sequences are listed in the Supplementary Table 3.

Data and code availability statement:

All raw scRNAseq data generated in this study will be deposited on dbGAP upon acceptance. All code used for genomics analysis will be available on github and figshare, along with intermediate analysis files upon acceptance.

Acknowledgement:

The authors thank Rahul Dhodapkar for helpful discussions on computational data analysis. Chuan He for technical support and Kathryn Miller-Jensen for experimental input and discussion. This work was supported by NIH grants (P01 AI073748, U19 AI089992 U24 AI11867, R01 AI22220, UM 1HG009390, P01 AI039671, P50 CA121974, R01 CA227473) and Race to Erase MS to D.A.H. and National MS Society to D.A.H. and M.R.L. and grants from Genentech and F. Hoffmann-La Roche Ltd as part of Integrative Neuroscience Collaborations Network.

Declaration of Interests

D.A.H. has received research funding from Bristol-Myers Squibb, Novartis, Sanofi, and Genentech. He has been a consultant for Bayer Pharmaceuticals, Repertoire Inc, Bristol Myers Squibb, Compass Therapeutics, EMD Serono, Genentech, Juno therapeutics, Novartis Pharmaceuticals, Proclara Biosciences, Sage Therapeutics, and Sanofi Genzyme. E.E.L. has received research support from Biogen, LabCorp, Intus and Genentech. She has been a consultant for Bristol Myers Squibb, EMD Serono, Genentech, Sanofi Genzyme and NGM Bio.

References

1 Lee, D. S. W., Rojas, O. L. & Gommerman, J. L. B cell depletion therapies in autoimmune
disease: advances and mechanistic insights. *Nat Rev Drug Discov* **20**, 179-199,
doi:10.1038/s41573-020-00092-2 (2021).

2 Pescovitz, M. D. *et al.* Rituximab, B-lymphocyte depletion, and preservation of beta-cell
function. *N Engl J Med* **361**, 2143-2152, doi:10.1056/NEJMoa0904452 (2009).

3 Hauser, S. L. *et al.* Ocrelizumab versus Interferon Beta-1a in Relapsing Multiple Sclerosis.
N Engl J Med **376**, 221-234, doi:10.1056/NEJMoa1601277 (2017).

4 Cao, Y. *et al.* Functional inflammatory profiles distinguish myelin-reactive T cells from
patients with multiple sclerosis. *Sci Transl Med* **7**, 287ra274,
doi:10.1126/scitranslmed.aaa8038 (2015).

5 Ota, K. *et al.* T-cell recognition of an immunodominant myelin basic protein epitope in
multiple sclerosis. *Nature* **346**, 183-187, doi:10.1038/346183a0 (1990).

6 Dominguez-Villar, M., Baecher-Allan, C. M. & Hafler, D. A. Identification of T helper type
1-like, Foxp3+ regulatory T cells in human autoimmune disease. *Nat Med* **17**, 673-675,
doi:10.1038/nm.2389 (2011).

7 Sumida, T. *et al.* Activated beta-catenin in Foxp3(+) regulatory T cells links inflammatory
environments to autoimmunity. *Nat Immunol* **19**, 1391-1402, doi:10.1038/s41590-018-
0236-6 (2018).

8 Viglietta, V., Baecher-Allan, C., Weiner, H. L. & Hafler, D. A. Loss of functional suppression
by CD4+CD25+ regulatory T cells in patients with multiple sclerosis. *J Exp Med* **199**, 971-
979, doi:10.1084/jem.20031579 (2004).

9 Jain, R. W. & Yong, V. W. B cells in central nervous system disease: diversity, locations and
pathophysiology. *Nat Rev Immunol* **22**, 513-524, doi:10.1038/s41577-021-00652-6
(2022).

10 Lucchinetti, C. F. *et al.* Inflammatory cortical demyelination in early multiple sclerosis. *N*
Engl J Med **365**, 2188-2197, doi:10.1056/NEJMoa1100648 (2011).

11 Magliozzi, R. *et al.* Meningeal B-cell follicles in secondary progressive multiple sclerosis
associate with early onset of disease and severe cortical pathology. *Brain* **130**, 1089-1104,
doi:10.1093/brain/awm038 (2007).

12 Comi, G. *et al.* Role of B Cells in Multiple Sclerosis and Related Disorders. *Ann Neurol* **89**,
13-23, doi:10.1002/ana.25927 (2021).

13 Reich, D. S., Lucchinetti, C. F. & Calabresi, P. A. Multiple Sclerosis. *N Engl J Med* **378**, 169-
180, doi:10.1056/NEJMra1401483 (2018).

14 Nissimov, N. *et al.* B cells reappear less mature and more activated after their anti-CD20-
mediated depletion in multiple sclerosis. *Proc Natl Acad Sci U S A* **117**, 25690-25699,
doi:10.1073/pnas.2012249117 (2020).

15 von Essen, M. R. *et al.* Ofatumumab Modulates Inflammatory T Cell Responses and
Migratory Potential in Patients With Multiple Sclerosis. *Neurol Neuroimmunol*
Neuroinflamm **9**, doi:10.1212/NXI.0000000000200004 (2022).

16 Burkhardt, D. B. *et al.* Quantifying the effect of experimental perturbations at single-cell
resolution. *Nat Biotechnol* **39**, 619-629, doi:10.1038/s41587-020-00803-5 (2021).

17   Esaulova, E. *et al.* Single-cell RNA-seq analysis of human CSF microglia and myeloid cells
in   neuroinflammation.   *Neurol   Neuroimmunol   Neuroinflamm*   **7**,
doi:10.1212/NXI.0000000000000732 (2020).
18   Farhadian, S. F. *et al.* Single-cell RNA sequencing reveals microglia-like cells in
cerebrospinal   fluid   during   virologically   suppressed   HIV.   *JCI   Insight*   **3**,
doi:10.1172/jci.insight.121718 (2018).
19   Ramesh, A. *et al.* A pathogenic and clonally expanded B cell transcriptome in active
multiple   sclerosis.   *Proc   Natl   Acad   Sci   U   S   A*   **117**,   22932-22943,
doi:10.1073/pnas.2008523117 (2020).
20   Munro, D. A. D., Movahedi, K. & Priller, J. Macrophage compartmentalization in the brain
and   cerebrospinal   fluid   system.   *Sci   Immunol*   **7**,   eabk0391,
doi:10.1126/sciimmunol.abk0391 (2022).
21   Trizzino, M. *et al.* EGR1 is a gatekeeper of inflammatory enhancers in human
macrophages. *Sci Adv* **7**, doi:10.1126/sciadv.aaz8836 (2021).
22   Ip, W. K. E., Hoshi, N., Shouval, D. S., Snapper, S. & Medzhitov, R. Anti-inflammatory effect
of IL-10 mediated by metabolic reprogramming of macrophages. *Science* **356**, 513-519,
doi:10.1126/science.aal3535 (2017).
23   Wculek, S. K. *et al.* Oxidative phosphorylation selectively orchestrates tissue macrophage
homeostasis. *Immunity* **56**, 516-530 e519, doi:10.1016/j.immuni.2023.01.011 (2023).
24   Grosche, L. *et al.* The CD83 Molecule - An Important Immune Checkpoint. *Front Immunol*
11, 721, doi:10.3389/fimmu.2020.00721 (2020).
25   Burke, J. R. *et al.* Autoimmune pathways in mice and humans are blocked by
pharmacological stabilization of the TYK2 pseudokinase domain. *Sci. Transl. Med.* **11**,
doi:10.1126/scitranslmed.aaw1736 (2019).
26   Peckert-Maier, K. *et al.* CD83 expressed by macrophages is an important immune
checkpoint molecule for the resolution of inflammation. *Front Immunol* **14**, 1085742,
doi:10.3389/fimmu.2023.1085742 (2023).
27   Aran, D. *et al.* Reference-based analysis of lung single-cell sequencing reveals a
transitional profibrotic macrophage. *Nat Immunol* **20**, 163-172, doi:10.1038/s41590-018-
0276-y (2019).
28   He, L. *et al.* NEBULA is a fast negative binomial mixed model for differential or co-
expression analysis of large-scale multi-subject single-cell data. *Commun Biol* **4**, 629,
doi:10.1038/s42003-021-02146-6 (2021).
29   Dijkstra, S. *et al.* Targeting the tetraspanin CD81 blocks monocyte transmigration and
ameliorates EAE. *Neurobiol Dis* **31**, 413-421, doi:10.1016/j.nbd.2008.05.018 (2008).
30   Geladaris, A. *et al.* IL-10-providing B cells govern pro-inflammatory activity of
macrophages and microglia in CNS autoimmunity. *Acta Neuropathol* **145**, 461-477,
doi:10.1007/s00401-023-02552-6 (2023).
31   Yasumizu, Y. *et al.* Single-cell transcriptome landscape of circulating CD4(+) T cell
populations   in   autoimmune   diseases.   *Cell   Genom*   **4**,   100473,
doi:10.1016/j.xgen.2023.100473 (2024).
32   Kimmig, S. *et al.* Two subsets of naive T helper cells with distinct T cell receptor excision
circle   content   in   human   adult   peripheral   blood.   *J   Exp   Med*   **195**,   789-794,
doi:10.1084/jem.20011756 (2002).

33   Dominguez-Villar, M. & Hafler, D. A. Regulatory T cells in autoimmune disease. *Nat*
Immunol **19**, 665-673, doi:10.1038/s41590-018-0120-4 (2018).
34   Axisa, P. P. *et al.* A multiple sclerosis-protective coding variant reveals an essential role
for HDAC7 in regulatory T cells. *Sci Transl Med* **14**, eabl3651,
doi:10.1126/scitranslmed.abl3651 (2022).
35   Chen, X. *et al.* TNFR2 is critical for the stabilization of the CD4+Foxp3+ regulatory T. cell
phenotype in the inflammatory environment. *J Immunol* **190**, 1076-1084,
doi:10.4049/jimmunol.1202659 (2013).
36   Torrey, H. *et al.* A novel TNFR2 agonist antibody expands highly potent regulatory T cells.
Sci Signal **13**, doi:10.1126/scisignal.aba9600 (2020).
37   Hao, Y. *et al.* Integrated analysis of multimodal single-cell data. *Cell* **184**, 3573-3587
e3529, doi:10.1016/j.cell.2021.04.048 (2021).
38   Joller, N. *et al.* Treg cells expressing the coinhibitory molecule TIGIT selectively inhibit
proinflammatory Th1 and Th17 cell responses. *Immunity* **40**, 569-581,
doi:10.1016/j.immuni.2014.02.012 (2014).
39   Lucca, L. E. *et al.* TIGIT signaling restores suppressor function of Th1 Tregs. *JCI Insight* **4**,
doi:10.1172/jci.insight.124427 (2019).
40   Markovic-Plese, S. *et al.* T cell recognition of immunodominant and cryptic proteolipid
protein epitopes in humans. *J Immunol* **155**, 982-992 (1995).
41   Raddassi, K. *et al.* Increased frequencies of myelin oligodendrocyte glycoprotein/MHC
class II-binding CD4 cells in patients with multiple sclerosis. *J Immunol* **187**, 1039-1046,
doi:10.4049/jimmunol.1001543 (2011).
42   Park, M. D., Silvin, A., Ginhoux, F. & Merad, M. Macrophages in health and disease. *Cell*
185, 4259-4279, doi:10.1016/j.cell.2022.10.007 (2022).
43   Mrdjen, D. *et al.* High-Dimensional Single-Cell Mapping of Central Nervous System
Immune Cells Reveals Distinct Myeloid Subsets in Health, Aging, and Disease. *Immunity*
48, 380-395 e386, doi:10.1016/j.immuni.2018.01.011 (2018).
44   Pappalardo, J. L. *et al.* Transcriptomic and clonal characterization of T cells in the human
central nervous system. *Sci Immunol* **5**, doi:10.1126/sciimmunol.abb8786 (2020).
45   Goverman, J. Autoimmune T cell responses in the central nervous system. *Nat Rev*
Immunol **9**, 393-407, doi:10.1038/nri2550 (2009).
46   Herz, J., Filiano, A. J., Wiltbank, A. T., Yogev, N. & Kipnis, J. Myeloid Cells in the Central
Nervous System. *Immunity* **46**, 943-956, doi:10.1016/j.immuni.2017.06.007 (2017).
47   Monney, L. *et al.* Th1-specific cell surface protein Tim-3 regulates macrophage activation
and severity of an autoimmune disease. *Nature* **415**, 536-541, doi:10.1038/415536a
(2002).
48   Li, R. *et al.* Proinflammatory GM-CSF-producing B cells in multiple sclerosis and B cell
depletion therapy. *Sci Transl Med* **7**, 310ra166, doi:10.1126/scitranslmed.aab4176
(2015).
49   Belge, K. U. *et al.* The proinflammatory CD14+CD16+DR++ monocytes are a major source
of TNF. *J Immunol* **168**, 3536-3542, doi:10.4049/jimmunol.168.7.3536 (2002).
50   Li, P., Zheng, Y. & Chen, X. Drugs for Autoimmune Inflammatory Diseases: From Small
Molecule Compounds to Anti-TNF Biologics. *Front Pharmacol* **8**, 460,
doi:10.3389/fphar.2017.00460 (2017).

733 51 Kunchok, A. *et al.* Association Between Tumor Necrosis Factor Inhibitor Exposure and
734   Inflammatory Central Nervous System Events. *JAMA Neurol* **77**, 937-946,
735   doi:10.1001/jamaneurol.2020.1162 (2020).

736 52 Gregory, A. P. *et al.* TNF receptor 1 genetic risk mirrors outcome of anti-TNF therapy in
737   multiple sclerosis. *Nature* **488**, 508-511, doi:10.1038/nature11307 (2012).

738 53 Harroud, A. & Hafler, D. A. Common genetic factors among autoimmune diseases. *Science*
739   **380**, 485-490, doi:10.1126/science.adg2992 (2023).

740 54 Montfort, A. *et al.* The TNF Paradox in Cancer Progression and Immunotherapy. *Front*
741   *Immunol* **10**, 1818, doi:10.3389/fimmu.2019.01818 (2019).

742 55 Bertrand, F. *et al.* TNFalpha blockade overcomes resistance to anti-PD-1 in experimental
743   melanoma. *Nat Commun* **8**, 2256, doi:10.1038/s41467-017-02358-7 (2017).

744 56 Bjornevik, K. *et al.* Longitudinal analysis reveals high prevalence of Epstein-Barr virus
745   associated with multiple sclerosis. *Science* **375**, 296-301, doi:10.1126/science.abj8222
746   (2022).

747 57 Lanzavecchia, A. Antigen-specific interaction between T and B cells. *Nature* **314**, 537-539,
748   doi:10.1038/314537a0 (1985).

749 58 Pisetsky, D. S. Pathogenesis of autoimmune disease. *Nat Rev Nephrol* **19**, 509-524,
750   doi:10.1038/s41581-023-00720-1 (2023).

751 59 Butler, A., Hoffman, P., Smibert, P., Papalexi, E. & Satija, R. Integrating single-cell
752   transcriptomic data across different conditions, technologies, and species. *Nat Biotechnol*
753   **36**, 411-420, doi:10.1038/nbt.4096 (2018).

754 60 Korsunsky, I. *et al.* Fast, sensitive and accurate integration of single-cell data with
755   Harmony. *Nat Methods* **16**, 1289-1296, doi:10.1038/s41592-019-0619-0 (2019).

756 61 Monaco, G. *et al.* RNA-Seq Signatures Normalized by mRNA Abundance Allow Absolute
757   Deconvolution of Human Immune Cell Types. *Cell Rep* **26**, 1627-1640 e1627,
758   doi:10.1016/j.celrep.2019.01.041 (2019).

759 62 Cribari-Neto, F. & Zeileis, A. Beta Regression in R. *J Stat Softw* **34**, 1-24 (2010).

760 63 Stephens, M. False discovery rates: a new deal. *Biostatistics* **18**, 275-294,
761   doi:10.1093/biostatistics/kxw041 (2017).

762 64 Bausch-Fluck, D. *et al.* A mass spectrometric-derived cell surface protein atlas. *PLoS One*
763   **10**, e0121314, doi:10.1371/journal.pone.0121314 (2015).

764 65 Browaeys, R., Saelens, W. & Saeys, Y. NicheNet: modeling intercellular communication by
765   linking ligands to target genes. *Nat Methods* **17**, 159-162, doi:10.1038/s41592-019-0667-
766   5 (2020).

767 66 Kang, J. B. *et al.* Efficient and precise single-cell reference atlas mapping with Symphony.
768   *Nat Commun* **12**, 5890, doi:10.1038/s41467-021-25957-x (2021).

769 67 Yasumizu, Y. *et al.* Myasthenia gravis-specific aberrant neuromuscular gene expression by
770   medullary thymic epithelial cells in thymoma. *Nat Commun* **13**, 4230,
771   doi:10.1038/s41467-022-31951-8 (2022).

Fig 1: Microglia-like CSF macrophages increase in frequency in MS patients post- B cell depletion therapy. **a,** Healthy donor and MS patient sample collection scheme for scRNAseq analysis (n = 6 healthy donors, n= 5 MS patients pre-treatment, n= 5 matched MS patients post-treatment). **b,** UMAP dimensionality reduction plot of immune cell clusters detected in CSF of healthy donor and MS patients (n= 60,704 single cells, 17 immune cell clusters). **c,** CSF macrophage cluster frequency pre- and post- B cell depletion therapy across all five MS patients. **d,** MELD likelihood enrichment heatmap and patient level summary values (mean+/- SEM, n= 5 MS patients) post- B cell depletion therapy of all immune clusters in the CSF. **e,** Heatmap of myeloid-related genes, showing average expression across all immune cell types. **f,** All myeloid clusters in the CSF scored against microglia, CNS macrophage, and pan- macrophage gene modules.

a

b

MHC Class I Module Score

MHC Class II Module Score

c

CD83 gene expression

d

e

Classical Monocyte Module Score

Intermediate Monocyte Module Score

Nonclassical Monocyte Module Score

Fig 2: Enriched CSF macrophages present anti-inflammatory phenotype in MS patients

post- B cell depletion therapy. Gene expression analyses of CSF macrophages were performed

by comparing cells from healthy donors (n=6) and pre- and post- B cell depletion therapy MS

patient samples (n=5). **a,** Volcano plot depicting differentially expressed genes in the Mac 1

cluster from MS patients pre- and post- B cell depletion. Blue: downregulated post-treatment, red:

upregulated post-treatment. -log10(p value) > 15 are capped to facilitate visualization (depicted

as triangles). **b,** MHC class I and class II gene module scores of healthy donor, MS pre-treatment,

and MS post-treatment Mac 1 cells. **c,** CD83 expression of healthy donor and MS patient Mac 1

cells. **d,** Dot plot depicting myeloid inflammatory and anti-inflammatory gene expression in healthy

donors and MS patients pre- and post- treatment. **e,** Gene module scores of all CSF myeloid

clusters against peripheral monocyte gene signatures. Top: classical monocyte module score,

middle: intermediate monocyte module score, bottom: nonclassical monocyte module score.

Fig 3: Increased CD16$^+$ monocytes abundance after anti-CD20 treatment. a, experimental design of pre- and post-treatment (anti-CD20) PBMC samples from MS patients (n=18) for droplet-based scRNAseq using 10x genomics platform. **b,** UMAP of annotated cell types (left) and overlayed MELD likelihoods for post-treatment status (right). **c,** MELD likelihood patient level summary values (mean+/- SEM) per fine-grained clusters and main cell types. **d,** fine-grained community frequency changes post-treatment (log fold change mean estimate +/-SE from beta regression, see methods). **E,** Flow cytometry validation of CD16$^+$ monocyte frequency changes in MS patients PBMCs.

Fig 4: Differential gene expression in CD16+ monocyte post-treatment. A, Mean abundance (MA) plot of gene expression changes, differentially expressed genes (DGE) are highlighted in red (upregulated post-treatment), or blue (downregulated post-treatment). **B,** Flow cytometry of HLA-DR and CD81 expression in CD16+ monocytes (n=16). **C,** GeneSet Enrichment Analysis (GSEA) using the Hallmark genesets in CD16+ monocytes. **D,** Custom GSEA analysis of PBMC monocytes post-treatment signature genesets (up- and downregulated genes) tested on CSF macrophage Mac 1 dataset (From Fig 1).

Fig 5: Geneset enrichment analysis (GSEA) of anti-CD20 gene expression alterations
across cell types. a, Heatmap of normalized enrichment scores (NES) from post-treatment
GSEA analyses run for each cluster shows ubiquitous increase in TNFα-NFκB pathway.
Differentially enriched genesets are highlighted with a *. "ID count" depicts the number of times a
geneset is found enriched across communities. **b,** Overlap graph-analysis of leading edge genes
for the "TNFα signaling via NFκB" geneset across cell types highlights two sets of signatures: B
and myeloid cells vs. T cells. **c,** Pre- and post- treatment fold change of *TNFA* transcript across
clusters (differential expression is highlighted in red). **d,** *In vitro* validation of TNFα upregulation
pre- and post- B cell depletion at the protein level in MS patient monocytes (n= 18) by intracellular
flow cytometry staining after LPS stimulation.

Fig 6: Detailed Analysis of CD4⁺ T Cell Alterations Following Anti-CD20 Treatment. a, Schematic illustration of the analysis of CD4⁺ T cells using a reference mapping and NMFproj. From CSF and PBMC samples, CD4⁺ T cells were extracted using Azimuth, and detailed CD4⁺ T clusters were predicted using Symphony. The 12 gene programs were calculated using NMFproj. **b,** Inferred CD4⁺ T cell clusters on UMAP plot. The clusters were assigned to either a major cluster (L1) or a detailed cluster (L2) level. **c, d,** Cell frequency changes after anti-CD20 treatment in CSF **(c)** and PBMC **(d)**. Coefficients of cell frequency change per cluster L2 quantified GLM (method) are visualized on the UMAP plot (left). The populations with cell frequency increases post-B cell depletion treatment are shown in red. CD4⁺ T cluster frequency pre- and post-B cell depletion therapy (right). Significantly altered clusters are shown. See Supplementary Figure 8c and 9c for additional details. **e, f,** Alterations of gene programs extracted by NMFproj after anti-CD20 treatment in CSF **(e)** and PBMC **(f)**. Dot plots depicting NMF cell feature changes in each cell type (left). Dot colors show coefficients, and sizes show the significance of GLM (method). The coefficient of gene program change per cluster for some gene programs was shown on UMAP plots (right). Annotations and representative genes of gene programs are following; NMF0 (Cytotoxic-Feature or Cytotoxic-F; *GZMB*, *CX3CR1*), NMF1 (Treg-F; *FOXP3*, *IL2RA*), NMF2 (Th17-F; *RORC*, *CCR6*), NMF3 (Naive-F; *CCR7*, *BACH2*), NMF4 (Activation-F or Act-F; *DACT1*, *CDK6*), NMF5 (TregEff/Th2-F; *HLA-DR*s, *CCR10*), NMF6 (Tfh-F; *MAF*, *CXCR5*), NMF7 (Interferon-F or IFN-F; *OAS1*, *MX1*), NMF8 (Central Memory-F; *CRIP2*, *PLP2*), NMF9 (Thymic Emigrant-F; *SOX4*, *PECAM1*), NMF10 (Tissue-F; *JUNB*, *NFKBIA*) NMF11 (Th1-F; *GZMK*, *EOMES*)

Fig 7: B cell depletion induces an increase in TIGIT$^+$ Tregs and reduces autoreactive T
cells. a, Visualization of Treg population extraction and changes after B cell depletion treatment.
Predicted CD4$^+$ T clusters and Tregs (dotted line) on UMAP plot (left). Re-embedding of extracted
Tregs using UMAP (top right). B cell depletion treatment associated relative likelihood in Treg
populations calculated using MELD (bottom right). **b,** Frequency changes of each subpopulation
within the Treg group. **c,** Volcano plot depicting differentially expressed genes in Tregs,
particularly highlighting genes encoding surface proteins. **d,** Heatmap displaying predicted
interactions between myeloid cell-derived ligands (limited to genes differentially regulated with B
cell depletion treatment) and Treg-derived receptors, weighted by prior interaction potential. **e,f,**
Flow cytometry data of Tregs frequency **(e)** and TIGIT protein expression of Tregs **(f)** in MS patient
PBMC (n=11) after B cell depletion treatment, **g,** Flow cytometry analysis of myelin tetramer-
reactive CD4$^+$ T cell frequency at pre-treatment and 6-month post-treatment timepoints (n=7). **h,**
Tfh (CD45RA$^-$CXCR5$^+$) cells in tetramer-reactive CD4$^+$ T cells and Th17 (CCR6$^+$CXCR3$^-$) cells in
tetramer-reactive CD4$^+$ T cells frequencies at pre-treatment and 6-month post-treatment
timepoints (n=7). **i.** Longitudinal kinetic analysis of the frequency of autoreactive CCR6$^+$CXCR3$^+$
CD4$^+$ T cells using flow cytometry at pre-treatment (n=4), 24-week post-treatment (n=4) and 52
week post-treatment (n=3) timepoints.