Single-cell trait diversity explains niche and fitness differences in aquatic microbial communities

Simone Fontana[a,b,c,#], Désirée A. Schmitz[d,e,f], Michael Daniels[d,e], Francesco Danza[g], Thomas Röösli[d,e], Hannah Bruderer[d,e], Jean-Claude Walser[h], Julien Dekaezemacker[i], Stephane Escrig[j], Anders Meibom[j,k], Francesco Pomati[a], Frank Schreiber[d,e,l,#]

[a]Eawag, Swiss Federal Institute for Aquatic Science and Technology, Dept. of Aquatic Ecology, Dübendorf, Switzerland.

[b]University of Freiburg, Nature Conservation and Landscape Ecology, Freiburg, Germany.

[c]Kanton St. Gallen, Amt für Natur, Jagd und Fischerei, Abteilung Natur und Landschaft, St. Gallen, Switzerland.

[d]ETH Zurich, Dept. of Environmental Systems Science, Institute for Biogeochemistry and Pollutant Dynamics, Zurich, Switzerland.

[e]Eawag, Swiss Federal Institute for Aquatic Science and Technology, Dept. of Environmental Microbiology, Dübendorf, Switzerland.

[f]University of Zurich, Department of Quantitative Biomedicine, Zurich, Switzerland.

[g]University of Applied Sciences and Arts of Southern Switzerland (SUPSI), Department of Environment, Constructions and Design, Institute of Microbiology, Via Flora Ruchat-Roncati 15, 6850 Mendrisio, Switzerland.

[h]ETH Zurich, Dept. of Environmental Systems Science, Genetic Diversity Centre (GDC), Zurich, Switzerland.

[i]Max Planck Institute for Marine Microbiology, Dept. of Biogeochemistry, Bremen, Germany.

[j]Laboratory for Biological Geochemistry, School of Architecture, Civil and Environmental

Engineering (ENAC), École Polytechnique Fédérale de Lausanne (EPFL), Lausanne,

Switzerland.

[k]Center for Advanced Surface Analysis, Institute of Earth Sciences, University of Lausanne,

Lausanne, Switzerland.

[l]Federal Institute for Materials Research and Testing (BAM), Department of Materials and

Environment, Division of Biodeterioration and Reference Organisms, Berlin, Germany.

#Corresponding authors: Simone Fontana (simonefontana.j@gmail.com) and Frank Schreiber

(frank.schreiber@bam.de)

Abstract

Two fundamental questions in ecology are how biodiversity is maintained and how it affects ecosystem functioning. Until now, it has been difficult to study the above mechanisms in natural microbial communities, yet they are important drivers of biogeochemical ecosystem functions. Here, we use a new approach to define and measure biodiversity in complex lake microbial communities (Lake Cadagno, Switzerland) based on the cell-to-cell variation in multiple functionally relevant phenotypic traits. We use stable isotope probing coupled to correlative imaging using confocal laser scanning microscopy (CLSM) and nanometer-scale secondary ion mass spectrometry (NanoSIMS) to obtain morphological (size), physiological (pigments) and metabolic (carbon and nitrogen isotope uptake and sulfur content) traits for a large number of individual cells along the environmental gradient found across lake depth. We show that cell-to-cell trait variation is significantly correlated with cell densities as a proxy for ecosystem functioning, whereas genetic diversity measured at the level of 16S and 18S is not. Our single-cell analysis provides evidence for a simultaneous increase in niche partitioning (measured as increased evenness in pigment composition) and decrease in fitness differences (measured as decreased variability in sulfur content) due to light limitation and competition for nutrients in deep layers of the lake. This leads to a negative relationship between niche and fitness differences. Our results suggest that niche and fitness differences in natural microbial communities can be understood at the level of single-cell traits, providing a mechanistic understanding of the relationship between microbial diversity and ecosystem functioning.

Introduction

Modern and past biodiversity levels on Earth have raised two central questions in ecological research: how did this diversity arise and how is it maintained in natural ecosystems? The principle of competitive exclusion[1-3] and the closely related concept of ecological niche[4-6] initially led to the recognition that species have different requirements and use distinct resources, thus explaining their coexistence. However, the number of different types of organisms – especially in microbial communities – greatly exceeds the number of distinct limiting resources (e.g., 'the paradox of the plankton'[7]), and other mechanisms must also play a role. These mechanisms have broadly been divided into two categories: increasing niche partitioning and decreasing fitness differences[8-10], which are not mutually exclusive. Indeed, both niche and fitness effects can act simultaneously[11,12] and be considered interdependent[13]. In addition, evolutionary (genetic) mechanisms also act as sources of diversity in microbial communities[14,15]. For example, rapid evolution has been shown to maintain taxonomic diversity in aquatic communities, even in the absence of niche differentiation[16,17].

In addition to the drivers of diversity, ecological research has also addressed its consequences, i.e., biodiversity and ecosystem functioning relationships (e.g.[18,19]). The mechanisms that generate diversity are arguably related to those determining ecosystem functioning[20-22]. As microbial communities are fundamental drivers of biogeochemical functions, such as the remineralization of polymers and the nitrogen cycle, it is essential to elucidate these mechanisms to better understand the contribution of microbial communities to ecosystem functioning.

Interestingly, both niche partitioning and fitness differences can be the outcome of variation in functional traits that, by definition, are features measurable at the level of individual organisms[23]. In other words, the outcome(s) of coexistence observed at the community level are in fact the result of interactions occurring among individuals[24]. However, studies of niche and fitness

differences at the level of the effective competitive units, i.e., individual organisms are still lacking.

Addressing this gap is particularly challenging for microorganisms in the natural environment, as

it is difficult to measure their relevant traits at the single-cell level. Most research to date in this

field has used low specificity measures of trait diversity, typically including multiple traits that

provide limited mechanistic understanding, such as broad classes of functionally unrelated genes

from metagenomic datasets[25]. However, the advantage of this approach is that it often provides

good predictive power of ecosystem functioning[24]. We capitalize on recent advances in the

understanding of microbial function through single-cell trait-based approaches to investigate

niche and fitness differences in microbial communities based on single-cell phenotypes.

Modern high-throughput single-cell technologies allow for a more mechanistic and hypothesis-

driven investigation of microbial adaptive strategies based on individual-level traits[26]. The

integration of such technologies has the potential to reveal niche and fitness differences at the

level of the individual organisms and assess whether these traits drive stabilizing and equalizing

mechanisms at the population and community level. First, the definition of microbial species

(assuming they exist) is challenging and not trivial[27-29]. Second, the extreme phenotypic variability

observed even within clonal microbial populations[30] or closely related sub-taxonomic groups[31]

might jeopardize the promise of modern taxonomic approaches linked to rRNA gene sequencing

to elucidate relationships between biodiversity and ecosystem function[32-34]. Taken together, a

single-cell perspective will improve our understanding of how niche and fitness differences relate

to ecosystem functioning in nature, with possible implications for the monitoring of biodiversity

and related functions.

In this study, we aim to investigate niche and fitness differences within a natural microbial

community and their relationship with ecosystem functioning based on the characteristics of

individual microbial cells. In this context, it is crucial to identify which traits might cause niche

partitioning, and which may be used to quantify fitness differences. We use correlative

fluorescence microscopy and stable isotope probing coupled to nanometer-scale secondary ion

mass spectrometry (NanoSIMS)[35] to determine photosynthetic pigment composition, size, and

metabolic activity for carbon, nitrogen and sulfur simultaneously in thousands of individual

microbial cells from different samples. We leverage the idea that environmental gradients explain

the diversification of ecological strategies or niches[36], as well as fitness differences between

phenotypes[37]. On the one hand, pigment-related traits (i.e., photosynthesis/physiology) enable

niche partitioning, as cells can decrease competition for light by targeting slightly different

wavelengths through fine-tuning of the relative expression of different pigments[38]. On the other

hand, traits defining the assimilation or cell content of elements involved in fundamental metabolic

pathways (C, N, and S) - which often show great intraspecific variation[39-41] - can be considered

as proxies for fitness, as they influence the maintenance, growth and reproduction of

organisms[42,43]. Here, we sampled single cells living in structured microbial communities at

different depths in the meromictic Lake Cadagno (Switzerland), and considered decreasing levels

of trait integration to predict ecosystem functioning with multidimensional trait diversity indices

and detect niche and fitness differences determined by the variation in physiological and

metabolic traits[24]. Accordingly, we formulate the following expectations:

1) Ecosystem functioning (i.e., cell density as a proxy for primary production) is better

explained by trait diversity metrics that integrate all available individual-based traits

(morphological, physiological, and metabolic) rather than by taxonomic diversity metrics based

on rRNA gene sequencing[33,44]. **2)** In deep layers of the lake, competition for light drives increasing

niche partitioning in pigment-related traits (i.e., increasing trait evenness[38]), while limitation for

essential nutrients mediates decreasing fitness differences in metabolic traits (i.e., trait

convergence[45]), as compared to the upper layers of the lake. **3)** These simultaneous shifts in trait

distribution lead to a negative relationship between single-cell niche and fitness differences across

water depth.

Results

Environmental variables structure Lake Cadagno into distinct layers

First, we set out to characterize the environmental variables that shape the microbial communities in Lake Cadagno and that potentially drive trait variability, mediating niche and fitness differences. The vertical structure of physical, chemical and biological parameters of the meromictic Lake Cadagno during one day of our sampling campaign is shown in Fig. 1A (see Extended Data Fig. 1 for the profiles of the other two days). Lake Cadagno is perfectly suited to study niche and fitness differences in microbial communities because its steep gradients of light intensity, oxygen and sulfide concentration (Fig. 1A) create ideal conditions for very distinct microbial communities at different depths[46]. Differences in salt concentration stably stratify the lake into three main layers. The upper layer, called mixolimnion, is dominated by oxygenic photosynthetic organisms. Oxygen is consumed around 11 m water depth, below which light can still penetrate. A chemocline then forms in this largely anoxic water, which is dominated by high cell densities of anoxygenic phototrophic purple and green sulfur bacteria. As electron donors, these bacteria use sulfide and other reduced sulfur compounds that diffuse upward from the lake's bottom waters (Fig. 1A; see also[47]). Below the chemocline, in the bottom waters, light penetration is no longer measurable, and sulfide is produced by sulfate-reducing bacteria (Fig. 1A).

Fig. 1. Vertical gradients of physical, chemical and biological parameters in Lake Cadagno separate the ecosystem into three distinct layers. **(A)** Vertical profiles of conductivity (blue), temperature (green), oxygen (O_2, black), total sulfide (red), light (orange), and cell density (purple) measured on day C (19/08/2014) of the sampling campaign. Profiles from the other sampling days are shown in Extended Data Fig. 1. The gray area represents the chemocline, where sulfide and O_2 are depleted, the cell density increases, and light gets severely limited. Light values are presented as % photosynthetic active radiation of the light above the water. Light values shown as half circles below the chemocline were below the detection limit of the sensor. **(B)** Principal component analysis (PCA) of individual samples based on environmental factors (abbreviations: T=temperature; Cl^-=chloride; NH_4^+=ammonium; PO_4^{3-}=ortho-phosphate; DIC=dissolved inorganic carbon; SO_4^{2-}; sulfate; PN=particulate nitrogen; PC=particulate carbon). The PCA plot shows individual samples (circles) taken at different days distributed across environmental factors. Samples belonging to the same sampling layer (color) are grouped more closely in two-dimensional space. Samples for which individual-based trait diversity (*'Trait Onion Peeling'* TOP, *'Trait Even Distribution'* TED and *'Functional Dispersion'* FDis) were calculated for further analysis are marked by a black point. The surface layer is defined hereafter as 'upper layer' (as the gray dots are not considered for further analyses), and the layer below chemocline as 'lower layer'.

Our analysis of the lake stratification based on environmental factors (Fig. 1B) shows that Lake Cadagno is separated into distinct layers and that different environmental factors shape the structure and function of the microbial communities in each layer. Specifically, the upper layer is strongly separated from the chemocline and the lower layer by one axis of the principal component analysis (PCA; dimension 1), which explains 56.4% of the variance. The variables that most

strongly contribute to variation in this dimension are depth and light intensity. This indicates that

the microbial communities found in these samples are, as expected, stratified based on the

availability of light as a resource for photosynthesis. Samples in the chemocline are most strongly

separated from the lower layer by other environmental parameters (dimension 2 of the PCA,

explaining 18.9% of the variance): high turbidity, oxidation-reduction potential, particulate carbon,

and ammonium concentration. Samples in the lower layer correlate most strongly with the

concentration of ammonium and ortho-phosphate. These environmental factors thus separate

Lake Cadagno into different layers and provide the physio-chemical basis for the structuring of

the lake's microbial communities.

Single-cell trait diversity explains ecosystem functioning

Next, we tested whether ecosystem functioning can be better explained by integrative trait

diversity metrics (including physiological and morphological traits from CLSM and metabolic traits

from NanoSIMS; Fig. 2) than taxonomic diversity metrics based on rRNA gene sequencing. To

do that, we analyzed trait and taxonomic diversity data across the different depths of the lake and

time (days) with linear models to assess their explanatory power for ecosystem functioning,

represented here by the density of cells as a proxy for primary production. The analysis of our

large dataset across time and depths shows that trait diversity is a significant driver of ecosystem

functioning. Specifically, the relative regularity in distances between individual cells (TED, *'Trait*

Even Distribution') and their spread around the average phenotype (FDis, *'Functional Dispersion'*)

in high-dimensional trait space was significantly related to ecosystem functioning, whereas the

total trait space covered (TOP, *'Trait Onion Peeling'*) did not. Thus, the two best models retained

for model averaging explained 56.8% (explanatory variables: TED, FDis) and 71.9% (explanatory

variables: TED, FDis, Day of sampling) of the variance in cell density relative to the maximum

achieved in a given layer (Extended Data Table S1). Model averaging showed a negative effect

of TED on cell density (standardized regression coefficient=-0.195; 95% CI=[-0.339; -0.050]),

whereas FDis had a positive effect of similar strength (standardized regression coefficient=0.153;

95% CI=[0.028; 0.279]) (Fig. 3). Contrary to this, taxonomic diversity metrics based on rRNA gene

sequencing did not show any significant correlation with cell density. Indeed, the best model

according to AICc (Akaike information criterion corrected for small sample sizes) was the null

model (including the intercept only). The only other model retained for model averaging also

included the Simpson index based on 18S sequencing and only explained 5.7% of the variance

in cell density relative to the maximum achieved in a given layer (Extended Data Table S1). As a

consequence, the effect size of the Simpson index based on 18S sequencing is not significantly

different from 0 (standardized regression coefficient=0.026; 95% CI=[-0.079; 0.130]) (Fig. 3).

Taken together, these data show that trait diversity explains ecosystem functioning in Lake

Cadagno while taxonomic diversity does not.

Fig. 2. Correlative microscopy to determine single-cell trait-based diversity indices. Panels **(A)** and **(B)** show example images for the same region of interest (ROI) measured with confocal laser scanning microscopy and NanoSIMS, respectively. The lines around cells show the cell segmentation that was based on DNA stain and the $^{12}C^{14}N$ NanoSIMS signal. Cells that are at the edge of each ROI were not included in the analysis. The excitation (Ex.) and emission (Em.) wavelengths are shown above each microscopy image. The intensity scale for isotope counts ($^{12}C^{14}N$ and ^{32}S) and isotope fractions ($^{15}N/[^{14}N+^{15}N]$ and $^{13}C/[^{12}C+^{13}C]$) are shown below each NanoSIMS image. Scale bars are only shown in the NanoSIMS images, but they are also valid for the microscopy images since they show the same ROI. **(C)** Scheme summarizing how the measured single-cell traits are used to calculate the three components of trait diversity: richness

(TOP index), evenness (TED index) and divergence (FDis index). Note that the scheme shows the indices on a 2-

dimensional trait axis to better understand the principle. In this study, the trait axes form a multi-dimensional trait space

with seven dimensions represented by fluorescence data (three channels) and cell size from CLSM, and NanoSIMS

data (C and N assimilation, as well as S cell content).

Fig. 3: Single-cell-based trait diversity explains ecosystem functioning in Lake Cadagno, whereas taxonomic diversity

based on rRNA gene sequencing does not. Effect size of trait diversity metrics (TOP index of trait richness, TED index

of trait evenness, and FDis index of trait divergence, calculated using three fluorescence channels, cell size from CLSM,

as well as C assimilation, N assimilation, and S cell content) and genetic diversity metrics (Chao1 and Simpson index

using 16S and 18S sequencing data) on cell density. Only metrics included in the list of models used for model

averaging are shown here (predictors with a significant effect are in bold). Trait diversity (N=17) and genetic diversity

(N=24) were modeled separately (see Methods). Values are standardized model-averaged regression coefficients

shown as black circles with their 95% confidence intervals depicted as error bars (details about single models can be

found in Extended Data Table S1). Cell density (quantified by scanning flow cytometry) is relative to the maximum

achieved in a given layer.

Single-cell traits drive niche and fitness differences in distinct lake layers

We analyzed our extensive single-cell dataset to investigate whether the highly competitive

environment in deep layers of the lake drives increasing niche partitioning in pigment-related traits

and decreasing fitness differences in metabolic traits. The data showed that the evenness of

pigment-related traits was significantly higher in the chemocline (under low light intensity

combined with high cell density) as compared to the upper layer (under highest light availability)

(Fig. 4, Table 1A). The lower layer showed intermediate evenness of pigment-related traits, but

differences with the upper layer and chemocline were not significant (Table 1A). In contrast, the

evenness of metabolic traits did not show any significant difference between the three layers of

the lake (Fig. 4 and Table 1B). This suggests that the evenness in traits related to light acquisition

is positively related to increased niche partitioning under light limitation.

The analysis further revealed that metabolic traits exhibit a significantly lower multidimensional

metabolic trait divergence in the chemocline and lower layer than in the upper layer (Fig. 4 and

Table 1D). In contrast, no significant difference in the divergence of pigment-related traits was

observed between the three layers of the lake (Table 1C). This suggests that decreased

divergence in metabolic traits related to resource acquisition is linked to decreased fitness

differences. The latter occurred when competition for limiting resources (i.e., nutrients) increased.

Taken together, our data suggest that niche partitioning and fitness effects act simultaneously

between single-cell phenotypes in microbial communities along the depth gradient of Lake

Cadagno and are linked to different sets of traits.

Fig. 4: Niche and fitness differences are linked to different sets of traits. One-way ANOVA analyses and post hoc Tukey tests (see Table 1; significance level at p<0.05, N=17) show that the evenness of fluorescence traits (TED) is higher in the chemocline than in the upper layer, whereas trait divergence (FDis) does not show any significant difference (column **A**). Conversely, the evenness of metabolic traits measured with NanoSIMS does not change significantly between layers, but trait divergence is highest in the upper layer (column **C**). Fluorescence and metabolic trait distributions show weak and non-significant Spearman correlations (column **B**; function *cor.test*, N=17), further supporting the idea that fluorescence and metabolic traits reflect distinct mechanisms. Each boxplot reports minimum and maximum (whiskers), first and third quartile (lower and upper side, respectively), as well as median (horizontal black line). We note that, for the calculation of TED and FDis, cell size (from CLSM) was included along with three fluorescence channels from CLSM on one side (see Fig. 2A), as well as C assimilation, N assimilation and S cell content on the other side (see Fig. 2B).

265   **Table 1:** The results of the one-way ANOVA analyses and post hoc Tukey tests for the data shown in Fig. 4. All p-

266   values are adjusted for multiple comparisons.

	A) TED (fluorescence+size)		B) TED (NanoSIMS+size)	
	difference	p-value	difference	p-value
chemocline *vs.* upper layer	0.235 [0.118; 0.351]	<0.001	-0.015 [-0.119; 0.089]	0.924
lower layer *vs.* upper layer	0.151 [-0.001; 0.303]	0.052	-0.086 [-0.222; 0.050]	0.258
lower layer *vs.* chemocline	-0.084 [-0.230; 0.062]	0.319	-0.071 [-0.201; 0.059]	0.356

	C) FDis (fluorescence+size)		D) FDis (NanoSIMS+size)	
	difference	p-value	difference	p-value
chemocline *vs.* upper layer	0.123 [-0.451; 0.697]	0.842	-0.551 [-0.894; -0.207]	0.002
lower layer *vs.* upper layer	-0.117 [-0.868; 0.635]	0.913	-0.502 [-0.952; -0.052]	0.028
lower layer *vs.* chemocline	-0.240 [-0.960; 0.480]	0.666	0.049 [-0.382; 0.479]	0.953

Niche and fitness differences are negatively related because of resource limitation

After we confirmed that individual-level niche and fitness differences are evident in our dataset,

we analyzed how they are linked to each other across the main environmental gradient of the

lake. To this end, we formulated the hypothesis that single-cell niche and fitness differences are

negatively related across depth. More specifically, our analysis showed that the cell-to-cell

coefficient of variation in fitness proxies was negatively related to the evenness of traits related to

pigments. This is arguably not unexpected because, with increasing depth, both increased niche

partitioning of the light spectrum and decreased fitness differences due to more similar metabolic

traits should occur. The results of the linear model (estimate= -2.378, SE=0.752, t=-3.164, df=15,

R^2=0.400; p=0.006) suggested that the individual-level niche partitioning of the light spectrum in

deeper layers was strongly associated with more similar fitness as estimated by S content.

Accordingly, cells in the upper layer were characterized by a high coefficient of variation in S

content and uneven distribution of pigment-related traits (Fig. 5A, green dots), whereas cells in

the chemocline and lower layer showed the expected combination of high evenness of pigment-

related traits and low coefficient of variation in S content (Fig. 5A, red and blue dots). We also

investigated the relationship between the evenness of pigment-related traits and the coefficient

of variation in the single-cell assimilation rates of two other elements (C as CO_2 and N as NH_4^+;

Fig. 5B and C, respectively). We did not find any significant relationship between trait evenness

of pigments and C assimilation rate (estimate=0.221, SE=1.537, t=0.144, df=15, R^2=0.001;

p=0.888), which suggests that CO_2 assimilation is not a good proxy for fitness in this system. In

contrast, the coefficient of variation of NH_4^+ assimilation rates showed a significant and positive

relationship with the evenness of pigment-related traits (estimate=1.332, SE=0.584, t=2.280,

df=15, R^2=0.257; p=0.038). Cells in the upper layer (Fig. 5C, green dots) were generally more

similar to each other in terms of NH_4^+ assimilation rates than cells in the chemocline and lower

layer (Fig. 5C, red and blue dots).

Fig. 5: Relationship between niche and fitness differences across different layers of Lake Cadagno. Niche partitioning of the light spectrum was quantified using the TED index of trait evenness calculated with fluorescence data (three channels) plus cell size from CLSM data (see Fig. 2A & C). Fitness differences were quantified from NanoSIMS data (see Fig. 2B) using the coefficient of variation in S content (panel **A**), C assimilation (panel **B**) and N assimilation rates (panel **C**) among individual cells, whereby fitness differences decrease with decreasing coefficient of variation. Each circle represents a sample taken at a specific water depth and day of sampling. The color of the circles represents the association of the sample to a specific lake layer as defined in Fig. 1B. The size of the circle represents the association of each sample to a category of the magnitude of relative cell density in this layer as a proxy for ecosystem functioning. Black regression lines, R^2, and p-values were obtained by fitting simple linear regressions (N=17).

Discussion

In this study, trait diversity metrics integrating individual-level morphological, physiological and metabolic traits correlate significantly with ecosystem functioning, measured as cell densities, in contrast to taxonomic diversity metrics based on rRNA gene sequencing. This is consistent with previous findings: for example, multidimensional indices of functional diversity based on individual-based traits have been shown to outperform taxonomic diversity metrics in explaining phytoplankton biomass in natural communities[34]. Furthermore, the appropriate definition of microbial strategies (trait combinations) can help explain the contribution of microbial processes to carbon fluxes in soil ecosystems[48]. The lack of correlation between ecosystem functioning and taxonomic diversity, however, contradicts some previous findings[21]. One potential explanation for this is the importance of resource limitation in Lake Cadagno as compared to previously investigated systems. Previous studies have shown a positive relationship between resource limitation and intraspecific phenotypic heterogeneity in microbial populations[38,41,49]. Phenotypic heterogeneity can result from physiological acclimation and stochastic gene expression, and these physiological responses are not captured by taxonomic analysis. It is important to note that, in this case, physiology might have a key role in predicting ecosystem functioning, more than community composition or richness from taxonomic analysis. An alternative explanation could be that rRNA-based diversity has much greater interspecies redundancy than single-cell functional diversity for the traits we measured[50] or that the measured traits show only a shallow phylogenetic conservation[51]. Indeed, to detect differences in trait diversity indices, it was necessary to first determine the minimum number of single cells per sample required for a robust analysis (Extended Data Fig. 2). It is noteworthy that 250 cells are sufficient for this purpose despite the presence of a high taxonomic diversity. This suggests that many different species and strains rely on similar functions and metabolic pathways[52]. Another explanation could be the extremely high accuracy with which we can measure differences in traits; for example, the consideration of the

relative absorbance of multiple light wavelengths on a continuous scale allows us to distinguish

individual cells based on slight differences in pigment profiles.

It has been recognized that trait variation among individuals must be considered to understand

species coexistence[53,54]. These differences among individuals, attributable for example to

phenotypic plasticity or phenotypic heterogeneity, affect both niche and fitness differences and

therefore coexistence[30,55]. This may explain, for example, why the average cell or colony size of

freshwater cyanobacterial species, while affecting both their niche and fitness differences, does

not predict their coexistence even in simplified experimental communities[56]. This is consistent

with our hypothesis that increasing niche partitioning (evenness in pigment-related traits) and

decreasing fitness differences (convergence in metabolic traits) are better defined at a single-cell

level when seeking a mechanistic understanding of microbial diversity. Niche and fitness

differences are not necessarily independent but may be directly correlated, especially if they are

both involved in metabolic activity and more specifically in photosynthesis. This interdependence

should become apparent when using more specific and less integrative proxies for fitness. In this

study, we find the cell content of S to be a good proxy for fitness, as sulfur globules ensure the

availability of electron donors despite the extremely variable environmental conditions in the

chemocline[57]. Indeed, sulfur bacteria can accumulate sulfur globules through sulfide uptake, and

further use them as electron donors under conditions of sulfide limitation[58,59]. An even partitioning

of the light spectrum among cells could also be directly responsible for the more similar values of

cellular S (Fig. 5A). Indeed, this form of niche partitioning allows for a more uniform access to

photons by decreasing competition for light, and the light-dependent accumulation of sulfur

globules[58,59]. In contrast, N and C assimilation rates might not be good proxies for fitness, as a

species dominating the uptake of both elements (i.e. *Chromatium okenii*) is present at very low

densities in the chemocline of Lake Cadagno, whereas the most abundant species (i.e.,

Chlorobium clathratiforme) contributes much less to N and C cycles[57]. This may explain why we

did not find any relationship between the evenness of pigment-related traits, our measure of niche

partitioning, and the variability of C assimilation rates (Fig. 5B). On the contrary, N assimilation

rates became increasingly variable among cells with increasing evenness of pigment-related traits

(Fig. 5C). This suggests that sulfide (electron donor) limitation in the chemocline and lower layer

drives heterogeneity in N assimilation, as previously shown experimentally for the fixation of N_2

in a strain of the green sulfur bacterium *Chlorobium phaeobacteroides* isolated from Lake

Cadagno[60].

Our results support the idea that niche and fitness differences, even at the individual level, can

be associated with different sets of traits, which can be measured simultaneously on the same

single cells with the methods used in this study. In the chemocline and lower layer, cells show a

higher evenness of pigment-related traits (i.e., niche partitioning of the light spectrum under light

limitation, as observed in experimental phytoplankton populations[38]) and at the same time a lower

variation in cell S content (i.e., more similar fitness). In other words, the environment of deep

water layers, which we assume to be highly competitive because of high cell densities along with

light and nutrient limitation, requires both high niche partitioning and low fitness differences for

organisms to coexist. It should be noted that the strong correlation between these two

mechanisms of coexistence (Fig. 5) does not imply a causal relationship. The fact that the cells

increasingly partition the light spectrum might not necessarily cause the very similar cell S content

but could otherwise indicate that S is a metabolic intermediate of an essential metabolic pathway

in the chemocline and lower layer as opposed to the upper layer. Low S content variation in deep

layers of the lake might reflect the role of S as an electron donor in anoxygenic photosynthesis.

In order to survive there, cells need both enough photons to sustain photosynthesis (leading to

high evenness in pigment-related traits) and enough S to channel their metabolism through this

element (leading to low variation in cell S content).

Our findings have important implications for microbial ecology, biogeochemistry and

community ecology. By measuring focal functional traits at the single-cell level, we connect the

three above-mentioned fields in a novel conceptual framework that uses niche and fitness

differences among individual cells to explain emerging properties at the community and

ecosystem level, such as cell densities and standing biomass that sustains natural food webs.

Fluorescence and metabolic traits represent fundamentally different axes of variation (see lack of

correlation in Fig. 4), which only in combination can explain the occurrence of the corresponding

phenotypes in a highly competitive environment. Our results further imply that, despite the high

taxonomic complexity of natural microbial communities, single-cell trait-based approaches can

explain community-level functions such as the nutrient cycling by reducing this complexity to a

few relevant traits[24], i.e., morphological, physiological and metabolic. Such strong links between

traits and ecosystem properties across different environments have also been shown to exist at

the level of microbiomes, aiding the identification of functional groups or guilds in metagenomic

datasets[61,62], as well as ecological clusters based on representative genomes[63]. Our approach

might benefit the understanding of ecological communities in general, including those composed

of larger organisms, and their links to broad-scale ecosystem functions[64]. Our findings support

the need for manipulative experiments and emerging theoretical concepts that explain species

coexistence using the fundamental unit at which interactions and selection occur in ecosystems,

namely the individual organisms. Such concepts should eventually encompass all levels at which

eco-evolutionary processes linked to environmental change occur: i.e. intraspecific trait variation

resulting from phenotypic heterogeneity[30] and phenotypic plasticity[55,65], as well as rapid

evolutionary   adaptations[16].

Methods

Study site, design, and sampling

The study was conducted in August 2014 in the meromictic lake 'Lake Cadagno' located in the Swiss Alps (1923 m above sea level) in Ticino, Switzerland. The maximum depth of the lake is 21 m. The water of the lake is infiltrated through gypsum-rich dolomite rock. The infiltrated water therefore transports salts, including sulfate, to the bottom water of the lake. This process establishes an anaerobic, euxinic monimolimnium with high salinity, and an aerobic, low-salinity mixolimnion separated by a permanent chemocline at ca. 10-14 m depth. The chemocline is characterized by sharp gradients and a strong turbidity maximum (ca. 1 m vertical thickness) dominated by populations of purple and green sulfur bacteria, which grow by anoxygenic photosynthesis with sulfide[57]. The exact depth of the turbidity maximum can fluctuate (between days or even within hours) as internal waves build up depending on the wind conditions.

Fieldwork was conducted for 3 consecutive days in August 2014. The days had varying sunlight exposure, introducing differences in photosynthetic activity. Each day, a multiparametric profile (temperature, conductivity, pH, salinity, dissolved oxygen, redox potential, and turbidity) was measured in the water column using a YSI 6000 profiler (Yellow Spring, Inc., Yellow Spring, OH). Water was retrieved through gas-tight tubing (Viton, d=6 mm) mounted to the multiparametric probe with a vacuum-driven pump (0.3 bar, 200 mL min^{-1}) during the profile measurements. Samples were retrieved throughout the water column in 1 m steps from 1 to 10 m depth and in 0.5 m steps from 10.5 to 15 m depth. An overview on all samples and the associated analysis conducted with them is shown in Supplementary Dataset 1. Samples for stable isotope incubations were filled anoxically into N_2-rinsed, septum-sealed, 250 mL serum bottles without head-space. Samples for chemical and flow cytometric analyses were sampled into 250 mL serum bottles that were cleaned by consecutive rinsing with 0.1% HCl and double-

deionized water. Samples for DNA extraction and amplicon sequencing were sampled into 1 L

bottles that were cleaned by consecutive rinsing with 2.5% NaClO, 0.1% HCl and 0.2 µm-filtered

double-deionized water. Moreover, we measured a profile for photosynthetically active radiation

as the percentage of transmission (PAR; Two LI-193SA spherical quantum sensors from LI-COR

Ltd, Lincoln, NE) in the afternoon of each day after samples had been retrieved and stable isotope

incubations had been placed in the lake.

Chemical analysis

Samples for chemical analyses were filtered onsite through 0.47 µm nitrocellulose filters

(Sartorius), stored cold and dark in autoclaved, crimp-sealed serum bottles without headspace,

and analyzed within 10 days after sampling. Samples for ammonium were treated differently;

these samples were filtered through 0.2 µm polycarbonate filters (GTTP, Millipore) and stored in

50 mL polypropylene tubes (consecutively rinsed with 0.1% HCl and 0.2 µm-filtered double-

deionized water). Sulfate, chloride and nitrate were analyzed by ion chromatography (Metrohm

930 Compact IC Flex with chem. suppression; Metrohm Metrosep A Supp 5 100/4 mm column;

conductivity detector). Ammonium was measured with Berthelot's reaction (DIN 38406-5:1983-

10). Ortho-phosphate was measured after the formation of a phosphorous-molybdenum blue-

complex (DIN EN ISO 6878, 2004). Dissolved inorganic carbon was measured with TOC-L CSH

(Shimadzu, Japan). Sulfide was measured one day from 12 mL samples that were immediately

transferred to screw-capped tubes containing 0.8 mL 4% zinc acetate solution. These samples

were stored in the dark and analyzed colorimetrically using a commercial kit (Spectroquant Sulfide

kit, Merck, Schaffhausen, Switzerland). Total particulate nitrogen and carbon were determined by

filtration of 100 to 220 mL lake water on pre-combusted (400°C, 3 h) 0.7-µm-pore glass microfibre

filters (GF/F, Whatman) and subsequent measurement with isotope ratio mass spectrometry

(IRMS). Briefly, the glass microfibre filters for IRMS were immediately stored at -20°C after

filtration. After the fieldwork, samples were dried for at least 3 h at 65°C and decalcified in an

atmosphere generated by 37% hydrochloric acid. The filters were kept dry in a desiccator at room

temperature until further processing. The filters were measured by N_2 and CO_2 released by flash

combustion in excess oxygen in an automated Thermo Flash EA 1112 elemental analyzer

coupled to an isotopic ratio mass spectrometer (Thermo Delta Plus XP, Thermo Fisher Scientific).

Standards were prepared from caffeine and total particulate carbon and nitrogen were calculated

from the summed peak areas of masses 44, 45, and 46 for carbon and masses 28, 29, and 30

for nitrogen in each sample.

Definition of lake layers

In order to determine the contribution of environmental factors to the distribution of samples in

our dataset, we performed a principal component analysis (PCA; Fig. 1B). Individual samples

were grouped into three major layers based on their position relative to the chemocline: upper

layer, chemocline and lower layer. Samples from above the chemocline were further divided into

a surface layer (initial 6 m of Lake Cadagno) and a layer just above the chemocline. This further

subdivision was based on the fact that many environmental factors such as conductivity,

temperature and oxygen concentrations were stable in the surface layer and gradually changed

thereafter (Fig. 1A). In order to maximize differences between layers, we excluded from further

analyses the layer between surface and chemocline, which shows intermediate properties and is

therefore less clearly defined (Fig. 1B). For the PCA the following environmental factors were

considered: depth [m], light [% PAR above the water], temperature [°C], conductivity [S/m], salinity

[practical salinity units, psu], oxygen [mg L^{-1}], pH, oxidation-reduction potential [mV], turbidity

[formazine turbidity unit, FTU], ammonium [µmol N/L], dissolved inorganic carbon (DIC) [mg/L],

ortho-phosphate [µg/L], chloride [mg/L], sulfate [mg/L], particulate nitrogen [µmolN/L], particulate

carbon [µmolC/L]. For a small subset of points, individual values were not collected due to device

failure. For these points, we calculated the mean of the respective variables using the samples

directly above and below the missing value.

Flow cytometry

Using the scanning flow cytometer Cytobuoy (www.cytobuoy.com, Woerden,The Netherlands), we counted and characterized phytoplankton cells from water sampled at different depths during each sampling day. Samples were fixed after stable isotope incubation with a fixative solution containing 0.01% paraformaldehyde and 0.1% glutaraldehyde at pH=7 and were stored at 4°C for 14 weeks until flow cytometric measurements. Cytobuoy contains two laser beams (coherent solid-state sapphire, wavelengths 488 and 635 nm) and three different detectors to measure the fluorescence emitted by photosynthetic pigments (ranges: 668–734 nm, 601–668 nm, 536–601 nm). Measurements were triggered by the sideward scattering signal. More details on the instrument can be found elsewhere[34].

Cytometry data were successively cleaned by separating living phytoplankton cells from organic debris, suspended solids, and non-photosynthetic (heterotrophic) bacteria. First, a clustering algorithm was applied to a random subset of particles using the library *flowPeaks*[66], and junk clusters were identified visually (characterized by low fluorescence in all pigments). Second, a random forest classifying algorithm[67,68] was trained using the library *randomForest*[66] to identify junk particles in the whole datasets and remove them. In addition, only particles in the size range between 1 μm and 1 mm are reliably characterized by Cytobuoy and were therefore retained in the dataset. The number of living phytoplankton cells identified was divided by the volume processed from each sample to calculate the cell density of photosynthetic cells as a proxy for primary production (ecosystem functioning).

Stable isotope incubations

Stable isotope incubations were performed with anoxically-sampled, freshly-retrieved lake water as described under 'Study site, design, and sampling'. The samples in the serum bottles were spiked with 100 μL stock solutions of ^{15}N-NH_4Cl (25 mM, Sigma) and ^{13}C-$NaHCO_3$ (0.5 M,

Sigma) to a final concentration of 10 µM and 200 µM, respectively. The final labeling percentage

(^{13}C = 10 to 30%; ^{15}N = 30 to 100%) was calculated for each sample retrospectively based on

total NH_4 and DIC measured through chemical analysis in separate water samples. After the

addition of the isotopically-labeled substances, serum bottles were attached to a rope that was

intruding into the water column and that was mounted on a stably-anchored research platform.

The bottles were attached at different positions on the rope in a way that they incubated at the

depth from which they had been retrieved. The bottles incubated for an entire day/night/day cycle

(approximately 22 h) starting from about noon each day. The serum bottles were retrieved from

the lake at the end of the incubations and kept in the dark and on ice until further processing (2

to 3 hours). Samples from the incubations were fixed with a fixative solution for later analysis with

correlative imaging and flow cytometry (30 to 50 mL).

Correlative imaging for autofluorescence and isotopic enrichment

Single-cell correlative imaging was performed with confocal laser scanning microscopy

(CLSM, Leica SP5) for autofluorescence and nanometer-scale secondary ion mass spectrometry

(NanoSIMS 50L, CAMECA) for isotopic enrichment within the Laboratory for Biological

Geochemistry at EPFL and University of Lausanne. The cells were placed on polycarbonate filters

(0.2 µm GTTP, Millipore) by vacuum filtration. The filters had a diameter of 5 mm. The diameter

of the filtration area was 3 mm. Etched marks were distributed across the filtration area with laser

micro-dissection (PALM micro-dissection, Zeiss 200M equipped with a 355 nm pulsed UV laser)

before filtration. The filtered volumes of fixed cells were adjusted between 0.2 to 2 mL to have an

evenly distributed, dense single-cell layer. Filtration of cells was immediately followed by filtration

of 200 µL double-deionized water to wash the filters. Air-dried filters were coated with a thin film

of low melting agarose by dipping them into a 0.1% low melting agarose solution at 35°C,

removing excess agarose with a tissue by carefully touching the filter edge with the tissue, and

air-drying the filters. The filters were stored dry and in the dark at room temperature until CLSM

and NanoSIMS measurements.

First, the filters were imaged with CLSM to detect single-cell autofluorescence signals. To this

end, cells were initially stained with a general DNA stain to enable cell detection. The filters were

embedded in a solution containing the general DNA stain Hoechst (10 µg mL^{-1}) and the antifading

reagents Citifluor AF1 (Citifluor Ltd., PA; containing phosphate-buffered saline and glycerol) at 3

parts and Vectashield (Vecta Laboratories, CA) at 1 part. The cells on the embedded filters were

imaged with CLSM by measuring z-stacks that were adjusted to extend from the filter surface until

the top of the biggest cell including large algae and diatoms. For each sample, we obtained

mosaics with 4 tiles from about 3 marks. The acquired images had the following dimensions: z-

stack step-size of 2 µm (open pinhole) and xy-dimensions between 140 to 145 µm with a

resolution of 1024×1024 pixels, resulting in a pixel size of about 0.14 µm (similar to the pixel size

of NanoSims images). Images were recorded with bidirectional scanning at 400 Hz with a signal

depth of 16 bit and averaging between 3 frames. Fluorescence was recorded in 17 channels

varying in excitation (ex.; λ in nm) and emission (em.; λ in nm) during 9 sequential scans.

Channels were selected based on (i) lambda scans of emitted light from samples obtained from

different depths, (ii) spectra (in nm) of known photopigments (chlorophyll-a, chl-a; phycocyanin,

PC; phycoerythrin, PE; fucoxanthin, Fuxo; accessory and decaying pigments, ADP) and (iii) equal

fluorescence emission as recorded by the flow cytometer Cytobuoy. Scan 1: ex.: 633 / em.: 660-

688 (chl-a/PC) & 725-750; scan 2: ex.: 633 / em.: 668-734 (Cytobuoy 1; chl-a/PE); scan 3: ex.:

543 / em.: 660-700 (PE/Fuxo); scan 4: ex.: 543 / em.: 625-650 & 700-750; scan 5: ex.: 488 / em.:

601-668 (Cytobuoy 2; PC) & 536-601 (Cytobuoy 3; ADP); scan 6: ex.: 488 / em.: 670-734 (chl-

a/Fuxo) & 735-775; scan 7: ex.: 488 / em.: 518-548 & 565-605; scan 8: ex.: 458 / em.: 525-600 &

675-725 (chl-a) & reflected light; scan 9: ex.: 405 / em.: 423-503 (Hoechst stain) & reflected light.

Imaging was done from higher to lower excitation wavelengths to reduce photobleaching. Gains

were kept constant between all images. After imaging, the filters were washed in 100% ethanol

and double-deionized water.

Next, the filters were prepared for NanoSIMS analysis by sputter-coating with a conductive

gold layer (ca. 20 nm). Individual images or image mosaics were measured in the same regions

that were imaged with CLSM previously. The respective areas were located using the etched

marks and the CCD camera built into the NanoSIMS 50L. The measured areas were pre-

sputtered with a Cs^+ primary ion beam of about 4 pA to remove surface contamination, implant

Cs^+ ions, and achieve an approximately stable secondary ion emission rate. A primary Cs^+ ion

beam with a beam current between 1 and 1.2 pA and a beam diameter of around 150 nm was

rastered across the cells for analysis with a dwell time of 5 ms per pixel. Secondary ion images

for $^{12}C^{12}C^-$, $^{13}C^{12}C^-$, $^{12}C^{14}N^-$, $^{12}C^{15}N^-$, and $^{32}S^-$ were simultaneously recorded from analysis areas

of 30 μm × 30 μm with a resolution of 256×256 pixels, resulting in a pixel size of 0.117 μm. Ten

planes from each individual area were measured. Mass resolving power was around 10,000

(Cameca definition), enough to resolve all measured isotopes from potential interferences in the

mass spectrum.

Images were analyzed with a custom-written Matlab code that integrated Schnitzcells[69] with

Look@NanoSIMS[70]. First, a maximum intensity projection (MIP) image was created from the z-

stack of the individual Hoechst images (scan 9). These MIP images, in which all cells were

stained, were aligned with the corresponding NanoSIMS image in Look@NanoSIMS. The

alignment was applied to all fluorescence images. Next, the aligned MIP image of scan 9 was

used for automated segmentation of cells using Schnitzcells. This segmentation was manually

corrected with an overlay image of Hoechst fluorescence and the $^{12}C^{14}N^-$ (biomass) NanoSIMS

image for 268 to 662 cells per analyzed image. The segmentation was applied to all fluorescence

images and the fluorescence intensities and cell dimensions for all channels were compiled.

Finally, the segmentation was imported into Look@NanoSIMS to compile the corresponding

isotope counts, isotopic fractions, and cell dimensions for each segmented cell. C and N

assimilation for each segmented cell were calculated from their isotopic enrichment $^{15}N/(^{14}N+^{15}N)$

and $^{13}C/(^{12}C+^{13}C)$. Overall, images from 17 different samples coming from different days and

depths were analyzed corresponding to a total of 6440 cells.

Calculation of trait diversity

To obtain single-cell trait data, we combined stable isotope incubations - by adding ^{15}N-NH_4+

and ^{13}C-CO_2 into samples from different water depths and days - with correlative fluorescence

microscopy and NanoSIMS (see previous two sections, as well as Fig. 2A and B). For the

calculation of trait diversity, we first selected seven traits in total to avoid collinearity (Spearman

correlation <0.7) and thus assure trait complementarity. Four traits were extracted from CLSM

images: the mean fluorescence intensity of 3 channels: (i) excitation 543 nm; emission 700-750

nm, (ii), excitation 458 nm; emission 525-600 nm, (iii) excitation 458 nm; emission 675-725 nm,

and (iv) cell size. Three traits were derived from NanoSIMS data: C and N assimilation, as well

as S cell content. Those traits were then transformed to the same unit (mean=0, SD=1) and used

to calculate three complementary indices of trait diversity, representing the distribution of

individual cells of a microbial community in a high-dimensional trait space (Fig. 2C): TOP (*'Trait*

Onion Peeling')[71], TED (*'Trait Even Distribution'*)[71] and FDis (*'Functional Dispersion'*)[72]. TOP, as

a measure of trait richness, is sensitive to expansions and contractions of the trait space covered

by the individuals of a community. TED, as a measure of trait evenness, is designed to respond

to competitive interactions by quantifying niche partitioning as relative regularity in distances

between individuals in the trait space. FDis, as a measure of trait divergence, reflects the degree

to which individuals are spread around the average phenotype of the community. Please note

that the list of traits included in the calculation of trait diversity varied according to the aim (and

underlying hypotheses) of each single analysis, always being a selection of the above-mentioned

seven traits (see 'Statistical analysis' below for more details). To robustly perform such analyses

for a large array of samples and make trait diversity completely independent of the number of

cells, we first had to determine the number of single cells in each sample that were required to

detect differences in trait diversity indices between upper layer and chemocline (sensitivity

analysis). To do this, we selected the two samples (A3 and A13) with the highest number of

measured cells (662 and 540, respectively), and bootstrapped 100 times 50 to 450 cells (in

intervals of 50) to calculate trait diversity indices (including all above mentioned seven traits). We

then fitted linear regressions to test differences in trait diversity between the two samples. The

data show that differences for each trait diversity index can be resolved by measuring

approximately 250 cells (Extended Data Fig. 2). More specifically, goodness of fit and p-values

indicate that at least 200 cells are necessary to distinguish upper layer and chemocline

communities with a reasonable degree of confidence, especially with respect to the TED index of

trait evenness (Extended Data Fig. 2). As the analyzed 17 samples had at least 269 cells, we

decided to bootstrap 250 cells from each of them to further increase the representativeness of

our subsamples. For further analyses, we considered the average trait diversity values of the 999

bootstrapped communities.

Community analysis and calculation of genetic diversity

Samples for DNA extraction and amplicon sequencing were immediately filtered on 0.2 μm

polycarbonate filters with a diameter of 25 mm (Millipore). The filtration volume was adjusted to

filter approximately 10^7 to 10^8 cells on each filter, corresponding to 0.2 to 0.4 L per filter. Filters

were stored at -20°C until further processing. An empty polycarbonate filter was processed as a

negative control (DNA extraction and sequencing).

DNA was extracted from the filters using hot phenol/chloroform extraction. Cell lysis was

performed with glass beads (106 μm) in 450 μL SDS-lysis buffer (35 mM sodium acetate, 7 mM

EDTA, 2.9% SDS, pH 8) mixed with 1 mL phenol:chloroform:isoamyl alcohol (25:24:1 at pH 8) at

70°C for 12 min and bead beating for 2 min (Tissue Lyser II, Qiagen). The mixture was centrifuged

at 4°C for 3 min at maximum speed and the aqueous phase was transferred to a new reaction

tube. An equal volume of phenol:chloroform:isoamyl alcohol (25:24:1) was added to this tube.

This extraction and centrifugation process was repeated for two more rounds. The final round of

extraction was performed by adding chloroform:isoamyl alcohol (24:1) and again collecting the

aqueous phase (~200 µL). DNA was then precipitated from the recovered aqueous phase by

adding 0.1 volume of 3 M sodium acetate (pH 8), gently mixing, and adding 2 volumes of 100 %

ethanol and 1 µL of glycogen (20 mg/mL). Samples were incubated at 20°C for at least 3 hours,

centrifuged at 4°C for 60 min at maximum speed and the supernatant was removed. The pellet

was then washed twice with 200 µL of 80% ethanol, centrifuged at 4°C for 5 min and the

supernatant removed. The pellet was dried under vacuum for 7 min and dissolved in 25 µL

nuclease-free water. The DNA concentration was determined fluorometrically (Qbit, Invitrogen)

and the concentration was normalized between samples to a maximum of 20 ng/µL by the addition

of appropriate volumes of nuclease-free water.

Sequencing library preparation and DNA amplicon sequencing were performed by MR DNA

(Molecular Research LP, Shallowater, TX) using a modified bTEFAP® library preparation

protocol[73] and sequencing on an Illumina MiSeq platform with a minimum of 20,000 reads per

sample and read lengths of 2x300 bp. The 16S sequencing library was prepared with Bact341F:

5'-CCTACGGGNGGCWGCAG-3' and Bact785R:5'-GACTACHVGGGTATCTAATCC-3' [74], and

the 18S sequencing library was prepared with euk516f:GGAGGGCAAGTCTGGT and

euk1055r:CGGCCATGCACCACC[75]. A mock bacterial community was sequenced as a positive

control (BEI Resources, Genomic DNA from Microbial Mock Community B, staggered, high

Concentration, v5.2H, for Whole Genome Shotgun Sequencing, HM-277D).

We had to use two slightly different methods to process the sequence data of the two genes.

The detailed protocol files describing the application, version, and parameters of the data

processing for both genes can be found in the Appendix. In the first step, the sequence data were

filtered to remove PhiX residues and sequences of low complexity. For the 16S amplicon data,

read pairs were merged and primer site sequences were removed. For the 18S amplicon data,

only the primer site trimmed forward reads were used as the reads could not be merged. We used

UNOISE3[76] to obtain zOTUs (zero-radius OTUs), which are considered equivalent to amplicon

sequence variants (ASVs). We used SINTAX[76] in combination with the reference SILVA SSU

(v138 - https://www.arb-silva.de/) for 16S and pr2 SSU (v4.14.0 - https://pr2-database.org/) for

18S to predict the taxonomic affiliation of the zOTUs. The sequence data is deposited at

NCBI/ENA under the BioProject number PRJNA1014378.

Finally, we calculated genetic diversity indices (Chao1 and Simpson) using the library *phyloseq*[66]

from unclustered zOTUs data.

Statistical analysis

To test our first hypothesis, that ecosystem functioning is better explained by trait diversity than

taxonomic diversity, we used a model averaging approach[77]. To this end, we formulated a list of

linear models including as predictors all possible combinations (=16) of three trait diversity metrics

(TOP, TED and FDis, calculated using three pigment-related traits, cell size, C assimilation, N

assimilation, and S cell content) and day of sampling. In addition, in a second list of linear models

we included as predictors all possible combinations (=32) of four genetic diversity metrics (Chao1

and Simpson index using 16S and 18S sequencing data) and the day of sampling. For both lists

of models, the response variable was cell density relative to the maximum achieved in a given

layer. Models with a difference in the Akaike information criterion corrected for small sample sizes

(delta AICc<2) were defined as the most parsimonious set of models[78] and were consequently

used for model averaging (separately for trait and taxonomic diversity). Model selection and

averaging were performed using the *MuMIn* R-package[66]. To address our second hypothesis that

the highly competitive environment in deep layers of the lake drives increasing niche partitioning

in pigment-related traits, and decreasing fitness differences in metabolic traits, we performed one-

way ANOVA analyses and post hoc Tukey tests using the *TukeyHSD* function of the R-package

stats. This procedure takes multiple comparisons into account when calculating p-values[66]. This

allowed us to assess differences in the evenness of pigment-related traits and the divergence of

metabolic traits between the three layers. To test our third hypothesis that single-cell niche and

fitness differences are negatively related across depth, we fitted three simple linear models (*lm*

function) with the evenness of pigment-related traits as a predictor and the coefficient of variation

in S content and C and N assimilation rates as response variables.

Acknowledgements

We thank the personnel of Piora Centro Biologia Alpina, especially Mauro Tonolla, for making it

possible to use the laboratory facilities and housing during field work at lake Cadagno. We thank

Martin Ackermann, Jukka Jokela and Dave Johnson (all Eawag) for discussions and support

during the initial stages of the project. We thank Ilona Szivak (Eawag) for acquiring CLSM images.

We thank the AUA chemical analysis teaching laboratory at Eawag for chemical analysis of

sulfate, chloride, nitrate, ortho-phosphate, and dissolved inorganic carbon. We thank Nicola

Storelli for reviewing the manuscript. This study was supported by Eawag, by the Swiss National

Science Foundation Project 31003A_144053, an Eawag academic transition grant to F.S., a

Synthesis Grant of the ETH Zurich Center for Adaptation to a Changing Environment (ACE) to

F.S, and a Marie-Curie-Intra-European fellowship for career development (FP7-MC-

IEF;271929;Phenofix) provided by the European Union to F.S. The NanoSIMS instrument in the

Laboratory for Biological Geochemistry was funded in part by the European Union through a

European Research Council Advanced Grant 246749 (BIOCARB) to A.M.

Credit Authorship

Simone Fontana: Conceptualization, Methodology, Software, Formal analysis, Investigation,

Data Curation, Writing - Original Draft, Writing - Review & Editing, Visualization, Project

administration

Désirée A. Schmitz: Formal analysis (correlative imaging data), Data Curation, Writing -

Original Draft, Writing - Review & Editing, Visualization

Michael Daniels: Formal analysis (PCA analysis), Writing - Review & Editing, Visualization

Francesco Danza: Investigation (field work), Resources, Writing - Review & Editing

Thomas Röösli: Software (for correlative image analysis), Writing - Review & Editing

Hannah Bruderer: Methodology, Investigation, Writing - Review & Editing

Jean-Claude Walser: Formal analysis (sequencing data), Data Curation, Writing - Review &

Editing

Julien Dekaezemacker: Investigation (IRMS measurements), Writing - Review & Editing

Stephane Escrig: Investigation (NanoSIMS measurements), Writing - Review & Editing

Anders Meibom: Resources (NanoSIMS), Writing - Review & Editing, Supervision

Francesco Pomati: Conceptualization, Writing - Review & Editing, Supervision

Frank Schreiber: Conceptualization, Methodology, Formal analysis, Data Curation, Writing -

Original Draft, Writing - Review & Editing, Visualization, Supervision, Project administration,

Funding acquisition

References

1   Darwin, C. *On the origin of species*. (John Murray, 1859).

2   Grinnell, J. The Origin and Distribution of the Chest-Nut-Backed Chickadee. *The Auk* **21**,

726   364-382 (1904).

727   3   Gause, G. F. Experimental Studies on the Struggle for Existence: I. Mixed Population of

728   Two Species of Yeast. *Journal of Experimental Biology* **9**, 389-402 (1932).

729   4   Grinnell, J. The Niche-Relationships of the California Thrasher. *The Auk* **34**, 427-433

730   (1917).

731   5   Elton, C. *Animal Ecology*. (Sidgwick and Jackson, 1927).

732   6   Hutchinson, G. E. Concluding remarks. *Cold Spring Harbor Symposia on Quantitative*

733   *Biology* **22**, 415-427 (1957).

734   7   Hutchinson, G. E. The paradox of the plankton. *The American Naturalist* **XCV**, 137-145

735   (1961).

8   Chesson, P. Mechanisms of maintenance of species diversity. *Annual Review of*
Ecology and Systematics **31**, 343-366 (2000).

9   Chesson, P. Updates on mechanisms of maintenance of species diversity. *Journal of*
Ecology **106**, 1773-1794, doi:10.1111/1365-2745.13035 (2018).

10   Barabás, G., D'Andrea, R. & Stump, S. M. Chesson's coexistence theory. *Ecological*
Monographs **88**, 277-303 (2018).

11   Cadotte, M. W. Concurrent niche and neutral processes in the competition-colonization
model of species coexistence. *Proceedings of the Royal Society B: Biological Sciences*
274, 2739-2744, doi:10.1098/rspb.2007.0925 (2007).

12   Letten, A. D., Hall, A. R. & Levine, J. M. Using ecological coexistence theory to
understand antibiotic resistance and microbial competition. *Nature Ecology and*
Evolution **5**, 431-441, doi:10.1038/s41559-020-01385-w (2021).

13   Song, C., Barabás, G. & Saavedra, S. On the Consequences of the Interdependence of
Stabilizing and Equalizing Mechanisms. *The American Naturalist* **194**, 627-639,
doi:10.5061/dryad.j27h930 (2019).

14   Cordero, O. X. & Polz, M. F. Explaining microbial genomic diversity in light of
evolutionary ecology. *Nature Reviews Microbiology* **12**, 263-273,
doi:10.1038/nrmicro3218 (2014).

15   Krismer, J., Tamminen, M., Fontana, S., Zenobi, R. & Narwani, A. Single-cell mass
spectrometry reveals the importance of genetic diversity and plasticity for phenotypic
variation in nitrogen-limited Chlamydomonas. *The ISME Journal* **11**, 988-998,
doi:10.1038/ismej.2016.167 (2017).

16   Hart, S. P., Turcotte, M. M. & Levine, J. M. Effects of rapid evolution on species
coexistence. *Proceedings of the National Academy of Sciences of the United States of*
America **116**, 2112-2117, doi:10.1073/pnas.1816298116 (2019).

17   Ward, B. A. & Collins, S. Rapid evolution allows coexistence of highly divergent lineages
within the same niche. *Ecology Letters* **25**, 1839-1853, doi:10.1111/ele.14061 (2022).

18   Reiss, J., Bridle, J. R., Montoya, J. M. & Woodward, G. Emerging horizons in biodiversity
and ecosystem functioning research. *Trends in Ecology and Evolution* **24**, 505-514,
doi:10.1016/j.tree.2009.03.018 (2009).

19   Eisenhauer, N. *et al.* Biodiversity–ecosystem function experiments reveal the
mechanisms underlying the consequences of biodiversity change in real world
ecosystems. *Journal of Vegetation Science* **27**, 1061-1070, doi:10.1111/jvs.12435
(2016).

20   Turnbull, L. A., Levine, J. M., Loreau, M. & Hector, A. Coexistence, niches and
biodiversity effects on ecosystem functioning. *Ecology Letters* **16**, 116-127,
doi:10.1111/ele.12056 (2013).

21   Patsch, D., van Vliet, S., Marcantini, L. G. & Johnson, D. R. Generality of associations
between biological richness and the rates of metabolic processes across microbial
communities. *Environmental Microbiology* **20**, 4356-4368, doi:10.1111/1462-2920.14352
(2018).

22   Godoy, O., Gómez-Aparicio, L., Matías, L., Pérez-Ramos, I. M. & Allan, E. An excess of
niche differences maximizes ecosystem functioning. *Nature Communications* **11**, 4180-
4180, doi:10.1038/s41467-020-17960-5 (2020).

23   Violle, C. *et al.* Let the concept of trait be functional! *Oikos* **116**, 882-892,
doi:10.1111/j.2007.0030-1299.15559.x (2007).

24   Fontana, S., Rasmann, S., de Bello, F., Pomati, F. & Moretti, M. Reconciling trait based
perspectives along a trait-integration continuum. *Ecology* **102**, e03472-e03472,
doi:10.1002/ecy.3472 (2021).

25   Johnson, D. R. & Pomati, F. A brief guide for the measurement and interpretation of
microbial functional diversity. *Environmental Microbiology* **22**, 3039-3048,
doi:10.1111/1462-2920.15147 (2020).

26   Lajoie, G. & Kembel, S. W. Making the Most of Trait-Based Approaches for Microbial
Ecology. *Trends in Microbiology* **27**, 814-823, doi:10.1016/j.tim.2019.06.003 (2019).

27   Achtman, M. & Wagner, M. Microbial diversity and the genetic nature of microbial
species. *Nature Reviews Microbiology* **6**, 431-440, doi:10.1038/nrmicro1872 (2008).

28   Philippot, L. *et al.* The ecological coherence of high bacterial taxonomic ranks. *Nature*
Reviews Microbiology **8**, 523-529, doi:10.1038/nrmicro2367 (2010).

29   Chase, A. B. *et al.* Emergence of soil bacterial ecotypes along a climate gradient.
Environmental Microbiology **20**, 4112-4126, doi:https://doi.org/10.1111/1462-2920.14405
(2018).

30   Ackermann, M. A functional perspective on phenotypic heterogeneity in microorganisms.
Nature Reviews Microbiology **13**, 497-508, doi:10.1038/nrmicro3491 (2015).

31   Larkin, A. A. & Martiny, A. C. Microdiversity shapes the traits, niche space, and
biogeography of microbial taxa. *Environmental Microbiology Reports* **9**, 55-70,
doi:https://doi.org/10.1111/1758-2229.12523 (2017).

32   Bolnick, D. I. *et al.* Why intraspecific trait variation matters in community ecology. *Trends*
in ecology & evolution **26**, 183-192, doi:10.1016/j.tree.2011.01.009 (2011).

33 Krause, S. *et al.* Trait-based approaches for understanding microbial biodiversity and
ecosystem functioning. *Frontiers in Microbiology* **5**, 1-10, doi:10.3389/fmicb.2014.00251
(2014).
34 Fontana, S., Thomas, M. K., Moldoveanu, M., Spaak, P. & Pomati, F. Individual-level
trait diversity predicts phytoplankton community properties better than species richness
or evenness. *ISME Journal* **12**, doi:10.1038/ismej.2017.160 (2018).
35 Hoppe, P., Cohen, S. & Meibom, A. NanoSIMS: Technical Aspects and Applications in
Cosmochemistry and Biological Geochemistry. *Geostandards and Geoanalytical*
Research **37**, 111-154, doi:https://doi.org/10.1111/j.1751-908X.2013.00239.x (2013).
36 Margalef, R. Life-forms of phytoplankton as survival alternatives in an unstable
environment. *Oceanologica Acta* **1**, 493-509, doi:10.1007/BF00202661 (1978).
37 Laughlin, D. C. & Messier, J. Fitness of multidimensional phenotypes in dynamic
adaptive landscapes. *Trends in Ecology and Evolution* **30**, 487-496,
doi:10.1016/j.tree.2015.06.003 (2015).
38 Fontana, S., Thomas, M. K., Reyes, M. & Pomati, F. Light limitation increases
multidimensional trait evenness in phytoplankton populations. *ISME Journal* **13**, 1159-
1167, doi:10.1038/s41396-018-0320-9 (2019).
39 Lemmen, K. D., Butler, O. M., Koffel, T., Rudman, S. M. & Symons, C. C. Stoichiometric
traits vary widely within species: A meta-analysis of common garden experiments.
Frontiers in Ecology and Evolution **7**, 339-339, doi:10.3389/fevo.2019.00339 (2019).
40 Zaoli, S. *et al.* Generalized size scaling of metabolic rates based on single-cell
measurements with freshwater phytoplankton. *Proceedings of the National Academy of*
Sciences of the United States of America **116**, 17323-17329,
doi:10.1073/pnas.1906762116 (2019).
41 Schreiber, F. *et al.* Phenotypic heterogeneity driven by nutrient limitation promotes
growth in fluctuating environments. *Nature Microbiology* **1**, 16055-16055,
doi:10.1038/nmicrobiol.2016.55 (2016).
42 Delong, J. P. & Hanson, D. T. Metabolic rate links density to demography in
Tetrahymena pyriformis. *ISME Journal* **3**, 1396-1401, doi:10.1038/ismej.2009.81 (2009).
43 Pettersen, A. K., White, C. R. & Marshall, D. J. Metabolic rate covaries with fitness and
the pace of the life history in the field. *Proceedings of the Royal Society B: Biological*
Sciences **283**, 20160323-20160323, doi:10.1098/rspb.2016.0323 (2016).

44   Saini, J. S. *et al.* Bacterial, Phytoplankton, and Viral Distributions and Their
Biogeochemical Contexts in Meromictic Lake Cadagno Offer Insights into the
Proterozoic Ocean Microbial Loop. *mBio* **13**, 1-20, doi:10.1128/mbio.00052-22 (2022).
45   Fox, J. W. & Vasseur, D. A. Character convergence under competition for nutritionally
essential resources. *American Naturalist* **172**, 667-680, doi:10.1086/591689 (2008).
46   Tonolla, M. *et al.* in *Ecology of Meromictic Lakes*   (eds Ramesh D. Gulati, Egor S.
Zadereev, & Andrei G. Degermendzhi) 155-186 (Springer International Publishing,
2017).
47   Tonolla, M., Peduzzi, S., Hahn, D. & Peduzzi, R. Spatio-temporal distribution of
phototrophic sulfur bacteria in the chemocline of meromictic Lake Cadagno
(Switzerland). *FEMS Microbiology Ecology* **43**, 89-98, doi:10.1111/j.1574-
6941.2003.tb01048.x (2003).
48   Malik, A. A. *et al.* Defining trait-based microbial strategies with consequences for soil
carbon cycling under climate change. *ISME Journal* **14**, 1-9, doi:10.1038/s41396-019-
0510-0 (2020).
49   Schreiber, F. & Ackermann, M. Environmental drivers of metabolic heterogeneity in
clonal microbial populations. *Curr Opin Biotechnol* **62**, 202-211,
doi:10.1016/j.copbio.2019.11.018 (2020).
50   Louca, S. *et al.* Function and functional redundancy in microbial systems. *Nature
Ecology and Evolution* **2**, 936-943, doi:10.1038/s41559-018-0519-1 (2018).
51   Martiny, J. B. H., Jones, S. E., Lennon, J. T. & Martiny, A. C. Microbiomes in light of
traits: A phylogenetic perspective. *Science* **350**, doi:10.1126/science.aac9323 (2015).
52   Johnson, D. R. *et al.* Association of biodiversity with the rates of micropollutant
biotransformations among full-scale wastewater treatment plant communities. *Applied
and Environmental Microbiology* **81**, 666-675, doi:10.1128/AEM.03286-14 (2015).
53   Hart, S. P., Schreiber, S. J. & Levine, J. M. How variation between individuals affects
species coexistence. *Ecology Letters* **19**, 825-838, doi:10.1111/ele.12618 (2016).
54   Edwards, K. F. *et al.* Evolutionarily stable communities: a framework for understanding
the role of trait evolution in the maintenance of diversity. *Ecology Letters* **21**, 1853-1868,
doi:https://doi.org/10.1111/ele.13142 (2018).
55   Turcotte, M. M. & Levine, J. M. Phenotypic Plasticity and Species Coexistence. *Trends
in Ecology & Evolution* **31**, 803-813, doi:10.1016/j.tree.2016.07.013 (2016).

56   Gallego, I., Venail, P. & Ibelings, B. W. Size differences predict niche and relative fitness
differences between phytoplankton species but not their coexistence. *ISME Journal* **13**,
1133-1143, doi:10.1038/s41396-018-0330-7 (2019).

57   Musat, N. *et al.* A single-cell view on the ecophysiology of anaerobic phototrophic
bacteria. *Proceedings of the National Academy of Sciences of the United States of*
America **105**, 17861-17866, doi:10.1073/pnas.0809329105 (2008).

58   Danza, F., Storelli, N., Roman, S., Lüdin, S. & Tonolla, M. Dynamic cellular complexity of
anoxygenic phototrophic sulfur bacteria in the chemocline of meromictic Lake Cadagno.
PLoS ONE **12**, e0189510-e0189510, doi:10.1371/journal.pone.0189510 (2017).

59   Di Nezio, F. *et al.* Anoxygenic photo-and chemo-synthesis of phototrophic sulfur bacteria
from an alpine meromictic lake. *FEMS Microbiology Ecology* **97**, fiab010-fiab010,
doi:10.1093/femsec/fiab010 (2021).

60   Zimmermann, M. *et al.* Substrate and electron donor limitation induce phenotypic
heterogeneity in different metabolic activities in a green sulphur bacterium.
Environmental Microbiology Reports **10**, 179-183, doi:10.1111/1758-2229.12616 (2018).

61   Shan, X. & Cordero, O. X. Identifying microbial guilds on the basis of ecological patterns.
Nature Ecology & Evolution **7**, 651-652, doi:10.1038/s41559-023-02030-y (2023).

62   Shan, X., Goyal, A., Gregor, R. & Cordero, O. X. Annotation-free discovery of functional
groups in microbial communities. *Nature Ecology & Evolution* **7**, 716-724,
doi:10.1038/s41559-023-02021-z (2023).

63   Guerra, C. A. *et al.* Global projections of the soil microbiome in the Anthropocene.
Global Ecology and Biogeography **30**, 987-999, doi:10.1111/geb.13273 (2021).

64   Eisenhauer, N. *et al.* The heterogeneity-diversity-system performance nexus. *National*
Science Review **10**, nwad109-nwad109, doi:10.1093/nsr/nwad109 (2023).

65   Hess, C., Levine, J. M., Turcotte, M. M. & Hart, S. P. Phenotypic plasticity promotes
species coexistence. *Nature Ecology & Evolution* **6**, 1256-1261, doi:10.1038/s41559-
022-01826-8 (2022).

66   Team, R. C. in *R Foundation for Statistical Computing*   (R Foundation for Statistical
Computing, Vienna, Austria, 2022).

67   Breiman, L. Random forests. *Machine Learning* **45**, 5-32, doi:10.1023/A:1010933404324
(2001).

68   Liaw, A. & Wiener, M. Classification and Regression by randomForest. *R News* **2**, 18-22,
doi:10.1177/154405910408300516 (2002).

69 Young, J. W. *et al.* Measuring single-cell gene expression dynamics in bacteria using

fluorescence time-lapse microscopy. *Nature Protocols* **7**, 80-88,

doi:10.1038/nprot.2011.432 (2011).

70 Polerecky, L. *et al.* Look@NanoSIMS-a tool for the analysis of nanoSIMS data in

environmental microbiology. *Environmental Microbiology* **14**, 1009-1023,

doi:10.1111/j.1462-2920.2011.02681.x (2012).

71 Fontana, S., Petchey, O. L. & Pomati, F. Individual-level trait diversity concepts and

indices to comprehensively describe community change in multidimensional trait space.

Functional Ecology **30**, 808–818-808–818, doi:10.1111/1365-2435.12551 (2016).

72 Laliberté, E. & Legendre, P. A distance-based framework for measuring functional

diversity from multiple traits. *Ecology* **91**, 299-305 (2010).

73 Dowd, S. E., Sun, Y., Wolcott, R. D., Domingo, A. & Carroll, J. A. Bacterial Tag–

Encoded FLX Amplicon Pyrosequencing (bTEFAP) for Microbiome Studies: Bacterial

Diversity in the Ileum of Newly Weaned Salmonella-Infected Pigs. *Foodborne Pathogens*

and Disease **5**, 459-472, doi:10.1089/fpd.2008.0107 (2008).

74 Klindworth, A. *et al.* Evaluation of general 16S ribosomal RNA gene PCR primers for

classical and next-generation sequencing-based diversity studies. *Nucleic Acids*

Research **41**, e1-e1, doi:10.1093/nar/gks808 (2013).

75 Stern, R. *et al.* Composition and Patterns of Taxa Assemblages in the Western Channel

Assessed by 18S Sequencing, Microscopy and Flow Cytometry. *Journal of Marine*

Science and Engineering **11**, 480-480, doi:10.3390/jmse11030480 (2023).

76 Edgar, R. C. SINTAX: a simple non-Bayesian taxonomy classifier for 16S and ITS

sequences. *bioRxiv*, 074161-074161, doi:10.1101/074161 (2016).

77 Burnham, K. P. & Anderson, D. R. *Model Selection and Multimodel Inference*. (Springer,

2002).

78 Symonds, M. R. E. & Moussalli, A. A brief guide to model selection, multimodel inference

and model averaging in behavioural ecology using Akaike's information criterion.

Behavioral Ecology and Sociobiology **65**, 13-21, doi:10.1007/s00265-010-1037-6 (2011).