Dual action of sphinganine in the plant disease resistance to bacteria

Huby Eloïse[1,2], Villaume Sandra[1], Chemotti Catherine[2], Dorey Stéphan[1], Cordelier Sylvain[1], Crouzet Jérôme[1], Gilliard Guillaume[2], Terryn Christine[3], Berquand Alexandre [4], Herrfurth Cornelia[5,6], Feussner Ivo[5,6,7], Jacquard Cédric[1], Fontaine Florence[1], Clément Christophe[1], Baillieul Fabienne[1], Deleu Magali[2*], Dhondt-Cordelier Sandrine[1*]

[1]Université de Reims Champagne-Ardenne, INRAE, RIBP USC 1488, 51100 Reims, France.
[2]Laboratoire de Biophysique Moléculaire aux Interfaces (LBMI), Gembloux Agro-Bio Tech, Université de Liège, 2 Passage des déportés, 5030 Gembloux, Belgium.
[3]Université de Reims Champagne-Ardenne, PICT Platform, 51100 Reims, France.
[4]Université de Reims Champagne-Ardenne, Laboratoire de Recherche en NanoScience EA4682/Nano'Mat, 51100 Reims, France.
[5]University of Goettingen, Albrecht-von-Haller-Institute for Plant Sciences, Department of Plant Biochemistry, Justus-von-Liebig-Weig 11, D-37077, Goettingen, Germany.
[6]University of Goettingen, Goettingen Center for Molecular Biosciences (GZMB), Service Unit for Metabolomics and Lipidomics, Justus-von-Liebig-Weg 11, D-37077, Goettingen, Germany.
[7]University of Goettingen, Goettingen Center for Molecular Biosciences (GZMB), Department of Plant Biochemistry, Justus-von-Liebig-Weg 11, D-37077, Goettingen, Germany.

* MD and SDC contributed equally

* Corresponding authors: Sandrine Dhondt-Cordelier (sandrine.cordelier@univ-reims.fr) and Magali Deleu (Magali.Deleu@uliege.be)

Short Title: Sphingolipids in plant defense responses

ABSTRACT

Sphingolipids are ubiquitous, highly diverse molecules constituting at least 40% of plant plasma membranes. Initially known as modulators of membrane integrity, they now emerge as important players in plant responses to (a)biotic stresses. The interaction between *Arabidopsis thaliana* and the bacterium *Pseudomonas syringae* pv. *tomato* DC3000 *AvrRpm1* (*Pst AvrRpm1*) culminates in the activation of a programmed cell death known as the hypersensitive response, which is part of the plant immune response. In this study, we showed that the co-infiltration of *Pst AvrRpm1*

and sphinganine (d18:0) in *Arabidopsis* leaves suppress the hypersensitive response.

This suppression phenotype is also observed with bacteria carrying the effectors AvrB

and AvrPphB but not with the ones carrying AvrRpt2 and AvrRps4. Sphingolipid-

induced hypersensitive response suppression by *Pst AvrRpm1* is correlated with the

down-regulation of the gene *AtNMT1* encoding a *N*-myristoyltransferase. d18:0 does

not have a direct antibacterial effect and its co-infiltration in plants does not display

typical signs of immune response such as activation of salicylic acid signaling pathway

and extracellular reactive oxygen species production. Biophysical studies showed that

d18:0 interacts with plant plasma membrane lipids. More specifically, d18:0 disturbs

plant plasma membrane organization and mechanical properties. Our results

demonstrate that sphingolipids play an important role in plant resistance, especially by

interfering with the plasma membrane organization and effector localization and thus

disturbing their function and subsequent immune responses.

Keywords: sphingolipids, biophysics, plant immunity, effectors, plasma membrane

INTRODUCTION

Recognition of pathogenic microorganisms leading to plant immunity is commonly divided into two pathways that can mutually potentiate each other (Ngou *et al*., 2021). Pathogen perception involves the recognition of pathogen-associated molecular patterns (PAMPs) through plant cell surface-anchored pattern recognition receptors (PRRs), which induces pattern-triggered immunity (PTI) (Saijo and Loo, 2020). However, some bacteria can interfere with PTI by delivering effectors into the plant cell to block signaling mechanisms at different steps (Desveaux *et al*., 2006; Kamoun, 2006; Dodds *et al*., 2009). The recognition of microbial effectors or virulence factors by the plant cells through intracellular nucleotide-binding leucine-rich repeat receptors (NLRs) induces effector-triggered immunity (ETI). PTI and ETI share many signals and components, but their signaling pathways are induced at different scales and time (Bjornson and Zipfel, 2021; Kadota *et al*., 2019; Yuan *et al*., 2021). Typical overlaps in signaling components are e.g., the production of reactive oxygen species (ROS), the influx of calcium, or mitogen-activated protein kinase (MAPK) signaling cascades. They also share some immune responses, such as transcriptional reprogramming and alterations in stress-related hormone homeostasis, including salicylic acid (SA), jasmonic acid (JA), and/or ethylene (ET) (Bari and Jones, 2009; Betsuyaku *et al*., 2018; Guan *et al*., 2015). Rapid programmed cell death (PCD) at the site of infection, known as the hypersensitive response (HR), usually is a hallmark of the ETI pathway (Couto and Zipfel, 2016; Henry *et al*., 2013; Kadota *et al*., 2019; Ngou *et al*., 2021; Yuan *et al*., 2021).

The bacterial pathogen *Pseudomonas syringae* pv. *tomato* DC3000 (*Pst* DC3000) has been extensively studied due to its capacity to cause bacterial speck in both tomato and *Arabidopsis thaliana* (hereafter *Arabidopsis*) (Lee *et al*., 2015; Wei *et al*., 2018),

as well as the availability of its complete genome sequence (Buell *et al.*, 2003).

Transformation of *Pst* DC3000 to express effectors has been essential for the

discovery of mechanisms associated with ETI (Boyle and Martin, 2015; Hicks and

Galán, 2013; Nimchuk *et al.*, 2000). AvrRpm1 effector was originally identified in *P.*

syringae pv. *maculicola* strain m2 (Debener *et al.*, 1991) and harbors an N-terminal

eukaryotic consensus site for myristoylation required for its full avirulence function. This

allows the effector, once acylated by the plant host, to be redirected to the plant plasma

membrane (PM) (Boyle and Martin, 2015; Hicks and Galán, 2013; Nimchuk *et al.*,

2000). Although the host acylation apparatus is initially exploited by bacteria to interfere

with PTI by binding to the kinase domain of PRRs (Boyle and Martin, 2015), in resistant

plants, it can actually turn against the pathogen. For instance, the NLR RPM1

(RESISTANCE TO *PSEUDOMONAS SYRINGAE* PV *MACULICOLA*) in *Arabidopsis*,

is monitored by the PM-associated protein RPM1-INTERACTING PROTEIN-4 (RIN4),

which guards it against effector-mediated perturbations (Boyes *et al.*, 1998; Jones and

Dangl, 2006; Mackey *et al.*, 2002). Therefore, once relocated to the PM, AvrRpm1 will

target RIN4, inducing its phosphorylation. This activates RPM1 which triggers the ETI

and the HR, together interrupting the bacterial spread of the and leading to the

degradation of RPM1 (Boyes *et al.*, 1998; Mackey *et al.*, 2002; Xin *et al.*, 2018).

Sphinganine is a sphingolipid (SL), belonging to the long-chain base (LCB) class,

which are one major building block of SLs (Pata *et al.*, 2010). It consists of a C18 core

with an amino group at C2 and two hydroxylations at C2 and C3 (Luttgeharm *et al.*,

2016; Mamode Cassim *et al.*, 2019). SLs are key components of the plant PM

structure, accounting for up to 40 mol% of its external leaflet (Cacas *et al.*, 2016). They

are found mostly in lipid microdomains, and play an important part in signaling, protein

aggregation, and stress response (Cacas *et al.*, 2016; Grennan, 2007; Mamode

Cassim *et al.*, 2019; Pata *et al.*, 2010). Recent studies have shown that SLs are key

players in PCD regulation occurring either during plant development or immunity (Ali

et al., 2018; Berkey *et al.*, 2012; Huby *et al.*, 2020; Lenarčič *et al.*, 2017). For instance,

Bax-inhibitor-1 interacts with enzymes of the SL biosynthetic pathway FAH1 and 2

(FATTY ACID HYDROXYLASE 1 and 2). It leads to changes in lipid content of

membrane microdomains (Lenarčič *et al.*, 2017). This causes the loss of proteins

normally located in these microdomains and which are essential for plant defense,

especially for cell death by oxidative stress or SA (Ishikawa *et al.*, 2015). Further links

between PCD, sphingolipids and the SA pathway have recently been highlighted. In

Arabidopsis, triple mutations of enzymes of the SL biosynthetic pathway, FAH1, FAH2

and LOH2 (LAG ONE HOMOLOG 2) led to the accumulation of sphinganine and

ceramides, another class of SLs. These mutants also exhibited a PCD that was

dependent on SA and was probably linked to d18:0 accumulation (König *et al.*, 2021,

2012).

Sensing of pathogen-derived lipid molecules is far from being elucidated. A recent

study demonstrated that the clivage product of the *Phytophthora infestans* ceramide D

by the neutral ceramidase 2 is recognized by the PRR RESISTANT TO DFPM-

INHIBITION OF ABSCISSIC ACID SIGNALING 2 (RDA2) (Kato *et al.*, 2022). Some

studies suggest that lipidic elicitors can directly interact with lipids in the plant PMs

(Cordelier *et al.*, 2022). For instance, ergosterol, the main sterol of fungi, is described

as a general elicitor of plant defenses (Klemptner *et al.*, 2014). Due to its lipidic nature

and its capacity to form rafts, it is thought to interact with the PM. In *Beta vulgaris,* it

can inhibit its H^+-ATPase activity, supposedly by tempering with the NADPH oxidase

present in the plant microdomains (Hao *et al.*, 2014; Rossard *et al.*, 2010).

While there are abundant reports on the links between LCBs and PCD (Ali *et al*., 2018;

Huby *et al*, 2020), only few data on the use of exogenously applied LCBs and their

mode of action are available (Glenz *et al*., 2019; Gutiérrez-Nájera *et al*., 2020; Magnin-

Robert *et al*., 2015; Peer *et al*., 2011; Saucedo-García *et al*., 2011, 2023; Shi *et al*.,

2007). In previous studies, we and others have shown that the co-infiltration of

Arabidopsis leaves with an LCB and *Pst AvrRpm1* does not induce an HR (Glenz *et*

al., 2022; Magnin-Robert *et al*., 2015). In order to decipher the underlying mechanism,

the activation of canonical defense responses and antibacterial effect of the LCB were

analyzed. Moreover, we showed similar phenotype with other effectors that share the

myristoylation sequence with AvrRpm1. Finally, biophysical analyses demonstrated

that sphinganine can interact with the plant biomimetic PM and disturb its organization

as well as its mechanical features.

RESULTS

Effect of *Pst AvrRpm1* and sphinganine co-infiltration on *Arabidopsis* cell death

We and others previously demonstrated that the co-infiltration of *Pst AvrRpm1* and d18:0 or t18:0, respectively, does not induce an HR in contrast to what happens with an infiltration of an avirulent bacteria or LCB alone (Magnin-Robert *et al.*, 2015; Glenz *et al.*, 2022). Therefore, *Pst* expressing effectors other than AvrRpm1 were selected to elucidate whether the absence of visible HR was dependent on the effector. AvrRpt2 was chosen because, like AvrRpm1, it is associated to the PM with RIN4 (Boyes *et al.*, 1998; Mackey *et al.*, 2003; Xin *et al.*, 2018). AvrRps4 was also tested because of the cytoplasmic localization of its target, RPS4 (Gassmann *et al.*, 1999; Wirthmueller *et al.*, 2007).

As expected, treatment with *Pst* DC3000 co-infiltrated or not with d18:0 triggered typical disease symptoms and infection with *Pst AvrRpm1*, *Pst AvrRpt2* or *Pst AvrRps4* alone, gave rise to a typical HR. Whereas co-infiltration of *Pst AvrRpm1* and d18:0 did not induce a visible HR as expected, the other tested bacteria displayed an HR when co-infiltrated with the sphingolipid (Fig. 1A). Changes in conductivity were then used as a marker of plant cell death, as PM damages result in strong electrolyte leakage (Kawasaki *et al.*, 2005) (Fig. 1B). All treatments triggered an increase in conductivity between 4 and 24 hpi, except for co-infiltration with *Pst AvrRpm1* and d18:0, displaying a conductivity stable over time and similar to the control plants (Fig. 1B). These findings are consistent with our previous study (Magnin-Robert *et al.*, 2015) and corroborated with phenotypic observations in Fig. 1A.

***In vitro* and *in planta* growth of *Pst AvrRpm1* in presence of sphinganine**

As a potential deleterious effect of d18:0 on the bacteria could not be excluded, *in vitro*

bacterial growth measurements were undertaken with or without the sphingolipid in the

culture medium (Fig. 2A). Ethanol had no effect on bacterial growth of all tested strains

(data not shown). Interestingly, d18:0 had no antibacterial effect on *Pst AvrRpm1* and

out of all the bacteria tested, d18:0 slightly reduced the growth rate of *Pst AvrRpt2*.

Surprisingly, this bacterium does provoke cell death when co-infiltrated into leaves.

To determine the effect of the co-infiltration on bacterial growth *in planta*, bacterial

populations were quantified at 0 and 48 hpi (Fig. 2B). The growth of the avirulent strain

Pst AvrRpm1 was slightly reduced compared to the virulent strain, *Pst* DC3000 at 48

hpi. Furthermore, while the co-infiltration had no consequence on the development of

Pst DC3000, it slightly reduced *Pst AvrRpm1* growth at 48 hpi. This bacterial count was

however two-fold higher at 48 hpi than at 0 hpi, like in the other conditions (Fig. 2B).

Collectively, these data suggest that d18:0 does not have an antibacterial effect that

could explain the absence of HR when *Pst AvrRpm1* is co-infiltrated with d18:0.

Moreover, while *Pst AvrRpm1* continues its development inside the plant, it is however

incapable of triggering an HR.

Extracellular ROS production by *Arabidopsis* in response to infection by *Pst*

Strong and sustained extracellular ROS production is an early defense event generally

linked to HR. We therefore monitored their production by *Arabidopsis* in response to

Pst AvrRpm1, *Pst AvrRpt2* and *Pst AvrRps4* (Fig. 3). No significant difference was

found between ROS production triggered by bacteria alone or mixed with ethanol (data

not shown). As already observed by Magnin-Robert *et al.* (2015), co-treatment of *Pst*

AvrRpm1 but also of *Pst AvrRpt2* and *Pst AvrRps4* with d18:0 reduced extracellular

ROS production by approximatively 6-, 2- and 1.7-fold, respectively (Fig. 3). Only the

co-infiltration of *Pst AvrRpm1* and d18:0 did not result in an HR (Fig. 1) but ROS levels

appear to be dramatically affected even with other bacteria like *Pst AvrRps4*. These

results suggest that reduced ROS production is not directly linked to the absence of

HR.

Effects of *Pst AvrRpm1* and sphinganine co-infiltration on *Arabidopsis* late

defense responses

Considering the effects of d18:0 on ROS production and cell death development, other

markers of plant defense were studied in response to co-infiltration. The accumulation

of the phytoalexin camalexin as well as the hormones SA and JA-Ile was followed. The

expression of *PATHOGENESIS-RELATED GENE 1* (*PR1*) and *VEGETATIVE*

STORAGE PROTEIN 1 (*VSP1*), marker genes of SA and JA-Ile signaling pathways,

respectively, was also analyzed. Control infiltration did not induce camalexin

production (Fig. 4A). At 48 hpi, *Pst* DC3000 inoculation with or without d18:0 triggered

similar camalexin accumulation. However, d18:0 co-infiltration significantly reduced

camalexin production in response to *Pst AvrRpm1* since it decreased from ~250 nmol/g

FW to a similar level observed in controls (~3 nmol/g FW) at 48 hpi (Fig. 4A).

Overexpression of *PR1* was observed 48h after infiltration with each *Pst* strain alone

(Fig. 4C), which correlated with the accumulation of SA (Fig. 4E). Neither SA

accumulation nor *PR1* expression was induced at earlier time points. The use of an

Arabidopsis reporter line *PR1::GUS* confirmed the induction of *PR1* expression in

treated leaves (Fig. 4B). Both SA accumulation and *PR1* overexpression are repressed

when leaves are co-infiltrated with d18:0, leading to similar levels as in control plants

(Fig. 4, C and E). *VSP1* is weakly expressed in response to *Pst* DC3000 but

overexpressed when this bacterium is co-infiltrated with d18:0 (~ 9-fold and ~ 6-fold

more at 6 and 48 hpi, respectively) (Fig. 4D). JA-Ile levels are however higher with the

bacterium alone than co-infiltrated (Fig. 4F). When *Arabidopsis* was inoculated with

Pst AvrRpm1, *VSP1* was slightly overexpressed at 48 hpi but repressed when co-

infiltrated with d18:0 (Fig. 4D). JA-Ile levels were concomitantly low (Fig. 4F), as

expected from plants challenged with avirulent *P. syringae* (Mur *et al.*, 2006). Overall,

co-infiltration of *Pst AvrRpm1* with d18:0 significantly lowers the amount of camalexin

and both defense hormone signaling pathways.

Involvement of RPM1 and RIN4 in the absence of HR

Once injected into the cytoplasm of the host, AvrRpm1 triggers the phosphorylation of

RIN4, inducing the activation of RPM1 and subsequently the ETI leading to an HR

(Nimchuck *et al.*, 2000). To elucidate whether RPM1 and / or RIN4 were involved when

the bacterium is co-infiltrated with d18:0, inoculations of *Pst AvrRpm1* were performed

on *Arabidopsis* mutants *rpm1-3,* and *rpm1-rps2-rin4,* as well as the complemented line

RPM1-myc rpm1. Infiltration of the bacteria alone induced cell death and co-infiltration

of the complemented line displayed no symptoms, as expected (Fig. 5A). Despite

lacking RPM1 and / or RIN4, all mutants showed the same phenotype as the WT when

co-infiltrated with d18:0 as they did not exhibit visible cell death (Fig. 5A). It has been

previously demonstrated that RPM1 is degraded during the first hpi, coincidently with

the onset of the HR (Boyes *et al.*, 1998). To further elucidate the role of RPM1 and

RIN4, *RPM1-myc rpm1* plant was used to monitor the degradation of RPM1 following

inoculation (Fig. 5B). Western blot analysis showed that while RPM1 was degraded

from 4 to 6 hpi when *Pst AvrRpm1* was infiltrated alone, the protein was still detected

24 hpi when the bacteria was co-infiltrated (Fig. 5B). Altogether, these data suggest

that the mechanism underlying the absence of HR occurs upstream of RPM1

degradation.

Effect of *Pst AvrB* and *Pst AvrPphB* co-infiltration with sphinganine on

***Arabidopsis* defense response**

The three effectors AvrRpm1, AvrB and AvrPphB share an eukaryotic *N*-myristoylation

site in their protein sequences that allows them to be redirected to the PM once they

have entered their host (Nimchuk *et al.*, 2000). Like AvrRpm1, AvrB interacts with

RPM1 and RIN4 (Grant *et al.*, 1995; Lee *et al.*, 2004), and AvrPphB with RPS5 and

PBS1, both located at the PM (Ade *et al.*, 2007; Pottinger and Innes, 2020). As already

demonstrated (Warren *et al.*, 1998; Whalen *et al.*, 1991), infiltration of *Pst AvrB* or *Pst*

AvrPphB triggers an HR. No cell death was visible when both *Pst AvrB* and *Pst*

AvrPphB were co-infiltrated with d18:0, suggesting HR was not triggered in treated

plants, similarly to *Pst AvrRpm1* (Fig. 6A). Further tests were conducted to determine

if d18:0 had an *in vitro* effect on these bacteria. Their growth rate remained unchanged

after d18:0 addition (Fig. 6B), confirming that d18:0 has no antibacterial effect on these

two bacteria. Extracellular ROS production by *Arabidopsis* in response to *Pst AvrB* and

Pst AvrPphB was also monitored (Fig. 6C). As for *Pst AvrRpm1*, d18:0 significantly

reduced this production in response to both bacteria (~ 2-fold for *Pst AvrB* and ~ 3-fold

for *Pst AvrPphB*).

***NMT1* relative expression following *Pst AvrRpm1* and sphinganine co-infiltration**

In *Arabidopsis*, two enzymes, NMT1 and NMT2, are responsible for *N*-myristoylation

(Pierre *et al.*, 2007). As the corresponding mutants displaying severe phenotypes, they

were thus not suitable for our experiments. We used qRT-PCR to determine

expression patterns of *NMT1* and *NMT2* in response to infiltration of *Arabidopsis* WT

leaves with *Pst* DC3000 or *Pst* AvrRpm1, either alone or with d18:0 (Fig. 7).

Unfortunately, the expression of *NMT2* was too low to be detectable. At 6 hpi, *NMT1*

expression in plants infiltrated with *Pst* DC3000 alone or with d18:0 was similar to that

of control samples but was induced at 48 hpi (~ 2-fold). When plants were infiltrated

with *Pst* AvrRpm1 alone, a stimulation of *NMT1* expression is observed during the time-

course experiment to reach a ~2.2-fold induction at 48 hpi (Fig. 7). However, when

plants were co-infiltrated with *Pst* AvrRpm1 and d18:0, no induction could be observed

during the time-course experiment (Fig. 7). Thus, a decrease of ~1.7-fold is observed

48 hpi between *Pst* AvrRpm1 infiltrated alone and with d18:0. Overall, these results

suggest that co-infiltration with d18:0 modifies *NMT1* expression in response to

infection with *Pst* AvrRpm1.

Interaction capacity of sphinganine with plant plasma membrane-mimicking

models

It has been shown that LCB sphingosine (d18:1) can increase the permeability of

liposomal and erythrocyte ghost membranes to aqueous solutes (Contreras *et al.*,

2006). Based on this finding, d18:0 was hypothesized to interact with the lipids of plant

PM. To investigate this hypothesis, we conducted *in silico* and *in vitro* biophysical

analyses using membrane plant models. Thanks to their defined structure and

controlled lipid composition, these models were found to be useful in understanding

how external molecules affect the organization and mechanical properties of the

membranes (Deleu *et al.*, 2014; Furlan *et al.*, 2020; Cordelier *et al.*, 2022).

The Impala modeling method was used to predict the ability of d18:0 to insert into an

implicit model lipid bilayer and to determine the preferable location and positioning of

d18:0 within a membrane bilayer (Fig. 8). The more negative the energy is, the more

favorable the insertion will be. A sharp drop of energy was observed as d18:0 moved

from the aqueous phase to the membrane surface (up to ~-10 Kcal/mol), indicating that

it could easily insert into membranes. The energy was close to 0 Kcal/mol in the

hydrophobic region of the membrane, suggesting that d18:0 has a less favorable

interaction with this region compared with the lipid bilayer hydrophilic region (Fig. 8A).

Nevertheless, the non-positive energy value in the hydrophobic region of the bilayer

suggests that d18:0 could flip-flop between the two bilayer leaflets. A large increase in

energy (from ~-10 Kcal/mol to +6.7 Kcal/mol) is also observed when d18:0 moves from

the inner bilayer leaflet to the aqueous medium, which strongly suggests that d18:0

has no propensity to cross the membrane and enter into the cytoplasm. The preferred

positioning of d18:0 within the membrane is shown in Fig. 8B. Its lipid chain lies within

the hydrophobic part of the membrane while the amine and alcohol groups are located

in the hydrophilic part of the bilayer. Altogether, this indicates that d18:0 can interact

with membranes and remains there. To support these simulation results, experimental

analyses were performed using large unilamellar vesicles (LUVs) that mimic plant PM,

consisting of palmitoyl-2-linoleoyl-*sn*-glycero-3-phosphocholine (PLPC),

glucosylceramide (GluCer) and β-sitosterol (sito) as the plant PM model phospholipid,

sphingolipid and sterol respectively (Deleu *et al*., 2014; Rondelli *et al*., 2021; Deboever

et al., 2022).

Results from ITC experiments showed a negative and decreasing heat flow after each

LUV injection into the d18:0 solution (Fig. 9A), indicating that d18:0 interacts with plant

PM-mimicking LUVs. The thermodynamic parameters calculated from the fitting of raw

data are presented in Fig. 9B. The binding coefficient K was relatively high (>80 mM^{-1})

suggesting that d18:0 has a strong affinity for plant-mimicking lipid membranes

confirming the low probability that it emerges in the intracellular medium. This affinity

for plant PM-like liposomes is higher than that observed for other plant-eliciting

molecules such as linoleic and linolenic acid hydroperoxides (<3 mM^{-1}; Deleu *et al.*,

2019), and rhamnolipid bolaforms (7.5 mM^{-1}; Luzuriaga-Loaiza *et al.*, 2018). The

binding reaction of d18:0 to LUVs was spontaneous as the molar free energy ΔG was

negative, exothermic as the molar enthalpy change ΔH was negative and generated a

positive change of the molar entropy (TΔS>0). The higher absolute value of the molar

entropy (TΔS>0) compared to the absolute value of the molar enthalpy change (ΔH),

indicates that the interactions is entropy-driven, i.e. mostly of hydrophobic nature.

Altogether, these results suggest that d18:0 is attracted by and can interact with plant

lipid membranes.

Lipid specificity of the interaction between sphinganine and plant plasma

membrane-mimicking models

To investigate specific interactions between d18:0 and lipids representative of plant

PM, the Langmuir trough technique was used to analyze the adsorption capacity of

d18:0 into monolayers of each individual lipid. In this technique, a lipid monolayer is

initially formed at an air-water interface, and then d18:0 is injected into the subphase

beneath the monolayer. The increase in surface pressure (ΔΠ) at a constant trough

area was monitored to determine the adsorption of d18:0 into the lipid monolayer at

different initial surface pressures (Π$_i$) (Supplemental Fig. S1). Two parameters were

determined from these data (Fig. 10A): (i) the differential surface pressure (dΠ$_0$), which

provides information on the attractive effect of the lipids on d18:0, and (ii) the maximal

insertion pressure (MIP), which reflects the ability of d18:0 to penetrate into the lipid

monolayer (Deleu *et al.*, 2014). Upon insertion of d18:0 onto monolayers formed with

individual plant PM lipids, $d\Pi_0$ values were all positive and similar for all compositions,

indicating that all tested lipids exert a similar attraction on d18:0 (Fig. 10A). MIP values

were all higher than the lateral pressure supposed to prevail into biological membranes

(30 – 35 mN/m) (Marsh, 1996), meaning that d18:0 may be able to adsorb into natural

plant membranes.

Lipid specificity of the interaction was also studied by an *in silico* approach using the

Hypermatrix docking method. This method calculates the energy of interaction

between d18:0 and individual plant lipids at a hydrophilic/hydrophobic interface,

indicating whether this interaction is energetically favorable or not (Brasseur *et al.*,

1987; Deleu *et al.*, 2014). Results showed that the interaction between d18:0 and all

plant lipids was favorable as all interaction energies are negative (Fig. 10B).

Interestingly, the interaction energies between d18:0 and plant lipids were more

favorable than those observed for d18:0 with itself. With this method, interaction

between d18:0 and PLPC or GluCer appeared to be more favorable than with β-

sitosterol.

Altogether, these data showed that d18:0 is attracted by and could interact with lipids

of plants. There is however no clear trend to determine a preferential interaction of

d18:0 with a particular lipid.

Modification of the organization and mechanical properties of plant plasma

membrane-mimicking models induced by sphinganine

Since d18:0 can interact with lipids found in plant PMs, additional experiments were

conducted to investigate how these interactions could impact the mechanical

properties and the organization of plant PMs.

Two aspects of membrane mechanics were considered: membrane fluidity and

membrane elasticity. Membrane fluidity refers to the degree of lipid packing and the

mobility or dynamics of lipids within the membrane, which can be characterized by the

order level parameter (Gerbeau-Pissot *et al*., 2014). Membrane elasticity

corresponding to the resistance of a material to deformation when subjected to an

applied force was determined by measuring the Young's modulus. The higher the

Young's modulus, the stiffer and less deformable the object is (Magazzu and

Marcuello, 2023). Although membrane fluidity can be influenced by the membrane

elasticity, there is no direct quantitative relationship between Young's modulus and

membrane fluidity or order level and therefore must be considered as separate aspects

of membrane behavior.

To analyze the order level of *Arabidopsis* PM in response to d18:0, *Arabidopsis* leaf

protoplasts were stained with the dye di-4-ANEPPDHQ. The order level of the

membrane, revealed by fluorescence imaging, was quantified using the red/green ratio

of the membrane (RGM) (Dinic *et al*., 2011; Gerbeau-Pissot *et al*., 2014). Observations

of protoplasts indicated a significant decrease in RGM compared to controls (from 0.99

to 0.80 and 0.74 for concentrations of 2 µM and 5 µM d18:0, respectively) (Fig. 11A-

D), with no statistical difference between the two concentrations of d18:0. The RGM

profile of a 4 µm fragment of PM from controls appeared homogenous whereas those

from sphingolipid-treated protoplasts is more heterogenous (Fig. 11E-G). These

findings suggest that the order level increased in response to the LCB. This increase

in membrane order level in the presence of d18:0 was confirmed by the greater

variation in the generalized polarization of Laurdan on biomimetic liposomes compared

to the control ethanol, and this as soon as d18:0 came into contact with the lipid bilayer

(Fig. 11H). Laurdan is a fluorescent probe sensitive to membrane lateral packing, a

higher generalized polarization corresponding to a greater lipid packing (Parasassi *et*

al., 1991).

Young's modulus was measured by recording force-distance curves using AFM on leaf

protoplasts. In the presence of d18:0, Young's modulus decreased (Fig. 11I) indicating

a reduction in membrane elasticity and a lower resistance to membrane deformation.

Again, there is no concentration dependence.

Using the Big Monolayer simulation method, which is useful for studying the lateral

distribution of molecules based on their mutual interaction energies, we observed that

d18:0 preferentially interacted with the lipid domains initially formed by β-sitosterol and

GluCer (Fig. 11J left), resulting in their fragmentation as shown in Fig. 11J right.

Accordingly, fluorescence images of di-4-ANEPPDHQ-stained protoplasts (Fig. 11 A

to C and E to G) demonstrated an increase in the number of small ordered domains in

the presence of d18:0 compared to its absence.

DISCUSSION

The roles of sphingolipids during PCD in response to biotic stress are still not fully understood (Berkey *et al*., 2012; Huby *et al*., 2020). Previous studies showed that co-infiltration of *Pst AvrRpm1* with the LCB d18:0 or t18:0 on *Arabidopsis* leaves did not trigger an HR on the treated leaves, with no evidence of cell death, indicating the conservation of plant PM integrity and absence of cell lysis following the co-infiltration (Glenz *et al*., 2022; Magnin-Robert *et al*., 2015). In the present work, we show that the levels of all plant immunity markers in response to co-infiltration, including those related to the plasma membrane, are similar to the control plants, despite the fact that the bacterium still develops inside plant tissues. We also report a similar effect with *Pst* expressing the effector AvrB or AvrPphB. AvrRpm1, AvrB and AvrPphB need to be myristoylated to be active. This process is disturbed during co-infiltration leading to an absence of plant defense response activation. Finally, interaction of d18:0 with lipids of the plant PM was demonstrated.

In vitro tests showed that this LCB does not have an antibacterial effect on *Pst AvrRpm1*, *Pst AvrB* and *Pst AvrPphB*, similarly to results obtained with *P. aeruginosa* (Fischer *et al*., 2012). t18:0 has been shown to inhibit *Pst* growth (Glenz *et al*., 2022). However, the authors used a smaller bacterial concentration and LB medium to perform their tests. Here, by using a higher bacterial concentration and a medium more favorable to *Pst* growth, the bacterial pressure was more important in our tests, which could explain the discrepancy with the data published by Glenz and colleagues (2022). *In planta* experiments showed that, even though *Pst AvrRpm1* is not capable of inducing an HR when co-infiltrated with d18:0, it is still present inside the plant and even continues to thrive, confirming the non-detrimental effect of d18:0 on the bacterium. In contrast, a pre-treatment with FB1, in order to promote endogenous LCB

accumulation, inhibit *Pst* DC3000 but not *Pst AvrRpm1* growth (Saucedo-García *et al*., 2023). Moreover, *loh1* plants that accumulate endogenous levels of LCBs together with *PR1* enhanced expression, displays spontaneous cell death (Ternes *et al*., 2011). The localization (apoplastic *vs* cytoplasmic) of the LCB and the fact that FB1 or *loh1* induces or displays higher LCB but also LCB-P, ceramide or GluCer accumulation (Chen *et al*., 2008; Kimberlin *et al*., 2016; Saucedo-García *et al*., 2011) could explain this discrepancy. This could also suggest that interaction of d18:0 with the PM and its presence within the PM when infiltrated could be essential in the cell death induction. Previous results showed that co-treatment of d18:0 with *Pst* DC3000 or with *Pst AvrRpm1* drastically reduced extracellular ROS production by *Arabidopsis* (Magnin-Robert *et al*., 2015). Our results demonstrate that d18:0 reduces ROS production in response to bacteria that induce symptoms *in planta* but also to bacteria that do not induce symptoms (*Pst AvrRps4* and *Pst AvrRpm1, Pst AvrB, Pst AvrPphB,* respectively). This production remained unchanged when plants are treated with *Pst AvrRpt2.* The hypothesis that LCBs could impact ROS production by plants has already been put forward in several studies, although some of them are somewhat contradictory. Coursol *et al*. (2015) showed that exogenous addition of 20 µM LCBs reduced tobacco ROS production in response to cryptogein. In *Arabidopsis*, LCBs can induce both intracellular as well as extracellular ROS production (Glenz *et al*., 2019; Peer *et al*., 2011; Shi *et al*., 2007; Saucedo-Garcia *et al*., 2023). ROS production can be detected during both PTI and ETI (Torres, 2010). In *Arabidopsis,* it is mainly related to RbohD located in lipid microdomains of the plant PM (Hao *et al*., 2014; Mongrand *et al*., 2004). Several studies have suggested that RbohD-derived ROS production is not directly involved in plant cell death, but rather in the signaling pathways associated with plant defense responses (Lherminier *et al*., 2009; Torres *et al*., 2005). This means

that even though d18:0 could alter ROS production, it may not have any impact on

PCD and would therefore not be responsible for the absence of HR. Additionally, it has

been already demonstrated that a lipidic elicitor is able to interact with lipid membrane

microdomains (Rossard *et al*., 2010). These data suggest that LCBs might interact with

and thus disturb the PM organization of *Arabidopsis* leading to the reduction of RbohD

activity.

SA pathway is triggered in response to hemibiotrophic pathogens, such as *Pst*,

whereas JA and its active form JA-Ile are involved in infection by necrotrophic

pathogens (Li *et al*., 2019). In our study, *PR1* is significantly down-regulated upon co-

infiltration with d18:0 and *Pst* DC3000, and even more so upon *Pst AvrRpm1,* which

coincides with lower SA accumulation. SA repression is commonly associated with

promotion of bacterial development and/or enhancement of a JA-dependent pathway

(Brooks *et al*., 2005; Moore *et al*., 1989; Zhou *et al*., 1998). *P. syringae* produces

coronatine, a phytotoxin that mimics JA-Ile and acts by inhibiting the accumulation of

SA in order to promote bacterial growth (Zheng *et al*., 2012). Conversely, in

Arabidopsis, coronatine-insensitive mutants that are resistant to *Pst* display higher

levels of SA (Block *et al*., 2005). In *Arabidopsis*, the SA and JA pathways antagonize

each other (Li *et al*., 2019; Mur *et al*., 2006). However, JA pathway markers are also

down-regulated in response to d18:0/*Pst AvrRpm1* infiltration. Considering these

results, we could conceive that co-infiltration represses or does not activate both SA

and JA pathways in response to the avirulent bacterium because of a lowered bacterial

virulence, a decreased susceptibility, and/or an altered perception of the pathogen.

AvrRpm1, AvrB, and AvrPphB effectors all possess a consensus N-terminal

myristoylation sites which are required for full avirulence function (Nimchuk *et al*.,

2000). Myristoylation process occurs inside the plant cell (Nimchuk *et al*., 2000) and is

catalyzed by NMT (*N*-myristoyltransferase) enzymes (Pierre *et al*., 2007). Here, *NMT1* expression is significantly reduced at 24 and 48 hpi when *Pst* AvrRpm1 is co-infiltrated with d18:0. d18:0 co-infiltration could thus interfere with the myristoylation of these effectors. As a consequence, it prevents them from being redirected to the PM where their targets are and consequently preventing their activity in *Arabidopsis*. Recently, a mis-localization of RPM1 at the PM, due to a depletion in anionic phospholipids in the PM, inhibits its cell death activity (Saile *et al*., 2021). Since cell death is still induced, such depletion does not affect RPS5 function, which is constitutively PM localized due to the presence of myristoylation motifs in its sequence. RPS5 mediates the recognition of AvrPphB. Co-infiltration of this effector with d18:0 results in an absence of HR, thus excluding the involvement of a PM phospholipid dependent mechanism. PM localization and stability of NLRs is essential for their activity in the induction of cell death and immune responses (Saile *et al*., 2021). Effector localization into plant cell is also essential for their interaction with their targets and subsequent activity (Nimchuk *et al*., 2000). The inability for AvrRPM1 to interact with its target implies not only that all the downstream plant defense responses are not activated but also the absence of RPM1 degradation after co-infiltration. Moreover, mutation of consensus myristoylation sites of *AvrRPM1* or *AvrB* leads to a highly reduction of HR events and an unhindered pathogen growth (Nimchuk *et al*., 2000), consistent with our results.

All the effects of d18:0 previously mentioned, and its amphiphilic nature suggest that the molecule can interact with the plant PM. It was thus hypothesized that it could insert into the lipid fraction of the PM, as already suggested for other molecules such as fatty acid hydroperoxides, rhamnolipids, lipopeptides that interact with lipids of plant PMs (Cordelier *et al*., 2022; Deboever *et al*., 2022; Deleu *et al*., 2013, 2008; Henry *et al*., 2011; Nasir *et al*., 2010; Schellenberger *et al*., 2021). Here, biophysical studies have

shown that d18:0 was attracted by, interacted with and stayed in biomimetic plant PMs.

These findings were further supported by *in silico* and *in vitro* analyses as well as

microscopic observations that notably showed d18:0 could disturb the lateral

organization and mechanical properties of plant PM and more specifically of its

microdomains. While it was suggested that exogenous LCB could cross the PM by

passive diffusion (Glenz *et al.*, 2019), our results suggested that d18:0 has no

propensity to go into the intracellular medium. It actually possesses a flip-flop potential

within the PM meaning that exogenous addition of d18:0 could disturb the PM. As

previously stated, this interaction could eventually disturb defense-related proteins

located within its domains (Gronnier *et al.*, 2018, 2016; Huby *et al.*, 2020; Mamode

Cassim *et al.*, 2019; Nagano *et al.*, 2016), which could explain the absence of

extracellular ROS production observed in response to *Pst*. Lipid composition and

organization of the PM influence PM localization and stability of NLRs and

consequently their activity in cell death inducing (Saile *et al.*, 2021). By disturbing the

PM organization, exogenous addition of d18:0 could also influence NLR localization

and function.

Collectively, our results demonstrated that d18:0 could have a dual effect on the PM

when co-infiltrated with *Pst AvrRpm1* by disturbing the organization and mechanical

properties of the PM and the myristoylation process, and thus the redirection of

effectors to the PM. Both actions prevent the interactions of NLR with its effector and

subsequent defense responses.

MATERIAL AND METHODS

Plant material and growth conditions

Seeds of the *Arabidopsis* T-DNA insertion mutants *rpm1-3* (CS68739), *rpm1-rps2-rin4*

(CS68760), and transgenic line *RPM1-myc rpm1* (CS68778) were obtained from the

Nottingham *Arabidopsis* Stock Center (http://Arabidopsis.info). Mutants, transgenic lines and

wild-type seeds Columbia-0 (Col-0) were grown in soil under 12 h-light/12 h-dark conditions

(150 µmol/m^2/s, 20°C, and 60% humidity) for 5 weeks.

Chemicals

Sphinganine (d18:0), palmitoyl-2-linoleoyl-*sn*-glycero-3-phosphocholine (PLPC), sitosterol

(sito), *D*-glucosyl-ß-1,1′-*N*-palmitoyl-*D*-erythro-sphingosine (d18:1/16:0 - GluCer or

Glucosylceramide) were purchased from Avanti Polar Lipids. Sphingolipids were dissolved in

ethanol (100%); PLPC, sitosterol and GluCer were dissolved in chloroform/methanol (2:1, v:v),

and used without further purification. Protamine sulphate was purchased from Sigma-Aldrich

(France).

Bacterial cultures and inoculations

Pst DC3000 strains transformed to express the effectors AvrRpm1 (*Pst AvrRpm1*), AvrRpt2

(*Pst AvrRpt2*), AvrRps4 (*Pst AvrRps4*), AvrB (*Pst AvrB*) or AvrPphB (*Pst AvrPphB*) were

provided by Prof. Jeff Dangl (University of North Carolina, USA) and Dr. Farid El-Kasmi

(University of Tübingen, Germany), Prof. Brian Staskawicz (University of California, USA),

Prof. Jane Parker (Max-Planck Institute, Cologne, Germany) and Dr. Brad Day (Michigan State

university, USA), respectively. Bacteria were cultured overnight under agitation (180 rpm) at

28°C in liquid King's B medium, supplemented with rifampicin (50 µg/mL) and kanamycin (50

µg/mL). Bacterial cells were collected by centrifugation, washed twice, and resuspended in 10

mM MgCl$_2$. Bacterial solutions were infiltrated at 10^7 CFU/mL on the abaxial side of *Arabidopsis*

leaves using a 1 mL syringe without needle. Co-treatment of bacteria with 100 µM d18:0 was

performed likewise. Control inoculations were conducted with 10 mM MgCl$_2$. At least four

leaves per plant were treated for each condition, with a minimum of three different plants.

***In vitro* antibacterial assays**

Bacteria were cultured as previously described before being distributed in 96 wells plates at

10^7 CFU/mL in King's B medium supplemented with appropriate antibiotics, under agitation

and at 28°C. Bacteria were plated either alone, with ethanol (0.1%, v/v) or with d18:0 (100 µM).

OD_{600} measurements were performed using the TECAN Spark® (USA) absorbance reader,

every hour for 50 h. The presented data are obtained from growth curve experiments, from

which the growth rate was extracted.

Pathogen assay *in planta*

For bacterial count experiments, bacterial cells were cultured and inoculated as described.

Four inoculated leaves were weighed and ground in 1 mL of 10 mM $MgCl_2$ using a mortar and

pestle. Appropriate dilutions were plated on King's B medium with suitable antibiotics, and

bacterial colonies were counted. Data are reported as means ±SD of the log (CFU/mg fresh

weight) of three replicates.

RNA extraction and qRT-PCR

Isolation of total RNA and real-time PCR were performed as described by Magnin-Robert et

al. (2015). Gene-specific primers are available in the Supplemental Table S1. For each

experiment, PCR was performed in triplicates. Transcript levels were normalized against those

of the UBIQUITIN5 (*UBQ5*) gene, used as an internal control. Fold induction compared with

untreated sample was calculated using the $2^{-\Delta\Delta Ct}$ method, where $\Delta\Delta Ct=$

$(Ct_{GI[\text{unknown sample}]}-Ct_{GI[\text{reference sample}]})-(Ct_{UBQ5[\text{unknown sample}]}-Ct_{UBQ5[\text{reference sample}]})$ where GI is the gene

of interest.

Immunoblotting assay

Total proteins from pre-treated leaves of *RPM1-myc rpm1* plants collected 0, 2, 6 and 24 hpi

were extracted and western-blotted according to Luzuriaga-Loaiza *et al*. (2018). Membrane

hybridization was carried out through overnight incubation with recombinant monoclonal Anti-

c-myc epitope tag [Clone 9E10] Mouse IgG1 kappa (Cliniscience, 1:1000) at 4°C, followed by

a 1 h incubation with anti-mouse IgG HRP-conjugated secondary antibodies (Bio-Rad, 1:3000)

at room temperature.

Electrolyte leakage

Measurement of electrolyte leakage were performed as described in Magnin-Robert *et al*.

(2015), except for leaf discs placed in each well of a 24-well plate containing 1 mL of distilled

water. Conductivity measurements were then conducted over time using a B-771 LaquaTwin

(Horiba) conductivity meter.

Extracellular ROS production

Measurements of extracellular ROS production were performed as described in Magnin-Robert

et al. (2015). Bacteria were added to a final concentration of 10^8 or 10^9 CFU/mL, as specified

further. In tests involving d18:0, the molecule was added at a final concentration of 100 μM

concomitantly with the elicitation solution.

Phytohormone and camalexin quantifications

The quantification of SA, JA-isoleucine (JA-Ile) and camalexin was performed at 0 and 48 hpi

by UPLC-nano-ESI-MS/MS as previously described in Herrfurth *et al*. (2020). Phytohormones

were extracted with methyl-tert-butyl ether (MTBE), reversed phase-separated using an

ACQUITY UPLC® system (Waters Corp., USA) and analysed by nanoelectrospray ionization

(nanoESI) (TriVersa Nanomate®; Advion BioSciences, USA) coupled with an AB Sciex 4000

QTRAP® tandem mass spectrometer (AB Sciex, USA) employed in scheduled multiple

reaction monitoring mode. For quantification, 10 ng D_4-SA (C/D/N Isotopes Inc., Canada),

20 ng D_5-IAA (Eurisotop, Germany), and 10 ng D_3-JA-Leu (kindly provided by Otto Miersch,

Halle/Saale, Germany) were added at the beginning of the extraction procedure. For detection,

the following mass transitions were used: 137/93 (declustering potential (DP) −25 V, entrance

potential (EP) -6 V, collision energy (CE) −20 V) for SA, 141/97 (DP −25 V, EP −6 V, CE −22

V) for D_4-SA, 179/135 (DP −35 V, EP −9 V, CE -14 V) for D_5-indole-acetic acid, 199/141 (DP

−60 V, EP −4 V, CE -32 V) for camalexin, 322/130 (DP −45 V, EP −5 V, CE -28 V) for JA-Ile

and 325/133 (DP −65 V, EP −4 V, CE -30 V) for D_3-JA-Leu.

GUS reporter assays

To determine induction of *PR1*, GUS enzyme activity of *PR1::GUS Arabidopsis* plants was

determined histochemically on leaves of the reporter line, treated as previously described,

collected at 48 hpi, and placed in the GUS staining solution (50 mM sodium phosphate buffer

- pH 7, 10mM EDTA, 1 mM $K_3Fe(CN)6$, 2 mM X-GlcA, and 0,1% Triton X-100). The leaves

were vacuum-infiltrated for 5 min and incubated for 4 h at 37°C. Next, they were washed with

Milli-Q water and left overnight at 4°C in a discoloration / fixation solution (ethanol:acetic acid,

3:1). Once the discoloration process was complete, leaves were placed in 95% (v/v) ethanol.

Lipid composition for plant biomimetic membranes

To mimic the lipid composition of plant PM, PLPC, β-sitosterol and GluCer were used, either

in a mixture (60:20:20, molar ratio) or alone, depending on the experiment. This lipid

composition was previously used to mimic plant PM (Deleu *et al.*, 2014, 2019; Deboever *et al.*,

2022).

Adsorption of sphinganine onto lipid monolayers using Langmuir trough

Adsorption experiments at constant surface area were performed in a KSV (Helsinki, Finland)

Minitrough (190 cm^3) equipped with a Wilhelmy plate, as previously described (Nasir *et al.*,

2016; Deleu *et al.*, 2019). The maximal insertion pressure (MIP) and the differential Π_0 ($d\Pi_0$)

were determined as previously described (Deboever *et al.*, 2020).

Preparation of large unilamellar vesicles

PLPC/ β-sitosterol/GluCer (60:20:20 molar ratio) large unilamellar vesicles (LUVs) were

prepared as described previously (Lebecque et al., 2019). Tris-HCl buffer (10 mM, pH 7.4) was

used for lipid film hydration and dispersion.

Extrusion was done *via* 13 passages through a 0.1 µM membrane filter using Liposofast

extruder (AVESTIN®). Diameter and stability of the LUVs were confirmed by dynamic light

scattering (DLS) measurements.

Thermodynamic parameters of sphinganine interaction with lipid bilayers by isothermal

titration calorimetry

For isothermal titration calorimetry (ITC) measurements, LUVs were prepared as described

and resuspended to reach a final concentration of 1 mM in Tris-HCl buffer (10 mM, pH 7.4).

Experiments were performed on a A VP-ITC microcalorimeter (Microcal Inc., Northampton,

MA, USA) as previously described (Lebecque *et al.*, 2018). Details can be found in supporting

information section (Supplemental Material S1).

Laurdan polarization on lipid vesicles LUVs

LUVs suspension at a concentration of 100 µM in lipids was incubated with 1 µM of Laurdan

(Sigma-Aldrich) for 1h15. Then, wells of black 96-well microplates (Greiner Bio-

One™ CellStar™, Fischer Scientific) were loaded with 100 µL of liposome solution per well.

The fluorescence was recorded using Spark® microplate reader (Tecan) with an excitation

wavelength at 360 (±35) nm and emission wavelengths at 430 (±20) nm and 485 (±20) nm.

The fluorescence was recorded once before treatment and every 5 min for up to 20 minutes,

after the addition of 100 µL of 2-times concentrated treatment.

The generalized polarization (GP) was defined as GP = $\frac{(I_{430nm}-I_{485nm})}{(I_{430nm}+I_{485nm})}$, where I_{430nm} and I_{485nm}

represents the blank-subtracted fluorescence intensities at emission wavelengths of 430 nm

and 485 nm respectively. Variation of GP (ΔGP) is defined as the subtraction of GP measured

at each time point following treatment from GP measured before treatment.

Propensity of sphinganine to insert into a bilayer determined by the Impala procedure

The most energetically favorable position and orientation of d18:0 in an implicite lipid bilayer was determined with the same procedure described in Ducarme *et al*. (1998) and Franche *et al*. (2020). For more details, see Supplementary Material S1.

Interaction energies between sphinganine and lipids determined by the Hypermatrix docking method

The hypermatrix docking method was employed, as previously described (Deleu *et al*., 2014; Deleu *et al*., 2019) to dock a molecule of sphinganine to a lipid system composed of either plant or bacterial lipids.

Modeling of sphinganine/plant lipids monolayer interactions with Big Monolayer method

The big monolayer method, as described by Deleu *et al.* (2019), was used to visualize lipid domains. Three repetitions of the system were calculated. The simulations were performed using plant lipid mixture with or without d18:0 (88% lipids, 12% d18:0).

Protoplast isolation

The isolation of mesophyll protoplasts was carried out by an enzymatic method as previously described (Yoo *et al*., 2007). The protoplasts were resuspended at 10^6 protoplasts/mL in a storage solution (20 mM MES, 400 mM mannitol, 20 mM KCl, 10 mM $CaCl_2$) until further use.

Atomic force microscopy

Protoplasts were adhered to the bottom of a glass-bottom Petrie dish by coating with positive-charge-bearing protamine sulphate 1 % in water: the bottom of the Petri dish was covered with a thin layer of this solution during 2 min, rinsed 3 times with the protoplast storage solution and the remaining solution was discarded. Protoplast suspension was incubated for 5 min with final

concentration 2 µM or 5 µM of d18:0 or control ethanol in protoplasts storage solution for 5

min. Then, pre-treated or not protoplasts suspension was gently deposited to the coating Petri

dish, and the protoplasts were settled for 5 min.

QNM-LC-CAL probes (Bruker, Billerica, USA), having a resonance frequency of 45 kHz, a tip

radius of 65 nm, a tip height of 16 µm and a nominal spring constant below 100 mN/m were

calibrated on a non-compliant sample to extract the deflection sensitivity which turned out to

be around 10 nm/V. Afterwards the tip was pulled off (of 100 Um) and tuned in liquid to

determine the spring constant.

Optics (Nikon Ti 2000, Nikon, Tokyo, Japan) were used in bright field mode to browse through

all protoplasts and focus on those having a 100 µm diameter. Therefore, the tip was engaged

at the very center of each and forces curves were captured in single force spectroscopy mode.

The atomic force microscopy (AFM) used was a Bioscope Catalyst (Bruker, Billerica, USA). A

z ramp output was used, with a ramp size of 5 µm, a ramp rate of 0.25 Hz, a contact time of

50 msec, and a forward and backward velocities of 5 µm/sec. Each force curves contained

2048 points. The Poisson's ratio was arbitrary set at 0.3. We used a relative trigger mode, a

tip threshold of 30 nm and plotted the deflection error as data type. It was verified that there

was no apparent change in diameter or x,y motion of the object during the acquisition.

The Young's modulus was extracted according to the Hert's theory and normalized as a

function of the protoplasts' volumes. A minimum of 50 curves per condition were recorded and

experiments were repeated three times.

Confocal microscopy

To determine the membrane order, protoplasts were labelled with 300 nM final concentration

of fluorescent probe di-4-ANEPPDHQ (Invitrogen, stock solution 1.5 mM in DMSO) in

protoplast storage solution, during 5 min. The labelled protoplasts suspension was gently

deposited to the bottom of a glass bottom Petri dish and the protoplasts were settled for 5 min.

The labelled protoplast suspension was incubated for 5 min with final concentration 2 µM or 5

µM of d18:0 or control ethanol for 5 min.

Fluorescence images were acquired using a laser scanning microscope LSM 880 from ZEISS

(Carl ZEISS SAS, Rueil Malmaison, France) through 63x (ON 1.4) oil immersion objective with

ZEN software. Di-4-ANEPPDHQ excitation was done at 488 nm and emission intensities were

acquired between 540–560 nm (green image) and between 630–650 nm (red image).

Ratiometric imaging was performed using the Fiji software (Schindelin *et al*., 2021).

ACKNOWLEDGEMENT

The authors thank Prof. Jeff Dangl (University of North Carolina, USA), Dr. Farid El-Kasmi
(University of Tübingen, Germany), Prof. Brian Staskawicz (University of California, USA),
Prof. Jane Parker (Max-Planck Institute, Cologne, Germany) and Dr. Brad Day (Michigan State
university, USA) for generously providing the different bacterial strains used in this study.
This work was supported by the National Fund for Scientific Research (FNRS, Belgium), the
SFR Condorcet (18ARC107), and the "Fondation Universitaire de Belgique" (Belgian
University Foundation). Some of the experiments were carried out within the Nanomat platform
(www.nanomat.eu) supported by the Ministère de l'Enseignement Supérieur et de la
Recherche, the Région Grand Est, and FEDER funds from the European Community. GG was
supported by the Foundation for Training in Industrial and Agricultural Research (FRIA, FNRS,
grant 1.E.069.20F), and MD is Senior Research Associates of the FNRS. I.F. acknowledges
funding by the Deutsche Forschungsgemeinschaft (INST 186/822-1).

AUTHOR CONTRIBUTIONS

741 SDC and MD designed the research. EH, SV, AB, CT, CCh, CH and GG performed

742 the research. EH, MD, SDC, FB analyzed and discussed the data. SC, SD, JC, CJ, FF,

743 CCl, IF provided critical feedback on the manuscript. EH, MD and SDC wrote the paper

744 and incorporated the input of the rest of the authors.

SUPPLEMENTAL DATA

Supplemental Figure S1. Adsorption behaviour of d18:0 into lipid

monolayer mimicking plant plasma membrane

Supplemental Table S1. Gene-specific primers used in real-time reverse transcription

PCR.

Supplementary Material S1. Methods in biophysics.

Conflict of interest statement

The authors declare no conflict of interest.

Orcid

Sylvain Cordelier: https://orcid.org/0000-0001-7723-4145
Jérôme Crouzet: https://orcid.org/0000-0003-0458-830X
Magali Deleu: https://orcid.org/0000-0001-8255-2965
Sandrine Dhondt-Cordelier: https://orcid.org/0000-0002-3778-6727
Stéphan Dorey: https://orcid.org/0000-0002-9583-2824
Ivo Feussner: https://orcid.org/0000-0002-9888-7003
Florence Fontaine: https://orcid.org/ 0000-0003-2918-8812
Guillaume Gilliard: https://orcid.org/0000-0002-4651-2920
Cornelia Herrfurth: https://orcid.org/0000-0001-8255-3255
Cédric Jacquard: https//orcid.org/0000-0003-4284-6082

References

Ade, J., DeYoung, B.J., Golstein, C., Innes, R.W., 2007. Indirect activation of a plant nucleotide binding
site-leucine-rich repeat protein by a bacterial protease. Proceedings of the National Academy of
Sciences U.S.A. 104, 2531–2536.
Ali, U., Li, H., Wang, X., Guo, L., 2018. Emerging roles of sphingolipid signaling in plant response to
biotic and abiotic stresses. Molecular Plant 11, 1328–1343.
Bari, R., Jones, J.D.G., 2009. Role of plant hormones in plant defence responses. Plant Molecular
Biology 69, 473–488.
Berkey, R., Bendigeri, D., Xiao, S., 2012. Sphingolipids and Plant Defense/Disease: The "Death"
Connection and Beyond. Frontiers in Plant Science 3.
Betsuyaku, S., Katou, S., Takebayashi, Y., Sakakibara, H., Nomura, N., Fukuda, H., 2018. Salicylic Acid
and Jasmonic Acid Pathways are Activated in Spatially Different Domains Around the Infection Site
During Effector-Triggered Immunity in *Arabidopsis thaliana*. Plant and Cell Physiology 59, 8–16.
Bjornson, M., Zipfel, C., 2021. Plant immunity: Crosstalk between plant immune receptors. Current
Biology 31, R796–R798.
Block, A., Schmelz, E., Jones, J.B., Klee, H.J., 2005. Coronatine and salicylic acid: the battle between
Arabidopsis and *Pseudomonas* for phytohormone control. Molecular Plant Pathology 6, 79–83.
Boyes, D.C., Nam, J., Dangl, J.L., 1998. The *Arabidopsis thaliana* RPM1 disease resistance gene
product is a peripheral plasma membrane protein that is degraded coincident with the hypersensitive
response. Proceedings of the National Academy of Sciences U.S.A. 95, 15849–15854.
Boyle, P.C., Martin, G.B., 2015. Greasy tactics in the plant-pathogen molecular arms race. Journal of
Experimental Botany 66, 1607–1616.
Brasseur, R., Killian, J.A., De Kruijff, B., Ruysschaert, J.M., 1987. Conformational analysis of gramicidin-
gramicidin interactions at the air/water interface suggests that gramicidin aggregates into tube-like
structures similar as found in the gramicidin-induced hexagonal HII phase. Biochimica et Biophysica
Acta (BBA) - Biomembranes 903, 11–17.
Brooks, D.M., Bender, C.L., Kunkel, B.N., 2005. The *Pseudomonas syringae* phytotoxin coronatine
promotes virulence by overcoming salicylic acid-dependent defences in *Arabidopsis thaliana*.
Molecular Plant Pathology 6, 629–639.

Buell, C.R., Joardar, V., Lindeberg, M., Selengut, J., Paulsen, I.T., Gwinn, M.L., Dodson, R.J., Deboy, R.T., Durkin, A.S., Kolonay, J.F., Madupu, R., Daugherty, S., Brinkac, L., Beanan, M.J., Haft, D.H., Nelson, W.C., Davidsen, T., Zafar, N., Zhou, L., Liu, J., Yuan, Q., Khouri, H., Fedorova, N., Tran, B., Russell, D., Berry, K., Utterback, T., Van Aken, S.E., Feldblyum, T.V., D'Ascenzo, M., Deng, W.-L., Ramos, A.R., Alfano, J.R., Cartinhour, S., Chatterjee, A.K., Delaney, T.P., Lazarowitz, S.G., Martin, G.B., Schneider, D.J., Tang, X., Bender, C.L., White, O., Fraser, C.M., Collmer, A., 2003. The complete genome sequence of the *Arabidopsis* and tomato pathogen *Pseudomonas syringae* pv. *tomato* DC3000. Proceedings of the National Academy of Sciences U.S.A. 100, 10181–10186.

Cacas, J.-L., Buré, C., Grosjean, K., Gerbeau-Pissot, P., Lherminier, J., Rombouts, Y., Maes, E., Bossard, C., Gronnier, J., Furt, F., Fouillen, L., Germain, V., Bayer, E., Cluzet, S., Robert, F., Schmitter, J.-M., Deleu, M., Lins, L., Simon-Plas, F., Mongrand, S., 2016. Revisiting plant plasma membrane lipids in tobacco: A focus on sphingolipids. Plant Physiology 170, 367–384.

Chen, M., Markham, J.E., Dietrich, C.R., Jaworski, J.G., Cahoon, E.B., 2008. Sphingolipid long-chain base hydroxylation is important for growth and regulation of sphingolipid content and composition in *Arabidopsis*. The Plant Cell 20, 1862–1878.

Contreras, F.-X., Sot, J., Alonso, A., Goñi, F.M., 2006. Sphingosine Increases the Permeability of Model and Cell Membranes. Biophysical Journal 90, 4085–4092.

Cordelier, S., Crouzet, J., Gilliard, G., Dorey, S., Deleu, M., Dhondt-Cordelier, S., 2022. Deciphering the role of plant plasma membrane lipids in response to invasion patterns: how could biology and biophysics help? Journal of Experimental Botany 73, 2765-2784.

Coursol, S., Fromentin, J., Noirot, E., Brière, C., Robert, F., Morel, J., Liang, Y.-K., Lherminier, J. and Simon-Plas, F., 2015. Long-chain bases and their phosphorylated derivatives differentially regulate cryptogein-induced production of reactive oxygen species in tobacco (*Nicotiana tabacum*) BY-2 cells. New Phytologist 205: 1239-1249.

Couto, D., Zipfel, C., 2016. Regulation of pattern recognition receptor signalling in plants. Nature Review Immunology 16, 537–552.

Debener, T., Lehnackers, H., Arnold, M., Dangl, J.L., 1991. Identification and molecular mapping of a single *Arabidopsis thaliana* locus determining resistance to a phytopathogenic *Pseudomonas syringae* isolate. The Plant Journal 1, 289–302.

Deboever, E., Lins L, Ongena M, De Clerck C, Deleu M, Fauconnier ML. 2020. Linolenic fatty acid hydroperoxide acts as biocide on plant pathogenic bacteria: Biophysical investigation of the mode of action. Bioorganic Chemistry 100, 10387.

Deboever, E., Van Aubel, G., Rondelli, V., Koutsioubas, A., Mathelie-Guinlet, M., Dufrene, Y.F., Ongena, M., Lins, L., Van Cutsem, P., Fauconnier, M., Deleu, M., 2022. Modulation of plant plasma membrane structure by exogenous fatty acid hydroperoxide is a potential perception mechanism for their eliciting activity. Plant Cell & Environment 45, 1082–1095.

Deleu, M., Crowet, J.-M., Nasir, M.N., Lins, L., 2014. Complementary biophysical tools to investigate lipid specificity in the interaction between bioactive molecules and the plasma membrane: A review. Biochimica et Biophysica Acta (BBA) - Biomembranes 1838, 3171–3190.

Deleu, M., Deboever, E., Nasir, M.N., Crowet, J.-M., Dauchez, M., Ongena, M., Jijakli, H., Fauconnier, M.-L., Lins, L., 2019. Linoleic and linolenic acid hydroperoxides interact differently with biomimetic plant membranes in a lipid specific manner. Colloids and Surfaces B: Biointerfaces 175, 384–391.

Deleu, M., Lorent, J., Lins, L., Brasseur, R., Braun, N., El Kirat, K., Nylander, T., Dufrêne, Y.F., Mingeot-Leclercq, M.-P., 2013. Effects of surfactin on membrane models displaying lipid phase separation. Biochimica et Biophysica Acta (BBA) - Biomembranes 1828, 801–815.

Deleu, M., Paquot, M., Nylander, T., 2008. Effect of Fengycin, a Lipopeptide Produced by *Bacillus subtilis*, on Model Biomembranes. Biophysical Journal 94, 2667–2679.

Desveaux, D., Singer, AU., Dangl JL., 2006. Type III effector proteins: doppelgangers of bacterial virulence. Current Opinion in Plant Biology 9, 376-382.

Dinic, J., Biverståhl, H., Mäler, L., Parmryd, I., 2011. Laurdan and di-4-ANEPPDHQ do not respond to membrane-inserted peptides and are good probes for lipid packing. Biochimica Biophysica Acta 1808, 298–306.

Dodds, P.N., Rafiqi, M., Gan, P.H.P., Hardham, A.R., Jones, D.A. and Ellis, J.G. (2009), Effectors of biotrophic fungi and oomycetes: pathogenicity factors and triggers of host resistance. New Phytologist, 183: 993-1000.

Ducarme, P., Rahman, M., Brasseur, R., 1998. IMPALA: A simple restraint field to simulate the biological membrane in molecular structure studies. PROTEINS: Structure, Function, and Genetics 30:357–371

Fischer, C.L., Drake, D.R., Dawson, D.V., Blanchette, D.R., Brogden, K.A., Wertz, P.W., 2012. Antibacterial activity of sphingoid bases and fatty acids against gram-positive and gram-negative bacteria. Antimicrobial Agents and Chemotherapy 56, 1157–1161.

Franche, A., Fayeulle, A., Lins, L., Billamboz, M., Pezron, I., Deleu, M., Léonard, E., 2020. Amphiphilic
azobenzenes: Antibacterial activities and biophysical investigation of their interaction with bacterial
membrane lipids. Bioorganic Chemistry 94, 103399.
Furlan, A.L., Laurin, Y., Botcazon, C., Rodríguez-Moraga, N., Rippa, S., Deleu, M., Lins, L., Sarazin, C.,
Buchoux, S., 2020. Contributions and limitations of biophysical approaches to study of the interactions
between amphiphilic molecules and the plant plasma membrane. Plants 9, 648.
Gassmann, W., Hinsch, M.E., Staskawicz, B.J., 1999. The *Arabidopsis* RPS4 bacterial-resistance gene
is a member of the TIR-NBS-LRR family of disease-resistance genes. The Plant Journal 20, 265–277.
Gerbeau-Pissot, P., Der, C., Thomas, D., Anca, I.-A., Grosjean, K., Roche, Y., Perrier-Cornet, J.-M.,
Mongrand, S., Simon-Plas, F., 2014. Modification of plasma membrane organization in tobacco cells
elicited by cryptogein. Plant Physiology 164, 273–286.
Glenz, R., Kaiping, A., Göpfert, D., Weber, H., Lambour, B., Sylvester, M., Fröschel, C., Mueller, M.J.,
Osman, M., Waller, F., 2022. The major plant sphingolipid long chain base phytosphingosine inhibits
growth of bacterial and fungal plant pathogens. Scientific Reports 12, 1081.
Glenz, R., Schmalhaus, D., Krischke, M., Mueller, M.J., Waller, F., 2019. Elevated levels of
phosphorylated sphingobases do not antagonize sphingobase- or fumonisin B1-induced plant cell
death. Plant and Cell Physiology 60, 1109–1119.
Grant, M., Godiard, L., Straube, E., Ashfield, T., Lewald, J., Sattler, A., Innes, R., Dangl, J., 1995.
Structure of the *Arabidopsis* RPM1 gene enabling dual specificity disease resistance. Science 269,
843–846.
Grennan, A.K., 2007. Lipid Rafts in Plants. Plant Physiology 143, 1083–1085.
Gronnier, J., Gerbeau-Pissot, P., Germain, V., Mongrand, S., Simon-Plas, F., 2018. Divide and rule:
Plant plasma membrane organization. Trends in Plant Science 23, 899–917.
Gronnier, J., Germain, V., Gouguet, P., Cacas, J.-L., Mongrand, S., 2016. GIPC: Glycosyl Inositol
Phospho Ceramides, the major sphingolipids on earth. Plant Signaling & Behavior 11, e1152438.
Guan, R., Su, J., Meng, X., Li, S., Liu, Y., Xu, J., Zhang, S., 2015. Multilayered regulation of ethylene
induction plays a positive role in *Arabidopsis* resistance against *Pseudomonas syringae*. Plant
Physiology 169, 299–312.
Gutiérrez-Nájera, N.A., Saucedo-García, M., Noyola-Martínez, L., Vázquez-Vázquez, C., Palacios-
Bahena, S., Carmona-Salazar, L., Plasencia, J., El-Hafidi, M., Gavilanes-Ruiz, M., 2020. Sphingolipid
effects on the plasma membrane produced by addition of fumonisin B1 to maize embryos. Plants 9,
150.
Hao, H., Fan, L., Chen, T., Li, R., Li, X., He, Q., Botella, M.A., Lin, J., 2014. Clathrin and membrane
microdomains cooperatively regulate RbohD dynamics and activity in *Arabidopsis*. Plant Cell 26,
1729–1745.
Henry, E., Yadeta, K.A., Coaker, G., 2013. Recognition of bacterial plant pathogens: local, systemic and
transgenerational immunity. New Phytologist 199, 908–915.
Henry, G., Deleu, M., Jourdan, E., Thonart, P., Ongena, M., 2011, The bacterial lipopeptide surfactin
targets the lipid fraction of the plant plasma membrane to trigger immune-related defence responses.
Cellular Microbiology 13, 1824-1837.
Herrfurth, C., Feussner, I., 2020. Quantitative jasmonate profiling using a high-throughput UPLC-
NanoESI-MS/MS method. In: Champion, A., Laplaze, L. (eds) Jasmonate in Plant Biology. Methods
in Molecular Biology, vol 2085. Humana, New York, NY.
Hicks, S.W., Galán, J.E., 2013. Exploitation of eukaryotic subcellular targeting mechanisms by bacterial
effectors. Nature Reviews Microbiology 11, 316–326.
Huby, E., Napier, J.A., Baillieul, F., Michaelson, L.V., Dhondt-Cordelier, S., 2020. Sphingolipids: towards
an integrated view of metabolism during the plant stress response. New Phytologist 225, 659–670.
Ishikawa, T., Aki, T., Yanagisawa, S., Uchimiya, H., Kawai-Yamada, M., 2015. Overexpression of BAX
INHIBITOR-1 links plasma membrane microdomain proteins to stress. Plant Physiology 169, 1333–
1343.
Jones, J.D.G., Dangl, J.L., 2006. The plant immune system. Nature 444, 323–329.
Kadota, Y., Liebrand, T.W.H., Goto, Y., Sklenar, J., Derbyshire, P., Menke, F.L.H., Torres, M., Molina,
A., Zipfel, C., Coaker, G., Shirasu, K., 2019. Quantitative phosphoproteomic analysis reveals common
regulatory mechanisms between effector- and PAMP-triggered immunity in plants. New Phytologist
221, 2160–2175.
Kamoun, S., 2006. A catalogue of the effector secretome of plant pathogenic oomycetes. Annual Review
of Phytopathology 44, 41-60.
Kato, H., Nemoto, K., Shimizu, M., Abe, A., Asai, S., Ishihama, N., Matsuoka, S., Daimon, T., Ojika, M.,
Kawakita, K., Onai, K., Shirasu, K., Yoshida, M., Ishiura, M., Takemoto, D., Takano, Y., Terauchi, R.,
2022. Recognition of pathogen-derived sphingolipids in Arabidopsis, Science 376, 857-860.

Kawasaki, T., Nam, J., Boyes, D.C., Holt, B.F., Hubert, D.A., Wiig, A., Dangl, J.L., 2005. A duplicated pair of *Arabidopsis* RING-finger E3 ligases contribute to the RPM1- and RPS2-mediated hypersensitive response: Transmembrane RING-E3 ligases and the HR. The Plant Journal 44, 258–270.

Kimberlin, A.N., Han, G., Chen, M., Cahoon, R.E., Luttgeharm, K.D., Stone, J.M., Markham, J.E., Dunn, T.M., Cahoon, E.B., 2016. ORM expression alters sphingolipid homeostasis and differentially affects ceramide synthase activities. Plant Physiology 172, 889-900.

Klemptner, R.L., Sherwood, J.S., Tugizimana, F., Dubery, I.A., Piater, L.A., 2014. Ergosterol, an orphan fungal microbe-associated molecular pattern (MAMP): Ergosterol as a MAMP. Molecular Plant Pathology 15, 747–761.

König, S., Feussner, K., Schwarz, M., Kaever, A., Iven, T., Landesfeind, M., Ternes, P., Karlovsky, P., Lipka, V., Feussner, I., 2012. *Arabidopsis* mutants of sphingolipid fatty acid α-hydroxylases accumulate ceramides and salicylates. New Phytologist 196, 1086–1097.

König, S., Gömann, J., Zienkiewicz, A., Zienkiewicz, K., Meldau, D., Herrfurth, C., Feussner, I., 2021. Sphingolipid-induced programmed cell death is a salicylic acid and EDS1-dependent phenotype in *Arabidopsis*. Biochemistry 63, 317-325.

Lebecque, S., Crowet, J.-M., Lins, L., Delory, B.M., du Jardin, P., Fauconnier, M.-L., Deleu, M., 2018. Interaction between the barley allelochemical compounds gramine and hordenine and artificial lipid bilayers mimicking the plant plasma membrane. Scientific Reports 8, 9784.

Lee, C.C., Wood, M.D., Ng, K., Andersen, C.B., Liu, Y., Luginbühl, P., Spraggon, G., Katagiri, F., 2004. Crystal structure of the type III effector AvrB from *Pseudomonas syringae*. Structure 12, 487–494.

Lee, D., Bourdais, G., Yu, G., Robatzek, S., Coaker, G., 2015. Phosphorylation of the plant immune regulator RPM1-INTERACTING PROTEIN4 enhances plant plasma membrane H + -ATPase activity and inhibits flagellin-triggered immune responses in *Arabidopsis*. Plant Cell 27, 2042–2056.

Lenarčič, T., Albert, I., Böhm, H., Hodnik, V., Pirc, K., Zavec, A.B., Podobnik, M., Pahovnik, D., Žagar, E., Pruitt, R., Greimel, P., Yamaji-Hasegawa, A., Kobayashi, T., Zienkiewicz, A., Gömann, J., Mortimer, J.C., Fang, L., Mamode-Cassim, A., Deleu, M., Lins, L., Oecking, C., Feussner, I., Mongrand, S., Anderluh, G., Nürnberger, T., 2017. Eudicot plant-specific sphingolipids determine host selectivity of microbial NLP cytolysins. Science 358, 1431–1434.

Lherminier, J., Elmayan, T., Fromentin, J., Elaraqui, K.T., Vesa, S., Morel, J., Verrier, J.-L., Cailleteau, B., Blein, J.-P., Simon-Plas, F., 2009. NADPH oxidase-mediated reactive oxygen species production: Subcellular localization and reassessment of its role in plant defense. Molecular Plant-Microbe Interactions 22, 868–881.

Li, N., Han, X., Feng, D., Yuan, D., Huang, L.-J., 2019. Signaling crosstalk between salicylic acid and ethylene/jasmonate in plant defense: Do we understand what they are whispering? International Journal of Molecular Sciences 20, 671.

Luttgeharm, K.D., Kimberlin, A.N., Cahoon, E.B., 2016. Plant sphingolipid metabolism and function, in: Nakamura, Y., Li-Beisson, Y. (Eds.), Lipids in Plant and Algae Development. Springer International Publishing, Cham, pp. 249–286.

Luzuriaga-Loaiza, W.P., Schellenberger, R., De Gaetano, Y., Obounou Akong, F., Villaume, S., Crouzet, J., Haudrechy, A., Baillieul, F., Clément, C., Lins, L., Allais, F., Ongena, M., Bouquillon, S., Deleu, M., Dorey, S., 2018. Synthetic rhamnolipid bolaforms trigger an innate immune response in *Arabidopsis thaliana*. Scientific Reports 8, 8534.

Mackey, D., Belkhadir, Y., Alonso, J.M., Ecker, J.R., Dangl, J.L., 2003. *Arabidopsis* RIN4 is a target of the type III virulence effector AvrRpt2 and modulates RPS2-mediated resistance. Cell 112, 379–389.

Mackey, D., Holt III, B.F., Wiig, A., Dangl, J.L., 2002. RIN4 interacts with *Pseudomonas syringae* type III effector molecules and is required for RPM1-mediated resistance in *Arabidopsis*. Cell 108, 743–754.

Magazzù, A., Marcuello, C., 2023. Investigation of soft matter nanomechanics by atomic force microscopy and optical tweezers: A comprehensive review. Nanomaterials 13, 963.

Magnin-Robert, M., Le Bourse, D., Markham, J.E., Dorey, S., Clément, C., Baillieul, F., Dhondt-Cordelier, S., 2015. Modifications of sphingolipid content affect tolerance to hemibiotrophic and necrotrophic pathogens by modulating plant defense responses in *Arabidopsis*. Plant Physiology 169, 2255-2274.

Mamode Cassim, A., Gouguet, P., Gronnier, J., Laurent, N., Germain, V., Grison, M., Boutté, Y., Gerbeau-Pissot, P., Simon-Plas, F., Mongrand, S., 2019. Plant lipids: Key players of plasma membrane organization and function. Progress in Lipid Research 73, 1–27.

Marsh, D., 1996. Lateral pressure in membranes. Biochimica et Biophysica Acta (BBA) - Reviews on Biomembranes 1286, 183–223.

Mongrand, S., Morel, J., Laroche, J., Claverol, S., Carde, J.-P., Hartmann, M.-A., Bonneu, M., Simon-Plas, F., Lessire, R., Bessoule, J.-J., 2004. Lipid rafts in higher plant cells: purification and characterization of triton x-100-insoluble microdomains from tobacco plasma membrane. Journal Biological Chemistry 279, 36277–36286.

Moore, R.A., Starratt, A.N., Ma, S.-W., Morris, V.L., Cuppels, D.A., 1989. Identification of a chromosomal region required for biosynthesis of the phytotoxin coronatine by *Pseudomonas syringae* pv. tomato. Canadian Journal of Microbiology 35, 910–917.

Mur, L.A.J., Kenton, P., Atzorn, R., Miersch, O., Wasternack, C., 2006. The outcomes of concentration-specific interactions between salicylate and jasmonate signaling include synergy, antagonism, and oxidative stress leading to cell death. Plant Physiology 140, 249–262.

Nagano, M., Ishikawa, T., Fujiwara, M., Fukao, Y., Kawano, Y., Kawai-Yamada, M., Shimamoto, K., 2016. Plasma membrane microdomains are essential for Rac1-RbohB/H-mediated immunity in rice. The Plant Cell 28, 1966–1983.

Nasir, M.N., Crowet, J.-M., Lins, L., Obounou Akong, F., Haudrechy, A., Bouquillon, S., Deleu, M., 2016. Interactions of sugar-based bolaamphiphiles with biomimetic systems of plasma membranes. Biochimie 130, 23–32.

Nasir, M.N., Thawani, A., Kouzayha, A., Besson, F., 2010. Interactions of the natural antimicrobial mycosubtilin with phospholipid membrane models. Colloids and Surfaces B: Biointerfaces 78, 17–23.

Ngou, B.P.M., Ahn, H.-K., Ding, P., Jones, J.D.G., 2021. Mutual potentiation of plant immunity by cell-surface and intracellular receptors. Nature 592, 110–115.

Nimchuk, Z., Marois, E., Kjemtrup, S., Leister, R.T., Katagiri, F., Dangl, J.L., 2000. Eukaryotic fatty acylation drives plasma membrane targeting and enhances function of several type III effector proteins from *Pseudomonas syringae*. Cell 101, 353–363.

Parasassi, T., De Stasio, G., Ravagnan, G., Rusch, R.M., Gratton, E., 1991. Quantitation of lipid phases in phospholipid vesicles by the generalized polarization of Laurdan fluorescence. Biophysical Journal 60, 179–189.

Pata, M.O., Hannun, Y.A., Ng, C.K.-Y., 2010. Plant sphingolipids: decoding the enigma of the Sphinx: Tansley review. New Phytologist 185, 611–630.

Peer, M., Bach, M., Mueller, M.J., Waller, F., 2011. Free sphingobases induce RBOHD-dependent reactive oxygen species production in *Arabidopsis* leaves. FEBS Letters 585, 3006–3010.

Pierre M., Traverso J.A., Boisson B., Domenichini S., Bouchez D., Giglione C., Meinnel T., 2007. *N*-myristoylation regulates the SnRK1 pathway in *Arabidopsis*, The Plant Cell 19, 2804–2821

Pottinger, S.E., Innes, R.W., 2020. RPS5-Mediated Disease Resistance: Fundamental Insights and Translational Applications. Annual Review of Phytopathology 58, 139-160.

Rondelli, V., Mollica, L., Koutsioubas, A., Nasir, N., Trapp, M., Deboever, E., Brocca, P., Deleu, M., 2022. Carbohydrate-carbohydrate interaction drives the preferential insertion of dirhamnolipid into glycosphingolipid enriched membranes. Journal of Colloid and Interface Science, 616, 739-748.

Rossard, S., Roblin, G., Atanassova, R., 2010. Ergosterol triggers characteristic elicitation steps in Beta vulgaris leaf tissues. Journal of Experimental Botany 61, 1807–1816.

Saijo, Y., Loo, E.P. iian, 2020. Plant immunity in signal integration between biotic and abiotic stress responses. New Phytologist 225, 87–104.

Saile, S.C., Ackermann, F.M., Sunil, S., Keicher, J., Bayless, A., Bonardi, V., Wan, L., Doumane, M., Stöbbe, E., Jaillais, Y., Caillaud, M.-C., Dangl, J.L., Nishimura, M.T., Oecking, C. and El Kasmi, F. 2021. Arabidopsis ADR1 helper NLR immune receptors localize and function at the plasma membrane in a phospholipid dependent manner. New Phytologist 232: 2440-2456.

Saucedo-García, M., González-Solís, A., Rodríguez-Mejía, P., Lozano-Rosas, G., Olivera-Flores, T. de J., Carmona-Salazar, L., Guevara-García, A.A., Cahoon, E.B., Gavilanes-Ruíz, M., 2023. Sphingolipid long-chain base signaling in compatible and non-compatible plant pathogen interactions in *Arabidopsis*. International Journal of Molecular Sciences 24, 4384.

Saucedo-García, M., Guevara-García, A., González-Solís, A., Cruz-García, F., Vázquez-Santana, S., Markham, J.E., Lozano-Rosas, M.G., Dietrich, C.R., Ramos-Vega, M., Cahoon, E.B., Gavilanes-Ruíz, M., 2011. MPK6, sphinganine and the LCB2a gene from serine palmitoyltransferase are required in the signaling pathway that mediates cell death induced by long chain bases in *Arabidopsis*. New Phytologist 191, 943–957.

Schellenberger, R., Crouzet, J., Nickzad, A., Shu, L.-J., Kutschera, A., Gerster, T., Borie, N., Dawid, C., Cloutier, M., Villaume, S., Dhondt-Cordelier, S., Hubert, J., Cordelier, S., Mazeyrat-Gourbeyre, F., Schmid, C., Ongena, M., Renault, J.-H., Haudrechy, A., Hofmann, T., Baillieul, F., Clément, C., Zipfel, C., Gauthier, C., Déziel, E., Ranf, S., Dorey, S., 2021. Bacterial rhamnolipids and their 3-hydroxyalkanoate precursors activate *Arabidopsis* innate immunity through two independent mechanisms. Proceedings of the National Academy of Sciences U.S.A. 118, e2101366118.

Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T., Preibisch, S.,
Rueden, C., Saalfeld, S., Schmid, B., Tinevez, J.-Y., White, D.J., Hartenstein, V., Eliceiri, K.,
Tomancak, P., Cardona, A., 2012. Fiji: an open-source platform for biological-image analysis. Nature
Methods 9, 676–682.
Shi, L., Bielawski, J., Mu, J., Dong, H., Teng, C., Zhang, J., Yang, X., Tomishige, N., Hanada, K.,
Hannun, Y.A., 2007. Involvement of sphingoid bases in mediating reactive oxygen intermediate
production and programmed cell death in *Arabidopsis*. Cell Research 17, 1030.
Ternes, P., Feussner, K., Werner, S., Lerche, J., Iven, T., Heilmann, I., Riezman, H., Feussner, I., 2011.
Disruption of the ceramide synthase LOH1 causes spontaneous cell death in *Arabidopsis thaliana*.
New Phytologist 192, 841–854.
Torres, M.A., 2010. ROS in biotic interactions. Physiologia Plantarum 138, 414–429.
Torres, M.A., Jones, J.D.G., Dangl, J.L., 2005. Pathogen-induced, NADPH oxidase–derived reactive
oxygen intermediates suppress spread of cell death in *Arabidopsis thaliana*. Nature Genetics 37,
1130–1134.
Warren, R.F., Henk, A., Mowery, P., Holub, E., Innes, R.W., 1998. A mutation within the leucine-rich
repeat domain of the *Arabidopsis* disease resistance gene RPS5 partially suppresses multiple
bacterial and downy mildew resistance genes. The Plant Cell 10, 1439–1452.
Wei, H.-L., Zhang, W., Collmer, A., 2018. Modular study of the type III effector repertoire in
Pseudomonas syringae pv. *tomato* DC3000 reveals a matrix of effector interplay in pathogenesis. Cell
Reports 23, 1630–1638.
Whalen, M.C., Innes, R.W., Bent, A.F., Staskawicz, B.J., 1991. Identification of *Pseudomonas syringae*
pathogens of *Arabidopsis* and a bacterial locus determining avirulence on both *Arabidopsis* and
soybean. The Plant Cell 3, 49–59.
Wirthmueller, L., Zhang, Y., Jones, J.D.G., Parker, J.E., 2007. Nuclear accumulation of the *Arabidopsis*
immune receptor RPS4 is necessary for triggering EDS1-dependent defense. Current Biology 17,
2023–2029.
Xin, X.-F., Kvitko, B., He, S.Y., 2018. *Pseudomonas syringae*: what it takes to be a pathogen. Nature
Reviews Microbiology 16, 316–328.
Yoo, SD., Cho, YH., Sheen, J., 2007. *Arabidopsis* mesophyll protoplasts: a versatile cell system for
transient gene expression analysis. Nature Protocols 2, 1565–1572.
Yuan, M., Jiang, Z., Bi, G., Nomura, K., Liu, M., Wang, Y., Cai, B., Zhou, J.-M., He, S.Y., Xin, X.-F.,
2021. Pattern-recognition receptors are required for NLR-mediated plant immunity. Nature 592, 105–
109.
Zheng, X., Spivey, N.W., Zeng, W., Liu, P.-P., Fu, Z.Q., Klessig, D.F., He, S.Y., Dong, X., 2012.
Coronatine promotes *Pseudomonas syringae* virulence in plants by activating a signaling cascade that
inhibits salicylic acid accumulation. Cell Host & Microbe 11, 587–596.
Zhou, N., Tootle, T.L., Tsui, F., Klessig, D.F., Glazebrook, J., 1998. PAD4 functions upstream from
salicylic acid to control defense responses in *Arabidopsis*. The Plant Cell 10, 1021–1030.
1074
1075

 FIGURE LEGENDS

 Figure 1. Effect of co-infiltration of *Pst* and d18:0 on cell death. (A) Representative

 leaves of Col-0 WT 48 hours after infiltration with either *Pst* DC3000, *Pst AvrRpm1*,

 Pst AvrRpt2, *Pst AvrRps4* alone or co-infiltrated with d18:0. The experiment was

 repeated at least 5 times with similar results (n=5). (B) Ion leakage measurements over

 24 hours following infiltration with *Pst AvrRpm1*, *Pst AvrRpt2* or *Pst AvrRps4*, alone or

 with d18:0. Data are reported as means ± SD (n=3). The experiment was repeated

 three times with similar results.

 Figure 2. *In vitro* and *in planta* bacterial growth in presence of sphinganine. (A)

 Growth rate was estimated from bacterial growth curve data (OD_{600}) over 48 hours,

 ±SD in King's B medium. Asterisks indicate significant differences between each

 bacterium co-infiltrated with d18:0 or not (Wilcoxon-Mann-Whitney, *, $p<0.05$). (B)

 Bacterial growth ±SD (n = 3) at 0 and 48 hpi in *Arabidopsis* WT Col-0 leaves infected

 by bacteria alone or with d18:0.

 Figure 3. Extracellular ROS production in response to bacterial infection.

 Extracellular ROS production measured in presence of *Pst AvrRpm1*, *Pst AvrRpt2 or*

 Pst AvrRps4 with or without d18:0 in the medium. Data represent the sum of RLU

 (Relative Light Units) ± SEM (n = 6) over 90 min. The experiment was performed three

 times with similar results. Asterisks indicate significant differences between each

 bacterium co-infiltrated with d18:0 or not (Wilcoxon-Mann-Whitney; *, $p<0.05$).

 Figure 4. Late plant defense events in response to bacterial infiltration. *Pst*

 DC3000 or *Pst AvrRpm1* were co-infiltrated with or without d18:0. (A) Camalexin was

quantified at 0 and 48 hpi (FW: fresh weight). Different letters indicate significant differences (ANOVA, p<0.05, one way, n=5). (B) Representative leaves of *PR1::GUS* lines 48 hpi. (C) and (D) Relative expression of *PR1* and *VSP1*, respectively, followed 6 and 48 hpi. Results are expressed as the fold increase in transcript level compared with the untreated control (0h), referred to as the x1 expression level. Experiments were repeated 3 times with similar results. (E) and (F) SA and JA-Isoleucine (JA-Ile) were quantified at 0 and 48 hpi. Different letters indicate significant differences (ANOVA, p<0.05, one way, n = 5).

Figure 5. Involvement of RPM1 in the absence of defense responses. (A) Photographs represent symptoms observed 72 hpi in WT, *rpm1-3*, *rpm1-rps2-rin4* and *RPM1-myc rpm1* lines treated with or without d18:0. (B) Western blot of RPM1 protein in WT and *RPM1-myc rpm1* line at 2, 6 or 24 hpi. Normalization was carried out with Actin. NT refers to non-treated samples. All experiments were repeated at least 3 times with similar results.

Figure 6. Effect of *Pst AvrB* and *Pst AvrPphB* co-infiltration with d18:0 on plant defense responses. (A) Photographs represent symptoms observed 48 hours after WT plant infiltration with either *Pst AvrB* or *Pst AvrPphB* alone or with d18:0. The experiment was repeated at least 3 times with similar results (n = 3). (B) Growth rate estimated from bacterial growth curve data (OD_{600}) over 48 hours, ±SD in King's B medium. (C) ROS production measurement in presence of *Pst AvrB* or *AvrPphB* with or without d18:0. Data represent the sum of RLU (Relative Light Units) ± SEM (n = 6) over 90 min. Experiments were performed three times with similar results. Asterisks indicate statistical differences (Wilcoxon-Mann-Whitney; *, p<0.05).

1126

Figure 7. *NMT1* expression in response to co-infiltration of bacteria and sphinganine. Relative expression of *NMT1*, (N-*myristoyltransferase 1*) was followed at 6 and 48 hpi of *Pst* DC3000 or *Pst AvrRpm1* with or without d18:0. Results are expressed as the fold increase in transcript level compared with the untreated control (0h), referred to as the x1 expression level. Experiments were repeated three times with similar results. Asterisks indicate significant differences between each bacterium co-infiltrated with d18:0 or not (Wilcoxon-Mann-Whitney; *, $p<0.05$).

Figure 8. Interaction of sphinganine with an implicit modeled bilayer. (A) The Impala simulation calculates the energy restraints of a lipid bilayer as d18:0 goes through it. The X-axis corresponds to the position of the center of mass along the Z-axis. From left to right: the interface between the bilayer and the aqueous phase (orange), the interface between the polar head and the alkyl chain (pink) and the center of the bilayer (green). (B) The most energetically favorable orientation of d18:0 in the modeled bilayer. The lines represent the same planar surfaces as in A.

Figure 9. Interaction of sphinganine with plant-like liposomes. (A) Raw data from ITC experiments at 26°C. Each peak corresponds to a single injection of a 1 mM PLPC, glucosylceramide and sitosterol (60:20:20, molar ratio) LUV suspension into a solution containing 20 µM of d18:0. (B) Calculated thermodynamic parameters for this interaction. These values are means of two independent experiments. K represents the binding coefficient (the affinity of d18:0 for the LUVs), ΔG represents the molar free energy, ΔH, the molar enthalpy change, and TΔS, the molar entropy change.

1150

Figure 10. Specificity of sphinganine interaction with various lipids representative of the plant plasma membrane. (A) Adsorption of d18:0 into monolayers of plant PM lipids: PLPC, glucosylceramide (GluCer) and β-sitosterol (Sito). d18:0 was injected at 1.25 µM in a subphase (Tris HCl, 10 mM, pH 7.4), underneath the lipid monolayer. The maximal insertion pressure (MIP) reflects the penetration power of d18:0 into the lipid monolayer and the differential Π (dΠ0) indicates the attracting effect the lipids have on it. (B) Total energies of interaction of d18:0 alone or with lipids representative of the plant plasma membrane were calculated with the Hypermatrix docking method. Lipids used were PLPC, GluCer, and Sito.

Figure 11. Mechanical properties and organization of the plant plasma membrane in response to sphinganine treatment. (A) to (C) Fluorescence red/green ratio images by multispectral confocal microscopy of a single-cell PM labeled with di-4-ANEPPDHQ after control (A), 2 µM d18:0 (B) or 5 µM d18:0 (C) treatment. (D) Red to green ratio of the PM (RGM) labeled with di-4-ANEPPDHQ treated with control (Ctrl), 2 µM or 5 µM d18:0. The asterisk indicates a significant difference compared to the control (P value < 0.05). (E) to (G) RGM profile on a 4 µm distance on PM from protoplasts after control (E), 2 µM d18:0 (F) or 5 µM d18:0 (G) treatment. (H) Variation of laurdan generalized polarization (ΔGP) measured in PM-mimicking liposomes (50 µM PLPC:Glucosylceramide: β-sitosterol (60:20:20 molar ratio)) at several time points after control (white bars) and 20 µM d18:0 treatment (grey bars). A positive ΔGP corresponds to a higher lipid packing within the bilayer. Mean ± SD of two replicates from four independent experiments. Asterisks indicate statistically significant differences with control, *P<0.05, **P<0.01, two-way ANOVA and Sidak's multiple comparison test. (I) Young's modulus measured on mesophyll protoplasts

before or after 5 min incubation of d18:0 at two concentrations. Mean ± SD of at least

50 analyses from three independent experiments. (J) Monolayer simulations of lipid-

d18:0 interactions performed with Big Monolayer where each pixel represents a

molecule. (Left) Plant lipid model with PLPC in green, glucosylceramide (GluCer) in

blue and sitosterol (sito) in orange (60:20:20, molar ratio). (Right) The same lipids after

addition of d18:0 in pink (88% lipids, 12% d18:0).

1182

FIGURE 1 Effect of co-infiltration of *Pst* and d18:0 on cell death.
(A) Representative leaves of Col-0 WT 48 hours after infiltration with either *Pst* DC3000, *Pst AvrRpm1*, *Pst AvrRpt2*, *Pst AvrRps4* alone or co-infiltrated with d18:0. The experiment was repeated at least 5 times with similar results (n=5). (B) Ion leakage measurements over 24 hours following infiltration with *Pst AvrRpm1*, *Pst AvrRpt2* or *Pst AvrRps4*, alone or with d18:0. Data are reported as means ± SD (n=3). The experiment was repeated three times with similar results.

FIGURE 2 *In vitro* and *in planta* **bacterial growth in presence of sphinganine.**
(A) Growth rate was estimated from bacterial growth curve data (OD_{600}) over 48 hours, ±SD in King's B medium. Asterisks indicate significant differences between each bacterium co-infiltrated with d18:0 or not (Wilcoxon-Mann-Whitney, *, $p<0.05$). (B) Bacterial growth ±SD (n = 3) at 0 and 48 hpi in *Arabidopsis* WT Col-0 leaves infected by bacteria alone or with d18:0.

FIGURE 3 Extracellular ROS production in response to bacterial infection.
Extracellular ROS production measured in presence of *Pst AvrRpm1*, *Pst AvrRpt2 or Pst AvrRps4* with or without d18:0 in the medium. Data represent the sum of RLU (Relative Light Units) ± SEM (n = 6) over 90 min. The experiment was performed three times with similar results. Asterisks indicate significant differences between each bacterium co-infiltrated with d18:0 or not (Wilcoxon-Mann-Whitney; *, p<0.05).

FIGURE 4 Late plant defense events in response to bacterial infiltration.
Pst DC3000 or *Pst AvrRpm1* were co-infiltrated with or without d18:0. (A) Camalexin
was quantified at 0 and 48 hpi (FW: fresh weight). Different letters indicate significant
differences (ANOVA, p<0.05, one way, n=5). (B) Representative leaves of *PR1::GUS*
lines 48 hpi. (C) and (D) Relative expression of *PR1* and *VSP1*, respectively, followed 6
and 48 hpi. Results are expressed as the fold increase in transcript level compared
with the untreated control (0h), referred to as the x1 expression level. Experiments
were repeated 3 times with similar results. (E) and (F) SA and JA-Isoleucine (JA-Ile)
were quantified at 0 and 48 hpi. Different letters indicate significant differences
(ANOVA, p<0.05, one way, n = 5).

FIGURE 5 Involvement of RPM1 in the absence of defense responses.
(A) Photographs represent symptoms observed 72 hpi in WT, *rpm1-3*, *rpm1-rps2-rin4* and *RPM1-myc rpm1* lines treated with or without d18:0. (B) Western blot of RPM1 protein in WT and *RPM1-myc rpm1* line at 2, 6 or 24 hpi. Normalization was carried out with Actin. NT refers to non-treated samples. All experiments were repeated at least 3 times with similar results.

FIGURE 6 Effect of *Pst AvrB* and *Pst AvrPphB* co-infiltration with d18:0 on plant defense responses.
(A) Photographs represent symptoms observed 48 hours after WT plant infiltration with either *Pst AvrB* or *Pst AvrPphB* alone or with d18:0. The experiment was repeated at least 3 times with similar results (n = 3). (B) Growth rate estimated from bacterial growth curve data (OD_{600}) over 48 hours, ±SD in King's B medium. (C) ROS production measurement in presence of *Pst AvrB* or *AvrPphB* with or without d18:0. Data represent the sum of RLU (Relative Light Units) ± SEM (n = 6) over 90 min. Experiments were performed three times with similar results. Asterisks indicate statistical differences (Wilcoxon-Mann-Whitney; *, $p < 0.05$).

FIGURE 7 *NMT1* **expression in response to co-infiltration of bacteria and sphinganine.**

Relative expression of *NMT1*, (N-*myristoyltransferase 1*) was followed at 6 and 48 hpi of *Pst* DC3000 or *Pst AvrRpm1* with or without d18:0. Results are expressed as the fold increase in transcript level compared with the untreated control (0h), referred to as the x1 expression level. Experiments were repeated three times with similar results. Asterisks indicate significant differences between each bacterium co-infiltrated with d18:0 or not (Wilcoxon-Mann-Whitney; *, p<0.05).

Total constraint energy (Kcal/mol)

FIGURE 8 Interaction of sphinganine with an implicit modeled bilayer.
(A) The Impala simulation calculates the energy restraints of a lipid bilayer as d18:0 goes through it. The X-axis corresponds to the position of the center of mass along the Z-axis. From left to right: the interface between the bilayer and the aqueous phase (orange), the interface between the polar head and the alkyl chain (pink) and the center of the bilayer (green). (B) The most energetically favorable orientation of d18:0 in the modeled bilayer. The lines represent the same planar surfaces as in A.

B	Mean Values	Standard deviations
K (m/M)	86.80	18.99
ΔG (J/mol)	-16.09	0.55
ΔH (J/mol)	-1.39	0.26
TΔS (J/mol)	14.70	0.29

FIGURE 9 Interaction of sphinganine with plant-like liposomes. (A) Raw data from ITC experiments at 26°C. Each peak corresponds to a single injection of a 1 mM PLPC, glucosylceramide and sitosterol (60:20:20, molar ratio) LUV suspension into a solution containing 20 µM of d18:0. (B) Calculated thermodynamic parameters for this interaction. These values are means of two independent experiments. K represents the binding coefficient (the affinity of d18:0 for the LUVs), ΔG represents the molar free energy, ΔH, the molar enthalpy change, and TΔS, the molar entropy change.

A

B

FIGURE 10 Specificity of sphinganine interaction with various lipids representative of the plant plasma membrane.
(A) Adsorption of d18:0 into monolayers of plant PM lipids: PLPC, glucosylceramide (GluCer) and b-sitosterol (Sito). d18:0 was injected at 1.25 µM in a subphase (Tris HCl, 10 mM, pH 7.4), underneath the lipid monolayer. The maximal insertion pressure (MIP) reflects the penetration power of d18:0 into the lipid monolayer and the differential Π (dΠ0) indicates the attracting effect the lipids have on it. (B) Total energies of interaction of d18:0 alone or with lipids representative of the plant plasma membrane were calculated with the Hypermatrix docking method. Lipids used were PLPC, GluCer, and Sito.

Figure 11. Mechanical properties and organization of the plant plasma membrane in response to sphinganine treatment.
(A) to (C) Fluorescence red/green ratio images by multispectral confocal microscopy of a single-cell PM labeled with di-4-ANEPPDHQ after control (A), 2 μM d18:0 (B) or 5 μM d18:0 (C) treatment. (D) Red to green ratio of the PM (RGM) labeled with di-4-ANEPPDHQ treated with control (Ctrl), 2 μM or 5 μM d18:0. The asterisk indicates a significant difference compared to the control (P value < 0.05). (E) to (G) RGM profile on a 4 μm distance on PM from protoplasts after control (E), 2 μM d18:0 (F) or 5 μM d18:0 (G) treatment. (H) Variation of laurdan generalized polarization (ΔGP) measured in PM-mimicking liposomes (50 μM PLPC:Glucosylceramide:β-sitosterol (60:20:20 molar ratio)) at several time points after control (white bars) and 20 μM d18:0 treatment (grey bars). A positive ΔGP corresponds to a higher lipid packing within the bilayer. Mean ± SD of two replicates from four independent experiments. Asterisks indicate statistically significant differences with control, *P<0.05, **P<0.01, two-way ANOVA and Sidak's multiple comparison test. (I) Young's modulus measured on mesophyll protoplasts before or after 5 min incubation of d18:0 at two concentrations. Mean ± SD of at least 50 analyses from three independent experiments. (J) Monolayer simulations of lipid-d18:0 interactions performed with Big Monolayer where each pixel represents a molecule. (Left) Plant lipid model with PLPC in green, glucosylceramide (GluCer) in blue and sitosterol (sito) in orange (60:20:20, molar ratio). (Right) The same lipids after addition of d18:0 in pink (88% lipids, 12% d18:0).