Cortistatin exerts an immunomodulatory and neuroprotective role in a

preclinical model of ischemic stroke

Castillo-González J[1], Buscemi L[2,3], Vargas-Rodríguez P[1], Serrano-Martínez I[1], Forte-

Lago I[1], Price M[2,3], Hernández-Cortés P[4], Hirt L[2,3*], González-Rey E[1*].

[1]Institute of Parasitology and Biomedicine Lopez-Neyra (IPBLN), CSIC, PT Salud,

18016 Granada, Spain; [2]University of Lausanne, Lausanne, Switzerland; [3]Lausanne

University Hospital, Lausanne, Switzerland; [4]Medicine School, University of Granada,

Granada, Spain

*Correspondence: elena.g@csic.es; lorenz.hirt@chuv.ch; Tel.: +34-958-181670

Abstract

Ischemic stroke is the result of a permanent or transient occlusion of a brain artery,

leading to irreversible tissue injury and long-term sequelae. Despite ongoing

advancements in revascularization techniques, stroke remains the second leading

cause of death worldwide. A comprehensive understanding of the complex and

interconnected mechanisms, along with the endogenous mediators that modulate

stroke responses is essential for the development of effective interventions. Our study

investigates cortistatin, a neuropeptide extensively distributed in the immune and

central nervous systems, known for its immunomodulatory properties. With

neuroinflammation and peripheral immune deregulation as key pathological features

of brain ischemia, cortistatin emerges as a promising therapeutic candidate. To this

aim, we evaluated its potential effect in a well-established middle cerebral artery

occlusion (MCAO) preclinical stroke model. Our findings indicate that the peripheral

administration of cortistatin at 24 hours post-stroke significantly reduces neurological

damage and enhances recovery. Importantly, cortistatin-induced neuroprotection was

multitarget, as it modulated the glial reactivity and astrocytic scar formation, facilitated blood-brain barrier recovery, and regulated local and systemic immune dysfunction. Surprisingly, administration of cortistatin at immediate and early post-stroke time points proved to be not beneficial and even detrimental. These results emphasize the importance of understanding the spatio-temporal dynamics of stroke pathology to develop innovative therapeutic strategies. Premature interruption of certain neuroinflammatory processes might inadvertently compromise neuroprotective mechanisms. In summary, our study highlights cortistatin as a novel pleiotropic therapeutic approach against ischemic stroke, offering new treatment options for patients for whom early revascularization intervention is unsuccessful.

Graphical abstract

Keywords: blood-brain barrier breakdown, cortistatin, ischemic stroke, glial response, neuroimmune regulation, neuroinflammation, neuroprotection.

1. <u>INTRODUCTION</u>

Ischemic stroke results from a permanent or transient occlusion of a cerebral artery.

The abrupt cessation of cerebral blood flow and subsequent reperfusion triggers a

rapid and complex cascade of molecular and cellular events that may evolve over

hours to days and weeks after the onset [1]. These pathophysiological events are

diverse and include energy failure, oxidative stress, excitotoxicity, cell death, microglia

reactivity, glial scar formation, blood-brain barrier (BBB) breakdown, and peripheral

immune responses [2]. Notably, these processes are intricately interlinked, often

unfolding concurrently within a brief time frame.

Despite significant efforts, stroke remains the second leading cause of death

globally. Current treatments are limited to intravenous recombinant tissue

plasminogen activator (rtPA) or endovascular thrombectomy. However, the narrow

therapeutic window and rather limited efficacy sometimes restrict the utility of these

treatments [3]. Furthermore, the intricate and still not fully understood spatiotemporal

pathophysiology of stroke, with a plethora of underlying cellular and molecular

processes involving the brain, vascular system, and peripheral organs, complicates its

management. Consequently, there is an urgent need to thoroughly understand the

neuroimmunological context and the factors involved at the different stages of

ischemic pathology to develop effective therapeutic approaches.

Cortistatin (CST) is a cyclic neuropeptide broadly distributed in the central

nervous system (CNS) and the immune system [4,5]. Despite sharing considerable

homology, functions, and receptors (Sst1-5) with somatostatin, cortistatin exhibits

unique functions in the nervous system and peripheral tissues [6,7]. These particular

functions are mediated through interactions with receptors distinct from those of

somatostatin, including the ghrelin receptor, the Mas gene-related receptor X-2, and

a yet unidentified cortistatin receptor [5]. Notably, cortistatin displays potent anti-

inflammatory properties, protecting against the development and progression of several inflammatory and autoimmune disorders (reviewed in [5]). Additionally, cortistatin treatment demonstrates immunomodulatory and neuroprotective effects in models of neuroinflammatory and neuroimmune disorders, such as multiple sclerosis [8], meningoencephalitis [9], excitotoxicity [10], neuropathic pain [11], and Parkinson's disease [12]. In these conditions, cortistatin consistently reduces pro-inflammatory mediators, modulates glial cell responses, and maintains tissue-surveillance activities. Importantly, the absence of cortistatin in mice is associated with an exacerbated pro-inflammatory basal state in the CNS and periphery, along with decreased expression of neurotrophic factors [5,8]. Recently, our group has demonstrated a key role for cortistatin in maintaining BBB integrity, mediating anti-inflammatory endothelial functions, and facilitating BBB repair by stabilizing tight-junction proteins (TJs) and mitigating exaggerated immune responses after ischemic-like *in vitro* conditions [13]. However, no data has been reported about the influence of cortistatin on *in vivo* ischemic damage.

Therefore, given its neuroprotective, anti-inflammatory, and immunomodulatory properties, as well as its ability to regulate the BBB, we hypothesize that cortistatin could be a promising therapeutic agent affecting the intricate glial, vascular and peripheral responses underlying stroke pathophysiology.

2. <u>MATERIALS AND METHODS</u>

2.1. *Peptides*

Mouse CST-29 was acquired from Bachem. The lyophilized peptide was dissolved in distilled water (final concentration 10^{-4} M).

2.2. *Animals*

The procedures described in this study were approved by the Animal Care and Use

Board and the Ethical Committee of the Spanish National Research Council (Animal

Care Unit Committee IPBLN-CSIC # protocol CEEA OCT/2017.EGR), and the Vaud

Cantonal Veterinary Office (protocol number 2017x6a) (Switzerland). All experimental

protocols adhered to the guidelines specified in Directive 2010/63/EU of the European

Parliament and the corresponding Swiss regulations regarding animal protection for

scientific purposes. Mice were housed in groups of five animals per cage under

controlled environmental conditions (60-70% relative humidity, 22 ± 1°C), and

maintained on a 12-hour light/dark cycle (lights on at 7:00 a.m.). Food and water were

provided *ad libitum*.

99   *2.3. Transient middle cerebral artery occlusion model*

Cerebral ischemia was induced in 12-week-old male C57BL/6J mice using the

transient middle cerebral artery occlusion (MCAO) model, as adapted from Buscemi

et al. [14]. Briefly, mice were anesthetized with isoflurane (1.5-2% in 70% N_2O/30%

O_2) using a facemask. For post-surgical analgesia buprenorphine (0.025 mg/kg) was

subcutaneously administered. A heating pad was used to maintain body temperature

throughout the surgery. Regional cerebral blood flow (CBF) was monitored using laser

Doppler flowmetry (moorVMS-LDF1, Moor Instruments) by fixing a probe on the skull

over the core area supplied by the middle cerebral artery (approximately 1mm

posterior and 6mm lateral from the bregma). Transient focal ischemia was induced by

introducing a silicon-coated filament (701756PK5Re, Doccol) through the left common

carotid artery into the internal carotid artery until it occluded the middle cerebral artery.

The filament was left in place for 20 minutes before being withdrawn to allow

reperfusion. Surgeries were considered successful when blood dropped below 20%

of the baseline during occlusion, and raised above 50% of the baseline after filament

withdrawal. Post-surgery, mice were housed with *ad libitum* access to food and water,

with each cage containing water-softened chow and recovery gel.

Cortistatin (144µg/kg) was administered intraperitoneally at 0h and 24h

(immediate treatment), 4h and 24h (early treatment), or 24h (late treatment) post-

reperfusion. Control mice received an equivalent volume of saline solution.

2.4. Neurological score and behavioural tests

Neurological assessments were conducted at 24h following MCAO and prior to

sacrifice at 48h. Stroke severity was assessed on a scale from 0 (no signs) to 3 (death)

determining motor, balance, reflex, and sensory functions using a compendium of

scales previously published [15,16].

To evaluate motor function and ataxia, the rotarod test was performed using a

rotating cylinder apparatus (Ugo Basile, Gemonio, Italy) with a gradual acceleration

protocol. The cylinder accelerated from 4rpm to 40rpm over 216s and then maintained

a constant speed of 40rpm until reaching a 300s time limit. To ensure reliable results

and exclude motivational factors, such as early falls, a training phase was conducted

[17]. Mice underwent three consecutive daily training trials (with a 10-minute gap

between them) for two days before MCAO (training trials, baseline). The longest

latency to fall from the rod in three trials was recorded for each mouse. Mice underwent

daily test trials at 24h and 48h after MCAO. For the test trials, the longest latency to

fall from the rod or to complete two consecutive full turns while gripping the rod within

the 300 seconds time limit was recorded for each mouse.

Additionally, muscle coordination and endurance were assessed using the wire-

hanging test. Animals were suspended on a custom-made apparatus consisting of a

single wire stretched between two posts above a soft ground. Mice were trained 24h

before surgery and scored based on their performance in scaping (touching and reaching either the right or left post) or falling at 24 hours and 48 hours after MCAO [15]. The time elapsed until they escaped or fell was recorded. The test concluded either after 180 seconds or when a mouse had fallen off the wire ten times.

2.5. Sacrifice and infarct volume measurement

Mice were euthanized by intraperitoneal injection overdose of ketamine (100mg/kg, Ritcher Pharma)/ xylazine (10mg/kg, Fatro Iberica) at 48h post-MCAO surgery, followed by intracardial perfusion with 4% paraformaldehyde (PFA). Brains were collected and post-fixed in 4% PFA for 24h at 4°C, and then cryoprotected in 30% sucrose for 48h at 4°C. Subsequently, brains were frozen in isopentane on dry ice at -25°C and stored at -80°C until further processing. Coronal sections of 20μm thickness were obtained using a cryostat.

Cresyl violet (CV)-stained sections were prepared by gradual hydration in ethanol, stained with 1mg/ml of CV-acetate in distilled water, gradual dehydration in ethanol, and clearing in xylene [15]. Sections were imaged under a Leica DM2000 Transmit stereomicroscope at 2.5x magnification. The infarct volume was determined by multiplying the sum of the infarcted areas on each section (measured using the ImageJ Fiji software) by the section spacing.

2.6. Immunofluorescence

For immunofluorescence labelling, 20μm free-floating sections were washed with PBS and blocked and permeabilized with PBS containing 1% BSA, 0.1% Triton X-100, and 5% goat serum for 60min at room temperature (RT). Following this, sections were incubated overnight (O/N) at 4°C with primary antibodies anti-MAP2 (1:1000, Sigma), anti-Iba1 (1:1000, Wako), anti-GFAP (1:5000, Dako), anti-CD31/PECAM-1 (1:300,

BD), anti-ZO-1 (1:100, Fisher), or APC-labelled anti-CD45 (1:200, BD) in PBS

containing 1% BSA, 0.1% Triton X-100, and 5% goat serum. After extensive washing

with PBS, sections were incubated with the corresponding secondary Alexa Fluor-

conjugated antibodies (1:1000, Sigma) and DAPI in PBS containing 1% BSA, 0.1%

Triton X-100, and 2% goat serum for 60min. Sections were mounted with Mowiol. For

IgG staining, an Alexa Fluor 594-conjugated goat anti-mouse IgG antibody was utilized

without a primary antibody. Images were captured using a LEICA DMi8 S Platform

Life-cell microscope at 20x magnification and a confocal Leica TCS SP8 STED

microscope at 63x magnification.

2.7.   Morphometric characterization

To comprehensively characterize morphological alterations in microglia and

astrocytes post-stroke, we employed different morphometric analyses. Images of

selected microglia and astrocytes within specific brain regions (*i.e.,* periinfarct or core

area) were randomly selected, ensuring no overlap between cells, and that the entire

nucleus and branches were included. A total of 15-30 cells per mouse (n=15-30) from

3-6 mice (N=3-6) were analysed. Image processing was conducted using ImageJ Fiji

software, following a systematic protocol adapted from others [12,18,19]. Images were

converted to 8-bit grayscale, pre-processed to minimize background noise and

enhance contrast, and binarized by applying a pre-established threshold. Extraneous

pixels not corresponding to cells were eliminated using the *Remove outliers* and

Despeckle tools, with manual editing performed as necessary to address significant

background noise.

Following image pre-processing, the Skeleton and Fractal Analysis plugins of

ImageJ Fiji software [20] were used. Specific parameters assessed included the

number of junctions (Skeleton Analysis) and the convex hull area (*i.e.,* the area

containing the entire cell shape enclosed by the smallest convex polygon), fractal

dimension (*i.e.,* index of complexity), span ratio (*i.e.,* the ratio of the axes of the convex

hull from major to minor), and circularity (*i.e.,* how closely the shape of an object

resembles a circle) (Fractal Analysis). For detailed procedures, refer to the user

manual of the respective plugins [20].

2.8. Determination of the peripheral immune response

To evaluate the peripheral immune response, we determined the levels of systemic

immune mediators and the spleen cellularity. First, to quantify the levels of peripheral

inflammatory factors, blood samples were collected by cardiac puncture, allowed to

clot for 2h at RT, and then centrifuged at 2,000rpm for 10min. The concentration of

inflammatory mediators in the serum was determined by sandwich ELISA assay.

Briefly, maxisorb plates (Millipore) were coated O/N at 4°C with the capture antibodies

(anti-IL-6, anti-IL-12 from BD Pharmigen, or anti-MCP-1 from PeproTech). The plates

were then washed with PBS containing 0.05% Tween20 and blocked with PBS

containing 10% FBS for 2h. The samples and the recombinant standards were added

to the plates and incubated O/N at 4°C. Following incubation, biotinylated detection

antibodies (Serotec) were added for 1h at RT. After washing, plates were incubated

with avidin-HRP (Sigma) (1:500 in blocking buffer) for 1h and then developed with

ABTS substrate (Sigma). Absorbance was measured at 405nm using a

spectrophotometer (Molecular Devices) and concentrations of inflammatory factors

were determined based on the standard curve generated simultaneously.

Next, spleens were collected and weighed. Spleen tissue was gently ground

with the frosted part of two slides onto a 100mm culture dish. The homogenized tissue

was then transferred to RPMI medium supplemented with 10% FBS, 2mM L-

glutamine, 1% P/S, and 50μM 2-mercaptoethanol. After allowing the suspension to sit

for 5min on ice, the cell suspension was carefully collected and centrifuged at

1,670rpm for 10min at 4°C. To the resulting pellet, Ack buffer (150mM NH_4Cl, 10mM

$KHCO_3$, 0.1mM Na_2EDTA) was subsequently added for 10min at RT to lyse red blood

cells. Next, RPMI medium was added, and cells were centrifuged at 1,670rpm for

10min at RT and counted.

2.9. RNA extraction and evaluation of gene expression

Following MCAO, brains were collected and divided into the contralateral and

ipsilateral sides in both control and cortistatin-treated mice. For RNA isolation, brain

tissues were placed in TriPure reagent (Roche). Then, cDNA synthesis was carried

out using the RevertAid First Strand cDNA Synthesis Kit (ThermoFisher) with random

hexamers in a CFX Connect QPCR System (Biorad) under the following conditions:

incubation at 25°C/5 min, reverse transcription at 42°C/60 min, inactivation at 70°C/5

min. Gene expression analysis was conducted using conventional qPCR. SensiFast

Sybr No-Rox kit 1x (Bioscience) was added to the cDNA corresponding to 20ng of

RNA, along with the corresponding reverse and forward primers for each gene, and

nuclease-free water to a final volume of 20µl. The CFX96 thermocycler (BioRad) was

used to conduct the reaction under the following conditions: 94°C/5min (polymerase

activation) followed by 40 cycles at 94°C/30s (denaturalization), 58°C/60°C/62°C/30s

(primer annealing temperature depending on the gene), and 72°C/30s (extension).

The comparative threshold cycle (Ct) method was performed for quantification. Each

gene Ct values were normalized to those of the housekeeping gene *Rplp0* within each

PCR reaction. Fold change was determined using the ΔΔCt method. The specific

primers for each gene are provided in Additional file 1: Tables S1.

2.10. Statistical analysis

Data are presented as the mean ± SEM. All experiments were conducted in a blinded

and randomized manner. The number of independent animals (N), and tissue sections

and cell types (n) are indicated. Statistical differences between two groups were

conducted using the unpaired Student's t test for normal distribution and parametric

data, or the non-parametric Mann-Whitney U-test. For comparing three or more groups

that display normal distribution and parametric data, one-way ANOVA with Fisher´s

LSD test *post-hoc* tests for multiple comparisons was conducted. Three or more

groups that display non-parametric data were evaluated by Kruskal-Wallis test with an

uncorrected Dunn *post-hoc* test. Brown-Forsythe and Welch ANOVA with unpaired t-

test and Welch´s correction *post-hoc* test were performed in the case standard

deviations were not assumed to be equal. All statistical analyses were performed using

GraphPad Prism v8.3.0 software, with P-values < 0.05 (one-tailed) considered

statistically significant.

3. <u>RESULTS</u>

3.1. Identification of the optimal administration time of cortistatin in an

acute stroke model

To investigate the potential therapeutic effect of the neuropeptide in ischemic stroke,

we chose the MCAO model, a well-established preclinical model that closely mimics

the human ischemic condition [21]. To consider the different phases of the pathology

[2], we first administered cortistatin across different temporal contexts during stroke

progression (Fig. 1A).

We administered the neuropeptide at three distinct time points: immediately

post-MCAO (0h and 24h post-reperfusion), early post-MCAO (4h and 24h post-

reperfusion), and late post-MCAO (24h post-reperfusion). The immediate treatment

aimed to influence neuroinflammatory and neurodegenerative responses immediately

after ischemia. The early treatment targeted pathological processes occurring within

the first hours after hypoxia-reperfusion, while the late treatment focused on processes

already established during the acute phase of stroke.

To determine the effect of cortistatin, we assessed neurological scores,

behavioural impairments, and brain damage in C57BL/6J mice after administering

cortistatin at the specified time points. First, neurological deficits resulting from stroke

were evaluated within 24-48h after MCAO using a neurological scoring system

encompassing motor, balance, reflex, and sensory functions graded from 0 to 3

(modified from [15]). As shown in Fig. 1B, MCAO control mice (saline-treated)

exhibited a mean score of 0.5-1, indicating minor motor incoordination and slight

hypomobility. Immediate cortistatin treatment significantly worsened neurological

scores to 1-2, related to moderate intermittent circling, tremor twitches, and motor

incoordination. However, early and late treatments did not significantly alter the

neurological score compared to controls, despite the significant reduction in lesion size

with late treatment (Fig. 1E,F).

We further evaluated motor function and muscle coordination using the rotarod

and wire-hanging tests (Fig. 1C,D; Supplementary Fig. 1). In both tests, mice

performance was measured 24h before the surgery (baseline, BL), and at 24 and 48h

post-surgery.

Fig. 1. Optimization of the administration time of cortistatin in a preclinical stroke model. A.
Schematic representation of the experimental timeline. C57BL/6J mice were subjected to MCAO and

received either a saline solution (MCAO) or cortistatin (144µg/kg; MCAO+CST) at three distinct time
intervals: immediately following reperfusion (0h + 24h), early after reperfusion (4h + 24h), and at late
time post-reperfusion (24h). **B.** Neurological scores at 48h were assessed on a scale from 0 (no signs)
to 3 (death). **C,D.** Behavioural performances were assessed by rotarod and wire-hanging tests, both at
baseline (BL, pre-MCAO) and 48h post-MCAO. The time on the rod was evaluated in the rotarod test
(**C**). In the wire-hanging test (**D**), the number of falls (**left**) and escapes (**right**) toward one of the posts
were evaluated. Data are represented as the percentage of the total values based on the legend
classification. N = 6-13 mice/group. **E.** Representative coronal brain sections from mice treated with
either saline or CST at the specified times, stained with cresyl violet (CV). Ischemic lesions are outlined
with dashed red lines. Scale bar: 1500µm **F.** Ischemic volume across the brain was determined by
measuring ten CV-stained brain slices at 2-mm intervals per mouse. Data are expressed as
mean ± SEM. *vs.* MCAO-saline treated mice, *$p \leq 0.05$, **$p \leq 0.01$, ***$p \leq 0.001$. CST: cortistatin,
MCAO: middle cerebral artery occlusion.

As expected, both tests showed worsened performance in MCAO mice at 24h-

48h post-surgery compared to BL (Fig. 1C,D, Supplementary Fig. 1). Specifically at

48h, saline-treated MCAO mice displayed a reduced time on the rod compared to BL

(46.2% lasting less than 200 seconds *vs.* 90% of mice performing beyond this time

point at BL), which was similar in the early treatment but reduced with the immediate

cortistatin administration (50% and 71.4% of mice falling before 200 seconds,

respectively) (Fig. 1C; Supplementary Fig. 1A). In the wire-hanging test, stroke control

mice displayed a worsened performance and increased number of falls (only 53.8%

falling less than 3 times *vs.* 89.5% of BL, and only 23.1% escaping more than 6 times

vs. 84.2% of BL) (Fig. 1D, Supplementary Fig. 1B). Immediate treatment resulted in

poorer performance compared to saline and other treatments (66.6% falling more than

6 times *vs.* 23.1% saline, and 83.3% escaping less than 3 times *vs.* 53.8% saline) (Fig.

1D, Supplementary Fig. 1B,C). Early treatment also showed similar trends in

performance metrics (Fig. 1D, Supplementary Fig. 1C). However, although late

treatment showed no beneficial effect on the rotarod test, it improved performance in

the wire-hanging assay, with fewer animals falling more than 6 times compared to

saline and other treatments (14.3% vs. 23.1% saline) (Fig. 1D, Supplementary Fig.

1B,C). Similarly, the percentage of animals escaping more than 6 times increased

(28.6% vs. 23.1% saline or 0% for the other treatments) (Fig. 1D, Supplementary Fig.

1B,C).

Next, neurological lesions were quantified using CV staining, which

differentiates intact neuronal cytoplasm (intense purple), from infarcted areas with

dead neurons (faded purple) [22]. We observed that mice treated immediately after

MCAO displayed larger infarct volumes when compared to MCAO controls (Fig. 1E,F).

Remarkably, this treatment caused significant damage in the cortex and the striatum,

spreading to the hippocampus area (data not shown), consistent with the observed

worsened scores. Early treatment also led to larger lesions compared to MCAO-saline

mice but to a lesser extent than the immediate treatment (Fig. 1E,F). In contrast, late

treatment with cortistatin significantly decreased infarct volume, with only 23% of mice

treated at 24h showing infarct volumes above 20mm³, compared to 44% of controls,

100% of immediately treated mice, and 66% of early-treated mice (Figure 1C,F).

3.2.   ***Differential time administration of cortistatin affects stroke-derived***

328   ***glial responses***

To investigate the underlying mechanisms by which cortistatin might differentially

affect ischemic lesions and neurological deficits when administered at different times,

we analysed the local neuroinflammatory response. In ischemic stroke, microglia are

very early responders. These cells swiftly migrate to the infarct core, proliferating and

adopting an amoeboid phenotype to engulf dead neurons [23]. Interestingly,

immediate and early treatment with cortistatin resulted in reduced microglia density in

the ischemic core compared to MCAO controls, while late treatment increased

microglia density, potentially indicating a more physiological response to the injury

(Fig. 2A,B).

Since morphological alterations in microglia closely correlate with their reactivity

states and serve as indicators of the severity of brain injury [18,24], we conducted

Skeleton Analysis to evaluate microglia morphotypes. In the infarct core, microglia in

saline-treated MCAO mice displayed a typical reactive amoeboid-like shape (Fig.

2C,D). Remarkably, immediate treatment with cortistatin led to a more pronounced

reactive phenotype than controls, characterized by fewer junctions, and larger area.

Conversely, early treatment resulted in a phenotype similar to MCAO mice, but with

an increased number of ramifications, while late treatment was associated with an

increased number of junctions and a significantly reduced soma, suggesting a

transition to a less ameboid microglia state (Fig. 2C,D).

Microglia in the ischemic penumbra also undergo intricate morphological and

functional changes [25]. We observed that immediate and early treatments increased

the number of junctions and overall microglial size, indicating a hyper-ramified

morphotype that was particularly pronounced in the early treatment. In contrast, late

treatment resulted in smaller soma sizes, while the number of junctions and microglia

area were similar to those in the MCAO control group, suggesting a different reactive

state (Fig. 2E,F).

Fig. 2. Time-dependent cortistatin treatment differentially regulates microglia response after stroke. A. Brain coronal sections of C57BL/6J mice were obtained after MCAO and subsequent treatment with either saline solution (MCAO) or cortistatin (144μg/kg; MCAO+CST) at three distinct time intervals: immediately following reperfusion (0h + 24h), early after reperfusion (4h + 24h), and at a late time post-reperfusion (24h). Microglia was identified by Iba1 staining. The ischemic core is represented by dashed red lines. Scale bar: 1,500μm. **B.** Density of microglia (cells/mm^2) was measured in the ischemic core in the above-mentioned mice. **C-E.** Representative binary microglia from the different experimental groups from the ischemic core (red square) (**C**), or the penumbra (yellow square) (**E**). Scale bar: 30μm. Morphological parameters analysed (**D,F**). Skeleton Analysis was used to assess the number of junctions (**D,F** left panel). Cell area (**D,F** middle panel) and soma size (**D,F** right panel) were manually measured. n = 15-20 cells/mice; N = 3 mice/group. Data represent the mean ± SEM. *vs. saline *$p ≤ 0.05$, **$p ≤ 0.01$, ***$p ≤ 0.001$, ****$p ≤ 0.0001$. CST: cortistatin, Iba1: ionized calcium-binding adaptor molecule 1, MCAO: middle cerebral artery occlusion.

Parallel to the microglia response, astrocytes also become reactive and establish a glial scar surrounding the wound, containing the injury and preventing its expansion to adjacent still-viable regions [26] (Fig. 3A)

Remarkably, our study found that the immediate cortistatin treatment impaired glial scar formation, resulting in decreased scar thickness when compared to MCAO control mice. Nevertheless, early and especially late treatments significantly increased glial scar formation (Fig. 3A,B). Of note, the disorganized glial scar in the immediate treatment match with the severe neuronal lesion and the reduced microglial density in the core (Fig. 1E,F; 2B). We further analysed astrocyte morphology using Skeleton Analysis to explore the influence of cortistatin treatment in the astrocytic response. While astrocytes from MCAO mice displayed a high number of junctions, immediate treatment resulted in a significant reduction, suggesting a more quiescent astrocytic phenotype (Fig. 3C,D). Astrocytes from mice with early treatment were similar to the saline-treated MCAO mice, while late treatment showed astrocytes with reduced junctions and area, but increased soma size (Fig. 3C,D), suggesting a distinct reactive phenotype, which can be linked to the enhanced glial scar formation (Fig. 3A,B).

Fig. 3. Time-dependent cortistatin treatment differentially impacts astroglial scar formation. A. Brain coronal sections of C57BL/6J mice were obtained after MCAO and subsequent treatment with either saline solution (MCAO) or cortistatin (115µg/kg; MCAO+CST) at three distinct time intervals: immediately following reperfusion (0h + 24h), early after reperfusion (4h + 24h), and at a late time post-reperfusion (24h). Astrocytes were identified by GFAP staining. Glial scar is outlined with dashed red lines. Scale bar: 1,500 µm. **B.** Glial scar thickness was quantified. N = 4-6 mice/group. **C.** Representative images of astrocytes in binary were extracted from the glial scar in the different experimental groups. Scale bar: 30µm. **D.** Evaluation of morphological parameters of astrocytes. Number of junctions (**D,** left panel) was evaluated by Skeleton Analysis. Cell (**D.** middle panel) and soma size (**D,** right panel) were manually measured. n = 15-20 cells/mice; N = 3 mice/group. Data are presented as the mean ± SEM. *vs.* saline, *$p \leq 0.05$, **$p \leq 0.01$, ***$p \leq 0.001$, ****$p \leq 0.0001$. CST: cortistatin, GFAP: glial fibrillary acidic protein, MCAO: middle cerebral artery occlusion.

Collectively, these results suggest that during the hyperacute stroke phase, the

initial glial response may be partially or fully deactivated by the potent anti-

inflammatory effects of the neuropeptide when administered immediately or early. This

deactivation leads to diminished protection of the damaged area, exacerbating

neuronal lesions and motor deficits. In contrast, late cortistatin administration

modulates the neuroinflammatory response differently, regulating glial cell

hyperreactivity and/or proliferation, which results in reduced neuronal lesions and

improved outcomes.

3.3.   Treatment with cortistatin at later stages of ischemic stroke exerts

neuroprotective effects

Based on these findings, we conducted a detailed investigation into the effects of

cortistatin administered at 24h on neurodegeneration, glial population, BBB integrity,

brain vasculature alterations, and local as well as peripheral immune responses at 48h

following MCAO.

Our initial observations revealed that cortistatin-treated mice display reduced

area of neuronal damage in both cortex and striatum compared to MCAO mice, as

evidenced by MAP-2 neuronal staining (Fig. 4A,B), which was consistent with CV

staining (Fig. 1E,F). Notably, ischemic injury coincided with decreased expression of

neurotrophic factors such as BDNF and GDNF. Following treatment with cortistatin,

the levels of these factors were significantly restored to baseline levels (Fig. 4C).

Fig. 4. Late administration of cortistatin promotes neuroprotection. **A**. Brain coronal sections from C57BL/6J mice subjected to MCAO and treated with either saline (MCAO) or cortistatin (144µg/kg) (MCAO+CST) at 24 hours post-reperfusion and sacrificed at 48 hours. Sections were stained with MAP-2 to visualize neurons. Ischemic lesions, corresponding to a lack of MAP-2 staining, are delineated with dashed red lines. Scale bar: 1,500µm. **B**. Quantification of the ischemic lesion area. N = 13 (MCAO), 8 (MCAO+CST) mice/group. **C**. Gene expression levels in the contralateral (CL, uninjured) and ipsilateral (ischemic) brain hemispheres of mice treated with saline or cortistatin following MCAO were analysed by real-time qPCR and normalized to *Rplp0* expression. N = 4-6 mice/group. Data are presented as mean ± SEM with dots representing individual mice. *vs*. saline, *$p \leq 0.05$, **$p \leq 0.01$. BDNF: brain-derived neurotrophic factor, CST: cortistatin, GDNF: glial cell line-derived neurotrophic factor, MAP-2: microtubule-associated protein 2, MCAO: middle cerebral artery occlusion.

3.4. Late treatment with cortistatin induces a neuroprotective microglial response in the acute stroke model

Next, we aimed to delve deeper into the role of cortistatin in modulating microglial morphotypes and functions, particularly focusing on its impact when administered at a later time post-stroke. To this aim, we complemented our initial morphometric analysis with a Fractal Analysis to evaluate various complexity parameters of microglia

from the ischemic core and penumbra in MCAO mice treated with cortistatin (Fig.5;

Supplementary Fig.2).

MCAO animals exhibited a reactive microglial phenotype in the ischemic

penumbra, characterized by a smaller territory area, enhanced complexity (as

indicated by span ratio and fractal dimension), and decreased circularity compared to

contralateral microglia (Fig. 5B,C). However, treatment with cortistatin seemed to

mitigate this reactivity, evidenced by reductions in the soma size, junctions, territory

area, and span ratio, along with an increase in complexity relative to saline-treated

mice (Fig. 2E,F; Fig. 5C). These changes suggest a potential shift towards a less

reactive phenotype or a different form of reactivity. In the ischemic core, although

microglia from MCAO and cortistatin-treated mice looked similar, the treatment still led

to distinct microglia with reduced soma size, increased junctions, and decreased span

ratio (Fig. 2C,D; Supplementary Fig. 2A,B).

To further understand the functional implications of these microglial changes,

we analysed the expression of different microglia markers in the ipsilateral hemisphere

of saline and cortistatin-treated mice compared to the contralateral hemisphere (Fig.

5D). Following MCAO, there was enhanced expression of microglial reactivity markers

such as *CD68* and *Trem2*. Remarkably, cortistatin treatment further upregulated the

expression of *CD68*, *Trem2*, and *Iba1* compared to the contralateral hemisphere, and

compared to MCAO mice (although the latter was not significant) (Fig. 5D).

Interestingly, although no changes were observed for the gene expression of CD206

between healthy and ischemic brain samples (Supplementary Fig. 3A), there was a

significant decrease in *CD163* expression in cortistatin-treated animals, advocating a

potential promotion of both anti-inflammatory and increased phagocytic activity of

these cells, consistent with the morphological findings and increased microglial density

observed in the ischemic core (Fig. 2).

Fig. 5. Late administration of cortistatin induces neuroprotective microglia. A. Brain coronal
sections from C57BL/6J mice subjected to MCAO and treated with either saline (MCAO) or cortistatin
(144µg/kg) (MCAO+CST) at 24 hours post-reperfusion and sacrificed at 48 hours. Microglia was
identified by Iba1 staining. Ischemic core is represented by dashed red lines. Scale bar: 1,500 µm. B.
Representative binary images of microglia from the ischemic penumbra (yellow square) and the
contralateral side (CL) (white square) from the distinct experimental groups. Scale bar: 30 µm. C.
Morphological parameters of microglia were analysed using Fractal Analysis including convex hull area,

fractal dimension/complexity, span ratio, and circularity. n = 15-20 cells/mice; N = 3 mice/group. **D**.
Gene expression levels of *CD68, TREM-2, Iba1,* and *CD163* in the contralateral and ipsilateral brain
hemispheres of saline and CST-treated MCAO mice were measured by real-time qPCR and normalized
to *Rplp0* expression (N=4-6 mice/group). Data are presented as mean ± SEM with dots representing
individual mice. *vs.* MCAO, *$p \leq 0.05$, **$p \leq 0.01$, ***$p \leq 0.001$, ****$p \leq 0.0001$. CD68: cluster of
differentiation 68, CD163: cluster of differentiation 163, CST: cortistatin, Iba1: ionized calcium-binding
adaptor molecule 1, MCAO: middle cerebral artery occlusion, TREM-2: triggering receptor expressed
on myeloid cells 2.

3.5.   Late administration with cortistatin enhances glial scar formation and

astrocyte reactivity

Given the collaborative role of microglia and astrocytes in the context of stroke to limit

the damage and prevent its spread [26], we assessed the impact of late cortistatin

treatment on the neuroinflammatory response of astrocytes (Fig. 6). Cortistatin

treatment resulted in thicker glial scars with astrocytes exhibiting a reduced territory

area, complexity index (fractal dimension), and circularity, and an increased span ratio,

compared to control MCAO mice (Fig. 6A-C).

Along with the fewer junctions, increased area, and enlarged soma shown in

Fig.3, these parameters indicate that cortistatin induces a distinct morphotype of

hypertrophic reactive astrocytes. To validate our findings, we examined the expression

of *GFAP,* a well-known marker of astrocyte reactivity [27]. Following MCAO, both

saline and cortistatin-treated mice showed increased *GFAP* expression compared to

the contralateral side. However, cortistatin treatment significantly reduced *GFAP*

expression relative to MCAO control mice. Thus, these results coupled with the

upregulation of neurotrophic and glial factors after cortistatin treatment (Fig. 4,5),

suggest that the reactive morphotype exhibited by these astrocytes may exert a role

in the establishment of an efficient glial scar and potentially linked to reduced neuronal

damage.

Fig. 6. Late administration of cortistatin promotes protective astrocytic responses in ischemic stroke. A. Brain coronal sections from C57BL/6J mice subjected to MCAO and treated with either saline (MCAO) or cortistatin (144µg/kg) (MCAO+CST) at 24 hours post-reperfusion and sacrificed at 48 hours. Sections were stained with GFAP antibody to identify astrocytes. The astrocyte scar is indicated by dashed red lines. Scale bar: 1,500µm. **B.** Representative binary images of astrocytes from the glial scar were evaluated. Scale bar: 30 µm. **C.** Morphological parameters of astrocytes (convex hull area, fractal dimension/complexity, span ratio, and circularity) were quantified using Fractal Analysis. n = 15-20 cells/mice; N = 3 mice/group. **D**. Gene expression levels of *GFAP* in the contralateral and ipsilateral brain hemispheres of saline and CST-treated MCAO mice were quantified by real-time qPCR and normalized to *Rplp0* expression (N=4-6 mice/group). Data are presented as mean ± SEM with dots representing individual mice. *vs MCAO, *$p \leq 0.05$, ***$p \leq 0.001$, ****$p \leq 0.0001$. CST: cortistatin, GFAP: glial fibrillary acidic protein, MCAO: middle cerebral artery occlusion.

513   Together, these findings highlight cortistatin as a key regulator of glial responses

514   in stroke pathology supporting neuroprotective properties.

515   **_3.6.   Cortistatin protects against BBB breakdown and improves vascular_**

516   **_dynamics after stroke_**

Stroke is accompanied by the breakdown of the BBB, contributing to neurovascular

damage and complications. To thoroughly explore the extent of BBB dysfunction after

ischemia in late cortistatin-treated mice, we evaluated IgG leakage. This is one of the

key indicators of BBB breakdown, reflecting the extravasation of endogenous blood

proteins. In our model, IgG leakage was prominently observed in the ipsilateral

hemisphere after stroke, but significantly reduced following cortistatin treatment (Fig.

7A,B).

Fig. 7. Cortistatin diminishes BBB disruption after stroke. A. Brain coronal sections from C57BL/6J mice subjected to MCAO and treated with either saline (MCAO) or cortistatin (144µg/kg) (MCAO+CST) at 24 hours post-reperfusion and sacrificed at 48 hours. Sections were stained with a fluorescently-labelled antibody against mouse IgGs. IgG leakage areas are outlined with dashed white lines. Scale bar: 1,500 µm. **B.** IgG$^+$ area was measured as a quantification of IgG leakage from the blood. N = 4-8 mice/group. **C.** Representative images taken in the ipsilateral striatum of mice of the two experimental groups following MCAO and treated with CST, stained with CD31 antibody for brain endothelial cells (BECs, green) and ZO-1 (red) for TJs. ZO-1 gaps are highlighted with an asterisk, while ZO-1 coverage recovery is indicated with arrows. Scale bar: 30 µm. **D.** Quantification of the number of ZO-1 gaps per vessel area. n = 3 ROIs/section/mice; N = 3-5 mice/group. **E.** Representative images of BECs (CD31, green) from the ipsilateral striatum of MCAO or cortistatin-treated mice. Scale bar: 30 µm. **F.** Vessel diameter was manually measured, and vascular density was calculated as the % of positive CD31 staining of the total area of the image. n = 10-20 measures/vessel with 5-10 vessels/mice; N = 3-6 mice/group. Data are presented as the mean ± SEM. *$vs.$ MCAO, *$p \leq 0.05$, **$p \leq 0.01$, ****$p \leq 0.0001$. CD31: cluster of differentiation 31, CST: cortistatin, IgG: immunoglobulin G, MCAO: middle cerebral artery occlusion, ZO-1: zonula-occludens 1.

After occlusion and reperfusion processes, hypoxic conditions particularly affect brain endothelial cells (BECs), disrupting TJs, leading to increased permeability and contributing to the development of vasogenic edema [28]. We observed that following MCAO, ZO-1 expression in the ischemic core showed disorganized patterns with increased gaps (Fig. 7C,D). Among TJs, disruption of ZO-1 has been extensively documented following stroke [29,30], playing a crucial role in BBB breakdown and subsequent recovery. Conversely, cortistatin treatment mitigated this disorganization, leading to a reduction in the number of ZO-1 gaps, thereby potentially improving TJ integrity (Fig. 7D). We also observed a decrease in the expression of *claudin-1* after MCAO *vs.* the contralateral side, with no alterations after cortistatin treatment (Supplementary Fig. 3B). Additionally, alterations in basement membrane proteins like collagen type I (Col1a2) were observed post-stroke, with cortistatin treatment showing a trend towards increasing *Col1a2* expression levels compared to MCAO alone (Supplementary Fig. 3B). In response to stroke, vascular dynamics undergo significant changes to promote angiogenesis and the modification of the vascular tree [31]. To investigate the importance of this process in stroke recovery, we examined the area occupied by vessels (CD31$^+$) and their diameter in the striatal core (Fig. 7E).

Cortistatin treatment enhanced vessel diameter and vascular density in the ischemic

striatum compared to saline-treated MCAO mice (Fig. 7F). This enhancement in

vascular parameters may facilitate improved reperfusion to the damaged area,

supporting post-stroke recovery mechanisms.

These findings suggest that cortistatin not only protects against BBB

permeability breakdown but also promotes angiogenic and vascular remodelling

events that aid in vascular repair and regeneration post-stroke.

3.7. Cortistatin modulates central and peripheral immune responses

following stroke, reducing brain infiltration and immunodepression

After ischemia-reperfusion injury, the brain undergoes oxidative stress and neuronal

death, triggering glial cell activation and the secretion of cytokines and other immune

factors. Following MCAO, we observed increased expression of *TNF-α* and *IL-6* in the

untreated ischemic brain hemisphere compared to the contralateral side, with no

differences in *IL-10* expression. Interestingly, cortistatin treatment significantly

elevated *TNF-α* and *IL-10* levels (Fig. 8A). Levels of *IL-1β* showed no significant

differences between groups. Additionally, ischemia-induced arginase 1 (*Arg1*)

expression, indicative of brain immune infiltration and stroke severity [32,33], was

significantly downregulated by cortistatin (Supplementary Fig. 3C). This suggests that

cortistatin influences the polarization of immune cells towards a neuroprotective

phenotype.

Cytokines and DAMPs derived from brain damage can be released into the

systemic circulation, getting to lymphoid organs and initiating the peripheral

inflammatory response [34]. This triggers the quick adhesion of platelets and

leukocytes to the endothelium facilitated by BBB disruption, enabling peripheral

immune cells to infiltrate the brain [35]. To explore how cortistatin influences peripheral

immune response dynamics, we measured cytokine levels in serum and assessed the

infiltration of immune cells into the ischemic core. We observed that in MCAO mice,

significant immune infiltration (CD45+ cells) occurred in the ischemic region, which

was markedly reduced following cortistatin treatment (Fig. 8B,C). This suggests that

cortistatin limits the infiltration of peripheral immune cells into the injured brain,

possibly through its effects on BBB integrity.

The initial, short-lasting activation of systemic immunity by DAMPs and cytokines

released in the bloodstream is swiftly followed by long-lasting immunodepression,

characterized by lymphopenia and reduced functional activity of T cells and monocytes

among others [36]. In our model, stroke-induced immunodepression was

characterized by an oedematous and fibrotic spleen with decreased splenocyte

numbers. Cortistatin treatment reversed these effects, significantly increasing the total

number of splenocytes and reducing spleen weight after stroke (Fig. 8D). Furthermore,

cortistatin treatment restored the reduced stroke-associated serum levels of IL-12 and

IL-6 to homeostatic conditions, and increased MCP1 levels, indicating a systemic

modulation of inflammatory responses (Fig. 8E).

Fig. 8. Cortistatin modulates immune infiltration and peripheral immune response after ischemic stroke. **A.** Gene expression levels for *TNF-α*, *IL-6*, *IL-1β*, and *IL-10* in the contralateral and ipsilateral brain hemispheres, of saline and CST-treated MCAO mice were quantified using real-time qPCR and normalized to *Rplp0* expression. N = 4-6 independent mice. **B.** Representative brain coronal sections stained with CD45 antibody (red) for leukocytes from C57BL/6J mice subjected to MCAO and treated with saline (MCAO) or cortistatin (MCAO+CST) at 24h post-reperfusion and sacrificed at 48h. Scale bar: 100 μm. **C.** Leukocyte density (number of cells/mm^2) was measured in the striatum and the contralateral side (CL). **D.** Spleen weight and the number of splenocytes/ml were measured. N = 4-8

independent mice. **E.** Serum levels of IL-12, IL-6, and MCP-1 (ng/ml) in serum were measured using
an ELISA assay. Dashed lines represent cytokine concentrations in healthy mice. N = 3-6 independent
mice. Data are the mean ± SEM with dots representing individual mice. *vs.* MCAO, #*vs.* healthy mice
*,#$p \leq 0.05$, **,##$p \leq 0.01$, ***$p \leq 0.001$, ****$p \leq 0.0001$. CD45: cluster of differentiation 45, CST: cortistatin,
IL-6: interleukin 6, IL-1β: interleukin 1β, IL-10: interleukin 10, IL-12: interleukin 12, MCAO: middle
cerebral artery occlusion, MCP-1: Monocyte Chemoattractant Protein-1, TNF-α: tumour necrosis factor-
α.

In summary, cortistatin administered with an appropriate post-stroke timing

exerts multifaceted beneficial effects. It enhances neuroprotective cytokine profiles in

the brain, reduces neuronal damage, potentially through the modulation of microglia

reactivity and enhancing the formation of the astrocytic scar, regulates BBB integrity

preventing immune cell infiltration, and modulates systemic inflammatory responses.

Additionally, cortistatin counteracts stroke-induced immunodepression by restoring

splenic immune function. These findings underscore the therapeutic potential of

cortistatin in stroke by not only protecting neuronal tissue and promoting vascular

recovery but also by regulating immune responses to support overall recovery

mechanisms.

4. <u>DISCUSSION</u>

The pathogenesis of stroke involves a complex interplay among nervous, circulating,

and cerebral immune cells, which can play dual roles influenced by their spatial

location, context, and timing [34]. Despite the significance of these factors, many

cytoprotective therapeutic approaches have overlooked them, instead focusing on

time windows that do not align with the underlying pathophysiology. Typically,

interventions target the hyper-acute stage to fit the time range of rtPA, or slightly later,

without discriminating the different phases. Therefore, an increased awareness of the

intricate spatiotemporal relationships between the nervous, vascular, and immune

systems may prove essential for the design of novel therapeutic approaches to

improve stroke prognosis.

In this context, our findings advocate that the neuropeptide cortistatin, which

provides a crucial link between the CNS and the immune system, could be a promising

therapeutic agent, positively influencing different aspects of stroke pathophysiology

when administered at appropriate stages. Surprisingly, when cortistatin was given

immediately after blood flow restoration, it worsened neuronal damage, motor

performance, and neurological scores. The decreased microglial density with

increased ameboid-like microglia in the core and the hyper-ramified microglial

phenotype in the penumbra might be related to the potent immunomodulatory effects

of cortistatin. The neuropeptide administered shortly after reperfusion may blunt the

initial endogenous response of microglia that tries to limit the ischemic damage.

Cortistatin may have prompted microglia to respond with an initially protective but later

excessively reactive phenotype. Furthermore, the immediate cortistatin treatment

appeared to hinder the initial astrocytic response, as evidenced by the disorganized

astrocytic scar, preventing the formation of well-defined astrocytic borders, and

potentially facilitating the expansion of neuronal damage.

Conversely, treatment with cortistatin at a slightly later stage (4h + 24h) did not

reduce infarct volume or improve neurological outcomes. This treatment did not affect

microglia density or morphotype in the core but induced a hyper-ramified microglia

phenotype in the perilesional area. The early treatment with cortistatin enhanced the

formation of the glial scar, possibly because it coincided with the initiation of the

astrocytic response [37], modulating astrocytes differently than at the 0-hour time-point

treatment. These findings underscore the critical importance of timing in therapeutic

intervention, as even a few hours can have a strong effect on the response to the

treatment. Indeed, late cortistatin administration (24h) appeared to lead to reduced

neuronal lesions and improved outcomes.

The lack of protection observed with the immediate administration might seem

contradictory to the beneficial effect shown at 24h and to the reported beneficial effects

of cortistatin administration in other neurodegenerative/neuroinflammatory disorders

[8,11,12]. However, the latter studies involved cortistatin injection at later stages of the

disorder (*i.e.,* when the injury is well-established), while in our investigation the

immediate administration coincided with the initial stages of injury. This suggests that

the observed harmful effects on outcome are not likely due to cortistatin being

neurotoxic but rather that when administered immediately or early after stroke,

cortistatin interferes with the initial local neuroinflammatory response that is essential

for containing the damage and removing debris and dead cells (*i.e.,* glial scar

formation and microglia response, respectively), thereby leading to detrimental

neurological outcome. The timing of cortistatin administration therefore appears vital

to the subsequent progression of brain healing processes. Indeed, blocking or

modifying the important early response leads to adverse outcomes in other disorders.

For instance, Szalay *et al.* demonstrated that depletion of microglia resulted in a

significant enhancement of lesion size, which was reversed by microglial repopulation

[38]. These findings may also explain the lack of clinical success reported with some

anti-inflammatory/immunomodulatory drugs administered at too early phases [3].

Collectively, these data underscore the fundamental role of the inflammatory response

in the regenerative process after stroke and suggest that these mechanisms should

not be completely halted. Instead, modulating the inflammatory response, rather than

attempting to stop it entirely, seems to be a more effective approach.

Given these first results, we focused on a deeper investigation of the effect of

cortistatin administration at the late stage. This approach, besides providing more

flexibility in treatment application, also enables concurrent administration with other

therapies such as rtPA. First, we confirmed the reduced striatal lesion size following

late (24h) cortistatin treatment based on CV and MAP-2 immunodetection. Notably,

MAP-2 provides a better insight into the first phases of neuronal death, even if the

process is not yet concluded [39]. This indicated that cortistatin efficiently targeted

neuronal death, protecting neuronal bodies, also evidenced by the increased

expression of neuroprotective factors such as *BDNF* or *GDNF*. GDNF is important for

the maintenance of dopaminergic neurons and their protection against injury [40],

while BDNF protects neurons from ischemia-reperfusion-induced apoptosis [41]. The

neuroprotective role of cortistatin conforms with previous research, where we and

others demonstrated that treatment with the neuropeptide protected neurons from

distinct insults *in vivo* and *in vitro* [12,42]. However, it is not clear yet the underlying

mechanisms (*i.e.,* inhibition of apoptosis, neuronal degeneration, or oxidative

damage). Moreover, we can hypothesize that cortistatin might also exert an indirect

beneficial effect on neurons by regulating the responses of glial cells.

For decades, ischemic stroke research has primarily focused on rescuing

neurons in the penumbra region to prevent the expansion of the infarct core. However,

given that excitotoxicity, oxidative stress, neuroinflammation, or peripheral immune

response contribute to secondary neuronal cell death [43] and that focusing solely on

neuronal death is oversimplistic [44], this study aimed to explore the impact of

cortistatin on the stroke-induced neuroinflammatory response. Microglia are the initial

responders to ischemic damage, swiftly undergoing dynamic morphological and

molecular changes in the infarct core and the peri-infarct zone [45]. Importantly,

microglia undergo multiple alterations after injury (*e.g.,* degree and length of

branching, branch configuration, shape and soma size, cell territory, receptor

distribution, or cytoskeletal rearrangement), that go beyond the oversimplified

paradigm of M1/M2 or the traditional morphological characterization categorizing microglia as either ramified (homeostatic) or reactive (short processes) [46]. Our findings showed an increased number of microglial cells in the ischemic core after late treatment with cortistatin. Moreover, elevated expression of markers related to phagocytosis (*i.e., CD68, Trem2*) was also found in the ipsilateral hemisphere of cortistatin-treated mice, consistent with the enhanced phagocytic phenotype and neuroprotective microglial function reported in early stages of stroke [47,48]. Since cortistatin significantly reduced brain lesions, these findings suggest an enhanced phagocytosis of dead neurons by the microglia from cortistatin-treated mice. Indeed, previous studies have demonstrated that endogenous cortistatin is essential for phagocytosis in physiological and demyelinating environments [49]. However, hyperreactive microglia over time can engulf live or stressed neurons, resulting in worsened neurological deficits [50], emphasizing the need to maintain a careful balance to ameliorate stroke prognosis. Microglia in the ischemic core exhibited reduced soma size and a larger number of junctions following cortistatin treatment, suggesting a shift to a more protective phenotype, diverging from the ameboid-like shape phenotype observed in control mice. While most microglia are engaged in engulfing dead neurons and secreting neurotrophic factors in the ischemic core, those in the peri-infarct area exhibit a different phenotypic polarization, influenced by surrounding factors [51,52]. Cortistatin treatment shifted peri-infarct microglia to a distinct reactive phenotype characterized by a reduced territory and soma but increased complexity. Moreover, elevated levels of *TNF-α*, *IL-10*, and neuroprotective factors such as *BDNF* or *GDNF* were detected. These results suggest a potential transformation to a distinct subtype of protective/reactive microglia, capable of phagocytosing cellular debris while aiding brain regeneration [53]. This is confirmed

by previous findings in other neuroinflammatory models (*i.e.,* multiple sclerosis,

Parkinson's disease, and sepsis) [8,12,54], where cortistatin-modulated microglia

markedly correlated with a better disease prognosis. Conversely, we observed a

decreased expression of the scavenger receptor *CD163*, involved in the clearance of

haemoglobin [55]. Interestingly, this factor is upregulated in patients with intracerebral

haemorrhage and edema development [56], suggesting a potential reduction of these

events following cortistatin treatment. Overall, the unique combination of phagocytic

function, inflammatory cytokines, and neurotrophic/neuroprotective factors secreted

by these cells may play a crucial role in promoting neuron survival and modulating

other neuroinflammatory responses. Although many questions remain unanswered

regarding the transitions between the different subsets of glial cells and the key

molecular cues that drive these transitions [23], our study indicates that cortistatin may

play an important role in this context. However, acknowledging the limitations of our

study, we could not determine the exact molecular and functional morphotype of these

cells in terms of spatial and temporal dynamics, since the gene expression analysis

could not discriminate between different brain areas within the same hemisphere,

different cell populations, or different cell subsets. Additionally, we cannot dismiss the

fact that more than one cell type produces the same factor, further complicating the

interpretation of our results. To address these limitations, further studies are needed

to explore the exact mechanisms through which cortistatin-modulated microglia exert

this immunomodulatory role.

Astrocytes also play essential roles in stroke, modulating the

neuroinflammatory response, protecting neurons by releasing neuroprotective factors,

forming barriers to limit damage, and promoting brain repair [57]. Although the

molecular and cellular mechanisms that underlie the formation of the glial scar and

astrocyte reactivity are not completely understood, they likely involve a complex and balanced interplay of molecular signals that can simultaneously boost phagocytosis and debris clearance, which would be essential for the protection and conservation of the still-healthy tissue [37,57,58]. Similar to microglia, an A1/A2 paradigm for astrocyte morphology and function has been proposed [59,60]. However, the spatiotemporal heterogeneity of reactive astrocytes has demonstrated that this classification is too simple, so additional parameters should be considered [61]. Our study revealed that cortistatin-treated mice exhibited a well-organized and thicker glial scar than control mice. This was accompanied by an increased expression of neurotrophins, which have been described to be involved in synaptic plasticity and neuronal growth regulation that tend to decrease in pathological CNS processes [62]. Thus, cortistatin promotes a distinct reactive phenotype that supports tissue repair and neuroprotection. An effective glial scar is critical as defective scar formation has been associated with larger lesions, demyelination, and reduced neuronal survival and functional recovery [37,58,63]. However, an excessive glial scar can be detrimental by exacerbating inflammation and interfering with synaptic and axonal growth [37]. The observed localization and cortistatin-dependent timing of astrocyte activation in our study indicate a controlled and neuroprotective response to limit damage following cortistatin treatment.

In addition, BBB integrity is needed for better stroke outcomes [64]. The traditional classification of BBB changes during stroke progression is restricted to "opening" or "closing" (*i.e.,* "bad" or " good") making this dichotomy outdated [65]. Instead, BBB permeability follows a sequence across different phases (*i.e.,* hyperacute, acute, subacute, and chronic phases), each of them characterized by distinct biological events [65]. Our study specifically focused on the acute phase in

which BBB permeability peaks, facilitating neuroinflammation progression and exacerbating injury [66]. Our findings demonstrated that cortistatin treatment helped preserve BBB integrity by reducing endothelial dysfunction and maintaining tight junctions, as evidenced by the decreased IgG leakage and reduced number of ZO-1 gaps. These results align with previous studies showing that cortistatin protects the BBB in models of mild neuroinflammation and *in vitro* models of BBB disruption [13]. Interestingly, the lack of changes in claudin-1 and claudin-5 gene expression suggests that TJ alterations may occur at the protein localization level rather than at the gene expression level [67–69]. Indeed, cortistatin has demonstrated a role in modulating protein levels, cellular redistribution, and functional organization of BBB proteins such as ZO-1, claudin-1, claudin-5, and VE-cadherin without significantly affecting their transcript levels [13].

Finally, it is important to acknowledge that stroke is a complex, organism-wide process that not only targets the brain but also peripheral tissues. Shortly after the interruption of the CBF and subsequent hypoxia, ECs express selectins in response to oxidative damage, quickly engaging circulating innate immune cells, and triggering a rapid pro-inflammatory response in the periphery [2]. The breakdown of various components of the BBB enables the extravasation of these cells, with different cadences and at variable times. In general, a well-regulated inflammatory response is inherently self-limiting, facilitating the repair, turnover, and tissue adaptation, and prioritizing processes that restore homeostasis over those that are harmful [70,71]. The immunomodulatory effects of cortistatin were evident as it reduced immune cell infiltration into the brain and modulated systemic immune responses, consistent with its effects observed in other neuroinflammatory models [8,13]. Specifically, cortistatin treatment modulated the expression of typically pro-inflammatory cytokines (*e.g.,* IL-

6, TNF-α, IL-12) in the brain. These cytokines exert biphasic or even polyphasic roles

in ischemic stroke, and depending on the timing, spatial location, and severity of the

injury, they can play vital roles in repair processes or have detrimental effects. For

example, it has been shown that the deletion of IL-6 in MCAO mice does not confer

protection against cerebral ischemia [70]. Indeed, IL-6 serves as a neurotrophic factor,

improving functional outcomes, reducing stroke volume in mice [72], and promoting

post-stroke angiogenesis [73]. Similarly, numerous studies have reported a

neuroprotective role of TNF-α in reducing the production of nitric oxide and free

radicals, maintaining neuronal calcium homeostasis, and inducing the synthesis of

neurotrophic factors [74]. Hence, these data, along with the increased levels of anti-

inflammatory factors (*i.e.,* IL-10), the reduced number of infiltrating cells, and the

decreased neuronal damage suggest that the elevated inflammatory factors observed

after the treatment with the neuropeptide might be crucial for developing an

appropriate response to the acute damage and subsequent resolution rather than

playing a detrimental role. This underscores the complexity of the immune response

in stroke and, again, the importance of the timing of therapeutic intervention.

This initial activation of systemic immunity is quickly succeeded by a prolonged

state of immunodepression, which enhances the susceptibility to infection or other

pathologies [36]. Depression of the immune system not only occurs in patients with

ischemic stroke but also in other acute CNS injuries such as cerebral haemorrhage,

traumatic brain injury, or spinal cord trauma (reviewed in [36]). In this sense, the

observed potent immunomodulatory role of cortistatin, especially regarding spleen

function and cytokine recovery to homeostatic levels, is consistent with other studies

[8,75,76]. Interestingly, we found elevated levels of MCP-1 after cortistatin treatment.

Although this factor is systemically upregulated in stroke patients and is known to

promote leukocyte chemotaxis, some reports also demonstrate that MCP-1 is crucial

for recruiting neural stem cells after stroke, which is necessary for better neurological

outcomes [77]. Along these lines, we also found that cortistatin reduced the elevated

levels of *Arg1*, which is consistently upregulated in stroke patients [32], and known to

induce lymphopenia in a murine model of stroke [78]. Indeed, Arg1 inhibition

ameliorates stroke outcomes [79] and it is already been used in clinical trials in patients

with coronary artery disease who have suffered from ischemia-reperfusion [80].

In summary, with timely administration, the neuroimmune-modulatory role of

cortistatin in stroke extends beyond its direct neuroprotective effects on brain damage,

glial responses, and BBB modulation, to include the regulation of systemic immune

responses and to positively affect detrimental behavioural and functional outcomes.

This multifaceted mechanism of action offers several advantages over therapies

targeting single aspects of stroke pathophysiology. Furthermore, the intriguing

possibility that this neuropeptide has a role in the main players in this condition (such

as neurons, astrocytes, microglia, BECs, pericytes, or immune cells) could confer an

advantage over other potential therapies. Notably, the physiological nature—and thus

intrinsically non-toxic properties—of this neuropeptide provide assurances of its

potential use. In fact, it has been successfully used at clinical levels, both systemically

and locally, demonstrating its therapeutic efficacy in patients with acromegaly,

prolactinoma [81], and Cushing's disease [82]. Being beyond the scope of this work,

further research should consider sex as an independent variable that must be

addressed to properly interpret the beneficial action of cortistatin. Future studies

should also focus on extending the investigation of the spatiotemporal dynamics of

cortistatin action, the molecular pathways involved, and its effects on other cell types

within the CNS and the peripheral immune system.

CONCLUSION

Our data underscore the importance of comprehending the spatio-temporal dynamics of ischemic stroke when devising therapeutic interventions to optimize stroke outcomes. The findings indicating that a too-potent anti-inflammatory effect of the neuropeptide is deleterious when given immediately after stroke, emphasize the need to shift therapeutic goals from halting specific processes to modulating and balancing them, thereby mitigating possible chronic adverse responses. We showed that cortistatin administered at later stages appears to be a very promising candidate drug to enhance recovery after stroke, alone or in combination with other treatments, mainly targeting the outcome of patients that underwent recanalization or spontaneous reperfusion. Our study further highlighted the multifaceted roles of cortistatin in stroke pathophysiology, emphasizing its potential as a therapeutic agent. Its capacity to simultaneously act on various pathological features, such as neuronal damage, glial over-reaction, astrocytic scar formation, and BBB breakdown with an overall amelioration of the neurological outcome, offers significant advantage over interventions targeting single aspects of this condition. Moreover, cortistatin stands as a key factor regulating the complex interplay between the CNS and the immune system, modulating exacerbated molecular and cellular responses from both systems that may affect the brain's recovery processes. Finally, the valuable insights gained from this research have the potential to advance the development of novel therapeutic approaches, not only targeting stroke pathology but also other CNS disorders linked to glial over-reactivity, BBB disruption, or dysfunctional CNS-immune system interactions. Future research should continue to explore the complex interplay between this neuropeptide, its receptors, and the multitarget effects in stroke to develop more effective therapeutic strategies.

ACKNOWLEDGEMENTS

The authors would like to thank the Animal Care Unit and Microscopy Service Unit

staff from IPBLN for the technical support.

FUNDING

This work was supported by the Spanish Ministry of Science and Innovation

(MCIN)/AEI/ 10.13039/501100011033 and by "ERDF A way of making Europe" grants:

SAF2017-85602-R and PID2020-119638RB-I00 (to E.G-R.), by the Swiss Science

Foundation grant N°310030_212233 and Biaggi Foundation grant (to L.H), and by

FPU-program FPU17/02616 and EMBO-STFL8942 (to J.C-G.), FPI-program

PRE2018-084824 (to I.S-M.) and PRE2021-100172 (to P.V-R.).

CONTRIBUTIONS

E.G-R conceived, designed and supervised the whole study. J.C-G and E.G-R

outlined the experiments and analysed all the data. J.C-G, L.B, P.V-R, I.S-M, and I.F-

L performed the experiments. J.C-G, L.B, M.P, P.H-C, L.H, and E.G-R interpreted the

data, and discussed the results. J.C-G and E.G-R wrote and drafted the manuscript.

J.C-G, E.G-R, L.B, M.P, and L.H revised the manuscript. All authors have read and

agreed to the published version of the manuscript.

ETHICS DECLARATION

Ethical approval

All procedures were approved by the Animal Care and Use Board and the Ethical

Committee of the Spanish National Research Council (Spain), and the Vaud Cantonal

Veterinary Office (Switzerland). All experimental protocols adhered to the guidelines

specified in Directive 2010/63/EU of the European Parliament and the corresponding

Swiss regulations regarding animal protection for scientific purposes.

Competing interests

The authors declare no competing interests.

DATA AVAILABILITY

All data used or analysed will be made available upon reasonable request to the

corresponding author.

ABBREVIATIONS

Arg1: arginase 1

BBB: blood-brain barrier

BDNF: brain-derived neurotrophic factor

BECs: brain endothelial cells

BL: baseline

CBF: cerebral blood flow

Cl: claudin

CD: cluster of differentiation

CL: contralateral

CNS: central nervous system

CST: cortistatin

CV: cresyl violet

DAMPs: danger-associated molecular pattern molecules

GDNF: glial cell-derived neurotrophic factor

GFAP: glial fibrillary acidic protein

Iba1: ionized calcium-binding adaptor molecule 1

IgG: immunoglobulin G

IL: Interleukin

MAP-2: microtubule-associated protein 2

MBP: myelin basic protein

MCAO: middle cerebral artery occlusion

MCP-1: monocyte chemoattractant protein-1

PFA: paraformaldehyde

Rplp0: ribosomal protein lateral stalk subunit P0

rtPA: recombinant tissue plasminogen activator

TJ: tight-junctions

942   TNF: tumour necrosis factor- α

943   Trem2: triggering receptor expressed on myeloid cells 2

944   ZO-1: zonula occludens 1

BIBLIOGRAPHY

1. Lapchak PA, Yang G-Y, editors. Translational Research in Stroke [Internet]. Singapore:
Springer Singapore; 2017. Available from: http://link.springer.com/10.1007/978-981-10-5804-
2

2. Campbell BCV, De Silva DA, Macleod MR, Coutts SB, Schwamm LH, Davis SM, et al.
Ischaemic stroke. Nat Rev Dis Primers. 2019;5:1–22.

3. Drieu A, Levard D, Vivien D, Rubio M. Anti-inflammatory treatments for stroke: from bench
to bedside. Ther Adv Neurol Disord. 2018;11:1756286418789854.

4. de Lecea L, Criado JR, Prospero-Garcia O, Gautvik KM, Schweitzer P, Danielson PE, et al.
A cortical neuropeptide with neuronal depressant and sleep-modulating properties. Nature.
1996;381:242–5.

5. Delgado M, Gonzalez-Rey E. Role of Cortistatin in the Stressed Immune System. In: Savino
W, Guaraldi F, editors. Frontiers of Hormone Research. S. Karger AG; 2017. p. 110–20.
Available from: https://www.karger.com/Article/FullText/452910

6. Córdoba-Chacón J, Gahete MD, Pozo-Salas AI, Martínez-Fuentes AJ, de Lecea L, Gracia-
Navarro F, et al. Cortistatin is not a somatostatin analogue but stimulates prolactin release
and inhibits GH and ACTH in a gender-dependent fashion: potential role of ghrelin.
Endocrinology. 2011;152:4800–12.

7. Deghenghi R, Papotti M, Ghigo E, Muccioli G. Cortistatin, but not somatostatin, binds to
growth hormone secretagogue (GHS) receptors of human pituitary gland. J Endocrinol Invest.
2001;24:RC1-3.

8. Souza-Moreira L, Morell M, Delgado-Maroto V, Pedreño M, Martinez-Escudero L, Caro M,
et al. Paradoxical Effect of Cortistatin Treatment and Its Deficiency on Experimental
Autoimmune Encephalomyelitis. J Immunol. 2013;191:2144–54.

9. Chiu C-T, Wen L-L, Pao H-P, Wang J-Y. Cortistatin Is Induced in Brain Tissue and Exerts
Neuroprotection in a Rat Model of Bacterial Meningoencephalitis. J Infect Dis.
2011;204:1563–72.

10. Braun H, Schulz S, Becker A, Schröder H, Höllt V. Protective effects of cortistatin (CST-
14) against kainate-induced neurotoxicity in rat brain. Brain Res. 1998;803:54–60.

11. Falo CP, Benitez R, Caro M, Morell M, Forte-Lago I, Hernandez-Cortes P, et al. The
Neuropeptide Cortistatin Alleviates Neuropathic Pain in Experimental Models of Peripheral
Nerve Injury. Pharmaceutics. 2021;13:947.

12. Serrano-Martínez I, Pedreño M, Castillo-González J, Ferraz-de-Paula V, Vargas-
Rodríguez P, Forte-Lago I, et al. Cortistatin as a Novel Multimodal Therapy for the Treatment
of Parkinson's Disease. Int J Mol Sci. 2024;25:694.

13. Castillo-González J, Ruiz JL, Serrano-Martínez I, Forte-Lago I, Ubago-Rodriguez A, Caro
M, et al. Cortistatin deficiency reveals a dysfunctional brain endothelium with impaired gene
pathways, exacerbated immune activation, and disrupted barrier integrity. J
Neuroinflammation. 2023;20:226.

14. Buscemi L, Price M, Bezzi P, Hirt L. Spatio-temporal overview of neuroinflammation in an
experimental mouse stroke model. Sci Rep. 2019;9:507.

15. Buscemi L, Price M, Castillo-González J, Chatton J-Y, Hirt L. Lactate Neuroprotection
against Transient Ischemic Brain Injury in Mice Appears Independent of HCAR1 Activation.
Metabolites. 2022;12:465.

16. Lee S, Hong Y, Park S, Lee S-R, Chang K-T, Hong Y. Comparison of surgical methods of
transient middle cerebral artery occlusion between rats and mice. J Vet Med Sci.
2014;76:1555–61.

17. Balkaya M, Kröber JM, Rex A, Endres M. Assessing post-stroke behavior in mouse models
of focal ischemia. J Cereb Blood Flow Metab. 2013;33:330–8.

18. Fernández-Arjona M del M, Grondona JM, Granados-Durán P, Fernández-Llebrez P,
López-Ávalos MD. Microglia Morphological Categorization in a Rat Model of
Neuroinflammation by Hierarchical Cluster and Principal Components Analysis. Front Cell
Neurosci. 2017;11:235.

19. Young K, Morrison H. Quantifying Microglia Morphology from Photomicrographs of
Immunohistochemistry Prepared Tissue Using ImageJ. J Vis Exp. 2018;57648.

20. Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, et al. Fiji: an
open-source platform for biological-image analysis. Nat Methods. 2012;9:676–82.

21. Rousselet E, Kriz J, Seidah NG. Mouse Model of Intraluminal MCAO: Cerebral Infarct
Evaluation by Cresyl Violet Staining. J Vis Exp. 2012;e4038.

22. Türeyen K, Vemuganti R, Sailor KA, Dempsey RJ. Infarct volume quantification in mouse
focal cerebral ischemia: a comparison of triphenyltetrazolium chloride and cresyl violet
staining techniques. J Neurosci Methods. 2004;139:203–7.

23. Qin C, Zhou L-Q, Ma X-T, Hu Z-W, Yang S, Chen M, et al. Dual Functions of Microglia in
Ischemic Stroke. Neurosci Bull. 2019;35:921–33.

24. Karperien A, Ahammer H, Jelinek H. Quantitating the subtleties of microglial morphology
with fractal analysis. Front Cell Neurosci 2013;7.

25. Peng L, Hu G, Yao Q, Wu J, He Z, Law BY-K, et al. Microglia autophagy in ischemic stroke:
A double-edged sword. Front Immunol. 2022;13.

26. Linnerbauer M, Rothhammer V. Protective Functions of Reactive Astrocytes Following
Central Nervous System Insult. Front Immunol. 2020;11.

27. Jurga AM, Paleczna M, Kadluczka J, Kuter KZ. Beyond the GFAP-Astrocyte Protein
Markers in the Brain. Biomolecules. 2021;11:1361.

28. Thal S, Luh C, Schaible E-V, Timaru-Kast R, Hedrich J, Luhmann H, et al. Volatile
Anesthetics Influence Blood-Brain Barrier Integrity by Modulation of Tight Junction Protein
Expression in Traumatic Brain Injury. PloS one. 2012;7:e50752.

29. Abdullahi W, Tripathi D, Ronaldson PT. Blood-brain barrier dysfunction in ischemic stroke:
targeting tight junctions and transporters for vascular protection. Am J Physiol Cell Physiol.
2018;315:C343–56.

30. Jiao H, Wang Z, Liu Y, Wang P, Xue Y. Specific Role of Tight Junction Proteins Claudin-
5, Occludin, and ZO-1 of the Blood–Brain Barrier in a Focal Cerebral Ischemic Insult. J Mol
Neurosci. 2011;44:130–9.

31. Rust R. Insights into the dual role of angiogenesis following stroke. J Cereb Blood Flow
Metab. 2020;40:1167–71.

32. Petrone AB, O'Connell GC, Regier MD, Chantler PD, Simpkins JW, Barr TL. The Role of
Arginase 1 in Post-Stroke Immunosuppression and Ischemic Stroke Severity. Transl Stroke
Res. 2016;7:103–10.

33. Li T, Zhao J, Gao H. Depletion of Arg1-Positive Microglia/Macrophages Exacerbates
Cerebral Ischemic Damage by Facilitating the Inflammatory Response. Int J Mol Sci.
2022;23:13055.

34. Iadecola C, Anrather J. The immunology of stroke: from mechanisms to translation. Nat
Med. 2011;17:796–808.

35. Iadecola C, Buckwalter MS, Anrather J. Immune responses to stroke: mechanisms,
modulation, and therapeutic potential. J Clin Invest. 2020;130:2777–88.

36. Westendorp WF, Dames C, Nederkoorn PJ, Meisel A. Immunodepression, Infections, and Functional Outcome in Ischemic Stroke. Stroke. 2022;53:1438–48.

37. Wanner IB, Anderson MA, Song B, Levine J, Fernandez A, Gray-Thompson Z, et al. Glial Scar Borders Are Formed by Newly Proliferated, Elongated Astrocytes That Interact to Corral Inflammatory and Fibrotic Cells via STAT3-Dependent Mechanisms after Spinal Cord Injury. J Neurosci. 2013;33:12870–86.

38. Szalay G, Martinecz B, Lénárt N, Környei Z, Orsolits B, Judák L, et al. Microglia protect against brain injury and their selective elimination dysregulates neuronal network activity after stroke. Nat Commun. 2016;7:11499.

39. Kharlamov A, LaVerde GC, Nemoto EM, Jungreis CA, Yushmanov VE, Jones SC, et al. MAP2 immunostaining in thick sections for early ischemic stroke infarct volume in non-human primate brain. J Neurosci Methods. 2009;182:205–10.

40. Duarte EP, Curcio M, Canzoniero LM, Duarte CB. Neuroprotection by GDNF in the ischemic brain. Growth Factors. 2012;30:242–57.

41. Endres M, Fan G, Hirt L, Fujii M, Matsushita K, Liu X, et al. Ischemic brain damage in mice after selectively modifying BDNF or NT4 gene expression. J Cereb Blood Flow Metab. 2000;20:139–44.

42. Rubio A, Sánchez-Mut JV, García E, Velasquez ZD, Oliver J, Esteller M, et al. Epigenetic control of somatostatin and cortistatin expression by β amyloid peptide. J Neurosci Res. 2012;90:13–20.

43. Fan P, Zhang, L., He, Z., Shao, P., Ding, L., Wang, G., Niu, W., et al. Reduced Severity of Outcome of Recurrent Ipsilateral Transient Cerebral Ischemia Compared with Contralateral Transient Cerebral Ischemia in Rats. J Stroke Cerebrovasc Dis. 2017;26, 2915–2925.

44. Ermine CM, Bivard A, Parsons MW, Baron J-C. The ischemic penumbra: From concept to reality. Int J Stroke. 2021;16:497–509.

45. Jia J, Yang L, Chen Y, Zheng L, Chen Y, Xu Y, et al. The Role of Microglial Phagocytosis in Ischemic Stroke. Front Immunol. 2022;12.

46. Paolicelli RC, Sierra A, Stevens B, Tremblay M-E, Aguzzi A, Ajami B, et al. Microglia states and nomenclature: A field at its crossroads. Neuron. 2022;110:3458–83.

47. Cai W, Liu H, Zhao J, Chen LY, Chen J, Lu Z, et al. Pericytes in Brain Injury and Repair After Ischemic Stroke. Transl Stroke Res. 2017;8:107–21.

48. Wang, H., Li, X., Wang, Q., Ma, J., Gao, X., Wang, M. TREM2, microglial and ischemic stroke. J Neuroimmunol. 2023;381.

49. Pérez Falo C. Papel del neuropéptido cortistatina en la fisiopatología de los procesos de des- y remielinización en el sistema nervioso [Internet]. Universidad de Granada; 2021. Available from: https://dialnet.unirioja.es/servlet/tesis?codigo=295627

50. Neher JJ, Emmrich JV, Fricker M, Mander PK, Théry C, Brown GC. Phagocytosis executes delayed neuronal death after focal brain ischemia. Pro Natl Acad Sci. 2013;110:E4098–107.

51. Jolivel V, Bicker F, Binamé F, Ploen R, Keller S, Gollan R, et al. Perivascular microglia promote blood vessel disintegration in the ischemic penumbra. Acta Neuropathol. 2015;129:279–95.

52. Xu X, Gao W, Li L, Hao J, Yang B, Wang T, et al. Annexin A1 protects against cerebral ischemia–reperfusion injury by modulating microglia/macrophage polarization via FPR2/ALX-dependent AMPK-mTOR pathway. J Neuroinflammation. 2021;18:119.

53. Li Q, Barres BA. Microglia and macrophages in brain homeostasis and disease. Nat Rev Immunol. 2018;18:225–42.

54. Wen Q, Ding Q, Wang J, Yin Y, Xu S, Ju Y, et al. Cortistatin-14 Exerts Neuroprotective Effect Against Microglial Activation, Blood-brain Barrier Disruption, and Cognitive Impairment in Sepsis-associated Encephalopathy. J Immunol Res. 2022;2022:3334145.

55. Han X, Liu Y-J, Liu B-W, Ma Z-L, Xia T-J, Gu X-P. TREM2 and CD163 Ameliorate
Microglia-Mediated Inflammatory Environment in the Aging Brain. J Mol Neurosci.
2022;72:1075–84.

56. Roy-O'Reilly M, Zhu L, Atadja L, Torres G, Aronowski J, McCullough L, et al. Soluble
CD163 in intracerebral hemorrhage: biomarker for perihematomal edema. Ann Clin Transl
Neurol. 2017;4:793–800.

57. Shen X-Y, Gao Z-K, Han Y, Yuan M, Guo Y-S, Bi X. Activation and Role of Astrocytes in
Ischemic Stroke. Front Cell Neurosci. 2021;15.

58. Burda JE, Sofroniew MV. Reactive Gliosis and the Multicellular Response to CNS Damage
and Disease. Neuron. 2014;81:229–48.

59. Zamanian JL, Xu L, Foo LC, Nouri N, Zhou L, Giffard RG, et al. Genomic Analysis of
Reactive Astrogliosis. J Neurosci. 2012;32:6391–410.

60. Liddelow SA, Barres BA. Reactive Astrocytes: Production, Function, and Therapeutic
Potential. Immunity. 2017;46:957–67.

61. Escartin C, Galea E, Lakatos A, O'Callaghan JP, Petzold GC, Serrano-Pozo A, et al.
Reactive astrocyte nomenclature, definitions, and future directions. Nat Neurosci.
2021;24:312–25.

62. Karantali E, Kazis D, Papavasileiou V, Prevezianou A, Chatzikonstantinou S, Petridis F,
et al. Serum BDNF Levels in Acute Stroke: A Systematic Review and Meta-Analysis. Medicina
(Kaunas). 2021;57:297.

63. Faulkner JR, Herrmann JE, Woo MJ, Tansey KE, Doan NB, Sofroniew MV. Reactive
astrocytes protect tissue and preserve function after spinal cord injury. J Neurosci.
2004;24:2143–55.

64. Candelario-Jalil E, Dijkhuizen RM, Magnus T. Neuroinflammation, Stroke, Blood-Brain
Barrier Dysfunction, and Imaging Modalities. Stroke. 2022;53:1473–86.

65. Castillo-González J, Gonzalez-Rey E. Beyond wrecking a wall: Revisiting the concept of
blood-brain barrier breakdown in ischemic stroke.Neural Regen Res. 2024;2.

66. Bernardo-Castro S, Sousa JA, Brás A, Cecília C, Rodrigues B, Almendra L, et al.
Pathophysiology of Blood–Brain Barrier Permeability Throughout the Different Stages of
Ischemic Stroke and Its Implication on Hemorrhagic Transformation and Recovery. Front
Neurol. 2020;11.

67. Ballabh P, Braun A, Nedergaard M. The blood–brain barrier: an overview: Structure,
regulation, and clinical implications. Neurobiol Dis. 2004;16:1–13.

68. Fischer S, Wobben M, Marti HH, Renz D, Schaper W. Hypoxia-induced hyperpermeability
in brain microvessel endothelial cells involves VEGF-mediated changes in the expression of
zonula occludens-1. Microvasc Res. 2002;63:70–80.

69. Sajja RK, Prasad S, Cucullo L. Impact of altered glycaemia on blood-brain barrier
endothelium: an in vitro study using the hCMEC/D3 cell line. Fluids Barriers CNS. 2014;11:8.

70. Famakin BM. The Immune Response to Acute Focal Cerebral Ischemia and Associated
Post-stroke Immunodepression: A Focused Review. Aging Dis. 2014;5:307–26.

71. Jayaraj RL, Azimullah S, Beiram R, Jalal FY, Rosenberg GA. Neuroinflammation: friend
and foe for ischemic stroke. J Neuroinflammation. 2019;16:142.

72. Grønhøj MH, Clausen BH, Fenger CD, Lambertsen KL, Finsen B. Beneficial potential of
intravenously administered IL-6 in improving outcome after murine experimental stroke. Brain,
Behav Immun. 2017;65:296–311.

73. Gertz K, Kronenberg G, Kälin RE, Baldinger T, Werner C, Balkaya M, et al. Essential role
of interleukin-6 in post-stroke angiogenesis. Brain. 2012;135:1964–80.

74. Figiel I. Pro-inflammatory cytokine TNF-alpha as a neuroprotective agent in the brain. Acta
Neurobiol Exp (Wars). 2008;68:526–34.

75. Gonzalez-Rey, E., Varela, N., Sheibanie, A.F., Chorny, A., Ganea, D., Delgado, M.,
Cortistatin, an antiinflammatory peptide with therapeutic action in inflammatory bowel
disease. Proc Natl Acad Sci U S A. 2006;103, 4228–4233.

76. Gonzalez-Rey E, Chorny A, Moral RGD, Varela N, Delgado M. Therapeutic effect of
cortistatin on experimental arthritis by downregulating inflammatory and Th1 responses. Ann
Rheum Dis. 2007;66:582–8.

77. Andres RH, Choi R, Pendharkar AV, Gaeta X, Wang N, Nathan JK, et al. The CCR2/CCL2
Interaction Mediates the Transendothelial Recruitment of Intravascularly Delivered Neural
Stem Cells to the Ischemic Brain. Stroke. 2011;42:2923–31.

78. Sippel TR, Shimizu T, Strnad F, Traystman RJ, Herson PS, Waziri A. Arginase I Release
from Activated Neutrophils Induces Peripheral Immunosuppression in a Murine Model of
Stroke. J Cereb Blood Flow Metab. 2015;35:1657–63.

79. Barakat W, Fahmy A, Askar M, El-Kannishy S. Effectiveness of arginase inhibitors against
experimentally induced stroke. Naunyn Schmiedebergs Arch Pharmacol. 2018;391:603–12.

80. Tratsiakovich Y, Yang J, Gonon AT, Sjöquist P-O, Pernow J. Arginase as a target for
treatment of myocardial ischemia-reperfusion injury. Eur J Pharmacol. 2013;720:121–3.

81. Giordano R, Picu A, Bonelli L, Broglio F, Prodam F, Grottoli S, et al. The activation of
somatostatinergic receptors by either somatostatin-14 or cortistatin-17 often inhibits ACTH
hypersecretion in patients with Cushing's disease. Eur J Endocrinol. 2007;157:393–8.

82. Grottoli S, Gasco V, Broglio F, Baldelli R, Ragazzoni F, Gallenca F, et al. Cortistatin-17
and Somatostatin-14 Display the Same Effects on Growth Hormone, Prolactin, and Insulin
Secretion in Patients with Acromegaly or Prolactinoma. J Clin Endocrinol Metab.
2006;91:1595–9.

1159
1160