Potent AMA1-specific human monoclonal antibody against P. vivax Pre-erythrocytic and Blood Stages

Anna C. Winnicki [1,+], Melanie H. Dietrich [6,8,+], Lee M. Yeoh [7,9], Lenore L. Carias [1], Wanlapa Roobsoong [5], Chiara L. Drago [7,12], Alyssa N. Malachin [1], Karli R. Redinger [1], Lionel Brice Feufack-Donfack [4], Lea Baldor [4], Nicolai C. Jung [6], Olivia S. McLaine [1], Yelenna Skomorovska-Prokvolit [1], Agnes Orban [4], D. Herbert Opi [7,10,11], Jetsumon Sattabongkot [5], Wai-Hong Tham [6,8], Jean Popovici [4], James G. Beeson [7,10,11], Jürgen Bosch*[1,3], and Christopher L. King*[1,2]

* Co-senior authors

+ Co-first authors

[1] Center for Global Health and Diseases, Department of Pathology, Case Western Reserve University School of Medicine, [2] Veterans Affairs Medical Center, Cleveland, OH, [3] InterRayBio LLC, [4] Malaria Research Unit, Institut Pasteur du Cambodge, Phnom Penh, Cambodia, [5] Mahidol Vivax Research Unit, Faculty of Tropical Medicine, Mahidol University, Bangkok, Thailand, [6] Walter and Eliza Hall Institute of Medical Research, Parkville, Victoria, Australia, [7] Burnet Institute, Melbourne, Victoria, Australia, [8] Department of Medical Biology, The University of Melbourne, Parkville, Victoria, Australia, [9] Department of Medicine, The University of Melbourne, Parkville, Victoria, Australia, [10] Department of Infectious Diseases, The University of Melbourne, Parkville, Victoria, Australia, [11] Central Clinical School and Department of Microbiology, Monash University, Clayton, Victoria, Australia. [12] Department of Microbiology, Monash University, Clayton, Victoria, Australia.

Abstract

New therapeutics are a priority for preventing and eliminating *Plasmodium vivax* (Pv) malaria because of its easy transmissibility and dormant stages in the liver. Relapses due to the dormant liver stages are the major contributor to reoccurring Pv. Therefore, therapies that reduce the establishment of dormant parasites and blood-stage infection are important for controlling this geographically widespread parasite. Here, we isolated 12 human monoclonal antibodies (humAbs) from the plasma of a Pv-exposed individual that recognized Pv apical membrane antigen 1 (PvAMA1). PvAMA1 is important for both sporozoite invasion of hepatocytes and merozoite invasion of reticulocytes. We identified one humAb, 826827, that blocked invasion of human erythrocytes using a transgenic *P. falciparum* line expressing PvAMA1 (IC_{50} = 3 µg/mL) and all *Pv* clinical isolates *in vitro*. This humAb also inhibited sporozoite invasion of a human hepatocyte cell line and primary human hepatocytes (IC_{50} of 0.3 – 3.7 µg/mL). The crystal structure of recombinant PvAMA1 with the antigen-binding fragment of 826827 at 2.4 Å resolution shows that the humAb partially occupies the highly conserved hydrophobic groove in PvAMA1 that binds its known receptor, RON2. HumAb 826827 binds to PvAMA1 with higher affinity than RON2, accounting for its potency. To our knowledge, this is the first reported humAb specific to PvAMA1, and the PvAMA1 residues it binds to are highly conserved across different isolates, explaining its strain-transcendent properties.

Word count: 224

Introduction

Half of the world's population is at risk of malaria, with 247 million cases and 619,000 deaths occurring in 2022 [1]. *Plasmodium falciparum* (Pf) and *Plasmodium vivax* (Pv) account for most human malaria cases. Pf predominates in sub-Saharan Africa, whereas Pv accounts for 80% of malaria in Asia and the Americas [2]. Pv causes significant disease, particularly in low-middle-income countries burdened with poor nutrition, anemia, and co-infections [3]. During the erythrocytic or blood stage, Pv merozoites infect reticulocytes, primarily at the sites of erythropoiesis in the bone marrow and spleen [4-6]. Pv forms a dormant pre-erythrocytic or liver phase (hypnozoites), causing frequent relapses, further contributing to anemia and other complications [7, 8]. The frequent relapses produce gametocytes that drive transmission in populations [9]. Presently, there is no vaccine available for Pv. Targeted therapies that inhibit hepatocyte and erythrocyte invasion are imperative in reducing the overall disease burden and enabling Pv elimination [10-13].

An important piece of the invasion machinery used by sporozoites and merozoites is the interaction between Apical Membrane Antigen 1 (AMA1) and an extracellular ß-hairpin loop in the C-terminal portion of Rhoptry Neck Protein 2 (RON2) [14]. Utilized by all members of the Apicomplexa, AMA1 originates in the parasite's micronemes and is later translocated to the parasite's membrane. In human hosts, AMA1 is expressed during the late merozoite and sporozoite stages [15-17]. Immediately preceding invasion, the RON complex (composed of RON2, 4, 5 and 8), is secreted from the parasite's rhoptries and embeds into the target cell's membrane. The extracellular ß-hairpin loop near the C-terminus of RON2 can then interact with AMA1 by nestling into the hydrophobic groove of Domain 1 of AMA1 [18]. For the RON2-loop to properly engage with AMA1, a mobile loop of Domain 2 of AMA1 is displaced by the incoming ligand to reveal the entire hydrophobic groove [16, 19]. This protein-protein interaction is important for forming a tight junction that allows the merozoite to move across the extracellular

space and into the erythrocyte through connections between the parasite surface and its myosin motor along actin filaments [7, 14, 15, 20-22]. Blocking the interaction of PfAMA1 or PkAMA1 with PfRON2 or PkRON2 by antibodies [14, 23-25] or peptides [14, 26, 27] inhibits invasion, confirming that the interaction of AMA1-RON2 is important for the *Plasmodium* life cycle. While blocking this interaction can prevent invasion, there is evidence that AMA1 has additional interactions directly with the surface of the erythrocyte [28] or with other receptors [29], which are yet to be defined.

Malaria-infected individuals acquire partial immunity to infection and disease primarily directed toward blood-stage parasites [30]. Antibodies play a key role in this partial immunity, as demonstrated by transferring human IgG from immune adults to non-immune children, resulting in protection from malaria [31]. The relative contribution of different malaria blood-stage antigens to producing Pv- and Pf-specific antibodies is yet to be fully understood. The antibodies may protect against malaria by one or more of the following mechanisms: 1) blocking merozoite invasion into erythrocytes, 2) complement activation, and 3) opsonic phagocytosis by monocytes and neutrophils [32, 33]. Elevated antibodies to specific *Plasmodium* antigens, such as AMA1, are associated with protection against infection and disease [31, 34, 35].

PvAMA1, a three-domain protein, is under strong immune selection, posing a problem for developing vaccines and therapeutic monoclonal antibodies to this antigen [33, 36-38]. Domains 1 and 2 contain two clusters of disulfide-bonded cysteines, with the crystal structure revealing regions comprised of many long loops. Extending from the core of Domain 1, these long loops allow for significant variation and protein flexibility. This aids parasite evasion of AMA1-specific protective human antibody responses [39]. The loops form a scaffold for the numerous polymorphisms on the surface of AMA1, primarily located on the one side [34, 40-44]. The hydrophobic RON2 binding groove is highly conserved across Pv clinical isolates (97.8% over 110 residues). However, this groove is surrounded by highly polymorphic residues, presumably due to selective pressure from host immune responses [45]. Since the

AMA1:RON2 interaction plays a role at multiple points of the parasite's life cycle, including a role for sporozoite infection of the liver and transmission [8, 15-17], it presents an opportunity for a multi-stage target.

Recently, the development of therapeutic monoclonal antibodies to sporozoite antigens has shown significant clinical efficacy against malaria in endemic populations [46]. Human monoclonals to Pf circumsporozoite protein (CSP) have been isolated from individuals following immunization with attenuated sporozoite. These humAbs have been shown to protect against infection in human clinical trials [47, 48] by reducing the risk of Pf infection by 88% in adults living in malaria-endemic areas of Africa [19]. These studies show the promise of using human humAbs specific to sporozoite antigens to protect against malaria infection and disease. Previously, a PfAMA1-specific human-derived monoclonal antibody was isolated, characterized, and showed blocking activity at an IC_{50} of 35 μg/mL [49] to blood-stage parasites *in vitro*. To our knowledge, no human-derived monoclonal antibody specific to PvAMA1 has been documented.

Here, we have isolated 12 PvAMA1-specific human-derived monoclonal antibodies (humAbs) from the plasma of a Cambodian individual with prior Pv infection(s) whose antibodies inhibit PvAMA1 binding to PvRON2. The biophysical properties of these 12 humAbs were characterized. We then tested the growth and invasion inhibition properties of the humAbs against merozoite and sporozoite invasion *in vitro*. In summary, we have identified a potent, strain-transcending humAb that blocks RON2-loop binding and can inhibit both blood stage and sporozoite infection. Our structural data shows that this humAb occupies the RON2 hydrophobic binding groove and recognizes conserved amino acids in its epitope.

Results

Isolation and expression of human monoclonal antibodies

We tested plasma from seven malaria-exposed Cambodian adults for antibodies that could inhibit the binding of PvRON2 to PvAMA1 [29]. Using the donor with the highest RON2 binding inhibition activity (**Supplemental Figure 1**), we isolated 169 PvAMA1-specific B cells that were PCR-amplified and sequenced for the IgG heavy chain V-D-J region. Based on a high sequence identity of the V-D-J region, the B cells were placed into 67 clonal groups from which we generated 12 human PvAMA1-specific monoclonal antibodies that represent 12 clonal groups. The clones selected for the generation of humAbs were based on whether there was a corresponding IgG light chain, the quality of sequence, the degree of somatic hypermutations (SHM), and the selection of one B cell H+L pair from the clonal group (**Figure 1A,B**).

To determine the specificity of 12 PvAMA1 humAbs, we tested the binding capabilities to recombinant AMA1 corresponding to two strains of PvAMA1 (Palo Alto the variant used to sort B cells, and PvAMA1_PNG16, a sequence with significant polymorphisms compared to Palo Alto), PkAMA1, PfAMA1_3D7, and TgAMA1 (**Figure 1C**). Two different published versions of PvAMA1 were used to represent the variance in protein sequence across Pv strains. The sequence identity of the other AMA1 constructs with respect to PvAMA1_Palo Alto was 97.2%, 85.6%, 60.3%, and 30.3% for PvAMA1_PNG16, PkAMA1, PfAMA1_3D7, and TgAMA1 respectively. All humAbs recognized PvAMA1_Palo Alto, ten recognized PvAMA1_PNG16, seven recognized PkAMA1, two recognized PfAMA1_3D7, and none recognized TgAMA1 in a multiplex immunoassay [50]. As expected, our isotype control humAb 043038, specific to tetanus toxoid C-terminal fragment, did not bind to any of these AMA1 recombinant proteins (**Figure 1C**). The humAb titers recognizing the different recombinant AMA1 constructs varied among the different humAbs and were associated with differences in humAb avidity (**Figure 1C,D**). The avidity to PvAMA1 was highest for humAbs 804805, 826827, 832833, and 838839 (**Supplemental Table 1**). It is of note that humAb 826827 has a particularly high avidity to PkAMA1. The calculated Avidity Index 50 (AI_{50}) represents the molar concentration of the chaotropic reagent where 50% of the binding of the humAb to AMA1 is lost (**Supplemental**

Table 1). To further characterize the biophysical properties of the humAbs, we measured the affinity to PvAMA1_Palo Alto using Surface Plasmon Resonance (SPR). Affinities of humAbs to PvAMA1 ranged from 10.7×10^{-9} to 47.7×10^{-9} M (**Figure 1E**, **Supplemental Figure 2**). Three humAbs, 800801, 804805 and 808809 did not have significant PvAMA1 binding in this assay. Thus, we isolated PvAMA1-specific humAbs with various biophysical characteristics that may have further therapeutic potential.

HumAb inhibition of Pf-PvAMA1 Transgenic Parasites

Because long-term *in vitro* culture of *P. vivax* is not currently possible, we used a Pf parasite line engineered to express PvAMA1 [28, 29, 51] as a model to evaluate merozoite invasion-inhibitory activity of the humAbs. Five humAbs, 808809, 826827, 828829, and 830831 displayed IC_{50}s between 2.6–11.5 µg/mL, with 826827 being the most potent. HumAb 043038 was used as a negative control and displayed no invasion inhibition at the concentrations tested **(Figure 2A, Supplemental Figure 3**).

HumAb inhibition of Pv clinical isolates from Cambodia

We performed *in vitro* reticulocyte invasion assays to evaluate the susceptibility of different Pv clinical isolates to anti-PvAMA1 humAbs (**Figure 2B)**. 826827 showed the highest percent inhibition with a mean of 67.4% (±8.6% SEM, N=7), while 043038 showed inhibition of 4.72% (±3.8% SEM, one-way ANOVA, *P* = <0.0001) (**Figure 2B**). HumAb 816817 had the second highest average inhibition at 41.1% (±11.4% SEM, one-way ANOVA compared to 043038, *P* = 0.4449). We performed dose response invasion assays using humAb 826827 from 7.8 to 1000 µg/mL against four additional Pv clinical isolates to estimate the IC_{50}. The average IC_{50} obtained using four clinical isolates was 48 µg/mL (± 6.6 SEM) (**Figure 2C**). This data, combined with the *in vitro* Pf-PvAMA1 transgenic parasite experiments results, indicates that human 826827 is highly effective at inhibiting AMA1-dependent erythrocyte invasion.

HumAb inhibition of Pv sporozoites in human hepatocytes

We examined humAb ability to inhibit Pv sporozoite invasion of hepatocytes *in vitro* (**Figure 3**). Sporozoites were isolated from the salivary glands of *Anopheles dirus* mosquitoes fed on Pv-infected subjects from Thailand, and humAbs were tested for their ability to inhibit sporozoite invasion of HC04 hepatocytes. Five humAbs displayed IC_{50}s (0.38–2.6 µg/mL) that were comparable to the positive control anti-CSP murine monoclonal 2F2 (IC_{50} CSP210 = 0.2 µg/mL, **Figure 3A, Supplemental Figure 4,** [52]). For three biological replicates of humAb 826827 (**Figure 3B),** IC_{50}s ranged from less than 0.01 to 1.8 µg/mL. We also performed the *Pv* sporozoite invasion assay on a Cambodian isolate using primary human hepatocytes and the humAb 826827 (**Figure 3C**). In this experiment, the IC_{50} was 3.7 µg/mL. Thus, humAb 826827 can inhibit AMA1-dependent sporozoite invasion from multiple clinical isolates.

Structural studies defining humAb 826827 interaction with PvAMA1

We obtained a 2.4 Å resolution structure of humAb 826827 bound to the PvAMA1 ectodomain with an R_{work}/R_{free} of 19.2% and 23.5% (**Supplemental Table 3**). A 1:1 complex is present in the asymmetric unit (**Figure 4A**). The resulting electron density map allowed the tracing of PvAMA1 from residues 46 to 474 with main chain gaps at residues 211–215, 296–303, 328–334, and 402–415. Heavy and light chains of the antigen-binding fragment of 826827 were fully traced from 1–231 and 1–214, respectively. Our structure shows that 826827 interacts with PvAMA1 residues of Domain 1 and the mobile loop of Domain 2 (**Figure 4A, panel i and ii**). Five of the six complementarity-determining regions (CDR, namely: L1, L2, H1, H2, and H3) form direct contacts with PvAMA1 with a buried interaction surface of 1392 $Å^2$, with the light chain contributing 432 $Å^2$ and the heavy chain contributing 959 $Å^2$ (**Figure 4B**). The CDR-H3 loop of 826827 forms a disulfide bridged β-hairpin that binds to the hydrophobic groove on PvAMA1 Domain 1, which constitutes part of the RON2-loop receptor binding site (**Figure

4A-C). CDR-H3 binding to the PvAMA1-hydrophobic groove involves 53 interatomic contacts

with distances < 3.8 Å of which six are hydrogen bonds (**Supplemental Table 4**). The mobile

Domain 2 loop of PvAMA1 is contacted by residues of CDR-H3 and CDR-H1, forming one salt

bridge and six hydrogen bonds that stabilize its position on Domain 1 (**Figure 4**). CDR-L2 and

H2 interact with PvAMA1 residues of Domain 1 loops that surround the mobile Domain 2 loop,

and CDR-L1 forms contacts with a Domain 1 loop next to the hydrophobic groove (**Figure

4**). Two residues located in the Domain 2 loop of AMA1, Arg317 and Lys321, provide a

positively charged patch at the bottom of the RON2-loop binding groove. Otherwise, this binding

pocket is largely hydrophobic (**Supplemental Figure 5**). One salt bridge between PvAMA1

Lys321 and humAb 826827 CDR-H3 Glu103 provides an anchoring and orientation point for the

observed interaction between the two proteins. The remainder of the interactions are

hydrophobic and hydrogen bonding interactions in nature (**Figure 4i, Supplemental Movie 1,

Supplemental Movie 2**).

Compared to structures of unbound PvAMA1 (PDB ID: 1W8K, [42]) and the PvAMA1

bound to a peptide representing the ß-hairpin loop of RON2 (PDB ID 5NQG, [26]), our PvAMA1

structure overlays with low root mean square (r.m.s.) deviation values of 0.463 Å (over 2137

atoms) and 0.417 Å (over 2121 atoms) respectively (**Figure 4D,E**). A major difference exists in

our structure in which the mobile Domain 2 loop with residues 304–327 is visible while it is

unstructured in the other PvAMA1 structures. Structural analysis with *Plasmodium* AMA1

homologues showed that the Domain 2 loop of our structure adopts a similar 'closed'

conformation as in unbound PkAMA1 (PDB ID: 4UV6, [43]) which share 90% sequence identity

in this loop (**Figure 4F,G**). While Akter et al. [53] observed an open or semi-open conformation

in their PfAMA1 structure complex with a cyclized RON2 peptide (**Supplemental Movie 3**), our

crystal structure clearly shows that in the presence of 826827, the Domain 2 loop of PvAMA1 is

stabilized and remains bound to Domain 1, effectively blocking the RON2 binding site. The

displacement of the Domain 2 loop is required to expose the complete RON2 binding site

(**Figure 4F,G**). Our proposed model is in line with molecular dynamics simulations performed on PfAMA1-PfRON2 and TgAMA1-TgRON2 complexes [54, 55]. Our co-crystallization structure confirms the humAb 826827 binds to an important region of PvAMA1, providing a mechanistic explanation for its high potency.

HumAb 826827 binds in a highly conserved binding pocket of PvAMA1

Previous co-crystal structures of Pf- and PvAMA1 reveal a hydrophobic binding pocket on AMA1 that interacts with the extracellular RON2-loop [23, 26, 45, 56]. Visually, our co-crystallization structure of humAb 826827 with PvAMA1 indicates that this antibody may interact with the same PvAMA1 residues as the RON2-loop. **Figure 5A** shows that the contact residues between PvAMA1 and RON2-loop overlap with the residues of PvAMA1 that contact humAb 826827 CDR-H3 residues. The combined buried surface area is 1392.5 $Å^2$ large, while the major contribution of 70% is provided by the heavy chain 826. Sequence and structural analysis of the contact residues indicate that humAb 826827's epitope is highly conserved with respect to PkAMA1 and PcAMA1. However, little epitope conservation is observed in PfAMA1, accounting for the lack of humAb recognition of the recombinant protein (**Figure 1C, 5A; Supplemental Figure 6**). Comparing the RON2 and 826 CDR3 binding site to other available *Plasmodium* species and model systems reveals that *P. cynomolgi* AMA1 is 100% conserved and would therefore serve as a predictive non-human primate model for Pv challenge infections to evaluate humAb 826827 (**Figure 5A, Supplemental Figure 6**).

HumAb 826827 competes for the same epitope on PvAMA1 as PvRON2

To confirm that the binding of humAb 826827 to PvAMA1 inhibits the extracellular RON2-loop from binding, we performed a dose-response competition assay between the two proteins. We found that humAb 826827 at a concentration as low as 2.5 μg/mL inhibits the binding of the biotinylated PvRON2-loop peptide to PvAMA1 (**Figure 5B**). This result demonstrates that humAb

826827 competes for the same epitope as the RON2-loop, further confirming our proposed mechanism of humAb 826827-induced invasion inhibition.

Sequence conservation across 390 clinical PvAMA1 isolates

To evaluate how sequence conservation and polymorphisms in PvAMA1 may impact humAb efficacy, we examined 390 published PvAMA1 sequences from clinical isolates. We found 98% conservation of PvAMA1's total amino acid sequence across the 484 residues. Domain 1 is the largest and most polymorphic of the domains of PvAMA1 [50]. Domain 1 (1–248) contains 21 polymorphic amino acid residues, Domain 2 (248–385) contains 6, and Domain 3 (386–484) contains 3. All polymorphic residues are 44–80% conserved across 390 PvAMA1 sequences (**Supplemental Figure 7A,** [50]).

There is 97% conservation of PvAMA1 residues that contact the CDR3 loop of 826827 (**Figure 6A**). We evaluated whether the PvAMA1 polymorphisms of the Pv clinical isolates used in the invasion assays affected the potency of 826827. We classified the invasion data by assessing the impact of the polymorphisms of AMA1 within the epitope recognized by humAb 826827 (**Supplemental Table 4**). None of the 7 isolates tested against 826827 (**Figure 2B**) shared the same haplotype with the wild type (WT) reference Sal1 strain (PVX_092275) and had unique amino acid sequences. Among the 7 isolates, 4 were WT for the residues recognized by 826827; 2 had two mutations, and 1 had three mutations. All mutations were observed in residues 117, 130, and 132. As predicted, these polymorphisms did not impact the invasion inhibition by 826827 in terms of number of mutations compared to the WT or when considering individually each mutation compared to the WT, supporting our structural interpretation of the interaction (**Figure 6B,C; Supplemental Table 2,4**).

Structural analysis of the two residues N130 (N 86.4%, K 13.9%) and N132 (D 52.9%, N 46.7%, G 0.5%) with the highest variability between PvAMA1 sequences are unlikely to disrupt humAb 826827 binding significantly as the interactions are either maintained (backbone

interactions) or do not directly lie in the groove of the binding pocket (**Figure 6B**). An additional

mutation within hydrogen bonding distance in the proximity of the CDR3 loop is G117 (G 99.0%,

R 1%) which also does not likely alter the interaction as it is pointing towards the solvent and is

not interacting with other residues on the protein (**Figure 6B; Supplemental Movie 2**). These

data indicate that humAb 826827 may be strain transcendent.

Discussion

We have isolated a PvAMA1-specific humAb 826827 that inhibits sporozoite invasion of

hepatocytes and merozoite invasion of reticulocytes *in vitro*. This is the first reported humAb that

recognizes PvAMA1 and inhibits invasion of pre-erythrocytic and blood-stage parasites. HumAb

826827 is potent and strain-transcending. The dual activity against sporozoites and merozoites

is especially valuable because Pv disease is driven by new infections and relapses from

dormant liver hypnozoites [57]. Therefore, blocking both sporozoites and merozoites might

prevent infection, further transmission, and disease from reactivation of hypnozoites.

HumAb 826827 was the most potent humAb that inhibited merozoite invasion into

erythrocytes using a Pf transgenic parasite line expressing PvAMA1 (IC_{50} = 3.0 μg/mL) [51] and

in short-term invasion experiments into reticulocytes using Pv clinical isolates (IC_{50} = 48

μg/mL). Three additional humAbs 808809, 828829, and 830831 inhibited Pf invasion of

PvAMA1-expressing transgenic parasites (IC_{50} = 5.9–8.2 μg/mL) and appear to have

overlapping epitopes with 826827 (*data not shown*). However, only humAb 826827 showed

substantial and significant inhibition of Pv clinical isolates. The difference in results between the

two experiments may be because of the strain-transcendent properties of humAb 826827,

whereas other growth-inhibiting humAbs may target epitopes that include more polymorphic

residues that differ among the clinical isolates from the PvAMA1_Palo Alto. The transgenic

parasites express PvAMA1_Palo Alto, which was the strain used to sort B cells, which may

explain the increased efficacy of humAbs other than 826827 in these experiments [51]. These humAbs are more potent than previously reported PfAMA1-specific murine mAbs IF9 and 4G2, which display IC_{50}s of 292 and 105 µg/mL respectively against the Pf WT strain 3D7 in blood stage invasion studies [24, 45, 49, 58, 59] and rat mAb R31C2 [43, 56]. They are also exhibit similar potency compared to a PfAMA1 humAb that was previously produced based on an IgG sequence isolated from a Ghanaian individual with an IC_{50} of 35 µg/mL against a single Pf 3D7 variant *in vitro* [49]. Of note, a single-component PfAMA1-RON2L immunogen was developed as a vaccine candidate to produce an antibody response to complexed PfAMA1-RON2. Although the elicited polyclonal antibodies displayed Pf strain-transcending properties like our humAb 826827, they had poor potency with IC_{50}s ranging from 1.5 to 4.5 mg/mL against Pf 3D7 [60].

HumAb 826827 also blocked sporozoite invasion into human hepatocytes, supporting recent studies that show AMA1 is utilized during sporozoite penetration of hepatocytes and entry into mosquito salivary glands [16, 17]. It has been previously demonstrated that mouse polyclonal antibodies generated to either PfAMA1 or PvAMA1 inhibit blood sporozoite invasion into human hepatocytes but require a concentration of 0.5 to 1 mg/mL [17]. By contrast, our humAb 826827 has an IC_{50} of 0.3 µg/mL when preventing sporozoite invasion of human hepatocyte cell line HC04 and an IC_{50} of 3.7 µg/mL in primary human hepatocytes. Using different clinical isolates across the different assays and observing little variation in inhibition suggests that humAb 826827 is strain-transcendent with high potency. Indeed, the potency of humAb 826827 was comparable to the murine anti-circumsporozoite protein mAb 2F2 [52], which was used as a positive control in these experiments. This is notable as passive transfusion of a suboptimal dose of mAb 2F2 in liver-humanized mice, that were then challenged with Pv sporozoites, reduced parasite relapse by 62%, which was associated with a corresponding reduction of hypnozoite numbers [12]. This suggests that humAb 826827, when

used in that same model, could achieve a similar or greater reduction in hypnozoites though this would require future testing. Of note, we validated humAb 826827 ability to inhibit sporozoite invasion into primary human hepatocytes, with a model that yields much higher infection rates than HC04 cell line and allows for the development of both hypnozoites and schizonts [61].

The difference in humAb 826827 concentrations required to inhibit sporozoites (IC_{50} = 0.3–3.7 µg/mL) in hepatocytes and merozoite invasion (IC_{50} = 48 µg/mL) may be attributed to variations in levels of antigen availability or the nature of molecular interactions that occur during invasion. Parasite load, and thus, the number of PvAMA1 molecules expressed, differs significantly between the different *in vitro* assays. While 1×10^5 sporozoites are used in the liver invasion assay, the number of merozoites present during the clinical isolate blood stage assay is more than ten-fold higher. Furthermore, the structural conformation of different epitopes displayed may vary between pre-erythrocytic and erythrocytic stages of invasion. However, the combination of our pre- and erythrocytic invasion inhibition data confirms that PvAMA1 plays an important role during sporozoite invasion into hepatocytes and merozoite invasion into reticulocytes. This confirms that AMA1 is a viable multi-stage therapeutic target against *Plasmodium* infection and requires further research.

The crystal structure of recombinant PvAMA1_Palo Alto bound to humAb 826827 reveals why this humAb is potent and strain transcendent. The CDR3 of the heavy chain (826) recognizes a conformational epitope that overlaps with the RON2-loop binding site in Domain 1 and displays a higher affinity for PvAMA1 than PvRON2, 30.5 nM and 50.0 nM respectively. Previous structural studies suggested that a mobile loop of PvAMA1 Domain 2 partially obstructs the RON2 binding grove. This mobile loop must be displaced for a successful PvAMA1:RON2 interaction to occur [18, 43, 44]. HumAb 826827 binds the Domain 2 loop, thus preventing displacement and further interfering with PvRON2 engagement of PvAMA1. The PvAMA1 contact residues that directly interact with humAb 826827 are conserved or possess

single nucleotide polymorphisms (SNP) that do not affect humAb potency. For example, the SNP (N132D) frequently occurs in Pv clinical isolates. The Pv clinical isolates data indicates this mutation does not impair the hydrogen bonding capabilities between PvAMA1 and humAb 826827 (**Figure 6C**). Similarly, D133N maintains hydrogen bonding capabilities and does not impact the IC_{50} of 826827 (**Figure 6C**). The mutation N130K does change the interaction between PvAMA1 and 826827. However, the sidechains point towards the solvent, and only the backbone of the residue interacts with the light chain of 826827. As expected, there is no variation in humAb activity (**Figure 6C**). Of note, 826827 displays a much higher avidity to PkAMA1 than to PvAMA1_Palo Alto. Two amino acid changes (M153L & M171I) occur in 826827's binding epitope from the PvAMA1 to PkAMA1 amino acid sequences. This variation in amino acid sequence creates a more hydrophobic and presumably stronger interaction between 826827 and PkAMA1, likely leading to the observed increase in avidity.

In conclusion, we have identified a highly conserved epitope of PvAMA1 that can be targeted by humAb 826827 which inhibits AMA1-dependent sporozoite invasion into hepatocytes and merozoite invasion into reticulocytes. HumAb 826827 could be translated into a new therapeutic to prevent Pv infection, disease and transmission. Furthermore, identifying a conserved inhibitory epitope of PvAMA1 that is effectively targeted by 826827 could inform structure-based vaccine design for Pv. Future studies using related non-human primate models would provide valuable data to inform the development of humAb 826827 as a therapeutic/prophylaxis.

Methods

Blood samples

Peripheral blood mononuclear cells (PBMC) were obtained from Cambodians with documented Pv malaria residing in Pursat Province, as described previously [49]. Samples were

screened for blocking antibodies to PvAMA1 (see below). Institutional review boards from the National Institutes of Health (National Institute of Allergy and Infectious Diseases protocol 08-N094, clinicaltrials.gov identifier NCT00663546), Cambodian Ministry of Health, and University Hospitals of Cleveland Medical Center Institutional Review Board (no. 04-14-19), approved the protocols. Written informed consent was obtained from all study participants or their parents/guardians.

Protein expression and purification

The protein sequences for each of the antigens were selected: PfAMA1 (XP_001348015.1; 3D7 genotype), PvAMA1 (ACB42433.1; Palo Alto genotype), PvAMA1 (PNG 20 genotype) and PkAMA1 (XP_002259339.1; Strain H) as previously described [29]. At the N-terminus of the sequences, the native signal peptide was removed and replaced with a signal peptide for tissue plasminogen activator, followed by a 6-histidine tag. At the C-terminus of all sequences, the sequence was truncated to remove the transmembrane domain and cytoplasmic tail. All sequences were assessed for potential glycosylation sites (https://services.healthtech.dtu.dk/services/NetNGlyc-1.0/). The AMA1 sequences for each species were then modified to prevent potential glycosylation (PfAMA1, six changes; PvAMA1, three changes; PkAMA1, seven changes) [29].

Preparation of the PvAMA1 probe and cell sorting

The addition of biotin to PvAMA1_Palo Alto was done using EZ-Link NHS-PEG12-Biotin based on the manufacturers' instructions. Using a Zeba spin 7 kDa cutoff buffer exchange spin column eliminated free biotin and changed to Bicine buffer, 50 mM pH 8.3. PvAMA1_Palo Alto biotin was quantified by BCA Protein assay (Pierce) and biotin/molecule calculated as 2 biotin per molecule using FluoReporter ® Biotin Quantitation Assay, Invitrogen.

CD19+ human B cells were isolated from 30 x 10^6 cryopreserved PBMCs using Miltenyi CD19 microbeads and then stained at 4°C for LIVE/DEAD, CD20, IgG, PvAMA1_Palo

Alto_biotin tetramers prepared with Streptavidin-FITC and Streptavidin-Brilliant Violet 421. Single cells were sorted as a 96 well sort to a chilled wells on a FACS_ARIA_SORP. 252 single cells were isolated into 4 μL of catch buffer. Stained CD19 cells were stored at 4°C overnight and 252 more cells were collected as a dry catch.

Cell staining and sorting of PvAMA1-specific Memory B cells

Single cells were identified and sorted according to previously described techniques [62] from cryopreserved PBMC without activation. Briefly, staining and single-cell sorting of PvAMA1-specific IgG+ MBCs were performed as follows: B cells were enriched using immunomagnetic positive selection with anti-CD19 magnetic MACS beads (Miltenyi Biotec) and stained with mouse anti-human CD20 (PE-Cy5.5; Invitrogen) and anti-human IgG Abs (PE-Cy7 clone G18-145; Becton Dickinson) along with biotinylated PvAMA1 using Streptavidin coupled with allophycocyanin (Becton Dickinson) and SYTOX Green Dead Cell Stain (Invitrogen) to gate out dead cells. Stained CD19$^+$ cells were sorted on a BD FACSAria II equipped with chilled stage sorting based on size and complexity. Doublet discrimination was performed to exclude aggregated cells. Individual AMA1-specific CD20$^+$, IgG+ MBCs were single cell sorted directly into 4 μL of mRNA extraction buffer on a cooled 96-well metal block. After cells were collected, plates were frozen immediately on dry ice and stored at -80°C until further processing.

cDNA synthesis

The 96-well plates with single cells were thawed on ice; a cold volume of 7 μL containing 300 ng of random hexamers (Qiagen Operon), 12 U Rnasin (Promega), and 0.9% NP-40 (Thermo Scientific Pierce) was added to each well. After thorough pipetting and rinsing, wells were capped, centrifuged at 4°C, heated to 68°C in a thermal cycler for 5 min, and placed on ice for at least 1 min. Reverse transcription was performed with the addition of 7 μL containing 3.6 μL of 53 reverse transcriptase buffer, 10 U RNAsin (Promega), 62 U Superscript III reverse transcriptase (Invitrogen), 0.62 μL dNTPs 25 mM each (V Bio-Tek), and 1.25 μL of 0.1 M DTT

(Sigma- Aldrich). All wells were capped, and the plate placed in a cold rack and vortexed for 10 sec before centrifugation at 300 x g. Thermal cycler conditions for reverse transcription were as follows: 42°C for 5 min, 25°C for 10 min, 50°C for 60 min, 94°C for 5 min, and 4°C hold. When completed, 10 µL of nuclease-free PCR water was added to each well.

Ig gene amplification

Immediately following cDNA synthesis, IgG genes (Igg) were amplified in a total of 20 µL per well for the first round of nested PCR for IgG H chain (Iggh), IgG k (Iggk) and IgG l (Iggl), using master mix (**Table 1**) and primers (**Table 2**) as previously described REF. cDNA from individual sorted B cells were added and Igg amplified under the following conditions: thermal cycle PCR at 94°C for 15 min; 50 cycles at 94°C for 30 sec, then 58°C (Iggh and Iggk) or 60°C (Iggl) for 30 sec, then 72°C for 55 sec; then one cycle at 72°C for 10 min. Second round of nested PCR for Iggh, Iggk, and Iggl used master mix (**Table 1**), 2 µL of first-round PCR product with second-round primers (**Table 3**) and the same master mix protocol, with the following conditions; thermal cycle PCR at 94°C for 15 min; 50 cycles at 94°C for 30 sec, then 58°C (Iggh and Iggk) or 60°C (Iggl) for 30 sec, then 72°C for 45 sec; then one cycle at 72°C for 10 min. The PCR product generated was purified and sequenced, with V(D)J genes determined using IMGT/V-Quest.

Table 1 – PCR Round 1 Master Mix Specifics

Component	Volume for 1 well (µL)
RNAse Free Water	10.79
5x Q-Solution	4.0
10x HotStar Taq PCR Buffer	2.0
25 mM MgCl2	0.55
dNTP Mix (25 mM each dNTP)	0.20
5' LV-Primer Group Mix	0.130
3' Primer (50 mM)	0.130
HotStar Taq (5 units/µL)	0.20
DNA Template	2.00
Total Volume (w/out DNA)	18.00

Table 2 – PCR Round 1 Primers

	5'	3'
Gamma	LVH Group Mix	CGammaCH1
Kappa	LVKappa Group Mix	CK543
Lambda	LVLambda Group Mix	Clambda

Table 3 – PCR Round 2 Primers

	5'	3'
Gamma	AgeVH Group Mix	IgGInternal
Kappa	PanVK	CK494
Lambda	AgeVLambda Group Mix	CLambdaXho

Specific V(D)J region amplification and cloning

Primers specific with restriction enzyme sites for V and J regions were used to amplify the first-round PCR product to generate a fragment for cloning based on previously described primers [62]. PCR product was purified, and restriction enzyme digested, cloned into Iggh, Iggk, or Iggl expression vectors, and chemically transformed into 5 µL of aliquots of TOP10 *E. coli* cells (Thermo Fisher Scientific). Successful transformants were screened by PCR amplification (**Table 4**) using a vector-specific primer paired with an insert-specific primer with the following conditions; thermal cycle PCR at 94°C for 15 min; 50 cycles at 94°C for 30 sec, then 57°C (Mu/Gamma/Kappa) or 60°C (Lambda) for 30 sec, then 72°C for 45 sec; then one cycle at 72°C for 10 min. Products were sequenced and compared with the second-round PCR product sequence.

Table 4 – Specific V(D)J Region Amplification Master Mix Specifics

Component	Volume for 1 well (µL)
RNAse Free Water	18.0
5x Q-Solution	8.0
10x HotStar Taq PCR Buffer	4.0
25 mM MgCl2	1.0
dNTP Mix (25 mM each dNTP)	0.4
5' primer at 5 µm	4.0
3' primer at 5 µm	4.0
HotStar Taq (5 units/µL)	0.4
DNA Template	4.2

Total Volume (w/out DNA)	39.8

Definition of clonal groups

Clonal groups were based on H-chain nucleotide sequences. Any PCR product with 0.8% nucleotide sequences with a Phred score of 20 was excluded. We determined H chain alleles from PCR-amplified sequences using IMGT/V-QUEST (http://www.imgt.org, international ImMunoGeneTic information system). Because of primer mixture ambiguities, the first 20–22 nt of IgGH variable regions were designated germline; thus, this region was not evaluated for somatic hypermutations. IMGT/V-QUEST was used to assign V(D)J organization and shared IgHV genes and CDR3 length-grouped sequences. Clonal grouping was determined using Sequence Manipulation Suite: Ident and Sim using and single-cell sorting of PvAMA1-specific IgG+ memory B cells were Ab-specific clusters as previously defined. A clonal group is defined by the same IGHV gene and CDR3 length, along with 72% similarity of each CDR3 aa sequence.

HumAb expression and purification

Briefly, following transformation and sequencing to confirm that the plasmid contains the desired nucleotide sequence, glycerol stocks of *E. coli* containing the proper plasmid were stored at -80°C. Plasmids were harvested from ampicillin grown *E. coli* cultures (50 - 250 mL) using the ZymoPURE Midiprep Purification Kit. Purified DNA was quantified using a NanoDrop and stored at 4°C until use. Transfection sizes were then dependent on DNA yield. Cells were transfected at a density of 3 - 5 x 10^6 viable cells per mL. On the day before transfection (Day −1), Expi293F™ culture was split to a final density of 2.5 – 3.0 × 10^6 viable cells/mL and allowed to grow overnight. The cells reached a density of approximately 4.5 – 5.5 × 10^6 viable cells/mL at the day of transfection. Cells were diluted with Expi293™ Expression Medium, pre-warmed to 37°C to a final density of 3 × 10^6 viable cells/mL. Plasmid DNA was diluted in Opti-MEM™ I

Reduced Serum Medium at 1 μg/mL of transfected cells. ExpiFectamine™ 293 Reagent was diluted with Opti-MEM™ I Reduced SerumMedium and incubated at room temperature for 5 minutes prior to initiating the plasmid DNA complexation reaction. Diluted ExpiFectamine™ 293 Reagent was added to diluted plasmid DNA and mixed gently by swirling. ExpiFectamine™ 293/plasmid DNA complexes were incubated at RT for 10 - 20 minutes and slowly transferred to the cell culture with continuous mixing. Cells were incubated at 37°C with a humidified atmosphere of 8% CO_2 in the air with continuous orbital shaking. 18 - 22 hours post-transfection, ExpiFectamine™ 293 Transfection Enhancer 1 and ExpiFectamine™ 293 Transfection Enhancer 2 were added to the flask per company instruction and gently mixed. The flask was returned to the 37°C incubator with a humidified atmosphere of 8% CO_2 with shaking. Six to seven days post transfections cells are centrifuged at 4,000 x g for 20 minutes at RT. The supernatant is decanted and filtered through a 0.22 μm sterile filter and stored at 4°C until purification. The volume of the cleared cell supernatant was supplemented with 1 M Tris pH 8.0 and 5 M NaCl solutions to a final concentration of 0.1 M Tris pH 8.0 and 0.5 M NaCl and loaded onto a 5 mL HiTrap MabSelect PrismA column. Once the complete sample was loaded onto the column, the unbound sample was washed out with 10 column volumes (CV) 1x PBS. Elution was carried out with 10 CV using a commercial IgG Elution buffer pH 2.5 (Thermo Fisher) into a collection tube containing 1 mL of 1M Tris pH 8.0 for immediate neutralization. Following the elution, the column was washed with 10 CV of 50 mM NaOH and re-equilibrated to 1x PBS pH 7.4. Eluted humAbs were dialyzed overnight in 1x PBS at 4°C. Protein concentrations were determined by Nanodrop (Thermo Fisher Scientific) using OD 1.3 for a 1 mg/mL concentration.

Antibody Titration Curve (MagPix) Methods

3.6 μl of 1 mg/mL PvAMA1_PaloAlto was conjugated to 300 μL of MagPix bead 19 following the manufacturer's protocol (Luminex, Corp, Austin, TX). 1 μl of 4 mg/mL PfAMA1_3D7 was conjugated to 200 μL of MagPix bead 20, 4 μL of 1 mg/mL PvAMA1_PNG16

was conjugated to 200 µL of MagPix bead 18, 4 µL of 1 mg/mL PkAMA1 was conjugated to 200 µL of MagPix bead 56, 4 µL of 1 mg/mL TgAMA1 was conjugated to 200 µL of MagPix bead 12. 50 µL of humAb at the following concentrations (1.0, 0.5, 0.25, 0.125, 0.0625, 0.03125, 0.015625, 0.0078125 µg/mL) were mixed with 50 µL of bead master mix (diluted 1:1000 in 1% BSA-TBST). Samples were incubated for 10 min at RT while shaking, and an additional 20 min while sitting. Plate was placed on a magnet for 5 min and excess liquid was removed. Plate was washed two times with 1x TBST for 2 min with shaking, followed by an additional 5 min on a magnet. Donkey anti-Human IgG (PE conjugated, Jackson ImmunoResearch) secondary antibody was diluted 1:400. 25 µL of secondary antibody was added to the plate and samples were incubated for 10 min at RT while shaking, and an additional 20 min while sitting. Plate was placed on a magnet for 5 min and excess liquid was removed. Plate was washed two times with 1x TBST for 2 min with shaking, followed by an additional 5 min on a magnet. 150 µL of 1% BSA-TBST was added to the plate, shaken for 2 min, and read using the MagPix machine.

Antibody Avidity (MagPix) Methods

The beads used in the humAb titration were also used to measure antibody avidity. 50 µL of 0.2 µg/mL of humAb was mixed with 50 µL of bead master mix (diluted 1:1000 in 1% BSA-TBST). Samples were incubated for 10 min at RT while shaking, and an additional 20 min while sitting. Plate was placed on a magnet for 5 min and excess liquid was removed. Plate was washed two times with 1x TBST for 2 min with shaking, followed by an additional 5 min on a magnet. 50 µL of Ammonium Thiocyanate at the following final concentrations of 0, 0.9, 1.2, 1.5, 1.8, 2.1, 2.4, 2.7, 3.0, 3.3, 3.6, 3.9 M was added to the plate. Samples were incubated for 5 min at RT while shaking, and an additional 5 min while sitting. The plate was placed on a magnet for 5 min and excess liquid was removed. The plate was washed two times with 1x TBST for 2 min with shaking, followed by an additional 5 min on a magnet. Donkey anti-Human IgG (PE conjugated, Jackson ImmunoResearch) secondary antibody was diluted 1:400. 25 µL of

secondary antibody was added to the plate and samples were incubated for 10 min at RT while shaking, and an additional 20 min while sitting. Plate was placed on a magnet for 5 min and excess liquid was removed. Plate was washed two times with 1x TBST for 2 min with shaking, followed by an additional 5 min on a magnet. 150 μL of 1% BSA-TBST was added to the plate, shaken for 2 min, and read using the MagPix machine. From this, GraphPad Prism was used to calculate the IC_{50} of each humAb.

Measurement of Antibody Affinity

To determine the binding affinity of each humAb to PvAMA1, Surface Plasmon Resonance (SPR) was utilized to perform single cycle kinetics. One CM5 chip (GE Healthcare) flow cell was coated with 275 RUs of anti-Human IgG antibody (Abcam). An untreated cell was used as a blank reference. Data was collected at 25°C and 10 Hz on a Biacore T200 system. Pilot studies of the interaction and follow-up regeneration protocols were established before data collection. Three start-up cycles were run with 1x HBSS Running Buffer (GE Healthcare). A humAb, at a concentration of 80 nM, was passed over all the flow cells at a rate of 10 μL/min for a contact time of 30 sec and allowed to stabilize for 120 sec. Increasing concentrations (5 nM, 10 nM, 20 nM, 40 nM, 80 nM) of rPvAMA1 were passed over all flow cells at a rate of 30 μL/min. Each injection was allowed to stabilize for 60 sec and was given 120 sec for dissociation. Anti-Human Fc chip was regenerated to baseline using IgG Elution Buffer (Thermo Scientific) and the process was repeated for all humAbs in the panel. Data was processed using the double referencing method using Scrubber 2.0 (BioLogic Software).

Transgenic Pf-PvAMA1 Growth Inhibition Assays

Invasion assays were performed as previously described [63], with minor changes. Parasites were synchronized with sorbitol, then inoculated at late trophozoite and schizont stages at 0.5% parasitemia in 4% hematocrit in 45 μL RPMI media, with 5 μL antibodies in 1x

PBS. Cultures were left to invade, then harvested at the ring stage by fixation with 0.25% glutaraldehyde. Parasites were stored in 1x PBS at 4°C. Cultures were later stained with 1× SYBR Green I (S7563, Invitrogen), and parasitemia assayed by flow cytometry in a BD FACSCanto™ II system, with further analysis in FlowJo™ v10.9.0 (BD Life Sciences). Parasites were gated from uninfected erythrocytes using the FITC-A and PE-A channels. Net growth rates were calculated by subtracting the parasitemia of parasites inoculated with 1 mg/mL heparin, then dividing by the parasitemia of parasites inoculated with 5 µL PBS. Growth percentages were subtracted from 100% to give invasion inhibition.

IC_{50} calculations were performed in R [R Core Team (2023). R: A Language and Environment for Statistical Computing. R Foundation for Statistical Computing, Vienna, Austria. https://www.R-project.org/] using the packages drc [64] and ggplot2 [65]. Assays were performed in biological triplicate, and IC_{50}s calculated independently for each replicate. The parasite lines were based on W2mef, with the AMA1 locus replaced with the *P. vivax* locus [51].

Parasite culturing and transgenic PfPvAMA1 inhibition experiments

P. falciparum asexual stage parasites were maintained in culture in human erythrocytes (blood group type O +) at a hematocrit of 4% in RPMI-HEPES supplemented with 0.25% (w/v) Albumax™ (Invitrogen) and 5% (v/v) heat-inactivated human serum. *P. falciparum* was synchronized using sorbitol and heparin treatments as described previously [48]. Transgenic W2mef *P. falciparum* strains expressing PfAMA1 W2mef allele (W2-W2), PfAMA1 3D7 allele (W2- 3D7) and PvAMA1 Palo Alto allele (W2-PvAMA1) were generated in previous studies [51].

In vitro invasion inhibition assay using Pv clinical isolates, DNA extraction amplification and sequencing of *PvAMA1*

Clinical isolates of *P. vivax* were either used fresh after blood collection or cryopreserved and infected RBCs were thawed and cultured in IMDM medium (Gibco) supplemented with

0.5% Albumax II (Gibco), 2.5% heat-inactivated human serum, 25mM HEPES (Gibco), 20 µg/mL gentamicin (Sigma) and 0.2mM hypoxanthine (C-C Pro) for ~24 or ~48 hr until a majority of schizont stage parasites were observed as previously described [66]. The schizont-infected erythrocytes were enriched using KCl-Percoll density gradient [67], then mixed at a ratio of 1 erythrocyte to 1 reticulocyte enriched from cord blood and previously labeled with CellTrace Far Red dye following manufacturer's instructions. The cultures were incubated for ~10 hr in a final volume of 50 µL in 384 well plates, in presence of the humAbs. The mouse monoclonal anti-Duffy 2C3 at 100 µg/mL was used as positive invasion inhibition control [66]. Cells were stained with DNA stain Höchst 33342 post-invasion and examined by flow cytometry. Reticulocytes, which were Höchst 33342 and Far Red positive, were scored as new invasion events. For quantification, data were normalized against parasites mock-treated with 1x PBS. Invasion of reticulocytes in absence of antibodies ranged from 0.2 % to 12.2% (mean 3.37%). The work presented here was approved by the National Ethics Committee for Health Research of Cambodia (192NECHR). All patients and/or their parents/guardians provided informed written consent.

Genomic DNA was extracted from an aliquot of the parasitized blood samples using the QIAamp DNA Blood Minit Kit (QUIAGEN), according to manufacturer's instructions. PvAMA1 sequences were determined by PCR and Sanger sequencing (Macrogen, Seoul, South Korea) using the following conditions. The PCR was conducted in 20 µL reaction consisting of 2 µL of DNA, 0.25 µM of primers (Forward 5'- AAGCTGCTCACCCGTTAGTG -3'; Reverse 5'-GGGTGGGAAGGTGCATTCTG-3') and 1X HOT FirePol Blend MasterMix (Solis BioDyne, Tartu, Estonia) under the following conditions: 95°C for 1 min, followed by 34 cycles 95°C for 20 sec, 63.8°C for 30 sec, 72°C for 2 min and a final extension at 72°C for 5 min. Nucleotides and corresponding amino acids were analyzed using MEGA 11 software. The sequences generated were compared with the reference strain Sal1 (PVX_092275).

RON2 peptide Competition

We conducted a RON2 binding assay as previously described [29] with a few modifications. 96-well plates were coated with PvAMA1. HumAb was serially diluted in 0.1% casein in 1x PBS by a factor of two, with concentrations ranging from 40–0.039 μg/mL. 50 μL of humAb was added to each well. 50 μL 0.1% casein in 1x PBS was added to additional wells as blank controls. The optical density (OD) of these were later subtracted from each of the other wells to correct for background. Each well, except for the blank control wells, was incubated with 50 μL of 1 μg/mL biotinylated PvRON2 peptide (Asp 2050 – Thr 2088) [29] in 1x PBS. Streptavidin–horse-radish-peroxidase conjugated protein (Thermo Fisher Scientific, 21130) was added at 1/500. Plates were incubated with ABTS for 20 min before reading.

In vitro invasion into HC04 hepatocytes using Pv Sporozoite in Thailand and primary human hepatocytes in Cambodia

Anopheles dirus mosquitoes were fed on blood collected from symptomatic patients attending malaria clinics in Tak, Songkla, and Ubon-Ratchathani Provinces in Thailand and in Kampong Speu province in Cambodia. Samples were confirmed positive for only P. vivax via microscopy and RT-PCR. P. vivax-infected blood samples were collected from patients under the approved protocol by the Ethics Committee of the Faculty of Tropical Medicine, Mahidol University (MUTM 2018-016). Sample collection in Cambodia was approved by the National Ethics Committee for Health Research of Cambodia (192NECHR). Written informed consent was obtained prior to blood collection.

Briefly, P. vivax infected blood was drawn into heparinized tubes and kept at 37°C until processing. Infected blood was washed once with RPMI 1640 incomplete medium. Packed infected blood was resuspended in warm, non-heat inactivated naive human AB serum (Thailand) or in heat-inactivated naive human AB serum (Cambodia) for a final hematocrit of 50%. Resuspended blood was fed to laboratory-reared female Anopheles dirus mosquitoes for

30 min via an artificial membrane attached to a water-jacketed glass feeder kept at 37°C. Engorged mosquitoes were kept on 10% sugar at 26°C under 80% humidity at the designated insectary at the Mahidol Vivax Research Unit or the Malaria Research Unit of Institut Pasteur du Cambodge. Sporozoites were dissected from the salivary glands of infected mosquitoes 14-21 days after blood feeding and pooled in DMEM supplemented with 200 mg/mL penicillin-streptomycin (Thailand) or in RPMI without NaHCO3 (Cambodia) [8].

In assays performed in Thailand, a 96-well plate was seeded with HC04 hepatocytes (50,000 cells/well). Isolated sporozoites (100,000 per well) were pre-incubated for 30 min with the humAbs at various concentrations (0.1, 1, 10, 100, and 1000 µg/mL). Sporozoite/humAb mixture was added to hepatocyte culture and incubated for 4 days at 37°C in duplicate. Murine anti-PvCSP polyclonal antibodies CSP 210 (clone 2F2) and 247 were used as positive controls [52]. Anti-tetanus toxoid humAb 043038 was used as a negative control.

In the assay performed in Cambodia a 384-well plate was seeded with primary human hepatocytes (18,000 cells/well). Isolated sporozoites (16,000 per well) were pre-incubated for 20 min with the humAb 826827 at various concentrations (0.1, 1, 10, 100, and 1000 µg/mL), with the humAb 043038 at 1000 µg/mL and with non-treated cell culture medium (latter two as negative controls). Each condition was tested in five technical replicates. Cell culture medium was changed to non-treated medium after 24h of infection and the plate was incubated for 6 days at 37°C.

Cells were fixed and stained using sporozoite specific polyclonal antibodies UIS4 and DAPI. Wells were then analyzed using fluorescent microscopy and invaded hepatocytes were quantified.

Protein expression and purification of PvAMA1 and humAb 826827 for crystallization

For crystallization purposes, PvAMA1 (ACB42433.1; Palo Alto genotype, [68]) and humAb 826827 were expressed using transient transfection of Expi293 cells at a density of 2.5

x10^6 cells/mL following the manufacturer's protocol. Five days after transfection, supernatants were harvested by centrifugation at 7,000 x g and dialyzed overnight into 30 mM Tris pH 7.5, 300 mM NaCl. Dialyzed PvAMA1-supernatant was incubated with Ni-NTA resin (Qiagen) for 1hr at 4°C followed by stepwise elution using gravity flow chromatography. Size exclusion chromatography (SD200 increase, in 20 mM HEPES pH 7.5, 150 mM NaCl) was used as the final purification step. Supernatant of humAb 826827 was loaded onto a 5 mL Mabselect PrismA column (Cytiva) equilibrated in 1x PBS. After washing the column with 1x PBS, humAb 826827 was eluted using 0.1 M citric acid pH 3.0 and neutralized with 1 M Tris pH 9.0 and buffer exchanged into 1x PBS.

Complex formation, crystallization, and structure determination

HumAb 826827 was incubated with enzyme FabALACTICA (IgdE) at a ratio of 50 μg enzyme/mg humAb for 24 - 48 hr at 37ºC. Fab fragments were purified by affinity chromatography using a 1 mL HiTrap KappaSelect column (Cytiva), eluted with 0.1 M glycine pH 1.5 and immediately neutralized with 1 M Tris pH 9.0.

For crystallization, PvAMA1- Fab 826827 complexes were formed by incubating the PvAMA1 and Fab 826827 at a molar ratio of 1.2:1 for 1 hr on ice. This complex was separated from excess Fab using size exclusion chromatography in 20 mM HEPES pH 7.5, 150 mM NaCl. Crystallization screens were set up at the Monash Macromolecular Crystallization Platform (MMCP, Clayton, VIC, Australia) with 3 - 6 mg/mL at 20°C. Initial PvAMA1-Fab 826827 crystals grew in 22% PEG Smear Broad, 0.1 M Tris pH 8.5 (BCS screen, Molecular Dimensions) and were optimized by additive screening in a sitting drop vapor diffusion experiment (Additive Screen HT, Hampton Research). A PvAMA1-Fab 826827 crystal obtained in 22% PEG Smear Broad, 0.1 M Tris pH 8.5, 0.3 M Glycyl-glycyl-glycine was harvested with 30% glycerol in mother liquor as cryo-protectant and used for data collection.

X-ray diffraction data was collected at the MXII beamline of the Australian Synchrotron at 100 K [69]. The XDS package [70] was used for data processing. Molecular replacement with Phaser was used to solve the phase problem using structural coordinates of PvAMA1 (PDB ID 5NQG, [26]) and Fab coordinates of PBD ID 6FG1 [71]. Iterative cycles of structure building and refinement was carried out using Coot [72] and Phenix [73]. Figures of the structure were prepared with PyMOL (Version 2.5.2 Schrödinger, LLC) or VIDA 4.4 (OpenEye, Cadence Molecular Science). The atomic coordinates and structure factor files have been deposited in the Protein Data Bank under PDB ID 8U9D.

Data Availability

Coordinates and structure factors of rPvAMA1_Palo Alto in complex with 826827 have been deposited in the Protein Data Bank (PDB) under PDB ID 8U9D.

References

1. Poespoprodjo, J.R., et al., *Malaria.* Lancet, 2023.
2. Menkin-Smith, L. and W.T. Winders, *Plasmodium vivax Malaria*, in *StatPearls*. 2023: Treasure Island (FL).
3. Battle, K.E. and J.K. Baird, *The global burden of Plasmodium vivax malaria is obscure and insidious.* PLoS Med, 2021. **18**(10): p. e1003799.
4. Kho, S., et al., *Evaluation of splenic accumulation and colocalization of immature reticulocytes and Plasmodium vivax in asymptomatic malaria: A prospective human splenectomy study.* PLoS Med, 2021. **18**(5): p. e1003632.
5. Kho, S., et al., *Hidden Biomass of Intact Malaria Parasites in the Human Spleen.* N Engl J Med, 2021. **384**(21): p. 2067-2069.
6. Molina-Franky, J., et al., *The Black Box of Cellular and Molecular Events of Plasmodium vivax Merozoite Invasion into Reticulocytes.* Int J Mol Sci, 2022. **23**(23).
7. Douglas, N.M., et al., *The anaemia of Plasmodium vivax malaria.* Malar J, 2012. **11**: p. 135.
8. Mancio-Silva, L., et al., *A single-cell liver atlas of Plasmodium vivax infection.* Cell Host Microbe, 2022. **30**(7): p. 1048-1060 e5.
9. Commons, R.J., et al., *Estimating the Proportion of Plasmodium vivax Recurrences Caused by Relapse: A Systematic Review and Meta-Analysis.* Am J Trop Med Hyg, 2020. **103**(3): p. 1094-1099.

10.   Chu, C.S. and N.J. White, *The prevention and treatment of Plasmodium vivax malaria.* PLoS Med, 2021. **18**(4): p. e1003561.

11.   Gisele Tatiane Soares da Veiga, M.R.M., Jean Franciesco Vettorazzi, Marcelo Muller-Santos, Letusa Albrecht, *Plasmodium vivax vaccine: What is the best way to go?* Frontiers in Immunology, 2023. **13**.

12.   Schafer, C., et al., *Partial protection against P. vivax infection diminishes hypnozoite burden and blood-stage relapses.* Cell Host Microbe, 2021. **29**(5): p. 752-756 e4.

13.   Wai-Hong Tham, J.G.B., Julian C. Rayner, *Plasmodium vivax vaccine research – we've only just begun.* International Journal of Parasitology, 2017. **47**(2-3): p. 111-118.

14.   Lamarque, M., et al., *The RON2-AMA1 Interaction is a Critical Step in Moving Junction-Dependent Invasion by Apicomplexan Parasites.* Plos Pathogens, 2011. **7**(2).

15.   Mitchell, G.H., et al., *Apical membrane antigen 1, a major malaria vaccine candidate, mediates the close attachment of invasive merozoites to host red blood cells.* Infect Immun, 2004. **72**(1): p. 154-8.

16.   Cowman, A.F., C.J. Tonkin, W.H. Tham, and M.T. Duraisingh, *The Molecular Basis of Erythrocyte Invasion by Malaria Parasites.* Cell Host Microbe, 2017. **22**(2): p. 232-245.

17.   Fernandes, P., et al., *The AMA1-RON complex drives Plasmodium sporozoite invasion in the mosquito and mammalian hosts.* PLoS Pathog, 2022. **18**(6): p. e1010643.

18.   Silvie, O., et al., *A role for apical membrane antigen 1 during invasion of hepatocytes by Plasmodium falciparum sporozoites.* J Biol Chem, 2004. **279**(10): p. 9490-6.

19.   Kayentao, K., et al., *Safety and Efficacy of a Monoclonal Antibody against Malaria in Mali.* N Engl J Med, 2022. **387**(20): p. 1833-1842.

20.   Srinivasan, P., et al., *Disrupting malaria parasite AMA1-RON2 interaction with a small molecule prevents erythrocyte invasion.* Nat Commun, 2013. **4**: p. 2261.

21.   Tyler, J.S., M. Treeck, and J.C. Boothroyd, *Focus on the ringleader: the role of AMA1 in apicomplexan invasion and replication.* Trends Parasitol, 2011. **27**(9): p. 410-20.

22.   Boucher, L.E. and J. Bosch, *The apicomplexan glideosome and adhesins - Structures and function.* J Struct Biol, 2015. **190**(2): p. 93-114.

23.   Srinivasan, P., et al., *Binding of Plasmodium merozoite proteins RON2 and AMA1 triggers commitment to invasion.* Proc Natl Acad Sci U S A, 2011. **108**(32): p. 13275-80.

24.   Collins, C.R., et al., *Fine mapping of an epitope recognized by an invasion-inhibitory monoclonal antibody on the malaria vaccine candidate apical membrane antigen 1.* J Biol Chem, 2007. **282**(10): p. 7431-41.

25.   Deans, J.A., et al., *Rat monoclonal antibodies which inhibit the in vitro multiplication of Plasmodium knowlesi.* Clin Exp Immunol, 1982. **49**(2): p. 297-309.

26.   Vulliez-Le Normand, B., et al., *Cross-reactivity between apical membrane antgen 1 and rhoptry neck protein 2 in P. vivax and P. falciparum: A structural and binding study.* PLoS One, 2017. **12**(8): p. e0183198.

27.   Mannuthodikayil, J., et al., *A Chimeric Peptide Inhibits Red Blood Cell Invasion by Plasmodium falciparum with Hundredfold Increased Efficacy.* Chembiochem, 2023. **24**(7): p. e202200533.

28.   Lee, S.K., et al., *The direct binding of Plasmodium vivax AMA1 to erythrocytes defines a RON2-independent invasion pathway.* Proc Natl Acad Sci U S A, 2023. **120**(1): p. e2215003120.

29. Drew, D.R., et al., *Defining species-specific and conserved interactions of apical membrane protein 1 during erythrocyte invasion in malaria to inform multi-species vaccines.* Cell Mol Life Sci, 2023. **80**(3): p. 74.

30. Beeson, J.G., et al., *Merozoite surface proteins in red blood cell invasion, immunity and vaccines against malaria.* FEMS Microbiol Rev, 2016. **40**(3): p. 343-72.

31. Hodder, A.N., P.E. Crewther, and R.F. Anders, *Specificity of the protective antibody response to apical membrane antigen 1.* Infect Immun, 2001. **69**(5): p. 3286-94.

32. Lopez, C., Yepes-Perez, Yoelis, Hincapie-Escobar, Natalia, Diaz-Arevalo, Diana, Patarroyo, Manuel A., *What Is Known about the Immune Response Induced by Plasmodium vivax Malaria Vaccine Candidates?* Frontiers in Immunology, 2017. **8**.

33. Antonelli, L.R., et al., *The immunology of Plasmodium vivax malaria.* Immunol Rev, 2020. **293**(1): p. 163-189.

34. Udhayakumar, V., et al., *Longitudinal study of natural immune responses to the Plasmodium falciparum apical membrane antigen (AMA-1) in a holoendemic region of malaria in western Kenya: Asembo Bay Cohort Project VIII.* Am J Trop Med Hyg, 2001. **65**(2): p. 100-7.

35. Bailey, J.A., et al., *Microarray analyses reveal strain-specific antibody responses to Plasmodium falciparum apical membrane antigen 1 variants following natural infection and vaccination.* Sci Rep, 2020. **10**(1): p. 3952.

36. Rodrigues, M.H., et al., *Antibody response of naturally infected individuals to recombinant Plasmodium vivax apical membrane antigen-1.* Int J Parasitol, 2005. **35**(2): p. 185-92.

37. Polley, S.D., et al., *Human antibodies to recombinant protein constructs of Plasmodium falciparum Apical Membrane Antigen 1 (AMA1) and their associations with protection from malaria.* Vaccine, 2004. **23**(5): p. 718-28.

38. Salinas, N.D., W.K. Tang, and N.H. Tolia, *Blood-Stage Malaria Parasite Antigens: Structure, Function, and Vaccine Potential.* J Mol Biol, 2019. **431**(21): p. 4259-4280.

39. Suman Sinha, A.B., Jagannath Mondal, Kalyaneswar Mandel, *Computational and experimental studies of the breathing motion of a protein loop: implications in PfAMA1-PfRON2 late-stage binding event.* bioRxiv, 2022.

40. Swearingen, K.E., et al., *Proteogenomic analysis of the total and surface-exposed proteomes of Plasmodium vivax salivary gland sporozoites.* PLoS Negl Trop Dis, 2017. **11**(7): p. e0005791.

41. Alicia Arnott, I.M., Paul A. Ramsland, Peter M. Siba, John C. Reeder, Alyssa E. Barry, *Global Population Structure of the Genes Encoding the Malaria Vaccine Candidate, Plasmodium vivax Apical Membrane Antigen 1 (PvAMA1).* PLoS Neglected Tropical Diseases, 2013. **7**(10).

42. Pizarro, J.C., et al., *Crystal structure of the malaria vaccine candidate apical membrane antigen 1.* Science, 2005. **308**(5720): p. 408-11.

43. Vulliez-Le Normand, B., et al., *Crystal structure of Plasmodium knowlesi apical membrane antigen 1 and its complex with an invasion-inhibitory monoclonal antibody.* PLoS One, 2015. **10**(4): p. e0123567.

44. Vulliez-Le Normand, B., et al., *Structural and functional insights into the malaria parasite moving junction complex.* PLoS Pathog, 2012. **8**(6): p. e1002755.

45. Coley, A.M., et al., *The most polymorphic residue on Plasmodium falciparum apical membrane antigen 1 determines binding of an invasion-inhibitory antibody.* Infect Immun, 2006. **74**(5): p. 2628-36.

46. Aleshnick, M., M. Florez-Cuadros, T. Martinson, and B.K. Wilder, *Monoclonal antibodies for malaria prevention.* Mol Ther, 2022. **30**(5): p. 1810-1821.

47. Gaudinski, M.R., et al., *A Monoclonal Antibody for Malaria Prevention.* N Engl J Med, 2021. **385**(9): p. 803-814.

48. Lyke, K.E., et al., *Low-dose intravenous and subcutaneous CIS43LS monoclonal antibody for protection against malaria (VRC 612 Part C): a phase 1, adaptive trial.* Lancet Infect Dis, 2023. **23**(5): p. 578-588.

49. Maskus, D.J., et al., *Characterization of a novel inhibitory human monoclonal antibody directed against Plasmodium falciparum Apical Membrane Antigen 1.* Sci Rep, 2016. **6**: p. 39462.

50. Crawford, J., M.L. Tonkin, O. Grujic, and M.J. Boulanger, *Structural characterization of apical membrane antigen 1 (AMA1) from Toxoplasma gondii.* J Biol Chem, 2010. **285**(20): p. 15644-15652.

51. Damien R. Drew, P.R.S., Gretchen Weiss, Paul R. Gilson, Brendon S. Crabb, James G. Beeson, *Functional Conservation of the AMA1 Host-Cell Invasion Ligand Between P. falciparum and P. vivax: A Novel Platform to Accelerate Vaccine and Drug Development.* The Journal of Infectious Diseases, 2017. **217**: p. 498-507.

52. Kucharska, I., et al., *Structural basis of Plasmodium vivax inhibition by antibodies binding to the circumsporozoite protein repeats.* Elife, 2022. **11**.

53. Akter, M., et al., *Identification of the Binding Site of Apical Membrane Antigen 1 (AMA1) Inhibitors Using a Paramagnetic Probe.* ChemMedChem, 2019. **14**(5): p. 603-612.

54. Delgadillo, R.F., et al., *Stability of the Plasmodium falciparum AMA1-RON2 Complex Is Governed by the Domain II (DII) Loop.* PLoS One, 2016. **11**(1): p. e0144764.

55. Parker, M.L. and M.J. Boulanger, *An Extended Surface Loop on Toxoplasma gondii Apical Membrane Antigen 1 (AMA1) Governs Ligand Binding Selectivity.* PLoS One, 2015. **10**(5): p. e0126206.

56. Tonkin, M.L., et al., *Host cell invasion by apicomplexan parasites: insights from the co-structure of AMA1 with a RON2 peptide.* Science, 2011. **333**(6041): p. 463-7.

57. Ashley, E.A., J. Recht, and N.J. White, *Primaquine: the risks and the benefits.* Malar J, 2014. **13**: p. 418.

58. Coley, A.M., et al., *Rapid and precise epitope mapping of monoclonal antibodies against Plasmodium falciparum AMA1 by combined phage display of fragments and random peptides.* Protein Eng, 2001. **14**(9): p. 691-8.

59. Coley, A.M., et al., *Structure of the malaria antigen AMA1 in complex with a growth-inhibitory antibody.* PLoS Pathog, 2007. **3**(9): p. 1308-19.

60. Patel, P.N., et al., *Structure-based design of a strain transcending AMA1-RON2L malaria vaccine.* Nat Commun, 2023. **14**(1): p. 5345.

61. Valenciano, A.L., et al., *In vitro models for human malaria: targeting the liver stage.* Trends Parasitol, 2022. **38**(9): p. 758-774.

62. Carias, L.L., et al., *Identification and Characterization of Functional Human Monoclonal Antibodies to Plasmodium vivax Duffy-Binding Protein.* J Immunol, 2019. **202**(9): p. 2648-2660.

63. Persson, K.E., C.T. Lee, K. Marsh, and J.G. Beeson, *Development and optimization of high-throughput methods to measure Plasmodium falciparum-specific growth inhibitory antibodies.* J Clin Microbiol, 2006. **44**(5): p. 1665-73.

64. Ritz, C., F. Baty, J.C. Streibig, and D. Gerhard, *Dose-Response Analysis Using R.* PLoS One, 2015. **10**(12): p. e0146021.

65. Wickham, H. and SpringerLink, *Ggplot2 : elegant graphics for data analysis*. 1st 2009. ed. Use R! 2009, Dordrecht ; New York

New York, NY: Springer

Springer New York.

66. Popovici, J., et al., *Amplification of Duffy binding protein-encoding gene allows Plasmodium vivax to evade host anti-DBP humoral immunity.* Nat Commun, 2020. **11**(1): p. 953.

67. Rangel, G.W., et al., *Enhanced Ex Vivo Plasmodium vivax Intraerythrocytic Enrichment and Maturation for Rapid and Sensitive Parasite Growth Assays.* Antimicrob Agents Chemother, 2018. **62**(4).

68. Ntumngia, F.B., et al., *Genetic variation among Plasmodium vivax isolates adapted to non-human primates and the implication for vaccine development.* Am J Trop Med Hyg, 2009. **80**(2): p. 218-27.

69. Aragao, D.A., J.; Cherukuvada, H.; Clarken, R.; Clift, M.; Philip Cowieson, N.; Jesper Ericsson, D.; Gee, C.L.; Macedo, S.; Mudie, N.; Panjikar, S.; Roy Price, J.; Riboldi-Tunnicliffe, A.; Rostan, R.; Williamson, R.; Tudor Caradoc-Davies, T., *MX2: a high-flux undulator microfocus beamline serving both the chemical and macromolecular crystallography communities at the Australian Synchrotron.* Journal of Synchrotron Radiation, 2018. **25**: p. 885-891.

70. Kabsch, W., *XDS.* Structural Biology, 2010. **66**: p. 125-132.

71. Cassotta, A., et al., *A single T cell epitope drives the neutralizing anti-drug antibody response to natalizumab in multiple sclerosis patients.* Nat Med, 2019. **25**(9): p. 1402-1407.

72. Emsley, P., B. Lohkamp, W.G. Scott, and K. Cowtan, *Features and development of Coot.* Acta Crystallogr D Biol Crystallogr, 2010. **66**(Pt 4): p. 486-501.

73. Afonine, P.V., et al., *Towards automated crystallographic structure refinement with phenix.refine.* Acta Crystallogr D Biol Crystallogr, 2012. **68**(Pt 4): p. 352-67.

Acknowledgments

The work was funded by The National Institutes of Health (R01AI143694) and Veterans Affairs Research Service (I01 BX001350). JGB is supported by a NHMRC Investigator Grant (1173046). WHT is supported by a NHMRC Investigator Grant (2016908). Burnet Institute is supported by

infrastructure grants from the NHMRC and Victorian State Government. We thank Damien Drew for database analysis and designing the PNG16 sequence for expression as a recombinant protein. We thank the individuals who donated their blood for this project and Monash Macromolecular Crystallization Platform (MMCP, Australia) for assistance with setting up the crystallization screens. This research used the MX2 beamline at the Australian Synchrotron, part of ANSTO, and used the Australian Cancer Research Foundation (ACRF) detector. We thank Martin Boulanger for recombinant TgAMA1. We thank the Cytometry & Imaging Microscopy Shared Resource of the Case Comprehensive Cancer Center (P30CA043703 and S10OD021559) for their support. The views are of the authors and not those of NIH and VA Research Service.

Author information

These authors contributed equally: Anna C. Winnicki and Melanie H. Dietrich

These authors jointly supervised the work: Jürgen Bosch and Christopher L. King

Author and Affiliations

Center for Global Health and Diseases, Department of Pathology, Case Western Reserve University, Cleveland, OH, USA
Anna C. Winnicki, Lenore L. Carias, Alyssa N. Malachin, Karli R. Redinger, Olivia McLaine, Yelenna Skomorovska-Prokvolit, Jürgen Bosch, and Christopher L. King

Veterans Affairs Research Service, VA Medical Center, Cleveland, OH
Christopher L. King

InterRayBio LLC
Jürgen Bosch

Malaria Research Unit, Institut Pasteur du Cambodge, Phnom Penh, Cambodia
Jean Popovici, Lionel Brice Feufack-Donfack, and Lea Baldor

Mahidol University, Salaya, Thailand
Wanlapa Roobsoong and Jetsumon Prachumsri

Walter and Eliza Hall Institute of Medical Research, Parkville, Victoria, Australia
Melanie H. Dietrich, Nicolai C. Jung, and Wai-Hong Tham

Department of Medical Biology, The University of Melbourne, Parkville, Victoria, Australia
Melanie H. Dietrich and Wai-Hong Tham

Burnet Institute, Melbourne, Victoria, Australia
Lee M. Yeoh, Chiara L. Drago, D. Herbert Opi, and James G. Beeson

Department of Medicine, The University of Melbourne, Parkville, Victoria, Australia
Lee M. Yeoh

Department of Infectious Diseases, The University of Melbourne, Parkville, Victoria, Australia
D. Herbert Opi and James G. Beeson

Central Clinical School and Department of Microbiology, Monash University, Clayton, Victoria, Australia
D. Herbert Opi and James G. Beeson

Department of Microbiology, Monash University, Clayton, Victoria, Australia.
Chiara L. Drago

Contributions

JB, CLK, JGB, WHT, and JP conceived and designed the experiments. LLC, ANM, KRR, OM, YS-P isolated and generated the humAbs. ACW, WR, LBF-D, LB, CD, AO, and LMY performed the functional experiments. LMY, DHO, JGB, JB, CLK, JP, AO, and ACW conducted data analysis and interpretation. LMY, DHO, and JGB generated and provided key reagents (recombinant proteins, antibodies, RON2 peptides, parasite lines). MHD and NCJ expressed and purified the recombinant proteins required for protein crystallography, performed crystallization experiments, and collected diffraction data at the Australian Synchrotron. MHD determined the crystal structure. All authors contributed to the writing of manuscript and ACW, JB, and CLK completed the first draft of the manuscript. All authors have read and approved the final manuscript.

Figure 1. *Sequence characterization of 12 humAbs and their selectivity, avidity, and affinity towards AMA1*. **A)** Sequences of CDR3 *Iggh* (orange) and corresponding Iggλ or Iggκ (yellow) from individual B cells from which humAbs were generated. The number of somatic hypermutations (SHM) that differ from germline sequences is shown for each clone. **B)** 67 clonal groups were identified, from which clonal groups PvAMA1-specific humAbs were isolated. **C)** HumAb reactivity to PvAMA1_Palo Alto, PvAMA1_PNG16, PkAMA1, PfAMA1_3D7, and TgAMA1 at varying concentrations (1000, 500, 250, 125, 62.5, 31.25, 15.6, 7.8 ng/mL). **D)** HumAb avidity as measured by reduction in binding (MFI) to AMA1 with varying concentrations of NH_4SCN (0.0, 0.9, 1.2, 1.5, 1.8, 2.1, 2.4, 2.7, 3.0, 3.3, 3.6, 4.0 M). HumAbs were used at a concentration of 0.2 µg/mL. **E)** Affinities (k_ds) of PvAMA1-specific humAbs determined using

SPR single-cycle kinetics. HumAbs 800801, 804805, and 808809 affinities could not be

determined (N/A) in this assay. Standard error of the mean was calculated using Prism.

Figure 2. *HumAbs inhibition of blood-stage infection.* **A)** Dose response of PvAMA1-specific

humAbs against Pf-PvAMA1 transgenic parasites. Each symbol represents the average of three

replicates for every concentration. **B)** The mean percentage (+ SEM) of reticulocytes infected

using Pv clinical isolates in short-term invasion inhibition with different humAbs at 100 μg/mL.

Each dot represents a biological replicate from a different clinical isolate. The flow cytometry

background of target cells (reticulocytes) without parasites (mean 9%, range 5-15% invasion)

was subtracted from each experiment. Mouse mAb, 2C3 (100 μg/mL) binds to Duffy antigen on

reticulocytes, thus blocking Pv invasion of reticulocytes (positive control). Only humAb 826827

significantly inhibited reticulocyte invasion compared to the negative control. HumAb 043038

was used as a negative control for experiments represented in panel A and B. **C)** Dose

response of humAb 826827 against four different Pv clinical isolates in short-term invasion inhibition cultures (Isolate 1, 2, 3, and 4 have IC_{50}s of 39.96, 66.78, 25.22, and 61.04 μg/mL respectively. *Note: These are different isolates than those used for Figure 2B*).

Figure 3. *PvAMA1-specific humAbs inhibition of sporozoite invasion of human hepatocyte HCO4 cell line and primary human hepatocytes.* **A)** IC_{50} of different humAbs for Pv sporozoite invasion of human HC04 hepatocytes performed at five concentrations (0.1–1000 µg/mL). Values represent the mean (SEM) of three biological replicates, with each biological replicate performed in duplicate. 043038 was used as a negative control. Murine anti-CSP 2F2 was used as a positive control. **B)** Dose response of 826827 inhibition with 3 different clinical isolates performed in duplicate with IC_{50} of 0.07 (isolate 1), 0.5 (isolate 2), and 1.8 (isolate 3) µg/mL. All dilutions were compared using a multi-variant one-way ANOVA, indicating significant differences in % inhibition at 100 µg/mL and 0.1 µg/mL. **C)** Percent inhibition of Pv sporozoites into human primary hepatocytes using 826827 (white bars) at various concentrations. 043038 (shaded bar) was used as a negative control and was only tested at one concentration. A multi-variant one-way ANOVA and Tukey's secondary test was used to calculate the P-values compared to the negative control using Prism.

Figure 4. *Crystal structure of the PvAMA1–Fab 826827 complex.* **A)** Ribbon representation of PvAMA1–Fab 826827 complex (PDB ID: 8U9D). The heavy chain of 826827 is orange, and the light chain is yellow. PvAMA1 with N-terminal extension (residues 46-62) is white, Domain 1 (residues 63-248) light blue, Domain 2 (residues 249-385) dark blue, and Domain 3 purple (residues 386-474) [50]. Close-up views show interactions between humAb 826827 and (**panel**

i) the mobile Domain 2 loop of PvAMA1 and **(panel ii)** the hydrophobic groove. Due to clarity, only polar interactions between side chains are indicated with dotted lines. **B)** Five CDR loops (L1, L2, H1, H2, and H3) are involved in PvAMA1 binding. PvAMA1 is shown in surface representation and colored as described in panel A. The size of the buried interaction surface on PvAMA1 is indicated. **C)** Comparison of PvAMA1 when bound by CDR-H3 of Fab 826827 or PvRON2 peptide (PDB ID: 5NQG). PvAMA1 is shown in surface representation, with transparency around the Domain 2 loop (dark blue). CDR-H3 and PvRON2 peptide (magenta) form disulfide-linked β-hairpin loops that bind to the hydrophobic groove on Domain 1. HumAb

826827 stabilizes the Domain 2 loop in a closed position on Domain 1 which is dislocated when PvRON2 peptide is bound to PvAMA1. **D)** Ribbon representation of overlaid PvAMA1 structures, comprising our structure of PvAMA1 in complex with Fab 826827 (blue, PDB ID: 8U9D), PvAMA1 in complex with PvRON2 peptide (light blue, PDB ID: 5NQG), and unbound PvAMA1 (grey, PDB ID: 1W8K). This indicates that 826827 stabilizes the Domain 2 loop and allows for the refinement shown. Residues between T296 and F335 are absent in the electron density map of PDB 5NQG and 1W8K but residues 304–327 are well defined in our structure. Overlays were generated using Pymol, and refined root mean square deviation values are indicated. **E)** Top view of the PvAMA1 overlays. **F)** Overlay of PvAMA1 (blue, PDB ID: 8U9D) with PkAMA1 (green, PDB ID: 4UV6). The mobile Domain 2 loops adopt a similar conformation in both structures. **G)** Top view of PvAMA1 and PkAMA1 overlay. (see **Supplemental Table 3** for data collection and refinement statistics).

Figure 5. *Interaction residues of PvAMA1 with RON2 and 826827.* **A)** Sequence alignment of published AMA1 amino acid sequences with the corresponding binding residues for RON2 (magenta) and humAb heavy chain (826; orange), light chain (827; yellow), or both chains (blue) arrows. The first line of the alignment represents the deposited structure 8U9D corresponding to the PvAMA1 Palo Alto strain (ACB42438). Pv is the Sal1 variant (PVX_092275). The red box outlines the Domain 2 loop (304–327), highlighting 826827's interactions with this region of PvAMA1. **B)** Competition assay using 50 μg/mL of PvRON2 (Asp 2050 – Thr 2088) that competed with varying concentrations of humAb 826827 (40–0.039 μg/mL) to bind recombinant PvAMA1. 043038 was used as a negative control.

Figure 6. *Analysis of conservation and common PvAMA1 polymorphisms regarding 826827 binding and activity.* **A)** PvAMA1 residues contacting humAb heavy chain 826 (orange circles), light chain 827 (yellow squares), or both humAb chains (dark blue hexagons) and their mutations are displayed on the X-axis while the Y-axis shows the conservation of that position within 390 clinical isolate sequences. Highlighted amino acids represent those depicted in **Figure 6B**. **B)** Structural analysis of the interaction between 826 (orange) 827 (yellow) and PvAMA1 (blue) shows three observed polymorphisms in the binding epitope (G117R, N132D, and N130K; pink). **C)** The *pvama1* gene was sequenced in the seven Pv clinical isolates (**Supplemental Table 2**). Amino acids 117, 130, and 132 represent polymorphic contact residues of humAb 826827 to PvAMA1 Sal1 reference strain (PVX_092275). Bars represent the mean (+/- SEM) of humAb invasion inhibition of WT compared to an isolate expressing a SNP at each of the three polymorphic residues.

A.

HumAb	Heavy V-GENE & allele	SHM	Heavy Chain CDR3	Light V-GENE & allele	SHM	Light Chain CDR3	Clones in group
800801	IGHV1-46	15	CARVYSSGWPPTKAFDPW	IGLV3-9	9	CQLWDSSTSGWVF	1
804805	IGHV3-30-5	21	CAKELYAYSTSPLDNW	IGKV1-5	7	CQEYNTYAISF	1
806807	IGHV4-31	30	CAREDRNYGKSCFDYW	IGKV1-39	9	CQQSYSAPRTF	2
808809	IGHV3-33	6	CARDGLGYCLGGRCYPSLDYW	IGLV1-44	8	CAAWDDSLNGYVF	2
810811	IGHV3-15	21	CATDLVGATGYYGLDVW	IGLV3-9	17	CQLWDSSTSSWVF	3
814815	IGHV3-23	16	CARDLAPPEFWSSYSYFGLGVW	IGKV3-11	8	CQQLSNWPPLTF	6
816817	IGHV4-59	24	CARETLNQRRFDLW	IGKV1-39	18	CQQSYSIPRTF	2
826827	IGHV4-61	24	CARSRGEGYCSFGTCYTLFYYFDYW	IGKV3-15	18	CQQYNEWPPRPTF	11
828829	IGHV4-39	9	CARRSAEGYCSGGSCYSLWGVGFGPW	IGKV3-15	0	CQQYNNWPPITF	1
830831	IGHV4-31	19	CARSVGDGYCSGGACFFLYYTGLDVW	IGKV3-15	7	CQQYSKWPPLTF	1
832833	IGHV1-3	20	CAREYCDTGRCLAGVVVIPAEAFDIW	IGKV3-20	17	CQQYGSSQWTF	1
838839	IGHV5-10-1	13	CARITLTWGGTFDFW	IGLV3-25	13	CQTSHSLAAYVVF	4

B. Clonal Group Distribution

# of Isolates: 169

C.

D.

E.

	k_d (nM)	Std Error (nM)
830831	10.73	9.10
810811	11.33	1.04
806807	12.01	9.78
816817	15.30	7.19
828829	15.89	4.72
832833	21.44	8.90
838839	21.96	13.75
826827	30.53	10.23
814815	47.66	3.63
PvRON2L	50.00	N/A
800801	N/A	N/A
804805	N/A	N/A
808809	N/A	N/A

A 826827

i

ii

B

C PvRON2

D PvAMA1 (r.m.s.d: 0.4-0.5 Å)

PDB ID 8U9D
PDB ID 5NQG
PDB ID 1W8K

E PvAMA1

F PvAMA1 vs PkAMA1 (r.m.s.d: 0.5 Å)

PDB ID 8U9D
PDB ID 4UV6

G PvAMA1 vs PkAMA1

HumAb	Heavy V-GENE & allele	SHM	Heavy Chain CDR3	Light V-GENE & allele	SHM	Light Chain CDR3	Clones in group
800801	IGHV1-46	15	CARVYSSGWPPTKAFDPW	IGLV3-9	9	CQLWDSSTSGWVF	1
804805	IGHV3-30-5	21	CAKELYAYSTSPLDNW	IGKV1-5	7	CQEYNTYAISF	1
806807	IGHV4-31	30	CAREDRNYGKSCFDYW	IGKV1-39	9	CQQSYSAPRTF	2
808809	IGHV3-33	6	CARDGLGYCLGGRCYPSLDYW	IGLV1-44	8	CAAWDDSLNGYVF	2
810811	IGHV3-15	21	CATDLVGATGYYGLDVW	IGLV3-9	17	CQLWDSSTSSWVF	3
814815	IGHV3-23	16	CARDLAPPEFWSSYSYFGLGVW	IGKV3-11	8	CQQLSNWPPLTF	6
816817	IGHV4-59	24	CARETLNQRRFDLW	IGKV1-39	18	CQQSYSIPRTF	2
826827	IGHV4-61	24	CARSRGEGYCSFGTCYTLFYYFDYW	IGKV3-15	18	CQQYNEWPPRPTF	11
828829	IGHV4-39	9	CARRSAEGYCSGGSCYSLWGVGFGPW	IGKV3-15	0	CQQYNNWPPITF	1
830831	IGHV4-31	19	CARSVGDGYCSGGACFFLYYTGLDVW	IGKV3-15	7	CQQYSKWPPLTF	1
832833	IGHV1-3	20	CAREYCDTGRCLAGVVVIPAEAFDIW	IGKV3-20	17	CQQYGSSQWTF	1
838839	IGHV5-10-1	13	CARITLTWGGTFDFW	IGLV3-25	13	CQTSHSLAAYVVF	4

B. Clonal Group Distribution

# of Isolates: 169

10.73	9.10
11.33	1.04
12.01	9.78
15.30	7.19
15.89	4.72
21.44	8.90
21.96	13.75
30.53	10.23
47.66	3.63
50.00	N/A
N/A	N/A
N/A	N/A
N/A	N/A

A 826827

C_L
C_H
V_H
PvAMA1
V_L
mobile loop
domain 2
domain 1
domain 3

i
Y117
Y49
R54
Y34
R101
E103
Y54
Y55
Y105
D118
T116
N315
N316
E83
K321

ii
R101
F115
E103
L114
N132
Y112
H134
Y105
R146
P129
F128
F169
R146
Y196
H181
F108
S168
Y147

B
L1
L2
H2
H3
mobile loop
H1
1392.5 Å²

C
H3
PvRON2
mobile loop
hydrophobic groove

D PvAMA1
(r.m.s.d: 0.4-0.5 Å)
domain 2 loop
T296
F335

PDB ID 8U9D
PDB ID 5NQG
PDB ID 1W8K

E PvAMA1
T296
F335
domain 2 loop

F PvAMA1 vs PkAMA1
(r.m.s.d: 0.5 Å)
domain 2 loop

PDB ID 8U9D
PDB ID 4UV6

G PvAMA1 vs PkAMA1
domain 2 loop

A.