Synchronous Ensembles of Hippocampal CA1 Pyramidal Neurons Associated with Theta but not Ripple Oscillations During Novel Exploration

En-Li Chen[1,2], Tsai-Wen Chen[1,2], Eric R. Schreiter[3], Bei-Jung Lin[1,2*]

[1]Institute of Neuroscience, National Yang Ming Chiao Tung University, Taipei, Taiwan
[2]Brain Research Center, National Yang Ming Chiao Tung University, Taipei, Taiwan
[3]Janelia Research Campus, Howard Hughes Medical Institute, Ashburn, VA 20147, USA.
*Corresponding author: beijunglin@nycu.edu.tw

Abstract

Synchronous neuronal ensembles play a pivotal role in the consolidation of long-term memory in the hippocampus. However, their organization during the acquisition of spatial memory remains less clear. In this study, we used neuronal population voltage imaging to investigate the synchronization patterns of CA1 pyramidal neuronal ensembles during the exploration of a new environment, a critical phase for spatial memory acquisition. We found synchronous ensembles comprising approximately 40% of CA1 pyramidal neurons, firing simultaneously in brief windows (~25ms) during immobility and locomotion in novel exploration. Notably, these synchronous ensembles were not associated with contralateral ripple oscillations but were instead phase-locked to theta waves recorded in the contralateral CA1 region. Moreover, the subthreshold membrane potentials of neurons exhibited coherent intracellular theta oscillations with a depolarizing peak at the moment of synchrony. Among newly formed place cells, pairs with more robust synchronization during locomotion displayed more distinct place-specific activities. These findings underscore the role of synchronous ensembles in coordinating place cells of different place fields.

Introduction

The establishment of long-term memory involves the initial acquisition and subsequent consolidation of memory traces, processes intricately connected to the modification of synaptic transmission[1,2]. Synchronous ensembles, characterized by the simultaneous firing of multiple neurons over tens of milliseconds (referred to as population synchrony), play a pivotal role in influencing synaptic plasticity[3-5]. The hippocampus, a crucial region for long-term memory formation[6,7], has been a focal point of synchronous ensemble studies. Within the hippocampus, synchronous ensembles are hypothesized to play roles in reactivating and consolidating labile memory traces[8]. However, their organization during memory acquisition remains less understood.

In the hippocampus, synchronous ensembles among CA1 pyramidal cells (CA1PCs) are predominantly observed during sharp-wave ripples, the brain waves associated with memory consolidation[8-13]. During these events, 10-18% of CA1PCs generate spikes, particularly during offline consummatory behavioral states or sleep[14,15]. These synchronous ensembles are biased toward experience-dependent reactivation, consolidating labile memory traces[16,17]. Furthermore, synchronous ensembles firing action potentials in short windows of 10-30 milliseconds enhance information processing[6] and discriminate distinct behavioral contingencies[18]. Although synchronous ensembles of CA1PCs during offline memory consolidation are well-characterized, their organization during online acquisition of spatial memory, such as when animals enter a new environment and new place cells form, remains unclear.

Previous studies have identified correlated activities among place cells during theta oscillation, where 1-3% of place cells' spikes co-activate in a forward order matching animals' trajectory in the theta cycles when animals' locations overlap in the place fields of these cells[19-21]. Because these place cells fire at distinct theta phases in the same cycle, this coordinated firing operates on the timescale of theta periods (~125ms)[22,23]. Population synchrony on shorter timescales (10-30 ms) is

speculated to be reduced by the presence of theta oscillations[22].

The subthreshold membrane voltage (subVm), influenced by synaptic inputs and intrinsic

conductances, directly triggers spiking activity[24,25]. Additionally, properties of subVm support the

organization of neuronal representation[26]. Although correlated membrane potentials have been

proposed to underlie the synchrony of neuronal ensembles[27,28], multicellular recordings of subVms

have not yet provided direct experimental evidence. Given the intricate interaction among subVm,

spiking, and place cell formation, it is crucial to investigate the temporal relationships of sub- and

suprathreshold neuronal activities among CA1PCs during novel exploration.

In this study, we used voltage imaging to investigate the temporal dynamics of sub- and

suprathreshold neuronal activities in CA1PCs during novel exploration. Additionally, we implanted

electrodes in the contralateral CA1 region to monitor theta and ripple oscillations, which are known

to co-occur across hemispheres[29-31]. We found synchronous ensembles involving approximately 40%

of CA1PCs exhibited transient population synchrony and correlated subthreshold membrane

potentials when mice ran and stopped in a new maze. Surprisingly, these synchronous ensembles

occurred outside of contralateral ripples and were phase-locked to intracellular theta oscillations as

well as extracellular theta oscillations recorded from the contralateral CA1 region. Notably, cell pairs

with stronger synchronous activities established more distinct place fields that overlapped less. In

essence, synchronous ensembles facilitated temporal association among place cells representing

diverse features of spatial memory. This plastic network, driven by synchronous ensembles, may

contribute to the stabilization of place cells, establishing place fields that effectively tile the

environment.

Results

To investigate the temporal relationships between neuronal activities during novel exploration, we

simultaneously recorded many neurons as animals explored a new environment for the first time.

To achieve this, we employed voltage imaging in conjunction with an air-lifted plastic track that

allowed mice to explore an environment while remaining head-fixed under a microscope (Figure

1A). To ensure a genuinely novel experience, all pre-training and habituation were performed in a

separate room distinct from the one where the imaging experiments were conducted. On the day of

imaging, the mice were introduced to the recording room for the first time and initially confined to

a specific corner of the track (Figure 1A top view). Subsequently, as the imaging session

commenced, the confinement was lifted, granting mice the freedom to explore the track while

imaging was conducted.

In novel exploration, mice moved at a speed of 8.3±0.9 cm/s during locomotion (speed>3 cm/s) and

spent 53±7% and 36±6% of their time in locomotion and immobility (speed<1 cm/s), respectively

(n=5 mice). Meanwhile, they completed 10±2.5 laps during the 180-s imaging periods (n=5 mice).

For voltage imaging, we expressed the voltage indicator Voltron2[32] in CA1 pyramidal cells (CA1PCs)

that were born on embryonic day (E) 14.5, allowing us to measure both suprathreshold and

subthreshold membrane potentials from multiple CA1PCs (14±4 cells imaged per field of view

(mean±s.d.) and, in total, 71 cells imaged from 5 fields of view in 5 mice; Figure 1B, Supplementary

Figure 1A, 1B). Simultaneously, we monitored the animals' positions and speed within the track

while recording the local field potential (LFP) from the contralateral side of the hippocampus.

Consistent with previous studies, neurons labeled on E14.5 located more on the deep side of the

pyramidal layer than those labeled on E17.5 ($t_{(601)}$=22.8, p<0.0001, Student's t-test; Supplementary

Figure 1C, D)[33-37].

The mean firing rates of cells ranged from 2.3 to 4.3 Hz (25th-75th percentiles) with a median of 3.2

Hz (n=71 cells, Supplementary Figure 2A). Additionally, most CA1PCs exhibited elevated mean firing

rates during immobility relative to locomotion, and their firing rates showed a negative correlation

with the animals' speed (Supplementary Figure 2B, 2C), consistent with previous research[38].

Synchronous ensembles among many CA1 pyramidal neurons

Figure 1C shows the fluorescence traces of all neurons in a session as an example. Instances of

synchronous activity, where many neurons emitted spikes within a narrow time window, were

frequently observed (Figure 1C). To investigate synchronous activity within the neuronal population,

we selected a reference cell and counted the number of spikes across all other cells that were

aligned with the spikes of the reference cell. This approach yielded a grand average cross-

correlogram (CCG) capable of elucidating the timing of spikes from the reference cell relative to the

combined spikes of the rest of the population. Notably, the grand average CCG exhibited heightened

spike counts around zero lags, unlike flattened counts calculated from randomly selected cells of

different sessions (Figure 1D and 1E). The full width at half maximum (FWHM) of the peak in the

grand average CCG measured 23±14 ms (mean±s.d., n=71 cells from 5 sessions; Figure 1F),

indicating the timescale of the population synchrony. Furthermore, pairwise CCGs of many cell pairs

showed significant peaks compared to jittered data (percentage of significant pairs: 63% (median)

n=497 pairs, Figure 1-figure supplement 1A and 1B). Among pairs with significant peaks, the

absolute values of the peak lags ranged from 0.8 to 4.3 ms (25th-75th percentiles) with a median of

2.1 ms, and the FWHM of the peaks ranged from 20 to 46 ms (25th-75th percentiles) with a median

of 30 ms (n=315 pairs, Figure 1-figure supplement 1C and 1D). These results highlight the

synchronous spiking between neurons on the millisecond timescale.

To systematically reveal individual synchronous events, we counted the number of spikes from all

simultaneously recorded neurons within sliding windows of 25 ms. To compare the summed spikes

with controls, we randomly jittered the spike trains to perturb spike timings while maintaining spike

rates. We identified moments of population synchrony when spike sums exceeded the average of

the jittered spike sums plus four standard deviations (Figure 1C). Using the same criterion for event

detection, we observed over a 7.8-fold increase in synchronous events in the original data

compared to jittered controls (mean event rate: 0.94±0.17 Hz for the original data, 0.12±0.01 Hz for

the jittered data, *n*=10 segments, *p*=0.002, *W*=55, Wilcoxon signed-rank test; Figure 1G). The

observed population synchrony was not attributable to spikes in complex bursts, as synchronous

event rates did not differ significantly with or without the inclusion of later spikes in bursts

(Supplementary Figure 3). Furthermore, we calculated the percentage of neurons participating in

these synchronous events to examine the synchronous ensemble sizes. The ensemble sizes ranged

from 32% to 53% (25th-75th percentiles) with a median of 42%, which were larger than the

synchronous ensemble sizes associated with ripples[14] (Figure 1H). These results demonstrate

synchronous ensembles of CA1PCs engaging many neurons.

Figure 1. Population synchrony during novel exploration.

(A) Schematic of the behavioral setup for voltage imaging. Mice explored a novel one-way track

while fluorescence images were captured at the CA1 pyramidal layer. Left: side view; right: top view

of the setup.

(B) Example time-averaged image of CA1 pyramidal cells expressing Voltron2-ST labeled with JF_{552}-

HaloTag ligand. Scale bar, 100 μm.

(C) Fluorescence traces of cells in an example session with identified synchronous ensembles

(labeled with gray vertical bars). Right: an example of magnified fluorescence traces within the

bracket, pinpointing the occurrence of the identified synchronous ensemble. Spikes are indicated by

dots on the top.

(D) The grand average cross-correlogram (CCG) averaged across all 71 cells. The gray line represents

the mean grand average CCG between reference cells and randomly selected cells from different

sessions.

(E) Distribution of the grand average CCG peak lags of all cells (*n*=71).

(F) Distribution of the full width at half maximum (FWHM) of the grand average CCGs (*n*=71 cells).

(G) Pairwise comparison of the event rates of population synchrony between original and jittered

data. Bar heights indicate group means. **$p<0.01$

(**H**) Histogram of the ensemble sizes as percentages of cells participating in the synchronous

ensembles.

Synchronous ensembles during locomotion and immobility of novel exploration

The hippocampus exhibits state-dependent activities when rodents are engaged in locomotion and

immobility[38,39]. In novel exploration, CA1PCs showed synchronous spiking during locomotion and

immobility (Figure 2A). To investigate whether these synchronous activities differed between these

behavioral states, we compared grant average CCGs and synchronous ensembles under these

conditions. During both immobility and locomotion, grant average CCGs showed prominent peaks

compared to jittered data (Figure 2B). Moreover, synchronous ensembles were rich in both

immobility and locomotion periods (Figure 2A, 2B and 2D). Nevertheless, the FWHMs of grand

average CCGs were broader in the immobility group compared to the locomotion group (Figure 2B

and 2C). Synchronous events occurred more frequently during immobility compared to locomotion

(Figure 2D). Additionally, ensemble sizes were significantly larger in the immobility group compared

to the locomotion group (Figure 2E). Taken together, with differences in sizes and kinetics, there

were synchronous ensembles during both immobility and locomotion periods of novel exploration.

A

LFP ⌇0.2 mV

Cell

Speed
0

3AP⌇
80ms

3cm/s⌇

LFP ⌇0.1 mV
0.3 s

LFP ⌇0.1 mV
0.3 s

LFP ⌇0.2mV

Cell

Speed
0

3AP⌇
80ms

5cm/s⌇

B
Normalize spike count
2
1.5
1
-100 0 100
2.5
2
1
-100 0 100
Lag (ms)

C
FWHM (ms)
60
40
20
0
Immo. Loco.
***

D
Event rate (Hz)
2
1
Immo. Loco.
**

E
Ensemble size (percent of cells)
60
40
20
Immo. Loco.
***

Figure 2. Comparison of synchronous ensembles between immobility and locomotion.

(A) Example traces of cells in a session with identified synchronous ensembles (gray vertical bars)

during periods of immobility (red rectangular frame) and locomotion (blue rectangular frame).

Spikes are indicated by dots at the top of individual traces. Top: the simultaneously recorded LFP

trace. Bottom: simultaneously recorded speed of the animal. Inset: the band-pass filtered LFP trace

at the theta frequency range (black line) overlaid on the raw LFP trace (gray line).

(B) The grand average cross-correlograms (CCGs) during immobility (red) and locomotion (blue)

averaged across all cells (n=71). The gray lines represent the mean grand average CCGs calculated

from jittered data.

(C) Pairwise comparison of peak widths (FWHM) in the grand average CCGs between immobility

(red) and locomotion (blue). Vertical bars represent means and standard deviations for both groups

(FWHM: 27±12 ms for immobility, 18±9 ms for locomotion, mean±s.d., $t_{(70)}$=5.45, $p<0.001$, paired t-

test, n=71 cells). ***$p<0.001$

(D) Pairwise comparison of event rates of population synchrony during immobility and locomotion.

Bar heights indicate group means (immobility: 1.4±0.7 Hz, locomotion: 0.6±0.5 Hz, mean±s.d., n=10

segments, W=55, p=0.002, Wilcoxon signed-rank test). **$p<0.01$

(E) Boxplot of ensemble sizes for synchronous ensembles occurring during immobility (red) and

locomotion (blue) (median ensemble size: 47% for immobility, n=446 events, 32% for locomotion,

n=313 events, $t_{(757)}$=13.54, $p<0.001$, Student's t-test). ***$p<0.001$

198
199

Synchronous ensembles occur outside ripple episodes

CA1PCs exhibit synchronous activities when the hippocampal network emits ripple waves[8,14]. To test

whether there is a co-occurrence of synchronous ensembles and contralateral ripples, we first

detected ripples from LFP recordings. Ripples were detected in three of the five recording sessions,

and all occurred during immobility (mean ripple rate: 0.04±0.02 Hz during immobility, 0 Hz during

locomotion). In the two recording sessions where we did not detect any ripples, we verified the

recording quality of the LFP signal by running multiple tests after the recording sessions. Ripples

were identified in the same animals during quality tests, indicating that the absence of ripples

during novel immobility in the recording session was not due to the deterioration of LFP recordings

(Figure 3-figure supplement 1).

To visualize both ripples and synchronous ensembles, we plotted voltage traces of neurons

alongside simultaneously recorded ripple oscillations. It was typical for synchronous ensembles to

occur far away from ripple episodes (Figure 3A). To delve into the relationship between neuronal

activities and ripples, we aligned traces of ripple power with population synchrony and ripples. In a

recording session where 64 synchronous ensembles and 12 ripples were detected during

immobility, the LFP power at ripple frequency (120-240 Hz) exhibited no discernable peaks within

any of the synchronous events (Figure 3B, left upper and middle panels). Nevertheless, the same

LFP power displayed substantial peaks during every ripple event (Figure 3B, right upper and middle

panels). A striking contrast between synchronous events and ripples was also evident when spiking

probabilities during these events were examined. During population synchrony, the mean spiking

probability averaged across all cells was high, while the spiking probability was almost zero during

ripple events (Figure 3B lower panel). In the three sessions where ripples were identified, we

calculated the percentages of ripple events coinciding with synchronous ensembles and vice versa.

Remarkably, there was no co-occurrence of ripples and synchronous ensembles (Figure 3C).

Furthermore, we quantified ripple-modulated spiking by calculating the ratio of the difference in

mean firing rates during in-ripple and out-ripple periods to the sum of the mean firing rates during

both periods. Indexes of most cells were negative, and a striking 85% of cells displayed ripple

modulation indexes of -1, indicating complete suppression of their spiking activities during ripples

(Figure 3D). Taken together, during novel exploration, synchronous ensembles involved many

neurons with millisecond-synchronized spikes and occurred outside the time frames of ripples.

Figure 3. Synchronous ensembles occur outside ripple episodes.

(A) Example traces with ripples and synchronous ensembles. Top: simultaneously recorded LFP

displaying an episode of ripple oscillation. A red vertical line indicates the timing of the peak ripple

power in the LFP trace. A yellow vertical bar marks the duration of the ripple episode. Inset at the

top: a magnified view of the ripple oscillation. Middle: fluorescence traces of 9 CA1PCs. Gray

vertical bars mark the timings of the synchronous ensembles on all traces. Bottom: animals' speed

recorded simultaneously with voltage imaging.

(B) LFP ripple power and spiking probability aligned to synchrony ensembles and ripple events.

Upper panel: color-coded LFP power at ripple frequency (120-240Hz) aligned to the timings of 64

synchronous ensembles (left) and 12 ripple events (right). Middle panel: Average LFP power of

ripple frequency aligned to the timings of the synchronous ensembles (left) and the ripple events

(right). Lower panel: Average spike counts of all cells aligned to the timings of the synchronous

ensembles (left) and ripple events (right).

(C) Percentages of co-occurrences between population synchrony and ripples. The co-occurrences

in the "sync to ripples" group represent the percentages of ripple episodes co-occurring with

synchronous ensembles in the same episodes. The co-occurrences in the "ripples to sync" group

indicate the percentages of synchronous ensembles co-occurring with ripple episodes. Each gray

circle represents an individual session.

(D) Distribution of the ripple modulation indexes (n=41 cells). Gray bar: modulation indexes with

significant p values (n=35 cells); white bars: modulation indexes without significance (n=6 cells).

It was puzzling that these CA1PCs exhibited robust spiking activities outside of ripples yet generated

few spikes during ripples. To further investigate neuronal activities during ripples, we established a

recording condition that allowed us to capture more ripple episodes. Specifically, we immobilized

mice in a tube to promote behaviors favoring ripple generation. The mice were habituated to head

fixation in a tube in a room distinct from the one where imaging experiments were conducted. On

the imaging day, the mice were introduced to the recording room and head-fixed under the

microscope for the first time.

CA1PCs were labeled *in utero* on embryonic day (E) 14.5 (n=56 cells from 3 sessions in 3 mice) and

E17.5 (n=55 cells from 3 sessions in 3 mice) and imaged in adult brains. Both neuronal populations

exhibited prominent peaks in their grand average CCGs and significantly higher synchronous event

rates compared to jittered data (Figure 3-figure supplement 2A, B). Approximately 40% of the

recorded neurons participated in synchronous ensembles, indicating robust synchronous activity

involving a substantial proportion of the recorded cells (Figure 3-figure supplement 2C).

In total, 1052 synchronous ensembles and 174 ripple episodes were detected across these imaging

sessions. Consistent with findings from walking animals, few synchronous ensembles occurred

during ripples when animals were immobilized in a tube (Figure 3-figure supplement 3A, B).

Moreover, no distinguishable ripple oscillations were observed in synchronous events, and the

average firing rates during ripple episodes were near zero (Figure 3-figure supplement 3C, D). At the

single-cell level, 90% of neurons showed significant negative spiking modulation during ripples, with

ripple modulation indexes close to -1, indicating strong suppression of spiking (Figure 4Ai). This

suppression extended to subthreshold membrane potentials, as nearly all cells exhibited decreased

fluorescence during ripples compared to baseline (Figure 4Bi, Ci). These results demonstrate that

spiking activity and subthreshold membrane potentials are robustly suppressed during ripples.

Contextual novelty plays a critical role in shaping hippocampal neuronal activities[40-46]. To assess its

influence, we trained mice to become familiar with the imaging procedure and the recording

environment over five days and recorded CA1PC activities on the final day. Mean firing rates were

significantly reduced in the familiar group compared to the novel group (Familiar group: 1.1 to 5.2

Hz (25th-75th percentiles), median=2.3 Hz, n=66 cells, 6 sessions, 4 mice; Novel group: 1.7 to 6.0 Hz

(25th-75th percentiles), median=4.2 Hz, n=111 cells, 6 sessions, 6 mice, p=0.0083, Wilcoxon signed-

rank test). Additionally, 15% of the neurons in the familiar group exhibited significantly positive

spiking modulation by ripples, while fewer cells showed negative modulation compared to the novel

group (Figure 4A). During ripples, neurons in the novel group predominantly displayed

hyperpolarizing membrane voltage responses, whereas a subset of neurons in the familiar group

exhibited prominent depolarizing responses (Figure 4B). The mean fluorescence changes in the

familiar group shifted toward depolarization compared to the novel group (Figure 4C). Finally,

synchronous event frequencies were significantly lower in the familiar context, indicating weaker

synchronous activities under familiar conditions (Figure 4D). These results demonstrate that

hippocampal neuronal activities, particularly synchronous ensembles, are strongly influenced by

contextual novelty.

Figure 4. Comparison of ripple-associated neuronal activities and synchronous event rates

between novel and familiar recording contexts.

(A) Histograms of ripple modulation indices for the novel (Ai) and familiar (Aii) recording contexts.

Gray bars represent modulation indices with significant *p*-values, while white bars indicate non-

significant modulation indices.

(B) Color-coded representations of mean fluorescence changes relative to baseline fluorescence for

individual neurons, averaged across ripple events, in the novel (Bi) and familiar (Bii) recording

contexts.

(C) Histograms of mean fluorescence changes within a $\pm$10-ms window centered on the peak

power of ripple events. Ci corresponds to the novel context, while Cii corresponds to the familiar

context.

(D) Boxplots of synchronous event rates for the novel and familiar recording contexts. Mean event

rates were 0.98±0.16 Hz for the novel context and 0.13±0.04 Hz for the familiar context (t(22)=5.20,

p<0.001, Student's t-test). ***p<0.001

Synchronous ensembles are associated with theta oscillations

To compare the network dynamics associated with synchronous ensembles during immobility and

locomotion, we aligned LFP traces triggered to the timings of synchronous ensembles. The average

LFP traces of a representative session showed prominent theta oscillations around the timings of

both immobility and locomotion population synchrony (Figure 5A). Spectral analysis of the mean

synchrony-triggered LFP traces during immobility and locomotion revealed theta frequency

components of 4-12 Hz, with a slightly higher peak frequency in locomotion LFP compared to

immobility LFP (Figure 5B). Furthermore, we computed LFP theta modulation and identified

preferred phases of synchronous ensembles. In line with the previous result, synchronous

ensembles displayed a high level of modulation by LFP theta oscillation (modulation strength: 0.61$\pm$

0.05 during immobility and 0.74$\pm$0.04 during locomotion, n=5 sessions). Although the preferred

phases varied from session to session due to differences in recording sites along the proximal-distal

axis of the hippocampus, the timings of individual ensembles were consistently locked to the

preferred phase of each session (Figure 5C). These results establish a strong link between

synchronous ensembles and LFP theta oscillation.

Subthreshold membrane voltage (subVm) is one of the critical parameters supporting the

generation of action potentials. To investigate the subthreshold signals underlying population

synchrony, we aligned subVm traces with the timings of synchronous ensembles and observed that

both immobility and locomotion synchrony rode on a subVm waveform oscillating at theta

frequency (Figure 5D). To elucidate further the relationship between subVm theta oscillation and

spikes involved in population synchrony, we calculated the subVm theta modulation and preferred

phases of spikes. Spikes participating in both immobility and locomotion synchrony were strongly

coupled to the subVm theta oscillation of neurons, with the mean preferred phase being in the

latter half of the rising phase closer to subVm theta peaks (Figure 5E). Conversely, spikes not

participating in synchronous ensembles had weaker modulation strength compared to spikes

involved in synchrony (Figure 5-figure supplement 1A, B). Furthermore, the membrane voltages

triggered by these two categories of spikes showed a significant difference in the hyperpolarization

phase of the theta oscillation. Synchronous spikes were situated within cycles of theta oscillations

featuring more hyperpolarized troughs than spikes outside synchronous ensembles (Figure 5-figure

supplement 1C, D). Thus, compared to spikes not participating in synchronous ensembles,

synchronous spikes showed a more robust modulation by theta oscillation of membrane voltage

during immobility and locomotion in the context of novel exploration.

Notably, phase locking of spikes to their subVm theta oscillation and the synchronous spikes among

many neurons suggest that subVms of different neurons would be correlated at the theta frequency.

To explore the possibility, we divided the 180-second subVm traces into 1-second segments with

50% overlap and calculated the cross-correlation and magnitude-squared coherence function for

each pair of cell segments. Segments were sorted into the immobility or locomotion group based on

the animals' speed during each segment (kept below 1 cm/s for immobility and above 3 cm/s for

locomotion for more than 90% of the time). In total, we analyzed 511 cell pairs from 5 sessions.

Indeed, the cross-correlation of subVm between cell pairs revealed a high correlation at zero lag and

periodic modulation of the correlation at theta frequency (Figure 5F). Furthermore, the subVm

theta coherences were increased in both groups (immobility: 0.50, locomotion: 0.40, median, n=511

pairs) and were negatively correlated with soma distances (immobility: -0.46, locomotion: -0.48,

n=511 pairs, Figure 5G). Taken together, during novel exploration, CA1PCs generate synchronous

activity associated with theta oscillation and exhibit correlated and theta-coherent subVms.

Correlated intracellular theta and theta-phase locking of the synchronous ensembles raise the

question of whether population synchrony among CA1PCs extends beyond synchrony derived from

these effects. To address this, we analyzed population synchrony after randomizing the theta cycles

during which neurons spiked, while keeping their theta phases unchanged. Supplementary Figure 4

illustrates a significant reduction in synchronous event rates following theta cycle randomization.

The finding indicates spiking at specific theta cycles plays a major role in driving population

synchrony.

To further investigate synchronous ensembles across different datasets, we analyzed publicly

available hippocampal recordings 'hc-11' from the CRCNS repository[46,47], where rats navigated novel

mazes for water rewards (see Method). Using our algorithm, we identified a significant number of

synchronous ensembles during the first three minutes of novel navigation. On average, the rates of

synchronous events were 6.4-fold higher than those detected in jittered controls (mean event rate:

2.0 ± 0.3 Hz for the original data vs. 0.32 ± 0.03 Hz for jittered data, n = 8 sessions, p = 0.0078, W =

36, Wilcoxon signed-rank test; Supplementary Figure 5A). To assess whether ripple oscillations were

associated with these synchronous ensembles, we analyzed ripple event rates and their relationship

to population synchrony. During this period, ripple events were infrequent (mean ripple rate: 0.02 ±

0.01, n = 8 sessions), and ripple power peaked during ripple episodes but remained low at the

timings of population synchrony (Supplementary Figure 5B). Nevertheless, LFP traces aligned to

population synchrony revealed prominent theta oscillations (Supplementary Figure 5C).

Synchronous ensembles were modulated by LFP theta oscillation (modulation strength: 0.30 ± 0.04,

n = 8 sessions, $p < 0.001$), and the timings of individual ensembles were consistently locked to the

preferred phase of each session, suggesting a functional coupling of synchronous ensembles to

theta oscillations important for information processing (Supplementary Figure 5D).

Figure 5. Theta oscillations in the LFP and subthreshold membrane potential were linked to

population synchrony.

(A) LFP waveforms triggered by timings of synchronous ensembles. Red: triggered by timings of

immobility synchrony; blue: triggered by timings of locomotion synchrony.

(B) Power spectral densities of the triggered LFP waveforms. Red: immobility; blue: locomotion.

(C) Distributions of the differences between LFP theta phases of individual synchronous ensembles and their preferred phases of the session. Note that most ensembles show minor differences within ±30 degrees. Red: immobility; blue: locomotion.

(D) Fluorescence traces triggered by timings of synchronous ensembles. Traces with spikes are excluded. Left panel: traces triggered by timings of immobility synchrony. Right panel: traces triggered by timings of locomotion synchrony. Thin gray lines: mean triggered fluorescence traces averaged across triggers for representative cells. Thick red and blue lines: mean triggered fluorescence traces averaged across cells.

(E) Polar plot illustrating theta modulation of spikes participating in the immobility synchrony (red dots) and of spikes participating in the locomotion synchrony (blue dots). The angle of the dots indicates the preferred phase, and the distance from the center of the circle represents the modulation strength. Each dot represents the averages from a neuron.

(F) Pairwise cross-correlation of the subthreshold membrane voltages (subVm). Top: color-coded cross-correlation of subVm between cell pairs during immobility and locomotion periods. Bottom: averaged cross-correlation over all cell pairs for immobility (red) and locomotion (blue) subVm. Both show a central peak at zero lag flanked by theta oscillation.

(G) Scatter plots of theta coherence against soma distances of immobility subVm (red) and locomotion subVm (blue) between cell pairs.

Synchrony between place cells with distinct place tunings

When animals explore a novel maze, spatially tuned spiking activity can form within a few minutes[48,49]. However, how synchronous ensembles organize their spatially tuned activities during novel exploration remains to be determined. Synchronous ensembles, generating spikes coincidently in short intervals, might display correlated spiking at longer time windows that match behavioral timescales of exploration, potentially leading to shared place responses. On the other hand, CA1PCs are interconnected through inhibitory interneurons[50-52]. Modulation of activities in inhibitory interneurons by some place cells may influence the activities of other place cells, resulting in divergent spatial tunings[48,53,54].

To compare the spatial tunings of synchronous cell pairs, we first calculated the spatial tuning curves of individual neurons by computing the mean firing rates in every spatial bin. On average, 40% of cells displayed selective spatial tunings of their spiking activities and were qualified as place cells[55] (Figure 6A). Among place cell pairs, some showed similar spatial tunings, while others did not (Figure 6B). For instance, in an illustrative session depicted in Figure 6B, the seventh and eighth place cells had highly similar spatial tunings with a correlation coefficient as high as 0.7. In contrast, the fifth and tenth place cells exhibited vastly different spatial tunings with a correlation coefficient as low as -0.5.

We further explored the relationship between the similarity of spatial tunings and synchronization strength among pairs of place cells. We measured synchronization strength as the peak of the CCG normalized by the averaged jittered spike count within the same bin as the CCG peak. We segregated spikes based on these behavioral states to accommodate differences between locomotion and immobility and calculated the corresponding CCG and synchronization strength.

During locomotion, we observed a robust negative correlation between synchronization strength

and the similarity of spatial tunings among cell pairs. In other words, cell pairs with more precisely

co-active spikes during locomotion exhibited more distinct place-tuning profiles (Figure 6C, D). This

anti-correlation between synchronization strength and place tunings indicates coordination

between place cells' temporal and spatial coding. On the other hand, synchronization strength

during immobility showed little correlation with the similarity of spatial tunings among place cell

pairs, suggesting that correlated activities at the millisecond timescale do not extend to slow,

behavioral-relevant timescales (Figure 6-figure supplement 1). Thus, place cells with distinct place

fields are linked by synchronous activity during novel locomotion.

Figure 6. Negative correlation between spatial tuning similarity and synchronization strength

during novel exploration.

(A) Spatially binned rate maps of place cells. High firing rates are indicated in yellow, and low firing

rates are noted in blue. Bottom: the mean firing rates along spatial bins averaged across laps.

(B) Spatial tunings of 10 place cells from an example session. Cells are sorted based on the locations

of their peak firing rates.

(C) Two example pairs of neurons with cross correlograms during locomotion and spatially binned

firing rate maps.

(D) A scatter plot of correlation coefficients of spatial tunings against normalized synchronization

strengths for place cell pairs. Each dot represents a pair of place cells. Synchronization strengths are

negatively correlated with similarities of spatial tunings (Spearman correlation coefficient=-0.4,

p<0.001, *n*=73 pairs).

Discussion

Our findings reveal the presence of synchronous ensembles comprising a substantial number of

CA1PCs during both locomotion and immobility in a novel environment. These ensembles

concurrently engage neurons representing diverse aspects of the environment and are rarely

observed during ripple events. Instead, they closely align with LFP and intracellular theta

oscillations. Our data demonstrate theta-associated synchronous ensembles during spatial memory

acquisition, coordinating numerous CA1 place cells that transmit information about different

segments of the environment.

The correlation between observed population synchrony and theta oscillations is intriguing. Theta

oscillations are typically associated with the sequential activation of neurons over time, resulting in

reduced neuronal synchrony[22]. However, theta oscillations come in different types, each

characterized by distinct pharmacological properties and behavioral correlates[56-58]. For example,

type 1 theta, or translational theta, occurs during voluntary movements and is sensitive to N-

methyl-D-aspartate receptor (NMDAR) blockers. On the other hand, type 2 theta, or attentional theta, is slightly slower and is blocked by muscarinic receptor antagonists, emerging during states of arousal or attention, such as when entering a new environment[59-61]. Consistent with these distinctions, the peak of the power spectrum density shows a distinctively slower theta frequency during immobility compared to locomotion (Figure 5B).

During theta oscillations, cholinergic neurons in the medial septum excite downstream CA1PCs, leading to increased firing rates and membrane potentials[62-64]. Given the widespread influence of cholinergic inputs on numerous CA1PCs, theta-coupled cholinergic inputs may drive depolarization and firing across many CA1PCs. Additionally, they could enhance the likelihood of synchronicity. Furthermore, GABAergic neurons in the medial septum, activated during theta oscillations, could activate interneurons in the CA1 region[58,65,66]. These CA1 interneurons, in turn, have the capability to entrain the activities of pyramidal neurons at both sub- and suprathreshold levels, thereby promoting synchronization among pyramidal neurons[67,68]. In summary, cholinergic and GABAergic neuron activation during theta waves may contribute to the excitation of CA1PCs and the consequent increase in their synchronicity.

Theta oscillations have long been implicated in memory acquisition. Specifically, theta power correlates positively with learning performance[69-71]. Perturbing generation of theta waves leads to impairment in memory acquisition[72-76], while electrical stimulation restoring theta rhythms rescues learning performance[77,78]. Furthermore, increased spike-theta coherence correlates with better memory, and stimulations that boost theta phase-locking of spikes enhance memory[79-81]. Despite abundant evidence supporting a link between theta oscillations and memory, the precise network mechanism by which theta oscillations support memory acquisition remains unclear. Theta oscillations are hypothesized to link spatially distributed neurons into functional ensembles to support memory acquisition[13]. The identified synchronous ensembles support the hypothesis,

providing a network substrate that links theta rhythms to the initial stage of memory acquisition.

Recent studies have demonstrated elevated coactivity of hippocampal principal cells during theta-

associated dentate spikes, events that are temporally coupled to theta oscillations and distinct from

sharp-wave ripples[82-84]. These findings highlight a network state characterized by rich synchronous

ensembles associated with theta, but not ripple, oscillations-a pattern consistent with our results.

Our analyses of publicly available data further support these observations, revealing theta-

associated population synchrony in a different dataset (Supplementary Figure 5). These results

suggest that hippocampal neurons achieve synchrony through network mechanisms distinct from

ripple-related events, reinforcing the hypothesis that this phenomenon is generalizable across

experimental conditions and behavioral contexts.

Previous studies have reported that a small percentage of CA1PCs generate spikes during ripple

episodes[8,14,17]. These measurements are typically conducted in familiar environments. In contrast,

when animals are introduced to a novel environment, ripple-associated spiking activities are

strongly suppressed, and CA1PCs exhibit prominent hyperpolarizing responses in their membrane

potentials (Figure 3, 4Ai-Ci, and Figure 3-figure supplement 3). Interestingly, after familiarization

training, a larger proportion of CA1PCs display positive ripple modulation of spiking activities and

depolarizing responses associated with ripples (Figure 4Aii-Cii). Consistent with the observation of

ripple-associated spiking activity in familiar environments, our study underscores the influence of

contextual novelty on hippocampal dynamics. Specifically, brain states associated with novelty may

modulate the activity of neurons upstream of CA1PCs, driving their membrane potentials toward

hyperpolarization during ripples. While the reduced correlation between ripples and spikes in novel

environments may initially appear counterintuitive, this unique neuronal activity pattern likely

reflects adaptive changes in hippocampal circuits related to the animals' behavioral states, such as

heightened vigilance in novel environments.

While contralateral LFP recordings reliably capture large-scale hippocampal theta and ripple oscillations[29-31], they may not fully account for ipsilateral-specific dynamics, such as variations in theta phase alignment or locally generated ripple events. Although contralateral recordings serve as a well-established proxy for large-scale hippocampal oscillatory states[30,85-87], incorporating simultaneous ipsilateral field potential recordings in future studies could refine our understanding of local-global network interactions. Despite these considerations, our findings provide robust evidence for the existence of synchronous neuronal ensembles and their role in coordinating newly formed place cells. These results advance our understanding of how synchronous neuronal ensembles contribute to spatial memory acquisition and hippocampal network coordination.

Although CA1PCs are known to exhibit weaker recurrent connectivity than that of CA3 pyramidal neurons, CA1PCs did communicate through monosynaptic connections[88]. These connections can be dynamically adjusted by the spike-timing-dependent plasticity mechanism[89-91], and therefore could be affected by the transient population synchrony. In addition, CA1PCs are known to communicate through inhibitory interneurons. Multiple pyramidal neurons converge onto a single interneuron, and individual connections are often weak and unreliable[51,92,93]. This connectivity scheme requires multiple pyramidal neurons spiking synchronously within the time window of synaptic integration to effectively activate postsynaptic neurons[51]. Furthermore, population synchrony among CA1PCs may propagate to downstream areas, synchronizing neurons outside the hippocampus. With their large ensemble sizes, these synchronous ensembles may synergistically convey relevant information or reorganize the connectivity of the neuronal networks.

Prior studies in sensory systems have commonly revealed that neurons with synchronous activities typically share similar sensory responses[94,95]. However, the synchronous ensembles we have identified often encompass neurons with distinct preferences for place fields. Pairwise synchrony

demonstrates a negative correlation with place field overlap (Figure 6). These findings indicate that

synchrony does not necessarily imply similar tunings between neurons. Furthermore, these

synchronous ensembles involve place cells encoding different spatial information. Recognizing the

crucial role of neuronal synchrony in binding distributed neurons in the brain[3,5], population

synchrony may serve as a mechanism to integrate diverse features of the same environment into a

cohesive mental map.

Methods

Animal

Time-pregnant wild-type C57BL/6J female mice underwent in utero virus injection of CamkII-cre

virus (105558-AAV1, Addgene) into the left side of the lateral ventricles. The resulting offspring

were raised in a breeding room under controlled temperature and humidity conditions and a 12-

hour light/dark cycle (lights on from 7:00 to 19:00). The mice used in the experiments were

between 12 and 18 weeks old at the time of the recordings. All surgical procedures, behavioral

training, and recording protocols were approved by the NYCU Institutional Animal Care and Use

Committee.

In-utero virus injection

Time-pregnant mice on embryonic day 14.5 or 17.5 were anesthetized with isoflurane (induction:4-

5%; maintenance:1-3%) before surgery. The animals were placed on a heating pad in a supine

position to maintain body temperature. Aseptic procedures were implemented to maintain sterile

conditions during the surgery. Uterine horns were exposed one at a time using spoons and fingers.

Warm and sterile phosphate-buffered saline (PBS) was used to rinse the uterus to prevent it from

drying out. Micropipettes loaded with CamkII-cre virus solution were used to inject 0.2 μ L of virus

suspension into the left side of the lateral ventricle of each embryo. The injection resulted in Cre

expression among neurons born on the day of injection, with earlier injection labeling neurons

located on the deeper side of the cell layer. The uterine horn was then gently returned to the

abdominal cavity. The exact process was repeated to the other uterine horn. After the surgery,

ketorolac (6mg/kg) and cefazolin (1g/kg) were administered for two days to minimize pain and

inflammation.

Cranial window and cannula implantation

The offspring of the mice that had undergone in-utero injection were subjected to chronic

hippocampal window surgery when they were between 2 and 4 months old. The animals were

initially anesthetized using isoflurane (induction:4-5%; maintenance:1-3%). Topical anesthesia was

applied by administering 0.5% lidocaine to the wound margins. After achieving deep anesthesia, an

incision was made in the skin, and a circular section of the skull was removed, centered 2 mm

caudal and 1.8 mm left to the bregma. Using 30-gauge needles and forceps, the dura was removed

from the exposed area, and cortical tissue within the craniotomy was aspirated. A serial dilution

experiment was conducted to determine an optimal titer of the virus carrying Voltron2 genes,

minimizing cell toxicity, for use in this and in previous imaging experiments[32]. A fine injection

pipette (tip diameter 10-60 um) was used to inject AAV2/1-CAG-flex-Voltron2-ST (2.7×10^{12} GC/ml, a

generous gift from Dr. Eric Schreiter and the GENIE team at HHMI Janelia Research Campus) into the

exposed regions at a depth of 200 μm (up to six injection sites and 100-200 nL of viral

suspension). An optical chamber was constructed by placing a cannula with a cover slip attached at

the bottom over the craniotomy and sealing it with dental acrylic or C&B Metabond (Sun Medical).

A custom-built titanium frame was then cemented to the animal's head using dental acrylic or C&B

Metabond. Mice received ketorolac (6mg/kg) and cefazolin (1g/kg) for two days following surgery to

minimize pain and inflammation.

Behavioral training

After at least two weeks of recovery from the window surgery, the mice underwent behavioral

training to ensure they were calm and attentive in the test environment. For mice subjected to

imaging experiments where they explored an air-lifted plastic track, the initial training phase

involved acclimating the mice to head fixation and treadmill running, which took place over 3-5 days

in a separate room distinct from the recording setup. Following this training, the imaging

experiments were conducted. On the day designated for imaging, the mice were introduced to the

recording room for the first time just before the imaging session. During the imaging experiments,

the mice were securely held in a head-fixed position beneath the microscope and on an air-lifted

plastic track that rested on an air table (AirLift, Neurotar; Figure 1A). The track was adorned with

various shapes of signs to help the mice navigate the track. Subdued blue lighting was provided to

illuminate the track. Before the initiation of imaging, the mice were confined in a corner of the track

using gates. Subsequently, the gates were opened upon the start of imaging, allowing the mice to

explore the new environment during their initial laps (Figure 1A Top view). The Mobile HomeCage

tracking system (Neurotar) was used to track the mice's position and speed. During the acquisition

of image frames, TTL pulses were sent by the high-speed camera to the Mobile HomeCage tracking

system to facilitate data alignment. Mice subjected to imaging experiments with head fixation in a

tube were first trained in a separate room, distinct from the recording setup, for 1-2 days to

habituate them to the head fixation procedure. Following this training, the mice were introduced to

the recording room for the first time immediately before the imaging session. During imaging, the

mice were securely positioned in a head-fixation setup within an acrylic tube and remained

stationary. To further familiarize the mice with the recording context, the imaging procedure was

repeated for 4 days without injecting the JF-HaloTag ligand. On the fifth day, the JF-HaloTag ligand

was injected prior to imaging, and voltage imaging was performed following the same procedure as

in the previous sessions.

Voltage imaging

JF$_{552}$-HaloTag ligand[96] (a generous gift from Dr. Luke Lavis) was first dissolved in DMSO (20 μ l,

Sigma) and then diluted in Pluronic^{TM} F-127 (20 μ l, P3000MP, Invitrogen) and PBS to achieve a

final concentration of 0.83 mM of JF$_{552}$-HaloTag ligand. The solution was then injected intravenously

through the retro-orbital sinus. Imaging sessions were initiated 3 hours after the injection of the

JF$_{552}$-HaloTag ligand. Fluorescence sensors were excited using a 532-nm laser (Opus 532, Laser

Quantum) with an excitation filter (FF02-520-28, Semrock), and the emitted light was separated

from the excitation light using a dichroic mirror (540lpxr Chroma) and passed through an emission

filter (FF01-596183 Semrock). A 16X, 0.8 NA objective (Nikon) was used to collect the emission light

which was then imaged on a CMOS camera (DaVinci-1K, RedShirt) with a 50-mm camera lens

(Nikkor 50mm f1.2, Nikon) as the tube lens. Images were acquired using Turbo-SM64 (Sci-Measure)

at 2kHz with a resolution of 190 x 160 pixels, corresponding to a field of view of 1.4 x 1.2 mm. The

number of frames in an image session was set at 360,000, corresponding to a duration of 180

seconds per session. Imaging depths ranged between 70 and 170 μm from the window surface,

where the CA1 pyramidal neurons were located. Time-lapse images were collected, and image

stacks of the same field of view along the z-axis were acquired. The image stacks consisted of 100

plans separated by two μm, covering the depth from the window surface to 200 um from the

surface.

LFP recording

The LFP signal from the contralateral CA1 region of the hippocampus was simultaneously recorded

during voltage imaging. To implant the LFP electrode, a small craniotomy was performed at the

dorsal CA1 coordinate (2mm caudal and 1.8 mm right to the bregma). A tungsten electrode

(#100211, 38 μm in diameter, polyimide insulated, California Fine Wire) was inserted into the

dorsal CA1 region, and the wire and pins were cemented in place with dental acrylic. In order to

target the CA1 pyramidal layer, the depth of the electrode was controlled and tracked by a

micromanipulator (Sutter Instrument, MP-285). Pyramidal layer-specific signatures of the LFP

signals, such as complex bursts and high-frequency ripple oscillations, were used to guide the

targeting. The electrode location was verified by using tungsten electrodes coated with DiI

(Invitrogen[TM], D282), followed by brain slicing after the recordings. The electrical signal was

amplified and filtered (1-1kHz) using a DAM80 amplifier (Word Precision Instruments). The

amplified signal was acquired by an I/O device (National Instruments USB-6341) and recorded using

WaveSurfer (Adam Taylor, HHMI Janelia Research Campus) at 10 kHz during implantation and

TurboSM64 (RedShirt) at 8 kHz during imaging.

Imaging somata and dendrites *in vivo* using a two-photon microscope

Immediately after voltage imaging, Voltron-labeled neurons were examined *in vivo* under a two-

photon microscope to assess the integrity of their membranes and dendrites. Images were acquired

using a two-photon scanning microscope equipped with a femtosecond-pulsed laser (Chameleon,

Coherent) and a resonant scanner controlled by ScanImage 4.2 software[97] (http://scanimage.org). A

16x water-immersion objective (Nikon, 0.8 NA, 3 mm working distance) was used for imaging. Z-

stacks of images (512 x 512 pixels, 600 μm x 600 μm) were collected with excitation light at 800

nm. The emission light was separated from the excitation light using a primary dichroic mirror

(FF735-Di02-25x36, Semrock), followed by a secondary dichroic mirror (565dclp, Chroma) and an

emission filter (FF01-609/57-30-D, Semrock). Each image stack consisted of 100 planes, separated

by 2 μm, covering depths from the window surface to 200 um below the surface.

Histological processing and quantification of soma depth

Mice with Voltron2$_{552}$ labeling were anesthetized with intraperitoneal injections of ketamine (250

mg/kg) and xylazine (25 mg/kg) and transcardially perfused with 4% paraformaldehyde in

phosphate-buffered saline (PBS; 2 mL/g). The brains were post-fixed in 4% paraformaldehyde

overnight and then washed in PBS. Coronal brain sections, 50 μm thick, were prepared using a

VT1200 vibratome (Leica). Sections were mounted with Vectashield mounting medium containing

DAPI (VectorLabs) to visualize nuclei. The soma depth of Voltron-labeled cells was measured relative

to the border between the *stratum pyramidale* (SP) and the *stratum radiatum*. Depth values were

normalized to the thickness of the SP, with larger depth ratios indicating somata located deeper

along the radial axis.

Voltage imaging preprocessing and spike detection

Fluorescence intensities were corrected for brain movement using rigid registration. Regions of

interest (ROIs) were manually selected by grouping pixels that cover individual somata. Previous

studies have described and validated the procedure for imaging preprocessing and spike

detection[31,32,98]. In short, the fluorescence intensities of individual neurons were calculated by

averaging the fluorescence intensities of pixels from the same ROIs. Bleaching was corrected by

calculating the baseline fluorescence (F_0) at each time point as an average of the fluorescence

intensities within ± 0.5 seconds around the time point. The dF/F was calculated as the F_0 minus the

fluorescence intensity of the same time point divided by F_0. Positive fluorescence transients were

detected to identify spikes from the high-passed dF/F traces created by subtracting the dF/F traces

from the median-filtered version with a 5-ms window. To simulate the noise of recordings, high-

passed dF/F traces were inverted, and the amplitudes of the transients detected from the inverted

traces were used to construct a noise distribution of the spike amplitudes. A threshold was set by

comparing the amplitudes of the detected transients with the noise distribution of the spike

amplitudes to minimize the sum of type I and type II errors. Spikes were first detected when

transients were larger than the threshold. Then, spike amplitudes smaller than half of the top 5%

spike amplitudes were excluded. The signal-to-noise ratio (SNR) was calculated for each neuron as a

ratio of the averaged spike amplitudes over the standard deviation of the high-passed dF/F traces,

excluding points 2 ms before and 4 ms after each detected spike to estimate the quality of the

recordings. The crosstalk between ROIs was routinely checked by examining the refractory periods

of neurons in auto-correlograms. Only neurons meeting the following criteria were included: (1)

SNR larger than 5, (2) full width at half maximum of the spike waveform longer than 0.8 ms, (3)

mean spike rates higher than 0.1 Hz, (4) distances between soma pairs at least 70 μm. Sessions

with at least nine neurons meeting the selection criteria were used for further analysis.

Behavioral states of locomotion and immobility

The periods of locomotion were defined as instances when the animals' speed exceeded 3 cm/s,

while periods of immobility were determined as instances when the animals' speed fell below 1

cm/s.

Mean firing rates during locomotion and immobility

To calculate the mean firing rates during locomotion, we counted the number of spikes during

locomotion periods and divided this number by the total duration of locomotion. Similarly, for mean

firing rates during immobility, we calculated the number of spikes that occurred during immobility

periods and divided this number by the total duration of immobility.

Correlation between instantaneous firing rate and animal speed

To estimate the instantaneous firing rates of individual neurons over time, we convolved their spike

trains with a Gaussian window of 250 ms. Subsequently, we calculated the Pearson correlation

coefficient between these instantaneous firing rates and the animal's speed for each neuron.

Grand average cross-correlogram (CCG)

Grand average CCGs were generated by constructing histograms of relative spike timings between

spikes of a reference cell and spikes of all other neurons in the same session as the target cell

population. Histograms were binned using 1-ms time bins. Spike counts were normalized by the

number of reference spikes times the number of all other neurons. An equal number of cells from

different sessions were randomly selected to form the shuffled target cell population to assess the

statistical significance of the CCG peaks. Subsequently, grand average CCGs were computed using

shuffled spike trains for comparison. Grand average CCGs were calculated using all spikes in the 180-

s recording periods, subsets of spikes during immobility periods (immobility grand average CCGs),

and subsets of spikes during locomotion periods (locomotion grand average CCGs).

Detection of the population synchrony

To assess population synchrony, we counted the number of spikes from all neurons within a sliding

window of 25 ms. To test the significance of the synchrony, we generated surrogate data by jittering

spike timings within a ±75-ms window 500 times. The means and standard deviations of each

sample point were calculated over the jitter population. Population synchrony was detected as

peaks when the spike count in the original data exceeded the mean plus four standard deviations of

the spike counts in the jittered data. Synchrony was also evaluated in modified spike trains,

including those where later spikes in bursts were removed and those where spike times were

shuffled across theta cycles while preserving phases. To estimate synchronous event rates, we

divided the entire 180-second recordings into halves and calculated the event rate in each of the 90-

s segments. Additionally, ensemble sizes were quantified by counting the percentage of neurons

participating in each synchronous event.

To detect population synchrony in a publicly available dataset, we analyzed the "hc-11" dataset

from CRCNS, which contains hippocampal recordings from rats running on novel mazes for water

rewards at designated locations[46,47]. Spikes recorded during the first three minutes of the novel-

maze running epochs were used to identify synchronous ensembles and generate jittered control

data. Spike counts for both the original and jittered data were calculated using a sliding 25-ms

window. The algorithm for testing the significance of synchrony was the same as described above.

Synchronous event rates were estimated using the entire 3-minute spike trains for both the original

and jittered data.

Pairwise cross-correlogram (CCG) of neuronal pairs

To compute the CCG, we first determined the time difference between spikes in the target and

reference neurons and then binned these values into 1-ms time bins. CCGs were only computed for

pairs with more than 100 counts within the CCG window of ±500 ms to ensure reliable correlation.

We defined the peak of the CCG as the local maxima within ±30 ms of lag. To assess the significance

of cross-correlations, we generated surrogate data by jittering the spike times of the target neuron

within a ± 75 ms window. We computed the CCG peak for a thousand iterations and calculated p-

values by comparing the percentile of the peak in the null distribution of the jittered peak values.

The synchronization strength was estimated as the ratio between the original CCG peak and the

mean spike count of the same bin as the original CCG peak averaged across the jittered histograms.

For display purposes, the CCGs were smoothed using a 5-point moving average. CCGs were

calculated using all spikes in the 180-s recording periods, subsets of spikes during immobility

periods (immobility pairwise CCGs), and subsets of spikes during locomotion periods (locomotion

pairwise CCGs).

LFP ripple analysis

To detect ripples, we performed band-pass filtering of the LFP signal between 120 and 240 Hz,

followed by computing the envelope using the absolute values of the Hilbert transform, which was

then low-pass filtered at 20 Hz. The start and end times of ripples were identified by setting the

upper and lower thresholds at the mean plus 7 and 3.5 times the standard deviation of the

envelope, respectively. For a ripple event to be included, three conditions had to be met: (1) values

within the event were higher than the lower threshold, (2) at least one value exceeded the upper

threshold, and (3) the minimum duration of the event was 30 ms. The start and end times of a

ripple event were defined as the positive and negative crossings of the lower threshold,

respectively. We defined in-ripple periods as the time between the start and end of detected LFP

ripple events, while out-ripple periods were periods outside the in-ripple periods. The envelope

peak values of the detected ripple events were used as the time of ripple occurrence. These timings

were utilized as trigger timings to calculate averages triggered by ripples. Two co-occurrences were

quantified: the percentages of LFP ripple oscillations that occurred with synchronous ensembles

and the percentages of synchronous ensembles that occurred with LFP ripple oscillations. The same

ripple detection algorithm was applied to the hc-11 dataset. To account for artifacts in the

recordings, raw traces and power spectra of the detected events were plotted and inspected for

false positives.

Ripple modulation index

We utilized the ripple modulation index to assess changes in firing rates during ripples compared to

periods outside ripples. This index was computed as the ratio of the difference in mean firing rates

during in- and out-ripple periods to the sum of the mean firing rates during both periods.

Specifically, we first calculated the mean firing rate during in-ripple periods and the mean firing rate

during out-ripple periods. Next, we subtracted the mean firing rate during out-ripple periods from

the mean firing rate during in-ripple periods and divided the result by the sum of the mean firing

rates during both periods. The resulting ripple modulation index indicates the magnitude and

direction of the firing rate modulation during ripple compared to periods outside ripples. To assess

the significance of the modulation index, we shuffled the spike timings and repeated the calculation

a thousand times to generate a distribution of modulation indices from the shuffled data. The

original modulation index was then compared to this distribution, and its significance was

determined if it exceeded the 95% confidence level of the shuffled indices.

LFP traces triggered by population synchrony and analysis

The midpoints of the synchronous event intervals were utilized as triggers to align LFP traces from

both our recordings and the hc-11 dataset. These traces were aligned within a $\pm$1-second window

around the trigger time points. To calculate the spectral density functions of the aligned LFP traces,

we performed a 16,384-point fast Fourier transform of the 2-second LFP traces. The power spectra

were then estimated as the squared absolute values of the Fourier coefficients. Ripple power was

calculated as the sum of the power distributed between 120 and 240 Hz. To calculate theta

modulation strength and preferred angles of the synchronous ensembles, LFP was first filtered

within the theta frequency range. Then, the phase $\varphi(t)$ and the instantaneous amplitude $A(t)$ of

the filtered LFP were computed using the Hilbert transform. A vector $V_k = A(t_k)e^{i\varphi(t_k)}$ was

assigned to each synchronous ensemble that occurred at time t_k. The modulation strength and the

preferred phase angle were determined as the absolute value and the phase angle of the summed

vector over all ensembles, respectively. Modulation strengths were normalized by the total length

of all vectors.

Subthreshold membrane voltage (subVm)

To compute the subVm of individual neurons, we began by determining the difference between the

raw dF/F signal and the high-pass version of dF/F. Next, we calculated the spike threshold for each

cell by identifying the voltage value corresponding to the peak of the first derivative of the spike

waveforms. Any fluorescence values in the slow dF/F time courses that surpassed the spike

thresholds were linearly interpolated.

SubVm traces triggered by population synchrony

The midpoint of the time intervals, during which synchronous ensembles were detected, served as

the reference points for aligning fluorescence traces within a $\pm$300 ms window around those

reference points. To eliminate any interference from membrane potential fluctuations associated

with spikes, we excluded aligned traces in cases where any spike occurred within the time intervals

of the aligned traces.

SubVm theta modulation and phase preference of spikes occurred inside and outside the

population synchrony events

The subVm was initially filtered within the theta frequency range. Subsequently, the phase $\varphi(t)$

and the instantaneous amplitude $A(t)$ of the filtered subVm were computed using the Hilbert

transform. Spikes were categorized into four groups: those occurring during or outside immobility

synchronous events and those occurring during or outside locomotion synchronous events. Within

each category, a vector $V_k = A(t_k)e^{i\varphi(t_k)}$ was assigned to each spike that occurred at time t_k.

The modulation strength and preferred phase angle were determined as the absolute value and

phase angle, respectively, of the summed vector over all spikes within a category. To facilitate

comparison between neurons, modulation strengths were normalized by the total length of all

vectors. The significance of the modulation strength was tested by shuffling the spike timings and

recalculating the modulation strength a thousand times to generate a distribution based on the

shuffled spike timings. The original modulation strength was then compared to the distribution,

with significance determined if it exceeded the 95% confidence interval of the shuffled values.

Significant modulation strengths were plotted and compared across groups.

Spike-triggered fluorescence traces

We extracted segments of fluorescence traces using a $\pm$300 ms time window centered on the spike

timings. To examine variations in fluorescence waveforms triggered by spikes within and outside

synchronous events, we categorized the fluorescence traces based on whether the spikes occurred

within or outside these events. Subsequently, we performed pairwise comparisons of the

fluorescence values from the same neuron, concentrating on spikes occurring during corresponding

behavioral states. Neurons with four or fewer triggering events in any of these categories were

omitted from the analysis.

Cross-correlation and coherence analysis to the subVm traces

To calculate cross-correlation and theta coherence of subVm between pairs of neurons during

locomotion and immobility, we initially divided the 180-second subVm traces into 1-second

segments with a 50% overlap. These segments were then categorized into the locomotion group if

the animal's speed remained above three cm/s for more than 90% of the time within that segment.

Similarly, segments were sorted into the immobility group if the animal's speed remained below

one cm/s for more than 90% of the time within that segment. Subsequently, we computed the

cross-correlation and magnitude-squared coherence function of frequency for pairs of co-occurred

segments belonging to a specific cell pair. Mean cross-correlations of individual cell pairs were

averaged across segments that occurred in the same behavioral states (i.e., locomotion or

immobility states). To estimate theta coherence, we averaged the values of the coherence function

within the theta frequency range (4-12 Hz). Theta coherences from the same cell pair during the

same behavioral state were pooled together, and the average was used as an estimate for the theta

coherence of that cell pair during a particular behavioral state. Finally, Spearman correlation

coefficients were calculated between the mean theta coherences of cell pairs and their soma

distances.

Spatial analysis

We partitioned a 90-cm-long track into 36 spatial bins to analyze spatial tuning properties. For each

cell, the spatial tuning curve was computed by determining the mean firing rate, defined as the

number of spikes within the bin when the animal's speed exceeded three cm/s, divided by the total

dwell times using the same speed threshold. The peak firing rates were identified as the maximal

rate on the spatial tuning curve. Spatial selectivity was calculated as one minus the circular variance

of the spatial tuning curve. Neurons were classified as place cells when their spatial selectivity

exceeded 0.25, and peak firing rates exceeded 1 Hz. To assess the similarity of spatial tuning

between pairs of neurons, Spearman correlation coefficients were calculated for the spatial tuning

curves. To explore the relationship between levels of synchronization and spatial tuning similarity

among pairs of place cells, Spearman correlation coefficients were computed between

synchronization strengths of cell pairs and correlation coefficients of their spatial tuning curves.

Data analysis and statistics

Data analysis was performed using MATLAB (Mathworks, R2019b), and results were presented as

standard error of the mean unless otherwise specified. Boxplots were used to depict the median,

first quartile, and third quartile, with the middle line, the upper edge, and the lower edge of the

box, respectively. The whiskers of the boxplots represented the 5th and 95th percentiles of the

distribution.

1   Takeuchi, T., Duszkiewicz, A. J. & Morris, R. G. The synaptic plasticity and memory hypothesis: encoding, storage and persistence. *Philos Trans R Soc Lond B Biol Sci* **369**, 20130288 (2014). https://doi.org:10.1098/rstb.2013.0288

2   Moser, M. B., Rowland, D. C. & Moser, E. I. Place cells, grid cells, and memory. *Cold Spring Harb Perspect Biol* **7**, a021808 (2015). https://doi.org:10.1101/cshperspect.a021808

3   Axmacher, N., Mormann, F., Fernandez, G., Elger, C. E. & Fell, J. Memory formation by neuronal synchronization. *Brain Res Rev* **52**, 170-182 (2006). https://doi.org:10.1016/j.brainresrev.2006.01.007

4   Paulsen, O. & Sejnowski, T. J. Natural patterns of activity and long-term synaptic plasticity. *Curr Opin Neurobiol* **10**, 172-179 (2000). https://doi.org:10.1016/s0959-4388(00)00076-3

5   Wang, X. J. Neurophysiological and computational principles of cortical rhythms in cognition. *Physiol Rev* **90**, 1195-1268 (2010). https://doi.org:10.1152/physrev.00035.2008

6   Harris, K. D., Csicsvari, J., Hirase, H., Dragoi, G. & Buzsaki, G. Organization of cell assemblies in the hippocampus. *Nature* **424**, 552-556 (2003). https://doi.org:10.1038/nature01834

7   Squire, L. R. & Wixted, J. T. The cognitive neuroscience of human memory since H.M. *Annu Rev Neurosci* **34**, 259-288 (2011). https://doi.org:10.1146/annurev-neuro-061010-113720

8   Buzsaki, G. Hippocampal sharp wave-ripple: A cognitive biomarker for episodic memory and planning. *Hippocampus* **25**, 1073-1188 (2015). https://doi.org:10.1002/hipo.22488

9   Nadasdy, Z., Hirase, H., Czurko, A., Csicsvari, J. & Buzsaki, G. Replay and time compression of recurring spike sequences in the hippocampus. *J Neurosci* **19**, 9497-9507 (1999). https://doi.org:10.1523/JNEUROSCI.19-21-09497.1999

10   Cheng, S. & Frank, L. M. New experiences enhance coordinated neural activity in the hippocampus. *Neuron* **57**, 303-313 (2008). https://doi.org:10.1016/j.neuron.2007.11.035

11   Dragoi, G. & Tonegawa, S. Preplay of future place cell sequences by hippocampal cellular assemblies. *Nature* **469**, 397-401 (2011). https://doi.org:10.1038/nature09633

12   Liu, C., Todorova, R., Tang, W., Oliva, A. & Fernandez-Ruiz, A. Associative and predictive hippocampal codes support memory-guided behaviors. *Science* **382**, eadi8237 (2023). https://doi.org:10.1126/science.adi8237

13   Colgin, L. L. Rhythms of the hippocampal network. *Nat Rev Neurosci* **17**, 239-249 (2016). https://doi.org:10.1038/nrn.2016.21

14   Buzsaki, G. & Silva, F. L. High frequency oscillations in the intact brain. *Prog Neurobiol* **98**, 241-249 (2012). https://doi.org:10.1016/j.pneurobio.2012.02.004

15   Csicsvari, J., Hirase, H., Czurko, A. & Buzsaki, G. Reliability and state dependence of pyramidal cell-interneuron synapses in the hippocampus: an ensemble approach in the behaving rat. *Neuron* **21**, 179-189 (1998). https://doi.org:10.1016/s0896-6273(00)80525-5

16   Gillespie, A. K. *et al.* Hippocampal replay reflects specific past experiences rather than a plan for subsequent choice. *Neuron* **109**, 3149-3163 e3146 (2021). https://doi.org:10.1016/j.neuron.2021.07.029

17 Yagi, S., Igata, H., Ikegaya, Y. & Sasaki, T. Awake hippocampal synchronous events are
incorporated into offline neuronal reactivation. *Cell Rep* **42**, 112871 (2023).
https://doi.org:10.1016/j.celrep.2023.112871

18 El-Gaby, M. *et al.* An emergent neural coactivity code for dynamic memory. *Nat Neurosci* **24**,
694-704 (2021). https://doi.org:10.1038/s41593-021-00820-w

19 Skaggs, W. E., McNaughton, B. L., Wilson, M. A. & Barnes, C. A. Theta phase precession in
hippocampal neuronal populations and the compression of temporal sequences.
Hippocampus **6**, 149-172 (1996). https://doi.org:10.1002/(SICI)1098-
1063(1996)6:2<149::AID-HIPO6>3.0.CO;2-K

20 Dragoi, G. & Buzsaki, G. Temporal encoding of place sequences by hippocampal cell assemblies.
Neuron **50**, 145-157 (2006). https://doi.org:10.1016/j.neuron.2006.02.023

21 Foster, D. J. & Wilson, M. A. Hippocampal theta sequences. *Hippocampus* **17**, 1093-1099
(2007). https://doi.org:10.1002/hipo.20345

22 Mizuseki, K. & Buzsaki, G. Theta oscillations decrease spike synchrony in the hippocampus and
entorhinal cortex. *Philos Trans R Soc Lond B Biol Sci* **369**, 20120530 (2014).
https://doi.org:10.1098/rstb.2012.0530

23 O'Keefe, J. & Recce, M. L. Phase relationship between hippocampal place units and the EEG
theta rhythm. *Hippocampus* **3**, 317-330 (1993). https://doi.org:10.1002/hipo.450030307

24 Lee, A. K. & Brecht, M. Elucidating Neuronal Mechanisms Using Intracellular Recordings during
Behavior. *Trends Neurosci* **41**, 385-403 (2018). https://doi.org:10.1016/j.tins.2018.03.014

25 Petersen, C. C. H. Whole-Cell Recording of Neuronal Membrane Potential during Behavior.
Neuron **95**, 1266-1281 (2017). https://doi.org:10.1016/j.neuron.2017.06.049

26 Giocomo, L. M., Zilli, E. A., Fransen, E. & Hasselmo, M. E. Temporal frequency of subthreshold
oscillations scales with entorhinal grid cell field spacing. *Science* **315**, 1719-1722 (2007).
https://doi.org:10.1126/science.1139207

27 Lampl, I., Reichova, I. & Ferster, D. Synchronous membrane potential fluctuations in neurons
of the cat visual cortex. *Neuron* **22**, 361-374 (1999). https://doi.org:10.1016/s0896-
6273(00)81096-x

28 Poulet, J. F. & Petersen, C. C. Internal brain state regulates membrane potential synchrony in
barrel cortex of behaving mice. *Nature* **454**, 881-885 (2008).
https://doi.org:10.1038/nature07150

29 Buzsaki, G. *et al.* Hippocampal network patterns of activity in the mouse. *Neuroscience* **116**,
201-211 (2003). https://doi.org:10.1016/s0306-4522(02)00669-3

30 Szabo, G. G. *et al.* Ripple-selective GABAergic projection cells in the hippocampus. *Neuron* **110**,
1959-1977 e1959 (2022). https://doi.org:10.1016/j.neuron.2022.04.002

31 Huang, Y. C. *et al.* Dynamic assemblies of parvalbumin interneurons in brain oscillations.
Neuron **112**, 2600-2613 e2605 (2024). https://doi.org:10.1016/j.neuron.2024.05.015

32 Abdelfattah, A. S. *et al.* Sensitivity optimization of a rhodopsin-based fluorescent voltage

974		indicator. *Neuron* (2023). https://doi.org:10.1016/j.neuron.2023.03.009
975	33	Angevine Jr, J. B. Time of neuron origin in the hippocampal region: an autoradiographic study
976		in the mouse. *Experimental Neurology* (1965).
977	34	Bayer, S. A. Development of the hippocampal region in the rat. I. Neurogenesis examined with
978		3H-thymidine autoradiography. *J Comp Neurol* **190**, 87-114 (1980).
979		https://doi.org:10.1002/cne.901900107
980	35	Cavalieri, D. *et al.* CA1 pyramidal cell diversity is rooted in the time of neurogenesis. *Elife* **10**
981		(2021). https://doi.org:10.7554/eLife.69270
982	36	Cembrowski, M. S. *et al.* Spatial Gene-Expression Gradients Underlie Prominent Heterogeneity
983		of CA1 Pyramidal Neurons. *Neuron* **89**, 351-368 (2016).
984		https://doi.org:10.1016/j.neuron.2015.12.013
985	37	Huszar, R., Zhang, Y., Blockus, H. & Buzsaki, G. Preconfigured dynamics in the hippocampus
986		are guided by embryonic birthdate and rate of neurogenesis. *Nat Neurosci* **25**, 1201-1212
987		(2022). https://doi.org:10.1038/s41593-022-01138-x
988	38	Adam, Y. *et al.* Voltage imaging and optogenetics reveal behaviour-dependent changes in
989		hippocampal dynamics. *Nature* **569**, 413-417 (2019). https://doi.org:10.1038/s41586-019-
990		1166-7
991	39	Klausberger, T. *et al.* Brain-state- and cell-type-specific firing of hippocampal interneurons in
992		vivo. *Nature* **421**, 844-848 (2003). https://doi.org:10.1038/nature01374
993	40	Wilson, M. A. & McNaughton, B. L. Dynamics of the hippocampal ensemble code for space.
994		*Science* **261**, 1055-1058 (1993).
995	41	Frank, L. M., Stanley, G. B. & Brown, E. N. Hippocampal plasticity across multiple days of
996		exposure to novel environments. *J Neurosci* **24**, 7681-7689 (2004).
997		https://doi.org:10.1523/JNEUROSCI.1958-04.2004
998	42	Leutgeb, S., Leutgeb, J. K., Treves, A., Moser, M. B. & Moser, E. I. Distinct ensemble codes in
999		hippocampal areas CA3 and CA1. *Science* **305**, 1295-1298 (2004).
1000		https://doi.org:10.1126/science.1100265
1001	43	Nitz, D. & McNaughton, B. Differential modulation of CA1 and dentate gyrus interneurons
1002		during exploration of novel environments. *J Neurophysiol* **91**, 863-872 (2004).
1003		https://doi.org:10.1152/jn.00614.2003
1004	44	Karlsson, M. P. & Frank, L. M. Network dynamics underlying the formation of sparse,
1005		informative representations in the hippocampus. *J Neurosci* **28**, 14271-14281 (2008).
1006		https://doi.org:10.1523/JNEUROSCI.4261-08.2008
1007	45	Jeewajee, A., Lever, C., Burton, S., O'Keefe, J. & Burgess, N. Environmental novelty is signaled
1008		by reduction of the hippocampal theta frequency. *Hippocampus* **18**, 340-348 (2008).
1009		https://doi.org:10.1002/hipo.20394
1010	46	Grosmark, A. D. & Buzsaki, G. Diversity in neural firing dynamics supports both rigid and
1011		learned hippocampal sequences. *Science* **351**, 1440-1443 (2016).

1012   https://doi.org:10.1126/science.aad1935

1013   47   Grosmark, A. D., Long, J. & Buzsáki, G. r. Recordings from hippocampal area CA1, PRE, during
1014    and POST novel spatial learning. *CRCNS. org* **10**, K0862DC0865 (2016).

1015   48   Rolotti, S. V. *et al.* Local feedback inhibition tightly controls rapid formation of hippocampal
1016    place fields. *Neuron* **110**, 783-794 e786 (2022). https://doi.org:10.1016/j.neuron.2021.12.003

1017   49   Sheffield, M. E. J., Adoff, M. D. & Dombeck, D. A. Increased Prevalence of Calcium Transients
1018    across the Dendritic Arbor during Place Field Formation. *Neuron* **96**, 490-504 e495 (2017).
1019    https://doi.org:10.1016/j.neuron.2017.09.029

1020   50   Takacs, V. T., Klausberger, T., Somogyi, P., Freund, T. F. & Gulyas, A. I. Extrinsic and local
1021    glutamatergic inputs of the rat hippocampal CA1 area differentially innervate pyramidal cells
1022    and interneurons. *Hippocampus* **22**, 1379-1391 (2012). https://doi.org:10.1002/hipo.20974

1023   51   English, D. F. *et al.* Pyramidal Cell-Interneuron Circuit Architecture and Dynamics in
1024    Hippocampal Networks. *Neuron* **96**, 505-520 e507 (2017).
1025    https://doi.org:10.1016/j.neuron.2017.09.033

1026   52   Maccaferri, G., Roberts, J. D., Szucs, P., Cottingham, C. A. & Somogyi, P. Cell surface domain
1027    specific postsynaptic currents evoked by identified GABAergic neurones in rat hippocampus
1028    in vitro. *J Physiol* **524 Pt 1**, 91-116 (2000). https://doi.org:10.1111/j.1469-7793.2000.t01-3-
1029    00091.x

1030   53   Geiller, T. *et al.* Local circuit amplification of spatial selectivity in the hippocampus. *Nature* **601**,
1031    105-109 (2022). https://doi.org:10.1038/s41586-021-04169-9

1032   54   Trouche, S. *et al.* Recoding a cocaine-place memory engram to a neutral engram in the
1033    hippocampus. *Nat Neurosci* **19**, 564-567 (2016). https://doi.org:10.1038/nn.4250

1034   55   Go, M. A. *et al.* Place Cells in Head-Fixed Mice Navigating a Floating Real-World Environment.
1035    *Front Cell Neurosci* **15**, 618658 (2021). https://doi.org:10.3389/fncel.2021.618658

1036   56   Kramis, R., Vanderwolf, C. H. & Bland, B. H. Two types of hippocampal rhythmical slow activity
1037    in both the rabbit and the rat: relations to behavior and effects of atropine, diethyl ether,
1038    urethane, and pentobarbital. *Exp Neurol* **49**, 58-85 (1975). https://doi.org:10.1016/0014-
1039    4886(75)90195-8

1040   57   Bland, B. H. The physiology and pharmacology of hippocampal formation theta rhythms. *Prog*
1041    *Neurobiol* **26**, 1-54 (1986). https://doi.org:10.1016/0301-0082(86)90019-5

1042   58   Buzsaki, G. Theta oscillations in the hippocampus. *Neuron* **33**, 325-340 (2002).
1043    https://doi.org:10.1016/s0896-6273(02)00586-x

1044   59   Sainsbury, R. S., Heynen, A. & Montoya, C. P. Behavioral correlates of hippocampal type 2 theta
1045    in the rat. *Physiol Behav* **39**, 513-519 (1987). https://doi.org:10.1016/0031-9384(87)90382-9

1046   60   Sainsbury, R. S. Hippocampal theta: a sensory-inhibition theory of function. *Neurosci Biobehav*
1047    *Rev* **22**, 237-241 (1998). https://doi.org:10.1016/s0149-7634(97)00011-0

1048   61   Barry, C., Heys, J. G. & Hasselmo, M. E. Possible role of acetylcholine in regulating spatial
1049    novelty effects on theta rhythm and grid cells. *Front Neural Circuits* **6**, 5 (2012).

https://doi.org:10.3389/fncir.2012.00005

62   Nakajima, Y., Nakajima, S., Leonard, R. J. & Yamaguchi, K. Acetylcholine raises excitability by inhibiting the fast transient potassium current in cultured hippocampal neurons. *Proc Natl Acad Sci U S A* **83**, 3022-3026 (1986). https://doi.org:10.1073/pnas.83.9.3022

63   Markram, H. & Segal, M. Acetylcholine potentiates responses to N-methyl-D-aspartate in the rat hippocampus. *Neurosci Lett* **113**, 62-65 (1990). https://doi.org:10.1016/0304-3940(90)90495-u

64   Fraser, D. D. & MacVicar, B. A. Cholinergic-dependent plateau potential in hippocampal CA1 pyramidal neurons. *J Neurosci* **16**, 4113-4128 (1996). https://doi.org:10.1523/JNEUROSCI.16-13-04113.1996

65   Toth, K., Freund, T. F. & Miles, R. Disinhibition of rat hippocampal pyramidal cells by GABAergic afferents from the septum. *J Physiol* **500 (Pt 2)**, 463-474 (1997). https://doi.org:10.1113/jphysiol.1997.sp022033

66   Colgin, L. L. Mechanisms and functions of theta rhythms. *Annu Rev Neurosci* **36**, 295-312 (2013). https://doi.org:10.1146/annurev-neuro-062012-170330

67   Cobb, S. R., Buhl, E. H., Halasy, K., Paulsen, O. & Somogyi, P. Synchronization of neuronal activity in hippocampus by individual GABAergic interneurons. *Nature* **378**, 75-78 (1995). https://doi.org:10.1038/378075a0

68   Stark, E. *et al.* Inhibition-induced theta resonance in cortical circuits. *Neuron* **80**, 1263-1276 (2013). https://doi.org:10.1016/j.neuron.2013.09.033

69   Landfield, P. W., McGaugh, J. L. & Tusa, R. J. Theta rhythm: a temporal correlate of memory storage processes in the rat. *Science* **175**, 87-89 (1972). https://doi.org:10.1126/science.175.4017.87

70   Berry, S. D. & Thompson, R. F. Prediction of learning rate from the hippocampal electroencephalogram. *Science* **200**, 1298-1300 (1978). https://doi.org:10.1126/science.663612

71   Belchior, H., Lopes-Dos-Santos, V., Tort, A. B. & Ribeiro, S. Increase in hippocampal theta oscillations during spatial decision making. *Hippocampus* **24**, 693-702 (2014). https://doi.org:10.1002/hipo.22260

72   Winson, J. Loss of hippocampal theta rhythm results in spatial memory deficit in the rat. *Science* **201**, 160-163 (1978). https://doi.org:10.1126/science.663646

73   Mitchell, S. J., Rawlins, J. N., Steward, O. & Olton, D. S. Medial septal area lesions disrupt theta rhythm and cholinergic staining in medial entorhinal cortex and produce impaired radial arm maze behavior in rats. *J Neurosci* **2**, 292-302 (1982). https://doi.org:10.1523/JNEUROSCI.02-03-00292.1982

74   Mizumori, S. J., Perez, G. M., Alvarado, M. C., Barnes, C. A. & McNaughton, B. L. Reversible inactivation of the medial septum differentially affects two forms of learning in rats. *Brain research* **528**, 12-20 (1990). https://doi.org:10.1016/0006-8993(90)90188-h

75   M'Harzi, M. & Jarrard, L. E. Effects of medial and lateral septal lesions on acquisition of a place
and cue radial maze task. *Behav Brain Res* **49**, 159-165 (1992). https://doi.org:10.1016/s0166-
4328(05)80160-3

76   Wang, Y., Romani, S., Lustig, B., Leonardo, A. & Pastalkova, E. Theta sequences are essential
for internally generated hippocampal firing fields. *Nat Neurosci* **18**, 282-288 (2015).
https://doi.org:10.1038/nn.3904

77   McNaughton, N., Ruan, M. & Woodnorth, M. A. Restoring theta-like rhythmicity in rats
restores initial learning in the Morris water maze. *Hippocampus* **16**, 1102-1110 (2006).
https://doi.org:10.1002/hipo.20235

78   Lee, D. J. *et al.* Medial septal nucleus theta frequency deep brain stimulation improves spatial
working memory after traumatic brain injury. *J Neurotrauma* **30**, 131-139 (2013).
https://doi.org:10.1089/neu.2012.2646

79   Hasselmo, M. E., Bodelon, C. & Wyble, B. P. A proposed function for hippocampal theta rhythm:
separate phases of encoding and retrieval enhance reversal of prior learning. *Neural Comput*
14, 793-817 (2002). https://doi.org:10.1162/089976602317318965

80   Rutishauser, U., Ross, I. B., Mamelak, A. N. & Schuman, E. M. Human memory strength is
predicted by theta-frequency phase-locking of single neurons. *Nature* **464**, 903-907 (2010).
https://doi.org:10.1038/nature08860

81   Siegle, J. H. & Wilson, M. A. Enhancement of encoding and retrieval functions through theta
phase-specific manipulation of hippocampus. *Elife* **3**, e03061 (2014).
https://doi.org:10.7554/eLife.03061

82   McHugh, S. B. *et al.* Offline hippocampal reactivation during dentate spikes supports flexible
memory. *Neuron* **112**, 3768-3781 e3768 (2024).
https://doi.org:10.1016/j.neuron.2024.08.022

83   Dvorak, D., Chung, A., Park, E. H. & Fenton, A. A. Dentate spikes and external control of
hippocampal function. *Cell Rep* **36**, 109497 (2021).
https://doi.org:10.1016/j.celrep.2021.109497

84   Farrell, J. S., Hwaun, E., Dudok, B. & Soltesz, I. Neural and behavioural state switching during
hippocampal dentate spikes. *Nature* **628**, 590-595 (2024). https://doi.org:10.1038/s41586-
024-07192-8

85   Dudok, B. *et al.* Alternating sources of perisomatic inhibition during behavior. *Neuron* **109**,
997-1012 e1019 (2021). https://doi.org:10.1016/j.neuron.2021.01.003

86   Terada, S. *et al.* Adaptive stimulus selection for consolidation in the hippocampus. *Nature* **601**,
240-244 (2022). https://doi.org:10.1038/s41586-021-04118-6

87   Geiller, T. *et al.* Large-Scale 3D Two-Photon Imaging of Molecularly Identified CA1 Interneuron
Dynamics in Behaving Mice. *Neuron* **108**, 968-983 e969 (2020).
https://doi.org:10.1016/j.neuron.2020.09.013

88   Deuchars, J. & Thomson, A. M. CA1 pyramid-pyramid connections in rat hippocampus in vitro:

dual intracellular recordings with biocytin filling. *Neuroscience* **74**, 1009-1018 (1996). https://doi.org:10.1016/0306-4522(96)00251-5

Caporale, N. & Dan, Y. Spike timing-dependent plasticity: a Hebbian learning rule. *Annu Rev Neurosci* **31**, 25-46 (2008). https://doi.org:10.1146/annurev.neuro.31.060407.125639

Markram, H., Lubke, J., Frotscher, M. & Sakmann, B. Regulation of synaptic efficacy by coincidence of postsynaptic APs and EPSPs. *Science* **275**, 213-215 (1997). https://doi.org:10.1126/science.275.5297.213

Bi, G. Q. & Poo, M. M. Synaptic modifications in cultured hippocampal neurons: dependence on spike timing, synaptic strength, and postsynaptic cell type. *J Neurosci* **18**, 10464-10472 (1998). https://doi.org:10.1523/JNEUROSCI.18-24-10464.1998

Ali, A. B., Deuchars, J., Pawelzik, H. & Thomson, A. M. CA1 pyramidal to basket and bistratified cell EPSPs: dual intracellular recordings in rat hippocampal slices. *J Physiol* **507 (Pt 1)**, 201-217 (1998). https://doi.org:10.1111/j.1469-7793.1998.201bu.x

Ali, A. B. & Thomson, A. M. Facilitating pyramid to horizontal oriens-alveus interneurone inputs: dual intracellular recordings in slices of rat hippocampus. *J Physiol* **507 (Pt 1)**, 185-199 (1998). https://doi.org:10.1111/j.1469-7793.1998.185bu.x

Engel, A. K., Konig, P., Kreiter, A. K. & Singer, W. Interhemispheric synchronization of oscillatory neuronal responses in cat visual cortex. *Science* **252**, 1177-1179 (1991). https://doi.org:10.1126/science.252.5009.1177

Alonso, J. M., Usrey, W. M. & Reid, R. C. Precisely correlated firing in cells of the lateral geniculate nucleus. *Nature* **383**, 815-819 (1996). https://doi.org:10.1038/383815a0

Zheng, Q. *et al.* Rational Design of Fluorogenic and Spontaneously Blinking Labels for Super-Resolution Imaging. *ACS Cent Sci* **5**, 1602-1613 (2019). https://doi.org:10.1021/acscentsci.9b00676

Pologruto, T. A., Sabatini, B. L. & Svoboda, K. ScanImage: flexible software for operating laser scanning microscopes. *Biomed Eng Online* **2**, 13 (2003). https://doi.org:10.1186/1475-925X-2-13

Abdelfattah, A. S. *et al.* Bright and photostable chemigenetic indicators for extended in vivo voltage imaging. *Science* **365**, 699-704 (2019). https://doi.org:10.1126/science.aav6416

Acknowledgments

We express our sincere gratitude to Dr. Luke Lavis for sharing the JF$_{552}$-HaloTag ligand. Special thanks to Mr. Yu-Ting Lin and Ms. Hui-Ching Chen for technical assistance. We are grateful to Drs. Shih-Chieh Lin, Kuo-Hua Huang, Shi-Bing Yang, and Fu-Chin Liu for their insightful comments on the manuscript.

Funding statements

National Science and Technology Council, Taiwan (B.-J.L. and T.-W.C.)

1164 Howard Hughes Medical Institute, USA (E.R.S.)

1165

Supplementary Figure 1. Morphology of Voltron-expressing neurons and their positions along the deep-superficial axis.

(A) Two-photon z-stack maximum intensity projection images of Voltron-expressing neurons obtained after imaging experiments. These neurons exhibit well-preserved membrane and dendritic morphology. Scale bars: 50 μm.

(B) Post-hoc histological images of a sparsely labeled neuron previously subjected to voltage imaging. Scale bars: 300 μm.

(C) Micrographs of labeled neurons from *in-utero* electroporation at embryonic day 14.5 (E14.5; upper panel) and E17.5 (lower panel). Scale bars: 300 μm.

(D) Histograms of normalized depths for E14.5 neurons (red) and E17.5 neurons (blue). A higher normalized depth indicates that the soma is located deeper within the pyramidal layer. Normalized depths for E14.5 neurons are significantly higher than those for E17.5 neurons ($t_{(601)}$=22.8, *p*<0.0001, Student's *t*-test).

1181

Supplementary Figure 2. Mean firing rate distribution and correlation with animals' speed.

(A) Cumulative distribution of mean firing rates averaged throughout the entire 180-second

recording period.

(B) Mean firing rates during locomotion plotted against mean firing rates during immobility for

individual cells. Each dot represents a cell. Mean firing rates are higher during immobility than

during locomotion (4.2±0.2 Hz for immobility, 2.8±0.2 Hz for locomotion, $t_{(70)}$=6.63, p<0.0001,

Student's t-test).

(C) Histogram of correlation coefficients between instantaneous firing rates and the animals' speed

for each neuron.

1191

1192

Figure 1-figure supplement 1. Pairwise cross-correlogram (CCG) between CA1 pyramidal cells.

(A) CCGs of two representative cell pairs exhibiting significant peaks compared to the jittered spike

trains of the target cells (gray lines).

(B) Distribution of the CCG peaks (n=497 pairs). Black: CCG peaks with significance (p<0.05). White:

CCG peaks without significance (p>=0.05).

(C) Histogram of peak lags of significant CCG peaks (n=315 pairs).

(D) Histogram of FWHMs of significant CCG peaks (n=315 pairs).

1200

Supplementary Figure 3. Population synchrony remains after removing the later spikes in bursts.

(A) Grand-average cross-correlogram (CCG) calculated using spike trains with the later spikes in

bursts removed. The gray line represents the mean grand-average CCG between reference cells and

randomly selected cells from different sessions.

(B) Pairwise comparison of population synchrony event rates between spike trains containing all

spikes and those with the later spikes in bursts removed. Bar heights represent group means ($n=10$

segments, $p=0.36$, Wilcoxon signed-rank test).

(C) Histogram of ensemble sizes showing the percentage of cells participating in synchronous

ensembles.

1213

Figure 3-figure supplement 1. LFP ripples recorded in the first and subsequent sessions in a novel

maze.

(A) An example LFP trace with ripples recorded in a subsequent session following the first session of

novel exploration, where no ripple was detected.

(B) Ripple rates in the first and subsequent sessions recorded in the same animals (n=5 mice) during

novel exploration.

1221

Figure 3-figure supplement 2. Population synchrony during novel immobility.

(A) Grand average cross-correlograms (CCGs) for neurons labeled on E14.5 (left) and on E17.5

(right). The gray line represents the mean grand-average CCG between reference cells and randomly

selected cells from different sessions.

(B) Pairwise comparison of synchronous event rates between the original and jittered data. Event

rates were significantly higher in the original data compared to jittered controls for E14.5 neurons

(left; mean event rate: 1.21 $\pm$ 0.24 Hz for the original data, 0.08 $\pm$ 0.009 Hz for jittered data, n=6

segments, W=21, p=0.03, Wilcoxon signed-rank test) and E17.5 neurons (right; mean event rate:

0.75 $\pm$ 0.17 Hz for the original data, 0.18 $\pm$ 0.03 Hz for jittered data, n=6 segments, W=21, p=0.03,

Wilcoxon signed-rank test). Bar heights represent group means. *p<0.05

(C) Histograms of ensemble sizes showing the percentage of cells participating in synchronous

ensembles for E14.5 neurons (left) and E17.5 neurons (right).

1234

Figure 3-figure supplement 3. Population synchrony outside ripple episodes during novel

immobility.

(A) Example traces illustrating the distinct timings of ripple episodes and synchronous ensembles.

Top: Simultaneously recorded LFP trace showing episodes of ripple oscillations. Red vertical lines

indicate the timings of peak ripple power in the LFP trace, while yellow vertical bars mark the

duration of ripple episodes. Bottom: Fluorescence traces from 25 CA1PCs. Gray vertical lines

indicate the timings of synchronous ensembles.

(B) Percentages of co-occurrence between synchronous ensembles and ripple events. The "sync to

ripples" group represents the percentage of ripple events that co-occur with synchronous

ensembles, while the "ripples to sync" group shows the percentage of synchronous ensembles that

co-occur with ripple events. Each gray circle represents an individual session.

(C) LFP ripple power and spiking probability aligned to synchronous ensembles and ripple events.

Upper panel: Color-coded LFP power at ripple frequency (120-240Hz), aligned to the timings of

1052 synchronous ensembles (left) and 174 ripple events (right). **Middle panel**: Average LFP power

at ripple frequencies aligned to the timings of synchronous ensembles (left) and ripple events

(right). **Lower panel**: Average spiking probability of all recorded cells aligned to the timings of

synchronous ensembles (left) and ripple events (right).

(D) Boxplots and comparisons of peak ripple power (upper panel) and peak spiking probability

(lower panel) at the timings (t=0) of population synchrony (left) and ripple events (right). For peak

ripple power: $p<0.001$, $t_{(1224)}=86.6$, Student's t-test; for peak spiking probability: $p<0.001$,

$t_{(1224)}=24.3$, Student's t-test. Gray circles indicate peak values for individual triggered traces.

1256

1257

Figure 5-figure supplement 1. SubVm theta modulation of spikes during immobility and

locomotion periods of novel exploration.

(A) Polar plot comparing subVm theta modulation between spikes participating in synchronous

ensembles (sync spikes) and spikes not participating in synchronous ensembles (other spikes) during

immobility. Each dot represents the averaged modulation of a cell. Cells with modulation strengths

that are not significant are excluded in the plot and in the comparison. The modulation strengths of

spikes participating in synchrony are significantly higher than those of spikes not participating in

synchrony (modulation strength for spikes participating in synchronous ensembles: 0.81±0.07, for

spikes not participating in synchronous ensembles: 0.64±0.11, mean±s.d., $t_{(64)}$=12.6, $p<0.001$, paired

t-test).

(B) Same as in (A), but during locomotion (modulation strength for spikes participating in

synchronous ensembles: 0.80 ± 0.09, for spikes not participating in synchronous ensembles: $0.60\pm$

0.12, mean±s.d., $t_{(53)}=12.3$, $p<0.001$, paired t-test).

(C) Average fluorescence triggered by cells' spikes participating in synchronous ensembles (red, sync

spikes) and not participating in synchronous ensembles during immobility (black, other spikes; $n=59$

cells). Inset: Pairwise comparison of average fluorescence 50 ± 5 ms before time of spikes (t=0)

between the two triggered groups ($n=59$ cells, $p<0.001$, paired t-test). ***p<0.001.

(D) Same as in (C), but during locomotion. Inset: Pairwise comparison of average fluorescence ($n=59$

cells, $p<0.001$, paired t-test).

1277

1278

Supplementary Figure 4. Significant reduction of population synchrony by randomizing spikes

across theta cycles while preserving phases

Pairwise comparison of population synchrony event rates between the original spike trains and

randomized spike trains, in which spike timings were shuffled across theta cycles while preserving

phases (n=10 segments, p=0.002, Wilcoxon signed-rank test). Bar heights represent group means.

** p<0.01

1285

1286

1287

1288 **Figure 5-figure supplement 1. Synchronization strengths during immobility show little correlation**

1289 **with the similarity of spatial tunings between place cells.** Synchronization strengths during

1290 immobility plotted against correlation coefficients of spatial tunings for pairs of place cells.

1291 Synchronization strengths show little correlation with similarities of spatial tunings (Spearman

1292 correlation coefficient=-0.003, *p*=0.98, *n*=88 cell pairs).

1293

1294

Supplementary Figure 5. Analyses of synchronous ensembles in the publicly available data

(A) Pairwise comparison of synchronous event rates between the original and jittered data. Event rates were significantly higher in the original data compared to jittered controls (n=8 sessions, p=0.008, Wilcoxon signed-rank test). Bar heights represent group means. **p<0.01.

(B) Dynamics of ripple power aligned to timings of ripples (black) and timings of synchronous ensembles (red).

(C) Average LFP trace aligned to timings of synchronous ensembles of a session.

(D) Polar histogram of the differences between theta phases of individual synchronous ensembles and the preferred phase of the session.