Transport mechanism of DgoT, a bacterial homolog of SLC17 organic anion transporters

Nataliia Dmitrieva[1], Samira Gholami[1,2], Claudia Alleva[1,7], Paolo Carloni[2,4,5,6],
Mercedes Alfonso-Prieto[2,3,*], Christoph Fahlke[1,*]

[1] Institute of Biological Information Processing (IBI-1), Molekular- und Zellphysiologie, Forschungszentrum Jülich, 52425 Jülich, Germany.

[2] Institute for Advanced Simulation (IAS-5) and Institute of Neuroscience and Medicine (INM-9), Computational Biomedicine, Forschungszentrum Jülich, 52425 Jülich, Germany.

[3] Cécile and Oskar Vogt Institute for Brain Research, University Hospital Düsseldorf, Medical Faculty, Heinrich Heine University Düsseldorf, 40225 Düsseldorf, Germany.

[4] JARA-HPC, Forschungszentrum Jülich, 54245 Jülich, Germany.

[5] Department of Physics, RWTH Aachen University, 52056 Aachen, Germany.

[6] JARA Institute Molecular Neuroscience and Neuroimaging (INM-11), Forschungszentrum Jülich, 52425 Jülich, Germany.

[7] Present address: Department of Biochemistry and Biophysics and Science for Life Laboratory, Stockholm University, Sweden.

* Contributing equally

Character count (Abstract, Introduction, Results, Discussion and Methods): 61048

Abstract

The solute carrier 17 (SLC17) family contains anion transporters that accumulate neuro-transmitters in secretory vesicles, remove carboxylated monosaccharides from lysosomes, or extrude organic anions from the kidneys and the liver. We combined classical molecular dynamics simulations, Markov state modeling and hybrid first principles quantum mechanical/classical mechanical (QM/MM) simulations with experimental approaches to describe the transport mechanisms of a model bacterial protein, the D-galactonate transporter DgoT, at atomic resolution. We found that protonation of D46 and E133 precedes galactonate binding and that substrate binding induces closure of the extracellular gate, with the conserved R47 coupling substrate binding to transmembrane helix movement. After isomerization to an inward-facing conformation, deprotonation of E133 and subsequent proton transfer from D46 to E133 opens the intracellular gate and permits galactonate dissociation either in its unprotonated form or after proton transfer from E133. After release of the second proton, apo DgoT returns to the outward-facing conformation. Our results provide a framework to understand how various SLC17 transport functions with distinct transport stoichiometries can be attained through subtle variations in proton and substrate binding/unbinding.

Keywords: major facilitator superfamily / molecular dynamics simulations / organic anion transporter / proton-coupled secondary transport /solid-supported membrane electrophysiology

Introduction

Solute carrier 17 (SLC17) transporters fulfil a variety of cellular functions (Reimer 2013; Omote et al. 2016; Li et al. 2022). They transport diverse anionic substrates; mostly electro-chemical proton (H^+) gradients as the driving force, with a large variability in transport stoi-chiometry ranging from electrogenic H^+-glutamate exchange by VGLUT (Kolen et al. 2023) to electroneutral H^+-sialic acid symport by sialin (Morin et al. 2004; Hu et al. 2023). Although this protein family has been studied for several decades (Kaback and Guan 2019), the mechanistic basis of SLC17 organic anion transport remains insufficiently understood.

The bacterial homolog DgoT (from *E. coli*) transports negatively charged galactonate in symport with more than one proton and, thus, differs from mammalian SLC17s in both its main substrate and transport stoichiometry (Leano et al. 2019). It belongs to the large major facilitator superfamily (MFS). MFS family members exhibit a distinct topology that includes 12 transmembrane (TM) helices, with the substrate-binding pocket located between two pseudo-symmetrical 6-TM bundles in the N-domain (TM1–TM6) and C-domain (TM7–TM12) (Quistgaard et al. 2016; Drew et al. 2021) . DgoT has been crystallized in two distinct conformations: (i) inward-facing and (ii) outward-facing with galactonate bound to the central binding site (Leano et al. 2019). These structures revealed four charged residues within the transmembrane domains, D46, R47, R126, and E133, with the last three conserved across the SLC17 family. The high structural similarity with vesicular glutamate transporters (VGLUTs) (Li et al. 2020) and lysosomal sialic acid transporter (Hu et al. 2023) (Supplementary Figs 1 and 2), together with differences in substrate selectivity and transport stoichiometry, makes DgoT an attractive model to study the atomistic basis of coupled transport in the SLC17 family. Here, we present a comprehensive study of DgoT transport using a combination of *in vitro* experiments and molecular dynamics (MD) simulations.

Results

DgoT mediates the coupled symport of two H⁺ and one D-galactonate

We studied DgoT transport using solid-supported membrane-based electrophysiology (SSME) with purified DgoT reconstituted into proteoliposomes. In such recordings, transport activity is induced by fast solution exchange. The resulting changes in proteoliposomal membrane potential are converted into an electrical signal and detected by a measuring electrode via capacitive coupling (Schulz et al. 2008; Bazzone et al. 2021).

Fig. 1a depicts SSME currents after galactonate concentration jumps at three different pHs. At neutral and alkaline pH, the time course for current decay decreased at higher lipid–protein ratios (LPRs; illustrated for pH 8.0 in the inset). This result indicates that the measured currents are caused by electrogenic DgoT transport (Bazzone et al. 2021). Transport currents assumed a maximum amplitude at pH 7.5 (blue line) and decreased at higher pH values (Fig. 1a; pH 9.0, black line). Under asymmetrical ionic conditions (i.e., with an alkaline pH outside, but not inside, the liposomes), the peak current was reduced (Fig. 1b bottom). This suggests that reduced transport at alkaline pH is caused by proton depletion at the binding site outside the liposomes. Under symmetrical pH conditions, the current decay was biphasic at pH values lower than 7.5 (Supplementary Fig. 3): the fast component represents conformational changes triggered by galactonate binding and the slow component is associated with the transport activity of the protein. At pH values lower than 6, only the pre-steady state, but not the transport component, was observed (Fig. 1a; red line). Acidic pH inside, but not outside, the liposomes inhibits transport activity, as demonstrated under asymmetrical ionic conditions (Fig. 1b top). We conclude that current reduction under conditions of acidic or alkaline pH is regulated by two different processes: proton binding from the outside and proton release to the inside of the proteoliposomes, respectively. Thus, analysis of the peak currents allowed the determination of apparent pKa values for proton binding and release (Fig. 1c).

The transport currents generated by DgoT were positive (Fig. 1a), indicating that at

least two protons are transported with each galactonate molecule, as previously reported

(Leano et al. 2019). We used a reversal potential assay to determine the transport stoichi-

ometry (Thomas et al. 2021). In SSME, no voltage is applied to the membrane and concen-

tration gradients are the only driving force for coupled transport. The free energy for the

coupled transport reaction is given by:

$\Delta G = n\Delta\mu_i + m\Delta\mu_s,$

where n and m are number of protons and galactonate molecules transported together. The

current reverses at $\Delta G = 0$:

$$\frac{n}{m} = -\frac{\Delta\mu_s}{\Delta\mu_i} = -\frac{RT\ln\left(\frac{[galactonate]_{in}}{[galactonate]_{out}}\right)}{RT\ln\left(\frac{[H^+]_{in}}{[H^+]_{out}}\right)} = \frac{\ln\left(\frac{[galactonate]_{in}}{[galactonate]_{out}}\right)}{\ln\left(\frac{[H^+]_{out}}{[H^+]_{in}}\right)}$$

Thus, plotting the transported charge against the ratio of the electrochemical gradi-

ents enables the transport stoichiometry to be determined (Thomas et al. 2021).

We preloaded proteoliposomes with galactonate-containing internal solutions and ap-

plied sets of external solutions to generate an outwardly directed proton concentration gra-

dient at various galactonate concentration gradients. The currents obtained with empty lip-

osomes—used as the negative control (Supplementary Fig. 4)—were subtracted. Null

transport occurred at the gradient ratios corresponding to a stoichiometry of 2 H$^+$:1 galacto-

nate (Fig. 1d, e).

To investigate the binding order of H$^+$ and galactonate, we measured changes to the

external pH of solutions containing purified detergent-solubilized protein upon the addition

of galactonate (Soskine et al. 2004). If galactonate binding precedes proton association, we

would expect the pH to increase upon galactonate addition. However, addition of 10 mM

galactonate to purified DgoT in unbuffered detergent-containing solutions did not elicit any

change in pH (Fig. 1f). This result suggests that the substrate binds to the protonated trans-

porter and does not induce protonation of DgoT. Therefore, we conclude that DgoT binds

two H^+ prior to galactonate association.

Protonation of key residues regulates extracellular gate dynamics

We studied H^+ and galactonate binding to outward-facing DgoT using unbiased all-atom MD

simulations. For this, we used the crystal structure of E133Q DgoT (Leano et al. 2019) (PDB

ID: 6E9O) as the starting conformation after reverting the mutation. First, we studied the

effect of protonation on *apo* protein dynamics. The crystallographic galactonate molecule

was removed from the binding site, and four different systems were generated with D46 and

E133 individually protonated or deprotonated. We observed reproducible movements of the

TM1 and TM7 transmembrane helices, with most flexibility in the most extracellular seg-

ments (around residues 48–52 of TM1 and 271–275 of TM7). This behavior resembles the

proposed role of TM1 and TM7 as gating helices that mediate transitions between open and

occluded conformations, as in other MFS members (Smirnova et al. 2014; Qureshi et al.

2020; Feng et al. 2021).

When D46 and E133 were unprotonated, the extracellular gate (defined as distance

between the center of mass of the Cα atoms of residues 48–52 and residues 271–275) was

mostly closed (Fig. 2a, d). Protonation of both D46 and E133 locked the gate in an open

conformation (Fig. 2b, d), and protonation of only one of these residues only partially opened

the gate (Fig. 2d). These changes in position of the gating helices were coupled to local

rearrangements in the N-terminal domain. Charged, but not protonated, D46 and E133 in-

teract with R126 and R47, respectively. After protonation, the side chain of D46 moves away,

allowing R126 to interact with N49. This results in rotation of TM1 and stabilization of the

open state of the extracellular gate (compare Fig. 2a, b). In addition, protonation of E133

releases R47, which then becomes available to interact with the carboxyl group of galacto-

nate.

We performed unbiased simulations with the *apo* outward-facing structure in pres-

ence of 100 mM galactonate in solution. Before starting the MD simulations, D46 and E133

were protonated or deprotonated individually. For each system, at least one of the two key

residues was protonated (five replicates were used). We observed spontaneous galacto-

nate-binding events in two simulations in which both D46 and E133 were protonated and in

one simulation in which only E133 was protonated. In all cases, opening of the extracellular

gate preceded the entry of galactonate into the binding site (Fig. 3d).

Substrate binding induces closure of the extracellular gate

Unbiased simulations of outward-facing DgoT with bound galactonate revealed the reduced

flexibility of gating helices TM1 and TM7 (Fig. 2c) and the extracellular gate is more likely to

be in a closed conformation (Fig. 2e). Although the gate could close regardless of the pro-

tonation state of key residues in substrate-bound DgoT, the equilibrium was shifted toward

the conformation with closed extracellular gate in simulations with only E133 protonated or

with both D46 and E133 protonated (Fig. 2e).

Protonation of E133 permits the R47 side chain to simultaneously interact with the

carboxyl groups of galactonate and E133, resulting in a more compact arrangement of the

charged residues (Fig. 2c). Substrate interaction with residues on both gating helices facili-

tates bending of the TM7 extracellular segment toward TM1. Hydrophobic interactions also

play a role in forming occluded conformations. In the outward-facing occluded state, the

substrate-binding site is separated from the extracellular solution by hydrophobic residues

located in TM1, TM2, and TM7 (F72, F274, and W277, respectively; Fig. 3a). When the

extracellular gate opens, the position of the side chains of the three hydrophobic residues

changes to allow more water molecules to flow between TM1 and TM7 (Fig. 3b) and galac-

tonate to enter the binding site (Fig. 3c). This resembles a "clamp-and-switch" mechanism

(Quistgaard et al. 2016), in which occlusion of the binding site precedes a rocker-switch-type

rotation on the N- and C-terminal domains, causing the binding site to become exposed to

the opposite side (Qureshi et al. 2020).

Structural basis of substrate release

We next ran unbiased simulations with the inward-facing structure of DgoT after placing a

galactonate molecule into the binding site. The position for the substrate molecule was de-

termined by aligning the crystal structures of the *apo* protein in the inward-facing state (PDB

ID: 6E9N) with the galactonate-bound outward-facing state (PDB ID: 6E9O). Galactonate in

solution is predicted to have a pKa of 3.9 and, therefore, should bind to outward-facing DgoT

in its deprotonated form. However, we also considered the possibility that galactonate could

be protonated in the binding pocket due to changes in the local environment and then re-

leased in its protonated form. In 7 out of 38 individual 500 ns unbiased MD simulations, we

observed spontaneous galactonate release (Supplementary Table 1). In all cases, the sub-

strate passed between TM4 and TM10, which act as gating helices on the intracellular side

of the protein. The intracellular segments of these helices (around residues 139–143 and

372–375, respectively) showed a high degree of flexibility in these simulations. The side

chains of F137 and W373 formed a hydrophobic lock similar to the one at the extracellular

gate (Fig. 4a), and galactonate could only pass through this neck region if a water-filled pore

was formed between the side chains of F137 and W373 (Fig. 4b, c).

All spontaneous release events were observed in systems containing deprotonated

D46, which suggests that the proton must be released from this residue prior to galactonate

unbinding. Fig. 4d shows probability densities for intracellular gate opening (measured as

the minimum distance between the side chains of F137 and W373) for various protonation

and substrate-binding states. The intracellular gate mostly assumed a closed conformation

in the substrate-bound double protonated system. Deprotonation of D46 favored the open

state of the gate in systems with a protonated or deprotonated substrate. When galactonate

is protonated, the intracellular gate has limited flexibility and becomes locked in the open

state, when only E133 is protonated, and in the closed state, when only D46 is protonated.

After release of the substrate and both H^+ ions, the intracellular gate closes, permitting re-

orientation of the *apo* transporter to the outward conformation. We conclude that D46 depro-

tonation increases the probability of galactonate release from the inward-facing DgoT by

promoting the opening of the intracellular gate.

Proton release from inward-facing DgoT

Galactonate spontaneously dissociated from inward-facing DgoT only when D46 was depro-

tonated (Supplementary Table 1). However, in MD simulations of DgoT in which both D46

and E133 were protonated, the D46 carboxyl group was sequestered, making direct depro-

tonation to the intracellular solution unlikely (Fig. 5a). In addition, the electrostatic potential

(calculated using *g_elpot* (Kostritskii et al. 2021; Kostritskii and Machtens 2023)) was higher

on the E133 carboxyl group than on D46 (Fig. 5b), indicating that E133 is likely the first

residue to release its proton. These observations suggest D46 donates its proton to E133

after the latter had released its proton.

Proton transfer from D46 requires a conformational change of its side chain, so as to

be in close proximity to E133; therefore, we calculated the free energy associated with con-

formational change as a function of the N-Cα-Cβ-*Cγ* (χ_1) dihedral angle of D46 using well-

tempered metadynamics (WT-MTD) (Tribello et al. 2014). Fig. 5c shows the presence of a

minimum at χ_1 = -70°, corresponding to the conformation sampled in unbiased MD simula-

tions. However, another minimum is present (hereafter denoted as *close*-D46, characterized

by χ_1 = +70°), which has a higher free energy (by ~8 kcal mol^{-1}) and is separated by a ~12

kcal mol^{-1} barrier. In this conformation, the carboxyl group of D46 is H-bonded to E133,

enabling proton transfer to take place.

E133 may release its proton to either the solvent or galactonate (Fig. 5a). Hydrated

excess protons can reorganize the surrounding water wires and facilitate further proton

translocation (Peng et al. 2015; Li and Voth 2021); therefore, we first studied proton transfer

from E133 to the adjacent water molecule to form a hydronium ion. Quantum mechanics/mo-

lecular mechanics (QM/MM) thermodynamics integration (TI) calculations showed that the

free energy barrier can be as small as 8 kcal mol^{-1} (Supplementary Fig. 5). Moreover, in the

presence of the ion pair between the negatively charged E133 and the adjacent hydronium

ion, the surrounding water molecules connect the excess proton with both galactonate and

the solution. To represent the diverse range of possible H-bond networks, we selected seven

structures (see Methods) for use in unbiased QM/MM MD simulations in which the quantum

part was treated at the density functional theory (DFT)-BLYP level (Becke 1988). In simula-

tions with three of the selected structures, the galactonate carboxyl group accepted the pro-

ton after a few picoseconds, either via water molecules only or via water molecules along

with the substrate hydroxyl group (Supplementary Figure 6 and Supplementary Table 2). In

simulations with another three structures, the proton was transferred to one of the molecules

of the water wire, without substrate protonation. In simulations with the seventh structure,

the proton was shared between the substrate and a neighboring water molecule. Therefore,

despite the limited statistics and underestimation of the proton transfer barriers due to the

BLYP functional (Mangiatordi et al. 2012), we suggest that proton transfer to galactonate or

the solvent is viable over a sub-nanosecond timescale. The released proton might eventually

leave the protein either bound to the substrate or through the solvent via a Grotthuss mech-

anism.

Finally, we performed QM/MM TI calculations to study the energetics associated with

a proton transfer from *close*-D46 to E133. There was a small free energy barrier (around

4 kcal mol^{-1}) associated with D46 deprotonation (Fig. 5d). Given the overall moderate en-

ergy cost of all processes involved (Supplementary Fig. 7), we conclude that the mechanism

of D46 deprotonation studied here, involving first proton transfer from E133 to the solvent

and/or galactonate and subsequent proton transfer from D46 to E133, appears feasible.

Free energy landscapes suggest that reorientation of the empty transporter is a rate-

limiting step of the cycle

Both inward- and outward-facing DgoT conformations display some flexibility; however, we

did not observe transitions between these two states in single trajectories. Since a quantita-

tive description requires multiple observations of these slow processes, we used Markov

state modeling (Husic and Pande 2018) to stitch together short simulation data and analyze

the transition probability between all conformational states. We featurized the trajectory data

using a set of interdomain Cα atomic distances and applied time-lagged independent com-

ponent analysis (tICA) to find the slowest components in the dataset. The first tICA-eigen-

vector discriminates between the inward- and outward-facing conformations (Fig. 6a), and

the second correlates with the (open or closed) state of the extracellular gate (Fig. 6b, Sup-

plementary Fig. 8). Two systems were chosen as most relevant for transport: (i) galactonate-

bound DgoT with protonated D46 and E133 (responsible for translocation of the substrates

across the membrane) and (ii) *apo* DgoT with deprotonated D46 and E133 (responsible for

reorientation of the empty transporter after substrate release). For both systems, we sam-

pled all relevant intermediate conformations with a total of ~26 μs (substrate-bound) and

~35 μs (*apo*) unbiased MD simulations, and constructed Markov state models. (Fig. 6a, b).

The free energy surfaces for both systems revealed a high energy barrier separating the

inward- and outward-facing conformations. In the presence of galactonate, the barrier was

lowered and the inward-facing state was the most energetically favorable. In contrast, the

system with the *apo* protein preferred outward-facing conformations such that DgoT favors

inward galactonate transport. Moreover, *apo* DgoT easily switches between outward-occluded and outward-open states, whereas substrate-bound DgoT adopts only occluded conformations. Perron-cluster cluster analysis (PCCA) (Deuflhard and Weber 2005), used to identify metastable states, revealed a difference in the direction of conformational changes between the *apo* and galactonate-bound states (Fig. 6c, d), as previously observed in the free energy surfaces (Fig. 6a, b). For substrate-bound DgoT, metastable states correspond to inward-facing, intermediate occluded, and outward-occluded states, with the highest probability for the inward conformation (Fig. 6c). For *apo* DgoT, PCCA identified inward-facing, outward-occluded, and outward-open states (lower to higher probability). The outward-open state was not adopted by substrate-bound DgoT: it was only adopted by *apo* DgoT, for which it represents the state with highest stationary probability (Fig. 6d). Given that substrate binding and release are fast processes compared with the conformational changes, our data show that reorientation of the empty transporter upon substrate release is the rate-limiting step in the transport cycle.

Neutralization of putative proton acceptors abolishes galactonate transport

To further investigate the function of the charged transmembrane residues, we next evaluated the effects of the corresponding neutralization mutations (D46N, E133Q, R47Q, and R126Q) on DgoT transport activity. We first monitored changes in the extracellular pH of a bacterial suspension using a micro pH electrode-based transport assay (Bosshart et al. 2019). The addition of galactonate to *E. coli* expressing WT DgoT resulted in a time-dependent increase in pH, which reached the maximum value in less than a minute. The addition of the same concentration of the galactonate epimer gluconate to DgoT-expressing *E. coli* nor of galactonate to vector-transformed (without DgoT) bacteria induced an increase in pH (Fig. 7a). All mutations tested abolished the galactonate-induced pH changes (Fig. 7b), indicating loss of transport function.

In SSME experiments, the application of galactonate elicited a positive fast pre-steady state, but not transport currents, for D46N and E133Q DgoT, indicating that these mutant transporters can bind galactonate but cannot complete the transport cycle. Galactonate-induced pre-steady state currents recorded with the E133Q mutant were pH independent (Fig. 7c), suggesting that protonation of this residue is responsible for the inhibition of WT DgoT at alkaline pH. At every pH tested, E133Q currents closely resembled WT currents under acidic pH conditions (Supplementary Fig. 9a), where transport is blocked by impaired proton release inside the vesicle. In contrast, D46N currents were pH dependent, with the largest peak currents observed under acidic pH conditions (Fig. 7d). Currents obtained with this mutant were biphasic (Supplementary Fig. 9b) and with slower decay than for E133Q. To determine whether the recorded currents represent the pre-steady state reaction or residual transport activity, we compared the time courses of current decay using liposomes reconstituted with different LPRs. Unlike the currents recorded with WT protein, the decay times for D46N DgoT did not systematically depend on the LPR (Supplementary Fig. 9c). Therefore, the observed reaction can be attributed to a slow pre-steady state process.

For D46N and E133Q DgoT, the peak current amplitudes changed with increasing galactonate concentrations in a saturating fashion (Supplementary Fig. 10), suggesting that these mutant transporters undergo conformational changes after rapid galactonate binding (Bazzone et al. 2021). Similar behavior was observed for WT DgoT under acidic conditions, where only the pre-steady state reaction (no transport activity) can be seen (Fig. 1a, Supplementary Fig. 3a). Fig. 7e shows that the observed rates (obtained as reciprocal decay time constants) depend on the galactonate concentration. Hyperbolic curve fitting provided the K_{DS} for galactonate binding, as well as the kinetic parameters characterizing the conformational changes (Table 1). Compared with WT, conformational changes are similar for E133Q and slower for D46N but with no difference in substrate affinity. We conclude that

deprotonation of D46 and E133 is necessary to complete the transport cycle; however, partial reactions that represent conformational changes can occur in the presence of galactonate.

R47 couples substrate binding and major conformational changes

For neutralizing mutations of both transmembrane arginine residues (R47Q and R126Q), galactonate application elicited fast negative currents in SSME experiments (Fig. 7f, g). For R126Q, current amplitudes were similar at pH 5–7, but lower at more alkaline pH. Peak currents changed with increasing galactonate concentrations with a hyperbolic concentration dependence (Fig. 7e), indicating conformational changes. In contrast, R47Q DgoT exhibited faster currents with amplitudes increasing linearly with increasing substrate concentration (Fig. 7h). Therefore, the observed pre-steady state reaction for R47Q differs from those observed for other transport-deficient mutants and likely represents electrogenic substrate binding rather than conformational changes (Bazzone et al. 2021).

The crystal structure of outward-facing DgoT (Leano et al. 2019), as well as our MD simulation data (Fig. 2), indicates that R47 interacts with the carboxyl group of bound galactonate. To better understand the role of this arginine residue, we carried out MD simulations with the outward-facing DgoT structure in which R47 was mutated to glutamine and D46 and E133 were protonated. In *apo* simulations, both WT and mutant DgoT assumed similar conformations with an open extracellular gate (Fig. 8a). However, galactonate failed to promote closure of the extracellular gate of R47Q DgoT (Fig. 8b). This difference is a consequence of changes in protein–substrate interactions. Galactonate is coordinated by multiple polar residues (Batarni et al. 2023) in helices from both the N- and C-domains, with the substrate located closer to TM7 in simulations with the mutant than with the WT transporter (Fig. 8c, d). Comparison of the substrate–protein interactions revealed more stable contacts for galactonate with C-domain residues such as Q264, T372, and N393 in WT; residue 47 from

the N-domain interacts with the substrate only in simulations with the WT protein (not the

mutant protein) (Fig. 8e).

Our combined experimental and computational results describe substrate binding as

a multiphasic process initiated by galactonate recognition and accommodation in the binding

site. Subsequent direct interaction between the carboxyl group of galactonate and R47 in-

duces closure of the extracellular gate. Both reactions are electrogenic; R47Q abolishes the

second step without affecting the initial substrate binding. The fast negative component cor-

responding to electrogenic galactonate binding also occurs with other mutants, along with

an additional slower component of positive (E133Q) or negative (R126Q) amplitude (Fig.

7c, f).

Discussion

We combined experimental and computational approaches to describe the transport cycle
of the bacterial SLC17 homolog DgoT. DgoT transport is based on an alternating access
mechanism, with the transporter cycling between inward- and outward-facing states through
occluded conformations in which the substrate-binding site is inaccessible from both mem-
brane sides (Fig. 9). In the outward-facing conformation, DgoT dynamically switches be-
tween states with a closed or open extracellular gate (states 1 and 2). Protonation of both
D46 and E133 stabilizes the open-gate conformation (state 2) and permits galactonate bind-
ing from the periplasmic side (state 3), followed by closure of the extracellular gate. For-
mation of the outward-occluded conformation is the initial step of major conformational
change (state 4) that brings the protein into an inward-facing conformation (states 5, 6, and
7). In this conformation, deprotonation of D46 (state 5) results in an open intracellular gate
that permits galactonate release (state 6). D46 deprotonation may occur via multiple path-
ways, as shown for proton transfer in other transporters (Swanson 2022). Our multiscale
simulations demonstrate that D46 deprotonation is possible via initial proton release from
E133, either to the intracellular solvent or to galactonate, followed by H^+ transfer from D46
to E133 (state 6). Galactonate can unbind in a protonated or non-protonated form; protona-
tion might promote galactonate release by weakening the electrostatic interaction with R47
and stabilizing intracellular gate opening. H^+ transfer from D46 to E133 (state 6) and subse-
quent release from E133 results in intracellular gate closure (state 7); reorientation of the
empty transporter to outward-facing conformation completes the cycle (state 8).

   Our classical MD simulations revealed how the protonation state of the two acidic
transmembrane residues is coupled to the conformational changes necessary for substrate
binding and release. In the *apo* state, deprotonated D46 and deprotonated E133 both serve
as "latches" by interacting with R126 (TM4) and R47 (TM1), respectively; this strategy is

commonly used by MFS transporters (Drew et al. 2021). Protonation of the two acidic residues unclasps these latches, albeit with distinct effects on extracellular gate mobility (Fig. 2d and Fig. 3d). For D46, H^+ association releases R126 and shifts the equilibrium toward a state with an open extracellular gate, whereas protonation of E133 alone stabilizes the gate in an only slightly opened state. Double protonation is necessary for full opening of the extracellular gate, which makes substrate binding most effective (Fig. 2d and Fig. 3d). Galactonate binding promotes closure of the extracellular gate for all four protonation state combinations (Fig. 2e). However, stable gate closure requires protonation of E133 (Fig. 2e), which disrupts the E133–R47 salt bridge and gives TM1 the flexibility to come into closer contact with TM7 upon galactonate binding (Fig. 2c).

Arginine residues at positions 47 and 126 are conserved within SLC17 transporters (Supplementary Fig. 2). R47 forms an electrostatic interaction with galactonate in the binding site that is suggested to be responsible for its recognition (Leano et al. 2019). However, R47Q DgoT exhibits galactonate-specific pre-steady state currents (Fig. 7g) that represent a binding reaction rather than rate-limiting conformational changes. Therefore, our results assign a role beyond substrate recognition to R47: this arginine connects the substrate-binding site to the protonation site and triggers closure of the extracellular gate, much like H322 in lactose permease LacY (Kumar et al. 2015). Pre-steady-state currents of R126Q DgoT indicate electrogenic conformational changes rather than a binding reaction. R126 might be involved in tuning the pKa of D46 at different stages of the transport cycle or be necessary for conformational changes in *apo* DgoT.

The proposed transport mechanism (Fig. 9) fully agrees with all published experimental data. It accounts for the stoichiometrically coupled symport of two H^+ and one galactonate, as determined experimentally (Fig. 1e). Solubilized DgoT does not bind protons after the addition of galactonate (Fig. 1f), as would be expected if protonation of the transporter

precedes substrate binding. Neutralization of residue D46 or E133 prevents coupled galactonate transport, as expected for the removal of obligatory protonation sites. Our transport scheme can also account for recently reported experiments demonstrating that E133Q, but not D46N, DgoT is capable of galactonate exchange (Batarni et al. 2023). The E133Q mutation has no effect on galactonate release and substrate exchange: it exclusively abolishes forward transport by preventing *apo* translocation back to the outward-facing conformation. In contrast, D46N prevents forward transport as well as galactonate exchange, since deprotonation of D46 is required for galactonate release (Fig. 4). The surprising result that the additional E133Q mutation restores galactonate exchange in D46N/E133Q DgoT might be explained if galactonate is predominantly released in a protonated form by WT DgoT. In D46N DgoT, galactonate may still be protonated via H^+ transfer from E133 although the closed intracellular gate would prevent its release (Fig. 4d). In D46N/E133Q DgoT, galactonate cannot be protonated and the slightly higher open probabilities for the intracellular gate with non-protonated galactonate make the double mutant competent for exchange.

E. coli can use galactonate as the sole carbon source (Deacon and Cooper 1977), demonstrating that DgoT transports as effectively as alternative glucose transporters, such as lactose permease LacY (Kaback and Guan 2019) or xylose symporter XylE (Madej et al. 2014). LacY and XylE transport uncharged sugar molecules with a stoichiometry of 1 proton:1 substrate, whereas DgoT co-transports two protons with the negatively charged galactonate, together resulting in a positive net transported charge. Most likely, the adjusted transport stoichiometry permits DgoT to utilize ΔpH and ΔΨ and provides high driving forces, thus optimizing bacteria for nutrient uptake when resources are limited.

Transport substrates and stoichiometries vary substantially among MFS transporters. LacY (Kaback and Guan 2019) and fucose transporter FucP (Dang et al. 2010) share a stoichiometry of 1 H^+:1 substrate. In LacY, protonation of a single site, E325 (equivalent to

E133 in DgoT (Kaback and Guan 2019)), permits lactose binding and transition to the oc-

cluded conformation. FucP has two protonation sites, D46 and E135, that are homologous

to D46 and E133 in DgoT. However, only D46 is accessible from the extracellular solution,

with E135 serving only as part of the proton transfer pathway. SLC17A5/sialin differs from

DgoT in having broad substrate specificity: it recognizes various sialic acids via electrostatic

interaction with two conserved arginines, R168 and R57. Two glutamates, E171 and E175,

serve as protonation sites. Protonation of E171 releases R168 and permits its interaction

with the substrate (Hu et al. 2023). Subsequent H^+ transfer to E175 is followed by substrate

transfer to a more cytoplasmic binding site close to R57 and translocation to the inward-

facing conformation. Such coupling of substrate translocation to proton transfer results in

the electroneutral co-transport of one H^+ and one sialic acid molecule.

The comparison of DgoT with FucP and SLC17A5/sialin reveals how small variations

in the arrangement of protonation sites can adjust transport stoichiometries of H^+-coupled

transporters with conserved architecture and transport mechanisms. The SLC17 family en-

compasses organic anion uniporters (Ishikawa et al. 2008; Iharada et al. 2010), H^+-gluta-

mate exchangers (Kolen et al. 2023), and a H^+-sialic acid symporter (Hu et al. 2023). Certain

SLC17 transporters can also function as a Na^+-PO_4^{3-} symporter (Iharada et al. 2010;

Preobraschenski et al. 2018). Therefore, our work may serve as a framework to understand

the mechanisms underlying the diversity of SLC17 transport mechanisms.

Methods

Expression and protein purification

The full-length DgoT gene (GenBank accession number AKK15832.2) was subcloned into a pQE60 vector through the NcoI and HindIII restriction sites in fusion with a C-terminal thrombin cleavage site and decahistidine tag. Mutant constructs were generated using PCR-based mutagenesis and verified by DNA sequencing. Protein expression and purification were performed using a published procedure, with modifications (Leano et al. 2019). *E. coli* C41 cells transformed with pQE60 DgoT WT were grown at 37 °C in TB medium supplemented with 2 mM $MgSO_4$. When an OD_{600} of 0.6–0.8 was reached, gene expression was induced with 1 mM IPTG and cells were grown for a further 4 h (typical yield of 15 g per 1 L culture). After sedimentation, cells were flash frozen in liquid nitrogen and stored at -80 °C for later use, Next, cells (15 g) were resuspended in 20 mM Tris (pH 7.4) and 300 mM NaCl (50 mL volume) containing complete protease inhibitor cocktail (Roche) and lysed by sonication. Debris was removed by centrifugation at 12,000 × *g* for 15 min, and membranes were collected at 200,000 × *g* for 1 h, flash frozen in liquid nitrogen, and stored at -80 °C until use.

A frozen membrane pellet from 15 g cells was resuspended in 20 mL membrane buffer (20 mM Tris (pH 7.4), 150 mM NaCl) containing cOmplete protease inhibitor cocktail, using a glass Dounce homogenizer, and then *n*-dodecyl-D-maltoside (DDM) was added to 1.4%. Membranes were solubilized for 2 h at 4 °C and the insoluble fraction was removed by ultracentrifugation at 75,000 g for 30 min at 4 °C. The supernatant was diluted 1:2 with solubilization buffer, imidazole was added to 15 mM, and pH was adjusted to 7.8–8.0. CoNTA resin (3 mL) was washed with 10 column volumes (CV) of membrane buffer, added to the supernatant, and incubated at 4 °C for 1.5 h under gentle agitation. The resin was then washed with 10 CV of wash1 buffer (20 mM Tris, 150 mM NaCl, 15 mM imidazole, 0.1% DDM, pH 8.0) and 10 CV of wash2 buffer (20 mM Tris, 150 mM NaCl, 0.1% lauryl

maltose neopentyl glycol (LMNG), pH 7.4). Protein was eluted with 4 CV of elution buffer

(20 mM Tris, 150 mM NaCl, 0.1% LMNG, 150 mM imidazole, pH 7.4) into 0.5 CV fractions,

and 5 mM EDTA was added immediately after collection. Protein concentration in the eluted

fractions was estimated by measuring the absorbance at 280 nm (NanoDrop), and fractions

with the highest protein concentrations were combined. Imidazole was removed using a PD-

10 desalting column, with protein eluted with buffer (20 mM Tris, 150 mM NaCl, 1 mM EDTA,

0.05% LMNG, pH 7.4) and stored overnight at 4 °C.

The purified protein was concentrated to 5–8 mg/ml using a 50 kDa molecular weight

cutoff (MWCO) centrifuge concentrator (Millipore) and loaded in 0.5 mL portions onto a Su-

perdex 200 Increase 10/300 GL column (GE Healthcare Life Sciences) preequilibrated with

size-exclusion chromatography (SEC) buffer (20 mM HEPES, 150 mM NaCl, 0.05% LMNG,

pH 7.4). The peak fractions were combined, flash frozen in liquid nitrogen, and stored at -

80 °C until use.

Reconstitution of proteoliposomes

E. coli polar lipids (Avanti Polar Lipids, *E.coli* polar lipid extract, 25 mg/ml solution in chloro-

form) were dried under nitrogen and then under vacuum overnight. The dried lipid film was

resuspended to a lipid concentration of 10 mg/ml by stirring in reconstitution buffer (1 mM

HEPES pH 7.4, 150 mM NaCl, 2 mM $MgSO_4$) for 1 h at room temperature (RT). After the

lipids were completely dissolved, the suspension was frozen in liquid nitrogen, stirred for

another 1 h at RT, and sonicated using a UP50H ultrasonic processor equipped with a mi-

crotip until clear (2 or 3 cycles of 30 s). The formed liposomes were destabilized by adding

0.6% Triton X-100 and incubated for 45 min at RT under gentle agitation. Purified DgoT was

added to the destabilized liposomes at a LPR of 5, 10, or 20 and then incubated for 1 h at

4 °C. To remove the detergent, 150 mg SM2 Bio-beads was added per 1 mL liposome sus-

pension. After incubation for 1 h at 4 °C, another 150 mg Bio-beads was added per 1 mL

suspension, followed by another 1 h incubation at 4 °C, and then the beads were removed

using a disposable column. A third volume of Bio-beads was then added (400 mg per 1 mL

suspension), incubated overnight at 4 °C, and removed using a disposable column. Empty

liposomes were prepared in parallel using the same procedure but with no added protein.

pH electrode-based transport measurements

DgoT was expressed in *E. coli* C41 cells as described for protein purification. After protein

expression, cells were pelleted, washed twice in assay solution (250 mM KCl, 1 mM MgSO$_4$,

2 mM CaCl$_2$), and resuspended in assay solution to an OD$_{600}$ 15. All centrifugations were

performed at 2,200 × *g* for 7–8 min, and cells were resuspended by vortexing. Substrates to

be assayed were dissolved in assay solution at a concentration of 160 mM and the pH was

adjusted to 6.5–6.7 with KOH such that the substrate pH was lower than the pH of the bac-

terial suspension for each experiment. A 800 µL volume of bacterial suspension was trans-

ferred to a 2 ml reaction tube (Eppendorf) and the pH of the suspension (i.e., of the extra-

cellular medium) was measured using a micro pH electrode with an integrated temperature

sensor (Xylem, SI Analytics) under constant stirring and adjusted to pH 6.7 using KOH and

HCl. After recording the baseline for 60 seconds, 50 µL of the corresponding compound was

added to the bacterial suspension (final concentration: 10 mM). All experiments were per-

formed at 20–21 °C. For experiments with DgoT in detergent, purified DgoT (stored at -

80 °C) was thawed and SEC buffer was replaced using a PD-10 desalting column with assay

solution containing 150 mM NaCl and 0.05% LMNG (pH ~7.4). Purified protein was used at

a concentration of 4 µM in all experiments. The experiment was repeated six times, with

comparable results.

SSM-based electrophysiology

Gold electrode sensors (1 or 3 mm) were prepared as previously described (Bazzone et al.

2017). Briefly, sensors were incubated for at least 30 min in an octadecane thiol solution,

and then rinsed thoroughly with isopropanol and water. The solid-supported membrane

(SSM) was prepared by pipetting 1.5 μL diphytanoyl phosphatidylcholine dissolved in *n*-dec-

ane onto the electrode surface, followed by 100 μL aqueous buffer. Immediately prior to

measurements, liposome samples were thawed, diluted to a final lipid concentration of

1 mg/mL and briefly sonicated. Each liposome sample (10 μL) was pipetted onto a SSM

sensor and adsorbed by centrifugation at 2,200 × *g* for 30 min at RT. All experiments were

repeated for at least three sensors, with each condition measured at least twice. All solutions

were buffered in 100 mM potassium phosphate (KPi) for each pH used.

For measurements with a single solution exchange protocol, three phases of 1 s du-

ration were applied: flow of nonactivating (NA) solution, activating (A) solution, and NA so-

lution. Only the A solution contained galactonate. In experiments with variable pH, pH of NA

and A solutions was kept constant within a single experiment. Between experiments using

different pH values, the sensor was incubated at the new pH for 5 min to equilibrate the

intraliposomal pH. In experiments with WT, D46N, E133Q, and R126Q DgoT, the NA solution

contained gluconate to compensate for galactonate in the A solution. For R47Q DgoT, glu-

tamate was used in NA solution instead, allowing a direct comparison of the responses to

galactonate and gluconate applications.

To determine the apparent pK values, normalized peak currents (measured using

3 mm sensors) were fitted with one of the following equations:

acidic deactivation (pH ≤ 8): $I_{norm}(pH) = \dfrac{I_{max}}{1 + 10^{pKa-pH}}$;

alkaline deactivation (pH ≥ 7.5): $I_{norm}(pH) = \dfrac{I_{max}}{1 + 10^{pH-pK_b}}$.

Currents were normalized to the I_{max} value obtained by the fit of peak currents measured

on the same sensor.

For the analysis of pre-steady state currents, 1 mm sensors were used. Rate con-

stants for the observed charge displacements ($k_{obs}= 1/\tau$) were derived from the transient

currents by fitting the decay with a monoexponential function $I = A * \exp(-t/\tau)$. This two-step

reversible reaction can be described as follows:

$$P + S \leftrightarrow PS \underset{k^-}{\overset{k^+}{\rightleftarrows}} P^*S,$$

with K_D above $P + S \leftrightarrow PS$ and k^+ above the rightward arrow.

where P is the protein, S is the substrate, and P^* is the protein after conformational changes.

The first step is the binding reaction described by the dissociation constant K_D, and the

second step is the substrate-induced conformational change characterized by the forward

and reverse rate constants. Assuming that substrate binding is rapid, the observed rate con-

stant has hyperbolic dependence on the substrate concentration (Smirnova et al. 2006):

$$k_{obs} = k^- + k^+ \frac{[gal]}{[gal]+K_D}.$$

Measurements under asymmetrical pH conditions were done using a double solution

exchange configuration. A resting (R) solution phase of 1 s and incubation period of 5–

20 min were added to the beginning of each measurement to allow the intraliposomal pH to

adjust to the pH of the R solution. Afterwards, a normal single solution exchange protocol

was used to establish the pH gradient (during NA phase) and substrate gradient (A phase).

To determine the transport stoichiometry, we used a reversal assay as previously

described (Thomas et al. 2021), with a single solution exchange protocol comprising three

phases (NA, A, and NA) extended to 2, 3, and 3 s, respectively. Between measurements,

the sensor was rinsed 3–5 times with NA solution; the current responses were recorded and

used as a baseline. The same protocol was used for samples with empty liposomes to ac-

count for solution exchange artifacts. The entire A phase was integrated to obtain the trans-

ported charge values. After subtraction of the negative control (integrated current recorded

with empty liposomes), the transported charge values were used to determine the transport

stoichiometry. The internal (NA) solution used for this experiment had a pH of 7.3 and con-

tained 0.5 mM galactonate and 7.5 mM gluconate, and applied external (A) solutions had a

pH of 7.6 and contained X mM galactonate and 8–X mM gluconate. This setup resulted in

$d\mu_{gal}/d\mu_{H+}$ values of 0–4.

MD simulations

DgoT protein coordinates obtained from the Protein Data Bank (inward-facing, PDB ID:

6E9N; outward-facing galactonate-bound, PDB ID: 6E9O (Leano et al. 2019)) were used as

the starting coordinates for MD simulations. The D46 and E133 protonation state and sub-

strate occupation were modified as described in the text. Standard protonation states at

neutral pH were assigned to all other residues (deprotonated aspartate and glutamate resi-

dues, singly protonated histidine residues) except for H56, which forms a salt bridge with

E180 and, therefore, was set as doubly protonated. Missing residues (235–242 in the in-

ward-facing structure, 231–243 and 277–290 in the outward-facing structure) were modeled

using the SWISS MODEL server (Waterhouse et al. 2018) with a final residue range of 27–

442 used for both protein structures. The N- and C-termini were capped with neutral acetyl

and methylamide groups, respectively. For modeling DgoT single point mutations, we used

PyMOL (Schrödinger, LLC 2015).

The initial protein orientation within the membrane was set to the corresponding DgoT

structure available in the Orientations of Proteins in Membranes database (Lomize et al.

2012). The protein was then embedded into a phosphatidylcholine (POPC) bilayer using

g_membed (Wolf et al. 2010) in GROMACS and solvated in a box with dimensions

~120 × 120 × 100 Å, which was chosen to ensure a minimum distance between periodic

copies of at least 30 Å. The simulation temperature was 310.15 K. The protein/membrane

system was surrounded by a ~100 mM solution of Na^+ and Cl^- ions. Ions were described

using default CHARMM parameters, and the CHARMM TIP3P model was used for water

molecules. MD simulations were performed using the GROMACS software package (versions 2018, 2020, and 2022) (Abraham et al. 2015) with a CHARMM36m force field (Huang et al. 2017). Galactonate parameters were obtained using the SwissParam server (Zoete et al. 2011) and added to the forcefield. An integration time step of 2 fs was used. Van der Waals interactions were calculated with the Lennard–Jones potential and a cutoff radius of 1.2 nm, with forces smoothly switched to zero in the range of 1.0–1.2 nm and no dispersion correction applied. Electrostatic interactions were calculated by the particle mesh Ewald method (Essmann et al. 1995), with a real-space cutoff distance of 1.2 nm. All simulations were done in an isothermal–isobaric ensemble, with the temperature set to 310 K using a v-rescale thermostat (Bussi et al. 2007) and a time constant of 0.5 ps. The thermostat was applied separately to the protein, lipid bilayer, and aqueous solution containing ions. The same groups were used for the removal of the center-of-mass linear motion.

The protein was equilibrated in three steps using the velocity-rescale thermostat and Berendsen (Berendsen et al. 1984) pressure coupling. The first step lasted 50 ns and was run with positional restraints on protein atoms with a harmonic potential with a force constant of 1000 kJ mol^{-1} nm^{-2} to allow for equilibration of water and ions. In the second step, only the backbone atoms of the protein were restrained to enable side chains to equilibrate. Lastly, the system was equilibrated for 1 ns without positional restraints to obtain the velocities used in the following production runs. Production MD simulations used Parrinello–Rahman (Parrinello and Rahman 1981) pressure coupling in a semi-isotropic manner with a time constant of 0.5 ps.

Classical WT-MTD simulations

The PLUMED v2.8.1 plugin (Tribello et al. 2014) interfaced with GROMACS v2020.4 (Abraham et al. 2015) was used for simulations, along with MD settings similar to those described for MD simulations. The collective variable used to describe the conformational change of

D46 is the N-Cα-Cβ-$C\gamma$ dihedral angle (χ_1). WT-MTD parameters were used as previously

reported (Chiariello et al. 2020; Chiariello et al. 2021): Gaussian height, 1.2 kJ/mol; Gauss-

ian sigma, 0.35 rad; bias factor, 15; and Gaussian deposition frequency, 500. Employing a

previously described algorithm (Tiwary and Parrinello 2015), we applied reweighting to the

dihedral angle serving as the CV to reconstruct the histogram of the sampled configurations.

WT-MTD simulations were run for 100 ns, which was sufficient to achieve a converged free

energy surface (shown in Supplementary Fig. 11).

QM/MM simulations

QM/MM simulations were carried out using GROMACS v2020.4 (Abraham et al. 2015) and

CPMD v4.3 (Hutter et al. 2000), coupled with MiMiC v0.2.0 (Olsen et al. 2019; Bolnykh et

al. 2019). MiMiC-QM/MM input files were generated using MiMiCPy v0.2.1 (Raghavan et al.

2023). Different QM partitions were used to investigate the different processes. Region QM1

was used to simulate proton transfer from protonated E133 to the immediate neighbor water

molecule and, therefore, encompassed the sidechains of E133, the galactonate molecule,

and the three water molecules bridging these. The initial QM/MM configuration was obtained

from WT-MTD simulations of the system in which both D46 and E133 were protonated. A

snapshot in which E133 and galactonate were connected through water molecules was

identified by visual inspection. Region QM2 included the hydronium ion formed by proton

release from E133, the substrate galactonate, and the first layer of surrounding water mol-

ecules, enabling the simulation of proton migration to either galactonate or solvent. The ini-

tial QM/MM configuration was taken from the window corresponding to the products (CV1 =

0.35 Å) of the first TI proton transfer simulation (see section "TI simulations of proton trans-

fer"). Lastly, QM3 included the sidechains of E133 and D46 to enable the study of proton

transfer from protonated D46 to deprotonated E133. Here, the initial QM/MM configuration

was extracted through visual inspection from the WT-MTD simulation of the system with

protonated D46 and now-deprotonated E133 in order to identify a snapshot with D46 in the

close conformation and H-bonded to E133. The total number of QM atoms for each partition

was 44 (QM1), 82 (QM2), and 18 (QM3).

The QM problem was solved using the DFT (Hohenberg and Kohn 1964; Kohn and

Sham 1965) using the BLYP functional (Becke 1988). Core electrons were described using

norm-conserving pseudopotentials of the Martins−Troullier type (Troullier and Martins 1991),

and the valence electrons were treated explicitly. Monovalent carbon pseudopotentials were

used to saturate the dangling bonds at the boundaries between the QM and MM regions

(von Lilienfeld et al. 2005). Isolated system conditions were achieved using the method of

Martyna and Tuckerman (Martyna and Tuckerman 1999). The MM system was treated with

the CHARMM36m force field (Klauda et al. 2010) for protein, lipid, and ions, and the TIP3P

model for waters, as in the classical MD simulations. Electrostatic interactions between the

QM and MM subsystems were described using the Hamiltonian electrostatic coupling

scheme (Laio et al. 2002; Olsen et al. 2019). The short-range electrostatic interactions be-

tween the QM partition and any MM atom within a cutoff of 30 a.u. from the QM region were

explicitly considered, with a seventh-order multipole expansion of the QM electrostatic po-

tential used for long-range interactions.

Constant temperature simulations were carried out at 300 K using a Nosé–Hoover

thermostat (Nosé 1984; Hoover 1985) (with a coupling frequency of 3500 cm^{-1}).

Initially, all systems underwent relaxation through simulated annealing, with gradual

reduction in the temperature of the QM/MM system by removing excess kinetic energy in

each step (velocity multiplied by a factor of 0.99). Subsequently, systems were linearly re-

heated to 300 K and the target temperature increased using a Berendsen thermostat cou-

pled to the system (coupling strength of 5000 a.u.). A Born–Oppenheimer molecular dynam-

ics (BOMD) scheme was used. A timestep of 10 a.u. (~ 0.24 fs) for the initial geometry opti-

mization using an annealing protocol. Finally, systems with QM1 and QM3 partitions under-

went 20 ps of QM/MM MD equilibration at 300 K with a timestep of 20 a.u. (~0.48 fs) before

running TI simulations; details of the QM/MM MD simulations with the QM2 partition are

provided in the section "QM/MM MD proton migration simulations".

TI simulations of proton transfer

The free energy profiles of the two proton transfers investigated here was determined

through TI in its Blue Moon sampling implementation (Ciccotti and Ferrario 2004; Ciccotti et

al. 2005). The collective variable (CV) used to describe proton transfer from E133[H] to its

adjacent water molecule (CV1) was the difference between (i) the distance between the

carboxylate oxygen atom and the transferred proton of E133[H] and (ii) the distance be-

tween the same proton and the oxygen atom of water. Similarly, the CV2 used to simulate

proton transfer from D46[H] to E133 was the difference between (i) the distance between

the carboxylate oxygen atom and the transferred proton of D46[H] and (ii) the distance be-

tween the same proton and the carboxylate oxygen atom of E133.

A total of 17 and 15 independent constrained MD simulations were conducted for

each TI calculation, respectively, with windows separated by 0.08 Å increments. At each CV

value (either CV1 or CV2), the systems were simulated for 2 ps and the last snapshot was

taken to set up the next window. The constraint force at each CV value was calculated by

averaging the Lagrange multiplier of the Shake algorithm after discarding the first 0.5 ps

(i.e., when it reached convergence). The free energy profiles were then obtained by inte-

grating the constraint force along the CV using the trapezoid method. For both TI profiles,

the correction term was found to negligible because fluctuations in the angle connecting

donor, proton, and acceptor are minimal.

QM/MM MD proton migration simulations

Starting with the final structure (CV1 = 0.35 Å) obtained from the first TI proton transfer

simulation, in which the proton was transferred from E133[H] to the adjacent water molecule,

we ran a 40 ps QM/MM MD simulation with a timestep of 20 a.u. (~0.48 fs), maintaining the

same constraint applied during TI. This served to sample an initial set of structures of the

solvated hydronium ion forming an ion pair with negatively charged E133. We then applied

a cluster analysis approach utilizing the *cluster* module of GROMACS. E133, D46, galacto-

nate, and the first layer of surrounding water molecules (within 4 Å) were considered in the

RMSD calculation, and clustering was performed with the *gromos* method(Daura et al. 1999)

and a 0.04 Å cutoff. This approach resulted in the identification of nine distinct clusters. Sub-

sequently, seven representative snapshots, corresponding to the centroids of the seven

most populated clusters and covering over 95% of the QM/MM MD trajectory, were selected

as starting configurations for the QM/MM MD simulations. Notably, these structures are

characterized by variations in their hydrogen bond networks and the number of connecting

water molecules among E133, hydronium ion, and galactonate. Statistical information per-

taining to the proton transfer dynamics of these QM/MM MD simulations is presented in

Supplementary Table 2.

Quantification of electrostatics

Electrostatics was quantified with g_elpot (Kostritskii et al. 2021; Kostritskii and Machtens

2023); source code, installation instructions, and usage recommendations can be found at

https://jugit.fz-juelich.de/computational-neurophysiology/g_elpot. The distribution of electro-

static potential was calculated via the smooth particle mesh Ewald (SPME) method. For our

system, the SPME potential was calculated on a grid of 256 × 256 × 208 points with an in-

verse Gaussian width β of 20 nm^{-1}. We exploited the extended functionality of the tool

(Kostritskii and Machtens 2023) to calculate the potential for the carboxyl groups of proto-

nated D46 and E133. The electrostatic potential is an average of the SPME potential in a

0.15-nm sphere around the following atom groups (CHARMM naming convention): OD1, OD2, and CG of D46 and OE1, OE2, and CD of E133. Since two selected groups are chemically identical, the residue-specific short-range part of the potential is the same and, therefore, comparison of the potential values is justified. The time course of the electrostatic potential was averaged across each trajectory.

Markov state model construction

Markov state models (MSMs) were constructed from multiple unbiased MD trajectories using PyEMMA 2 software (Scherer et al. 2015). To better capture the movement of N- and C-terminal domains relative to each other, we selected every 5th Cα atom of the TM helices and measured the distances between each selected atom located on the N-terminal (TM1–6) and C-terminal (TM7–12) domains, respectively. This resulted in 27 × 26 = 2809 pairwise distances. To reduce the dimensionality, we used a time-lagged independent component analysis (tICA) with a lag time of 50 ns on a set of unbiased MD simulations to obtain the slowest collective motions. For a description of the major conformational changes, we kept the first two independent components (ICs) because they had slower timescales than the other ICs (Supplementary Fig. 11b, c). Since the starting point of the unbiased simulations was the corresponding crystal structure (either inward- or outward-facing), only regions of constructed conformational space that are close to the starting conformations were sampled. To overcome the sampling gap between unbiased simulations that started with different crystal structures, we identified conformations that were closest to the central region of conformational space and, therefore, should represent different occluded states and initiated new unbiased simulations from them. This process was repeated iteratively until the implied timescales converged. A total of ~25 μs (double protonated substrate-bound DgoT) and ~37 μs (DgoT *apo*) simulation data were obtained and used for further data analysis, with

individual trajectories of 400–800 ns in length (Supplementary Fig. 11a). The implied time-scales plot demonstrates the Markovian behavior after a lag time of ~ 80 ns (Supplementary Fig. 11b, c). For constructing the MSM, a lag time of 100 ns was chosen from the implied timescales plot. For model validation, we performed a Chapman–Kolmogorov test with three metastable states for both substrate-bound and *apo* DgoT systems (Supplementary Fig. 11d, e). Metastable states were identified using Perron-cluster cluster analysis (PCCA) (Deuflhard and Weber 2005). For each system, the implied timescale plots (Supplementary Fig. 11b, c) revealed two slow processes; therefore, we clustered the microstates into three metastable states.

Data analysis and statistics

Each experimental dataset represented as means ± error bars was generated by performing the same experiment on multiple sensors (sensor numbers are given in the respective leg-ends). Error bars represent standard deviations of the average value, and individual data points are given for n < 5. Hydration profiles (Fig. 3c and 4c) were calculated as the number of water molecules in 2 Å sections along the z-coordinate in each frame; average values and standard deviations are plotted. Violin plots were used to visualize the shape of data distribution and highlight trends. For statistical analysis of two groups (Fig. 8e), we used Mann–Whitney test, one-sided; significance is indicated as *p < 0.05.

Data Availability

This study includes no data deposited in external repositories

Acknowledgement

We thank Drs Andre Bazzone, Bassam Haddad, Andrei Kostritskii, Piersilvio Longo and Jan-Philipp Machtens for helpful discussions and Meike Berndt for excellent technical support. This work was supported by the Deutsche Forschungsgemeinschaft (German Research

Foundation) to Ch.F. (FA 301/15–2), P.C. (CA 973/27-2) and M.A.P. (AL 2511/1-2) as part of

Research Unit FOR 2518, DynIon. The authors gratefully acknowledge computing time on

the supercomputer JURECA at Forschungszentrum Jülich under grants dgoth and vglut-pt.

Author contributions

N.D., P.C., M. A.-P., and Ch.F. conceived the study. P.C., M. A.-P., and Ch.F. supervised the

project, D.N. designed, performed and analyzed experiments, D.N., S.G. and C.A. designed,

performed and analyzed simulations, N.D. and Ch.F. wrote the manuscript with inputs from

other authors.

Conflict of interest

The authors declare no competing interests.

References

Abraham MJ, Murtola T, Schulz R, Páll S, Smith JC, Hess B, Lindahl E. 2015. GROMACS:
High performance molecular simulations through multi-level parallelism from laptops to su-
percomputers. SoftwareX. 1–2:19–25. https://doi.org/10.1016/j.softx.2015.06.001

Batarni S, Nayak N, Chang A, Li F, Hareendranath S, Zhou L, Xu H, Stroud R, Eriksen J,
Edwards RH. 2023. Substrate recognition and proton coupling by a bacterial member of
solute carrier family 17. J Biol Chem.:104646. https://doi.org/10.1016/j.jbc.2023.104646

Bazzone A, Barthmes M, Fendler K. 2017. SSM-Based Electrophysiology for Transporter
Research. Methods Enzymol. 594:31–83. https://doi.org/10.1016/bs.mie.2017.05.008

Bazzone A, Tesmer L, Kurt D, Kaback HR, Fendler K, Madej MG. 2021. Investigation of
sugar binding kinetics of the E. coli sugar/H+ symporter XylE using solid-supported mem-
brane-based   electrophysiology.   J   Biol   Chem.   298(2):101505.
https://doi.org/10.1016/j.jbc.2021.101505

Becke AD. 1988. Density-functional exchange-energy approximation with correct asymptotic
behavior. Phys Rev Gen Phys. 38(6):3098–3100. https://doi.org/10.1103/physreva.38.3098

Berendsen HJC, Postma JPM, van Gunsteren WF, DiNola A, Haak JR. 1984. Molecular
dynamics   with   coupling   to   an   external   bath.   J   Chem   Phys.   81(8):3684–3690.
https://doi.org/10.1063/1.448118

Bolnykh V, Olsen JMH, Meloni S, Bircher MP, Ippoliti E, Carloni P, Rothlisberger U. 2019.
Extreme Scalability of DFT-Based QM/MM MD Simulations Using MiMiC. J Chem Theory
Comput. 15(10):5601–5613. https://doi.org/10.1021/acs.jctc.9b00424

Bosshart PD, Kalbermatter D, Bonetti S, Fotiadis D. 2019. Mechanistic basis of L-lactate
transport in the SLC16 solute carrier family. Nat Commun. 10(1):2649.
https://doi.org/10.1038/s41467-019-10566-6

Bussi G, Donadio D, Parrinello M. 2007. Canonical sampling through velocity rescaling. J
Chem Phys. 126(1):014101. https://doi.org/10.1063/1.2408420

Chiariello MG, Alfonso-Prieto M, Ippoliti E, Fahlke C, Carloni P. 2021. Mechanisms Under-
lying Proton Release in CLC-type F−/H+ Antiporters. J Phys Chem Lett. 12(18):4415–4420.
https://doi.org/10.1021/acs.jpclett.1c00361

Chiariello MG, Bolnykh V, Ippoliti E, Meloni S, Olsen JMH, Beck T, Rothlisberger U, Fahlke
C, Carloni P. 2020. Molecular Basis of CLC Antiporter Inhibition by Fluoride. J Am Chem
Soc. 142(16):7254–7258. https://doi.org/10.1021/jacs.9b13588

Ciccotti G, Ferrario M. 2004. Blue Moon Approach to Rare Events. Mol Simul. 30(11–
12):787–793. https://doi.org/10.1080/0892702042000270214

Ciccotti G, Kapral R, Vanden-Eijnden E. 2005. Blue moon sampling, vectorial reaction coor-
dinates, and unbiased constrained dynamics. Chemphyschem Eur J Chem Phys Phys
Chem. 6(9):1809–1814. https://doi.org/10.1002/cphc.200400669

Dang S, Sun L, Huang Y, Lu F, Liu Y, Gong H, Wang J, Yan N. 2010. Structure of a fucose
transporter in an outward-open conformation. Nature. 467(7316):734–738.
https://doi.org/10.1038/nature09406

Daura X, Gademann K, Jaun B, Seebach D, van Gunsteren WF, Mark AE. 1999. Peptide
Folding: When Simulation Meets Experiment. Angew Chem Int Ed. 38(1–2):236–240.
https://doi.org/10.1002/(SICI)1521-3773(19990115)38:1/2<236::AID-ANIE236>3.0.CO;2-M

Deacon J, Cooper RA. 1977. D-Galactonate utilisation by enteric bacteria. The catabolic
pathway in Escherichia coli. FEBS Lett. 77(2):201–205. https://doi.org/10.1016/0014-
5793(77)80234-2

Deuflhard P, Weber M. 2005. Robust Perron cluster analysis in conformation dynamics. Lin-
ear Algebra Its Appl. 398:161–184. https://doi.org/10.1016/j.laa.2004.10.026

Drew D, North RA, Nagarathinam K, Tanabe M. 2021. Structures and General Transport
Mechanisms by the Major Facilitator Superfamily (MFS). Chem Rev. 121(9):5289–5335.
https://doi.org/10.1021/acs.chemrev.0c00983

Essmann U, Perera L, Berkowitz ML, Darden T, Lee H, Pedersen LG. 1995. A smooth par-
ticle mesh Ewald method. J Chem Phys. 103(19):8577–8593.
https://doi.org/10.1063/1.470117

Feng J, Selvam B, Shukla D. 2021. How do antiporters exchange substrates across the cell
membrane? An atomic-level description of the complete exchange cycle in NarK. Struct
Lond Engl 1993. 29(8):922-933.e3. https://doi.org/10.1016/j.str.2021.03.014

Hohenberg P, Kohn W. 1964. Inhomogeneous Electron Gas. Phys Rev. 136(3B):B864–
B871. https://doi.org/10.1103/PhysRev.136.B864

Hoover WG. 1985. Canonical dynamics: Equilibrium phase-space distributions. Phys Rev
A. 31(3):1695–1697. https://doi.org/10.1103/PhysRevA.31.1695

Hu W, Chi C, Song K, Zheng H. 2023. The molecular mechanism of sialic acid transport
mediated by Sialin. Sci Adv. 9(3):eade8346. https://doi.org/10.1126/sciadv.ade8346

Huang J, Rauscher S, Nawrocki G, Ran T, Feig M, de Groot BL, Grubmüller H, MacKerell
AD. 2017. CHARMM36m: an improved force field for folded and intrinsically disordered pro-
teins. Nat Methods. 14(1):71–73. https://doi.org/10.1038/nmeth.4067

Husic BE, Pande VS. 2018. Markov State Models: From an Art to a Science. J Am Chem
Soc. 140(7):2386–2396. https://doi.org/10.1021/jacs.7b12191

Hutter J, Alavi A, Deutsch T, Bernasconi M, Goedecker S, Marx D, Tuckerman M, Parrinello
M. 2000. CPMD program. MPI Für Festkörperforschung IBM Zurich Res Lab.

Iharada M, Miyaji T, Fujimoto T, Hiasa M, Anzai N, Omote H, Moriyama Y. 2010. Type 1
sodium-dependent phosphate transporter (SLC17A1 Protein) is a Cl(-)-dependent urate ex-
porter. J Biol Chem. 285(34):26107–26113. https://doi.org/10.1074/jbc.M110.122721

Ishikawa T, Nishikawa H, Gao Y, Sawa Y, Shibata H, Yabuta Y, Maruta T, Shigeoka S. 2008.
The pathway via D-galacturonate/L-galactonate is significant for ascorbate biosynthesis in
Euglena gracilis: identification and functional characterization of aldonolactonase. J Biol
Chem. 283(45):31133–31141. https://doi.org/10.1074/jbc.M803930200

Kaback HR, Guan L. 2019. It takes two to tango: The dance of the permease. J Gen Physiol.
151(7):878–886. https://doi.org/10.1085/jgp.201912377

Klauda JB, Venable RM, Freites JA, O'Connor JW, Tobias DJ, Mondragon-Ramirez C, Vo-
robyov I, MacKerell AD, Pastor RW. 2010. Update of the CHARMM all-atom additive force
field for lipids: validation on six lipid types. J Phys Chem B. 114(23):7830–7843.
https://doi.org/10.1021/jp101759q

Kohn W, Sham LJ. 1965. Self-Consistent Equations Including Exchange and Correlation
Effects. Phys Rev. 140(4A):A1133–A1138. https://doi.org/10.1103/PhysRev.140.A1133

Kolen B, Borghans B, Kortzak D, Lugo V, Hannack C, Guzman RE, Ullah G, Fahlke C. 2023.
Vesicular glutamate transporters are H+-anion exchangers that operate at variable stoichi-
ometry. Nat Commun. 14(1):2723. https://doi.org/10.1038/s41467-023-38340-9

Kostritskii AY, Alleva C, Cönen S, Machtens J-P. 2021. g_elpot: A Tool for Quantifying Bio-
molecular Electrostatics from Molecular Dynamics Trajectories. J Chem Theory Comput.
17(5):3157–3167. https://doi.org/10.1021/acs.jctc.0c01246

Kostritskii AY, Machtens J-P. 2023. Domain- and state-specific shape of the electric field
tunes voltage sensing in voltage-gated sodium channels. Biophys J. 122(10):1807–1821.
https://doi.org/10.1016/j.bpj.2023.04.013

Kumar H, Finer-Moore JS, Kaback HR, Stroud RM. 2015. Structure of LacY with an α-sub-
stituted galactoside: Connecting the binding site to the protonation site. Proc Natl Acad Sci.
112(29):9004–9009. https://doi.org/10.1073/pnas.1509854112

Laio A, VandeVondele J, Rothlisberger U. 2002. A Hamiltonian electrostatic coupling
scheme for hybrid Car–Parrinello molecular dynamics simulations. J Chem Phys.
116(16):6941–6947. https://doi.org/10.1063/1.1462041

Leano JB, Batarni S, Eriksen J, Juge N, Pak JE, Kimura-Someya T, Robles-Colmenares Y,
Moriyama Y, Stroud RM, Edwards RH. 2019. Structures suggest a mechanism for energy
coupling by a family of organic anion transporters.Dutzler R, editor. PLOS Biol.
17(5):e3000260. https://doi.org/10.1371/journal.pbio.3000260

Li C, Voth GA. 2021. A quantitative paradigm for water-assisted proton transport through
proteins and other confined spaces. Proc Natl Acad Sci U S A. 118(49):e2113141118.
https://doi.org/10.1073/pnas.2113141118

Li F, Eriksen J, Finer-Moore J, Chang R, Nguyen P, Bowen A, Myasnikov A, Yu Z, Bulkley
D, Cheng Y, et al. 2020. Ion transport and regulation in a synaptic vesicle glutamate trans-
porter. Science. 368(6493):893–897. https://doi.org/10.1126/science.aba9202

Li F, Eriksen J, Finer-Moore J, Stroud RM, Edwards RH. 2022. Diversity of function and
mechanism in a family of organic anion transporters. Curr Opin Struct Biol. 75:102399.
https://doi.org/10.1016/j.sbi.2022.102399

von Lilienfeld OA, Tavernelli I, Rothlisberger U, Sebastiani D. 2005. Variational optimization
of effective atom centered potentials for molecular properties. J Chem Phys. 122(1):14113.
https://doi.org/10.1063/1.1829051

Lomize MA, Pogozheva ID, Joo H, Mosberg HI, Lomize AL. 2012. OPM database and PPM
web server: resources for positioning of proteins in membranes. Nucleic Acids Res. 40(Da-
tabase issue):D370-376. https://doi.org/10.1093/nar/gkr703

Madej MG, Sun L, Yan N, Kaback HR. 2014. Functional architecture of MFS D-glucose
transporters. Proc Natl Acad Sci U S A. 111(7):E719-727.
https://doi.org/10.1073/pnas.1400336111

Mangiatordi GF, Brémond E, Adamo C. 2012. DFT and Proton Transfer Reactions: A Bench-
mark Study on Structure and Kinetics. J Chem Theory Comput. 8(9):3082–3088.
https://doi.org/10.1021/ct300338y

Martyna GJ, Tuckerman ME. 1999. A reciprocal space based method for treating long range
interactions in ab initio and force-field-based calculations in clusters. J Chem Phys.
110(6):2810–2821. https://doi.org/10.1063/1.477923

Morin P, Sagné C, Gasnier B. 2004. Functional characterization of wild-type and mutant
human sialin. EMBO J. 23(23):4560–4570. https://doi.org/10.1038/sj.emboj.7600464

Nosé S. 1984. A unified formulation of the constant temperature molecular dynamics meth-
ods. J Chem Phys. 81(1):511–519. https://doi.org/10.1063/1.447334

Olsen JMH, Bolnykh V, Meloni S, Ippoliti E, Bircher MP, Carloni P, Rothlisberger U. 2019.
MiMiC: A Novel Framework for Multiscale Modeling in Computational Chemistry. J Chem
Theory Comput. 15(6):3810–3823. https://doi.org/10.1021/acs.jctc.9b00093

Omote H, Miyaji T, Hiasa M, Juge N, Moriyama Y. 2016. Structure, Function, and Drug In-
teractions of Neurotransmitter Transporters in the Postgenomic Era. Annu Rev Pharmacol
Toxicol. 56:385–402. https://doi.org/10.1146/annurev-pharmtox-010814-124816

Parrinello M, Rahman A. 1981. Polymorphic transitions in single crystals: A new molecular
dynamics method. J Appl Phys U S. 52(12):7182–7190. https://doi.org/10.1063/1.328693

Peng Y, Swanson JMJ, Kang S, Zhou R, Voth GA. 2015. Hydrated Excess Protons Can
Create Their Own Water Wires. J Phys Chem B. 119(29):9212–9218.
https://doi.org/10.1021/jp5095118

Preobraschenski J, Cheret C, Ganzella M, Zander JF, Richter K, Schenck S, Jahn R, Ahnert-
Hilger G. 2018. Dual and Direction-Selective Mechanisms of Phosphate Transport by the
Vesicular Glutamate Transporter. Cell Rep. 23(2):535–545.
https://doi.org/10.1016/j.celrep.2018.03.055

Quistgaard EM, Löw C, Guettou F, Nordlund P. 2016. Understanding transport by the major
facilitator superfamily (MFS): structures pave the way. Nat Rev Mol Cell Biol. 17(2):123–
132. https://doi.org/10.1038/nrm.2015.25

Qureshi AA, Suades A, Matsuoka R, Brock J, McComas SE, Nji E, Orellana L, Claesson M,
Delemotte L, Drew D. 2020. The molecular basis for sugar import in malaria parasites. Na-
ture. 578(7794):321–325. https://doi.org/10.1038/s41586-020-1963-z

Raghavan B, Schackert FK, Levy A, Johnson SK, Ippoliti E, Mandelli D, Olsen JMH, Roth-
lisberger U, Carloni P. 2023. MiMiCPy: An Efficient Toolkit for MiMiC-Based QM/MM Simu-
lations. J Chem Inf Model. 63(5):1406–1412. https://doi.org/10.1021/acs.jcim.2c01620

Reimer RJ. 2013. SLC17: A functionally diverse family of organic anion transporters. Mol
Aspects Med. 34(2–3):350–359. https://doi.org/10.1016/j.mam.2012.05.004

Scherer MK, Trendelkamp-Schroer B, Paul F, Pérez-Hernández G, Hoffmann M, Plattner N,
Wehmeyer C, Prinz J-H, Noé F. 2015. PyEMMA 2: A Software Package for Estimation, Val-
idation, and Analysis of Markov Models. J Chem Theory Comput. 11(11):5525–5542.
https://doi.org/10.1021/acs.jctc.5b00743

Schrödinger, LLC. 2015. The PyMOL Molecular Graphics System, Version 1.8.

Schulz P, Garcia-Celma JJ, Fendler K. 2008. SSM-based electrophysiology. Methods San
Diego Calif. 46(2):97–103. https://doi.org/10.1016/j.ymeth.2008.07.002

Smirnova I, Kasho V, Kaback HR. 2014. Real-time conformational changes in LacY. Proc
Natl Acad Sci U S A. 111(23):8440–8445. https://doi.org/10.1073/pnas.1408374111

Smirnova IN, Kasho VN, Kaback HR. 2006. Direct sugar binding to LacY measured by res-
onance energy transfer. Biochemistry. 45(51):15279–15287.
https://doi.org/10.1021/bi061632m

Soskine M, Adam Y, Schuldiner S. 2004. Direct evidence for substrate-induced proton re-
lease in detergent-solubilized EmrE, a multidrug transporter. J Biol Chem. 279(11):9951–
9955. https://doi.org/10.1074/jbc.M312853200

Swanson JM. 2022. Multiscale kinetic analysis of proteins. Curr Opin Struct Biol. 72:169–
175. https://doi.org/10.1016/j.sbi.2021.11.005

Thomas NE, Feng W, Henzler-Wildman KA. 2021. A solid-supported membrane electro-
physiology assay for efficient characterization of ion-coupled transport. J Biol Chem.
297(4):101220. https://doi.org/10.1016/j.jbc.2021.101220

Tiwary P, Parrinello M. 2015. A Time-Independent Free Energy Estimator for Metadynamics.
J Phys Chem B. 119(3):736–742. https://doi.org/10.1021/jp504920s

Tribello GA, Bonomi M, Branduardi D, Camilloni C, Bussi G. 2014. PLUMED 2: New feathers
for an old bird. Comput Phys Commun. 185(2):604–613.
https://doi.org/10.1016/j.cpc.2013.09.018

Troullier N, Martins JL. 1991. Efficient pseudopotentials for plane-wave calculations. Phys
Rev B Condens Matter. 43(3):1993–2006. https://doi.org/10.1103/physrevb.43.1993

Waterhouse A, Bertoni M, Bienert S, Studer G, Tauriello G, Gumienny R, Heer FT, de Beer
TAP, Rempfer C, Bordoli L, et al. 2018. SWISS-MODEL: homology modelling of protein
structures and complexes. Nucleic Acids Res. 46(W1):W296–W303.
https://doi.org/10.1093/nar/gky427

Wolf MG, Hoefling M, Aponte-Santamaría C, Grubmüller H, Groenhof G. 2010. g_membed:
Efficient insertion of a membrane protein into an equilibrated lipid bilayer with minimal per-
turbation. J Comput Chem. 31(11):2169–2174. https://doi.org/10.1002/jcc.21507

Zoete V, Cuendet MA, Grosdidier A, Michielin O. 2011. SwissParam: a fast force field gen-
eration tool for small organic molecules. J Comput Chem. 32(11):2359–2368.
https://doi.org/10.1002/jcc.21816

995
996

Figure legends

Figure 1 - Functional characterization of DgoT.

A pH dependence of WT DgoT peak currents measured by SSME upon application of 10 mM D-galactonate concentration jump. The inset shows transport currents at pH 8.0 from liposomes reconstituted with different lipid to protein ratios (LPR). Currents were normalized to their peak value for comparison of the current decay.

B Transient currents under symmetrical (black line) or asymmetrical conditions. Peak current decreases if acidic pH is applied inside liposomes (top) or alkaline pH is applied outside (bottom).

C Peak currents and determination of apparent pK values. The solid lines are fits to the data using the equations described in Methods. Experiments were performed in duplicate on two independent sensors.

D Representative transport current traces used for the determination of the transport stoichiometry. The inset shows time dependence of transported charge obtained by integration of current traces.

E Plots of transported charge *versus* ratio of the galactonate and proton chemical potentials.

F Changes in pH after addition of substrate to purified DgoT in detergent. Additions of 10 mM of galactonate are indicated by the black arrows. 5 μM HCl or NaOH were added to induce a pH shift comparable with the expected changes in H^+ concentration due to proton binding to DgoT.

Figure 2 - Unbiased MD simulations reveal protonation- and substrate- dependent dynamics of extracellular gate.

A-C Top row: snapshots showing the arrangement of the gating helices TM1 and TM7 in unbiased MD simulations with DgoT in outward-facing conformation with D46 and E133 deprotonated (A), protonated (B) and galactonate bound (C). The red colored parts of the helices correspond to residues 48-52 (TM1) and residues 271-275 (TM7). Bottom row: representative snapshots showing interactions around D46 and E133.

D, E Probability densities for extracellular gate opening in *apo* (D) and galactonate-bound (E) MD simulations with DgoT in the outward-facing conformation with different protonation states of D46 and E133. The TM1-TM7 distance is measured as distance between center of mass of Ca atoms of residues 48-52 and residues 271-275.

Figure 3 - Hydrophobic interactions at the level of the extracellular gate.

A, B Snapshots showing water molecules distribution in simulations with DgoT in outward-facing conformation with closed (A) and open (B) extracellular gate. Representative snapshots were taken from unbiased MD simulations with both D46 and E133 either deprotonated (A) or protonated (B).

C Hydration profile of the pore represented by the number of water molecules in 2 Å sections along the pore axis. Trajectories with following parameters were used for analysis: *apo*, closed gate – D46 and E133 protonated; *apo*, open gate and substrate bound – D46 and E133 protonated. In *apo* simulations only frames with minimum distance between side chains of F72 and W277 < 12.5 Å (blue line) or > 12.5 Å (green line) were used.

D Time course of extracellular gate opening (measured as TM1-TM7 distance, as in Fig. 2), number of water molecules in 10 Å section near extracellular gate (z coordinate between 58 and 68 Å) and galactonate binding to DgoT (measured as minimum distance between galactonate molecule and guanidinium group of R47) in a trajectory with both D46 and E133 protonated. Shaded lines represent raw data from the trajectory, solid lines are moving averages.

Figure 4 - A hydrophobic lock within the intracellular gate regulates access to the binding site.

A, B Snapshots showing water molecule distribution in simulations with DgoT in inward-facing conformation with closed (A) and open (B) intracellular gate. Representative snapshots were taken from unbiased MD simulations with D46 and E133 protonated (A) or only E133 protonated (B).

C Hydration profile of the pore, represented by the number of water molecules in 2-Å sections along the pore axis. Trajectories with following parameters were used for analysis: closed gate – D46 and E133 protonated; open gate – D46 deprotonated, E133 protonated. Substrate was bound to the protein in both cases. Only frames with minimum distance between side chains of F137 and W373 < 8 Å (blue line) or > 8 Å (green line) were used.

D Probability densities for intracellular gate opening (measured as distance between side chains of F137 and W373) in MD simulations with DgoT in inward-facing conformation with different protonation states of D46 and E133 and galactonate and various substrate occupancies.

Figure 5 - Proton release from D46.

A Water occupancy map in unbiased simulations with D46 and E133 protonated, contoured at an occupancy level of 0.1.

B Electrostatic potential at the carboxyl groups of protonated D46 and E133 in the unbiased MD simulation.

C Classical MTD simulations of the conformational change of protonated D46.

D Free energy profile for the proton transfer between D46 and E133, computed at the QM (BLYP)/MM level. The insets show representative starting, transition state, and final configurations. Error bars are omitted since they are smaller than marker size

1074

Figure 6 - Energetical description of major conformational changes.

A, B Free energy landscape for DgoT in substrate-bound (A) or *apo* (B) state. Protein con-
formations captured in DgoT crystal structures were projected onto the tICA space (black
points). Representative snapshots illustrate how the value of tIC2 correlates with the degree
of opening of the extracellular gate.

C *Left:* coarse representation of intermediate metastable states obtained with PCCS for sub-
strate-bound DgoT. *Right:* Stationaty probabilities of such metastables states.

D Same representation as in (C), for apo DgoT.

1083

Figure 7 - Neutralization of transmembrane charged residues abolish DgoT galacto-
nate transport.

A pH changes induced by H+-coupled galactonate, but not gluconate transport into *E.coli*
cells expressing WT DgoT or transformed with the same vector without the DgoT gene.

B Maximum pH changes observed in experiments with E.coli cells expressing different DgoT
variants.

C, D, F, G pH dependency of SSME currents initiated by galactonate concentration jump (10
mM) at symmetrical pH conditions. Experiments were performed on n=4 (C), n=5 (D), n=5
(F) and n=3 (G) independent sensors.

E Concentration-dependent changes in kobs, obtained by monoexponential fit of the current
decay, for WT and mutant DgoT. Solid lines are fits to a three-states induced-fit model given
above the graph. Experiments were performed on n=7 (WT), n=4 (E133Q), n=4 (R126Q)
and n=3 (D46N) independent sensors.

H Substrate concentration dependence of the peak current values for WT and R47Q DgoT.
R47Q concentration dependences were fitted to a two-state binding model given above the
graph. Values of peak current values for WT DgoT (fitted to three-states model from E) are
given for comparison. Experiments were performed on n=7 (WT) and n=4 (R47Q) independ-
ent sensors.

1102

Figure 8 - Galactonate does not induce extracellular gate closure in R47Q DgoT.

A, B Probability densities for extracellular gate opening in *apo* (A) and galactonate-bound
(B) simulations with WT and R47Q DgoT in outward-facing conformation with D46 and E133
protonated.

C, D Position of galactonate in the binding site of WT (C) and R47Q DgoT (D).

E Fraction of frames, in which distance between the side chain of specified residue and
closest oxygen atom of galactonate was equal or less than 2 Å. Each data point represents
one of the four independent trajectories for WT (blue points) and R47Q mutant (red points)
DgoT. Significance was evaluated with the Mann–Whitney test, one-sided: *$p < 0.05$.

1112

Figure 9 - Schematic transport cycle of DgoT.

1113

1114 In outward-facing conformation (1) extracellular gate opening is stabilized by protonation of
1115 D46 and E133 (2). Galactonate binding (3) induces major conformational changes (4). In
1116 the inward-facing state (5) proton release from D46 leads to galactonate release to the in-
1117 tracellular solution (6), followed by the release of the second proton (7) and reorientation of
1118 the empty carrier to outward-facing conformation (8).

1119

Tables and their legends

1120

1121 Table 1. Overview of I_{max} values (50 mM galactonate) and kinetic parameters for transport-

1122 deficient mutants, averaged over n = 3 different sensors (errors reflect standard deviations).

1123 For R47Q analysis of current decays was limited by time resolution of the instrument, there-

1124 fore only I_{max} values are given.

Variant	I_{max} (nA)	k_{obs} (s^{-1}) (50 mM galactonate)	K_d (mM)	k^- (s^{-1})	k^+ (s^{-1})
WT at pH 5.8	0.7 ± 0.2	79 ± 14	15.1	23.8	70.6
D46N	2.7 ± 0.3	68 ± 4	11.0	5.8	19.2
E133Q	1.0 ± 0.3	105 ± 15	15.3	30.8	92.2
R47Q	-0.37 ± 0.05				
R126Q	-1.5 ± 0.4	74 ± 11	22	14.4	86.7

1125

1126

A   Apo, closed gate

F274
A51
W277
L275
F72

Neck region

B   Apo, open gate

W277
A51
F72
L275
F274

TM1   TM7

C

Substrate binding site   Neck region   Extracellular solution

water molecules

56   64   72

Pore axis, Å

Apo, closed gate   Apo, open gate

Substrate bound

D

equil   no restraints

Distance TM1–TM7, Å

water molecules in neck region

20
16
12
8

60
40
20

Distance R47–gal., Å

30
20
10

0   25   50

Time, ns

A IF, closed gate

TM4 TM10

galactonate

F137

W373

B IF, open gate

TM4 TM10

galactonate

F137

Neck region

W373

C

Intracellular solution Neck region Substrate binding site

water molecules

Pore axis, Å

○ Closed gate   □ Open gate

D

Distance F137-W373, Å

D46[-]
E133[-]

D46[-]
E133[H]

D46[H]
E133[-]

D46[H]
E133[H]

D46[-]
E133[-]

D46[H]
E133[H]

D46[-]
E133[-]

D46[-]
E133[H]

D46[H]
E133[H]

■ bound
■ bound, protonated substrate
■ apo