The nanoscale organization of the Nipah virus fusion protein informs new membrane fusion mechanisms.

Qian Wang[1], Jinxin Liu[1], Yuhang Luo[1], Vicky Kliemke[1], Giuliana Leonarda Matta[1], Jingjing Wang[1], Qian Liu[1,2,*]

[1] Institute of Parasitology, Faculty of Agricultural and Environmental Sciences, McGill University, Saint Anne de Bellevue, Canada.

[2] McGill Center for Viral Diseases, Lady Davis Institute, Montreal, Canada.

Abstract.

Paramyxovirus membrane fusion requires an attachment protein for receptor binding and a fusion protein for membrane fusion triggering. Nipah virus (NiV) attachment protein (G) binds to ephrinB2 or -B3 receptors, and fusion protein (F) mediates membrane fusion. NiV-F is a class I fusion protein and is activated by endosomal cleavage. The crystal structure of a soluble GCN4-decorated NiV-F shows a hexamer-of-trimer assembly. Here, we used single-molecule localization microscopy to quantify the NiV-F distribution and organization on cell and virus-like-particle membranes at a nanometer resolution. We found that NiV-F on biological membranes forms distinctive clusters that are independent of endosomal cleavage or expression levels. The sequestration of NiV-F into dense clusters favors membrane fusion triggering. The nano-distribution and organization of NiV-F are susceptible to mutations at the hexamer-of-trimer interface, and the putative oligomerization motif on the transmembrane domain. We also show that NiV-F nanoclusters are maintained by NiV-F-AP-2 interactions and the clathrin coat assembly. We propose that the organization of NiV-F into nanoclusters facilitates membrane fusion triggering by a mixed population of NiV-F molecules with varied degrees of cleavage and opportunities for interacting with the NiV-G/receptor complex. These observations provide insights into the in-situ organization and activation mechanisms of the NiV fusion machinery.

Introduction.

The family of *Paramyxoviridae* contains many clinically important viruses, such as human parainfluenza virus 1-3, measles virus, and Nipah virus (NiV). Paramyxoviruses code for two glycoproteins on the viral membranes for virus-cell membrane fusion. The receptor-binding proteins (RBP, HN/H/G) engage the host receptors and activate the refolding of the fusion protein (F) that merges the virus and cell membranes[1]. Paramyxovirus-F is a class I fusion protein, along with human immunodeficiency virus (HIV) envelope, influenza virus hemagglutinin, and severe acute respiratory syndrome coronavirus-2 (SARS-CoV-2) spike[1]. Paramyxovirus-F is a trimeric transmembrane protein with a "tree-like" structure at its prefusion conformation. Upon triggering by the RBP/receptor complex, paramyxovirus-F mediates fusion by inserting its N-terminal fusion peptides into the target membrane and undergoes a conformational change from pre- to post-fusion conformations, similarly to other class I fusion proteins. The refolding of the paramyxovirus-F overcomes the energy barrier for membrane fusion and leads to virus-cell and cell-cell membrane fusion.

NiV is a well-studied paramyxovirus due to its capability of human-human and human-animal transmissions, high mortality and morbidity rates, and the unavailability of vaccines or therapeutics for human use[2,3]. NiV is closely related to Hendra virus, Cedar virus, and the newly identified Langya virus[4]. NiV-F is expressed on the cell surface as an fusion-inactive form,

activated in the endosome by cathepsin B and L cleavage, and exocytosed to the cell surface as
a fusion-active form[5,6]. NiV-G binds to ephrinB2 and/or -B3 receptors in host cells and triggers
the refolding of F that leads to virus-cell membrane fusion[7]. Three complementary models of the
NiV fusion activation mechanism have been proposed in the past. *In the first model*, the NiV-G
and F form a complex before receptor engagement. The F/G interaction maintains F at a
prefusion conformation. The receptor binding to G induces conformational changes in G,
leading to the release and refolding of F. This model is supported by co-immunoprecipitation
and conformational antibody binding assays[8,9]. *In the second model*, NiV-F exists as hexamer-
of-trimers, and the fusion peptides are sequestered at the hexameric interface. The activation of
one single trimer by the NiV-G/ephrinB2 complex can be transmitted to all six trimers via the
interfaces of the hexamer-of-trimers, leading to efficient F-triggering and fusion pore formation.
This model is supported by the crystal structure of a GCN4-decorated NiV-F ectodomain and
mutagenesis analysis of key amino acid residues at the hexameric interfaces[10]. *In a third model*,
NiV-F and G do not form a complex before receptor binding. The NiV-G/ephrinB2 engagement
clusters ephrinB2 and triggers F. F triggering leads to reduced mobility of a portion of F
molecules that potentially contributes to fusion pore formation. This model is supported by the
nanoscale distribution of NiV-F and G on cell membranes resolved by super-resolution
microscopy and a single-particle tracking assay of NiV fusion protein at the interface between
supported lipid bilayer and live cell membranes[11,12]. Although these models explain the NiV
membrane fusion mechanisms from different perspectives, a link between the molecular
structure and the fusion machinery on biological membranes remains unclear.

We attempted to address this link by analyzing the nano-organization of NiV-F at the prefusion
state on biological membranes using single-molecule localization microscopy (SMLM) and
mutagenesis analyses. We have shown that NiV-F forms clusters on biological membranes that
are isolated from NiV-G[11]. Here, we show that NiV-F forms regular-sized nanoclusters on cell
membranes regardless of the expression level or endosomal cleavage. The estimated size of
NiV-F clusters on cell membrane is similar to that of the hexamer-of-trimer assemblies formed
by the GCN4-decorated soluble NiV-F. The NiV-F nano-organization is altered by mutations at
the hexameric interface and the putative oligomerization motifs at the transmembrane domain.
NiV-F molecules enriched in nanoclusters favor membrane fusion activation. The NiV-F
nanoclusters are stabilized by the interactions between NiV-F, the endocytosis adaptor complex
AP-2, and the clathrin coat at the cell membranes. In summary, our study reveals the NiV-F
nano-organization on the biological membranes and provides novel insights to the NiV fusion
machinery *in situ*.

NiV-F forms regular-sized clusters that are not affected by the surface expression level.

We recently published that NiV-F formed distinctive nanoclusters on the plasma membrane
regardless of the presence of NiV-G or ephrinB2[11]. To investigate the determinants of the NiV-F
nano-organization, we used SMLM to probe the NiV-F at various conditions. SMLM generates
super-resolution images with nanometer resolution by localizing blinks from individual
fluorophores with a distance greater than the diffraction limit of light[13]. PK13 cells have a flat
morphology that is suitable for SMLM imaging. PK13 cells express little endogenous ephrinB2
and -B3 receptors for NiV and thus can eliminate any potential receptor-dependent clustering.
Cell surface NiV-F was detected using a FLAG tag in its extracellular domain[11]. The FLAG tags
were employed to detect NiV-F because of the availability of high affinity and specificity anti-
FLAG antibodies for a high labeling density, which is vital for the reconstruction accuracy of

SMLM images[14]. To rule out the effect of the FLAG tag on NiV-F clustering, we compared NiV-F
clusters formed by NiV-F-FLAG to that of a NiV-F-HA construct. Both HA and FLAG tags were
inserted after amino acid residue 104, right before the fusion peptide (Fig. S1A). Visual
examination of the SMLM images reveals that NiV-F-FLAG and NiV-F-HA form similar clusters
(Fig. S1B). The clustering tendency, indicated by Hopkins' index, is comparable between NiV-F-
HA and NiV-F-FLAG (Fig. S1C), suggesting that NiV-F clustering is not affected by specific
epitope tags.

Next, we investigated whether the nano-organization of NiV-F depends on the cell surface
expression levels. The total cell surface expression level (CSE) of NiV-F in a selected cell was
measured before the cell was subjected to SMLM imaging. The SMLM images show that NiV-F
forms similar nanoclusters in both high-expression and low-expression cells (Fig. 1A and B).
Our data show that the clustering tendency of NiV-F is independent of its CSE in individual cells
(Fig. 1C). To gain a quantitative understanding of NiV-F nanoscale organization, we used a
cluster identification method DBSCAN (Density-Based Spatial Clustering of Applications with
Noise) that links the closely situated localizations in a propagative manner. We created cluster
and density maps using Clus-DoC for representative high-expression and low-expression cells
(Fig. 1D and E)[15]. Our data show that the estimated diameter of the NiV-F clusters ranges from
20-34 nm, with a peak at 24 nm (Fig. 1F). Nonetheless, the precise values of the cluster
diameter are indicative rather than definite in SMLM as it could be affected by the labeling
complex and the blinking property of the fluorophore. The distribution of the cluster diameter
shows that ~60% of NiV-F clusters have similar sizes, indicating that NiV-F clusters maintain a
quite uniform appearance. Interestingly, we noticed that the size of the clusters (Fig. 1G) and
localization density (Fig. 1H) in clusters were similar between high- and low-expression cells,
suggesting that the NiV-F organization in nanoclusters is not affected by the expression levels.
Moreover, our data show that the high-expression cells have more clusters than the low-
expression cells (Fig. 1I). These results suggest that changes in CSE may affect the number of
NiV-F clusters on the plasma membrane (Fig. 1I), but not the clustering tendency of NiV-F (Fig.
1C), the size of the clusters (Fig. 1G), or the localization density within clusters (Fig. 1H). These
data imply that the distribution and nano-organization of NiV-F are tightly regulated and may be
important for membrane fusion activation. We also analyzed NiV-F distribution in different cell
lines. HeLa cells express ephrinB2 and -B3, and thus are NiV permissive cells. The overall NiV-
F nano-organization seems quite similar on PK13 and HeLa cells, agreeing with our previous
results (Fig. S2A)[11]. The Hopkins' indices are also similar for NiV-F on PK13 and HeLa cells
(Fig. S2B). The DBSCAN analysis shows that the percentage localizations in clusters (Fig.
S2C), the relative density defined as the ratio of the localization density in clusters to the total
localization density in the selected region (Fig. S2D), and cluster diameters (Fig. S2E) are
comparable for NiV-F on PK13 and HeLa cells. These results suggest that NiV-F clusters are
minimally affected by the cell lines of expression or the presence of the ephrinB2 and/or -B3.

Fig. 1. NiV-F forms regular-sized clusters that are not affected by the surface expression level. *(A and B) Cross-section (Δz = 600 nm) of SMLM images of NiV-F in high-expression (A) and low-expression (B) PK13 cells. Scale bar: 1 µm. The yellow boxed region is enlarged to show the detailed distribution pattern. Scale bar: 0.2 µm. (C) Hopkin's index of the F localizations in low- and high-expression cells. n = 35 and 40. (D and E) Cluster maps (left) and localization density maps (right) of the enlarged regions in A and B. Cluster contours are highlighted with gray lines. Normalized relative density is pseudocolored according to the scale on the right. (F) The percentage distribution of the NiV-F cluster diameter from 13 cells. n = 133. (G) The cluster diameters in low- and high-expression cells. n = 58 and 56. (H) The localization density (# of localizations per µm²) within clusters in low- and high-expression cells. n = 58 and 50. (I) The number of clusters per µm² in low- and high-expression cells. n = 59 and 74. The cut-off fluorescence intensity for low- and high-expression cells is 8000 (Arb. Unit). Sample size n is the number of total regions from 6-8 cells. Bars represent mean ± SD. P value was obtained using the Mann-Whitney test. ns: p > 0.05; * p<0.01; ** p<0.001; *** p<0.0001.*

The NiV-F nanoclusters are not affected by NiV-F cleavage.

NiV-F is synthesized in host cells as inactive precursor F_0 and cleaved by cellular proteases
cathepsin L and cathepsin B in the endosomes at an acidic pH to generate the fusion-active,
disulfide-linked F_1-F_2 construct. The F_1-F_2 is subsequently recycled to the cell surface to induce
cell-cell fusion and incorporated into virus particles. We hypothesized that the F_0 precursor and
the F_1-F_2 active forms co-existed in the same clusters on the cell surface. To test this
hypothesis, we used a pan-cysteine cathepsin inhibitor E64d to inhibit the NiV-F cleavage. E64d
inhibited the cell-cell fusion induced by NiV-F and NiV-G in HeLa cells (Fig. S3), agreeing with
the previous fusion inhibitory effects in Vero and MDCK cells[5,6]. Our SMLM images show that
E64d treatment does not result in any significant change in the clustering of NiV-F in HeLa cells
(Fig. 2A). The clustering extent of NiV-F was not altered by the E64d treatment, as shown by
comparable Hopkin's indices (Fig. 2B). Additionally, quantitative parameters of the nano-
organization such as percentage of localizations in clusters (Fig. 2C), relative density (Fig. 2D),
cluster size (Fig. 2E), and the total density of the regions (Fig. 2F). Our data show that clusters
formed by non-cleaved NiV-F in cells treated by E64d are similar to those formed by a mixture
of cleaved and non-cleaved NiV-F in the control cells, suggesting that F cleavage does not alter
F clustering on the cell membrane. These data support our hypothesis that the non-cleaved
precursor F_0 and cleaved F_1-F_2 coexist in the same cluster on the cell membrane.

Fig 2. Endosomal cleavage does not affect the nanoscale distribution of NiV-F. *(A) First*
column: Cross-section (Δz = 600 nm) of SMLM images of NiV-F in HeLa cells untreated (NC)

 and treated with 20 µM E64d (E64d). HeLa cells were co-transfected by expression plasmids
coding for NiV-G and NiV-F. Twenty µM E64d or the same volume of solvent methanol was
added to cells at 2 hrs post-transfection. Scale bar: 1 µm. Second column: The yellow boxed
region in the first column is enlarged to show individual clusters. Scale bar: 0.2 µm. Third
column: Cluster maps from the enlarged regions. Fourth column: Localization density maps
show the normalized relative density of the enlarged regions. (B-F) Quantitative analyses of
NiV-F clusters formed in HeLa cells without (NC) and with (E64d) the E64d treatment: (B)
Hopkin's index, n = 40 and 40; (C) percentage of localizations in clusters, n = 101 and 101; (D)
relative density, n = 100 and 103; (E) average cluster diameters, n = 100 and 101; (F) total
density of the region, n = 101 and 102. Bars represent mean ± SD. p value was obtained using
*Mann-Whitney test. ns: p > 0.05; * p<0.01; ** p<0.001; *** p<0.0001. Sample size n is the*
number of total regions from 4-10 cells.

Mutations destabilizing the NiV-F hexamer-of-trimer assembly alter its nano-organization on plasma membrane.

We noticed that the uniformity in the NiV-F cluster morphology resembled the hexameric assembly of soluble NiV-F decorated by GCN4[10]. Next, we investigated whether NiV-F clusters were stabilized by the hexameric interface. As reported, L53D and V108D mutants destabilize the hexameric interface and demonstrate decreased membrane fusion ability; Q393L stabilizes the hexameric interface and demonstrates increased fusion ability. We inserted the FLAG tag into the ectodomains of these mutants at the same position as that of the NiV-F. We verified that the relative fusion ability to their F-WT counterpart was comparable between the FLAG-tagged and untagged mutants (Fig. S4A and S4B)[10]. The FLAG-tagged mutants showed similar CSE levels to NiV-F (Fig. S4C). All mutants showed some levels of processing, although the cleavage of L53D was the least efficient comparing to F-WT and other mutants (Fig. S4D).

NiV-F mutants L53D, V108D, and Q393L were expressed on PK13 cells and subjected to SMLM imaging. The images show that clusters formed by L53D and V108D are smaller and more dispersed than those of the F-WT and Q393L (Fig. 3A). The Hopkin's index for Q393L is significantly higher than that of F-WT, while the Hopkin's indices for L53D and V108D are slightly lower than that of F-WT (Fig. 3B). Consistently, a lower percentage of L53D and V108D localizations are segregated into clusters than that of F-WT, while a similar percentage of Q393L and F-WT localizations form clusters (Fig. 3C). Our analysis also shows that L53D and V108D form smaller clusters compared to the F-WT. The Q393L clusters are of a similar size to that of F-WT (Fig. 3D). In addition, clusters of L53D and V108D are less packed than that of F-WT, with 48-, 49-, and 55-fold denser than the total localization density in the region, respectively (Fig. 3E). The clusters of Q393L are not significantly denser than that of F-WT (Fig. 3E). The total density of localizations and the total CSE levels for all constructs are comparable (Fig. 3F and Fig. S4C), suggesting that differences in clustering are not due to variable numbers of localizations or levels of expression. Collectively, these data show that the mutations destabilizing the hexameric interface (L53D and V108D) make NiV-F localizations more dispersed and form smaller and less packed clusters. Q393L that stabilizes the hexameric interface promotes the clustering of NiV-F localizations (Fig. 3B), but does not significantly alter the organization and morphology of individual clusters (Fig. 3C-F). These data suggest that the NiV-F clusters are susceptible to mutations at the hexameric interface. Considering that the uniformed cluster size is similar to the assemblies formed by the soluble GCN4-NiV-F, we propose that NiV-F may organize into hexamer-of-trimer on the plasma membrane.

210
Fig. 3. Mutations at the NiV-F hexameric interface affect its nano-organization. *(A) First
column: Cross-section (Δz = 600 nm) of SMLM images of the FLAG-tagged NiV-F-WT (WT),
L53D, V108D, and Q393L on PK13 cell membrane. Scale bar: 1 μm. Second column: The
yellow boxed regions in the first column are enlarged to show individual clusters. Scale bar: 0.2
μm. Third row: Cluster maps from enlarged regions. Fourth column: Localization density maps
show normalized relative density of the enlarged regions. (B-F) Quantitative analyses of the
distribution of the FLAG-tagged NiV-F constructs: (B) Hopkin's index, n = 57-70; (C) Percentage
of localizations in clusters, n = 106-198; (D) Average cluster diameters, n = 106-196; (E)*

Relative density, n = 90-187; (F) total density of the region (a ratio of total localizations in a
region to the size of the region), n = 105-242. Bars represent mean ± SD. Sample size n is the
number of total regions from 11-20 cells. p value was obtained using Mann-Whitney test. ns: p >
0.05; * p<0.01; ** p<0.001; *** p<0.0001.

Mutations at the NiV-F hexameric interface affect its distribution on the VLP membrane.
To further investigate the organization of F mutants in viral membranes, we imaged the FLAG-
tagged NiV-F constructs on VLPs produced by 293T cells expressing NiV-M-GFP, NiV-G, and
NiV-F-WT, L53D, V108D, or Q393L mutants. The western blot analysis of NiV-F constructs on
the VLPs (Fig. 4A). The L53D and V108D on VLPs are less cleaved compared to that of the F-
WT and Q393L, as suggested by the weaker NiV-F_2 bands (Fig. 4A). The incorporation of NiV-F
is comparable among the NiV-F constructs (Fig. 4A). Notably, the incorporation of NiV-G is
largely abrogated in the VLPs that expressing L53D (Fig. 4A).

To probe the organization of NiV-F constructs on VLPs, the F-constructs on VLP membranes
were stained using Alexa Fluor 647 and subjected to SMLM imaging. The GFP fluorescence on
NiV-M was used to locate the VLPs. Fig. 4B shows that F-WT and F-Q393L form distinctive
clusters, while localizations of L53D and V108D are more dispersed on a z = 100 nm projection.
This agrees with their distributions on the plasma membrane (Fig. 3A). To gain a quantitative
insight into the distribution of the F constructs on the three-dimensional (3D) VLPs, we used an
algorithm named OPTICS (Ordering Points To Identify Clustering Structure)[16]. To identify
clusters, the OPTICS algorithm sorts out the localizations by calculating the reachability
distance between two adjacent localizations in a propagative manner and generates a
sequence of the ordered localizations[16]. In the plot of reachability distance vs. the sequence of
ordered localizations, a sudden increase in the reachability distance marks the cutoff of a cluster
(Fig. S5A). The shade between neighboring peaks represents localizations that are classified in
one cluster (Fig. S5A). OPTICS can identify sub-clusters on a 3D VLP that may not be
recognized by DBSCAN that uses one fixed parameter for the entire dataset. Fig. S5B shows
the OPTICS plots of the localizations of F-WT and mutants on representative VLPs (Fig. 4B).
The noticeable kinks suggest that F-WT and Q393L form distinctive clusters on VLPs, while the
irregular, small kinks indicate that L53D and V108D are more dispersed and form smaller
clusters on VLPs (Fig. 4B and Fig. S5B). To gain a population insight on the nano-organization
of the F constructs on VLPs, we developed a one-dimensional convolutional neural network (1D
CNN) to classify the ordered sequence of reachability distances of the NiV-F localizations
obtained by the OPTICS algorithm[17]. The distribution patterns of the localizations of the F-WT
and Q393L partition into one category, and L53D and V108D in another (Fig. 4C). These results
indicate that L53D and V108D form smaller and more dispersed clusters than F-WT and Q393L
on the VLP membranes, agreeing with that of the plasma membrane (Fig. 3). Our results
indicate that the trimer-trimer interaction at the hexameric interface is key in stabilizing the nano-
organization of NiV-F on both cell and viral membranes, and NiV-F do not seem to rearrange
during the incorporation into VLPs. Nonetheless, our data do not rule out the possibility that the
association of nucleocapsids with the plasma membrane during assembly may reorganize NiV-F
on the authentic virus membranes.

Fig. 4. The distribution and organization of NiV-F constructs in VLPs. *(A). The incorporation of F-WT and mutants in VLPs. NiV-M-GFP, G-HA, and FLAG-tagged F-WT or mutants were transfected to 293T cells. The supernatants were collected at 48 hrs post-transfection and analyzed on SDS-PAGE followed by western blotting. NiV-M was probed by polyclonal goat anti-GFP, NiV-G polyclonal rabbit anti-HA, F_0 and F_2 M2 monoclonal mouse anti-FLAG antibody. (B) Cross-section (Δz = 100 nm) of SMLM images of the FLAG-tagged NiV-F-WT (WT), L53D, V108D, and Q393L on individual VLPs. Scale bar: 0.2 µm.* (C) The classification of the ordered sequence of reachability distances of the NiV-F constructs localizations. Orange: F-WT and Q393L; Blue: L53D and V108D.

The NiV-F clusters are dispersed upon the disruption of the interactions between transmembrane domains (TMD). Evidence shows that TMD of class I fusion protein can self-associate in the absence of the rest of the protein and is important for membrane fusion[18]. Previous studies show that mutations in the Leucine-Isoleucine Zipper (LI zipper) motif lead to the dissociation of an HeV-F TMD-derived peptide, decreased stability, and the fusion ability of the whole HeV-F protein[19]. As the TMD domains of the NiV- and HeV-F demonstrate a 94% similarity, we mutated leucine (488 and 509) and isoleucine (495 and 502) residues of the LI zipper in NiV-F TMD to alanine (LI4A) (Fig. S6A). Similar to their counterparts of HeV-F, the NiV-F-LI4A showed a decreased F processing (Fig. S6B), cell-cell fusion activity (Fig. S6C and S6D), and CSE levels (Fig. S6E)[19]. Visual inspection of the images suggests that LI4A forms bigger clusters than F-WT (Fig. 5A). However, the LI4A seems less likely to cluster than F-WT, indicated by a lower Hopkin's index (Fig. 5B). Similarly, the portion of LI4A localizations partitioned into clusters is slightly lower than that of WT (Fig. 5C). Interestingly, we noticed that clusters formed by LI4A are significantly bigger (Fig. 5D) than that of F-WT and with a similar localization density within the clusters (Fig. 5E). The data suggest that interactions between TMDs of NiV-F monomers play a role in stabilizing its nano-organization. Dissociation of TMDs by mutations may result in increased space between F monomers and thus leads to bigger clusters on the cell membrane.

Fig. 5. Mutations in the LI zipper of the NiV-F transmembrane domain disturbs the NiV-F distribution. A) First column: Cross-section (Δz = 600 nm) of SMLM images of the FLAG-tagged NiV-F-WT (WT) and NiV-F-LI4A (LI4A) mutant on PK13 cell membrane. Scale bar: 1 μm. Second column: The yellow boxed region in the first column is enlarged to show individual clusters. Scale bar: 0.2 μm. Third column: Cluster maps from enlarged regions. Fourth column: Localization density maps show normalized relative density of the enlarged regions in the second row. (B-F) Quantitative analyses of the WT and LI4A clusters: (B) Hopkin's index, n=40 and 40 (C) Percentage of localizations in clusters, n = 171 and 211; (D) Average cluster diameters, n = 171 and 211; (E) Relative density, n = 166 and 211. Bars represent mean ± SD. p value was obtained using Mann-Whitney test. ns: p > 0.05; * p<0.01; ** p<0.001; *** p<0.0001. Sample size n is the number of total regions from 13-16 cells.

The NiV-F nanoclusters are stabilized by endocytosis components.

NiV-F can be endocytosed for endosomal cleavage. Next, we investigated whether NiV-F clusters are stabilized during endocytosis. NiV-F contains an endocytosis sorting signal YSRL and an additional YY motif at its cytoplasmic tail (Fig. S7A). Mutation of these tyrosine residues to alanine almost diminished NiV-F cleavage (Diederich et al., 2005). A FLAG-tagged NiV-F-YA construct containing the aforementioned mutations resulted in significantly less F_2 band than F-WT (Fig. S7B) and completely abrogated cell-cell fusion in 293T cells (Fig. S7C and S7D),

although the CSE levels were comparable to that of F-WT (Fig. S7E). By visually inspecting the
SMLM images, we noticed that F-YA was less clustered than that of F-WT (Fig. 6A). Indeed, the
Hopkin's analysis suggests that F-YA is less clustered than F-WT (Fig. 6B). The DBSCAN
analysis shows that a lower percentage of F-YA localizations segregate into clusters (Fig. 6C)
than that of the F-WT, and clusters formed by F-YA are smaller (Fig. 6D) and less dense (Fig.
6E) than that of the F-WT. Notably, there is no significant difference in the total density for F-YA
and F-WT SMLM images (Fig. 6F), indicating that the differences in cluster organizations do not
result from overall protein expression or the stochastic blinking properties of the fluorophore.
These results suggest that the endocytosis sorting signal of NiV-F may facilitate the enrichment
of NiV-F into clusters on the cell surface, preparing them for endosomal cleavage and/or
incorporation into viruses.

It is established that the YxxΦ sorting signal on the endocytosis cargo is bound and enriched by
the heterotetrameric adaptor protein (AP) complexes, which further recruit the assembly of the
clathrin coat[20,21]. We hypothesize that the clustering of NiV-F is promoted by the assembly of the
clathrin coat via the NiV-F-AP-2 interactions. As suggested by a yeast two-hybrid analysis, NiV-
F interacts with μ1, μ2, μ3 and μ4 subunits of the AP complexes AP-1, AP-2, AP-3, and AP-4,
and mutations of Y525, Y542, and Y543 almost completely abolished the interaction between
AP-2 and NiV-F[22]. Indeed, the co-immunoprecipitation assay shows that less AP2μ2 is pulled
down by F-YA than that of F-WT, suggesting that F-YA has a reduced interaction with AP-2 (Fig.
S7F). Next, we examined the NiV-F clusters in HeLa cells treated by the endocytosis inhibitor
pitstop2. Pitstop2 blocks clathrin-mediated endocytosis by obstructing the binding of the
accessory protein (*e.g.* AP-2) and the clathrin terminal domain, and potentially preventing the
assembly of the clathrin coat[23]. The SMLM images (Fig. 6G) show that the NiV-F localizations
are more dispersed upon pitstop2 treatment, agreeing with the lower Hopkin's index (Fig. 6H)
and a smaller percentage of localizations in clusters (Fig. 6I) in pitstop2 treated group compared
to that of the control group. Interestingly, we noticed that the NiV-F clusters in pitstop2 treated
cells were larger than the control cells (Fig. 6K), although with similar densities (Fig. 6J). The
total density is consistent between the pitstop2 treated and control groups (Fig. 6L). The overall
dispersed NiV-F distribution upon pitstop2 treamtnet may potentially be caused by the disrupted
assembly of clathrin coat. In combination, these results suggest that the enrichment of NiV-F by
AP-2 and the subsequent assembly of the clathrin-coated endosomes are important in
stabilizing the F clusters on cell membranes. We envision that the endocytosis components may
also facilitate the NiV-F clustering in virus particles as previous proteomic studies show the
presence of endocytosis related proteins, such as clathrin, AP-2, and dynamin in NiV VLPs[24–26].

Fig. 6. The NiV-F nanoclusters are stabilized by endocytosis components. *(A) First*
column: Cross-section (Δz = 600 nm) of SMLM images of the FLAG-tagged NiV-F-WT (WT) and
NiV-F-YA (YA) mutant on PK13 cell membrane. Scale bar: 1 µm. Second column: The yellow
boxed region in the first column is enlarged to show individual clusters. Scale bar: 0.2 µm. Third
column: Cluster maps from enlarged regions. Fourth column: Localization density maps show
normalized relative density of the enlarged regions in the second row. (B-F) Quantitative
analyses of the WT and YA clusters: (B) Hopkins index of WT and YA, n = 40 and n = 40; (C)
Percentage of localizations in clusters, n = 72 and 77; (D) Average cluster diameters, n = 72 and
77; (E) Relative density, n = 71 and 76; (F) total density of the region, n = 72 and 77. (G) First
column: Cross-section (Δz = 600 nm) of SMLM images of the FLAG-tagged NiV-F-WT treated
without (NC) and with pitstop2 (pitstop2) on HeLa cell membrane. Scale bar: 1 µm. Second
column: The yellow boxed region in the first column is enlarged to show individual clusters.
Scale bar: 0.2 µm. Third column: Cluster maps from enlarged regions. Fourth column:
Localization density maps show normalized relative density of the enlarged regions in the
second row. (H-L) Quantitative analyses of NiV-F without (NC) and with pitstop2 (pitstop2): (H)
Hopkins index, n = 40 and n = 40; (I) Percentage of localizations in clusters, n = 82 and 85; (J)
Relative density, n = 82 and 81; (K) Average cluster diameters, n = 82 and 81; (L) total density
of the region, n = 82 and 81. Bars represent mean ± SD. p value was obtained using Mann-
*Whitney test. ns: p > 0.05; * p<0.01; ** p<0.001; *** p<0.0001. Sample size n is the number of*
total regions from 4-9 cells.

Discussion.

Here we used SMLM to resolve the arrangement and organization of NiV-F on the biological
membranes at a resolution of 10 nm. Our data supports the following conclusions: 1) NiV-F is
organized into nanoclusters on the biological membranes and this organization is independent
of the protein expression level or endosomal cleavage; 2) the NiV-F nano-organization is

susceptible to mutations at the trimer interface on the NiV-F ectodomain and the putative
oligomerization motif on the transmembrane domain; 3) NiV-F sequestered in nanocluaters
favors membrane fusion activation; 4) the interactions among NiV-F, the AP-2 complex, and the
clathrin coat assembly stabilize the NiV-F clusters. We propose that the NiV-F nanoclusters are
the fundamental unit of the NiV fusion machinery, and this organization facilitates membrane
fusion triggering by a mixed population of NiV-F molecules with varied degrees of cleavage and
opportunities of interacting with NiV-G/receptor complex.

The nano-organization of NiV-F informs the coordinated membrane fusion triggering by a mixed
population of NiV-F. This notion is supported by the following observations from this and
previous studies: 1) NiV-F molecules form distinctive, regular-sized nanoclusters on the cell and
virus membranes (Fig. 1A-E); 2) the cleaved, active F co-exist with the noncleaved, inactive F in
a nanocluster, as suggested by that the NiV-F nanoclusters are resistant to the cathepsin
inhibitor, E64d (Fig. 2); 3) the NiV-F and G are segregated into different clusters [11]; and 4) the
full-length NiV-F and -G proteins do not show stable interactions either before or after ephrinB2
activation [12]. These observations indicate that the NiV-F triggering may be a result of transient
interactions between the F molecules at the edge of the clusters with the adjacent ephrinB2/G
complex. Moreover, we observed that the NiV-F sequestering into clusters is favorable for
membrane fusion activation (Fig. 3 and 5), highlighting the importance of the spatial
arrangement of NiV-F in its fusion activity. Therefore, it is likely that the triggered, fusion-active
NiV-F at the cluster peripheral can facilitate other F molecules in the same cluster to fulfill the
conformational changes, and thus maximize the energy to merge the opposing membranes.

The nano-organization of NiV-F reveals the fundamental unit of NiV fusion machinery. Previous
study revealed a hexamer-of-trimer assembly of soluble, GCN4-decorated, prefusion NiV-F.
Using SMLM, we estimated that the NiV-F clusters had a diameter between 24-26 nm (Fig. 1F).
This estimation agrees with the size of the hexameric soluble NiV-F-GCN4 assemblies[10].
Notably, the density of clusters (Fig. 1I), but not the size and localization density of individual
clusters (Fig. 1G and H), is affected by the surface expression levels of NiV-F, implying that the
NiV-F clusters are highly regulated and a fundamental functional unit. Nonetheless, a
collaborative effort of multiple copies of viral envelope proteins is favorable for overcoming the
energy barrier required for membrane fusion. Nanoclusters formed by viral glycoproteins have
been observed for HIV-1[27,28], Herpes virus[29], and Influenza virus[30]. The viral restriction factor,
serine incorporator protein 5 (SERINC5), inhibits HIV-1 fusion by disrupting the HIV env
nanoclusters, highlighting the importance of the nanoclusters in the function of the viral
glycoproteins[31]. More interestingly, the NiV-F distribution becomes more dispersed,
characterized by smaller and less dense clusters, upon the mutation of two key residues, L53
and V108, at the hexameric interface, suggesting that the NiV-F clusters are likely to be the
hexamer-of-trimer assemblies on the cell and viral membranes (Fig. 3 and 4).

We also identified that the endosomal components as key host factors in maintaining the NiV-F
nano-organization on the cell membrane. Our data suggest that both the disruption of the F/AP-
2 interaction and the inhibition of clathrin assembly by pitstop2 result in dispersed NiV-F
distribution. AP-2 is the key adaptor of clathrin-mediated endocytosis. It binds cargo and
PtdIns(4,5)P_2 -containing membrane via multiple interface, and undergoes conformational
changes to append to the clathrin lattice[23,32]. Our data indicate that AP-2-NiV-F interaction can
enrich and strengthen the F nanoclusters (Fig. 6A-F and Fig. S7F), and potentially facilitate the
uptake of NiV-F by endocytosis. It is reported that the clustering of CD44, modulated by *N-*

glycosylations, facilitates the uptake of CD44 by endocytosis. Furthermore, the assembly of the
clathrin coat at the plasma membrane may also strengthen the F clusters, because pitstop2 that
inhibits the clathrin-coat assembly by blocking the clathrin terminal domain[23] prevents the
cluster formation of NiV-F. It is plausible that both clathrin and AP-2 facilitate NiV-F clustering on
cell and virus membranes because of the similar organization of F on virus and cell membranes
and the presence of clathrin and AP-2 in VLPs[24,26].

In conclusion, our observations provide direct evidence on the nano-organization and
distribution of NiV-F on the cell and viral membranes and shed lights to the fusion activation
mechanisms. It would be interesting to analyze the reorganization of the NiV-F clusters upon
triggering by the NiV-G/receptor complex on live cell membranes by combining single-molecule
imaging and the supported lipid bilayer technologies[12]. Due to technical challenges, we cannot
obtain an exact number of NiV-F molecules in NiV-F clusters. It would be interesting to pinpoint
the numbers of NiV-F required for membrane fusion activation on cell and virus surface, and this
could potentially facilitate the vaccine design. Additionally, elucidating host factors that maintain
NiV-F clusters and the interacting motif in NiV-F may represent a new therapeutic strategy for
NiV infections.

Materials and Methods.

Reagent type	Designation	Source or reference	Identifiers	Additional information
Cell line	HEK293T	ATCC	CRL-3216	
Cell line	PK13	ATCC	CRL-6489	
Cell line	HeLa cells	ATCC	CCL-2	
DNA plasmid	NiV-F-FLAG and mutants	This paper		The FLAG tag was inserted after residue 104 of the codon-optimized NiV-F (genebank accession no. AY816748.1). The hexameric interface mutants (Fig. 3)[10], and LI4A mutant (Fig. 5)[19], and YA mutant (Fig. 6)[6], were constructed as previously published.
DNA plasmid	NiV-F-HA	This paper		The HA tag was inserted after residue 104 of the codon-optimized NiV-F (genebank accession no. AY816748.1).
DNA plasmid	NiV-G-HA	Liu et al. 2018[11]		

DNA plasmid	NiV-M-GFP	Liu et al. 2018, Wang et al. 2010 [11,33]		
DNA plasmid	AP2μ2-mCherry	Christien Merrifield	Addgene 27672	
Chemical compounds, drug	E64d	Sigma-Aldrich	E8640-250UG	20uM
Chemical compounds, drug	Pitstop2	Sigma-Aldrich	SML1169-5MG	30uM
Antibody	Anti-FLAG mouse monoclonal	Sigma-Aldrich	F1804	IF, SMLM: 1:100, Flow: 1:200; WB: 1:500-1:5,000
Antibody	Anti-HA rabbit polyclonal antibody	Biolegend	902301	IF: 1:900 WB: 1:2,000
Antibody	Anti-GFP goat polyclonal antibody	Abcam	Ab5450	WB: 1:1000
Antibody	Anti-mCherry rabbit polyclonal antibody	Abcam	Ab167453	WB: 1:2000
Antibody	Anti-mouse donkey polyclonal antibody, Alexa Fluor® 647 conjugated	Invitrogen	A31571	IF and SMLM: 1:400
Antibody	Anti-rabbit donkey polyclonal antibody, Alexa Fluor® 488 conjugated	Invitrogen	A21206	IF: 1:400
Antibody	Anti-mouse donkey polyclonal antibody, Alexa Fluor® 488 conjugated	Invitrogen	A21202	Flow: 1:400
Antibody	Anti-goat donkey polyclonal antibody, HRP conjugated	Jackson Immunoresearch	705-035-147	WB: 1:5,000
Antibody	Anti-mouse goat polyclonal antibody, HRP conjugated	Biorad	1705047	WB: 1:5,000

Antibody	Anti-rabbit goat polyclonal antibody, HRP conjugated	Biorad	1706515	WB: 1:5,000
Commercial assays and kits	µMACS anti-DYKDDDDK starting kit	Miltenyi Biotec	130-101-636	
software	SMLM image reconstruction	Liu et al. 2018[11]		MATLAB Codes are available upon request.
software	ClusDoc	Pageon et al. 2015[15]		Codes were modified to process the data generated in the custom-built SMLM microscope.
software	OPTICS [16]	This paper		C++ Codes are available upon request.
software	One-dimensional convolutional neural network (1D CNN)	This paper		PYTHON Codes are available upon request.
software	Prism Graphpad	Dotmatics		
software	Flowjo	BD		

Cell lines and plasmids.

PK13, HeLa, and HEK293T cells were cultured at 37°C and 5% CO_2 in DMEM (Sigma-Aldrich,
D6429) complemented with 10% fetal bovine serum (Invitrogen, 12483-020). Cells were
passaged using phosphate-buffered saline (PBS, Invitrogen, 10010-049) and 0.25% Trypsin-
EDTA solution (Invitrogen, 25002-072). Cells were monitored routinely for mycoplasma
contamination using a mycoplasma detection PCR kit (ABM, G238). A FLAG (NiV-F-FLAG) or
HA (NiV-F-HA) tag was inserted after residue 104 of codon-optimized NiV-F in pcDNA3.1 vector.
NiV-F-YA, L53D, V108D, Q393L, and LI4A were produced by site-directed mutagenesis using
the NiV-F-FLAG in pcDNA3.1 vector as a template[6,10,19]. AP2µ2-mcherry plasmid is a gift from
Christien Merrifield (Addgene # 27672). The HA-tagged NiV-G and GFP-tagged NiV-M were
constructed previously.

Immunofluorescence (IF) for SMLM.

For SMLM imaging on cells, 1×10^5 PK13 or HeLa cells were seeded on coverslips (Marienfeld
#1.5H, 18 mm) coated with 2.5 µg fibronectin (Sigma-Aldrich, F4759-2 mg) in a 12-well plate,
and transfected with 1 µg NiV-F variants using lipofectamine 3000 (Invitrogen, L3000015) on the
following day. E64d was dissolved in methanol and pitstop2 in DMSO as recommended by the
manufacturer. E64d was added to cells at 3 hrs post-transfection for a total treatment time of 15
hrs. Pitstop2 was added to cells at 10 hrs post-transfection for a total treatment time of 8 hrs. At
18 hrs posttransfection, cells were fixed with PBS containing 4% paraformaldehyde (PFA;
Electron Microscopy Sciences; 50980487) and 0.2% glutaraldehyde (Sigma-Aldrich, G5882-50

ml) for 90 min at room temperature. Cells were treated with signal enhancer image-IT-Fx (Life Technologies, I36933) for 30 min at room temperature, and then blocked using BlockAid (Life Technologies, B10710) for 1 hr at room temperature. The FLAG-tagged NiV-F and mutants were detected by the anti-FLAG mouse monoclonal antibody (Sigma-Aldrich, F1804) and an Alexa Fluor® 647 conjugated donkey anti-mouse secondary antibody (Invitrogen, A31571). Cells were incubated with primary antibody overnight at 4°C, and then with the secondary antibody for 1 hr at room temperature. Each antibody incubation is followed by five PBS washes, 5 min each time. Cells were then fixed in PBS containing 4% PFA for 10 min at room temperature.

SMLM setup, imaging, and data analysis.

Imaging was performed on a custom-built SMLM. Briefly, the microscope was built upon an apochromatic TIRF oil-immersion objective lens (Nikon, 60x; numerical aperture 1.49). Four lasers were used for excitation: a 639 nm laser (MRL-FN-639, 500 mW) for exciting Alexa Fluor® 647, a 532 nm laser (MGL-III-532-300 mW) for exciting cy3B, a 488 nm laser (MBL-F-473-300 mW) for exciting GFP, and a 405 nm laser (MDL-III-405-100 mW) for reactivating Alexa Fluor 647 and cy3B. The emission fluorescence was separated using appropriate dichroic mirrors and filters (Semrock) and detected by electron multiplying charge-coupled devices (EMCCD; Ixon, Andor). A feedback loop was employed to control the sample drift to <1 nm laterally and 2.5 nm axially.

Fluorescence beads (Life technologies, F8799) were added to samples as fiducial markers for drift control. Samples were immersed in oxygen-scavenging buffer supplemented with 50 mM mercaptoethylamine (Sigma Aldrich, 30070-10G). The expression level of the protein of interest in individual cells was determined by measuring the average emission fluorescence intensity of an area of 27×27 μm^2. For SMLM imaging, samples were exposed to a laser power density of 1 kW/cm^2 for the 639 nm lasers to activate Alexa Fluor® 647. A total of 40,000 images were acquired at 50 Hz for the reconstruction of one SMLM image. Custom-written software in MATLAB (Mathworks) was used to reconstruct SMLM images. Clusters of NiV-F localizations on cell surface resolved by SMLM were identified and characterized using ClusDoC. The min points and ε were set at 4 and 20, respectively.

VLP production and immunofluorescence.

To produce NiV VLPs, HEK293T cells were transfected by NiV-F-FLAG or mutants, NiV-G-HA, and NiV-M-GFP at a 9:1:5 ratio by using polyethylenimine (PEI) at 1 mg/ml (Polysciences, 23966-100). At 48hrs posttransfection, the supernatant of the cell culture was collected and subjected to ultracentrifugation on a 20% sucrose cushion at 125, 392 rcf for 90 min. The VLP-containing pellets were resuspended in 5% sucrose-NTE buffer. VLPs were bound to 2.5 μg fibronectin-coated 18 mm coverslips at 37°C for 4 hrs, followed by fixation using PBS containing 4% PFA and 0.2% glutaraldehyde. The immunofluorescence, SMLM setup, and SMLM imaging for cells were followed to stain and image NiV-F-FLAG and mutants on VLPs using SMLM. A widefield image of GFP was acquired and superimposed on the SMLM image of NiV-F. VLPs were identified as GFP-positive particles. The distribution of NiV-F localizations on VLPs was analyzed by using custom-written OPTICS algorithm based on C++ and Point Cloud Library. The classification of the ordered sequence of reachability distances was performed by using custom written one-dimensional convolutional neural network (1D CNN) based on Python and Tensorflow.

SDS-PAGE, and western blot analysis and co-immunoprecipitation.

For western blot analysis to confirm the expression of NiV-F-FLAG and mutants, HEK293T cells
were seeded in 6-well plate and transfected with 2.5 µg pcDNA3.1 empty vector, NiV-F-FLAG,
or mutants. At 28 or 48 hrs posttransfection, cells were lysed in RIPA buffer (Millipore-Sigma,
20-188) supplemented with protease inhibitor (Sigma-Aldrich, 11836170001) on ice for 30min.
The cell lysates were collected after centrifuge at 16,000 ×g for 20 min at 4°C. The cell lysates
were supplemented with 1x SDS loading dye [60 mM Tris-HCl (pH=6.8); 2% SDS; 10% glycerol,
0.025% Brophenol blue] and 15 mM DTT (Thermo Scientific, R0861), and heated at 95°C for
denature. The denatured cell lysates were loaded into 10% polyacrylamide gels for SDS-PAGE.
Proteins were transferred to PVDF membrane, pore size 0.45 µm (Cytiva, GE10600021).
Membrane was blocked with 1% Bovine serum albumin (Sigma-Aldrich, A9647-50G) in PBS
and incubated with anti-FLAG mouse monoclonal antibody (Sigma-Aldrich, F1804). An HRP-
conjugated goat anti-mouse secondary antibody and the clarity Western ECL substrate (Biorad,
1705060) were used for protein detection. Images were acquired using Chemic Doc MP
Imaging System (BioRad).

For co-immunoprecipitation experiments, HEK293T cells were transfected with the following
combinations 1) 2.3 µg empty pcDNA 3.1 vector and 0.2 µg AP2µ2-mcherry, 2) 0.2 µg AP2µ2-
mcherry and 2.3 µg NiV-F-FLAG, and 3) 0.2 µg AP2µ2-mcherry and 2.3 µg NiV-F-YA. At 48 hrs
post-transfection, cells from 1 well of a 6-well plate were washed with PBS and lysed in 200 µl
lysis buffer provided with the µMACS DYKDDDDK isolation kit, and supplemented with protease
inhibitors. Cells were isolated on ice for 30 min. Cell debris was removed by centrifuge at
16,000 x g for 20 min at 4°C. 60 µl of cell lysate was set aside for immunoblot analysis, and the
rest was used for immunoprecipitation, as recommended by the manufacturer. 6 µl anti-
DYKDDDDK microbeads (Miltenyi Biotec, 130-101-591) were added to 140 µl cell lysates and
incubated for 30min on ice. µ columns (Miltenyi Biotec, 130-101-591) were prepared according
to the manufacturer's instructions. Lysate was run over the columns, and microbeads were
washed according to the manufacturer's instruction. 20 µl preheated elution buffer (95°C) was
added to the column before eluting the bound immunoprecipitated protein in 50 µl elution buffer.
Elute was separated by 10% SDS-PAGE, and proteins were immunoblotted by mouse anti-
FLAG and rabbit anti-mcherry primary antibodies. The HRP-conjugated goat anti-mouse and
goat anti-rabbit secondary antibodies were used for protein detection.

Flow cytometry.

HEK293T cells were seeded in 6-well plate and transfected with 2.5 µg pcDNA3.1 empty vector,
NiV-F-FLAG, or mutants. Cells were collected in 1 mL phosphate-buffered saline (PBS)
containing 10 mM EDTA after 28hrs post-transfection. The collected cells were incubated on ice
for 1 h with primary antibody mouse anti-flag diluted in 1:200. Samples were washed twice in
fluorescence-activated cell sorting (FACS) buffer (0.1% FBS with PBS). After washing, samples
were incubated with fluorescent anti-mouse Alexa Fluor 647 antibody 1:400 on ice for 45 min.
After being washed twice again, samples were read on a flow cytometer (Attune® NxT™
Acoustic Focusing Cytometer, Thermo Fisher) The results were analyzed through FlowJo. The
mean fluorescent intensities (MFI) were normalized to the MFIs of the wt NiV F.

Cell-cell fusion.

HEK293T cells were seeded in the 12-well plate and transfected with 1µg total DNA of NiV-G
and wt-F or mutant-F with a 3:1 ratio using Lipofectamine 3000. Cells were fixed with 4% PFA
after 18 hrs post-transfection. Four or more nuclei within a common cytoplasm were considered

syncytium. Syncytia were quantified by counting the number of nuclei in the syncytium per 10 ×
filed (5 fields are counted per group) under microscope TE2000U.

Acknowledgements. We thank the multiscale imaging facility for instrument use. We thank Dr.
Youssef Chebli at McGill University and Dr. David Scriven at the University of British Columbia
for the support on laser scanning confocal microscopy and single-molecule imaging. **Funding.**
This work is supported by funding from the Canadian Institutes of Health Research to Q.L.
(183861) and the Coronavirus Variants Rapid Response Network to Q.L. (175622). **Author**
contribution: Conceptualization: Q.W. and Q.L.; Image acquisition: Q.W.; Data analysis and
code writing: Q.W., J.L. and Q.L.; Functional analysis: Q.W., Y.L., V.K., G.L.M., J.W.; Resources:
Q.L., Supervision: Q.L.; Funding Acquisition: Q.L.. Writing: original draft: Q.W., J.L., and Q.L.;
revision: all authors. Competing Interests: The authors declare that they have no competing
interests. Code, reagents, and data availability: Custom-written codes are deposited in GitHub
or available upon requests. Reagents can be provided by the corresponding author on a
completed material transfer agreement. Requests for the reagents and data should be
submitted to qian.liu3@mcgill.ca. **Declare of interests.**The authors declare no competing
interests.

1. Navaratnarajah, C.K., Generous, A.R., Yousaf, I., and Cattaneo, R. (2020). Receptor-
mediated cell entry of paramyxoviruses: Mechanisms, and consequences for tropism and
pathogenesis. Journal of Biological Chemistry *295*, 2771–2786.
10.1074/jbc.REV119.009961.

2. Haas, G.D., and Lee, B. (2023). Paramyxoviruses from bats: changes in receptor specificity
and their role in host adaptation. Current Opinion in Virology *58*, 101292.
10.1016/j.coviro.2022.101292.

3. Aditi, and Shariff, M. (2019). Nipah virus infection: A review. Epidemiology & Infection *147*,
e95. 10.1017/S0950268819000086.

4. Thibault, P.A., Watkinson, R.E., Moreira-Soto, A., Drexler, J.F., and Lee, B. (2017). Zoonotic
potential of emerging paramyxoviruses: knowns and unknowns. Adv Virus Res *98*, 1–55.
10.1016/bs.aivir.2016.12.001.

5. Diederich, S., Sauerhering, L., Weis, M., Altmeppen, H., Schaschke, N., Reinheckel, T.,
Erbar, S., and Maisner, A. (2012). Activation of the Nipah Virus Fusion Protein in MDCK
Cells Is Mediated by Cathepsin B within the Endosome-Recycling Compartment. J Virol *86*,
3736–3745. 10.1128/JVI.06628-11.

6. Diederich, S., Moll, M., Klenk, H.-D., and Maisner, A. (2005). The Nipah Virus Fusion Protein
Is Cleaved within the Endosomal Compartment *. Journal of Biological Chemistry *280*,
29899–29903. 10.1074/jbc.M504598200.

7. Negrete, O.A., Levroney, E.L., Aguilar, H.C., Bertolotti-Ciarlet, A., Nazarian, R., Tajyar, S.,
and Lee, B. (2005). EphrinB2 is the entry receptor for Nipah virus, an emergent deadly
paramyxovirus. Nature *436*, 401–405. 10.1038/nature03838.

8. Liu, Q., Bradel-Tretheway, B., Monreal, A.I., Saludes, J.P., Lu, X., Nicola, A.V., and Aguilar,
H.C. (2014). Nipah Virus Attachment Glycoprotein Stalk C-Terminal Region Links Receptor
Binding to Fusion Triggering. J Virol *89*, 1838–1850. 10.1128/JVI.02277-14.

9. Liu, Q., Stone, J.A., Bradel-Tretheway, B., Dabundo, J., Montano, J.A.B., Santos-Montanez,
J., Biering, S.B., Nicola, A.V., Iorio, R.M., and Lu, X. (2013). Unraveling a three-step
spatiotemporal mechanism of triggering of receptor-induced Nipah virus fusion and cell
entry. PLoS Pathog *9*, e1003770.

10. Xu, K., Chan, Y.-P., Bradel-Tretheway, B., Akyol-Ataman, Z., Zhu, Y., Dutta, S., Yan, L.,
Feng, Y., Wang, L.-F., Skiniotis, G., et al. (2015). Crystal Structure of the Pre-fusion Nipah
Virus Fusion Glycoprotein Reveals a Novel Hexamer-of-Trimers Assembly. PLoS Pathog *11*,
e1005322. 10.1371/journal.ppat.1005322.

11. Liu, Q., Chen, L., Aguilar, H.C., and Chou, K.C. (2018). A stochastic assembly model for
Nipah virus revealed by super-resolution microscopy. Nature Communications *9*, 3050.
10.1038/s41467-018-05480-2.

12. Wong, J.J., Chen, Z., Chung, J.K., Groves, J.T., and Jardetzky, T.S. (2021). EphrinB2
clustering by Nipah virus G is required to activate and trap F intermediates at supported lipid
bilayer–cell interfaces. Sci. Adv. *7*, eabe1235. 10.1126/sciadv.abe1235.

13. Coelho, S., Baek, J., Graus, M.S., Halstead, J.M., Nicovich, P.R., Feher, K., Gandhi, H.,
Gooding, J.J., and Gaus, K. (2020). Ultraprecise single-molecule localization microscopy
enables in situ distance measurements in intact cells. Science Advances *6*, eaay8271.
10.1126/sciadv.aay8271.

14. Dempsey, G.T., Vaughan, J.C., Chen, K.H., Bates, M., and Zhuang, X. (2011). Evaluation of
fluorophores for optimal performance in localization-based super-resolution imaging. Nature
Methods *8*, 1027–1036. 10.1038/nmeth.1768.

15. Pageon, S.V., Nicovich, P.R., Mollazade, M., Tabarin, T., and Gaus, K. (2016). Clus-DoC: a
combined cluster detection and colocalization analysis for single-molecule localization
microscopy data. Mol Biol Cell *27*, 3627–3636. 10.1091/mbc.E16-07-0478.

16. Ankerst, M., Breunig, M.M., Kriegel, H.-P., and Sander, J. (1999). OPTICS: ordering points
to identify the clustering structure. SIGMOD Rec. *28*, 49–60. 10.1145/304181.304187.

17. Kiranyaz, S., Avci, O., Abdeljaber, O., Ince, T., Gabbouj, M., and Inman, D.J. (2021). 1D
convolutional neural networks and applications: A survey. Mechanical Systems and Signal
Processing *151*, 107398. 10.1016/j.ymssp.2020.107398.

18. Barrett, C.T., and Dutch, R.E. (2020). Viral Membrane Fusion and the Transmembrane
Domain. Viruses *12*, 693. 10.3390/v12070693.

19. Webb, S., Nagy, T., Moseley, H., Fried, M., and Dutch, R. (2017). Hendra virus fusion
protein transmembrane domain contributes to pre-fusion protein stability. Journal of
Biological Chemistry *292*, 5685–5694. 10.1074/jbc.M117.777235.

20. Bonifacino, J.S., and Neefjes, J. (2017). Moving and positioning the endolysosomal system.
Curr Opin Cell Biol *47*, 1–8. 10.1016/j.ceb.2017.01.008.

21. Bonifacino, J.S., and Traub, L.M. (2003). Signals for sorting of transmembrane proteins to
endosomes and lysosomes. Annu Rev Biochem *72*, 395–447.
10.1146/annurev.biochem.72.121801.161800.

22. Mattera, R., Farías, G.G., Mardones, G.A., and Bonifacino, J.S. (2014). Co-assembly of
Viral Envelope Glycoproteins Regulates Their Polarized Sorting in Neurons. PLOS
Pathogens *10*, e1004107. 10.1371/journal.ppat.1004107.

23. von Kleist, L., Stahlschmidt, W., Bulut, H., Gromova, K., Puchkov, D., Robertson, M.J.,
MacGregor, K.A., Tomilin, N., Pechstein, A., Chau, N., et al. (2011). Role of the Clathrin
Terminal Domain in Regulating Coated Pit Dynamics Revealed by Small Molecule Inhibition.
Cell *146*, 471–484. 10.1016/j.cell.2011.06.025.

24. Johnston, G.P., Bradel-Tretheway, B., Piehowski, P.D., Brewer, H.M., Lee, B.N.R., Usher,
N.T., Zamora, J.L.R., Ortega, V., Contreras, E.M., Teuton, J.R., et al. (2019). Nipah Virus-
Like Particle Egress Is Modulated by Cytoskeletal and Vesicular Trafficking Pathways: a
Validated Particle Proteomics Analysis. mSystems *4*, 10.1128/msystems.00194-19.
10.1128/msystems.00194-19.

25. Pentecost, M., Vashisht, A.A., Lester, T., Voros, T., Beaty, S.M., Park, A., Wang, Y.E., Yun,
T.E., Freiberg, A.N., Wohlschlegel, J.A., et al. (2015). Evidence for ubiquitin-regulated
nuclear and subnuclear trafficking among Paramyxovirinae matrix proteins. PLoS Pathog
11, e1004739. 10.1371/journal.ppat.1004739.

26. Vera-Velasco, N.M., García-Murria, M.J., Sánchez Del Pino, M.M., Mingarro, I., and
Martinez-Gil, L. (2018). Proteomic composition of Nipah virus-like particles. J Proteomics
172, 190–200. 10.1016/j.jprot.2017.10.012.

27. Chojnacki, J., and Eggeling, C. (2018). Super-resolution fluorescence microscopy studies of
human immunodeficiency virus. Retrovirology *15*, 41. 10.1186/s12977-018-0424-3.

28. Muranyi, W., Malkusch, S., Müller, B., Heilemann, M., and Kräusslich, H.-G. (2013). Super-
Resolution Microscopy Reveals Specific Recruitment of HIV-1 Envelope Proteins to Viral
Assembly Sites Dependent on the Envelope C-Terminal Tail. PLOS Pathogens *9*, e1003198.
10.1371/journal.ppat.1003198.

29. Laine, R.F., Goodfellow, G., Young, L.J., Travers, J., Carroll, D., Dibben, O., Bright, H., and
Kaminski, C.F. (2018). Structured illumination microscopy combined with machine learning
enables the high throughput analysis and classification of virus structure. eLife *7*, e40183.
10.7554/eLife.40183.

30. McMahon, A., Andrews, R., Groves, D., Ghani, S.V., Cordes, T., Kapanidis, A.N., and Robb,
N.C. (2023). High-throughput super-resolution analysis of influenza virus pleomorphism
reveals insights into viral spatial organization. PLOS Pathogens *19*, e1011484.
10.1371/journal.ppat.1011484.

31. Chen, Y.-C., Sood, C., Marin, M., Aaron, J., Gratton, E., Salaita, K., and Melikyan, G.B.
(2020). Super-Resolution Fluorescence Imaging Reveals That Serine Incorporator Protein 5
Inhibits Human Immunodeficiency Virus Fusion by Disrupting Envelope Glycoprotein
Clusters. ACS Nano *14*, 10929–10943. 10.1021/acsnano.0c02699.

32. Kovtun, O., Dickson, V.K., Kelly, B.T., Owen, D.J., and Briggs, J.A.G. (2020). Architecture of
the AP2/clathrin coat on the membranes of clathrin-coated vesicles. Sci Adv *6*, eaba8381.
10.1126/sciadv.aba8381.

33. Wang, Y.E., Park, A., Lake, M., Pentecost, M., Torres, B., Yun, T.E., Wolf, M.C., Holbrook,
M.R., Freiberg, A.N., and Lee, B. (2010). Ubiquitin-Regulated Nuclear-Cytoplasmic
Trafficking of the Nipah Virus Matrix Protein Is Important for Viral Budding. PLOS Pathogens
6, e1001186. 10.1371/journal.ppat.1001186.

665