Neurons enhance blood-brain barrier function via upregulating claudin-5

and VE-cadherin expression due to GDNF secretion

Author names

Lu Yang[a], Zijin Lin[a], Ruijing Mu[a], Wenhan Wu[a], Hao Zhi[a], Xiaodong Liu[a, *], Hanyu

Yang[a, *], Li Liu[a, *]

Affiliations

[a]Department of Pharmacology, School of Pharmacy, China Pharmaceutical University,

Nanjing 210009, China

*Correspondence.

Tel./fax: +86 2583271060

Email address: xdliu@cpu.edu.cn (Xiaodong Liu), shenyhy@cpu.edu.cn (Hanyu

Yang), liulee@cpu.edu.cn (Li Liu)

Impact Statement

Neurons and astrocytes enhance the integrity of BBB by releasing GDNF. The

released GDNF upregulated claudin-5 and VE-cadherin expression by the activation

of PI3K/AKT and MAPK/ERK pathways.

Abstract

Blood-brain barrier (BBB) prevents neurotoxins from entering central nervous

system. We aimed to establish and characterize an *in vitro* triple co-culture BBB

model consisting of brain endothelial cells hCMEC/D3, astrocytoma U251 cells, and

neuroblastoma SH-SY5Y cells. Co-culture of SH-SY5Y and U251 cells markedly

enhanced claudin-5 and VE-cadherin expression in hCMEC/D3 cells, accompanied by

increased transendothelial electrical resistance and decreased permeability.

Conditioned medium (CM) from SH-SY5Y cells (S-CM), U251 cells (U-CM), and

co-culture of SH-SY5Y and U251 cells (US-CM) also promoted claudin-5 and

VE-cadherin expression. Glial cell line-derived neurotrophic factor (GDNF) levels in

S-CM and US-CM were significantly higher than CMs from hCMEC/D3 and U-CM.

Both GDNF and US-CM upregulated claudin-5 and VE-cadherin expression, which

were attenuated by anti-GDNF antibody and GDNF signaling inhibitors. GDNF

increased claudin-5 expression via the PI3K/AKT/FOXO1 and MAPK/ERK pathways.

Meanwhile, GDNF promoted VE-cadherin expression by activating PI3K/AKT/ETS1

and MAPK/ERK/ETS1 signaling. The roles of GDNF in BBB integrity were

validated using brain-specific *Gdnf* silencing mice. The developed triple co-culture

BBB model was successfully applied to predict BBB permeability. In conclusion,

neurons enhance BBB integrity by upregulating claudin-5 and VE-cadherin

expression through GDNF secretion and established triple co-culture BBB model may

be used to predict drugs' BBB permeability.

Keywords

Blood-Brain Barrier; *in vitro* BBB model; Glia-derived neurotrophic factor; IVIVC;
Co-culture; Tight junction

Introduction

As a dynamic interface between the blood circulatory system and the central
nervous system (CNS), the blood-brain barrier (BBB) maintains homeostasis and
normal function of CNS by strictly regulating material exchange between the blood
and brain (Palmiotti et al., 2014). The maintenance of BBB is mainly attributed to the
expression of tight junctions (TJ) and adherent junctions (AJ) between adjacent brain
endothelial cells as well as a variety of drug transporters. However, as a double-edged
sword that protects CNS function, BBB also restricts the transport of some drugs from
blood to brain, leading to poor CNS therapeutic effects and even CNS treatment
failure (Banks, 2016).

Several *in silico, in vitro, in situ,* and *in vivo* methods have been developed to
assess the permeability of drugs across BBB, but each method has its limitations
(Hanafy et al., 2021). The *in situ* brain perfusion (ISBP) is considered the "gold
standard" for assessing BBB permeability but there exist limits related to animal
ethics. Moreover, ISBP is not suitable for human or high-throughput workflows. Thus,
a suitable, accurate, and high-throughput *in vitro* BBB model is required to predict the
permeability of drug candidates through BBB.

BBB is formed by neurovascular units (NVU), composed of neural (neurons,
microglia, and astrocytes) and vascular components (vascular endothelial cells,
pericytes, and vascular smooth muscle cells) (Potjewyd et al., 2018). The constant
crosstalk and interactions among these cells contribute to the structural and

signaling-based regulation of transcellular and paracellular transport, control of BBB

permeability, and regulation of cerebral circulation (Arvanitis et al., 2020; Muoio et

al., 2014; Potjewyd et al., 2018). Although brain microvascular endothelial cells

(BMECs) are commonly utilized as an *in vitro* BBB model to assess drug

permeability across BBB, *in vitro* mono-culture of BMECs tends to lose some unique

characteristics without the support of other cell types. To overcome the drawbacks of

mono-culture BBB models, a range of multicellular co-culture BBB models

co-cultured with other NVU elements (such as astrocytes or pericytes) have been

developed. Astrocytes are the most abundant glial cell type in brain (Clasadonte et al.,

2017). Their terminal feet cover >80% of the surface of capillaries to form

interdigitating coverage without slits (Mathiisen et al., 2010). In addition, astrocytes

also contain several proteins related to the tight binding of the basement membrane.

The *in vitro* multicellular co-culture BBB models composed of astrocytes, pericytes,

and endothelial cells are considered to mimic the vascular structure and offer

paracellular tightness in BBB (Ito et al., 2019; Nakagawa et al., 2009; Watanabe et al.,

2021).

Neurons also play a crucial role in the NVU and may be involved in the

regulation of BBB function. The maturation of BBB in mice considerably overlaps

with the establishment of neuronal activity (Biswas et al., 2020). A study revealed that

the conditioned medium collected from primary rat neurons also attenuated cell death

caused by glucose-oxygen-serum deprivation (Lin et al., 2016). Furthermore, the

primary rat neurons were reported to affect the differentiation and formation of

BMECs (Savettieri et al., 2000; Schiera G. et al., 2003; Xue et al., 2013). Recent

research reported that, compared with the double co-culture of hCMEC/D3 and

1321N1 (human astrocytoma cells), the triple co-culture of hCMEC/D3, 1321N1, and

human neuroblastoma SH-SY5Y cells exhibited a higher transendothelial electrical

resistance (TEER) (Barberio et al., 2022). A similar report showed that co-cultivation

of RBE4.B (rat brain capillary endothelial cells) and neurons resulted in the lower

permeability of [^{3}H] sucrose than RBE4.B cells grown alone (Schiera Gabriella. et al.,

2005). All these studies indicate that neurons, as the elements of the NVU, may be

tightly connected to the formation and maintenance of BBB functions.

The aims of the study were: 1) to establish and characterize an *in vitro* triple

co-culture BBB model consisting of human brain endothelial cells (hCMEC/D3),

human astrocytoma cells (U251), and human neuroblastoma cells (SH-SY5Y); 2) to

investigate whether neurons were involved in the formation and maintenance of BBB

integrity and explore their underlying mechanisms. The integrity of BBB was

evaluated by quantifying the leakage of both fluorescein and FITC-Dextran 3–5 kDa

(FITC-Dex); 3) to validate the *in vitro* results through *in vivo* experiments. Finally, the

in vivo/*in vitro* correlation assay (IVIVC) was analyzed to prove that the triple

co-culture BBB model could better predict the BBB penetration of CNS drugs

compared with the mono-culture BBB model.

Result

Establishment and characterization of the *in vitro* triple co-culture BBB

model

Four types of BBB *in vitro* models were established to compare the contributions

of U251 and SH-SY5Y cells to hCMEC/D3 cells (Figure 1A). TEER values were

measured during the co-culture (Figure 1B). TEER values of the four *in vitro* BBB

models gradually increased until day 6. On day 7, the TEER values showed a

decreasing trend. Thus, six-day co-culture period was used for subsequent

experiments. The highest TEER values were observed in the triple co-culture BBB

model, followed by double co-culture with U251 cells and double co-culture with

SH-SY5Y cells BBB models, hCMEC/D3 cells mono-culture BBB model showed the

lowest TEER values. We also measured the TEER values of U251 monolayer cells,

and the results showed that U251 cells themselves also contributed to the physical

barrier of the model (Figure 1C). The apparent permeability coefficient (P_{app}) values

of fluorescein and FITC-Dex were measured to characterize the integrity of four BBB

models (Figure 1D, 1E). Consistent with the TEER values, the triple co-culture BBB

model showed the lowest permeability of fluorescein and FITC-Dex, followed by

double co-culture with U251 cells and double co-culture with SH-SY5Y cells BBB

models. It was noticed that monolayer of U251 cells itself also worked as a barrier,

preventing the leakage of permeability markers, which may explain why the

permeability of FITC-Dex in double co-culture model with U251 cells is lower than

that in double co-culture model with SH-SY5Y cells (Figure 1F). Co-culture with

SH-SY5Y, U251, and U251 + SH-SY5Y cells also enhanced the proliferation of

hCMEC/D3 cells. Moreover, the promoting effect of SH-SY5Y cells was stronger

than that of U251 cells (Figure 1G-1I). Furthermore, hCMEC/D3 cells were incubated

with basic fibroblast growth factor (bFGF), which promotes cell proliferation without

affecting both claudin-5 and VE-cadherin expression (Figure 2F). The results showed

that incubation with bFGF increased cell proliferation and reduced permeabilities of

fluorescein and FITC-Dex across hCMEC/D3 cell monolayer. However, the

permeability reduction was less than that by double co-culture with U251 cells or

triple co-culture. These results inferred that contribution of cell proliferation to the

barrier function of hCMEC/D3 was minor (Figure 1—figure supplement 1).

The paracellular barrier of BBB is also associated with TJs, AJs, and transporters

(Abbott, 2013). The mRNA levels of TJs (claudin-5, ZO-1, and occludin), AJs

(VE-cadherin and β-catenin), and transporters (P-gp, BCRP, OCT-1, OCT-2, OAT-3,

and OATP1A1) in hCMEC/D3 cells from the four BBB models were analyzed using

qPCR (Figure 1J and 1K). Compared with hCMEC/D3 cell mono-culture model,

double co-culture with SH-SY5Y, double co-culture with U251, and triple co-culture

BBB models showed markedly increases in claudin-5, VE-cadherin, β-catenin, and

BCRP mRNA expression. Expression of corresponding proteins was measured using

Western blot (Figure 1L-1O). Notably increased claudin-5 expression was detected in

double co-culture with SH-SY5Y cells and triple co-culture BBB models, while

VE-cadherin expression was markedly increased in double co-culture with U251 cells

and triple co-culture BBB models. Expression levels of other TJ proteins (ZO-1 and

occludin) and AJ protein (β-catenin) were unaltered. The expression of BCRP was

slightly affected by co-cultivation with U251 cells. Significant negative correlations

were found between P_{app} values of fluorescein and the expression of claudin-5 or

VE-cadherin. P_{app} values of FITC-Dex were also negatively correlated to the

expression levels of claudin-5 or VE-cadherin (Figure 1P-1S). These results indicate

that the decreased permeability of fluorescein and FITC-Dex mainly results from the

upregulated expression of both claudin-5 and VE-cadherin.

Neurons and astrocytes upregulated the expression of claudin-5 and

VE-cadherin by GDNF secretion.

The hCMEC/D3 cells did not direct contact with U251 or SH-SY5Y cells in the

double co-culture and triple co-culture BBB models, indicating that cell-cell

interaction between U251, SH-SY5Y, and hCMEC/D3 cells relied on secreted active

factors. To test this hypothesis, the effects of conditioned medium (CM) from

SH-SY5Y cells (S-CM), U251 cells (U-CM), and co-culture of SH-SY5Y and U251

cells (US-CM) on the expression of claudin-5 and VE-cadherin in hCMEC/D3 cells were analyzed (Figure 2A). Both S-CM, U-CM, and US-CM markedly increased the expression of claudin-5 and VE-cadherin. US-CM showed the strongest induction effects on claudin-5 and VE-cadherin.

To investigate which cytokines were involved in the promotion of hCMEC/D3 cell integrity by U251 and SH-SY5Y cells, the mRNA expression levels of various cytokines in these three types of cells were compared. The results showed that U251 or SH-SY5Y cells exhibited significantly higher expression levels of GDNF, nerve growth factor (NGF), insulin-like growth factor-1 (IGF-1), vascular endothelial growth factor (VEGF), bFGF, and transforming growth factor-β (TGF-β) compared to hCMEC/D3 cells (Figure 2B). Furthermore, the mRNA expression of GDNF, IGF-1, TGF-β, and bFGF in SH-SY5Y cells was higher than those in U251 cells

In these cytokines, GDNF (Dong et al., 2018; Igarashi et al., 1999; Shimizu et al., 2012), bFGF (Shimizu et al., 2011; Wang et al., 2016), IGF-1 (Ji-Ae K et al., 2009; Nowrangi et al., 2019), and TGF-β (Fu et al., 2021) have been reported to promote BBB integrity. Thus, the concentrations of GDNF, bFGF, IGF-1, and TGF-β in the CMs were measured (Figure 2C). The results showed that levels of GDNF, bFGF, and IGF-1 in S-CM and US-CM were significantly higher than CMs from hCMEC/D3 cells (H-CM) and U-CM, but levels of TGF-β in S-CM and U-CM were lower than those in H-CM. Interestingly, the level of IGF-1 in US-CM was remarkably lower than that in S-CM, indicating that U251 cells suppressed IGF-1 secretion from SH-SY5Y. The effects of GDNF, bFGF, IGF-1, and TGF-β on the expression of claudin-5 and VE-cadherin were investigated (Figure 2D-2G). Among the four tested neurotrophic factors, only GDNF induced the expression of claudin-5 and VE-cadherin in a concentration-dependent manner (Figure 2D). In contrast, a high

level (5 ng/mL) of TGF-β slightly downregulated claudin-5 expression (Figure 2G).

These results demonstrate that upregulation of claudin-5 and VE-cadherin expression

by US-CM are attributed to secreted GDNF.

To provide additional verification of the deduction, anti-GDNF antibody was

used to neutralize exogenous and endogenous GDNF in culture medium. Consistent

with our expectation, the anti-GDNF antibody concentration-dependent reversed the

US-CM-induced claudin-5 and VE-cadherin expression (Figure 2H). Furthermore,

GDNF-induced upregulation of claudin-5 and VE-cadherin expression was also

reversed by the anti-GDNF antibody (Figure 2I). The induction effects of US-CM and

GDNF on the claudin-5 and VE-cadherin expression in hCMEC/D3 cells were also

confirmed in primary rat brain microvascular endothelial cells (Figure 2J)

GDNF forms a heterohexameric complex with two GFRα1 molecules and two

RET receptors to activate the GDNF-GFRα1-RET signaling (Fielder et al., 2018). The

RET receptor tyrosine kinase inhibitor SPP-86 (Bhallamudi et al., 2021) and Src-type

kinase inhibitor PP2 (Morita et al., 2006) were used to further investigate whether

GDNF upregulated the expression of claudin-5 and VE-cadherin in hCMEC/D3 cells

by activating the GDNF-GFRα1-RET signaling pathway. Both SPP-86 and PP2

markedly attenuated claudin-5 and VE-cadherin expression induced by GDNF (Figure

2K) and US-CM (Figure 2L).

The contributions of GDNF-induced claudin-5 and VE-cadherin expression to

TEER and permeability were investigated using hCMEC/D3 cells mono-culture BBB

model. GDNF significantly increased TEER values (Figure 2M, 2O) and decreased

the permeability of fluorescein and FITC-Dex (Figure 2N, 2P), which were almost

abolished by SPP-86 or PP2. Furthermore, treatment with SPP-86 or PP2 completely

reversed the increased TEER values (Figure 2Q, 2S) and decreased permeability of

fluorescein and FITC-Dex (Figure. 2R, 2T) in the triple co-culture BBB model. These

results indicate that neurons but also astrocytes upregulate claudin-5 and VE-cadherin

expression in hCMEC/D3 cells by secreting GDNF. Subsequently, GDNF induces

claudin-5 and VE-cadherin expression by activating GDNF-GFRα1-RET signaling.

GDNF induced claudin-5 and VE-cadherin expression of hCMEC/D3 by

activating the PI3K/AKT and MAPK/ERK pathways

GDNF exerts its biological activities by activating several signaling pathways,

including the phosphatidylinositol-3-kinase (PI3K)/ protein kinase B (AKT),

mitogen-activated protein kinase (MAPK)/ extracellular regulated kinase (ERK),

MAPK/ c-Jun N-terminal kinase (JNK), and MAPK/ p38 pathways (Fielder et al.,

2018). The effects of the PI3K/AKT, MAPK/ERK, MAPK/JNK, and MAPK/p38

pathway inhibitors LY294002 (Figure 3A), U0126 (Figure 3B), SP600125 (Figure

3C), and SB203580 (Figure 3D), respectively, on GDNF-induced claudin-5 and

VE-cadherin expression in hCMEC/D3 cells were investigated. GDNF increased

claudin-5 and VE-cadherin expression, accompanied by the phosphorylation of AKT

(p-AKT) and ERK (p-ERK). However, it did not stimulate the phosphorylation of

JNK and p38. SPP-86, LY29002, and U0126 significantly suppressed GDNF-induced

claudin-5 and VE-cadherin expression, while SP600125 and SB203580 had almost no

effect on GDNF-induced claudin-5 and VE-cadherin expression. GDNF-induced

phosphorylation of AKT and ERK was also markedly attenuated by the anti-GDNF

antibody (Figure 3E). Similarly, US-CM remarkably upregulated the expression of

claudin-5, VE-cadherin, p-AKT, and p-ERK, which were also markedly reversed by

SPP-86, LY29002, U0126, or anti-GDNF antibody (Figure 3F-3J). These findings

indicate that GDNF induces the expression of claudin-5 and VE-cadherin in

hCMEC/D3 cells by activating both the PI3K/AKT and MAPK/ERK pathways.

GDNF upregulated the claudin-5 expression in hCMEC/D3 cells by activating the PI3K/AKT/FOXO1 pathway

Claudin-5 is negatively regulated by the transcriptional repressor forkhead box O1 (FOXO1) (Beard et al., 2020). FOXO1 is also an important target of PI3K/AKT signaling. FOXO1 phosphorylation results in FOXO1 accumulation in the cytoplasm (Zhang et al., 2011) and lowers its level in the nucleus. Here, we investigated whether GDNF-induced claudin-5 expression is involved in FOXO1 nuclear exclusion. As shown in Figure 4A, both GDNF and US-CM significantly enhanced FOXO1 phosphorylation (p-FOXO1). Similarly, GDNF and US-CM increased the levels of phosphorylated and unphosphorylated FOXO1 in the cytoplasm and decreased the levels of nuclear FOXO1 (Figure 4B). Whether FOXO1 was involved in the GDNF-induced regulation of claudin-5 and VE-cadherin expression was investigated in hCMEC/D3 cells transfected with *FOXO1* siRNA. Silencing *FOXO1* significantly decreased FOXO1 levels in both whole-cell lysates and nucleus of hCMEC/D3 cells (Figure 4C), demonstrating *FOXO1* silencing efficacy. Consistent with our expectation, silencing *FOXO1* upregulated the expression of claudin-5 rather than VE-cadherin expression (Figure 4D). In contrast, high levels of FOXO1 were observed in both whole-cell lysates and nucleus of hCMEC/D3 cells that were transfected with plasmids containing *FOXO1*. Meanwhile, FOXO1 overexpression resulted in a decrease in both basal and GDNF-induced claudin-5 expression (Figure 4E), consistent with the known role of FOXO1 on claudin-5 expression.

Several reports have demonstrated that nuclear localization of FOXO1 is modulated by multiple pathways, including the PI3K/AKT (Tang et al., 1999) and MAPK/ERK (Asada et al., 2007) pathways. The effects of LY294002 and U0126 on GDNF-induced FOXO1 phosphorylation were measured in hCMEC/D3 cells.

LY294002, but not U0126, significantly reversed the GDNF-induced alterations in

total p-FOXO1, cytoplasmic p-FOXO1, cytoplasmic FOXO1, and nuclear FOXO1

(Figure 4F). Furthermore, LY294002 reversed the upregulation of claudin-5

expression induced by *FOXO1* siRNA and GDNF (Figure 4G).

It was reported that VE-cadherin also upregulates claudin-5 via inhibiting

FOXO1 activities (Taddei et al., 2008). Effect of VE-cadherin on claudin-5 was

studied in hCMEC/D3 cells silencing VE-cadherin. It was not consistent with Taddei

et al. that silencing VE-cadherin only slightly decreased the mRNA level of claudin-5

without significant difference. Furthermore, basal and GDNF-induced claudin-5

protein levels were unaltered by silencing VE-cadherin (Figure 4—figure supplement

1). Thus, the roles of VE-cadherin in regulation of claudin-5 in BBB should be further

investigated.

GDNF upregulated VE-cadherin expression in hCMEC/D3 cells by

activating the PI3K/AKT/ETS1 and MAPK/ERK/ETS1 signaling

pathways.

E26 oncogene homolog 1 (ETS1) is a transcription factor that binds to the

ETS-binding site located in the proximal region of the VE-cadherin promoter, hence

regulating the expression of VE-cadherin (Lelie`vre et al., 2000; Luo et al., 2022).

Activation of the PI3K/AKT (He et al., 2023; Hui et al., 2018) and MAPK/ERK

(Watanabe et al., 2004) pathways was reported to upregulate the expression of ETS1.

The previous results showed that GDNF induced VE-cadherin and claudin-5

expression in hCMEC/D3 cells by activating the PI3K/AKT and MAPK/ERK

pathways. Therefore, we hypothesized that GDNF modulated ETS1 levels to promote

VE-Cadherin and claudin-5 expression through PI3K/AKT and MAPK/ERK

signaling pathways. Both US-CM and GDNF significantly increased total (Figure 5A)

and nuclear ETS1 expression (Figure 5B). LY294002 and U0126 markedly attenuated

GDNF-induced total (Figure 5C) and nuclear ETS1 expression (Figure 5D). To

further confirm the involvement of the PI3K/AKT/ETS1 and MAPK/ERK/ETS1

pathways in GDNF-induced VE-cadherin expression, ETS1 in hCMEC/D3 cells was

knocked down using *ETS1* siRNA. *ETS1* silencing saliently declined the expression

levels of total (Figure 5E) and nuclear (Figure 5F) ETS1 in hCMEC/D3 cells,

demonstrating silencing efficacy. In *ETS1* silencing hCMEC/D3 cells, GDNF no

longer induced the expression of total (Figure 5E) and nuclear (Figure 5F) ETS1.

Moreover, *ETS1* silencing substantially downregulated VE-cadherin expression and

attenuated GDNF-induced VE-cadherin expression (Figure 5G and 5H), while having

minimal impact on both basal and GDNF-induced claudin-5 expression (Figure 5G

and 5I).

Brain GDNF deficiency increased BBB permeability partly due to the

impairment of claudin-5 and VE-cadherin expression

To further demonstrate the positive effects of GDNF on BBB maintenance,

GDNF in mice brains was knocked down via intracerebroventricular injection of

AAV-PHP.eB packaged with *Gdnf* short hairpin RNA (shRNA) (Figure 6A).

Knockdown efficiency was confirmed through Western blotting (Figure 6B).

Consistent with *in vitro* results, GDNF knockdown greatly downregulated claudin-5

and VE-cadherin expression in the mice brains (Figure 6B). The integrity of BBB was

assessed by examining the brain distributions of fluorescein and FITC-Dex. The

results showed that specifically knocking down brain GDNF little affected plasma

levels of fluorescein (Figure 6C) and FITC-Dex (Figure 6F), but significantly elevated

the concentrations of fluorescein (Figure 6D) and FITC-Dex (Figure 6G) in the brains,

leading to notable increases in the brain-to-plasma concentration ratios of two probes

(Figure 6E and 6H). These alterations were consistent with the decline in claudin-5

and VE-cadherin expression. In addition, significant reductions in the levels of p-AKT

(Figure 6I), p-ERK (Figure 6J), p-FOXO1 (Figure 6K), and ETS1 expression (Figure

6L) were observed in the brains of *Gdnf* knockdown mice.

The triple co-culture BBB model better predicted the permeabilities of drugs across BBB

In this study, 18 drugs were utilized to further investigate the superiority of the

triple co-culture BBB model over the hCMEC/D3 mono-culture BBB model. The P_{app}

of 18 drugs from the apical to the basolateral side based on the hCMEC/D3

mono-culture ($P_{app, Mono}$) and triple co-culture ($P_{app, Triple}$) BBB models are measured

and listed in Table 1. The results showed that $P_{app, Triple}$ values of all tested drugs were

lower than the $P_{app, Mono}$ values. Significant differences were observed in 14 out of 18

drugs. The predicted permeability coefficient-surface area product values (PS) of the

tested drugs were respectively calculated based on their $P_{app, Mono}$ values ($PS_{Pre, Mono}$),

and $P_{app, Triple}$ values ($PS_{Pre, Triple}$). The predicted PS values were further compared to

their corresponding observations (PS_{obs}). The results showed that the predictive

accuracy of $PS_{Pre, Triple}$ was superior to $PS_{Pre, Mono}$. Except for verapamil, amitriptyline,

fluoxetine, and clozapine, the predicted $PS_{Pre, Triple}$ values of the other 14 drugs in the

triple co-culture BBB models were within the 0.5–2 folds of PS_{obs} (Figure 7B).

However, in the hCMEC/D3 mono-culture BBB model, only 7 predicted $PS_{Pre, Mono}$

values were within the 0.5–2 folds range of their observations (Figure 7A).

Discussion

The main findings of the study were to successfully develop an *in vitro* triple

co-culture BBB model consisting of hCMEC/D3, U251, and SH-SY5Y cells and to

confirm the involvement of neurons in BBB maintenance as well as the possible

mechanisms. Co-culture with U251 and/or SH-SY5Y cells markedly promoted the

TEER of hCMEC/D3 cells and reduced the leakage of fluorescein and FITC-Dex due

to the upregulation of claudin-5 and VE-cadherin expression.

The roles of claudin-5 and VE-cadherin in the maintenance of BBB function

have been demonstrated (Dejana et al., 2008; Hashimoto et al., 2023; Li et al., 2018;

Ohtsuki et al., 2007). It was reported that *Cld-5*–deficient mice exhibited BBB

impairment, allowing the transport of small molecules (<800 D) across BBB (Nitta et

al., 2003). In contrast, claudin-5 overexpression significantly restricted the

permeability of inulin across the conditionally immortalized rat brain capillary

endothelial cell monolayer (Ohtsuki et al., 2007). Moreover, the claudin-5 expression

in hCMEC/D3 is much lower than in human brain microvessels (Eigenmann et al.,

2013; Ohtsuki et al., 2013; Weksler et al., 2013), which may cause the low TEER

values. VE-cadherin, a major member of the cadherin family in endothelial cells, is

also required for BBB integrity (Dejana et al., 2008; Li et al., 2018). The absence of

VE-cadherin resulted in faulty cell-to-cell junctions and disrupted distribution of

ZO-1 (Sauteur et al., 2017; Tunggal et al., 2005). Strongly negative correlations

between P_{app} values of fluorescein or FITC-Dex and expression of claudin-5 or

VE-cadherin further demonstrated the importance of claudin-5 and VE-cadherin in

BBB integrity.

Neurons and astrocytes, as important components of the NVU, may be involved

in the formation and maintenance of BBB function. Several reports showed that

co-culture with astrocytes CC-2565 or SC-1810 enhanced the TEER of hCMEC/D3

cells (Hatherell et al., 2011), whereas co-culture of RBE4.B cells with primary rat

neurons and astrocytes showed a lower permeability of [³H] sucrose than

mono-culture of RBE4.B cells (Schiera Gabriella. et al., 2005). Similarly, co-cultured

with hCMEC/D3 cells and 1321N1 (astrocytes) or 1321N1+SH-SY5Y cells possess

higher TEER values and lower permeability to Lucifer yellow than hCMEC/D3 alone.

Compared with double co-culture of hCMEC/D3 and 321N1 cells, the triple

co-culture of hCMEC/D3, 1321N1, and SH-SY5Y cells showed higher TEER values

(Barberio et al., 2022). Our study also demonstrated that co-culture with U251 and/or

SH-SY5Y cells significantly lowered BBB permeability and upregulated VE-cadherin

or claudin-5 expression in hCMEC/D3 cells.

Next, we focused on the molecular mechanisms by which neurons and possibly

astrocytes upregulated the VE-cadherin and claudin-5 expression in BMECs.

Co-culture with SH-SY5Y cells significantly upregulated claudin-5 and VE-cadherin

expression in hCMEC/D3 cells. In the double co-culture with SH-SY5Y cells or triple

co-culture BBB models, hCMEC/D3 cells were not in direct contact with SH-SY5Y

cells, indicating that the interaction between SH-SY5Y and hCMEC/D3 cells

depended on the release of some active compounds. It was consistent with the above

deduction that the S-CM also markedly induced claudin-5 and VE-cadherin

expression. Different from S-CM, U-CM mainly upregulated the VE-cadherin

expression and just had a slight impact on claudin-5. In general, neurons but also

astrocytes secrete some neurotrophic factors (Lonka-Nevalaita et al., 2010; Sweeney

et al., 2019) that could contribute to the maintenance of structural stability of BBB.

High levels of bFGF, GDNF, IGF-1, and TGF-β were detected in U-CM, S-CM, and

US-CM. Further research showed that only GDNF induced VE-cadherin and

claudin-5 expression in hCMEC/D3 cells in a concentration-dependent manner, and

the anti-GDNF antibody attenuated claudin-5 and VE-cadherin expression induced by

US-CM or GDNF. These results indicated that neurons upregulated claudin-5 and

VE-cadherin expression through GDNF secretion. Levels of GDNF in S-CM were higher than those in U-CM, which seemed to partly explain why S-CM has a stronger promoting effect on claudin-5 than U-CM. The roles of GDNF in BBB maintenance and the regulation of claudin-5 and VE-cadherin expression were further confirmed using brain-specific *Gdnf* knockdown C57BL/6J mice. Consistent with our *in vitro* results, brain-specific *Gdnf* silencing greatly increased BBB penetration of fluorescein and FITC-Dex, accompanied by the downregulation of claudin-5 and VE-cadherin expression.

GDNF is mainly expressed in astrocytes and neurons (Lonka-Nevalaita et al., 2010; Pochon et al., 1997). In adult animals, GDNF is mainly secreted by striatal neurons rather than astrocytes and microglial cells (Hidalgo-Figueroa et al., 2012). The present study also shows that GDNF mRNA levels in SH-SY5Y cells were significantly higher than that in U251 cells. GDNF was also detected in conditioned medium from SH-SY5Y cells. All these results demonstrate that neurons may secrete GDNF.

Generally, GDNF activates several signal transduction pathways, such as the PI3K/AKT and MAPK signaling pathways (Fielder et al., 2018) by forming a heterohexameric complex with two GFRα molecules and RET receptors. It was also reported that GDNF improved BBB barrier function due to the activation of MAPK/ERK1 (Dong et al., 2018) and PI3K/AKT (Liu D. et al., 2022) signaling. Consistent with previous reports, we found that signaling inhibitors SPP-86, PP2, LY294002, and U0126 markedly attenuated US-CM- and GDNF-induced claudin-5 and VE-cadherin expression in hCMEC/D3 cells, inferring that GDNF promoted claudin-5 and VE-cadherin expression via activating both the PI3K/AKT and MAPK/ERK pathways.

Signal transduction pathways control gene expression by modifying the function

of nuclear transcription factors. The nuclear accumulation of FOXO1 negatively

regulates claudin-5 expression (Beard et al., 2020; Taddei et al., 2008). FOXO1 is an

important target of the PI3K/AKT signaling axis (Zhang et al., 2011), and

AKT-induced phosphorylation of FOXO1 results in cytoplasmic FOXO1

accumulation and decreases nuclear FOXO1 accumulation (Zhang et al., 2011). From

these results, we inferred that GDNF-induced claudin-5 expression in hCMEC/D3

cells may be involved in the activation of PI3K/AKT/FOXO1 pathway. Similarly,

both US-CM and GDNF increased cytoplasmic p-FOXO1 and decreased nuclear

FOXO1 in hCMEC/D3 cells, which was reversed by LY29002 rather than U0126.

Roles of FOXO1 in GDNF-induced claudin-5 were verified through silencing and

overexpressing *FOXO1* in hCMEC/D3 cells. *FOXO1* silencing enhanced claudin-5

but not VE-cadherin expression, accompanied by a decline in total and nuclear

FOXO1. In contrast, *FOXO1* overexpression significantly decreased claudin-5

expression. In hCMEC/D3 cells overexpressing *FOXO1*, GDNF lost its promotion

effect on claudin-5 expression. It was noticed that U0126 attenuated the

GDNF-induced upregulation of claudin-5 but had minimal impact on

GDNF-mediated FOXO1 phosphorylation and the decline of nuclear FOXO1. In

Sertoli cells, it was found that testosterone stimulated claudin-5 expression by

activating the RAS/RAF/ERK/CREB pathway (Bulldan et al., 2016). The

transcriptional regulation of claudin-5 by CREB was confirmed in bEnd.3 (mouse

brain endothelial cell). CREB overexpression significantly increased both gene and

protein expression of claudin 5. In contrast, depletion of CREB decreased claudin-5

expression in gene and protein levels (Li Y. et al., 2022). However, another report

showed that in human lung microvascular endothelial cells, U0126 attenuated

phosphorylation of ERK and lipopolysaccharide-stimulated claudin-5 damage,

indicating activation of MAPK/ERK pathway impaired rather than promoted

claudin-5 expression (Liu Y. et al., 2019). Thus, the real mechanisms that

GDNF-induced activation of the RET/MAPK/ERK pathway promotes claudin-5

expression need further investigation.

ETS1 is a member of the ETS family that plays an important role in cell

adhesion, migration, and blood vessel information. ETS1 binds to an ETS-binding site

located in the proximal region of the VE-cadherin promoter, controlling VE-cadherin

expression (Lelie`vre et al., 2001). Several studies have demonstrated the role of

ETS1 in the regulation of VE-cadherin expression. For example, IFN-γ and TNF-α

impaired BBB integrity by decreasing ETS1-induced VE-cadherin expression (Luo et

al., 2022). *ETS1* silencing reduced VE-cadherin expression in umbilical vein

endothelial cells (Colas-Algora et al., 2020). In contrast, *ETS1* overexpression

induced VE-cadherin expression in mouse brain capillary endothelial cells and

fibroblasts (Lelie`vre et al., 2000). As expected, *ETS1* silencing resulted in a decrease

in the expression of VE-cadherin in hCMEC/D3 cells, but claudin-5 expression

remained unaffected. Additionally, *ETS1* silencing removed the inductive effect of

GDNF on VE-cadherin expression while unaffecting the upregulation of claudin-5

induced by GDNF. The activation of the PI3K/AKT (He et al., 2023; Hui et al., 2018)

and MAPK/ERK (Watanabe et al., 2004) pathways promotes ETS1 expression. In our

findings, US-CM and GDNF significantly increased total and nuclear ETS1 levels,

which were eliminated by signaling inhibitors LY294002 and U0126. Both our results

and previous research provide evidence that GDNF upregulates ETS1 expression via

the activation of PI3K/AKT and MAPK/ERK signaling to promote VE-cadherin

expression.

Claudin-5 expression is also regulated by VE-cadherin (Taddei et al., 2008).

Differing from the previous reports, silencing VE-cadherin with siRNA only slightly

affected basal and GDNF-induced claudin-5 expression. The discrepancies may come

from different characteristics of the tested cells. Several reports have supported the

above deduction. In retinal endothelial cells, hyperglycemia remarkably reduced

claudin-5 expression (but not VE-cadherin) (Saker et al., 2014). However, in

hCMEC/D3 cells, hypoglycemia significantly decreased claudin-5 expression but

hyperglycemia increased VE-cadherin expression (Sajja et al., 2014).

The present study showed that characteristics of *in vitro* triple co-culture BBB

model were superior to those of hCMEC/D3 mono-culture BBB model. The

hCMEC/D3 mono-culture and triple co-culture BBB models were used to try to

predict the *PS* values of 18 drugs by comparing them with their observations. The

prediction success rate (14/18) of triple co-culture BBB model was greater than that

of hCMEC/D3 mono-culture BBB model (7/18). However, poor predictions were

observed for verapamil, amitriptyline, fluoxetine, and clozapine, which may partly be

due to inaccuracies in their $f_{u, brain}$ values. These four drugs are high protein-binding

drugs. Due to the methodological discordance and limitations of historic devices for

these drugs, the $f_u \leq 0.01$ maybe with low confidence and accuracy (Bowman et al.,

2018; Di et al., 2017). For example, $f_{u, brain}$ values of amitriptyline from different

reports display large differences (0.002 (Sanchez-Dengra et al., 2021), 0.0071 (Weber

et al., 2013)). The same large differences of $f_{u, brain}$ values were also found in

fluoxetine (0.0023 (Maurer et al., 2005), 0.004 (Summerfield et al., 2007), 0.00094

(Liu X. et al., 2005)), and clozapine (0.0056 (Bhyrapuneni et al., 2018), 0.014

(Cremers et al., 2012), 0.0094 (Summerfield et al., 2007)).

In summary, our triple co-culture BBB model outperformed the mono-culture or

double co-culture BBB models, mainly attributing to the fact that neurons and

possibly astrocytes upregulated claudin-5 and VE-cadherin expression by secreting

GDNF through activating PI3K/AKT and MAPK/ERK pathways. Additionally, the

developed *in vitro* triple co-culture BBB model accurately predicted the *in vivo* BBB

permeability of CNS drugs. This suggests the potential of our triple co-culture BBB

model for utilization in CNS candidate screening during the drug development

process (Figure 8).

However, the study also has some limitations. In addition to neurons and

astrocytes, other cells such as microglia, pericytes, and vascular smooth muscle cells,

especially pericytes, may also affect BBB function. How pericytes affect BBB

function and interaction among neurons, astrocytes, and pericytes needs further

investigation.

Materials and Methods

Key Resources Table				
Reagent type (species) or resource	**Designation**	**Source or reference**	**Identifiers**	**Additional information**
Antibody	*β*-actin (mouse monoclonal)	Proteintech	66009 RRID:AB_2883475	1:10000
Antibody	GAPDH (mouse monoclonal)	Absin	Abs830030ss RRID:AB_2811228	1:50000
Antibody	*β*-Tubulin (mouse monoclonal)	Fdbio Science	FD0064 RRID: AB_3076327	1:10000

Antibody	Lamin B (mouse monoclonal)	Proteintech	66095 RRID:AB_2721256	1:10000
Antibody	Claudin-5 (rabbit polyclonal)	Wanleibio	WL03731 RRID: AB_3076320	1:1000
Antibody	Occludin (rabbit polyclonal)	Wanleibio	WL01996 RRID:AB_3076325	1:500
Antibody	ZO-1 (mouse polyclonal)	Proteintech	21773-1-AP RRID:AB_10733242	1:5000
Antibody	VE-cadherin (rabbit polyclonal)	Wanleibio	WL02033 RRID:AB_3076321	1:1000
Antibody	β-catenin (rabbit polyclonal)	Wanleibio	WL0962a RRID: AB_3076323	1:5000
Antibody	BCPR (rabbit polyclonal)	CST	4477S RRID:AB_10544928	1:1000
Antibody	P-gp (rabbit monoclonal)	CST	13978S RRID:AB_2798357	1:1500
Antibody	p-AKT (mouse monoclonal)	Huaan Biotechnology	ET1607 RRID:AB_2940863	1:2000
Antibody	AKT (mouse monoclonal)	Huaan Biotechnology	ET1609 RRID:AB_3069857	1:2000
Antibody	p-ERK (rabbit polyclonal)	Proteintech	28733-1-AP RRID:AB_2881202	1:1000
Antibody	ERK (rabbit polyclonal)	Proteintech	11257-1-AP RRID:AB_2139822	1:1000
Antibody	p-p38 (rabbit monoclonal)	CST	4511S RRID:AB_10890701	1:250
Antibody	p38 (rabbit	CST	8690S RRID:AB_10999090	1:250

	monoclonal)			
Antibody	p-JNK (rabbit polyclonal)	Wanleibio	WL01813 RRID:AB_2910628	1:1000
Antibody	JNK (rabbit polyclonal)	Wanleibio	WL01295 RRID:AB_3064853	1:1000
Antibody	FOXO1 (rabbit polyclonal)	Proteintech	18592 RRID:AB_2934932	1:1000
Antibody	p-FOXO1 (rabbit polyclonal)	Wanleibio	WL03634 RRID: AB_3076326	1:1000
Antibody	ETS1 (mouse monoclonal)	Santa Cruz	sc-55581 RRID:AB_831289	1:500
Antibody	ETS1 (mouse monoclonal)	Proteintech	66598 RRID:AB_2881958	1:3000
cell line (*Homo sapiens*)	hCMEC/D3 cells	JENNIO Biological Technology	Cat#JNO-H0520	Authenticated (STR profiling)
cell line (*Homo sapiens*)	U251 cells	Cellcook Biological Technology	Cat#CC1701	Authenticated (STR profiling)
cell line (*Homo sapiens*)	SH-SY5Y cells	Cellcook Biological Technology	Cat#CC2101	Authenticated (STR profiling)
Software, algorithm	GraphPad Prism	Version 8.0.2	RRID:SCR_002798	
Software, algorithm	BioTek Cytation 5 Cell Imaging Multi-Mode Reader	BioTek Cytation 5	RRID:SCR_019732	
Software, algorithm	QuantStudio 3 Real Time PCR System	QuantStudio 3	RRID:SCR_018712	
commercial	GDNF-Elisa kit	R&D system	Cat#212-GD	

assay or kit				
commercial assay or kit	bFGF-Elisa kit	Elabscience	Cat#E-EL-H6042	
commercial assay or kit	IGF-1-Elisa kit	Elabscience	Cat#E-EL-H0086	
commercial assay or kit	TGF-β-Elisa kit	Elabscience	Cat#E-EL-0162	
peptide, recombinant protein	GDNF	R&D system	Cat#212-GD	
peptide, recombinant protein	bFGF	MedChemExpress	Cat#HY-P7331	
peptide, recombinant protein	IGF-1	MedChemExpress	Cat#HY-P70783	
peptide, recombinant protein	TGF-β	MedChemExpress	Cat#HY-P70543	
chemical compound, drug	SPP-86	MedChemExpress	Cat#HY-110193	
chemical compound, drug	PP2	MedChemExpress	Cat#HY-13805	
chemical compound, drug	LY294002	MedChemExpress	Cat#HY-10108	
chemical compound, drug	U0126	MedChemExpress	Cat#S1102	
chemical compound, drug	SP600125	Selleck	Cat#HY-12041	

chemical compound, drug	SB203580	MedChemExpress	Cat#HY-10256	

Cell culture and viability assay

Rat brain microvascular endothelial cells were isolated from Sprague-Dawley rats (male, 7–10 days old, Sino-British Sippr/BKLaboratory Animal Ltd, Shanghai, China) as the described method (Bian-Sheng Ji et al., 2013; Ming-Shan L et al., 2013) and cultured in Dulbecco's Modified Eagle Media (DMEM)/F12 (#12500-039, Gibco, Carlsbad, CA, USA) containing 10% fetal bovine serum (FBS) (#10100147C, Gibco, Carlsbad, CA, USA) and 62.5 µg/mL penicillin and 100 µg/mL streptomycin (SunShine Biotechnology Co., ltd. Nanjing, China). Then, hCMEC/D3, U251, and SH-SY5Y cells were cultured in DMEM/F12 containing 10% FBS, 62.5 µg/mL penicillin and 100 µg/mL streptomycin. Cell viability was assessed using a CCK-8 kit (Beyotime Biotechnology, Shanghai, China), and the results were expressed as the fold of control.

Establishment of the triple co-culture model

Although hCMEC/D3 cells have poor barrier properties and low TEER compared to human physiological BBB, the use of human BMECs may be restricted by the acquisition of materials and ethical approval. Isolation and purification of primary BMECs are time-consuming and laborious. Moreover, culture conditions can alter transcriptional activity (Qi et al., 2023). All limit the establishment of BBB models based on primary human BMECs for high-throughput screening. Here, hCMEC/D3 cells were selected to establish an *in vitro* BBB model.

The establishment process of the triple co-culture model is illustrated in Figure 9. SH-SY5Y cells were seeded at a density of 4.5×10^4 cells/cm^2 in plates and

differentiated with 10 μM retinoic acid (Sigma-Aldrich, St. Louis, MO, USA) for 72 h. The differentiated SH-SY5Y cells were cultured in the fresh DMEM/F12 medium containing 10% FBS. U251 cells were seeded at 2×10^4 cells/cm^2 on the bottom of Transwell inserts (PET, 0.4 μm pore size, SPL Life Sciences, Pocheon, Korea) coated with rat-tail collagen (Corning Inc., Corning, NY, USA). Next, the inserts were suspended in plate wells containing the culture medium after 5 h of incubation. After 24 h of incubation, hCMEC/D3 cells were seeded on the apical side of the inserts at 3 $\times 10^4$ cells/cm^2 and cultured for another 48 h. Then, the inserts seeded with U251 and hCMEC/D3 cells were suspended in plates seeded with differentiated SH-SY5Y cells and co-cultured for another 6 days. The culture medium was replaced every 24 h. TEER was periodically measured using a Millicell-ERS (MERS00002) instrument (Millipore, Billerica MA, USA) to monitor cell confluence and development of tight junctions.

EdU incorporation assay

The cells were incubated with medium containing 10 μM Edu for 2 hours. Then cells were washed by PBS and harvested by 0.25% trypsin-ethylenediaminetetraacetic acid (#25200072, Gibco, Carlsbad, CA, USA). The EdU incorporation assay was measured using the BeyoClick™ EdU Cell Proliferation Kit (Beyotime Biotechnology, Shanghai, China) according to the manufacturer's instructions. The samples were determined on the FACSCelesta flow cytometer (Becton, Dickins on and Company, USA, and data were analyzed by Flowjo 10.4 software.

In vitro **BBB permeability study**

On day 7, the TEER values of BBB models showed a decreasing trend. Therefore, the subsequent experiments were all completed on day 6. The culture medium was removed from the apical and basolateral sides of the inserts and washed

twice with preheated Hank's balanced salt solution (HBSS). Fresh HBSS was then added to both the apical and basolateral chambers. After 15 min of preincubation, HBSS in the apical and basolateral chambers was replaced with HBSS containing FITC-dextran 3–5 kDa (FITC-Dex) (Sigma-Aldrich, St. Louis, MO, USA), fluorescein sodium (Sigma-Aldrich, St. Louis, MO, USA), or other tested agents and blank HBSS, respectively. Next, 200 µL aliquots were collected from the basolateral chamber after 30 min of incubation at 37 ℃. The concentrations of the tested agents in the basolateral chamber were measured.

The apparent permeability coefficient (P_{app}, cm/s) values of the tested agents across the *in vitro* BBB model were calculated using the equation (Tavelin et al., 2002):

$$P_{app}=(Q/1800)/(S \times C_0) \tag{1}$$

where S is the surface area of the insert membrane (0.33 cm^2 for 6.5 mm inserts, 4.46 cm^2 for 24 mm inserts), Q is the transported amount of the tested agents transported from the donor chamber to the receiver chamber for 30 min (1800 s), and C_0 is the initial concentration of the tested agents in the donor chamber.

The quantification methods of prazosin, verapamil, lamotrigine

Prazosin, verapamil, and lamotrigine (Aladdin, Shanghai, China) were analyzed by high-performance liquid chromatography (Shimadzu, Kyoto, Japan) with YMC-Triart C18 column (5 µm, 150 × 4.6 mm, YMC America Inc., Allentown, PA, USA). Prazosin and verapamil were detected using the RF-20A fluorescence detector. Lamotrigine was detected using the SPD-20A ultraviolet detector. Samples were centrifugated at 12000 rpm for 10 mins, then 150 µL supernatant was taken and used for analysis. The run temperature was set at 40 °C, the injection volume was 20 µL and the flow rate was 1 mL/min. Initial concentrations in donor chamber and other

chromatographic conditions of drugs are summarized in Table 2.

The quantification methods of clozapine, venlafaxine, bupropion,
amantadine, carbamazepine, fluoxetine, amitriptyline, gabapentin,
midazolam, risperidone, olanzapine, mirtazapine, metoclopramide,
doxepin, donepezil

All compounds were purchased from Aladdin (Shanghai, China). Except for
prazosin, verapamil, and lamotrigine, the other compounds were analyzed by using
liquid chromatography-mass spectrometry (Shimadzu, Kyoto, Japan) with
YMC-Triart C18 column (5 μm, 150 × 2.0 mm, YMC America Inc., Allentown, PA,
USA). Each sample was mixed with 10 μL internal standard. Then 1 mL extraction
was added to each sample. The samples were vortex vibrated on the oscillator for 10
min, and then centrifuged at 4 ℃ and 12000 rpm for 10 mins. The supernatant solvent
was evaporated with nitrogen flow, then redissolved with 100 μL 40 % (v/v)
acetonitrile, and centrifuged at 4 ℃ and 15000 rpm for 10 min. The supernatants were
injected into LC-MS for analysis. The injection volume of each sample was 5 μL. The
mass charge ratio, extraction, and initial concentrations in donor chamber of drugs
were summarized in Table 3.

Cell density analysis

On day 6 of co-culture, hCMEC/D3 cells were fixed with 4% paraformaldehyde
for 15 min and washed with phosphate buffered saline (PBS) for three times. Next,
the fixed cells were blocked with 5% goat serum for 2 h and washed with PBS for
four times. The blocked cells were incubated with DAPI (Invitrogen, Carlsbad, CA,
USA) and washed with PBS for four times. Cell numbers were counted using
Cytation5 (BioTek, Winooski, VT, USA).

Reverse transcription and Quantitative real-time PCR (qPCR)

Total RNA of cells was extracted using RNAiso Plus reagent (Takara Bio Inc.

Otsu, Shiga, Japan) and reverse transcribed using HiScript III RT SuperMix (Vazyme,

Shanghai, China) as the described method (Yang et al., 2023). Paired primers were

synthesized by Tsingke Biotech Co., Ltd (Beijing, China), and their sequences were

listed in Table 4. The SYBR Master Mix was purchased from Yeasen (Shanghai,

China). Then, qPCR was performed on the Applied Biosystems QuantStudio 3

real-time PCR system (Thermo Fisher Scientific, Waltham, MA, USA). The mRNA

levels of related genes were normalized to *ACTB* or *GAPDH* using the comparative

cycle threshold method.

Western blotting analysis

Whole-cell and tissue lysates, nucleoprotein, and cytoplasmic protein were

prepared using RIPA Lysis Buffer (Beyotime, Shanghai, China) as the described

method (Wu et al., 2021). Proteins were separated through sodium dodecyl

sulfate-polyacrylamide gel electrophoresis and transferred onto nitrocellulose or

polyvinylidene difluoride membranes. The membranes were blocked with 5% skim

milk and incubated with corresponding primary antibodies at 4 °C overnight. After

being washed with Tris-buffered saline Tween buffer, the membranes were incubated

with secondary antibodies (Cell Signaling Technology, MA, USA) at 1:3000 dilution:

Anti-mouse IgG, HRP-linked Antibody (#7076), Anti-rabbit IgG, HRP-linked

Antibody (#7074). Protein levels were visualized using a highly sensitive ECL

western blotting substrate and a gel imaging system (Tanon Science & Technology,

Shanghai, China).

Preparation of conditioned medium

Conditioned medium (CM) of U251 cells (U-CM), SH-SY5Y cells (S-CM), or

co-culture of U251 and SH-SY5Y cells (US-CM) were prepared. U251 cells were

seeded at the top of the insert membrane and suspended on 6-well plates seeded with

differentiated SH-SY5Y cells to co-culture U251 cells with SH-SY5Y cells. The

medium was collected every 24 h. The CMs were subsequently used for hCMEC/D3

cell culture after filtrating with 0.2 μm filters for 144 h. The levels of glia-derived

neurotrophic factor (GDNF), basic fibroblast growth factor (bFGF), insulin-like

growth factor-1 (IGF-1), and transforming growth factor-β (TGF-β) in CMs were

measured using corresponding ELISA kits according to the manufacturers'

instructions.

Neutralization of GDNF with Anti-GDNF Antibody

Exogenous and endogenous GDNF in the medium was neutralized with

anti-GDNF antibody (#AF-212-NA, R&D system, Minneapolis, MN, USA). 0.25, 0.5,

1.0 μg/mL anti-GDNF antibody was added into the US-CM or medium containing

200 pg/mL GDNF, and then the medium was preincubated at 4 °C for 1 h. The

hCMEC/D3 cells were incubated with 200 pg/mL GDNF or US-CM containing

anti-GDNF antibody or not for 6 days, and then cell lysate was collected for western

blot. The medium was replaced every 24 h.

Transfection of hCMEC/D3

Here, hCMEC/D3 cells were plated in the plates or culture dishes at 6×10^4

cells/cm^2 and transfected with 10 nM of negative control or human *FOXO1* and *ETS1*

small interfering RNA (siRNA) (Tsingke Biotechnology, Beijing, China) for 12 h

using Lipofectamine™ 3000 (Invitrogen, Carlsbad, CA, USA) reagent according to

the manufacturer's instructions. Cells were then incubated with a medium containing

GDNF for 72 h. The siRNA sequences of human *FOXO1* and *ETS1* were summarized

in Table 5.

FOXO1 Overexpression by plasmids

The plasmids encoding FOXO1 (EX-Z7404-M02) were constructed by

GeneCopoeia (Rockville, MD, USA). The hCMEC/D3 cells were plated in plates or

dishes at 6×10^4 cells/cm^2. They were subsequently transfected with 1 µg of negative

control or plasmids encoding FOXO1 6 h using Lipofectamine™ 3000 reagent

according to the manufacturer's instructions. Transfected cells were then incubated

with the medium containing GDNF for 72 h.

Animals

C57BL/6J mice (male, 4–5 weeks old, 16–18 g, 12 mice) were obtained from

Sino-British Sippr/BKLaboratory Animal Ltd (Shanghai, China). Mice were

maintained in groups under standard conditions with free access to food and water.

Animal studies were performed in accordance with the Guide for the Care and Use of

Laboratory Animals (National Institutes of Health) and approved by the Animal

Ethics Committee of China Pharmaceutical University (Approval Number:

202307003).

Brain-specific *Gdnf* knockdown and evaluation of BBB permeability

Mice were randomly divided into control (shNC) and *Gdnf* silencing (sh*Gdnf*)

groups (6 mice each group). The shRNA sequence of mice *Gdnf* was listed in Table 5.

The 2×10^9 viral genome each of pAAV-U6-shRNA (NC2)-CMV-EGFP or pAAV-

U6-shRNA (*Gdnf*)-CMV-EGFP (OBio Technology, Beijing, China) were injected into

the bilateral lateral ventricle area (relative to the bregma: anterior-posterior -0.3 mm;

medial-lateral ±1.0 mm; dorsal-ventral -3.0 mm) through intracerebroventricular (*i.c.v*)

infusion. Three weeks following *i.c.v* injection, BBB permeability and expression of

corresponding targeted proteins were measured in the mice.

A mixture of FITC-Dex (50 mg/kg) and fluorescein sodium (10 mg/kg) was

intravenously administered to experimental mice. Thirty minutes after the injection,

the mice were euthanized under isoflurane anesthesia, and brain tissue and plasma

samples were obtained quickly. The concentrations of FITC-Dex and fluorescein in

the plasma and brain were measured as previously described (Li P. et al., 2022; Zhou

Y. et al., 2019). No blinding was performed in animal studies.

The prediction of drug permeability across BBB using the developed in

vitro BBB model

The P_{app} values of 18 drugs – prazosin, verapamil, lamotrigine, clozapine,

venlafaxine, bupropion, amantadine, carbamazepine, fluoxetine, amitriptyline,

gabapentin, midazolam, risperidone, olanzapine, mirtazapine, metoclopramide,

doxepin, and donepezil – across the hCMEC/D3 cells mono-culture and triple

co-culture models were measured. The predicted *in vivo* permeability-surface area

product (*PS*, μL/min/g brain) values across BBB were calculated using the following

equation:

689
$$PS_{Pre} = \frac{P_{app} \times 60 \times VSA \times 1000}{f_{u,brain}}$$
(2)

where VSA is the luminal area of the vascular space of brain, which was set to 150

cm^2/g (Fenstermacher et al., 1988), and $f_{u, brain}$ is the unbound fraction of brain. The

published *in vivo* brain permeability values were unified to observed *PS* (*PS$_{Obs}$*) by

multiplying by VSA equal to 150 cm^2/g. If *PS$_{Pre}$* values were within 0.5–2.0 folds of

observations, the prediction was considered successful.

Statistical analyses

All results are presented as mean ± SEM. The average of technical replicates

generated a single independent value that contributes to the n value used for

comparative statistical analysis. The data were assessed for Gaussian distributions

using Shapiro-Wilk test. Brown-Forsythe test was employed to evaluate the

homogeneity of variance between groups. For comparisons between two groups, statistical significance was determined by unpaired 2-tailed t-test. The acquired data with significant variation were tested using unpaired t-test with Welch's correction, and non-Gaussian distributed data were tested using Mann-Whitney test. For multiple group comparisons, one-way ANOVA followed by Fisher's LSD test was used to determine statistical significance. The acquired data with significant variation were tested using Welch's ANOVA test, and non-Gaussian distributed data were tested using Kruskal-Wallis test. $P < 0.05$ was considered statistically significant. The simple linear regression analysis was used to examine the presence of a linear relationship between two variables. Data were analyzed using GraphPad Prism software version 8.0.2 (GraphPad Software, La Jolla, CA, USA).

Acknowledgments

The authors would like to give special thanks to the support of the China Pharmaceutical University Pharmaceutical Animal Laboratory Center.

Funding information

Funder	Grant reference number	Author
National Natural Science Foundation of China	82373943	Xiaodong Liu
National Natural Science Foundation of China	82173884	Li Liu
National Natural Science Foundation of China	82204511	Hanyu Yang
"Double First-Class" university project	No. CPU2022QZ21	Li Liu
Jiangsu Funding Program for Excellent Postdoctoral Talent	No. 2022ZB305	Hanyu Yang
China Postdoctoral Science Foundation	2023M733897	Hanyu Yang
Postgraduate Research & Practice Innovation Program of Jiangsu Province	KYCX24_1019	Lu Yang
The funding sources were involved in study design, data collection, and interpretation, or the decision to submit the work for publication.		

Author Contributions

Lu Yang: Conceptualization, Methodology, Investigation, Data curation, and

Writing–original draft preparation, Funding acquisition; **Zijin Lin:** Data curation,

Formal analysis; **Ruijing Mu**: Visualisation, Software; **Wenhan Wu**: Formal analysis;

Hao Zhi: Data curation; **Xiaodong Liu**: Supervision, Conceptualization, Project

administration, Funding acquisition; **Hanyu Yang**: Supervision, Conceptualization,

Funding acquisition; **Li Liu**: Supervision, Conceptualization, Project administration,

Funding acquisition, Resources. All authors reviewed and approved the final version

of this manuscript.

Declaration of interests

The authors declare no competing interests.

Data availability

All data are available in the manuscript and supporting files; source data files for

western blots have been provided for all figures.

References

Abbott N J. (2013). Blood-brain barrier structure and function and the challenges for

CNS drug delivery. *J Inherit Metab Dis, 36*(3), 437-449.

https://doi.org/10.1007/s10545-013-9608-0

Arvanitis C D, Ferraro G B, Jain R K. (2020). The blood-brain barrier and

blood-tumour barrier in brain tumours and metastases. *Nat Rev Cancer, 20*(1), 26-41.

https://doi.org/10.1038/s41568-019-0205-x

Asada S, Daitoku H, Matsuzaki H, Saito T, Sudo T, Mukai H, Iwashita S, Kako K,

Kishi T, Kasuya Y, Fukamizu A. (2007). Mitogen-activated protein kinases, Erk and

p38, phosphorylate and regulate Foxo1. *Cell Signal, 19*(3), 519-527.

https://doi.org/10.1016/j.cellsig.2006.08.015

Avdeef A, Sun N. (2010). A New In Situ Brain Perfusion Flow Correction Method for

Lipophilic Drugs Based on the pH-Dependent Crone-Renkin Equation.

Pharmaceutical Research, 28(3), 517-530. https://doi.org/10.1007/s11095-010-0298-0

Banks W A. (2016). From blood-brain barrier to blood-brain interface: new

opportunities for CNS drug delivery. *Nat Rev Drug Discov, 15*(4), 275-292.

https://doi.org/10.1038/nrd.2015.21

Barberio C, Withers A, Mishra Y, Couraud P-O, Romero I A, Weksler B, Owens R M.

(2022). A human-derived neurovascular unit in vitro model to study the effects of

cellular cross-talk and soluble factors on barrier integrity. *Frontiers in Cellular*

Neuroscience, 16. https://doi.org/10.3389/fncel.2022.1065193

Beard R S, Jr., Hoettels B A, Meegan J E, Wertz T S, Cha B J, Yang X, Oxford J T,

Wu M H, Yuan S Y. (2020). AKT2 maintains brain endothelial claudin-5 expression

and selective activation of IR/AKT2/FOXO1-signaling reverses barrier dysfunction. *J*

Cereb   Blood   Flow   Metab,   40(2),   374-391.

https://doi.org/10.1177/0271678X18817512

Bhallamudi S, Roos B B, Teske J J, Wicher S A, McConico A, C M P, Sathish V,

Prakash Y S. (2021). Glial-derived neurotrophic factor in human airway smooth

muscle. *J Cell Physiol, 236*(12), 8184-8196. https://doi.org/10.1002/jcp.30489

Bhattacharya C, Masters A R, Bach C, Stratford R E, Jr. (2021). Population model

analysis of chiral inversion and degradation of bupropion enantiomers, and

application to enantiomer specific fraction unbound determination in rat plasma and

brain. *J Pharm Biomed Anal, 195*, 113872. https://doi.org/10.1016/j.jpba.2020.113872

Bhyrapuneni G, Thentu J B, Palacharla V R C, Muddana N, Aleti R R, Ajjala D R,

Nirogi R. (2018). A definite measure of occupancy exposures, seeking with

non-radiolabeled in vivo 5-HT2A receptor occupancy and in vitro free fractions. *J*

Recept Signal Transduct Res, 38(4), 359-366.

https://doi.org/10.1080/10799893.2018.1531888

Bian-Sheng Ji, Juan Cen, Ling He, Meng Liu, Yan-Qing Liu, Liu L. (2013).

Modulation of P-glycoprotein in rat brain microvessel endothelial cells under oxygen

glucose deprivation. *J Pharm Pharmacol, 65*(10), 1508-1517.

https://doi.org/10.1111/jphp.12122

Biswas S, Cottarelli A, Agalliu D. (2020). Neuronal and glial regulation of CNS

angiogenesis and barriergenesis. *Development, 147*(9).

https://doi.org/10.1242/dev.182279

Bowman C M, Benet L Z. (2018). An examination of protein binding and

protein-facilitated uptake relating to in vitro-in vivo extrapolation. *Eur J Pharm Sci,*

123, 502-514. https://doi.org/10.1016/j.ejps.2018.08.008

Bulldan A, Dietze R, Shihan M, Scheiner-Bobis G. (2016). Non-classical testosterone

signaling mediated through ZIP9 stimulates claudin expression and tight junction

formation in Sertoli cells. *Cellular Signalling, 28*(8), 1075-1085.

https://doi.org/10.1016/j.cellsig.2016.04.015

Clasadonte J, Prevot V. (2017). The special relationship: glia–neuron interactions in

the neuroendocrine hypothalamus. *Nature Reviews Endocrinology, 14*(1), 25-44.

https://doi.org/10.1038/nrendo.2017.124

Colas-Algora N, Garcia-Weber D, Cacho-Navas C, Barroso S, Caballero A, Ribas C,

Correas I, Millan J. (2020). Compensatory increase of VE-cadherin expression

through ETS1 regulates endothelial barrier function in response to TNFalpha. *Cell*

Mol Life Sci, 77(11), 2125-2140. https://doi.org/10.1007/s00018-019-03260-9

Cremers T I, Flik G, Hofland C, Stratford R E, Jr. (2012). Microdialysis evaluation of

clozapine and N-desmethylclozapine pharmacokinetics in rat brain. *Drug Metab*

Dispos, 40(10), 1909-1916. https://doi.org/10.1124/dmd.112.045682

Dejana E, Orsenigo F, Lampugnani M G. (2008). The role of adherens junctions and

VE-cadherin in the control of vascular permeability. *J Cell Sci, 121*(Pt 13), 2115-2122.

https://doi.org/10.1242/jcs.017897

Di L, Umland J P, Chang G, Huang Y, Lin Z, Scott D O, Troutman M D, Liston T E.

(2011). Species independence in brain tissue binding using brain homogenates. *Drug*

Metab Dispos, 39(7), 1270-1277. https://doi.org/10.1124/dmd.111.038778

Di L, Breen C, Chambers R, Eckley S T, Fricke R, Ghosh A, Harradine P, Kalvass J C,

Ho S, Lee C A, Marathe P, Perkins E J, Qian M, Tse S, Yan Z, Zamek-Gliszczynski M

J. (2017). Industry Perspective on Contemporary Protein-Binding Methodologies:

Considerations for Regulatory Drug-Drug Interaction and Related Guidelines on

Highly   Bound   Drugs.   *J   Pharm   Sci,   106*(12),   3442-3452.

https://doi.org/10.1016/j.xphs.2017.09.005

Di Marco A, Gonzalez Paz O, Fini I, Vignone D, Cellucci A, Battista M R, Auciello G,

Orsatti L, Zini M, Monteagudo E, Khetarpal V, Rose M, Dominguez C, Herbst T,

Toledo-Sherman L, Summa V, Munoz-Sanjuan I. (2019). Application of an in Vitro

Blood-Brain Barrier Model in the Selection of Experimental Drug Candidates for the

Treatment of Huntington's Disease. *Mol Pharm, 16*(5), 2069-2082.

https://doi.org/10.1021/acs.molpharmaceut.9b00042

Dong C, Ubogu E E. (2018). GDNF enhances human blood-nerve barrier function in

vitro via MAPK signaling pathways. *Tissue Barriers, 6*(4), 1-22.

https://doi.org/10.1080/21688370.2018.1546537

Eigenmann D E, Xue G, Kim K S, Moses A V, Hamburger M, Oufir M. (2013).

Comparative study of four immortalized human brain capillary endothelial cell lines,

hCMEC/D3, hBMEC, TY10, and BB19, and optimization of culture conditions, for

an in vitro blood–brain barrier model for drug permeability studies. *Fluids Barriers*

CNS, 10(1), 33. https://doi.org/10.1186/2045-8118-10-33

Esaki T, Ohashi R, Watanabe R, Natsume-Kitatani Y, Kawashima H, Nagao C,

Mizuguchi K. (2019). Computational Model To Predict the Fraction of Unbound Drug

in the Brain. *J Chem Inf Model, 59*(7), 3251-3261.

https://doi.org/10.1021/acs.jcim.9b00180

Fenstermacher J, Gross P, Sposito N, Acuff V, Pettersen S, Gruber K. (1988).

Structural and Functional Variations in Capillary Systems within the Brain. *Ann N Y*

Acad Sci, 529, 21-30. https://doi.org/10.1111/j.1749-6632.1988.tb51416.x

Fielder G C, Yang T W, Razdan M, Li Y, Lu J, Perry J K, Lobie P E, Liu D X. (2018).

The GDNF Family: A Role in Cancer? *Neoplasia, 20*(1), 99-117.

https://doi.org/10.1016/j.neo.2017.10.010

Friden M, Bergstrom F, Wan H, Rehngren M, Ahlin G, Hammarlund-Udenaes M,

Bredberg U. (2011). Measurement of unbound drug exposure in brain: modeling of

pH partitioning explains diverging results between the brain slice and brain

homogenate methods. *Drug Metab Dispos, 39*(3), 353-362.

https://doi.org/10.1124/dmd.110.035998

Fu J, Li L, Huo D, Zhi S, Yang R, Yang B, Xu B, Zhang T, Dai M, Tan C, Chen H,

Wang X. (2021). Astrocyte-Derived TGFβ1 Facilitates Blood–Brain Barrier Function

via Non-Canonical Hedgehog Signaling in Brain Microvascular Endothelial Cells.

Brain Sciences, 11(1). https://doi.org/10.3390/brainsci11010077

Hanafy A S, Dietrich D, Fricker G, Lamprecht A. (2021). Blood-brain barrier models:

Rationale for selection. *Adv Drug Deliv Rev, 176*, 113859.

https://doi.org/10.1016/j.addr.2021.113859

Hashimoto Y, Greene C, Munnich A, Campbell M. (2023). The CLDN5 gene at the

blood-brain barrier in health and disease. *Fluids and Barriers of the CNS, 20*(1).

https://doi.org/10.1186/s12987-023-00424-5

Hatherell K, Couraud P O, Romero I A, Weksler B, Pilkington G J. (2011).

Development of a three-dimensional, all-human in vitro model of the blood-brain

barrier using mono-, co-, and tri-cultivation Transwell models. *J Neurosci Methods,*

199(2), 223-229. https://doi.org/10.1016/j.jneumeth.2011.05.012

He F, Wang Q F, Li L, Yu C, Liu C Z, Wei W C, Chen L P, Li H Y. (2023). Melatonin

Protects Against Hyperoxia-Induced Apoptosis in Alveolar Epithelial type II Cells by

Activating the MT2/PI3K/AKT/ETS1 Signaling Pathway. *Lung, 201*(2), 225-234.

https://doi.org/10.1007/s00408-023-00610-0

Hidalgo-Figueroa M, Bonilla S, Gutierrez F, Pascual A, Lopez-Barneo J. (2012).

GDNF is predominantly expressed in the PV+ neostriatal interneuronal ensemble in

normal mouse and after injury of the nigrostriatal pathway. *J Neurosci, 32*(3), 864-872.

https://doi.org/10.1523/JNEUROSCI.2693-11.2012

Hui K, Wu S, Yue Y, Gu Y, Guan B, Wang X, Hsieh J T, Chang L S, He D, Wu K.

(2018). RASAL2 inhibits tumor angiogenesis via p-AKT/ETS1 signaling in bladder

cancer. *Cell Signal, 48*, 38-44. https://doi.org/10.1016/j.cellsig.2018.04.006

Igarashi Y, Utsumi H, Chiba H, Yamada-Sasamori Y, Tobioka H, Kamimura Y,

Furuuchi K, Kokai Y, Nakagawa T, Mori M, Sawada N. (1999). Glial cell line-derived

neurotrophic factor induces barrier function of endothelial cells forming the

blood-brain barrier. *Biochem Biophys Res Commun, 261*(1), 108-112.

https://doi.org/10.1006/bbrc.1999.0992

Ito R, Umehara K, Suzuki S, Kitamura K, Nunoya K I, Yamaura Y, Imawaka H, Izumi

S, Wakayama N, Komori T, Anzai N, Akita H, Furihata T. (2019). A Human

Immortalized Cell-Based Blood-Brain Barrier Triculture Model: Development and

Characterization as a Promising Tool for Drug-Brain Permeability Studies. *Mol

Pharm, 16*(11), 4461-4471. https://doi.org/10.1021/acs.molpharmaceut.9b00519

Ji-Ae K, RyojiI Y, Teruo N. (2009). IGF-1 released by corneal epithelial cells induces

up-regulation of N-cadherin in corneal fibroblasts. *J Cell Physiol, 221*(1), 254-261.

https://doi.org/10.1002/jcp.21850

Kodaira H, Kusuhara H, Fujita T, Ushiki J, Fuse E, Sugiyama Y. (2011). Quantitative

evaluation of the impact of active efflux by p-glycoprotein and breast cancer

resistance protein at the blood-brain barrier on the predictability of the unbound

concentrations of drugs in the brain using cerebrospinal fluid concentration as a

surrogate. *J Pharmacol Exp Ther, 339*(3), 935-944.

https://doi.org/10.1124/jpet.111.180398

Lelie`vre E, Mattot V, Huber P, Vandenbunde B, Soncin F. (2000). ETS1 lowers

capillary endothelial cell density at confluence and induces the expression of

VE-cadherin. *Oncogene, 19*(20), 2438-2446. https://doi.org/10.1038/sj.onc.1203563

Lelie`vre E, Lionneton F, Soncin F, Vandenbunder B. (2001). The Ets family contains

transcriptional activators and repressors involved in angiogenesis. *Int J Biochem Cell*

Biol, 33(4), 391-407. https://doi.org/10.1016/s1357-2725(01)00025-5

Li P, Yang Y, Lin Z, Hong S, Jiang L, Zhou H, Yang L, Zhu L, Liu X, Liu L. (2022).

Bile Duct Ligation Impairs Function and Expression of Mrp1 at Rat Blood-Retinal

Barrier via Bilirubin-Induced P38 MAPK Pathway Activations. *Int J Mol Sci, 23*(14).

https://doi.org/10.3390/ijms23147666

Li W, Chen Z, Chin I, Chen Z, Dai H. (2018). The Role of VE-cadherin in

Blood-brain Barrier Integrity Under Central Nervous System Pathological Conditions.

Current Neuropharmacology, 16(9), 1375-1384.

https://doi.org/10.2174/1570159x16666180222164809

Li Y, Li Y, Zhang Y, Zhao Q, Zhang P, Sun M, Liu B, Yang H, Li P. (2022). Muscone

and (+)-Borneol Cooperatively Strengthen CREB Induction of Claudin 5 in

IL-1β-Induced Endothelium Injury. *Antioxidants, 11*(8).

https://doi.org/10.3390/antiox11081455

Lin C H, Wang C H, Hsu S L, Liao L Y, Lin T A, Hsueh C M. (2016). Molecular

Mechanisms Responsible for Neuron-Derived Conditioned Medium (NCM)-Mediated

Protection of Ischemic Brain. *PLoS One, 11*(1), e0146692.

https://doi.org/10.1371/journal.pone.0146692

Liu D, Yang L, Liu P, Ji X, Qi X, Wang Z, Chi T, Zou L. (2022). Sigma-1 receptor

activation alleviates blood-brain barrier disruption post cerebral ischemia stroke by

stimulating the GDNF-GFRalpha1-RET pathway. *Exp Neurol, 347*, 113867.

https://doi.org/10.1016/j.expneurol.2021.113867

Liu X, Smith B J, Chen C, Callegari E, Becker S L, Chen X, Cianfrogna J, Doran A C,

Doran S D, Gibbs J P, Hosea N, Liu J, Nelson F R, Szewc M A, Van Deusen J. (2005).

Use of a physiologically based pharmacokinetic model to study the time to reach

brain equilibrium: an experimental analysis of the role of blood-brain barrier

permeability, plasma protein binding, and brain tissue binding. *J Pharmacol Exp Ther,*

313(3), 1254-1262. https://doi.org/10.1124/jpet.104.079319

Liu Y, Mu S, Li X, Liang Y, Wang L, Ma X. (2019). Unfractionated Heparin

Alleviates Sepsis-Induced Acute Lung Injury by Protecting Tight Junctions. *Journal*

of Surgical Research, 238, 175-185. https://doi.org/10.1016/j.jss.2019.01.020

Lonka-Nevalaita L, Lume M, Leppänen S, Jokitalo E, Peränen J, Saarma M. (2010).

Characterization of the Intracellular Localization, Processing, and Secretion of Two

Glial Cell Line-Derived Neurotrophic Factor Splice Isoforms. *The Journal of*

Neuroscience, 30(34), 11403-11413. https://doi.org/10.1523/jneurosci.5888-09.2010

Luo Y, Yang H, Wan Y, Yang S, Wu J, Chen S, Li Y, Jin H, He Q, Zhu D Y, Zhou Y,

Hu B. (2022). Endothelial ETS1 inhibition exacerbate blood-brain barrier dysfunction

in multiple sclerosis through inducing endothelial-to-mesenchymal transition. *Cell*

Death Dis, 13(5), 462. https://doi.org/10.1038/s41419-022-04888-5

Mathiisen T M, Lehre K P, Danbolt N C, Ottersen O P. (2010). The perivascular

astroglial sheath provides a complete covering of the brain microvessels: an electron

microscopic 3D reconstruction. *Glia, 58*(9), 1094-1103.

https://doi.org/10.1002/glia.20990

Maurer T S, Debartolo D B, Tess D A, Scott D O. (2005). Relationship between

exposure and nonspecific binding of thirty-three central nervous system drugs in mice.

Drug Metab Dispos, 33(1), 175-181. https://doi.org/10.1124/dmd.104.001222

Ming-Shan L, Juan C, Ling H, Lu L, Bian-Sheng J. (2013). CJY, an isoflavone,

interacts with ATPase of P-glycoprotein in the rat brain microvessel endothelial cells

(RBMECs). *J Chemother, 25*(6), 347-354.

https://doi.org/10.1179/1973947813Y.0000000094

Morita A, Yamashita N, Sasaki Y, Uchida Y, Nakajima O, Nakamura F, Yagi T,

Taniguchi M, Usui H, Katoh-Semba R, Takei K, Goshima Y. (2006). Regulation of

Dendritic Branching and Spine Maturation by Semaphorin3A-Fyn Signaling. *The*

Journal of Neuroscience, 26(11), 2971-2980.

https://doi.org/10.1523/jneurosci.5453-05.2006

Muoio V, Persson P B, Sendeski M M. (2014). The neurovascular unit - concept

review. *Acta Physiol, 210*(4), 790-798. https://doi.org/10.1111/apha.12250

Nakagawa S, Deli M A, Kawaguchi H, Shimizudani T, Shimono T, Kittel A, Tanaka K,

Niwa M. (2009). A new blood-brain barrier model using primary rat brain endothelial

cells, pericytes and astrocytes. *Neurochem Int, 54*(3-4), 253-263.

https://doi.org/10.1016/j.neuint.2008.12.002

Nitta T, Hata M, Gotoh S, Seo Y, Sasaki H, Hashimoto N, Furuse M, Tsukita S. (2003).

Size-selective loosening of the blood-brain barrier in claudin-5-deficient mice. *J Cell*

Biol, 161(3), 653-660. https://doi.org/10.1083/jcb.200302070

Nowrangi D S, McBride D, Manaenko A, Dixon B, Tang J, Zhang J H. (2019).

rhIGF-1 reduces the permeability of the blood-brain barrier following intracerebral

hemorrhage in mice. *Exp Neurol, 312*, 72-81.

https://doi.org/10.1016/j.expneurol.2018.11.009

Ohtsuki S, Sato S, Yamaguchi H, Kamoi M, Asashima T, Terasaki T. (2007).

Exogenous expression of claudin-5 induces barrier properties in cultured rat brain

capillary endothelial cells. *J Cell Physiol, 210*(1), 81-86.

https://doi.org/10.1002/jcp.20823

Ohtsuki S, Ikeda C, Uchida Y, Sakamoto Y, Miller F, Glacial F, Decleves X,

Scherrmann J M, Couraud P O, Kubo Y, Tachikawa M, Terasaki T. (2013).

Quantitative targeted absolute proteomic analysis of transporters, receptors and

junction proteins for validation of human cerebral microvascular endothelial cell line

hCMEC/D3 as a human blood-brain barrier model. *Mol Pharm, 10*(1), 289-296.

https://doi.org/10.1021/mp3004308

Palmiotti C A, Prasad S, Naik P, Abul K M, Sajja R K, Achyuta A H, Cucullo L.

(2014). In vitro cerebrovascular modeling in the 21st century: current and prospective

technologies. *Pharm Res, 31*(12), 3229-3250.

https://doi.org/10.1007/s11095-014-1464-6

Pochon N A, Menoud A, Tseng J L, Zurn A D, Aebischer P. (1997). Neuronal GDNF

Expression in the Adult Rat Nervous System Identified By In Sifu Hybridization.

European Journal of Neuroscience, 9(3), 463-471.

https://doi.org/10.1111/j.1460-9568.1997.tb01623.x

Potjewyd G, Moxon S, Wang T, Domingos M, Hooper N M. (2018). Tissue

Engineering 3D Neurovascular Units: A Biomaterials and Bioprinting Perspective.

Trends Biotechnol, 36(4), 457-472. https://doi.org/10.1016/j.tibtech.2018.01.003

Qi D, Lin H, Hu B, Wei Y. (2023). A review on in vitro model of the blood-brain

barrier (BBB) based on hCMEC/D3 cells. *J Control Release, 358*, 78-97.

https://doi.org/10.1016/j.jconrel.2023.04.020

Sajja R K, Prasad S, Cucullo L. (2014). Impact of altered glycaemia on blood-brain

barrier endothelium: an in vitro study using the hCMEC/D3 cell line. *Fluids Barriers*

CNS, 11(1), 8. https://doi.org/10.1186/2045-8118-11-8

Saker S, Stewart E A, Browning A C, Allen C L, Amoaku W M. (2014). The effect of

hyperglycaemia on permeability and the expression of junctional complex molecules

in human retinal and choroidal endothelial cells. *Exp Eye Res, 121*, 161-167.

https://doi.org/10.1016/j.exer.2014.02.016

Sanchez-Dengra B, Gonzalez-Alvarez I, Sousa F, Bermejo M, Gonzalez-Alvarez M,

Sarmento B. (2021). In vitro model for predicting the access and distribution of drugs

in the brain using hCMEC/D3 cells. *Eur J Pharm Biopharm, 163*, 120-126.

https://doi.org/10.1016/j.ejpb.2021.04.002

Sauteur L, Affolter M, Belting H-G. (2017). Distinct and redundant functions of Esam

and   VE-cadherin   during   vascular   morphogenesis.   *Development*.

https://doi.org/10.1242/dev.140038

Savettieri G, Di Liegro I, Catania C, Licata L, Pitarresi G L, D'Agostino S, Gabriella

S, De Caro V, Giandalia G, Giannola L I, Cestelli A. (2000). Neurons and ECM

regulate occludin localization in brain endothelial cells. *Neuroreport, 11*(5),

1081-1084. https://doi.org/10.1097/00001756-200004070-00035

Schiera G, Bono E, Raffa M P, Gallo A, Pitarresi G L, Di Liegro I, Savettieri G.

(2003). Synergistic effects of neurons and astrocytes on the differentiation of brain

capillary endothelial cells in culture. *J Cell Mol Med, 7*(2), 165-170.

https://doi.org/10.1111/j.1582-4934.2003.tb00215.x

Schiera G, Sala S, Gallo A, Raffa M P, Pitarresi G L, Savettieri G, Di Liegro I. (2005).

Permeability properties of a three-cell type in vitro model of blood-brain barrier.

Journal of Cellular and Molecular Medicine, 9(2), 373-379.

https://doi.org/10.1111/j.1582-4934.2005.tb00362.x

Shimizu F, Sano Y, Abe M A, Maeda T, Ohtsuki S, Terasaki T, Kanda T. (2011).

Peripheral nerve pericytes modify the blood-nerve barrier function and tight

junctional molecules through the secretion of various soluble factors. *J Cell Physiol,*

226(1), 255-266. https://doi.org/10.1002/jcp.22337

Shimizu F, Sano Y, Saito K, Abe M A, Maeda T, Haruki H, Kanda T. (2012).

Pericyte-derived glial cell line-derived neurotrophic factor increase the expression of

claudin-5 in the blood-brain barrier and the blood-nerve barrier. *Neurochem Res,*

37(2), 401-409. https://doi.org/10.1007/s11064-011-0626-8

Summerfield S G, Read K, Begley D J, Obradovic T, Hidalgo I J, Coggon S, Lewis A

V, Porter R A, Jeffrey P. (2007). Central nervous system drug disposition: the

relationship between in situ brain permeability and brain free fraction. *J Pharmacol*

Exp Ther, 322(1), 205-213. https://doi.org/10.1124/jpet.107.121525

Sweeney M D, Zhao Z, Montagne A, Nelson A R, Zlokovic B V. (2019). Blood-Brain

Barrier: From Physiology to Disease and Back. *Physiol Rev, 99*(1), 21-78.

https://doi.org/10.1152/physrev.00050.2017

Taddei A, Giampietro C, Conti A, Orsenigo F, Breviario F, Pirazzoli V, Potente M,

Daly C, Dimmeler S, Dejana E. (2008). Endothelial adherens junctions control tight

junctions by VE-cadherin-mediated upregulation of claudin-5. *Nat Cell Biol, 10*(8),

923-934. https://doi.org/10.1038/ncb1752

Tang E D, Nuñez G, Barr F G, Guan K-L. (1999). Negative Regulation of the

Forkhead Transcription Factor FKHR by Akt. *Journal of Biological Chemistry,*

274(24), 16741-16746. https://doi.org/10.1074/jbc.274.24.16741

Tavelin S, Gråsjö J, Taipalensuu J, Ocklind G, Artursson P. (2002). Applications of

Epithelial Cell Culture in Studies of Drug Transport. In C. Wise (Ed.), *Epithelial Cell*

Culture Protocols (pp. 233-272). Totowa, NJ: Humana Press.

https://doi.org/10.1385/1-59259-185-x:233

Tunggal J A, Helfrich I, Schmitz A, Schwarz H, Günzel D, Fromm M, Kemler R,

Krieg T, Niessen C M. (2005). E-cadherin is essential for in vivo epidermal barrier

function by regulating tight junctions. *The EMBO Journal, 24*(6), 1146-1156.

https://doi.org/10.1038/sj.emboj.7600605

Wang Z G, Cheng Y, Yu X C, Ye L B, Xia Q H, Johnson N R, Wei X, Chen D Q, Cao

G, Fu X B, Li X K, Zhang H Y, Xiao J. (2016). bFGF Protects Against Blood-Brain

Barrier Damage Through Junction Protein Regulation via PI3K-Akt-Rac1 Pathway

Following   Traumatic   Brain   Injury.   *Mol   Neurobiol,   53*(10),   7298-7311.

https://doi.org/10.1007/s12035-015-9583-6

Watanabe D, Takagi H, Suzuma K, Suzuma I, Oh H, Ohashi H, Kemmochi S, Uemura

A, Ojima T, Suganami E, Miyamoto N, Sato Y, Honda Y. (2004). Transcription factor

Ets-1 mediates ischemia- and vascular endothelial growth factor-dependent retinal

neovascularization.   *Am   J   Pathol,   164*(5),   1827-1835.

https://doi.org/10.1016/S0002-9440(10)63741-8

Watanabe D, Nakagawa S, Morofuji Y, Toth A E, Vastag M, Aruga J, Niwa M, Deli M

A. (2021). Characterization of a Primate Blood-Brain Barrier Co-Culture Model

Prepared   from   Primary   Brain   Endothelial   Cells,   Pericytes   and   Astrocytes.

Pharmaceutics, 13(9). https://doi.org/10.3390/pharmaceutics13091484

Weber M L, Hofland C M, Shaffer C L, Flik G, Cremers T, Hurst R S, Rollema H.

(2013). Therapeutic doses of antidepressants are projected not to inhibit human

alpha4beta2 nicotinic acetylcholine receptors. *Neuropharmacology, 72*, 88-95.

https://doi.org/10.1016/j.neuropharm.2013.04.027

Weksler B, Romero I A, Couraud P-O. (2013). The hCMEC/D3 cell line as a model of

the human blood brain barrier. *Fluids and Barriers of the CNS, 10*(1), 16.

https://doi.org/10.1186/2045-8118-10-16

Wu T, Sheng Y, Qin Y Y, Kong W M, Jin M M, Yang H Y, Zheng X K, Dai C, Liu M,

Liu X D, Liu L. (2021). Bile duct ligation causes opposite impacts on the expression

and function of BCRP and P-gp in rat brain partly via affecting membrane expression

of ezrin/radixin/moesin proteins. *Acta Pharmacol Sin, 42*(11), 1942-1950.

https://doi.org/10.1038/s41401-020-00602-3

Xue Q, Liu Y, Qi H, Ma Q, Xu L, Chen W, Chen G, Xu X. (2013). A Novel Brain

Neurovascular Unit Model with Neurons, Astrocytes and Microvascular Endothelial

Cells of Rat. *International Journal of Biological Sciences, 9*(2), 174-189.

https://doi.org/10.7150/ijbs.5115

Yang H, Su M, Liu M, Sheng Y, Zhu L, Yang L, Mu R, Zou J, Liu X, Liu L. (2023).

Hepatic retinaldehyde deficiency is involved in diabetes deterioration by enhancing

PCK1- and G6PC-mediated gluconeogenesis. *Acta Pharm Sin B, 13*(9), 3728-3743.

https://doi.org/10.1016/j.apsb.2023.06.014

Zhang X, Tang N, Hadden T J, Rishi A K. (2011). Akt, FoxO and regulation of

apoptosis. *Biochimica et Biophysica Acta, 1813*(11), 1978-1986.

https://doi.org/10.1016/j.bbamcr.2011.03.010

Zhou L, Schmidt K, Nelson F R, Zelesky V, Troutman M D, Feng B. (2009). The

effect of breast cancer resistance protein and P-glycoprotein on the brain penetration

of flavopiridol, imatinib mesylate (Gleevec), prazosin, and

2-methoxy-3-(4-(2-(5-methyl-2-phenyloxazol-4-yl)ethoxy)phenyl)propanoic acid

(PF-407288) in mice. *Drug Metab Dispos, 37*(5), 946-955.

https://doi.org/10.1124/dmd.108.024489

Zhou Y, Zhou J, Li P, Xie Q, Sun B, Li Y, Chen Y, Zhao K, Yang T, Zhu L, Xu J, Liu

X, Liu L. (2019). Increase in P-glycoprotein levels in the blood-brain barrier of partial

portal vein ligation /chronic hyperammonemia rats is medicated by ammonia/reactive

oxygen species/ERK1/2 activation: In vitro and in vivo studies. *Eur J Pharmacol, 846*,

119-127. https://doi.org/10.1016/j.ejphar.2019.01.005

____

1077

Figure 1. The effects of co-culture with U251 and/or SH-SY5Y cells on the integrity of hCMEC/D3 and BBB function.

(**A**): Four different types of BBB models were prepared from hCMEC/D3 cells (h), SH-SY5Y cells (S), and U251 cells (U). (**B**): The transendothelial electrical resistance (TEER) of four models, and the TEER values in day 6 were compared. Blank: no cells. Four biological replicates per group. (**C**): The TEER of hCMEC/D3 and U251 cells monolayer. Four biological replicates per group. (**D** and **E**): The apparent permeability coefficient (P_{app}, $\times$ 10^{-6} cm/s) of fluorescein (NaF) and FITC-Dextran 3–5 kDa (FITC-Dex) of four BBB models. Four biological replicates per group. (**F**): The P_{app} ($\times$ 10^{-6} cm/s) of NaF and FITC-Dex across the blank inserts, and hCMEC/D3 or U251 mono-culture models. Four biological replicates

per group. (**G** and **H**): The cell density (G), EdU incorporation (H) of hCMEC/D3 cells after

mono/co-culturing. Three biological replicates per group. (**I**): Cell viability of hCMEC/D3

cells after mono/co-culturing. Four biological replicates per group. (**J** and **K**): The mRNA

levels of tight junction proteins, adherent junction proteins, and transporters. Four biological

replicates per group. (**L-O**): The protein expression levels of claudin-5 (CLDN-5), ZO-1,

occluding (L and M), VE-cadherin (VE-Cad), β-catenin, and BCRP (N and O) in hCMEC/D3

cells. Four biological replicates per group. (**P** and **Q**): The correlations between the P_{app} ($\times$

10^{-5} cm/s) of NaF and claudin-5 expression (P), or VE-cadherin expression (Q). (**R** and **S**):

The correlation between P_{app} ($\times$ 10^{-5} cm/s) of FITC-Dex and claudin-5 expression (R), or

VE-cadherin expression (S). The above data are shown as the mean $\pm$ SEM. For J and K, two

technical replicates per biological replicate. One technical replicate per biological replicate

for the rest. * $p < 0.05$; ** $p < 0.01$ by one-way ANOVA test followed by Fisher's LSD test,

Welch's ANOVA test, or Kruskal-Wallis test. The simple linear regression analysis was used

to examine the presence of a linear relationship between two variables.

Figure 1-Source data1

The western blot raw images in Figure 1

Figure 1-Source data2

The labeled western blot images in Figure 1

Figure 1-Source data3

Excel file containing summary data and data analysis of Figure 1

1108
Figure 2. Neurons and astrocytes upregulated claudin-5 and VE-cadherin expression in

hCMEC/D3 cells due to GDNF secretion.

(**A**): Effects of conditioned medium (CM) on claudin-5 and VE-cadherin expression. Con: the

normal medium; S-CM: the CM from SH-SY5Y cells; U-CM: the CM from U251 cells;

US-CM: the CM from SH-SY5Y cells co-culture with U251 cells. (**B**): The mRNA expression

levels of neurotrophic factors in hCMEC/D3, U251, and SH-SY5Y cells. (**C**): Concentrations

of GDNF, bFGF, IGF-1, and TGF-β in the CMs. H-CM: the CM from hCMEC/D3 cells.

(**D**-**G**): Effects of GDNF (D), IGF-1 (E), bFGF (F), and TGF-β (G) on the expression of

claudin-5 and VE-cadherin. The dosages have been marked in the figure. (**H** and **I**): Effects of

anti-GDNF antibody on the upregulation of claudin-5 and VE-cadherin expression induced by

US-CM (H) or 200 pg/mL GDNF (I). (**J**): Effects of 200 pg/mL GDNF and US-CM on

claudin-5 and VE-cadherin expression in primary rat brain microvascular endothelial cells. (**K**

and **L**): Effects of 3 μM RET tyrosine kinase inhibitor SSP-86 (SPP), and 5 μM Src family

kinases inhibitor PP2 on the upregulation of claudin-5 and VE-cadherin induced by 200

pg/mL GDNF (K) and US-CM (L). (**M** and **N**): Effects of SPP on the TEER on day 6 (M), the

permeability of NaF, and FITC-Dex (N) of the hCMEC/D3 mono-culture BBB model treating

200 pg/mL GDNF. (**O** and **P**): Effects of PP2 on the TEER on day 6 (O), the permeability of

NaF, and FITC-Dex (P) of the hCMEC/D3 mono-culture BBB model treating 200 pg/mL

GDNF. (**Q** and **R**): Effects of SPP on the TEER on day 6 (Q), the permeability of NaF, and

FITC-Dex (R) of the triple co-culture BBB model. (**S** and **T**): Effects of PP2 on the TEER on

day 6 (S), the permeability of NaF, and FITC-Dex (T) of the triple co-culture BBB model.

The above data are shown as the mean ± SEM. Four biological replicates per group. For B

and C, two technical replicates per biological replicate. One technical replicate per biological

replicate for the rest. * $p < 0.05$; ** $p < 0.01$ by one-way ANOVA test followed by Fisher's

LSD test, Welch's ANOVA test, or Kruskal-Wallis test.

Figure 2-Source data1

The western blot raw images in Figure 2

Figure 2-Source data2

The labeled western blot images in Figure 2

Figure 2-Source data3

Excel file containing summary data and data analysis of Figure 2

1140

Figure 3. GDNF induced claudin-5 and VE-cadherin expression in hCMEC/D3 cells by activating the PI3K/AKT and MAPK/ERK signaling.

(**A**): Effects of 3 μM LY294002 (LY) on the levels of claudin-5, VE-cadherin, and p-AKT/AKT in hCMEC/D3 cells stimulated by 200 pg/mL GDNF. (**B**): Effects of 2 μM U0126 (U0) on the levels of claudin-5, VE-cadherin, and p-ERK/ERK in hCMEC/D3 cells stimulated by 200 pg/mL GDNF. (**C**): Effects of 5 μM SP600125 (SP) on the levels of claudin-5, VE-cadherin, and p-JNK/JNK in hCMEC/D3 cells stimulated by 200 pg/mL GDNF. (**D**): Effects of 2 μM SB203580 (SB) on the levels of claudin-5, VE-cadherin, and p-p38/p38 in hCMEC/D3 cells stimulated by 200 pg/mL GDNF. (**E**): Effects of anti-GDNF antibody on the GDNF-induced p-AKT/AKT and p-ERK/ERK ratios. (**F**): Effects of 3 μM LY on the levels of claudin-5, VE-cadherin, and p-AKT/AKT in hCMEC/D3 cells stimulated by US-CM. (**G**): Effects of 2 μM U0 on the levels of claudin-5, VE-cadherin, and p-ERK/ERK in hCMEC/D3 cells stimulated by US-CM. (**H**): Effects of 5 μM SP on the levels of claudin-5, VE-cadherin, and p-JNK/JNK in hCMEC/D3 cells stimulated by US-CM. (**I**): Effects of 2 μM SB on the levels of claudin-5, VE-cadherin, and p-p38/p38 in hCMEC/D3 cells

stimulated by 2 US-CM. (**J**): Effects of anti-GDNF antibody on the US-CM-induced p-AKT/AKT and p-ERK/ERK ratios. The above data are shown as the mean ± SEM. Four biological replicates per group. One technical replicate for each biological replicate. * $p < 0.05$; ** $p < 0.01$ by one-way ANOVA test followed by Fisher's LSD test or Welch's ANOVA test.

Figure 3-Source data1

The western blot raw images in Figure 3

Figure 3-Source data2

The labeled western blot images in Figure 3

Figure 3-Source data3

Excel file containing summary data and data analysis of Figure 3

Figure 4. GDNF induced the claudin-5 expression in hCMEC/D3 cells by activating the PI3K/AKT/FOXO1 pathway.

(**A** and **B**): Effects of US-CM and GDNF on the phosphorylated FOXO1 (p-FOXO1)/FOXO1 ratio, total FOXO1 expression (A), cytoplasmic p-FOXO1, cytoplasmic FOXO1, and nuclear

FOXO1 expression (B). (**C** and **D**): The expression levels of total and nuclear FOXO1 (C),

claudin-5, and VE-cadherin (D) in hCMEC/D3 cells transfected with *FOXO1* siRNA

(siFOXO1). NC: negative control. (**E**): Effects of *FOXO1* overexpression (FOXO1-OE) and

GDNF on the expression levels of claudin-5, total FOXO1, and nuclear FOXO1. FOXO1-NC:

negative control plasmids. (**F**): Effects of LY and U0 on GDNF-induced alterations of total

p-FOXO1/FOXO1 ratio, cytoplasmic p-FOXO1, cytoplasmic FOXO1, and nuclear FOXO1

expression. (**G**): Effects of LY on the claudin-5 expression upregulated by siFOXO1. The

above data are shown as the mean ± SEM. Four biological replicates per group. One technical

replicate for each biological replicate. * $p < 0.05$; ** $p < 0.01$ by one-way ANOVA test

followed by Fisher's LSD test, Welch's ANOVA test, or Kruskal-Wallis test.

Figure 4-Source data1

The western blot raw images in Figure 4

Figure 4-Source data2

The labeled western blot images in Figure 4

Figure 4-Source data3

Excel file containing summary data and data analysis of Figure 4

1189

1190
Figure 5. GDNF induced VE-cadherin expression in hCMEC/D3 cells by activating the
PI3K/AKT/ETS1 and MAPK/ERK/ETS1 pathways.

(**A** and **B**): Effects of US-CM and GDNF on total (A) and nuclear (B) ETS1 expression. (**C** and **D**): Effects of LY and U0 on 200 pg/mL GDNF-induced total (C) and nuclear (D) ETS1 expression. (**E** and **F**): Expression levels of total (E) and the nuclear ETS1 (F) in hCMEC/D3 cells after knocking down *ETS1* with siRNA (siETS1). (**G**-**I**): Effects of GDNF and siETS1 on the expression of VE-cadherin and claudin-5. The above data are shown as the mean ± SEM. Four biological replicates per group. One technical replicate for each biological replicate. * $p < 0.05$; ** $p < 0.01$ by one-way ANOVA test followed by Fisher's LSD test.

Figure 5-Source data1
The western blot raw images in Figure 5
Figure 5-Source data2
The labeled western blot images in Figure 5
Figure 5-Source data3
Excel file containing summary data and data analysis of Figure 5
1206

1207

Figure 6. The deficiency of brain GDNF in mice increased the permeability of BBB and reduced claudin-5 and VE-cadherin expression in mice brains.

(**A**): Experimental configuration of AAV-GFP (shNC) or AAV-sh*Gdnf* (sh*Gdnf*) intracerebroventricular injection. (**B**): Effects of brain-specific *Gdnf* silencing on the expression levels of GDNF, claudin-5, and VE-cadherin in the brains. (**C**-**E**): Effects of brain-specific *Gdnf* silencing on NaF levels in plasma (C), brain (D), and the ratio of brain to plasma (E). (**F**-**H**): Effects of brain-specific *Gdnf* silencing on FITC-Dex levels in plasma (F), brain (G), and the ratio of brain to plasma (H). (**I**-**K**): The expression ratios of p-AKT/AKT (I), p-ERK/ERK (J), and p-FOXO1/FOXO1 (K) in the brains of *Gdnf* silencing mice. (**L**): The expression level of ETS1 in the brains of *Gdnf* silencing mice. The above data are shown as the mean ± SEM. Six biological replicates per group. One technical replicate for each biological replicate. * $p < 0.05$; ** $p < 0.01$ by unpaired t-test, unpaired t-test with Welch's correction, or Mann-Whitney test.

Figure 6-Source data1

The western blot raw images in Figure 6

Figure 6-Source data2

The labeled western blot images in Figure 6

Figure 6-Source data3

Excel file containing summary data and data analysis of Figure 6

1227

1228

Figure 7. *In vitro/in vivo* correlation assay of BBB permeability.

(**A**): The comparison of the estimated permeability coefficient-surface area product (PS_{Mono}) recalculated from $P_{app, Mono}$ with the observed *in vivo* PS values (PS_{Obs}). (**B**): The comparison of the estimated permeability coefficient-surface area product (PS_{Triple}) recalculated from $P_{app, Triple}$ with the observed *in vivo* PS values (PS_{Obs}). The solid line represents a perfect prediction, and the dashed lines represent the 0.5-2 folds of their observations. The PS_{Obs} values were determined by *in situ* brain perfusion in rodents, which were collected from the literature.

1236
Figure 8. The mechanism of neurons and astrocytes induced the integrity of brain endothelial cells. Neurons but also astrocytes trigger the activation of PI3K/AKT and MAPK/ERK pathways in brain endothelial cells by GDNF secretion, which in turn regulates transcription factors of claudin-5 (FOXO1) and VE-cadherin (ETS1) to promote claudin-5 and VE-cadherin expression and leads to the enhancement of BBB integrity. Meanwhile, with the increase in barrier integrity, the *in vitro* BBB model also obtained a stronger *in vivo* correlation.

1244

1245

1246 **Figure 9. Schematic diagram of the establishment process of the triple co-culture BBB**

1247 **model.**

1248

Table 1. The unbound fraction in brain ($f_{u, brain}$), the observed PS_{Obs}, and the predicted PS (PS_{Pre}), P_{app} across the hCMEC/D3 mono-culture model ($P_{app, Mono}$) and triple co-culture model ($P_{app, Triple}$) of the tested drugs. One technical replicate of 4 biological replicates per group.

Compounds	$f_{u, brain}$	PS_{Obs} μl/min/g	$P_{app, Mono}$ cm/s×10^{-6}	$PS_{Pre, Mono}$ μl/min/g	$P_{app, Triple}$ cm/s×10^{-6}	$PS_{Pre, Triple}$ μl/min/g
Amantadine	0.1985[a]	116.10[b]	6.84±0.95	310.22	3.64±0.26	165.23
Amitriptyline	0.01[c]	4608.00[k]	15.24±0.64	13716.00	14.61±0.27	13149.00
Bupropion	0.12[d]	1519.20[b]	15.19±0.20	1139.58	11.34±0.44	850.58
Carbamazepine	0.116[e]	959.40[b]	34.37±1.26	2666.89	11.71±0.15	908.69
Clozapine	0.014[f]	2260.80[b]	38.97±0.54	25052.57	12.87±2.06	8272.72
Donepezil	0.07[g]	1581.30[b]	20.91±0.75	2688.43	14.47±0.84	1860.43
Doxepin	0.025[b]	2192.40[b]	16.88±1.08	6076.80	10.66±0.92	3837.60
Fluoxetine	0.004[b]	2698.20[b]	11.48±0.84	2583.00	9.97±1.03	2243.25
Gabapentin	0.782[b]	162.90[b]	16.75±1.62	192.77	8.78±0.23	101.05
Lamotrigine	0.273[b]	126.00[b]	14.26±0.37	470.11	5.97±0.11	196.88
Metoclopramide	0.365[b]	125.10[b]	14.14±1.44	348.66	6.63±0.42	163.41
Midazolam	0.045[h]	2727.00[b]	25.30±1.00	5060.00	19.09±0.24	3818.00
Mirtazapine	0.08[b]	1912.50[b]	23.44±0.44	2637.00	17.86±0.21	2009.25
Olanzapine	0.034[b]	2279.70[b]	22.91±3.80	6064.41	12.49±0.53	3306.18
Prazosin	0.09[i]	169.20[j]	5.61±0.38	560.93	2.81±0.52	280.99
Risperidone	0.099[b]	849.60[b]	16.10±2.87	1463.64	11.70±0.25	11063.64
Venlafaxine	0.205[a]	584.10[b]	9.58±0.28	420.6	8.25±0.36	362.02
Verapamil	0.033[c]	335.70[c]	7.21±0.41	1965.24	5.56±0.06	1517.67

[a](Esaki et al., 2019); [b](Summerfield et al., 2007); [c](Friden et al., 2011); [d](Bhattacharya et al., 2021); [e](Maurer et al., 2005); [f](Cremers et al., 2012); [g](Di et al., 2011); [h](Kodaira et al., 2011); [i](Zhou L. et al., 2009); [j](Di Marco et al., 2019); [k](Avdeef et al., 2010).

Table 1-Source data1

The apparent permeability coefficients of 18 tested drugs from mono or triple culture BBB model

1259   **Table 2** Initial concentrations in donor chamber and chromatographic conditions of prazosin,

1260   verapamil, and lamotrigine

Compound	Concentration	Wavelength
Prazosin	5 μM	Ex: 250 nm Em: 390 nm
Verapamil	5 μM	Ex: 280 nm Em: 310 nm
Lamotrigine	6 μM	220nm

1261

1262   **Table 3** The summary of mass charge ratio, extraction, initial concentrations in donor

1263   chamber

Compound	Concentration	Mass charge ratio $[M+H]^+$	Extraction
Amantadine	3 μM	181	Water-saturated N-butanol
Amitriptyline	1.5 μM	278	Ethyl acetate
Bupropion	3 μM	240	Ethyl acetate
Carbamazepine	3 μM	237	Ethyl acetate
Clozapine	4 μM	327	Ethyl acetate
Donepezil	3 μM	380	Methyl tert-butyl ether
Doxepin	4 μM	317	Methyl tert-butyl ether
Fluoxetine	3 μM	310	Ethyl acetate
Gabapentin	10 μM	172	Ethyl acetate
Metoclopramide	4 μM	301	Ethyl acetate
Midazolam	3 μM	327	Ethyl acetate
Mirtazapine	3 μM	266	Methyl tert-butyl ether
Olanzapine	3 μM	313	Methyl tert-butyl ether
Risperidone	4 μM	427	Methyl tert-butyl ether
Venlafaxine	10 μM	278	Ethyl acetate

1264

Table 4 Primer sequences for qPCR for indicted genes

Gene (protein)	Forwards Primer, 5'→3'	Reverse Primer, 3'→5'
ACTB (*β*-actin)	GGACTTCGAGCAAGAGATGG	AGCACTGTGTTGGCGTACAG
GAPDH (GAPDH)	TGTGGGCATCAATGGATTTGG	ACACCATGTATTCCGGGTCAAT
CLDN5 (claudin-5)	CTCTGCTGGTTCGCCAACAT	CAGCTCGTACTTCTGCGACA
OCLN (occludin)	ACAAGCGGTTTTATCCAGAGTC	GTCATCCACAGGCGAAGTTAAT
TJP1 (ZO-1)	ACCAGTAAGTCGTCCTGATCC	TCGGCCAAATCTTCTCACTCC
CDH5 (VE-cadherin)	AAGCGTGAGTCGCAAGAATG	TCTCCAGGTTTTCGCCAGTG
ABCB1 (P-gp)	TTGCTGCTTACATTCAGGTTTCA	AGCCTATCTCCTGTCGCATTA
ABCG2 (BCRP)	ACGAACGGATTAACAGGGTCA	CTCCAGACACACCACGGAT
SLC22A1 (OCT1)	ACGGTGGCGATCATGTACC	CCCATTCTTTTGAGCGATGTGG
SLC22A2 (OCT2)	CATCGTCACCGAGTTTAACCTG	AGCCGATACTCATAGAGCCAAT
SLC22A8 (OAT3)	ATGGCCCAGTCTATCTTCATGG	GACGGTGCTCAGGGTAATGC
SLCO1A1 (OATP1A1)	TAATGTGGGTGTACGTCCTAGT	GCTCCTGTTTCTACAAGCCCAA
GDNF (GDNF)	GCAGACCCATCGCCTTTGAT	CCACACCTTTTAGCGGAATGC
BDNF (BDNF)	CTACGAGACCAAGTGCAATCC	AATCGCCAGCCAATTCTCTTT
NGF (NGF)	TGTGGGTTGGGGATAAGACCA	GCTGTCAACGGGATTTGGGT
IGF1 (IGF-1)	GCTCTTCAGTTCGTGTGTGGA	GGTCATGGATGGACCTTACTGT
VEGFA (VEGF)	CCCACTGAGGAGTCCAACAT	AAATGCTTTCTCCGCTCTGA
FGF2 (bFGF)	AGAAGAGCGACCCTCACATCA	CGGTTAGCACACACTCCTTTG
TGFB1 (TGF-*β*)	GGCCAGATCCTGTCCAAGC	GTGGGTTTCCACCATTAGCAC

Table 5 The target sequences for siRNA or shRNA

Gene	Target sequence
ETS1	CGCTATACCTCGGATTACT
FOXO1	AATCTCCTAGGAGAAGAGCTG
Gdnf	GCCAGTGTTTATCTGATAC
CDH5	GCCTCTGTCATGTACCAAA

1269   **Figure 1—figure supplement 1**

1270

Figure 1—figure supplement 1: The induced proliferation of hCMEC/D3 cells by

bFGF slightly reduced the permeability of cell layers. EdU incorporation (A), cell viability

(B), and apparent permeability coefficient (P_{app}, $\times$ 10^{-6} cm/s) of fluorescein (C) or

FITC-Dextran 3–5 kDa (D) of hCMEC/D3 cells treated with bFGF (6, 20, 60 ng/mL) for 6

days. The above data are shown as the mean ± SEM. Four biological replicates per group.

One technical replicate for each biological replicate. Statistical significance was determined

using one-way ANOVA test followed by Fisher's LSD test or Welch's ANOVA test.

1278

1279 **Figure 4—figure supplement 1**

1280

Figure 4—figure supplement 1: The contribution of VE-Cadherin on the GDNF-induced

claudin-5 expression. Effects of the VE-Cadherin siRNA (siVE-Cad) on mRNA expression

of VE-cadherin (A) and claudin-5 (B). Effects of siVE-Cad and GDNF on claudin-5 and

VE-cadherin protein expression (C). NC: negative control plasmids. The above data are

shown as the mean ± SEM. Four biological replicates per group. Two technical replicates for

A and B and one technical replicate for C. Statistical significance was determined using

unpaired Student's t-test or one-way ANOVA test followed by Fisher's LSD test.

1288

Figure 1-Source data1

The western blot raw images in Figure 1

Figure 1-Source data2

The labeled western blot images in Figure 1

Figure 1-Source data3

Excel file containing summary data and data analysis of Figure 1

Figure 2-Source data1

The western blot raw images in Figure 2

Figure 2-Source data2

The labeled western blot images in Figure 2

Figure 2-Source data3

Excel file containing summary data and data analysis of Figure 2

Figure 3-Source data1

The western blot raw images in Figure 3

Figure 3-Source data2

The labeled western blot images in Figure 3

Figure 3-Source data3

Excel file containing summary data and data analysis of Figure 3

Figure 4-Source data1

The western blot raw images in Figure 4

Figure 4-Source data2

The labeled western blot images in Figure 4

Figure 4-Source data3

Excel file containing summary data and data analysis of Figure 4

Figure 5-Source data1

The western blot raw images in Figure 5

Figure 5-Source data2

The labeled western blot images in Figure 5

Figure 5-Source data3

Excel file containing summary data and data analysis of Figure 5

Figure 6-Source data1

The western blot raw images in Figure 6

Figure 6-Source data2

The labeled western blot images in Figure 6

Figure 6-Source data3

Excel file containing summary data and data analysis of Figure 6

Table 1-Source data1

The apparent permeability coefficients of 18 tested drugs from mono or triple culture BBB

model

ETS1

MEK → ERK

Raf

Ras

CDH5

VE-cadherin

FOXO1

CLDN5

PI3K → PDK1 → AKT

FOXO1 P

Claudin-5

GDNF Secretion

GDNF

GFRα1

RET

Permeability ↓

TEER ↑

Triple co-culture
BBB Model

IVIVC

$PS_{Pre, Triple}$ (µl/min/g)

PS_{Obs} (µl/min/g)

hCMEC/D3

U251

SH-SY5Y