1 **Geographically widespread and novel hemotropic mycoplasmas and bartonellae in**
2 **Mexican free-tailed bats and sympatric North American bat species**
3

4 Daniel J. Becker[1*], Kristin E. Dyer[1], Lauren R. Lock[1], Shawn A. Pladas[2], Anushka A. Sukhadia[1],
5 Bret Demory[1], Juliana Maria Nunes Batista[1,3], Micaela Pineda[2], Nancy B. Simmons[4], Amanda
6 M. Adams[5], Winifred F. Frick[5,6], M. Teague O'Mara[2,5,7,8], Dmitriy V. Volokhov[9]
7
[1] School of Biological Sciences, University of Oklahoma, Norman, OK, USA
[2] Department of Biological Sciences, Southeastern Louisiana University, Hammond, LA, USA
[3] Department of Preventive Veterinary Medicine and Animal Health, School of Veterinary
Medicine and Animal Science, University of São Paulo, São Paulo, Brazil
[4] Department of Mammalogy, Division of Vertebrate Zoology, American Museum of Natural
History, New York, NY, USA
[5] Bat Conservation International, Austin, TX, USA
[6] Department of Ecology and Evolutionary Biology, University of California Santa Cruz, Santa
Cruz, CA, USA
[7] Department of Migration, Max Planck Institute of Animal Behavior, Radolfzell, Germany
[8] Smithsonian Tropical Research Institute, Panama City, Panama
[9] Center for Biologics Evaluation and Research, U.S. Food and Drug Administration, Silver
Spring, MD, USA
*Corresponding author, danbeck@ou.edu

Keywords: hemoplasmas, *Tadarida brasiliensis*, migration, One Health

Running head: Bacterial pathogens in Mexican free-tailed bats
25
Abstract

Bacterial pathogens remain poorly characterized in bats, especially in North America. We
describe novel (and in some cases panmictic) hemoplasmas (12.9% positivity) and bartonellae
(16.7% positivity) across three colonies of Mexican free-tailed bats (*Tadarida brasiliensis*), a
partially migratory species that can seasonally travel hundreds of kilometers. Molecular analyses
identified three novel *Candidatus* hemoplasma species most similar to another novel *Candidatus*
species in Neotropical molossid bats. We also detected novel hemoplasmas in sympatric cave
myotis (*Myotis velifer*) and pallid bats (*Antrozous pallidus*), with sequences in the latter 96.5%
related to *C.* Mycoplasma haemohominis. We identified eight *Bartonella* genotypes, including
those in cave myotis, with 96.7% similarity to *C.* Bartonella mayotimonensis. We also detected
Bartonella rochalimae in migratory *Tadarida brasiliensis*, representing the first report of this
human pathogen in bats. The seasonality and diversity of these bacteria observed here suggest
that additional longitudinal, genomic, and immunological studies in bats are warranted.

39
40

Introduction

Bats have been intensively sampled for viral pathogens, with species in this mammalian order hosting multiple viruses with high virulence in humans (1). However, bats remain understudied for bacterial pathogens, which can be significant for their impacts on both human health and bat morbidity and even mortality (2,3). Hemotropic mycoplasmas (hemoplasmas) and bartonellae are facultative intracellular bacteria of special interest in bats, given their high prevalence and substantial genetic diversity (4,5). For example, sampling of Neotropical bat communities has identified many common and co-circulating genotypes of these bacteria (6–8). Surveys in Oceania and Europe have also supported plausible zoonotic transmission of these bacteria from bats to humans, including *Candidatus* Mycoplasma haemohominis and *C.* Bartonella mayotimonensis (9,10). Greater characterization of these bacteria across global bat diversity (over 1,470 species) is therefore warranted to inform infection risks for both bats and humans, although little surveillance has thus far been conducted in North American bats (11,12).

Although flight enables high mobility of bats, relatively few bat species undertake long-distance migrations (e.g., between maternity and wintering grounds) (13). In North America, Mexican free-tailed bats (*Tadarida brasiliensis*) display highly variable migratory strategies (14). The southwestern United States contains both non-migratory and migratory populations, with some individuals traveling hundreds to over 1,000 kilometers between wintering grounds in Mexico to northern maternity colonies in Oklahoma, Kansas, and Colorado (14–16). Other colonies across the species range include year-round residents (17,18). This variation in migratory behavior could shape patterns of infection, including the seasonal dispersal of bacterial pathogens across landscapes to naïve hosts. Hemoplasmas have not yet been detected in this bat species (4,19), and bartonellae have only been minimally described in the southernmost part of the bat species' geographic range (i.e., Chile and Argentina) (20).

Here, we conducted an initial characterization of hemoplasmas and bartonellae in Mexican free-tailed bats across multiple populations and seasons. Our goals were to identify novel pathogens in this bat species and to test for differences in prevalence among colonies that differ in migratory strategy and across the bat annual cycle. We also tested whether pathogen lineages were unique to each geography or if migration may facilitate panmixia. Lastly, we used this opportunity to perform a pilot characterization of these pathogens in sympatric bat species.

Material and Methods

Wild bat sampling

We sampled three North American colonies of Mexican free-tailed bats in 2021 and 2022 to compare infections among migratory strategies and provide an initial assessment of pathogen seasonality. We sampled non-migratory individuals in southeastern Louisiana (17), focusing on a colony in Pine Grove of approximately 1,000 bats, during the non-breeding season (October 2021, *n*=5) and maternity season (July 2022, *n*=9). We also sampled the partially migratory population of Bracken Cave near San Antonio, Texas, which hosts tens of millions of Mexican free-tailed bats in the maternity season and declines to approximately 10,000 bats in winter

(16,21). Sampling focused on the maternity season (August 2021, *n*=20) and mid-to-late winter
(December 2021 and March 2022; *n*=9 and *n*=19). Lastly, we sampled a fully migratory colony
at the Selman Bat Cave near Freedom, Oklahoma, where this maternity roost holds up to 100,000
bats in summer and is empty during winter (16,22). We sampled after spring arrival and into the
maternity period (April, May, and June 2022; *n=7*, *n*=46, and *n*=24). In the same Oklahoma site,
we also sampled four cave myotis (*Myotis velifer*), one hoary bat (*Lasiurus cinereus*), one
Townsend's big-eared bat (*Corynorhinus townsendii*), and two pallid bats (*Antrozous pallidus*).
Bats were captured with hand nets and mist nets while emerging from or returning to
roosts and placed in individual cloth bags. Bats were identified to species by morphology and
identified by sex, reproductive status, and age (23). Blood (<1% body mass) was sampled by
lancing the propatagial vein using 27G and 30G needles and collected with heparinized
capillaries. Blood was preserved on Whatman FTA cards and held at room temperature until
−20°C storage at the University of Oklahoma (OU). Sampling was approved by the Institutional
Animal Care and Use Committees of OU (2022-0198) and Southeastern Louisiana University
(0064), with permits from the Texas Parks and Wildlife Department (SPR-0521-063), Louisiana
Department of Wildlife and Fisheries (WDP-21-101), and Oklahoma Department of Wildlife
Conservation (ODWC, 10567389). All bats were released after sampling at the capture site.
98
Molecular diagnostics
We extracted genomic DNA from blood using QIAamp DNA Investigator Kits (Qiagen). To
determine hemoplasma presence, we used PCR targeting the partial 16S and 23S rRNA genes
(Table S1; (6,7,24,25), with amplicons purified and sequenced at Psomagen. For DNA samples
positive for the 16S or 23S rRNA genes, we also attempted to amplify the partial *rpoB* gene,
using primers newly designed for this study (Table S1). To determine the presence of
bartonellae, we used nested PCR targeting the *gltA* gene (Table S1; 26), with amplicons purified
using Zymo kits (DNA Clean & Concentrator-5, Zymoclean Gel DNA Recovery) and sequenced
at the North Carolina State University Genomic Sciences Laboratory. We included blank FTA
card punches and ultrapure water as extraction and negative controls, respectively, in all PCR
assays. Sequences are available on GenBank through accessions OQ407831–50, OR783323,
OQ359160–75, OQ554332–38, and PP101634.
111
Statistical analysis
We analyzed infection states using generalized linear models (GLMs) with binary response in R.
Given the relatively small sample sizes included here, all our GLMs were fit using mean bias
reduction methods using the *brglm2* package (27). For each of our two pathogens, we fit the
following five GLMs. The first model was fit to all data and included a main effect of site to
compare odds of infection among Louisiana, Texas, and Oklahoma bats. The second model was
limited to the early-to-mid non-breeding season and compared odds of infection between Texas
and Louisiana bats. The third model was limited to the late overwintering period and spring
arrival to compare odds of infection between Texas and Oklahoma bats. The fourth model was

limited to Texas and included main effects of sex, reproductive status, age, and month to test
demographic effects and seasonality (i.e., the maternity season through the late overwintering
period). Similarly, the fifth model included the same main effects but was limited to Oklahoma
to test a different aspect of seasonality (i.e., spring arrival into the late maternity season).
125
Phylogenetic analysis
We first used NCBI BLASTn to identify related mycoplasma (16S rRNA, 23S rRNA, *rpoB*) and
bartonellae sequences (*gltA*), which we aligned with our sequences and reference sequences
using MUSCLE. We then used MrBayes for phylogenetic analysis, with each gene tree run for
20,000,000 generations using a GTR+I+G model. BLASTn was implemented in Geneious,
whereas MUSCLE and MrBayes were implemented using the NGPhylogeny.fr platform (28).
We delineated genotypes of hemoplasmas and bartonellae based on pairwise similarity among
sequences and clustering on their phylogenies, using established criteria for defining novel
bacterial genotypes (6,29). For hemoplasmas, we also used our multi-loci data to propose novel
Candidatus species when the same genotype was identified in at least two individuals using 16S
rRNA and at least one other marker (i.e., 16S rRNA and 23S rRNA, 16S rRNA and *rpoB*) (7,30).
We conducted two tests to assess if pathogen lineages were unique to each Mexican free-
tailed bat colony, which would suggest geographically constrained transmission dynamics. First,
we used chi-squared tests with *p* values generated by a Monte Carlo procedure to quantify
associations between geography and genotype assignments. Next, we derived matrices of spatial
and phylogenetic distance among sequenced PCR-positive samples and used Mantel tests with
the *vegan* R package to assess isolation by distance (31). Both tests used 1,000 randomizations.
143
Results
Migratory and seasonal effects on bat bacterial infection
We detected hemoplasmas in 18 of 139 Mexican free-tailed bats when targeting the partial 16S
or 23S rRNA genes (12.9%, 95% CI: 8.4−19.5%). Sequencing of the 23S rRNA gene showed
three other bats (two from Texas and one from Oklahoma) had non-hemotropic mycoplasmas
(OQ359169–70, OQ359174) most related to *Mycoplasma muris* (97.5%). We also detected
bartonellae in 22 of 132 tested bats (16.7%, 95% CI: 11.3–23.9%; we did not have remaining
DNA for all bats). Only five Mexican free-tailed bats were coinfected by hemoplasmas and
bartonellae (3.6%, 95% CI: 1.5–8.1%). Hemoplasmas were detected in all three colonies, while
bartonellae were only detected in the Texas and Oklahoma colonies. PCR data are fully available
in the Pathogen Harmonized Observatory (PHAROS): https://pharos.viralemergence.org/.
Across all Mexican free-tailed bats (model 1), we did not detect colony differences in the
odds of infection with hemoplasmas ($\chi^2 = 0.40$, $p = 0.94$) or bartonellae ($\chi^2 = 3.65$, $p = 0.30$)
(Figure 1). When comparing only the resident and partially migratory populations during the
non-breeding season (model 2), Louisiana and Texas bats did not differ in the odds of either type
of infection (hemoplasmas: $\chi^2 = 1.21$, $p = 0.27$; bartonellae: $\chi^2 = 0$, $p = 1$). When comparing only
the partially and fully migratory populations in spring (model 3), we did not detect colony

differences for hemoplasmas ($\chi^2 = 0$, $p = 1$), although Texas bats were marginally more likely to
be infected with bartonellae than newly arrived Oklahoma bats ($\chi^2 = 1.79$, $p = 0.18$). When
assessing risk factors of infection in the partially migratory Texas colony (model 4), we found no
evidence of seasonal or demographic effects for either pathogen (Table S2). However, when
assessing the same risk factors for the fully migratory Oklahoma colony, we identified strong
seasonal effects (Table S3). Prevalence was greatest later in the maternity season for both
hemoplasmas (odds ratio = 13.24, $p = 0.02$) and bartonellae (odds ratio = 12.89, $p < 0.01$).

Genetic diversity of Mexican free-tailed bat hemoplasmas
Sequencing of 16S rRNA amplicons revealed four hemoplasma genotypes specific to Mexican
free-tailed bats (i.e., TB1–4; Figure 2, Table 1). The TB1 genotype was 97% similar to the MR1
genotype that we earlier isolated from another molossid bat (*Molossus nigricans*) in Belize (e.g.,
MH245174) (6), and TB1 was found in all three sampled populations. In contrast, TB2 was only
detected in the Texas colony and was ~96–97% similar to hemoplasmas from carnivores and
rodents and to the Belize bat MR1 genotype (6,32,33). Both TB3 and TB4 were only detected in
the Oklahoma colony and were ~99% similar to the MR1 genotype. Amplification of the 23S
rRNA gene from two Mexican free-tailed bats (one each from Texas and Oklahoma) also found
a genotype that we initially detected in cave myotis (i.e., MV1; Figure S1) as well as the above
Mycoplasma muris–like (non-hemotropic) genotype. These six mycoplasma genotypes were only
weakly associated with bat geography ($\chi^2 = 16.50$, $p = 0.08$), although we found general support
for isolation by distance in mycoplasma genetic diversity for both the 16S rRNA phylogeny
(Mantel $r = 0.72$, $p = 0.001$) and the 23S rRNA phylogeny (Mantel $r = 0.50$, $p = 0.01$).
Amplification of paired partial 23S rRNA (Figure S1) and/or *rpoB* (Figure S2) genes for
samples belonging to these 16S rRNA genotypes suggested at least three novel *Candidatus*
hemoplasma species circulate in Mexican free-tailed bats. Based on 100% identity of two *rpoB*
sequences (OQ554335–36) and 100% identity of paired 16S rRNA sequences included in the
TB1 genotype, first detected in Bracken Cave (OQ407846, OQ407848), we propose the name *C.*
Mycoplasma haematobrackenitadaridae sp. nov. Similarly, given 99.98% identity among seven
23S rRNA sequences (OQ359161–65, OQ359168, OQ359172) and 100% identity in paired 16S
rRNA sequences included in the TB3 genotype from the Selman Bat Cave (OQ407832,
OQ407836–39, OQ407843–44), we propose the name *C.* M. haematoselmanitadaridae sp. nov.
Lastly, given 100% identity of two 23S rRNA sequences (OQ359160, OQ359166) and 99.8%
identity in paired 16S rRNA sequences from the TB4 genotype (also identified from the Selman
Bat Cave; OQ407831, OQ407840), we propose the name *C.* M. haematotraderitadaridae sp. nov.
(Figures 2 and S1), based on the stream running adjacent to the bat cave (Traders Creek) (34).
Given the similarity of Mexican free-tailed bat hemoplasma 16S rRNA sequences to
those from molossid bats sampled in Belize (6), we also attempted to amplify the 23S rRNA and
rpoB genes from *Molossus nigricans* sampled in 2017 and 2018 in Belize that previously tested
positive for the MR1 and MR2 genotypes (6). We re-extracted DNA from four FTA cards and
applied the same additional PCR protocols described earlier. We obtained partial 23S rRNA and

rpoB sequences for two (OQ518943–44) and three (OQ554329–31) *M. nigricans*, respectively.
Based on 100% inter-sequence similarity of the *rpoB* sequences and high ($\bar{x}$ = 99.98%) identity
of paired 16S rRNA sequences (MH245122, MH245172, MH245174), we propose the name *C.*
Mycoplasma haematomolossi sp. nov. to designate this novel hemoplasma (Figures 2 and S2).
205
Genetic diversity of Mexican free-tailed bat bartonellae
Sequencing of *gltA* amplicons next revealed at least three bartonellae genotypes circulating in
Mexican free-tailed bats (Figure 3, Table 1). The first genotype was detected in the Texas and
Oklahoma populations, with sequences from all three sampled months across 2021 and 2022
from Bracken Cave. The corresponding *gltA* sequences identified here were ≥99.6% similar to
those recently detected in Mexican free-tailed bats in Argentina (KX986617) (20), such that we
therefore consider these sequences to all form the TB1 genotype. The TB2 genotype was only
found in the migratory Oklahoma population and was identical to bartonella from streblid bat
flies in Mexico (i.e., ≥99.7% identity to MF988072 and MF988082) (35). In addition to the TB1
and TB2 genotypes, the Oklahoma population also harbored g*ltA* sequences with 100% identity
to *Bartonella rochalimae* from red foxes (OQ834668) and humans (DQ683195) (36,37). These
three bartonellae genotypes were not associated with Mexican free-tailed bat geography (χ^2 =
3.00, *p* = 0.38), and we likewise found no strong support for isolation by distance in the genetic
diversity of bartonellae when using our *glt*A phylogeny (Mantel *r* = 0.22, *p* = 0.30).
220
Bacterial infections of sympatric bat species
Opportunistic sampling of other bats in Oklahoma revealed further bacterial diversity (Table 1).
Three of the four cave myotis and both pallid bats tested positive for hemoplasmas, whereas
three of three cave myotis, both pallid bats, and the single Townsend's big-eared bat tested
positive for bartonellae; the single hoary bat tested negative for both bacterial pathogens.
For hemoplasmas (Figure 2), we identified a single novel hemoplasma genotype in each
positive species (i.e., MV1 and AP1), with 16S rRNA sequences most closely related (i.e., ≥98%
similarity) to previously detected genotypes in vesper bats from Chile and Belize (e.g., EF1 and
MYE) (6,19). Notably, 16S rRNA sequences of the AP1 genotype were 96.5% similar to those
of *C.* Mycoplasma haemohominis (i.e., GU562823), whereas those of the MV1 genotype were
only ~94% similar. 23S rRNA and *rpoB* sequences from cave myotis (OQ359173, OQ554337)
were entirely novel (<85% similarity to GenBank sequences), with the former 100% similar to
select 23S rRNA sequences from our Mexican free-tailed bats in both Oklahoma and Texas.
For bartonellae (Figure 3), all three positive cave myotis had unique genotypes. One *gltA*
sequence was 98.5% similar to that first identified in little brown bats (*Myotis lucifugus*)
elsewhere in North America (i.e., KX807172), here denoted the ML1 genotype. A new genotype
first detected in our cave myotis (i.e., MV1) was distantly related (<91%) to bartonellae from
nycteribid bat flies in Europe (i.e., MW007693) and to *Bartonella quintana* (i.e., Z70014) (38).
The final cave myotis *gltA* sequence clustered within a clade of *Candidatus* Bartonella
mayotimonensis sequences (~96%), isolated from a human endocarditis patient in Iowa, USA

(FJ376732) and other little brown bats in North America (KX807177–9) (11,39). For pallid bats,
one sequence also belonged to the MV1 genotype, whereas the other formed the novel AP1
genotype most related (~95%) to bartonellae from common bent-wing bats (*Miniopterus*
schreibersii) and their bat flies in Eastern Europe (e.g., MK140349 and MK140263) (40,41).
Lastly, the single positive Townsend's big-eared bat hosted a unique genotype (CT1) only ~94%
similar to bartonellae identified from *Myotis nigricans* in Guatemala (e.g., MN529509) (5).

Discussion
Hemoplasmas and bartonellae are emerging as model systems for studying bacterial infections in
bats (5,7), but their infection dynamics and diversity remain poorly characterized, notably in
North American systems (11,12). We here demonstrate novel diversity of hemoplasmas and
bartonellae in bats in the south-central United States, including the circulation of lineages of both
pathogens with clear infection seasonality in a migratory colony of Mexican free-tailed bats.
Such work provides the foundation for further empirical studies to elucidate the transmission
dynamics of these bacteria, their pathogenicity in bats, and their possible zoonotic risk.
Our findings suggest relatively common infection with site-specific and panmictic
bacterial lineages. We found general support for pathogen isolation by distance, with most
bacterial genotypes found in a single site, indicating spatially constrained transmission.
However, we also detected bacterial genotypes in multiple sites, such as the TB1 hemoplasma
genotype in Louisiana, Texas, and Oklahoma; the MV1 hemoplasma genotype and *Mycoplasma*
muris–like genotype in Texas and Oklahoma; and the TB1 *Bartonella* genotype in Texas and
Oklahoma (for which sequences were nearly identical to those from *Tadarida brasiliensis* in
Argentina) (20). Such results may be explained by migratory connectivity in Mexican free-tailed
bats, such as between the Selman Bat Cave and Bracken Cave (14–16), suggesting migratory
behavior of this species can enhance bacterial dispersal. However, given the presence of these
genotypes in migratory and non-migratory populations (i.e., Louisiana) and at the extremes of
the bat range (e.g., over 5,000 km between the Selman Bat Cave and the Argentina site for the
TB1 *Bartonella* genotype), these results also suggest the ancestral spread of these bacteria.
Future studies are needed to identify the migratory routes of Mexican free-tailed bats,
especially for understanding the origins of possibly zoonotic bacterial lineages and the potential
for these bats to disperse infection during spring and fall migrations. Researchers could capitalize
on advances in tracking small vertebrates for long periods, such as use of absorbent sutures, to
ensure lightweight radiotags stay attached to bats for the duration of migration and winter (42).
Such work is also needed to assess if these infections negatively impact bat migration trajectory
and success, as observed for blood pathogens in migratory songbirds (43). Longitudinal studies
would also inform such analyses, as our data from the Oklahoma population in particular suggest
increasing bacterial prevalence into the maternity season. Future seasonal sampling is needed to
assess how infection risk varies across the migratory cycle, if prevalence tracks bat population
size, and whether infections are sufficiently common in autumn to facilitate dispersal with
migration. Further genetic analyses may also inform patterns of bat connectivity and studies of

pathogen spread. For example, the TB2 *Bartonella* genotype from Oklahoma Mexican free-tailed
bats showed 100% identity to bartonellae from blood-feeding streblid bat flies from Morelos,
Hidalgo, and Jalisco in central Mexico (35), all of which span the likely wintering grounds of
this bat species (14,15). Similarly, previous analyses of *Trypanasoma cruzi* from this same
Oklahoma bat population detected lineages similar to those along the Texas–Mexico border,
further showing possible southern origins of infection and high pathogen dispersal capacity (44).
Additional molecular and immunological studies are also needed to better characterize
these novel bat bacterial pathogens and their health impacts. We identified 16S rRNA and *gltA*
sequences with moderate-to-high similarity to zoonotic pathogens such as *C.* Mycoplasma
haemohominis, *C.* Bartonella mayotimonensis, and *Bartonella rochalimae* (9,37,39). For the
former two pathogens, our bat sequences were ~96% similar to zoonotic lineages, likely
indicating divergence from a common ancestor at least tens of millions of years ago (7,45).
Generation of whole genomes for our novel bat pathogens could inform their zoonotic risk, both
by better linking them to cryptic human infections (9) and by facilitating machine learning
models that predict zoonotic potential from genomic composition, as applied for viruses (46).
Other -omics analyses could also elucidate whether these bacterial infections are pathogenic in
bats themselves. In addition to assessing impacts of infection on migration outcomes as noted
above, approaches such as transcriptomics and proteomics could test if bats have a pronounced
immune response to these bacterial infections or appear largely tolerant (47). Such studies could
be especially informative when comparing immunity between migratory and non-migratory
periods, which could test whether long-distance migration may disrupt immune tolerance in bats.
Lastly, Mexican free-tailed bats and their sympatric bat species provide several important
ecosystem services, including but not limited to predating on crop pests and contributing to the
tourism economy from bat flight watching (48,49). Understanding the prevalence, genetic
diversity, and pathogenicity of bacterial pathogens in bats can inform One Health approaches
that emphasize conservation measures to promote bat, domestic animal, and human health (50).
307

Acknowledgements
This work was supported by the National Institute of General Medical Sciences of the National
Institutes of Health (P20GM134973), Louisiana Board of Regents Support Fund, Louisiana
Biomedical Research Network, and Research Corporation for Science Advancement (RCSA).
This work was conducted as part of Subawards No. 28365 and 29018, part of a USDA Non-
Assistance Cooperative Agreement with RCSA Federal Award No. 58-3022-0-005. Support was
also provided by the Vice President for Research and Partnerships of the University of Oklahoma
and the Data Institute for Societal Challenges. Bat Conservation International provided in-kind
support. We thank William Caire, Melynda Hickman, and the ODWC for site access; the Selman
Living Laboratory for fieldwork support; and Konstantin Chumakov for laboratory support.

Tables

Table 1. *Mycoplasma* spp. (A) and *Bartonella* spp. (B) genotypes identified from Louisiana,

Texas, and Oklahoma bats during this study (2021–2022). Genotypes are given with their host

species, locations, and mean intra-genotype sequence similarity from the partial 16S rRNA, 23S

rRNA, *rpoB*, or *gltA* gene sequences identified here.

	Genotype	States	Bat species	Mean intra-genotype similarity (%)
(A)	TB1[*]	LA, TX, OK	*Tadarida brasiliensis*	99.6[i], 100[iii]
	TB2[*]	TX	*Tadarida brasiliensis*	NA
	TB3[*]	OK	*Tadarida brasiliensis*	100[i], 100[ii]
	TB4[*]	OK	*Tadarida brasiliensis*	100[i], 100[ii]
	MV1[*]	TX, OK	*Myotis velifer, Tadarida brasiliensis*	100[i], 100[ii]
	AP1[*]	OK	*Antrozous pallidus*	100[i]
	M. muris–like[†]	TX, OK	*Tadarida brasiliensis*	99.3[ii]
(B)	TB1	TX, OK	*Tadarida brasiliensis*	96.5[iv]
	TB2[*]	OK	*Tadarida brasiliensis*	NA
	Bartonella rochalimae	OK	*Tadarida brasiliensis*	NA
	ML1	OK	*Myotis velifer*	NA
	MV1[*]	OK	*Myotis velifer, Antrozous pallidus*	97.9[iv]
	C. Bartonella mayotimonensis–like	OK	*Myotis velifer*	NA
	AP1[*]	OK	*Antrozous pallidus*	NA
	CT1[*]	OK	*Corynorhinus townsendii*	NA

[*] Novel genotypes

[†] Non-hemotropic mycoplasma

[i] 16S rRNA sequence

[ii] 23S rRNA sequence

[iii] *rpoB* sequence

[iv] *gltA* sequence

Figures

 Figure 1. (A) Sampled Mexican free-tailed bat (*Tadarida brasiliensis*) colonies relative to the
 host distribution in the Americas. (B) Hemoplasma and *Bartonella* spp. infection prevalence
 across months and colonies; segments denote 95% confidence intervals using Wilson's method.

346 Figure 3. Consensus Bayesian phylogeny of the partial *gltA Bartonella* spp. sequences from this
347 study (highlighted in bold and colored by geography; see Table 1 for genotype assignments) and
348 reference sequences from bats, other mammals, and ectoparasites. Nodes are colored by posterior
349 probability (nodes with less than 50% support are not shown).
350

References

1. Letko M, Seifert SN, Olival KJ, Plowright RK, Munster VJ. Bat-borne virus diversity, spillover and emergence. Nat Rev Microbiol. 2020 Aug;18(8):461–71.

2. Brook CE, Dobson AP. Bats as "special"reservoirs for emerging zoonotic pathogens. Trends Microbiol [Internet]. 2015; Available from: https://www.sciencedirect.com/science/article/pii/S0966842X14002480

3. Szentivanyi T, McKee C, Jones G, Foster JT. Trends in Bacterial Pathogens of Bats: Global Distribution and Knowledge Gaps. Transbound Emerg Dis [Internet]. 2023 Mar 27 [cited 2023 May 18];2023. Available from: https://www.hindawi.com/journals/tbed/2023/9285855/

4. Millán J, Di Cataldo S, Volokhov DV, Becker DJ. Worldwide occurrence of haemoplasmas in wildlife: Insights into the patterns of infection, transmission, pathology and zoonotic potential. Transbound Emerg Dis [Internet]. 2020 Nov 19;(tbed.13932). Available from: https://onlinelibrary.wiley.com/doi/10.1111/tbed.13932

5. McKee CD, Bai Y, Webb CT, Kosoy MY. Bats are key hosts in the radiation of mammal-associated Bartonella bacteria. Infect Genet Evol. 2021 Apr;89:104719.

6. Becker DJ, Speer KA, Brown AM, Fenton MB, Washburne AD, Altizer S, et al. Ecological and evolutionary drivers of haemoplasma infection and bacterial genotype sharing in a Neotropical bat community. Mol Ecol. 2020 Apr;29(8):1534–49.

7. Volokhov DV, Lock LR, Dyer KE, DeAnglis IK, Andrews BR, Simonis MC, et al. Expanded diversity of novel hemoplasmas in rare and undersampled Neotropical bats. One Health. 2023 Dec 1;17:100633.

8. Bai Y, Kosoy M, Recuenco S, Alvarez D, Moran D, Turmelle A, et al. Bartonella spp. in Bats, Guatemala. Emerg Infect Dis. 2011 Jul;17(7):1269–72.

9. Descloux E, Mediannikov O, Gourinat AC, Colot J, Chauvet M, Mermoud I, et al. Flying Fox Hemolytic Fever, Description of a New Zoonosis Caused by Candidatus Mycoplasma haemohominis. Clin Infect Dis. 2021 Oct 5;73(7):e1445–53.

10. Veikkolainen V, Vesterinen EJ, Lilley TM, Pulliainen AT. Bats as reservoir hosts of human bacterial pathogen, Bartonella mayotimonensis. Emerg Infect Dis. 2014 Jun;20(6):960–7.

11. Lilley TM, Wilson CA, Bernard RF, Willcox EV, Vesterinen EJ, Webber QMR, et al. Molecular Detection of Candidatus Bartonella mayotimonensis in North American Bats. Vector Borne Zoonotic Dis. 2017 Apr;17(4):243–6.

12. Mascarelli PE, Keel MK, Yabsley M, Last LA, Breitschwerdt EB, Maggi RG. Hemotropic mycoplasmas in little brown bats (Myotis lucifugus). Parasit Vectors. 2014 Mar 24;7:117.

13. Fleming TH, Eby P, Kunz TH, Fenton MB. Ecology of bat migration. Bat ecology.

387   2003;156:164–5.

388 14. Russell AL, Medellín RA, McCracken GF. Genetic variation and migration in the Mexican
389   free-tailed bat (Tadarida brasiliensis mexicana). Mol Ecol. 2005 Jun;14(7):2207–22.

390 15. Glass BP. Seasonal Movements of Mexican Freetail Bats Tadarida brasiliensis mexicana
391   Banded in the Great Plains. Southwest Nat. 1982;27(2):127–33.

392 16. Wiederholt R, López-Hoffman L, Cline J, Medellín RA, Cryan P, Russell A, et al. Moving
393   across the border: modeling migratory bat populations [Internet]. Vol. 4, Ecosphere. 2013.
394   p. art114. Available from: http://dx.doi.org/10.1890/es13-00023.1

395 17. LaVal RK. Observations on the Biology of Tadarida brasiliensis cynocephala in
396   Southeastern Louisiana. Am Midl Nat. 1973;89(1):112–20.

397 18. McCracken GF, Gassel MF. Genetic Structure in Migratory and Nonmigratory Populations
398   of Brazilian Free-Tailed Bats. J Mammal. 1997 May 21;78(2):348–57.

399 19. Millán J, Cevidanes A, Sacristán I, Alvarado-Rybak M, Sepúlveda G, Ramos-Mella CA, et
400   al. Detection and Characterization of Hemotropic Mycoplasmas in Bats in Chile. J Wildl
401   Dis. 2019 Oct;55(4):977–81.

402 20. Cicuttin GL, De Salvo MN, La Rosa I, Dohmen FEG. Neorickettsia risticii, Rickettsia sp.
403   and Bartonella sp. in Tadarida brasiliensis bats from Buenos Aires, Argentina. Comp
404   Immunol Microbiol Infect Dis. 2017 Jun;52:1–5.

405 21. Stepanian PM, Wainwright CE. Ongoing changes in migration phenology and winter
406   residency at Bracken Bat Cave. Glob Chang Biol. 2018 Jul;24(7):3266–75.

407 22. Ganow KB, Caire W, Matlack RS. Use of thermal imaging to estimate the population sizes
408   of Brazilian free-tailed bat, Tadarida brasiliensis, maternity roosts in Oklahoma. swna. 2015
409   Mar;60(1):90–6.

410 23. Morgan CN, Ammerman LK, Demere KD, Doty JB, Nakazawa YJ, Mauldin MR. Field
411   Identification Key and Guide for Bats of the United States of America. Occas Pap Tex Tech
412   Univ Mus [Internet]. 2019 Jan 17;360. Available from:
413   https://www.ncbi.nlm.nih.gov/pubmed/31148880

414 24. Volokhov DV, Becker DJ, Bergner LM, Camus MS, Orton RJ, Chizhikov VE, et al. Novel
415   hemotropic mycoplasmas are widespread and genetically diverse in vampire bats.
416   Epidemiol Infect. 2017 Nov;145(15):3154–67.

417 25. Volokhov DV, Norris T, Rios C, Davidson MK, Messick JB, Gulland FM, et al. Novel
418   hemotrophic mycoplasma identified in naturally infected California sea lions (Zalophus
419   californianus). Vet Microbiol. 2011 Apr 21;149(1-2):262–8.

420 26. Becker DJ, Bergner LM, Bentz AB, Orton RJ, Altizer S, Streicker DG. Genetic diversity,
421   infection prevalence, and possible transmission routes of Bartonella spp. in vampire bats.

422   PLoS Negl Trop Dis. 2018 Sep;12(9):e0006786.

423   27. Kosmidis I, Kenne Pagui EC, Sartori N. Mean and median bias reduction in generalized
424   linear models. Stat Comput. 2020 Feb 1;30(1):43–59.

425   28. Lemoine F, Correia D, Lefort V, Doppelt-Azeroual O, Mareuil F, Cohen-Boulakia S, et al.
426   NGPhylogeny.fr: new generation phylogenetic services for non-specialists. Nucleic Acids
427   Res. 2019 Jul 2;47(W1):W260–5.

428   29. La Scola B, Zeaiter Z, Khamis A, Raoult D. Gene-sequence-based criteria for species
429   definition in bacteriology: the Bartonella paradigm. Trends Microbiol. 2003 Jul;11(7):318–
430   21.

431   30. Volokhov DV, Simonyan V, Davidson MK, Chizhikov VE. RNA polymerase beta subunit
432   (rpoB) gene and the 16S--23S rRNA intergenic transcribed spacer region (ITS) as
433   complementary molecular markers in addition to the 16S rRNA gene for phylogenetic
434   analysis and identification of the species of the family Mycoplasmataceae. Mol Phylogenet
435   Evol. 2012;62(1):515–28.

436   31. Legendre P, Legendre L. Numerical Ecology. 3rd English ed. Development in
437   Environmental Modelling Vol. 24. Elsevier Science BV, Amsterdam (990 pp.); 2012.

438   32. Sacristán I, Acuña F, Aguilar E, García S, López MJ, Cevidanes A, et al. Assessing cross-
439   species transmission of hemoplasmas at the wild-domestic felid interface in Chile using
440   genetic and landscape variables analysis. Sci Rep. 2019 Nov 14;9(1):16816.

441   33. Hornok S, Földvári G, Rigó K, Meli ML, Gönczi E, Répási A, et al. Synanthropic rodents
442   and their ectoparasites as carriers of a novel haemoplasma and vector-borne, zoonotic
443   pathogens indoors. Parasit Vectors. 2015 Jan 15;8:27.

444   34. Mc Neely DL, Caire W. The fishes of traders creek in northwestern Oklahoma [Internet].
445   [cited 2024 Feb 4]. Available from: http://nanfa.org/education/sll/traderscreek.pdf

446   35. Moskaluk AE, Stuckey MJ, Jaffe DA, Kasten RW, Aguilar-Setién A, Olave-Leyva JI, et al.
447   Molecular Detection of Bartonella Species in Blood-Feeding Bat Flies from Mexico. Vector
448   Borne Zoonotic Dis. 2018 May;18(5):258–65.

449   36. Shiyi W, Cui N, Lv Z, Wang N, Liu G, Zhao S, et al. Molecular detection of vector-borne
450   pathogens in red foxes (Vulpes vulpes) in China [Internet]. Research Square. 2023 [cited
451   2024 Jan 7]. Available from: https://www.researchsquare.com/article/rs-3573995/latest

452   37. Eremeeva ME, Gerns HL, Lydy SL, Goo JS, Ryan ET, Mathew SS, et al. Bacteremia, fever,
453   and splenomegaly caused by a newly recognized bartonella species. N Engl J Med. 2007
454   Jun 7;356(23):2381–7.

455   38. Birtles RJ, Raoult D. Comparison of partial citrate synthase gene (gltA) sequences for
456   phylogenetic analysis of Bartonella species. Int J Syst Bacteriol. 1996 Oct;46(4):891–7.

39.   Lin EY, Tsigrelis C, Baddour LM, Lepidi H, Rolain JM, Patel R, et al. Candidatus
Bartonella mayotimonensis and endocarditis. Emerg Infect Dis. 2010 Mar;16(3):500–3.

40.   Urushadze L, Bai Y, Osikowicz L, McKee C, Sidamonidze K, Putkaradze D, et al.
Prevalence, diversity, and host associations of Bartonella strains in bats from Georgia
(Caucasus). PLoS Negl Trop Dis. 2017 Apr 11;11(4):e0005428.

41.   McKee CD, Krawczyk A, Sándor AD, Gorfol T, Foldvari M, Foldvari G, et al. Host
phylogeny, geographic overlap, and roost sharing shape parasite communities in European
bats. 2019; Available from:
https://repository.ubn.ru.nl/bitstream/handle/2066/203688/203688.pdf

42.   Castle KT, Weller TJ, Cryan PM, Hein CD, Schirmacher MR. Using sutures to attach
miniature tracking tags to small bats for multimonth movement and behavioral studies. Ecol
Evol. 2015 Jul;5(14):2980–9.

43.   Emmenegger T, Bensch S, Hahn S, Kishkinev D, Procházka P, Zehtindjiev P, et al. Effects
of blood parasite infections on spatiotemporal migration patterns and activity budgets in a
long-distance migratory passerine. Ecol Evol. 2021 Jan;11(2):753–62.

44.   Nichols MD, Lord WD, Haynie ML, Brennan RE, Jackson VL, Monterroso WS.
Trypanosoma cruzi in a Mexican Free-Tailed Bat (Tadarida brasiliensis) in Oklahoma,
USA. J Wildl Dis. 2019 Apr 9;55(2):444–8.

45.   Ochman H, Elwyn S, Moran NA. Calibrating bacterial evolution. Proc Natl Acad Sci U S
A. 1999 Oct 26;96(22):12638–43.

46.   Mollentze N, Babayan SA, Streicker DG. Identifying and prioritizing potential human-
infecting viruses from their genome sequences. PLoS Biol. 2021 Sep;19(9):e3001390.

47.   Vicente-Santos A, Lock L, Allira M, Dyer K, Dunsmore A, Tu W, et al. Serum proteomics
reveals a tolerant immune phenotype across multiple pathogen taxa in wild vampire bats.
Front Immunol [Internet]. 2023;14. Available from:
https://www.frontiersin.org/articles/10.3389/fimmu.2023.1281732

48.   Bagstad KJ, Wiederholt R. Tourism Values for Mexican Free-Tailed Bat Viewing. Hum
Dimensions Wildl. 2013 Jul 1;18(4):307–11.

49.   Cleveland CJ, Betke M, Federico P, Frank JD, Hallam TG, Horn J, et al. Economic value of
the pest control service provided by Brazilian free-tailed bats in south-central Texas. Front
Ecol Environ. 2006 Jun;4(5):238–43.

50.   Rocha R, Aziz SA, Brook CE, Carvalho WD. Bat conservation and zoonotic disease risk: a
research agenda to prevent misguided persecution in the aftermath of COVID-19. Animal
[Internet]. 2020; Available from: https://www.repository.utl.pt/handle/10400.5/20334