1 **Geographically widespread and novel hemotropic mycoplasmas and bartonellae in**
2 **Mexican free-tailed bats and sympatric North American bat species**
3

4 Daniel J. Becker[1*], Kristin E. Dyer[1], Lauren R. Lock[1], Beckett L. Olbrys[1], Shawn A. Pladas[2],
5 Anushka A. Sukhadia[1], Bret Demory[1], Juliana Maria Nunes Batista[1,3], Micaela Pineda[2], Nancy
6 B. Simmons[4], Amanda M. Adams[5], Winifred F. Frick[5,6], M. Teague O'Mara[2,5,7,8], Dmitriy V.
7 Volokhov[9]
8
[1] School of Biological Sciences, University of Oklahoma, Norman, OK, USA
[2] Department of Biological Sciences, Southeastern Louisiana University, Hammond, LA, USA
[3] Department of Preventive Veterinary Medicine and Animal Health, School of Veterinary
Medicine and Animal Science, University of São Paulo, São Paulo, Brazil
[4] Department of Mammalogy, Division of Vertebrate Zoology, American Museum of Natural
History, New York, NY, USA
[5] Bat Conservation International, Austin, TX, USA
[6] Department of Ecology and Evolutionary Biology, University of California Santa Cruz, Santa
Cruz, CA, USA
[7] Department of Migration, Max Planck Institute of Animal Behavior, Radolfzell, Germany
[8] Smithsonian Tropical Research Institute, Panama City, Panama
[9] Center for Biologics Evaluation and Research, U.S. Food and Drug Administration, Silver
Spring, MD, USA
*Corresponding author, danbeck@ou.edu

Keywords: hemoplasmas, *Tadarida brasiliensis*, migration, One Health
Running head: Bacterial pathogens in Mexican free-tailed bats

Abstract

Bacterial pathogens remain poorly characterized in bats, especially in North America. We
describe novel (and in some cases panmictic) hemoplasmas (10.5% positivity) and bartonellae
(25.5% positivity) across three colonies of Mexican free-tailed bats (*Tadarida brasiliensis*), a
partially migratory species that can seasonally travel hundreds of kilometers. Molecular analyses
identified three novel *Candidatus* hemoplasma species most similar to another novel *Candidatus*
species in Neotropical molossid bats. We also detected novel hemoplasmas in sympatric cave
myotis (*Myotis velifer*) and pallid bats (*Antrozous pallidus*), with sequences in the latter 96.5%
related to *C.* Mycoplasma haemohominis. We identified nine *Bartonella* genogroups, including
those in cave myotis with 96.7% similarity to *C.* Bartonella mayotimonensis. We also detected
Bartonella rochalimae in migratory Mexican free-tailed bats, representing the first report of this
human pathogen in the Chiroptera. The seasonality and diversity of these bacteria observed here
suggest that additional longitudinal, genomic, and immunological studies in bats are warranted.

40
41
42

Introduction

Bats have been intensively sampled for viral pathogens, with species in this mammalian order hosting multiple viruses with high virulence in humans (1). However, bats remain understudied for bacterial pathogens, which can be significant for their impacts on both human health and bat morbidity and even mortality (2, 3). Hemotropic mycoplasmas (hemoplasmas) and bartonellae are facultative intracellular bacteria of special interest in bats, given their high prevalence and substantial genetic diversity (4, 5). For example, sampling of Neotropical bat communities has identified many common and co-circulating genotypes of these bacteria (6–8). Surveys in Oceania and Europe have also supported plausible zoonotic transmission of these bacteria from bats to humans, including *Candidatus* Mycoplasma haemohominis and *C.* Bartonella mayotimonensis (9, 10). Greater characterization of these bacteria across global bat diversity (over 1,470 species) is therefore warranted to inform infection risks for both bats and humans, although little surveillance has thus far been conducted in North American bats (11, 12).

Although flight enables high mobility of bats, relatively few bat species undertake long-distance migrations (e.g., between maternity and wintering grounds) (13). In North America, Mexican free-tailed bats (*Tadarida brasiliensis*) display highly variable migratory strategies (14). The southwestern United States contains both non-migratory and migratory populations, with some individuals traveling hundreds to over 1,000 kilometers between wintering grounds in Mexico to northern maternity colonies in Oklahoma, Kansas, and Colorado (14–16). Other colonies across the species range include year-round residents (17, 18). This variation in migratory behavior could shape patterns of infection, including the seasonal dispersal of bacterial pathogens across landscapes to naïve hosts. Hemoplasmas have not yet been detected in this bat species (4, 19), and bartonellae have only been minimally described in the southernmost part of the bat species' geographic range (i.e., Chile and Argentina) (20).

Here, we conducted an initial characterization of hemoplasmas and bartonellae in Mexican free-tailed bats across multiple populations and seasons. Our goals were to identify novel pathogens in this bat species and to test for differences in prevalence among colonies that differ in migratory strategy and across the bat annual cycle. We also tested whether pathogen lineages were unique to each geography or if migration may facilitate panmixia. Lastly, we used this opportunity to perform a pilot characterization of these pathogens in sympatric bat species.

Material and Methods

Wild bat sampling

We sampled three North American colonies of Mexican free-tailed bats in 2021 and 2022 to compare infections among migratory strategies and provide an initial assessment of pathogen seasonality. We sampled non-migratory individuals in southeastern Louisiana (17), focusing on a colony in Pine Grove of approximately 1,000 bats, in the non-breeding season (October 2021, *n*=5) and maternity season (July 2022, *n*=10). We also sampled the partially migratory population of Bracken Cave near San Antonio, Texas, which hosts tens of millions of this bat species in the maternity season and declines to approximately 10,000 bats in winter (16, 21). We sampled the maternity season (August 2021, *n*=20) and mid-to-late winter (December 2021 and March 2022; *n*=9 and *n*=19). We also sampled a fully migratory colony at the Selman Bat Cave near Freedom, Oklahoma, where this maternity roost holds up to 100,000 bats during summer and is empty in winter (16, 22). We sampled bats monthly, from April to September 2022, spanning spring arrival, the maternity season, and fall migration (*n*=146). In the same Oklahoma

site, we also sampled four cave myotis (*Myotis velifer*), one hoary bat (*Lasiurus cinereus*), one
Townsend's big-eared bat (*Corynorhinus townsendii*), and two pallid bats (*Antrozous pallidus*).
Bats were captured with hand nets and mist nets while emerging from or returning to
roosts and placed in individual cloth bags. Bats were identified to species by morphology and
identified by sex, reproductive status, and age (23). Blood (<1% body mass) was sampled by
lancing the propatagial vein using 27G and 30G needles and collected with heparinized
capillaries. Blood was preserved on Whatman FTA cards and held at room temperature until
−20°C storage at the University of Oklahoma (OU). Sampling was approved by the Institutional
Animal Care and Use Committees of OU (2022-0198) and Southeastern Louisiana University
(0064), with permits from the Texas Parks and Wildlife Department (SPR-0521-063), Louisiana
Department of Wildlife and Fisheries (WDP-21-101), and Oklahoma Department of Wildlife
Conservation (ODWC, 10567389). All bats were released after sampling at the capture site.
100
Molecular diagnostics
We extracted genomic DNA from blood using QIAamp DNA Investigator Kits (Qiagen). To
determine hemoplasma presence, we used PCR targeting the partial 16S and 23S rRNA genes
(Table S1; (6, 7, 24, 25), with amplicons purified and sequenced at Psomagen. For DNA samples
positive for the 16S or 23S rRNA genes, we also attempted to amplify the partial *rpoB* gene,
using primers newly designed for this study (Table S1). To determine the presence of
bartonellae, we used nested PCR targeting the partial *gltA* gene (Table S1; (26)), with amplicons
purified with Zymo kits (DNA Clean & Concentrator-5, Zymoclean Gel DNA Recovery) and
sequenced at the North Carolina State University Genomic Sciences Laboratory. We included
blank FTA card punches and ultrapure water as extraction and negative controls, respectively, in
all PCRs. Hemoplasma PCRs used *Candidatus* Mycoplasma haemozalophi as a positive control,
but we did not include positive controls for *Bartonella* spp. to reduce cross-contamination risks
from nested PCR; instead, amplicons of expected size (~300 bp) were identified during gel
electrophoresis. Sequences are available on GenBank through accessions OQ407831–50,
OR783320–23, and PQ465198 (*Mycoplasma* spp. 16S rRNA); OQ359160–75 (*Mycoplasma* spp.
23S rRNA); OQ554332–38 (*Mycoplasma* spp. *rpoB*); and PP317862–72 (*Bartonella* spp. *gltA*).
117
Statistical analysis
We analyzed infection states using generalized linear models (GLMs) or generalized additive
models (GAMs) with binary response in R. All GLMs were fit using mean bias reduction
methods with the *brglm2* package (27), whereas GAMs were fit using restricted maximum
likelihood and the *mgcv* package (28). For each of our two pathogens, we fit the following four
GLMs. The first model was fit to all data and compared the odds of infection among Louisiana,
Texas, and Oklahoma bats. The second model was limited to the early-to-mid non-breeding
season and compared odds of infection between Texas and Louisiana bats. The third model was
limited to the late overwintering period and spring arrival to compare odds of infection between
Texas and that subset of Oklahoma bats. The fourth model was limited to Texas and included
reproductive status (only females were reproductive) and month to test demographic effects and
seasonality (i.e., maternity season through late overwintering). Lastly, for our GAMs, we fit a
similar model for each pathogen to the Oklahoma data, with a cyclic cubic smooth of month to
assess a different aspect of infection seasonality (i.e., spring arrival until onset of fall migration).
132
Phylogenetic analysis

We used NCBI BLASTn to identify related mycoplasma (16S rRNA, 23S rRNA, *rpoB*) and
bartonellae sequences (*gltA*), which we aligned with our sequences and reference sequences
using MUSCLE. We used MrBayes for phylogenetic analysis, with each gene tree run for
20,000,000 generations using a GTR+I+G model. BLASTn was implemented in Geneious,
whereas MUSCLE and MrBayes were implemented using NGPhylogeny.fr (29). We delineated
genotypes of hemoplasmas and genogroups of bartonellae based on pairwise similarity among
sequences and clustering on their phylogenies, using established criteria for defining novel
bacterial lineages (6, 30). For hemoplasmas, we also used multi-loci data to propose novel
Candidatus species when the same genotype was identified in at least two samples using 16S
rRNA and one other marker (i.e., 16S rRNA and 23S rRNA, 16S rRNA and *rpoB*) (7, 31).
We conducted two tests to assess if pathogen lineages were unique to each Mexican free-
tailed bat colony, which would suggest geographically constrained transmission dynamics. First,
we used chi-squared tests with *p* values generated by a Monte Carlo procedure to quantify
associations between geography and pathogen lineage assignments. Next, for any lineages
identified in multiple colonies with sufficient sample size, we derived matrices of spatial and
phylogenetic distance among sequenced PCR-positive samples and used Mantel tests with the
vegan R package to assess isolation by distance (32). These tests used 1,000 randomizations.
151
Results
Migratory and seasonal effects on bat bacterial infection
We detected hemoplasmas in 21 of 209 Mexican free-tailed bats when targeting the partial 16S
or 23S rRNA genes (10.5%, 95% CI: 7.1−15.4%). Sequencing of the 23S rRNA gene showed
three other bats (two from Texas and one from Oklahoma) had non-hemotropic mycoplasmas
(OQ359169–70, OQ359174) most related to *Mycoplasma muris* (97.5% sequence identity). We
detected bartonellae in 53 of 208 tested bats (25.5%, 95% CI: 20–31.8%). Only six Mexican
free-tailed bats were coinfected by bartonellae and any mycoplasmas (2.9%, 95% CI: 1.3–6.1%).
Hemoplasmas were detected in all three Mexican free-tailed bat colonies, while bartonellae were
only detected in the Texas and Oklahoma colonies. PCR positivity data are fully available in the
Pathogen Harmonized Observatory (PHAROS): https://pharos.viralemergence.org/ (33).
Across all Mexican free-tailed bats (model 1), the odds of infection differed by colony for
bartonellae ($\chi^2 = 9.72$, $p < 0.01$) but not hemoplasmas ($\chi^2 = 0.02$, $p = 0.99$; Figure 1). When
comparing only the resident and partially migratory populations in the non-breeding season
(model 2), Louisiana and Texas bats did not differ in the odds of either infection (hemoplasmas:
$\chi^2 = 1.21$, $p = 0.27$; bartonellae: $\chi^2 = 0$, $p = 1$). When comparing only the partially and fully
migratory populations in spring (model 3), we did not detect colony differences for hemoplasmas
($\chi^2 = 0$, $p = 1$) or bartonellae ($\chi^2 = 0.11$, $p = 0.74$). When assessing risk factors of infection in the
partially migratory Texas colony (model 4), we found no evidence of seasonal or demographic
effects for either pathogen (Table S2). However, when assessing these predictors for the fully
migratory Oklahoma colony across the full 2022 occupancy period, we identified significant
seasonality in infection (Table S3). Prevalence increased from spring arrival and peaked in the
maternity season (i.e., June 2022) for both bacteria, declining into fall migration (Figure 1).
175
Genetic diversity of Mexican free-tailed bat hemoplasmas
Sequencing of 16S rRNA amplicons revealed four hemoplasma genotypes specific to Mexican
free-tailed bats (i.e., TB1–4; Figure 2, Table 1). The TB1 genotype was 97% similar to the MR1
genotype that we earlier isolated from another molossid bat (*Molossus nigricans*) in Belize (e.g.,

MH245174) (6), and TB1 was found in all three sampled populations. In contrast, TB2 was only
detected in the Texas colony and was ~96–97% similar to hemoplasmas from carnivores and
rodents and to the Belize bat MR1 genotype (6, 34, 35). Both TB3 and TB4 were only detected
in Oklahoma and were ~99% similar to MR1. We also detected the PPM1 genotype, originally
found in *Pteronotus mesoamericanus* and *P. fulvus* in Belize (6, 7), in the Oklahoma colony.
Amplification of the 23S rRNA gene from two bats (one each from Texas and Oklahoma) also
found a genotype we initially detected in cave myotis (i.e., MV1; Figure S1) and the above
Mycoplasma muris–like (non-hemotropic) genotype. These seven mycoplasma genotypes were
not associated with geography (χ^2 = 13.07 p = 0.38; Figure S2). When considering the one
genotype observed in multiple colonies and in more than one bat per site (TB1), we also found
little support for isolation by distance with the 16S rRNA phylogeny (Mantel r = 0.35, p = 0.10).
Amplification of paired partial 23S rRNA (Figure S1) and/or *rpoB* (Figure S3) genes for
samples belonging to these 16S rRNA genotypes suggested at least three novel *Candidatus*
hemoplasma species circulate in Mexican free-tailed bats. Based on 100% identity of two *rpoB*
sequences (OQ554335–36) and 100% identity of paired 16S rRNA sequences included in the
TB1 genotype, first detected in Bracken Cave (OQ407846, OQ407848), we propose the name *C.*
Mycoplasma haematobrackenitadaridae sp. nov. Similarly, given 99.98% identity among seven
23S rRNA sequences (OQ359161–65, OQ359168, OQ359172) and 100% identity in paired 16S
rRNA sequences included in the TB3 genotype from the Selman Bat Cave (OQ407832,
OQ407836–39, OQ407843–44), we propose the name *C.* M. haematoselmanitadaridae sp. nov.
Lastly, given 100% identity of two 23S rRNA sequences (OQ359160, OQ359166) and 99.8%
identity in paired 16S rRNA sequences from the TB4 genotype (also identified from the Selman
Bat Cave; OQ407831, OQ407840), we propose the name *C.* M. haematotraderitadaridae sp. nov.
(Figures 2 and S1), based on the stream running adjacent to the bat cave (Traders Creek) (36).
Given the similarity of Mexican free-tailed bat hemoplasma 16S rRNA sequences to
those from molossid bats sampled in Belize (6), we also attempted to amplify the 23S rRNA and
rpoB genes from *Molossus nigricans* sampled in 2017 and 2018 in Belize that previously tested
positive for the MR1 and MR2 genotypes (6). We re-extracted DNA from four FTA cards and
applied the same additional PCR protocols described earlier. We obtained partial 23S rRNA and
rpoB sequences for two (OQ518943–44) and three (OQ554329–31) *M. nigricans*, respectively.
Based on 100% inter-sequence similarity of the *rpoB* sequences and high (x̄ = 99.98%) identity
of paired 16S rRNA sequences (MH245122, MH245172, MH245174), we propose the name *C.*
Mycoplasma haematomolossi sp. nov. to designate this novel hemoplasma (Figures 2 and S3).
213
Genetic diversity of Mexican free-tailed bat bartonellae
Sequencing of *gltA* amplicons next revealed at least six *Bartonella* genogroups circulating in
Mexican free-tailed bats (Figure 3, Table 1). The first genogroup was detected in both Texas and
Oklahoma, with sequences from all three sampled months across 2021 and 2022 from Bracken
Cave but only from May and June in the Selman Bat Cave. The *gltA* sequences were ≥99.6%
similar to those recently detected in Mexican free-tailed bats in Argentina (KX986617) (20),
such that we consider these sequences to all form the TB1 genogroup. The TB2–4 genogroups
were only found in the Oklahoma colony. TB2 was identical to sequences from streblid bat flies
in Mexico (e.g., ≥99.7% identity to MF988072) (37), while TB3 represents a novel genogroup
with ~92% similarity to *Bartonella vinsonii* (e.g., MK984790) (38). Likewise, TB4 was novel
and distantly related (~91%) to *Bartonella quintana* (i.e., Z70014) (39). Oklahoma bats also
harbored g*ltA* sequences with 97–100% identity to *Bartonella rochalimae* in fleas from foxes

(OQ436435) (40); these *B. rochalimae* sequences were only detected in summer months. Lastly,
Oklahoma bats had *gltA* sequences in the same clade as bartonellae originally found in vampire
bats (*Desmodus rotundus*; DR8 genogroup) (26). These six *Bartonella* genogroups were not
associated with geography ($\chi^2 = 8.20$, $p = 0.16$; Figure S4), and we also found no support for
isolation by distance for the TB1 genotype with our *glt*A phylogeny (Mantel $r = -0.04$, $p = 0.59$).
231
Bacterial infections of sympatric bat species
Opportunistic sampling of other bats in Oklahoma revealed further bacterial diversity (Table 1).
Three of the four cave myotis and both pallid bats tested positive for hemoplasmas, whereas
three of three cave myotis, both pallid bats, and the single Townsend's big-eared bat tested
positive for bartonellae; the single hoary bat tested negative for both bacterial pathogens.
For hemoplasmas (Figure 2), we identified a single novel hemoplasma genotype in each
PCR-positive species (i.e., MV1 and AP1), with 16S rRNA sequences most closely related (i.e.,
≥98% similarity) to previously detected genotypes in vesper bats from Chile and Belize (e.g.,
EF1 and MYE) (6, 19). Notably, 16S rRNA sequences of the AP1 genotype were 96.5% similar
to those of *Candidatus* Mycoplasma haemohominis (i.e., GU562823), whereas those of the MV1
genotype were only ~94% similar. 23S rRNA and *rpoB* sequences from cave myotis
(OQ359173, OQ554337) were entirely novel (<85% similarity to GenBank sequences), with the
former 100% similar to select 23S rRNA sequences identified from our Mexican free-tailed bats.
For bartonellae (Figure 3), all three positive cave myotis had their own genogroup. One
gltA sequence was ~98% similar to that first identified in little brown bats (*Myotis lucifugus*)
elsewhere in North America (i.e., KX807172), here denoted the ML1 genogroup. Another cave
myotis had the TB4 genotype. The final cave myotis *gltA* sequence clustered within a clade of
Candidatus Bartonella mayotimonensis sequences (~96%), isolated from a human endocarditis
patient in Iowa, USA (FJ376732) and other little brown bats (KX807177–9) (11, 41). For pallid
bats, one sequence also belonged to the TB4 genogroup, whereas the other formed the novel AP1
genogroup most related (~95%) to bartonellae from Natal long-fingered bats (*Miniopterus*
natalensis) and their bat flies in South Africa (e.g., MW007711 and MW007707) (42). Lastly,
the single positive Townsend's big-eared bat hosted a unique genogroup (CT1) only ~94%
similar to bartonellae found from *Myotis nigricans* in Guatemala (e.g., MN529509) (5).
256
Discussion
Hemoplasmas and bartonellae are emerging as model systems for studying bacterial infections in
bats (5, 7), but their infection dynamics and diversity remain poorly characterized, notably in
North American systems (11, 12). We here demonstrate novel diversity of hemoplasmas and
bartonellae in bats in the south-central United States, including the circulation of lineages of both
pathogens with clear infection seasonality in a migratory colony of Mexican free-tailed bats.
Such work provides the foundation for further empirical studies to elucidate the transmission
dynamics of these bacteria, their pathogenicity in bats, and their possible zoonotic risk.
Our findings suggest relatively common infection with site-specific and panmictic
bacterial lineages. Within both bacterial genera, most lineages found in Mexican free-tailed bats
were restricted to a single site, indicating spatially constrained transmission. However, we also
detected lineages in multiple sites, such as the TB1 hemoplasma genotype in Louisiana, Texas,
and Oklahoma; TB1 and MV1 hemoplasma genotypes as well as the *Mycoplasma muris*–like
genotype in Texas and Oklahoma; and TB1 *Bartonella* genogroup in Texas and Oklahoma (for
which sequences were nearly identical to those from *Tadarida brasiliensis* in Argentina) (20).

Such results may be explained by migratory connectivity in Mexican free-tailed bats, for which
regional migrations spanning hundreds to over 1,000 kilometers have been well-characterized in
North America, including between the Selman Bat Cave and Bracken Cave (14, 15, 43); this
suggests the migratory behavior of this species can enhance bacterial dispersal. However, given
the presence of these bacterial lineages in migratory and non-migratory populations (i.e.,
Louisiana) and at the extremes of the bat range (e.g., over 5,000 km between the Selman Bat
Cave and the Argentina site for the TB1 *Bartonella* genogroup), these results also suggest the
ancestral spread of these bacteria and limited selection pressure on lineages across the bat range.
Future studies are needed to identify the migratory routes of Mexican free-tailed bats,
especially for understanding the origins of possibly zoonotic bacterial lineages and the potential
for these bats to disperse infection during spring and fall migrations. Researchers could capitalize
on advances in tracking small vertebrates for long periods, such as use of absorbent sutures, to
ensure lightweight radiotags stay attached to bats for the duration of migration and winter (44).
Such work is also needed to assess if these infections negatively impact bat migration trajectory
and success, as observed for blood pathogens in migratory songbirds (45). Longitudinal studies
would also inform such analyses, as our data from one full occupancy season in the Oklahoma
colony suggest bacterial prevalence peaks in the maternity season. Additional seasonal sampling
is needed to assess how infection risk varies across the full migratory cycle, if prevalence tracks
bat population size and/or ectoparasite intensity, and whether infections are sufficiently common
in autumn to facilitate their dispersal with migration. Further genetic analyses could also inform
patterns of bat connectivity and pathogen spread. For example, the TB2 *Bartonella* genogroup
from Oklahoma Mexican free-tailed bats showed 100% identity to bartonellae from bat flies
from Morelos, Hidalgo, and Jalisco in central Mexico (37), spanning the likely wintering sites of
this bat species (14, 15). Similarly, previous analyses of *Trypanasoma cruzi* from this same
Oklahoma bat population detected lineages similar to those along the Texas–Mexico border,
further showing possible southern origins of infection and high pathogen dispersal capacity (46).
Additional molecular and immunological studies are also needed to better characterize
these novel bat bacterial pathogens and their health impacts. We identified 16S rRNA and *gltA*
sequences with moderate-to-high similarity to zoonotic pathogens such as *C.* Mycoplasma
haemohominis, *C.* Bartonella mayotimonensis, and *Bartonella rochalimae* (9, 41, 47). For the
former two pathogens, our bat sequences were ~96% similar to zoonotic lineages, likely
indicating divergence from a common ancestor at least tens of million of years ago (7, 48). *B.*
rochalimae has been found in cats and dogs (49), and our detections in Mexican free-tailed bats
indicate a broadening of host range into bats. Our sequences showed ~97–100% similarity to
those from fox fleas (40), relevant given likely flea transmission (49) and detection of fleas in the
Oklahoma bat population where these sequences were found (Dyer, personal communication).
Generation of whole genomes for our novel bat pathogens could inform their zoonotic risk, both
by better linking them to cryptic human infections (9) and by facilitating machine learning
models that predict zoonotic potential from genomic composition, as applied for viruses (50).
Other -omics analyses could also elucidate whether these bacterial infections are pathogenic in
bats themselves. In addition to assessing impacts of infection on migration outcomes as noted
above, approaches such as transcriptomics and proteomics could test if bats have a pronounced
immune response to these bacterial infections or appear largely tolerant (51). Such studies could
be especially informative when comparing immunity between migratory and non-migratory
periods, which could test whether long-distance migration may disrupt immune tolerance in bats.

Lastly, Mexican free-tailed bats and their sympatric bat species provide several important
ecosystem services, including but not limited to predating on crop pests and contributing to the
tourism economy from bat flight watching (52, 53). Understanding the prevalence, genetic
diversity, and pathogenicity of bacterial pathogens in bats can inform One Health approaches
that emphasize conservation measures to promote bat, domestic animal, and human health (54).
322

Acknowledgements
This work was supported by the National Institute of General Medical Sciences of the National
Institutes of Health (P20GM134973), Louisiana Board of Regents Support Fund, Louisiana
Biomedical Research Network, and Research Corporation for Science Advancement (RCSA).
This work was conducted via Subawards No. 28365 and 29018, part of a USDA Non-Assistance
Cooperative Agreement with RCSA Federal Award No. 58-3022-0-005. Additional support was
provided by the Edward Mallinckrodt, Jr. Foundation and University of Oklahoma (Data
Institute for Societal Challenges, Vice President for Research and Partnerships). Bat
Conservation International provided in-kind support. We thank William Caire, Melynda
Hickman, and the ODWC for site access; the Selman Living Laboratory for fieldwork support;
and Konstantin Chumakov for laboratory support. We thank one anonymous reviewer for helpful
feedback.

Tables

Table 1. *Mycoplasma* spp. (A) and *Bartonella* spp. (B) lineages identified from Louisiana, Texas, and Oklahoma bats during this study (2021–2022). Lineages are given with their host species, locations, and mean intra-genotype (mycoplasmas) or intra-genogroup (bartonellae) sequence similarity from the partial 16S rRNA, 23S rRNA, *rpoB*, or *gltA* gene sequences identified here.

	Lineage	States	Bat species	Mean intra-lineage similarity (%)
(A)	TB1[*]	LA, TX, OK	*Tadarida brasiliensis*	99.6[i], 100[iii]
	TB2[*]	TX, OK	*Tadarida brasiliensis*	100[i]
	TB3[*]	OK	*Tadarida brasiliensis*	100[i], 100[ii]
	TB4[*]	OK	*Tadarida brasiliensis*	100[i], 100[ii]
	PPM1	OK	*Tadarida brasiliensis*	NA[¶]
	MV1[*]	TX, OK	*Myotis velifer, Tadarida brasiliensis*	100[i], 100[iii]
	AP1[*]	OK	*Antrozous pallidus*	100[i]
	M. muris–like[†]	TX, OK	*Tadarida brasiliensis*	99.3[ii]
(B)	TB1	TX, OK	*Tadarida brasiliensis*	96.8[iv]
	TB2[*]	OK	*Tadarida brasiliensis*	NA[¶]
	TB3[*]	OK	*Tadarida brasiliensis*	97.2[iv]
	TB4[*]	OK, TX	*Tadarida brasiliensis, Myotis velifer, Antrozous pallidus*	97.4[iv]
	Bartonella rochalimae	OK	*Tadarida brasiliensis*	97.5[iv]
	DR8	OK	*Tadarida brasiliensis*	98.2[iv]
	ML1	OK	*Myotis velifer*	NA[¶]

C. Bartonella mayotimonensis–like	OK	*Myotis velifer*	NA[¶]
AP1[*]	OK	*Antrozous pallidus*	NA[¶]
CT1[*]	OK	*Corynorhinus townsendii*	NA[¶]

[*] Novel lineages
[†] Non-hemotropic mycoplasma
[i] 16S rRNA sequence
[ii] 23S rRNA sequence
[iii] *rpoB* sequence
[iv] *gltA* sequence
[¶] Single sequence
349

Figures

Figure 1. (A) Sampled Mexican free-tailed bat (*Tadarida brasiliensis*) colonies relative to the host distribution in the Americas. (B) Hemoplasma and *Bartonella* spp. infection prevalence across months and colonies; segments denote 95% confidence intervals using Wilson's method. For Oklahoma bats,

Figure 2. Consensus Bayesian phylogeny of the partial 16S rRNA hemoplasma sequences from this study (highlighted in bold and colored by geography; see Table 1 for genotype assignments) and reference sequences from bats and other mammals. Nodes are colored by posterior probability (nodes with less than 50% support are not shown). Hemoplasmas with *Candidatus* species names proposed here are indicated by asterisks and have paired 23S rRNA or *rpoB* sequences (see Figures S1 and S3).

369 Figure 3. Consensus Bayesian phylogeny of the partial *gltA Bartonella* spp. sequences from this
370 study (highlighted in bold and colored by geography; see Table 1 for genogroup assignments)
371 and reference sequences from bats, other mammals, and ectoparasites. Nodes are colored by
372 posterior probability (nodes with less than 50% support are not shown).

374
375
376

References

1. Letko M, Seifert SN, Olival KJ, Plowright RK, Munster VJ. 2020. Bat-borne virus diversity, spillover and emergence. Nat Rev Microbiol 18:461–471.
2. Brook CE, Dobson AP. 2015. Bats as "special"reservoirs for emerging zoonotic pathogens. Trends Microbiol.
3. Szentivanyi T, McKee C, Jones G, Foster JT. 2023. Trends in Bacterial Pathogens of Bats: Global Distribution and Knowledge Gaps. Transbound Emerg Dis 2023.
4. Millán J, Di Cataldo S, Volokhov DV, Becker DJ. 2020. Worldwide occurrence of haemoplasmas in wildlife: Insights into the patterns of infection, transmission, pathology and zoonotic potential. Transbound Emerg Dis https://doi.org/10.1111/tbed.13932.
5. McKee CD, Bai Y, Webb CT, Kosoy MY. 2021. Bats are key hosts in the radiation of mammal-associated Bartonella bacteria. Infect Genet Evol 89:104719.
6. Becker DJ, Speer KA, Brown AM, Fenton MB, Washburne AD, Altizer S, Streicker DG, Plowright RK, Chizhikov VE, Simmons NB, Volokhov DV. 2020. Ecological and evolutionary drivers of haemoplasma infection and bacterial genotype sharing in a Neotropical bat community. Mol Ecol 29:1534–1549.
7. Volokhov DV, Lock LR, Dyer KE, DeAnglis IK, Andrews BR, Simonis MC, Stockmaier S, Carter GG, Downs CJ, Fenton MB, Simmons NB, Becker DJ. 2023. Expanded diversity of novel hemoplasmas in rare and undersampled Neotropical bats. One Health 17:100633.
8. Bai Y, Kosoy M, Recuenco S, Alvarez D, Moran D, Turmelle A, Ellison J, Garcia DL, Estevez A, Lindblade K, Rupprecht C. 2011. Bartonella spp. in Bats, Guatemala. Emerg Infect Dis 17:1269–1272.
9. Descloux E, Mediannikov O, Gourinat A-C, Colot J, Chauvet M, Mermoud I, Desoutter D, Cazorla C, Klement-Frutos E, Antonini L, Levasseur A, Bossi V, Davoust B, Merlet A, Goujart M-A, Oedin M, Brescia F, Laumond S, Fournier P-E, Raoult D. 2021. Flying Fox Hemolytic Fever, Description of a New Zoonosis Caused by Candidatus Mycoplasma haemohominis. Clin Infect Dis 73:e1445–e1453.
10. Veikkolainen V, Vesterinen EJ, Lilley TM, Pulliainen AT. 2014. Bats as reservoir hosts of human bacterial pathogen, Bartonella mayotimonensis. Emerg Infect Dis 20:960–967.
11. Lilley TM, Wilson CA, Bernard RF, Willcox EV, Vesterinen EJ, Webber QMR, Kurpiers L, Prokkola JM, Ejotre I, Kurta A, Field KA, Reeder DM, Pulliainen AT. 2017. Molecular Detection of Candidatus Bartonella mayotimonensis in North American Bats. Vector Borne Zoonotic Dis 17:243–246.
12. Mascarelli PE, Keel MK, Yabsley M, Last LA, Breitschwerdt EB, Maggi RG. 2014. Hemotropic mycoplasmas in little brown bats (Myotis lucifugus). Parasit Vectors 7:117.
13. Fleming TH, Eby P, Kunz TH, Fenton MB. 2003. Ecology of bat migration. Bat ecology 156:164–165.
14. Russell AL, Medellín RA, McCracken GF. 2005. Genetic variation and migration in the Mexican free-tailed bat (Tadarida brasiliensis mexicana). Mol Ecol 14:2207–2222.
15. Glass BP. 1982. Seasonal Movements of Mexican Freetail Bats Tadarida brasiliensis mexicana Banded in the Great Plains. Southwest Nat 27:127–133.
16. Wiederholt R, López-Hoffman L, Cline J, Medellín RA, Cryan P, Russell A, McCracken G, Diffendorfer J, Semmens D. 2013. Moving across the border: modeling migratory bat populations. Ecosphere https://doi.org/10.1890/es13-00023.1.
17. LaVal RK. 1973. Observations on the Biology of Tadarida brasiliensis cynocephala in Southeastern Louisiana. Am Midl Nat 89:112–120.

18. McCracken GF, Gassel MF. 1997. Genetic Structure in Migratory and Nonmigratory
Populations of Brazilian Free-Tailed Bats. J Mammal 78:348–357.
19. Millán J, Cevidanes A, Sacristán I, Alvarado-Rybak M, Sepúlveda G, Ramos-Mella CA,
Lisón F. 2019. Detection and Characterization of Hemotropic Mycoplasmas in Bats in
Chile. J Wildl Dis 55:977–981.
20. Cicuttin GL, De Salvo MN, La Rosa I, Dohmen FEG. 2017. Neorickettsia risticii, Rickettsia
sp. and Bartonella sp. in Tadarida brasiliensis bats from Buenos Aires, Argentina. Comp
Immunol Microbiol Infect Dis 52:1–5.
21. Stepanian PM, Wainwright CE. 2018. Ongoing changes in migration phenology and winter
residency at Bracken Bat Cave. Glob Chang Biol 24:3266–3275.
22. Ganow KB, Caire W, Matlack RS. 2015. Use of thermal imaging to estimate the population
sizes of Brazilian free-tailed bat, Tadarida brasiliensis, maternity roosts in Oklahoma. swna
60:90–96.
23. Morgan CN, Ammerman LK, Demere KD, Doty JB, Nakazawa YJ, Mauldin MR. 2019.
Field Identification Key and Guide for Bats of the United States of America. Occas Pap Tex
Tech Univ Mus 360.
24. Volokhov DV, Becker DJ, Bergner LM, Camus MS, Orton RJ, Chizhikov VE, Altizer SM,
Streicker DG. 2017. Novel hemotropic mycoplasmas are widespread and genetically diverse
in vampire bats. Epidemiol Infect 145:3154–3167.
25. Volokhov DV, Norris T, Rios C, Davidson MK, Messick JB, Gulland FM, Chizhikov VE.
2011. Novel hemotrophic mycoplasma identified in naturally infected California sea lions
(Zalophus californianus). Vet Microbiol 149:262–268.
26. Becker DJ, Bergner LM, Bentz AB, Orton RJ, Altizer S, Streicker DG. 2018. Genetic
diversity, infection prevalence, and possible transmission routes of Bartonella spp. in
vampire bats. PLoS Negl Trop Dis 12:e0006786.
27. Kosmidis I, Kenne Pagui EC, Sartori N. 2020. Mean and median bias reduction in
generalized linear models. Stat Comput 30:43–59.
28. Wood SN. 2017. Generalized Additive Models: An Introduction with R, Second Edition.
CRC Press.
29. Lemoine F, Correia D, Lefort V, Doppelt-Azeroual O, Mareuil F, Cohen-Boulakia S,
Gascuel O. 2019. NGPhylogeny.fr: new generation phylogenetic services for non-
specialists. Nucleic Acids Res 47:W260–W265.
30. La Scola B, Zeaiter Z, Khamis A, Raoult D. 2003. Gene-sequence-based criteria for species
definition in bacteriology: the Bartonella paradigm. Trends Microbiol 11:318–321.
31. Volokhov DV, Simonyan V, Davidson MK, Chizhikov VE. 2012. RNA polymerase beta
subunit (rpoB) gene and the 16S--23S rRNA intergenic transcribed spacer region (ITS) as
complementary molecular markers in addition to the 16S rRNA gene for phylogenetic
analysis and identification of the species of the family Mycoplasmataceae. Mol Phylogenet
Evol 62:515–528.
32. Legendre P, Legendre L. 2012. Numerical Ecology. 3rd English ed. Development in
Environmental Modelling Vol. 24. Elsevier Science BV, Amsterdam (990 pp.).
33. Stevens T, Zimmerman R, Albery G, Becker D, Kading R, Keiser C, Khandelwal S,
Kramer-Schadt S, Krut-Landau R, McKee C, Montecino-Latorre D, O'Donoghue Z, Olson
S, Poisot T, Robertson H, Ryan S, Seifert S, Simons D, Vicente-Santos A, Wood C,
Graeden E, Carlson C. 2024. A minimum data standard for wildlife disease studies.
EcoEvoRxiv.

34.   Sacristán I, Acuña F, Aguilar E, García S, López MJ, Cevidanes A, Cabello J, Hidalgo-
Hermoso E, Johnson WE, Poulin E, Millán J, Napolitano C. 2019. Assessing cross-species
transmission of hemoplasmas at the wild-domestic felid interface in Chile using genetic and
landscape variables analysis. Sci Rep 9:16816.

35.   Hornok S, Földvári G, Rigó K, Meli ML, Gönczi E, Répási A, Farkas R, Papp I, Kontschán
J, Hofmann-Lehmann R. 2015. Synanthropic rodents and their ectoparasites as carriers of a
novel haemoplasma and vector-borne, zoonotic pathogens indoors. Parasit Vectors 8:27.

36.   Mc Neely DL, Caire W. The fishes of traders creek in northwestern Oklahoma.
http://nanfa.org/education/sll/traderscreek.pdf. Retrieved 4 February 2024.

37.   Moskaluk AE, Stuckey MJ, Jaffe DA, Kasten RW, Aguilar-Setién A, Olave-Leyva JI,
Galvez-Romero G, Obregón-Morales C, Salas-Rojas M, García-Flores MM, Aréchiga-
Ceballos N, García-Baltazar A, Chomel BB. 2018. Molecular Detection of Bartonella
Species in Blood-Feeding Bat Flies from Mexico. Vector Borne Zoonotic Dis 18:258–265.

38.   Beckmann S, Engelbrecht M, Chavez F, Rojas G. 2019. Prevalence of zoonotic Bartonella
among prairie rodents in Illinois. J Mammal 101:291–297.

39.   Birtles RJ, Raoult D. 1996. Comparison of partial citrate synthase gene (gltA) sequences for
phylogenetic analysis of Bartonella species. Int J Syst Bacteriol 46:891–897.

40.   Millán J, Sepúlveda-García P, Di Cataldo S, Canales N, Sallaberry-Pincheira N, Painean J,
Cevidanes A, Müller A. 2023. Molecular identification of Bartonella spp. and Rickettsia
felis in fox fleas, Chile. Comp Immunol Microbiol Infect Dis 96:101983.

41.   Lin EY, Tsigrelis C, Baddour LM, Lepidi H, Rolain JM, Patel R, Raoult D. 2010.
Candidatus Bartonella mayotimonensis and endocarditis. Emerg Infect Dis 16:500–503.

42.   Szentiványi T, Markotter W, Dietrich M, Clément L, Ançay L, Brun L, Genzoni E, Kearney
T, Seamark E, Estók P, Christe P, Glaizot O. 2020. Host conservation through their
parasites: molecular surveillance of vector-borne microorganisms in bats using ectoparasitic
bat flies. Parasite 27:72.

43.   Villa BR, Cockrum EL. 1962. Migration in the Guano Bat Tadarida Brasiliensis Mexicana
(Saussure). J Mammal 43:43–64.

44.   Castle KT, Weller TJ, Cryan PM, Hein CD, Schirmacher MR. 2015. Using sutures to attach
miniature tracking tags to small bats for multimonth movement and behavioral studies. Ecol
Evol 5:2980–2989.

45.   Emmenegger T, Bensch S, Hahn S, Kishkinev D, Procházka P, Zehtindjiev P, Bauer S.
2021. Effects of blood parasite infections on spatiotemporal migration patterns and activity
budgets in a long-distance migratory passerine. Ecol Evol 11:753–762.

46.   Nichols MD, Lord WD, Haynie ML, Brennan RE, Jackson VL, Monterroso WS. 2019.
Trypanosoma cruzi in a Mexican Free-Tailed Bat (Tadarida brasiliensis) in Oklahoma,
USA. J Wildl Dis 55:444–448.

47.   Eremeeva ME, Gerns HL, Lydy SL, Goo JS, Ryan ET, Mathew SS, Ferraro MJ, Holden
JM, Nicholson WL, Dasch GA, Koehler JE. 2007. Bacteremia, fever, and splenomegaly
caused by a newly recognized bartonella species. N Engl J Med 356:2381–2387.

48.   Ochman H, Elwyn S, Moran NA. 1999. Calibrating bacterial evolution. Proc Natl Acad Sci
U S A 96:12638–12643.

49.   Pérez-Martínez L, Venzal JM, González-Acuña D, Portillo A, Blanco JR, Oteo JA. 2009.
Bartonella rochalimae and other Bartonella spp. in fleas, Chile. Emerg Infect Dis 15:1150–
1152.

50.   Mollentze N, Babayan SA, Streicker DG. 2021. Identifying and prioritizing potential

human-infecting viruses from their genome sequences. PLoS Biol 19:e3001390.

51. Vicente-Santos A, Lock L, Allira M, Dyer K, Dunsmore A, Tu W, Volokhov D, Herrera C,
Lei G-S, Relich R, Janech M, Bland A, Simmons N, Neely B, Becker D. 2023. Serum
proteomics reveals a tolerant immune phenotype across multiple pathogen taxa in wild
vampire bats. Front Immunol 14.

52. Bagstad KJ, Wiederholt R. 2013. Tourism Values for Mexican Free-Tailed Bat Viewing.
Hum Dimensions Wildl 18:307–311.

53. Cleveland CJ, Betke M, Federico P, Frank JD, Hallam TG, Horn J, López JD Jr,
McCracken GF, Medellín RA, Moreno-Valdez A, Sansone CG, Westbrook JK, Kunz TH.
2006. Economic value of the pest control service provided by Brazilian free-tailed bats in
south-central Texas. Front Ecol Environ 4:238–243.

54. Rocha R, Aziz SA, Brook CE, Carvalho WD. 2020. Bat conservation and zoonotic disease
risk: a research agenda to prevent misguided persecution in the aftermath of COVID-19.
Animal.

529