Mapping Serotonergic Dynamics using Drug-Modulated Molecular

Connectivity

Ionescu TM[1], Amend M[1], Hafiz R[2], Maurer A[1], Biswal BB[2], Wehrl HF[1†], Herfert K[1†]

1 Werner Siemens Imaging Center, Department of Preclinical Imaging and Radiopharmacy,
Eberhard Karls University Tuebingen, Germany

2 Department of Biomedical Engineering, New Jersey Institute of Technology, University
Heights, Newark, New Jersey, USA

[†] contributed equally

Corresponding author:

Prof. Dr. Kristina Herfert (kristina.herfert@med.uni-tuebingen.de)

Address: Roentgenweg 13, 72076 Tuebingen, Germany,

Tel: +49 7071 29 87680

Abstract

Understanding the complex workings of the brain is one of the most significant challenges in neuroscience, providing insights into normal brain function, neurological diseases, and the effects of potential therapeutics. A major challenge in this field lies in the limitations of traditional brain imaging techniques, which often capture only fragments of the complex puzzle of brain function. Our research employs a novel approach termed "molecular connectivity" (MC), which combines the strengths of various imaging methods to provide a comprehensive view of how specific molecules, such as the serotonin transporter, interact across different brain regions and influence brain function.

This innovative technique bridges the gap between functional magnetic resonance imaging (fMRI), known for its ability to monitor brain activity by tracking blood flow, and positron emission tomography (PET), which visualizes specific molecular changes. By integrating these methods, we can better understand how drugs influence brain function. Our study focuses on the application of dynamic $[^{11}C]$DASB PET scans to map the distribution of serotonin transporters, key players in regulating mood and emotions and examines how these transporters are altered following exposure to methylenedioxymethamphetamine (MDMA), which is commonly known as ecstasy.

Through a detailed comparison of MCs with traditional measures of brain connectivity, we reveal significant patterns that closely align with physiological changes. Our results revealed clear changes in molecular connectivity after a single dose of MDMA, establishing a direct link between the effects of drugs on serotonin transporter occupancy and changes in the functional brain network.

This work offers a novel methodology for the in-depth study of brain function at the molecular level and opens new pathways for understanding how drugs modulate brain activity.

Keywords: Molecular Connectivity, PET/fMRI, Functional Connectivity, Pharmacological Imaging

Introduction

The field of neuroscience has experienced remarkable advancements over the past decade, specifically propelled by the advent of innovative imaging techniques. The simultaneous application of positron emission tomography (PET) and magnetic resonance imaging (MRI) has revolutionized our ability to assess brain function across various physiological dimensions [1, 2]. This is especially true in the context of drug effect evaluations. Combining PET with functional MRI (fMRI) has opened many investigative avenues. Techniques such as resting-state functional connectivity (rs-FC) [3] or pharmacological MRI (phMRI) [4] are now complemented by the ability of PET to quantify molecular changes, including alterations in receptor and transporter availability [5].

While fMRI provides high spatial and temporal resolution, the interpretation of its readout necessitates caution. The blood-oxygen-level-dependent (BOLD) signal in fMRI indirectly captures neuronal changes through neurovascular coupling [6], revealing only the hemodynamic consequences of molecular-level drug effects. The integration of simultaneous PET acquisition can bridge this interpretative gap by offering essential molecular insights, particularly regarding transporter or receptor alterations. Typically, PET, used independently or simultaneously with fMRI, has been used mainly in pharmacological studies to illustrate quantitative shifts in neuroreceptor or transporter availability [7]. Several studies have also explored the interregional coherence of PET tracer signals [8-10], an approach akin to fMRI-derived rs-FC. While subject-level metabolic connectivity using [18F]FDG-PET has been established through the temporal correlation of regional PET signals [1, 8], studies employing transporter or receptor tracers have focused predominantly on interregional binding coherence across subjects using static scans [9, 10]. The concept and feasibility of molecular connectivity (MC) [11, 12] through the temporal correlation of the dynamic binding potentials of transporter or receptor tracers has yet to be explored.

In our study, we investigated the feasibility of deriving MCs from dynamic [11C]DASB-PET scans acquired simultaneously with fMRI in rats. We divided the rats into two cohorts, a baseline group and a pharmacological application group, which were exposed to 3,4-methylenedeoxymetamphetamine (MDMA). The baseline cohort assessed the feasibility of the novel methodology, contrasting it with traditional fMRI-

derived rs-FC with a specific focus on temporal stability. We postulated that dynamic
[^{11}C]DASB PET temporal fluctuations could be harnessed for connectivity data like
that used for hemodynamic rs-FC using seed-based and independent component
analysis (ICA).

The second cohort, subjected to an MDMA challenge, allowed us to evaluate the
utility of our novel approach. We aimed to outline the effects of MDMA by integrating
the innovative MC concept with established analysis techniques. Our primary
objective in this research was to elucidate the potential of PET-derived MCs in
conjunction with simultaneous PET/fMRI, exploring the avenues this methodology
could open for future diagnostic and drug development studies.

Materials and Methods

88 Our study reevaluates two datasets previously published by our group [13, 14] to
89 explore FC and MC simultaneously at baseline [14] and following MDMA
90 administration [13]. Please refer to these earlier publications for detailed descriptions
91 of the animal handling methods, experimental setups, and data acquisition
92 procedures.

Animals

A total of 41 male Lewis rats were obtained from Charles River Laboratories
(Sulzfeld, Germany). Thirty rats (354 ± 37 g, corresponding to 11 weeks of age)
underwent baseline [^{11}C]DASB PET/fMRI scans without any pharmacological
intervention, whereas 11 rats (365 ± 19 g, corresponding to 11 weeks of age)
underwent [^{11}C]DASB PET/fMRI scans, including an acute MDMA challenge. All
experiments were in compliance with German federal regulations for experimental
animals and received approval from the Regierungspräsidium Tübingen.

Simultaneous PET/fMRI Experiments

The rats were subjected to simultaneous PET/fMRI experiments involving 1.3%
isoflurane anesthesia, tail vein catheterization, positioning on a temperature-
controlled bed, and monitoring of vital signs. The scans were performed using a 7T
small-animal ClinScan MRI scanner (Bruker Biospin, Ettlingen, Germany) with a
custom-developed PET insert **[15]**. For [^{11}C]DASB PET, we employed a bolus plus
constant infusion protocol (k_{bol} = 38.7 min) and reconstructed the scans in 1-minute

time frames. The scanning protocol and sequence parameters are outlined in detail in

our previous publication **[13]**. The MDMA cohort received a pharmacological MDMA

challenge of 3.2 mg/kg intravenously 40 minutes after tracer injection.

Data Preprocessing and Analysis

Data preprocessing followed established protocols, including steps such as

realignment, mask creation, coregistration, spatial normalization, signal cleaning, and

spatial smoothing, as detailed in our previous work [13]. For the MDMA dataset, PET

scans were analyzed for early and late effects post-challenge using the general

linear model (GLM) available in SPM. For both datasets, the baseline was defined as

30 to 40 minutes after the start of the scan. For the fMRI data, a first-level analysis

was applied to the individual scans without a high-pass filter (the filter was set to

"Inf"). Statistical parametric maps were generated after GLM parameter estimation

using contrast vectors. The group-level analysis involved a one-sample t-test on the

subject-level statistical parametric maps ($p < 0.05$, one-sided, familywise error/FWE

adjusted).

Static PET scans were generated by summing dynamic frames over defined periods

for 10-minute periods after the MDMA challenge (50-60 minutes to investigate early

effects and 70-80 minutes to investigate late MDMA effects). Two-sample t-maps

were calculated between the normalized [^{11}C]DASB uptakes of (1) the baseline scan

period and the early effect period and (2) the early effect period and the late effect

period ($p < 0.05$, FWE-adjusted).

All group-level t-maps were subjected to voxelwise signal quantification to determine

the regional contributions of 48 regions selected according to the Schiffer atlas [16].

The average t-scores and standard deviations of all voxels were calculated.

Functional Connectivity Analysis

FC was determined using a seed-based analysis approach. The mean time series of

the preprocessed BOLD-fMRI signals for each dataset across all regions (refer to

Supplementary Table 1 for the list of regions) were extracted using the SPM toolbox

Marseille Boîte À Région d'Intérêt (MarsBaR). Pairwise Pearson's correlation

coefficients were calculated between each pair of mean regional time series for every

dataset. The Pearson's r coefficients were converted into z values using Fischer's

transformation for group-level analysis. Fischer's z-transformed correlation
coefficients were then used to generate mean correlation matrices for both cohorts
[17].

Molecular connectivity analysis

The mean [¹¹C]DASB signal from the preprocessed PET datasets was extracted from
the designed regions, including the 48 regions used for fMRI data analysis and the
cerebellum using MarsBaR. Binding potentials were calculated framewise for all
dynamic PET scans using the DVR-1 (Equation 1) to generate regional BP_{ND} values
with the cerebellar gray matter as a reference region, which our earlier studies
demonstrated to be the most appropriate for this tracer in rats [18, 19]:

$$BP_{ND} = \frac{V_T - V_{ND}}{V_{ND}} = \frac{V_T}{V_{ND}} - 1 = DVR - 1 \text{ (Eq. 1)}$$

where:

• BP_{ND} is the binding potential
• V_T is the total volume of distribution
• V_{ND} is the volume of distribution in a reference tissue
• DVR is the relative volume of distribution

To calculate the MC, we discarded the first 20 minutes of every scan, which were
dominated by perfusion effects, and applied a detrending approach to the remaining
60 minutes to obtain temporally stable values (for further details, please refer to the
Supplementary Methods and Supplementary Figure 1). The BP_{ND} time courses were
then used to calculate the MC as described above for fMRI: ROI-to-ROI subject-level
correlation matrices between all regional time courses were generated, and z-
transformed correlation coefficients were used to calculate mean correlation
matrices.

Independent Component Analysis

Group ICA (GIFT toolbox, MIALAB, University of New Mexico, Albuquerque, NM,
USA) was used for ICA in the baseline group. The fMRI and PET preprocessed
datasets were investigated between 30 and 80 minutes after the start of data
acquisition. For fMRI, we selected ten independent components, while we started
with two components for PET and increased the number to ten components to

thoroughly dissect the varying components within the signal. The components were
thresholded at a z-value $\geq$ 1.96 (p-value $\leq$ 0.05) [20], and average z-scores and
standard deviations were calculated for each component. The physiological
significance of the components was further explored by contrasting them with the
regional [^{11}C]DASB changes induced by MDMA. Accordingly, the z scores of the
independent components generated in the baseline cohort were correlated with the
early and late regional [^{11}C]DASB changes induced by MDMA, measured using t
scores.

Results

Comparability of MC and FC in spatial contexts and over time

We first aimed to evaluate whether MCs align spatially with FC, possess similar
graph theory properties and provide consistent temporal readouts throughout the
scan duration in the baseline group (Figure 1).

Connectome comparison

Figure 1: Evaluation of the seed-based MC. (A) Correlation matrix indicating whole-brain FC (beneath the diagonal) and MC (above the diagonal). Correlations not significant with multiple comparison corrections were set to zero (p < 0.05, FWE correction). **(B)** Scatter plot and correlation between MC and FC edges. **(C)** Small-world coefficients for all subjects and group-level one-sample t-test against the value of 1 (SW > 1 indicates small-world properties). Comparison of **(D)** FC and **(G)** MC early (20-40 minutes after the start of the scan, below the diagonal) and late (60-80 minutes after the start of the scan, above the diagonal). The similarities of early and late readouts were quantified

for both **(E)** FC and **(H)** MC. The temporal stability of both **(F)** FC and **(I)** MC was evaluated using a sliding window approach, including 20-minute windows between 20 and 80 minutes after the start of the scan. Abbreviations: FC = fMRI-derived hemodynamic functional connectivity, MC = [^{11}C]DASB PET-derived molecular connectivity.

We found a moderate but significant correlation between the edge-level MC and FC ($r = 0.51$, $p < 0.001$; Figure 1A and B). Furthermore, both connectomes revealed small-world properties at the group level, with coefficients higher than 1 (Figure 1C). At the subject level, three molecular and four functional connectomes fell below the threshold of 1 for the small-world coefficient. Significant consistency was observed between early and late scan-derived connectomes (Figure 1D for FC, G for MC), with FC having a slight edge (Figure 1E, $r = 0.96$) over MC (Figure 1H, $r = 0.8$). While both FC and MC maintained steady correlation intensities, there was a negligible decline over the scan duration (Figure 1F and 1I).

Deciphering the spatial characteristics of FC and MC using ICA

After establishing the feasibility of obtaining temporally stable readouts using the ROI-to-ROI approach, we employed a data-driven approach using ICA to compare the spatial characteristics of FC and MC (Figure 2).

Figure 2: Group independent component analysis for FC and MC. (A) ICA was performed over 10 components for fMRI. **(B)** ICA performed over 2 components for [^{11}C]DASB PET and regional quantification of the two derived components. The ICA was repeated over 10 components. Four and three components showed good overlap with the two components defined above. All the components were thresholded at z > 1.96 (p ≤ 0.05). Abbreviations: FC = fMRI-derived hemodynamic functional connectivity, MC = [^{11}C]DASB PET-derived molecular connectivity.

We extracted ten group ICs from the fMRI data (Figure 2A), revealing known
canonical resting-state networks, such as the posterior default-mode-like network
(IC1-5, red), sensorimotor networks (IC1-5, green and purple), the anterior default-
mode-like network (IC6-10, yellow) and the visual network (IC6-10, red). Given the
unpredictability of the number of ICs suitable for MC IC extraction, we started with
two components (Figure 2B). IC1 (orange) comprised both subcortical and cortical
anterior brain regions, including the nucleus accumbens, amygdala, cingulate cortex,
caudate putamen, orbitofrontal cortex and medial prefrontal cortex, whereas IC2
(blue) primarily received contributions from deeper posterior areas, such as the
midbrain, thalamus, hypothalamus, periaqueductal gray cortex and medulla.
Interestingly, when we extracted 10 independent components to mimic the number of
components used for the FC data, the initial anterior component split into four
different ICs, and the initial posterior IC split into three different ICs. Relatively clear
spatial segregation can be observed for the newly formed ICs, for instance, the three
posterior components extracted from specific regions (green, mainly from the
medulla; red, from the hypothalamus; and part of the midbrain, blue, from the
midbrain and thalamus).

MDMA-induced changes in ICA-derived molecular connectivity

Next, we aimed to explore the relationship between molecular changes in SERT
availability and the molecular connectome derived from the ICA induced by acute
MDMA administration (Figure 3).

Figure 3: Comparison of MDMA-induced [¹¹C]DASB alterations. (A) Left panel: Dynamic binding
potentials of regions comprising the SERT subcortical network, defined by IC 1 in the validation cohort.
Right panel: Dynamic binding potentials of regions comprising the SERT salience network, defined by
IC 2 in the validation cohort (continuous lines indicate the means, and dotted lines indicate standard
deviations). **(B)** Overlap between independent components extracted from the validation cohort (IC 1 =
SERT subcortical network, IC 2 = SERT salience network;) and the early and late effects of MDMA.
(C) Pairwise correlations between regional z scores of the ICs extracted from the validation cohort and
regional t scores of early and late MDMA effects. (*** indicates p < 0.001, ns = nonsignificant).

Abbreviations: SERT = serotonin transporter, ICA = independent component analysis; for
abbreviations of regions, please refer to the *Supplementary Information*).

Two ICs extracted from the MC showed good overlap with regions associated with
the SERT salience network (IC1) and the SERT subcortical network, comprising
regions with intrinsically high SERT densities (IC2). Therefore, we defined the
regions contributing strongly to IC2 as the SERT salience network (CPu, Cg, NAc,
Amyg, Ins, mPFC) and those with strong signals in IC1 as the SERT subcortical
network (VTA, Th, MB, PAG, Hypo). Interestingly, MDMA induced immediate strong
decreases in all SERT subcortical network regions, and salience areas exhibited a
delay of approximately 10 minutes (Figure 3A). Voxel-level analysis revealed clear
spatial overlaps between early MDMA-responsive regions and those from the
posterior IC, with delayed regions mirroring the anterior IC reminiscent of the SERT
salience network (Figure 3B). To quantify the striking spatial similarity between the
baseline independent components and the spatiotemporal characteristics of
[11C]DASB changes after MDMA exposure, we found highly significant correlations
between the z scores of the posterior and anterior ICs and the t scores of late and
early MDMA effects on [^{11}C]DASB alterations ($p < 0.0001$, Figure 3C).

Finally, we investigated the relationships between SERT availability changes and MC
reductions following acute MDMA challenge (Figure 4).

Figure 4: Comparison of MC and BP$_{ND}$ changes following MDMA. A Reductions in BP$_{ND}$ following
MDMA (orange) and MC strength (blue) were compared. **B** The correlation coefficient of the regional T
scores was low ($r = 0.29$).

We found that the decreases in MC and [^{11}C]DASB BP$_{ND}$ following MDMA application
(Figure 4A) showed only a low correlation ($r = 0.29$, Figure 4B). While decreases in

SERT availability exhibited a strong anterior-posterior gradient, which was most
pronounced in areas with high SERT availability, such as the MB, VTA, Pons or PAG,
the MC encompassed regions across the brain to similar extents. Specifically, while
the significance of [^{11}C]DASB reductions in Ins was very low across the investigated
regions, Ins presented the greatest effects among regions for MC. In contrast, strong
SERT occupancy effects in IC and SC did not translate into very prominent
reductions in the respective global MCs of the two regions.

MDMA-induced changes in seed-based molecular connectivity

Next, we performed a seed-based analysis to compare changes in FC and MC after
an MDMA pharmacological challenge on the salience network and regions with high
SERT binding (Figure 5).

Figure 5: MDMA effects on seed-based FC and MC of the salience and subcortical networks.
(A) FC and **(C)** MC brain networks depicting the edge and node strengths of the salience network and subcortical network at baseline (20-40 min after the start of the scan) and after MDMA (60-80 min after the start of the scan). **(B)** FC and **(D)** MC time-resolved salience and subcortical network strengths computed by sliding windows. Individual values are provided in the bar graphs for the baseline and final post-MDMA periods. Asterisks indicate significant (p < 0.05, FDR-corrected) changes from baseline (time point zero, corresponding to 20–40 minutes after the start of the scan).

We observed a decrease in SN FC (16% at the end of the scan, p = 0.02, did not survive FDR correction) and almost constant FC of the subcortical network (<5% decrease, p = 0.79) following MDMA, as shown in Figure 5A at the edge and node levels and Figure 5C at the network level. For MCs, we observed profound reductions

in the subcortical network (Figure 5C and D, p = 0.009), emphasizing the acute and
spatially specific effect of MDMA on MCs.

Discussion

The mammalian brain operates on diverse physiological, spatial and temporal scales.
FC via BOLD-fMRI offers insights into coherent functional brain networks, but its
complexity and indirect link to neural activity highlight the need for more direct
methodologies. In this context, the concept of MC using [11C]DASB PET, as
introduced in our study, provides a more direct and complementary perspective on
brain organization and its response to external stimuli, such as MDMA.

Physiological basis

Our findings suggest that [11C]DASB binding reflects the interplay of serotonin levels
and SERT dynamics [21]. In support of the competition model, evidence indicates
that endogenous serotonin competes with tracers for binding sites, affecting tracer
binding [22]. However, contrasting results from various studies highlight the
complexity of this interaction [23] [24] [25]. The internalization model suggests that
serotonin levels influence SERT internalization, impacting [11C]DASB binding [21, 26,
27]. While supporting evidence, further exploration is needed to fully understand
these dynamics, especially during the resting state [28]. Some models on this topic
have proposed a regulatory function of the raphé nuclei in maintaining serotonin
fluctuations over several temporal scales at rest [29]. Remarkably, fast microdialysis
has resolved multiple spontaneous surges of up to 1500-fold of the basal serotonin
occurring during 30-minute intervals [30]. Additionally, the same study indicated that
SERT expression is essential for spontaneous surges and that reduced SERT
drastically decreases serotonin spiking. Thus, it is feasible that the correlated
temporal fluctuations captured by dynamic [11C]DASB at least partly reflect the role of
the serotonergic system in the resting activity of the brain. In addition, SERT
regulation occurs over multiple time scales, ranging from milliseconds to hours,
depending on the mechanism involved [31]. Rapid changes in SERT surface
expression can be mediated by protein-protein interactions or posttranslational
modifications [32, 33], such as phosphorylation, which occur on a timescale of
milliseconds to minutes. These processes dynamically modulate surface availability
and function, allowing fine-tuned regulation of serotonin uptake even under resting-

state conditions. Additionally, while slower processes involving endocytosis,
recycling, and degradation typically occur over minutes to hours, subtle fluctuations in
SERT trafficking and activity can still occur under basal conditions. These minor yet
biologically relevant changes likely reflect ongoing homeostatic regulation essential
for maintaining serotonergic balance. Therefore, tracer fluctuations observed during
resting-state measurements should not be dismissed, as they may represent
meaningful variations in SERT regulation that contribute to the fine control of
serotonin clearance.

Implications for the whole-brain serotonergic system

Our graph theory analysis revealed comparable whole-brain organization between
the hemodynamic (BOLD-derived) and serotonergic (PET-derived) connectomes,
although we found only a moderate correlation between the edges of the two
measurements. Notably, the [^{11}C]DASB-based MC results showed a distinct binding
pattern compared with BOLD-fMRI-derived FC. For FC, the ICA revealed classic
RSNs, including the default-mode network, sensory network and motor network,
consistent with previous studies in rats [34] and humans [35]. In contrast, the PET
data revealed two distinct anatomical components in the serotonergic system. One
component included key subcortical areas, such as the brainstem, parts of the
midbrain and the thalamus, regions rich in SERT availability and central to
serotonergic regulation. The other component comprises regions of the limbic
system, such as the striatum, amygdala, insula, cingulate cortex and prefrontal
cortex, critical targets of serotonergic projections from the raphe nuclei. These
components suggest a broader spatial reach of the serotonergic system than the
typically described RSN, reflecting its unique neurochemical architecture. In our
analysis, PET ICs appeared less bilateral than fMRI ICs. This is likely due to the
lower temporal resolution of PET (100 frames) than of fMRI (3000 frames), resulting
in a reduced signal☐to☐noise ratio (SNR) and potentially affecting the stability and
symmetry of the independent components.

Our results demonstrate that compared with FC, MDMA induces more pronounced
changes in MCs, particularly in regions associated with the SERT subcortical
network. The distinct temporal dynamics of BPnd variations between these
components may reflect the hierarchical organization of the serotonergic system.
Specifically, the raphe nuclei, as the primary source of serotonin, are likely to exert

more immediate modulation on posterior subcortical structures (IC2), whereas
downstream effects on limbic and cortical regions (IC1) may occur more gradually.
While these findings align with current neuroanatomical and molecular knowledge,
the precise biological mechanisms driving these temporal differences remain unclear.
Future investigations are warranted to elucidate these mechanisms. Future studies
combining direct measurements of serotonin levels with neuroimaging data will be
critical to fully understanding these components' distinct roles and temporal profiles in
regulating serotonergic function.

Relevance to prior research

Our findings align with Salvan et al. [36], who integrated molecular maps into fMRI
data and demonstrated how individual serotonergic receptors contribute to network-
level activity. Despite our differing methodologies, the receptor activity patterns
found, may also correspond to the independent components we found from the
[^{11}C]DASB PET data. The dual regression approach reported by Salvan et al. was
first reported in the context of pharmacology to delineate the receptor-specific effects
of MDMA in humans [37]. The authors found that MDMA specifically decreased FC in
the 5HT1A maps in areas that could be ascribed to the human salience network,
such as the insula and a collection of medial cortical regions. While not reaching
significance, we observed a reduced salience FC and MC in our data.

Previous studies evaluating the effects of MDMA on FC have indicated relatively few
effects on FC, as confirmed by the results of this study [38, 39]. Recently, a novel
approach [37] revealed decreased connectivity induced by MDMA in areas
associated with SERT and 5-HT$_{1A}$ availability. The increased activity in limbic and
cortical structures when controlling for vascular effects is accompanied by decreased
salience connectivity, indicating that, while neurons become more active through the
drug, they do so in an incoherent manner, which would be in line with the hyperactive
yet abnormal behavior reported for MDMA abuse. Importantly, the potent vascular
effects that play a role in modulating the amplitude of the BOLD-fMRI signal may also
influence FC readouts, although the extent of such effects is difficult to estimate [13].

Over the past decade, PET studies using [^{11}C]DASB have focused on the
associations of serotonin transporter (SERT) availability across different brain
regions, revealing altered interregional SERT connections in patient cohorts,

posttreatment changes, and a predictive capacity for treatment response [9, 10, 40]
[41] [42]. Our study builds upon this foundation, revealing significant within-subject
temporal associations in [^{11}C]DASB binding and demonstrates more pronounced
alterations in MC than in FC induced by drugs, thus highlighting the enhanced utility
of our method.

Study limitations and future directions

One limitation of our study is that our experimental protocols predate the recently
published consensus recommendations for rat fMRI [43], particularly concerning
anesthesia and preprocessing pipelines. Using isoflurane anesthesia, although
common at the time of data acquisition, introduces a potential confound due to its
known neuronal activity effects. However, we previously demonstrated that isoflurane
at 1.3% maintains stable physiological parameters and avoids burst suppression [44],
a concern at higher doses. Furthermore, other studies have reported that low-dose
isoflurane remains feasible for resting-state functional connectivity studies [45]. While
isoflurane, a GABA-A agonist, could theoretically interact with the mechanisms of
MDMA in the brain, we found no evidence in the literature suggesting significant
cross-talk between these substances. Future studies employing medetomidine-based
protocols may help minimize this potential confounding factor.

Additionally, the number of animals used for our pharmacological study with MDMA
was low compared to the resting-state cohort. While futures studies with higher group
sizes may be valuable to confirm and expand our findings, the consistency of effects
between subjects indicated in our data point towards the stability of our results even
at the employed group size.

With respect to data preprocessing, we retained the same pipeline used in our prior
publications [13, 14] to maintain methodological consistency. While we recognize the
advantages of adopting standardized preprocessing, as outlined in the consensus
guidelines, this approach ensures comparability with our previous analyses. To
facilitate further investigation, we have made the full dataset publicly available (see
Data Availability Statement), enabling reanalysis with updated pipelines or additional
explorations of this dataset.

Nonetheless, we provide a strong rationale for considering such analyses in future
studies. First, we show that at rest, the data are reliable, temporally stable and exhibit

similar graph theory metrics to traditionally calculated functional connectomes.
Second, despite comparable network-level organizational properties, at rest, MC is
only moderately correlated with FC, indicating the complementary nature of both
readouts. Third, resting-state MC ICs correlate well with SERT occupancy changes
induced by the MDMA challenge. Finally, we show that the changes that MDMA
elicits on the MC are complementary to standardly calculated [^{11}C]DASB BP$_{ND}$
alterations. Our data indicate that while changes in BP$_{ND}$ are more pronounced in
regions with higher baseline SERT availabilities, MC reveals a more globally
distributed measure of tracking serotonergic changes since regions, such as the
insula, with relatively low SERT expression, are strongly affected. Additionally, our
multimodal imaging approach, although powerful, cannot fully decipher the
mechanisms of interregional coherence in PET time courses. Furthermore, studies
employing more direct sensitive methods to measure neurotransmitter release could
provide deeper insights into the molecular processes underpinning our observations.

435 Conclusion

This study significantly contributes to integrating molecular data into connectomic
frameworks, demonstrating that subject-level MCs are reliable and complementary to
FC in both resting and pharmacologically challenged states. Our research provides a
strong foundation for future investigations into the value and generalizability of PET-
derived MCs, particularly for understanding drug-induced brain-wide molecular
network changes.

442 Bibliography

1.   Wehrl, H.F., et al., *Simultaneous PET-MRI reveals brain function in activated*
and resting state on metabolic, hemodynamic and multiple temporal scales.
Nat Med, 2013. **19**(9): p. 1184-9.
2.   Judenhofer, M.S., et al., *Simultaneous PET-MRI: a new approach for*
functional and morphological imaging. Nat Med, 2008. **14**(4): p. 459-65.
3.   Biswal, B., et al., *Functional connectivity in the motor cortex of resting human*
brain using echo-planar MRI. Magn Reson Med, 1995. **34**(4): p. 537-41.
4.   Jonckers, E., et al., *The power of using functional fMRI on small rodents to*
study brain pharmacology and disease. Frontiers in Pharmacology, 2015.
6(231).
5.   Sander, C.Y., et al., *Neurovascular coupling to D2/D3 dopamine receptor*
occupancy using simultaneous PET/functional MRI. Proceedings of the
National Academy of Sciences of the United States of America, 2013. **110**(27):
p. 11169-11174.

6.   Buxton, R.B., *Dynamic models of BOLD contrast.* NeuroImage, 2012. **62**(2): p.
953-961.
7.   Silberbauer, L.R., et al., *Serotonin Transporter Binding in the Human Brain*
After Pharmacological Challenge Measured Using PET and PET/MR.
Frontiers in Molecular Neuroscience, 2019. **12**(172).
8.   Amend, M., et al., *Functional resting-state brain connectivity is accompanied*
by dynamic correlations of application-dependent [(18)F]FDG PET-tracer
fluctuations. Neuroimage, 2019. **196**: p. 161-172.
9.   Lanzenberger, R., et al., *Prediction of SSRI treatment response in major*
depression based on serotonin transporter interplay between median raphe
nucleus and projection areas. Neuroimage, 2012. **63**(2): p. 874-81.
10. Vanicek, T., et al., *Altered interregional molecular associations of the serotonin*
transporter in attention deficit/hyperactivity disorder assessed with PET.
Human Brain Mapping, 2017. **38**(2): p. 792-802.
11. Hahn, A., R. Lanzenberger, and S. Kasper, *Making Sense of Connectivity.* Int
J Neuropsychopharmacol, 2019. **22**(3): p. 194-207.
12. Sala, A., et al., *Brain connectomics: time for a molecular imaging perspective?*
Trends Cogn Sci, 2023. **27**(4): p. 353-366.
13. Ionescu, T.M., et al., *Neurovascular Uncoupling: Multimodal Imaging*
Delineates the Acute Effects of 3,4-Methylenedioxymethamphetamine. J Nucl
Med, 2023. **64**(3): p. 466-471.
14. Ionescu, T.M., et al., *Striatal and prefrontal D2R and SERT distributions*
contrastingly correlate with default-mode connectivity. Neuroimage, 2021. **243**:
p. 118501.
15. Disselhorst, J.A., et al., *NEMA NU 4-2008 performance evaluation and MR*
compatibility tests of an APD-based small animal PET-insert for simultaneous
PET/MR imaging. Phys Med Biol, 2022. **67**(4).
16. Schiffer, W.K., et al., *Serial microPET measures of the metabolic reaction to a*
microdialysis probe implant. J Neurosci Methods, 2006. **155**(2): p. 272-84.
17. Xia, M., J. Wang, and Y. He, *BrainNet Viewer: A Network Visualization Tool*
for Human Brain Connectomics. PLOS ONE, 2013. **8**(7): p. e68910.
18. Walker, M., et al., *In Vivo Evaluation of 11C-DASB for Quantitative SERT*
Imaging in Rats and Mice. J Nucl Med, 2016. **57**(1): p. 115-21.
19. Walker, M., et al., *Imaging SERT Availability in a Rat Model of L-DOPA-*
Induced Dyskinesia. Mol Imaging Biol, 2020. **22**(3): p. 634-642.
20. Di, X., B.B. Biswal, and I. Alzheimer's Disease Neuroimaging, *Metabolic brain*
covariant networks as revealed by FDG-PET with reference to resting-state
fMRI networks. Brain Connect, 2012. **2**(5): p. 275-83.
21. Paterson, L.M., et al., *Measuring Endogenous 5-HT Release by Emission*
Tomography: Promises and Pitfalls. Journal of Cerebral Blood Flow &
Metabolism, 2010. **30**(10): p. 1682-1706.
22. Yamamoto, S., et al., *Effects of increased endogenous serotonin on the in vivo*
binding of [11C]DASB to serotonin transporters in conscious monkey brain.
Synapse, 2007. **61**(9): p. 724-731.
23. Lundquist, P., et al., *Effect on [11C]DASB binding after tranylcypromine-*
induced increase in serotonin concentration: Positron emission tomography
studies in monkeys and rats. Synapse, 2007. **61**(6): p. 440-449.
24. Talbot, P.S., et al., *Effects of reduced endogenous 5-HT on the in vivo binding*
of the serotonin transporter radioligand 11C-DASB in healthy humans.
Synapse, 2005. **55**(3): p. 164-175.

25. Lefevre, A., et al., *Oxytocin and Serotonin Brain Mechanisms in the*
Nonhuman Primate. The Journal of Neuroscience, 2017. **37**(28): p. 6741-
6750.
26. Ramamoorthy, S. and R.D. Blakely, *Phosphorylation and Sequestration of*
Serotonin Transporters Differentially Modulated by Psychostimulants. Science,
1999. **285**(5428): p. 763-766.
27. Blakely, R.D., et al., *Regulated phosphorylation and trafficking of*
antidepressant-sensitive serotonin transporter proteins. Biological Psychiatry,
1998. **44**(3): p. 169-178.
28. Raichle, M.E. and D.A. Gusnard, *Appraising the brain's energy budget.*
Proceedings of the National Academy of Sciences, 2002. **99**(16): p. 10237-
10239.
29. Salomon, R.M. and R.L. Cowan, *Oscillatory serotonin function in depression.*
Synapse, 2013. **67**(11): p. 801-820.
30. Yang, H., et al., *Physiologically relevant changes in serotonin resolved by fast*
microdialysis. ACS chemical neuroscience, 2013. **4**(5): p. 790-798.
31. Lau, T. and P. Schloss, *Differential regulation of serotonin transporter cell*
surface expression. Wiley Interdisciplinary Reviews: Membrane Transport and
Signaling, 2012. **1**(3): p. 259-268.
32. Jayanthi, L.D., et al., *Evidence for biphasic effects of protein kinase C on*
serotonin transporter function, endocytosis, and phosphorylation. Mol
Pharmacol, 2005. **67**(6): p. 2077-87.
33. Steiner, J.A., A.M. Carneiro, and R.D. Blakely, *Going with the flow: trafficking-*
dependent and -independent regulation of serotonin transport. Traffic, 2008.
9(9): p. 1393-402.
34. Becerra, L., et al., *Robust reproducible resting state networks in the awake*
rodent brain. PLoS One, 2011. **6**(10): p. e25701.
35. Fox, M.D., et al., *The human brain is intrinsically organized into dynamic,*
anticorrelated functional networks. Proc Natl Acad Sci U S A, 2005. **102**(27):
p. 9673-8.
36. Salvan, P., et al., *Serotonin regulation of behavior via large-scale*
neuromodulation of serotonin receptor networks. Nat Neurosci, 2023. **26**(1): p.
53-63.
37. Dipasquale, O., et al., *Receptor-Enriched Analysis of functional connectivity by*
targets (REACT): A novel, multimodal analytical approach informed by PET to
study the pharmacodynamic response of the brain under MDMA. NeuroImage,
2019. **195**: p. 252-260.
38. Carhart-Harris, R.L., et al., *The Effects of Acutely Administered 3,4-*
Methylenedioxymethamphetamine on Spontaneous Brain Function in Healthy
Volunteers Measured with Arterial Spin Labeling and Blood Oxygen Level–
Dependent Resting State Functional Connectivity. Biological Psychiatry, 2015.
78(8): p. 554-562.
39. Roseman, L., et al., *The effects of psilocybin and MDMA on between-network*
resting state functional connectivity in healthy volunteers. Frontiers in human
neuroscience, 2014. **8**: p. 204-204.
40. James, G.M., et al., *Effects of Selective Serotonin Reuptake Inhibitors on*
Interregional Relation of Serotonin Transporter Availability in Major
Depression. Frontiers in Human Neuroscience, 2017. **11**(48).
41. Hahn, A., et al., *Attenuated serotonin transporter association between dorsal*
raphe and ventral striatum in major depression. Human brain mapping, 2014.
35(8): p. 3857-3866.

42. Hahn, A., et al., *Escitalopram enhances the association of serotonin-1A*
autoreceptors to heteroreceptors in anxiety disorders. J Neurosci, 2010.
30(43): p. 14482-9.
43. Grandjean, J., et al., *A consensus protocol for functional connectivity analysis*
in the rat brain. Nat Neurosci, 2023. **26**(4): p. 673-681.
44. Ionescu, T.M., et al., *Elucidating the complementarity of resting-state networks*
derived from dynamic [(18)F]FDG and hemodynamic fluctuations using
simultaneous small-animal PET/MRI. Neuroimage, 2021. **236**: p. 118045.
45. Hutchison, R.M., et al., *Isoflurane induces dose-dependent alterations in the*
cortical connectivity profiles and dynamic properties of the brain's functional
architecture. Hum Brain Mapp, 2014. **35**(12): p. 5754-75.

Statements & Declarations

Funding

This research was supported by funds from the Eberhard Karls University Tübingen Faculty of Medicine (fortüne 2209-0-0 and 2409-0-0) to HFW, from Bundesministerium für Bildung und Forschung (BMBF, Grant No. 01GQ1415) to HFW and from the Carl Zeiss Foundation to KH.

Competing Interests

The authors declare that they have no conflicts of interest.

CRediT authorship contribution statement

Tudor M. Ionescu: Conceptualization, Methodology, Software, Validation, Formal analysis, Investigation, Data curation, Writing – original draft, Visualization. **Mario Amend:** Investigation, Writing – review & editing. **Rakibul Hafiz:** Methodology, Software, Writing – review & editing. **Bharat B. Biswal:** Conceptualization, Writing – review & editing, Supervision, Project administration, Funding acquisition. **Andreas Maurer:** Writing – review & editing. **Hans F. Wehrl:** Conceptualization, Methodology, Writing – review & editing, Supervision, Project administration, Funding acquisition. **Kristina Herfert:** Writing – review & editing, Visualization, Supervision, Project administration, Funding.

Data availability

The data are openly available at the DRYAD repository (https://doi.org/10.5061/dryad.6djh9w1bf).