1 **Mapping Serotonergic Dynamics using Drug-Modulated Molecular**

2 **Connectivity**

Ionescu TM[1], Amend M[1], Hafiz R[2], Maurer A[1], Biswal BB[2], Wehrl HF[1†], Herfert K[1†]

1 Werner Siemens Imaging Center, Department of Preclinical Imaging and Radiopharmacy,
Eberhard Karls University Tuebingen, Germany

2 Department of Biomedical Engineering, New Jersey Institute of Technology, University
Heights, Newark, New Jersey, USA

[†] contributed equally

Corresponding author:

Prof. Dr. Kristina Herfert (kristina.herfert@med.uni-tuebingen.de)

Address: Roentgenweg 13, 72076 Tuebingen, Germany,

Tel: +49 7071 29 87680

Abstract

Understanding the complex workings of the brain is one of the most significant
challenges in neuroscience, providing insights into the healthy brain, diseases, and
the effects of potential therapeutics. A major challenge in this field is the limitations of
traditional brain imaging techniques, which often deliver only a part of the complex
puzzle of brain function. Our research employs a novel approach named "Molecular
Connectivity" (MC), which merges the strengths of various imaging methods to offer
a comprehensive view of how molecules interact within the brain and affect its
function.

This innovative technique bridges the gap between functional magnetic resonance
imaging (fMRI), known for its ability to monitor brain activity by tracking blood flow,
and positron emission tomography (PET), which can depict specific molecular
changes. By integrating these methods, we can better understand the far-reaching
impacts of drugs on the brain. Our study focuses on the application of dynamic
$[^{11}C]$DASB PET scans to map the distribution of serotonin transporters, a key player
in regulating mood and emotions, and examines how these are altered following the
use of methylenedioxymethamphetamine (MDMA), commonly known as ecstasy.

Through a detailed analysis comparing MC with traditional measures of brain
connectivity, we uncover significant patterns that closely align with physiological
changes. Our results reveal clear changes in molecular connectivity after a single
dose of MDMA, establishing a direct link between the drug's effects on serotonin
transporter occupancy and changes in the brain's functional network.

This work not only offers a novel methodology for the in-depth study of brain function
at the molecular level but also opens new pathways for understanding how drugs
modulate brain activity.

Keywords: Molecular Connectivity, PET/fMRI, Functional Connectivity,
Pharmacological Imaging

Introduction

The field of neuroscience has witnessed remarkable advancements over the past decade, particularly propelled by the advent of innovative imaging techniques. The simultaneous application of positron emission tomography (PET) and magnetic resonance imaging (MRI) has revolutionized our ability to concurrently assess brain function across various physiological dimensions [1, 2]. This is especially true in the context of drug effect evaluations. The combination of PET with functional MRI (fMRI) has opened a plethora of investigative avenues. Techniques such as resting-state functional connectivity (rs-FC) [3] or pharmacological fMRI (phMRI) [4] are now complemented by the capability of PET to quantify molecular changes, including alterations in receptor and transporter availability [5].

While fMRI provides high spatial and temporal resolution, the interpretation of its readout necessitates caution. The Blood-Oxygen-Level-Dependent (BOLD) signal used in fMRI indirectly captures neuronal changes through neurovascular coupling [6], onling reveals the hemodynamic consequences of molecular-level drug effects. The integration of simultaneous PET acquisition can bridge this interpretative gap by offering essential molecular insights, particularly regarding transporter or receptor alterations. Typically, PET, used either independently or simultaneously to fMRI, has been mainly used in pharmacological studies for illustrating quantitative shifts in neuroreceptor or transporter availabilities [7]. Several studies have also explored the interregional coherence of PET tracer signals [8-10], an approach akin to fMRI-derived rs-FC. While subject-wise metabolic connectivity using [18F]FDG-PET has been established through the temporal correlation of regional PET signals [1, 8], studies employing transporter or receptor tracers have predominantly focused on interregional binding coherences across subjects using static scans [9, 10]. The concept and feasibility of molecular connectivity (MC) through the temporal correlation of dynamic binding potentials of transporter or receptor tracers has yet to be explored.

In our study, we investigate the feasibility of deriving MC from dynamic [11C]DASB-PET scans acquired simultaneously with fMRI in rats. We divided the rats into two cohorts: a baseline group and a pharmacological application group, exposed to 3,4-methyldeoxymetamphetamine (MDMA). The baseline cohort served to assess the feasibility of the novel methodology, contrasting it with traditional fMRI-derived rs-FC

with a specific focus on temporal stability. We postulated that dynamic [^{11}C]DASB
PET temporal fluctuations could be harnessed for connectivity data, in a manner
similar to hemodynamic rs-FC, using seed-based and independent component
analysis (ICA).

The second cohort, subjected to an MDMA challenge allowed us to evaluate the
utility of our novel approach. We aimed to outline the effects of MDMA by integrating
the innovative MC concept with established analysis techniques. Our primary
objective of this research was to elucidate the potential of PET-derived MC in
conjunction with simultaneous PET/fMRI, exploring the avenues this methodology
could open for future diagnostic and drug development studies.

Material and Methods

86 Our study reevaluates two distinct datasets previously published by our group [11,
87 12], to simultaneously explore FC and MC both at baseline conditions [12] and
88 following MDMA administration [11]. For detailed descriptions on animal handling,
89 experimental setups, and data acquisition procedures, please refer to these earlier
90 publications.

Animals

A total of 41 male Lewis rats were obtained from Charles River Laboratories
(Sulzfeld, Germany). Thirty rats underwent baseline [^{11}C]DASB PET/fMRI scans
without any pharmacological intervention, while 11 rats underwent [^{11}C]DASB
PET/fMRI scans including an acute MDMA challenge. All experiments were
conducted in compliance with German federal regulations for experimental animals
and received approval from the Regierungspräsidium Tübingen.

Simultaneous PET/fMRI Experiments

The rats were subjected to simultaneous PET/fMRI experiments involving 1.3%
isoflurane anesthesia, tail vein catheterization, positioning on a temperature-
controlled bed, and monitoring vital signs. The scans were performed using a 7T
small-animal ClinScan MRI scanner (Bruker Biospin, Ettlingen, Germany) with a
custom-developed PET insert [13]. Both the scanning protocol and the sequence
parameters have been outlined in detail in our previous publication [11]. The MDMA

cohort received a pharmacological MDMA challenge of 3.2 mg/kg intravenously 40

minutes after tracer injection.

Data Preprocessing and Analysis

Data preprocessing followed established protocols, including steps such as

realignment, mask creation, coregistration, spatial normalization, signal cleaning, and

spatial smoothing, as detailed in our previous work [11]. For the MDMA dataset, PET

scans were analyzed for early and late effects post-challenge using the general linear

model (GLM) available in SPM. For both datasets the baseline was defined 30 to 40

minutes after scan start. For the fMRI data a first level analysis was applied to the

individual scans without a high-pass filter (the filter was set to "Inf"). Statistical

parametric maps were generated post GLM parameter estimation using contrast

vectors. Group-level analysis involved a one-sample t-test on the subject-level

statistical parametric maps ($p < 0.05$, one-sided, family wise error / FWE-adjusted).

Static PET scans were generated by summing dynamic frames over defined periods

for 10-minute periods after the MDMA challenge (50-60 minutes to investigate early

effects, and 70-80 minutes to investigate late MDMA effects). Two-sample t-maps

were calculated between the normalized [^{11}C]DASB uptakes of (1) the baseline scan

period and the early effect time period and of (2) the early effect time period and the

late effect time period ($p < 0.05$, FWE-adjusted).

All group-level t-maps underwent voxel-wise signal quantification to determine the

regional contributions of 48 regions selected according to the Schiffer atlas [14].

Average t-scores and standard deviations of all voxels were calculated.

Functional Connectivity Analysis

Functional connectivity was determined using a seed-based analysis approach. The

mean time series of the pre-processed BOLD-fMRI signals for each dataset across

all regions (refer to *Supplementary Table 1* for the list of regions) were extracted

using SPM toolbox Marseille Boîte À Région d'Intérêt (MarsBaR). Pairwise Pearson's

correlation coefficients were calculated between each pair of mean regional time-

series for every dataset. The Pearson's r coefficients were converted into z-values

using Fischer's transformation for group-level analysis. The Fischer's z-transformed

correlation coefficients were then used to generate mean correlation matrices for
both cohorts [15].

Molecular connectivity analysis

138 The mean [^{11}C]DASB signal from the preprocessed PET datasets was extracted from
139 the designed regions, including the 48 regions used for fMRI data analysis and the
140 cerebellum using MarsBaR. Binding potentials were calculated frame-wise for all
141 dynamic PET scans using the DVR-1 (equation 1) to generate regional BP$_{ND}$ values
142 using the cerebellum as a reference region [16]:

143
$$BP_{ND} = \frac{V_T - V_{ND}}{V_{ND}} = \frac{V_T}{V_{ND}} - 1 = DVR - 1,$$

where:

• BP_{ND} is the binding potential
• V_T is the total volume of distribution
• V_{ND} is the volume of distribution in a reference tissue
• DVR is the relative volume of distribution

To calculate MC, we discarded the first 20 minutes of every scan, which were
dominated by perfusion effects and applied a detrending approach on the remaining
60 minutes in order to obtain temporally stable values (for further details, please refer
to Supplementary Methods and Supplementary Figure 1). The BP$_{ND}$ time courses
were then used to calculate MC as described above for fMRI: subject-level
correlation matrices between all regional time courses were generate and z-
transformed correlation coefficients were used to calculate mean correlation
matrices.

Independent Component Analysis

Group ICA (GIFT toolbox, MIALAB, University of New Mexico, Albuquerque, NM,
USA) was used for ICA of the baseline group. Both the fMRI and PET preprocessed
data sets were investigated between 30 and 80 minutes after start of data
acquisition. For fMRI, we selected 10 independent components, while we started with
two components for PET and increased the number to ten components to thoroughly
dissect the varying components within the signal. Components were thresholded at a
z-vlaue ≥ 1.96 (p-value ≤ 0.05) [17] and average z-scores and standard deviations

were calculated for each component. These components' physiological significance

was further explored by contrasting them with the regional [11C]DASB changes

induced by MDMA. Accordingly, the z-scores of the independent components

generated in the baseline cohort were correlated with the early and late regional

[^{11}C]DASB changes induced by MDMA, measured using t-scores.

Results

Comparability of MC and FC in spatial contexts and over time

172 We first aimed to evaluate whether MC aligns spatially with FC, possesses similar

173 graph theory properties and provides consistent temporal readouts throughout the

174 scan duration in the baseline group (Figure 1).

Figure 1: Evaluation of seed-based MC. (A) Correlation matrix indicating whole-brain FC (beneath diagonal) and MC (above diagonal). Correlations not surviving significance testing with multiple comparison corrections were set to zero (p < 0.05, FWE correction). **(B)** Scatter plot and correlation between MC and FC edges. **(C)** Small-world coefficients for all subjects and group-level one-sample t-test against the value of 1 (SW > 1 is considered as indicative of small-world properties). Copmarison of **(D)** FC and **(G)** MC early (20-40 minutes after scan start, below diagonal) and late (60-80 minutes after scan start, above diagonal). The similarties of early and late readouts were quantified for both **(E)** FC and **(H)** MC. Temporal stability of both **(F)** FC and **(I)** MC was evaluated using a sliding window approach including 20-minute windows between 20 and 80 minutes after scan start. Abbreviations: FC

= fMRI-derived hemodynamic functional connectivity, MC = [^{11}C]DASB PET-derived molecular

connectivity.

We found moderate, but significant correlation between the edge-level MC and FC (r

= 0.51, p < 0.001, Figure 1A and B). Furthermore, both connectomes revealed small-

world properties at group-level, with coefficients higher than 1 (Figure 1C). At subject

level, three molecular and four functional connectomes fell below the threshold of 1

for the small-world coefficient. A significant consistency was observed betweeen

early and late scan-derived connectomes (Figure 1D for FC, G for MC), with FC

having a slight edge (Figure 1E, r = 0.96) over MC (Figure 1H, r = 0.8). While both,

FC and MC maintained steady correlation intensities, there was a negligable decline

over the scan duration (Figure 1F and 1I).

Deciphering spatial characteristics of FC and MC using ICA

After establishing the feasibility of obtaining temporally stable readouts using the

ROI-to-ROI approach, we employed a data-driven approach, using ICA, to compare

spatial characteristics of FC and MC (Figure 2).

Figure 2: Group independent component analysis for FC and MC. (A) ICA performed over 10 components for fMRI. **(B)** ICA performed over 2 components for [^{11}C]DASB PET and regional quantification of the two derived components. The ICA was repeated over 10 components. Four and three components showed good overlap with the two components defined above. All components were thresholded at z > 1.96 (p ≤ 0.05). Abbreviations: FC = fMRI-derived hemodynamic functional connectivity, MC = [^{11}C]DASB PET-derived molecular connectivity.

We extracted ten group ICs from the fMRI data (Figure 2A), revealing known

canonical resting-state networks, such as the posterior default-mode-like network

(IC1-5, red), sensorimotor networks (IC1-5, green and purple), the anterior default-

mode-like network (IC6-10, yellow) or the visual network (IC6-10, red). With

unpredictability for the number of ICs suitable for MC IC extraction, we started with

two components (Figure 2B). IC1 (orange) comprised both subcortical and cortical

anterior brain regions, including the nucleus accumbens, amygdala, cingulate cortex,

caudate putamen, orbitofrontal cortex and medial prefrontal cortex, while IC2 (blue)

primarily received contributions from deeper posterior areas, such as the midbrain,

thalamus, hypothalamus, periaquaeductal gray and medulla. Interestingly, when we

extracted 10 independent components, to mimic the number of components used for

the FC data, we found that the initial anterior component split in four different ICs and

the initial posterior IC in three different ICs. A relatively clear spatial segregation can

be seen for the newly-formed ICs, for instance the three posterior components being

extracted from specific regions (green mainly from medulla, red from hypothalamus

and part of the midbrain, blue from midbrain and thalamus).

223   MDMA-induced changes of ICA-derived molecular connectivity

224   Next, we aimed to explore the relationship between molecular changes in SERT

225   availability and the molecular connectome derived from ICA, induced by an acute

226   MDMA administration (Figure 3).

Figure 3: Comparison of MDMA-induced [^{11}C]DASB alterations. (A) Left panel: Dynamic binding potentials of regions comprising SERT network, defined by IC 1 in validation cohort. Right panel: Dynamic binding potentials of regions comprising salience network, defined by IC 1 in validation cohort. **(B)** Overlap between independent components extracted from the validation cohort (IC 1 = SERT network; IC 2 = salience network) and the early and late effects of MDMA respectively. **(C)** Pair-

wise correlations between regional z-scores of the ICs extracted from validation cohort and regional t-
scores of early and late MDMA effects. (*** indicates p < 0.001, ns = non-significant). Abbreviations:
SERT = serotonin transporter, ICA = independent component analysis; for abbreviations of regions
please refer to *Supplementary Information*).

Two IC extracted from the MC showed good overlap with regions associated with the
salience network (IC1) and with those having high SERT densities (IC2). Therefore,
we defined the regions contributing strongly to IC1 as salience network (CPu, Cg,
NAc, Amyg, Ins, mPFC) and those with strong signals in IC2 as SERT network (VTA,
Th, MB, PAG, Hypo). Interestingly, MDMA induced immediate strong decreases in all
SERT network regions, salience areas exhibited a delay by approximately 10 minutes
(Figure 3A). Voxel-level analysis showed clear spatial overlaps between early MDMA
responsive regions and those from the posterior IC, with delayed regions mirroring
the anterior IC reminiscent of the salience network (Figure 3B). To quantify the
striking spatial similarity between the baseline independent components and the
spatiotemporal characteristics of $[^{11}C]$DASB changes after MDMA exposure, we
show highly significant correlations between the z-scores of the posterior and anterior
ICs and the t-scores of late and early MDMA effects on $[^{11}C]$DASB alterations,
respectively (p < 0.0001, Figure 3C).

Finally, we investigated relationships between SERT availability changes and MC
reductions following the acute MDMA challenge (Figure 4).

Figure 4: Comparison of MC and BP$_{ND}$ changes. A Reductions in BP$_{ND}$ following MDMA (orange)
and MC strength (blue) were compared. **B** T-scores derived from each approach show only low
correlations.

We found that the decreases in MC and in $[^{11}C]$DASB BP$_{ND}$ following MDMA
application (Figure 4A) showed only low correlation (r = 0.29, Figure 4B). While

decreases in SERT availability exhibited a strong anterior-posterior gradient, being
most pronounced in areas with high SERT availability, such as the MB, VTA, Pons or
PAG, MC encompassed regions across the brain to similar extents. Specifically,
while the significance of [^{11}C]DASB reductions in the Ins was very low across
investigated regions, the Ins showed highest effects among regions for MC, while
strong SERT occupancy effects in the IC and SC did not translate in very prominent
reductions in the respective global MC of the two regions.

MDMA-induced changes of seed based molecular connectivity

267 Next, we performed a seed-based analysis and aimed to compare changes in FC
268 and MC after an MDMA pharmacological challenge on the salience network and on
269 regions with high SERT binding (Figure 5).

Figure 5: MDMA effects on seed-based FC and MC of the salience and SERT networks. (A) FC
and **(C)** MC brain networks depicting edge and node strengths of the salience network and SERT
network at baseline (20-40 min after scan start) and after MDMA (60-80 min after scan start). **(B)** FC
and **(D)** MC time-resolved salience and SERT network strengths computed using sliding windows.
Asterisks indicate significant (p < 0.05, FDR-corrected) changes to baseline (time-point zero,

corresponding to 20-40 minutes after scan start). Abbreviations: SN = salience network, SERT =
SERT network.

We observed a small decrease in the SN for FC (16% at the end of the scan) and
almost constant FC of the SERT network (<5% decrease) following MDMA, as shown
in Figure 5A at edge and node level and Figure 5C at network level (p>0.05). For
MC, we observed profound reductions in the SERT network (Figure 5C and D),
emphasizing an acute and spatially specific effect of MDMA on MC.

Discussion

The mammalian brain operates on diverse physiological, spatial and temporal scales.
FC via BOLD-fMRI offers insights into coherent functional brain networks, but its
complexity and indirect link to neural activity highlight the need for more direct
methodologies. In this context, the concept of MC using [11C]DASB PET, as
introduced in our study, provides a more direct and complementary perspective on
brain organization and its response to external stimuli, such as MDMA.

Physiological basis

Our findings suggest that [11C]DASB binding reflect the interplay of serotonin levels
and SERT dynamics [18]. Supporting the competition model, evidence indicates that
endogenous serotonin competes with tracers for binding sites, affecting tracer
binding [19]. However, contrasting results in various studies highlight the complexity
of this interaction [20] [21] [22]. The internalization model suggests serotonin levels
influence SERT internalization, impacting [11C]DASB binding [18, 23, 24]. While
supporting evidence exists, further exploration is needed to fully understand these
dynamics, especially during resting-state [25]. Some models on this aspect have
proposed a regulatory function of the raphé nuclei in maintaining serotonin
fluctuations over several temporal scales at rest [26]. Remarkably, fast microdialysis
has resolved multiple spontaneous surges of up to 1500-fold of the basal serotonin
occurring during 30-minute intervals [27]. Additionally, the same study has indicated
that SERT expression is essential for the spontaneous surges, reduced SERT
drastically decreasing serotonin spiking. Thus, it is feasible that the correlated
temporal fluctuations captured by dynamic [11C]DASB at least partly reflect the role of
the serotonergic system in the resting activity of the brain.

Implications for whole-brain serotonergic system

Our graph-theory analysis revealed comparable whole-brain organization between hemodynamic and serotonergic connectomes, although we only found moderate correlation between the edges of the two measurements. Notably, the correlations in [^{11}C]DASB bindings differed from BOLD-fMRI-derived correlations. For BOLD-fMRI, ICA revealed classic RSNs, widely reported for both rats [28] and humans [29], including the default-mode network, the sensory network and the motor network. In contrast, the ICA analysis of PET data revealed two distinct anatomical modules. The first component included primarily the brainstem, parts of the midbrain and the thalamus. The second component comprised regions of the limbic system, such as the striatum, amygdala, insular, cingulate and prefrontal cortices, reflecting the diverse projections of the raphe nuclei. MDMA's interaction with SERT, as elucidated by [^{11}C]DASB, shows region-specific temporal effects, with immediate decreases in binding in the brainstem and posterior subcortical regions, and delayed effects in limbic pathways. This suggests that the serotonergic system's response to drugs can be predicted, to an extent, by the observed independent components.

Relevance to prior research

Our findings align with those by Salvan et al [31] on the integration of molecular maps into fMRI data demonstrating how individual serotonergic receptors contribute to network-level activity. Despite our differing methodologies, the receptor activity patterns they found, may also correspond to the different independent components we found from the [^{11}C]DASB PET data. The dual regression approach reported by Salvan et al. has first been reported in the context of pharmacology to delineate the receptor-specific effects of MDMA in humans [32]. The authors found that MDMA specifically decreased FC in the 5HT1A maps in areas which could be ascribed to the human salience network, such as the insula and a collection of medial cortical regions. While not reaching significance, we observed a reduction in salience FC and MC in our data.

Previous studies evaluating the effects of MDMA on FC indicated relatively sparse effects on FC, as confirmed by the readout of this study [33, 34]. Recently, a novel approach [32] has revealed decreased connectivity induced by MDMA in areas associated with SERT and 5-HT$_{1A}$ availability. The increased activity indicated in

limbic and cortical structures when controlling for vascular effects is accompanied
with decreased salience connectivity, indicating that, while neurons become more
active through the drug, they do so in an incoherent manner, which would be in line
with the hyperactive, yet abnormal behavior reported for MDMA abuse. Importantly,
the potent vascular effects which play a role in modulating the amplitude of the
BOLD-fMRI signal may also influence FC readouts, although the extent of such
effects are difficult to estimate [11].

Over the past decade, PET studies using [^{11}C]DASB have focused on the
associations of serotonin transporter (SERT) availabilities across different brain
regions, revealing altered interregional SERT connections in patient cohorts, post-
treatment changes, and predictive capacity for treatment response [9, 10, 35] [36]
[37]. Our study builds upon this foundation, uncovering significant within-subject
temporal associations in [^{11}C]DASB binding, and demonstrates more pronounced
alterations in MC compared FC changes induced by drugs, thus highlighting the
enhanced utility of our method.

Study limitations and future directions

While our study advances the understanding of MC, the use of anesthetized animal is
a limitation, potentially affecting PET and fMRI readouts and their interaction with
MDMA. Future studies should involve awake animals to validate our method.
Additionally, our multimodal imaging approach, though powerful, cannot fully
decipher the mechanisms of interregional coherences in PET timecourses. Future
studies employing sensitive methods to measure neurotransmitter release, could
provide deeper insights into the molecular processes underpinning our observations.

Nonetheless, we provide several strong arguments to consider such analyses as
presented here in future studies. First, we show that at rest the data are reliable,
temporally stable and exhibiting similar graph theory metrics to traditionally calculated
functional connectomes. Second, in spite of comparable network-level organizational
properties, already at rest MC is only moderately correlated to FC, indicating the
complementary nature of both readouts. Third, resting-state MC ICs correlate well
with SERT occupancy changes induced by the MDMA challenge. Finally, we show
that the changes MDMA elicits on MC are complementary to standardly-calculated
[^{11}C]DASB BP$_{ND}$ alterations. Our data indicate that, while changes in BP$_{ND}$ are more

pronounced in regions with higher baseline SERT availabilities, MC reveals a more
globally distributed measure of tracking serotonergic changes, since also regions,
such as the insula, with relatively low SERT expressions, were strongly affected.

Conclusion

This study significantly contributes to integrating molecular data into connectomic
frameworks, demonstrating that subject-level MC is reliable and complementary to
FC in both resting and pharmacologically challenged states. Our research lays a
strong foundation for future investigations into the value and generalizability of PET-
derived MC, particularly for understanding drug-induced brain-wide molecular
network changes.

Bibliography

1.   Wehrl, H.F., et al., *Simultaneous PET-MRI reveals brain function in activated*
and resting state on metabolic, hemodynamic and multiple temporal scales.
Nat Med, 2013. **19**(9): p. 1184-9.
2.   Judenhofer, M.S., et al., *Simultaneous PET-MRI: a new approach for*
functional and morphological imaging. Nat Med, 2008. **14**(4): p. 459-65.
3.   Biswal, B., et al., *Functional connectivity in the motor cortex of resting human*
brain using echo-planar MRI. Magn Reson Med, 1995. **34**(4): p. 537-41.
4.   Jonckers, E., et al., *The power of using functional fMRI on small rodents to*
study brain pharmacology and disease. Frontiers in Pharmacology, 2015.
6(231).
5.   Sander, C.Y., et al., *Neurovascular coupling to D2/D3 dopamine receptor*
occupancy using simultaneous PET/functional MRI. Proceedings of the
National Academy of Sciences of the United States of America, 2013. **110**(27):
p. 11169-11174.
6.   Buxton, R.B., *Dynamic models of BOLD contrast.* NeuroImage, 2012. **62**(2): p.
953-961.
7.   Silberbauer, L.R., et al., *Serotonin Transporter Binding in the Human Brain*
After Pharmacological Challenge Measured Using PET and PET/MR.
Frontiers in Molecular Neuroscience, 2019. **12**(172).
8.   Amend, M., et al., *Functional resting-state brain connectivity is accompanied*
by dynamic correlations of application-dependent [(18)F]FDG PET-tracer
fluctuations. Neuroimage, 2019. **196**: p. 161-172.
9.   Lanzenberger, R., et al., *Prediction of SSRI treatment response in major*
depression based on serotonin transporter interplay between median raphe
nucleus and projection areas. Neuroimage, 2012. **63**(2): p. 874-81.
10.   Vanicek, T., et al., *Altered interregional molecular associations of the serotonin*
transporter in attention deficit/hyperactivity disorder assessed with PET.
Human Brain Mapping, 2017. **38**(2): p. 792-802.

11. Ionescu, T.M., et al., *Neurovascular Uncoupling: Multimodal Imaging Delineates the Acute Effects of 3,4-Methylenedioxymethamphetamine.* J Nucl Med, 2023. **64**(3): p. 466-471.

12. Ionescu, T.M., et al., *Striatal and prefrontal D2R and SERT distributions contrastingly correlate with default-mode connectivity.* Neuroimage, 2021. **243**: p. 118501.

13. Disselhorst, J.A., et al., *NEMA NU 4-2008 performance evaluation and MR compatibility tests of an APD-based small animal PET-insert for simultaneous PET/MR imaging.* Phys Med Biol, 2022. **67**(4).

14. Schiffer, W.K., et al., *Serial microPET measures of the metabolic reaction to a microdialysis probe implant.* J Neurosci Methods, 2006. **155**(2): p. 272-84.

15. Xia, M., J. Wang, and Y. He, *BrainNet Viewer: A Network Visualization Tool for Human Brain Connectomics.* PLOS ONE, 2013. **8**(7): p. e68910.

16. Walker, M., et al., *In Vivo Evaluation of 11C-DASB for Quantitative SERT Imaging in Rats and Mice.* J Nucl Med, 2016. **57**(1): p. 115-21.

17. Di, X., B.B. Biswal, and I. Alzheimer's Disease Neuroimaging, *Metabolic brain covariant networks as revealed by FDG-PET with reference to resting-state fMRI networks.* Brain Connect, 2012. **2**(5): p. 275-83.

18. Paterson, L.M., et al., *Measuring Endogenous 5-HT Release by Emission Tomography: Promises and Pitfalls.* Journal of Cerebral Blood Flow & Metabolism, 2010. **30**(10): p. 1682-1706.

19. Yamamoto, S., et al., *Effects of increased endogenous serotonin on the in vivo binding of [11C]DASB to serotonin transporters in conscious monkey brain.* Synapse, 2007. **61**(9): p. 724-731.

20. Lundquist, P., et al., *Effect on [11C]DASB binding after tranylcypromine-induced increase in serotonin concentration: Positron emission tomography studies in monkeys and rats.* Synapse, 2007. **61**(6): p. 440-449.

21. Talbot, P.S., et al., *Effects of reduced endogenous 5-HT on the in vivo binding of the serotonin transporter radioligand 11C-DASB in healthy humans.* Synapse, 2005. **55**(3): p. 164-175.

22. Lefevre, A., et al., *Oxytocin and Serotonin Brain Mechanisms in the Nonhuman Primate.* The Journal of Neuroscience, 2017. **37**(28): p. 6741-6750.

23. Ramamoorthy, S. and R.D. Blakely, *Phosphorylation and Sequestration of Serotonin Transporters Differentially Modulated by Psychostimulants.* Science, 1999. **285**(5428): p. 763-766.

24. Blakely, R.D., et al., *Regulated phosphorylation and trafficking of antidepressant-sensitive serotonin transporter proteins.* Biological Psychiatry, 1998. **44**(3): p. 169-178.

25. Raichle, M.E. and D.A. Gusnard, *Appraising the brain's energy budget.* Proceedings of the National Academy of Sciences, 2002. **99**(16): p. 10237-10239.

26. Salomon, R.M. and R.L. Cowan, *Oscillatory serotonin function in depression.* Synapse, 2013. **67**(11): p. 801-820.

27. Yang, H., et al., *Physiologically relevant changes in serotonin resolved by fast microdialysis.* ACS chemical neuroscience, 2013. **4**(5): p. 790-798.

28. Becerra, L., et al., *Robust reproducible resting state networks in the awake rodent brain.* PLoS One, 2011. **6**(10): p. e25701.

29. Fox, M.D., et al., *The human brain is intrinsically organized into dynamic, anticorrelated functional networks.* Proc Natl Acad Sci U S A, 2005. **102**(27): p. 9673-8.

461   30.   Shih, J.-H., et al., *Evaluation of brain SERT occupancy by resveratrol against*
462   *MDMA-induced neurobiological and behavioral changes in rats: A 4-[18F]-*
463   *ADAM/small-animal PET study.* European Neuropsychopharmacology, 2016.
464   **26**(1): p. 92-104.
465   31.   Salvan, P., et al., *Serotonin regulation of behavior via large-scale*
466   *neuromodulation of serotonin receptor networks.* Nat Neurosci, 2023. **26**(1): p.
467   53-63.
468   32.   Dipasquale, O., et al., *Receptor-Enriched Analysis of functional connectivity by*
469   *targets (REACT): A novel, multimodal analytical approach informed by PET to*
470   *study the pharmacodynamic response of the brain under MDMA.* NeuroImage,
471   2019. **195**: p. 252-260.
472   33.   Carhart-Harris, R.L., et al., *The Effects of Acutely Administered 3,4-*
473   *Methylenedioxymethamphetamine on Spontaneous Brain Function in Healthy*
474   *Volunteers Measured with Arterial Spin Labeling and Blood Oxygen Level–*
475   *Dependent Resting State Functional Connectivity.* Biological Psychiatry, 2015.
476   **78**(8): p. 554-562.
477   34.   Roseman, L., et al., *The effects of psilocybin and MDMA on between-network*
478   *resting state functional connectivity in healthy volunteers.* Frontiers in human
479   neuroscience, 2014. **8**: p. 204-204.
480   35.   James, G.M., et al., *Effects of Selective Serotonin Reuptake Inhibitors on*
481   *Interregional Relation of Serotonin Transporter Availability in Major*
482   *Depression.* Frontiers in Human Neuroscience, 2017. **11**(48).
483   36.   Hahn, A., et al., *Attenuated serotonin transporter association between dorsal*
484   *raphe and ventral striatum in major depression.* Human brain mapping, 2014.
485   **35**(8): p. 3857-3866.
486   37.   Hahn, A., et al., *Escitalopram enhances the association of serotonin-1A*
487   *autoreceptors to heteroreceptors in anxiety disorders.* J Neurosci, 2010.
488   **30**(43): p. 14482-9.

Statements & Declarations

Funding

This research was supported by funds from the Eberhard Karls University Tübingen Faculty of Medicine (fortüne 2209-0-0 and 2409-0-0) to HFW, from Bundesministerium für Bildung und Forschung (BMBF, Grant No. 01GQ1415) to HFW and from the Carl Zeiss Foundation to KH.

Competing Interests

The authors authors declare no conflict of interest.

CRediT authorship contribution statement

Tudor M. Ionescu: Conceptualization, Methodology, Software, Validation, Formal analysis, Investigation, Data curation, Writing – original draft, Visualization. **Mario Amend:** Investigation, Writing – review & editing. **Rakibul Hafiz:** Methodology, Software, Writing – review & editing. **Bharat B. Biswal:** Conceptualization, Writing –

review & editing, Supervision, Project administration, Funding acquisition. **Andreas**
Maurer: Writing – review & editing. **Hans F. Wehrl:** Conceptualization,
Methodology, Writing – review & editing, Supervision, Project administration, Funding
acquisition. **Kristina Herfert:** Writing – review & editing, Visualization, Supervision,
Project administration, Funding.

Data availability

The data will be made available upon reasonable request.

508